1 **Dietary inclusion of nitrite-containing frankfurter exacerbates**

2 **colorectal cancer pathology, alters metabolism and causes gut**

3 **dybiosis in APCmin mice.**

5 **Authors:** William Crowe, Xiaobei Pan, James Mackle, Adam Harris, Gary Hardiman,
6 Chris T Elliott, Brian D Green*

8 Institute for Global Food Security, School of Biological Sciences, Queens University

9 Belfast, Biological Sciences Building, Chlorine Gardens, Belfast BT9 5DL, Northern

10 Ireland, UK.

11 *Correspondence to: b.green@qub.ac.uk

Abstract

Colorectal cancer (CRC) is the second most prevelant malignancy in Europe and diet is an important modifiable risk factor. Processed meat consumption, including meats with preservative salts such as sodium nitrite, have been implicated in CRC pathogenesis. This study investigated how the CRC pathology and metabolic status of adenomatous polyposis coli (APC) multiple intestinal neoplasia *(min)* mice was perturbed following 8 weeks of pork meat consumption. Dietary inclusions (15%) of either nitrite-free pork, nitrite-free sausage or nitrite-containing sausage (frankfurter) were compared against a parallel control group (100% chow). Comprehensive studies investigated: gastrointestinal tract histology (tumours, aberrant crypt foci (ACF) and mucin deplin foci (MDF), lipid peroxidation (urine and serum), faecal microbiota and serum metabolomics (599 metabolites). After 8 weeks mice consuming the frankfurter diet had 53% more (P=0.014) gastrointestinal tumours than control, although ACF and MDF did not differ. Urine and serum lipid peroxidation markers were 59% (P=0.001) and 108% (P=0.001) higher, respectively in the frankfurter group. Gut dysbiosis was evident in these mice with comparably fewer *Bacteriodes* and more *Firmicutes*. Fasting serum levels of trimethylamine N-oxide (TMAO) and numerous triglycerides were elevated. Various serum phosphotidylcholine species were decreased. These results demonstrate that nitrite-containing sausages may exaccerbate the development of CRC pathology in APC[Min] mice to a greater extent than nitrite-free sausages, and this is associated with greater lipid peroxidation, wide-ranging metabolic alternation and gut dysbiosis.

Keywords
Processed meat, sausages, colorectal cancer, microbiome, metabolism, metabolomics, gastrointestinal tumours.

Introduction

Colorectal cancer (CRC) is the third most prevalent malignancy worldwide. Over 1.9

million people were diagnosed with CRC in 2020 [1]. CRC always begins in the form

of abnormal growths called polyps in the colon or rectum. The majority of colorectal

tumours are adenocarcinomas that originate from polyps. Polyps are pedunculated

masses of tissue that may be pre-cancerous. Around 5% of CRC is attributable to

familial adenomatous polyposis[2]. The remaining cases of CRC are believed to be

influenced by environmental factors. One modifiable risk factor that has been

proposed as a leading candidate in the search for causes of CRC is dietary pattern. A

diet high in processed meat is suggested as a major risk factor for CRC development.

The International agency for Reseach on Cancer (IARC) have advised consumers to

avoid consuming processed meat and limit the quantity of red meat consumed.

Processed meat is classed as a group 1 carcinogen, and therefore carcinogenic to

humans. This conclusion is based on 9 case-control studies and 18 cohort studies and

of these 6 and 12, respectively, linked processed meat consumption to CRC 3[3].

However, these studies only represent 10% of the existing body of evidence, the

remainder of which was deemed uninformative due to methodological limitations.

IARC list examples of processed meat as hot dogs (frankfurters), ham, sausages,

corned beef, and biltong or beef jerky as well as canned meat and meat-based

preparations and sauces. Processed meat is often preserved with nitrite salts, and

recent scientific evidence suggests that it is this component that is responsible for

processed meats unfavourable link with CRC [4]. Sodium nitrite uniquely prevents the

growth of *Clostridium Botulinum*, it also contributes to the colour and taste of

processed meat. Ingested nitrite has the potential to form N-Nitroso compounds

(NOC) including nitrosamines, some of which are known to be carcinogenic. NOCs

have been proposed to be involved in the aetiology of several types of cancer,

particularly those in the gastrointestinal tract [5].

The composition of gut microbiota (consisting of more than 10^{13} bacteria), is

modulated by exposure to environmental factors such as diet and medication. These

microbes protect the host by inhibiting the growth of pathogens and contributing to

the production of short chain fatty acids (SCFA). SCFAs include acetate, butyrate,

and propionate are used an energy source and participate in the proliferation of

intestinal epithelial cells[6]. Dysbiosis has been implicated in the pathogenesis of

numerous health conditions, including CRC [7]. It has been reported in CRC that there

is a depletion of *Bacteroidetes* and *Firmicutes*, and at the order level there is a

reduction in *Clostridiales* [8]. Sodium nitrite is also known to prevent the colonisation

of *Clostridiales*.

A total of ten animal studies have investigated the effect of nitrite consumption on

CRC, 4 of these studies found that nitrite consumption had no effect on CRC, 5

studies found that nitrite consumption significantly increased the risk of CRC, and 1

study found that nitrite containing processed meat significantly decreased the risk of

CRC[4]. All studies reporting a significant causative effect supplied ≥50% processed

meat in the diet. This does not reflect typical human consumption and therefore

further studies employing more realistic levels of processed meat are required.

Current public health advice treats all processed meat equally, when in reality there

are many differences in their composition. Most notably there is the addition of

sodium nitrite as a preservative which is present in some products and not in others.

For example, nitrites are not traditionally an ingredient in the manufacture of

traditional British/Irish sausages but it is more common in continental European

sausages. Therefore, the primary aim of this investigation was to examine whether

cancer progression in a mouse model of CRC pathology differs when fed a moderate

amount of either nitrite-containing frankfurter sausage or nitrite-free sausage. A

secondary aim was to examine whether CRC development was impacted by the

consumption of either processed (nitrite-free sausage) or unprocessed (pork) meat. A

number of additional analyses were undertaken to ascertain what the wider

metabolic consequences of these meat diets.

Results

Intestinal pathology

In the duodenum the mean tumour count for mice fed a 15% frankfurter diet was 4.2

(Figure 1b). This was 75% higher than the mean tumour count of the control group

(p=0.022). No other inter-group differences were detected for duodenal tumours. The

quantity of tumours in the jejunum did not differ between any groups. Frankfurter

fed animals had 88.2% more tumours in their ilieum than the control (p=0.011), and

190.9% more than pork fed mice (p=0.001), but did not significantly different from

the nitrite-free sausage group. Sausage fed mice had 154.6% more tumours in their

ilieum than the pork fed mice (p=0.050). In the colon, the only significant difference

was that pork fed mice had 81.8% more tumours than the control mice (p=0.047). In

terms of the overall number of GI tumours, the frankfurter group had the highest

with a mean of 11.2 and this was significantly more (p=0.002) than the control which

had a mean of 7.3 tumours, it was also significantly higher (p=0.029) than the

sausage group which had a mean of 8.6 tumours, and significantly higher (p=0.019)

than the pork group which had a mean of 8.3 tumours.

As shown in Table 1 no significant differences were found in ACF or MDF between

any dietary group. There was a trend for pork-fed mice to have the highest quantities

of ACF (not significant), and for sausage-fed mice to have the highest quantities of

MDF (not significant).

Lipid peroxidation

Frankfurter feeding consistently led to the highest concentrations of urinary DHN-

MA (Figure 1c). By week 3 these were 54.30% higher than control (p=0.001), 47.39%

higher than sausage-fed (p=<0.001), and 65.66% higher than pork-fed (p=<0.001). By

week 3 Pork-fed mice had 72.17% higher urinary DHN-MA than sausage-fed

(p=0.048). By week 6 the frankfurter-fed group had DHN-MA levels that were

83.05% higher than the control (p=0.030), 80.26% higher than sausage-fed (p=0.009),

and 80.66% higher than pork-fed (p=0.013). At week 8 the frankfurter group had

DHN-MA levels that were 59.18% higher than the control (p=0.001), 67.92% higher

than sausage-fed (p=<0.001), and 69.07% higher than pork-fed (p=0.001).

Similar results were found when comparing serum MDA concentrations between

groups. Frankfurter-fed mice had the highest mean concentration (928.89 pg/ml) and

this was 48.07% higher than the control (p=<0.001), 42.90% higher than pork-fed

(p=<0.001), and 52.56% higher than sausage-fed (p=0.001). These were all

significantly different after controlling for mouse baseline values.

Bodyweight and food intake

As shown in Figure 1d, mice fed pork or frankfurter diet had significantly higher

body weight than the control and sausage groups at week 1. At almost every

timepoint the frankfurter group had the highest mean bodyweight and this was

significantly higher than the control. At weeks 6, 7, and 8 the sausage group had

significantly lower mean bodyweights than the control. At weeks 2-6 the pork group

had significantly higher mean bodyweights than the control, but not at weeks 7 and

8.

Altered concentrations of serum metabolites

Unsupervised Principal Component Analysis (PCA) of serum metabolomic data

explained 89.4% of the total variation (76.9%, 8.0% and 4.5% for PC1, PC2 and PC3,

respectively; Figure 2a). There was visible separation with a tendency for frankfurter

group profiles to separate from the other three groups along the PC1 axis. Profiles

from control, pork and sausages groups were more closely clustered together with

variation more restricted to the PC2 axis.

Serum metabolites were ranked by false-discovery rate (FDR) corrected p-values

based on one-way ANOVA. Out of 599 measured metabolites 232 significantly

differed between two or more groups. When each of the dietary interventions was

compared with Control it was the frankfurter diet which caused by far the greatest

number of metabolite alternations (Figure 2b). In total 218 metabolite alternations

were observed for frankfurter (165 of which were unique), whereas comparably

fewer were observed for pork and sausage, 52 and 42, respectively. Hierarchical

clustering analysis (HCA) was conducted by selecting the 50 most statistically

significant metabolites from the ANOVA found that the greatest distinction in

profiles was between the control and frankfurter-fed mice (Figure 2c). Metabolic

profiles obtained for pork-fed and sausage-fed mice were similar to one another and

were clustered together by HCA. From this it was evident that many triglyceride

species were significantly higher in the frankfurter group, and several certain

cholestrol esters and phosphotidylcholines were significantly lower.

Figure 3 shows that serum TMAO (trimethylamine N-oxide) levels were 184%

higher in mice receiving frankfurter diet. However, the levels of TMAO precursor

metabolites (choline, betaine, carnitine, acetylcarnitine propionylcarnitine) are not

similarly affected. Levels of choline and betaine did not differ between groups. Mice

consuming any of the meat-containing diets had similarly higher levels of carnitine

(71-119%), acetylcarnitine (114-217%) or propionylcarnitine (97-155%). In no cases

were these carnitine metabolites the highest overall in mice consuming the

frankfurter diet.

Faecal microbiota

181
As Figure 4 shows, there are clear differences between in the profiles (Figures 4a and

4c) and the diversity (Figure 4b) of the frankfurter fed mices microbiota compared

with all other groups. These differences exist on all phylogenetic levels (phylum,

class, order, family, genus, and species). An analysis of covariance (ANCOVA) was

used to compare the mean OTU levels of each phylum from post study between

dietary groups, whilst controlling for baseline levels of each phylum. There was no

significant difference in *Bacteroidetes* from the terminal sample between any groups,

when controlling for baseline. The highest levels of *Firmicutes* were seen in the

frankfurter consuming group, and the lowest levels were seen in the sausage

consuming group, however when controlling for baseline, the there was no

significant differences between any groups. The highest levels of *Actinobacteria* were

seen in the frankfurter consuming group, and this was significantly higher than the

control (p=0.002), sausage (0.001), and pork (<0.001) consuming groups. Similarly,

the highest levels of *Proteobacteria* were seen in the frankfurter group and these were

significantly higher than the pork (P=0.014), and sausage (P=0.011), but not the

control (P=0.551). The control group had significantly higher levels of *Tenericutes*

than the pork (P=0.001), sausage (P=0.001), and frankfurter (P=0.002) fed groups. The

control group had significantly lower levels of *Verrucomicrobia* than the pork

(P=0.551), sausage (P=0.551), and frankfurter groups (P=0.551).

The most abundant bacteria at the species level in the control, sausage and pork

group was *Bacteriodes thetaiotaomicron*, the gut microbiome consisted of a mean (SD)

relative abundance of 16.6 (9.37), 14.7 (9.07), and 19.7 (3.71)% respectively. The

frankfurter group tended to have lower abundance than all 3 other groups at 7.3

(2.9)% (not significant). *Duboisiella newyorkenesis* was the most abundant species in

the frankfurter group, making up a total of 28.8 (1.37)% of all species. This species

was not detectable in any other group. All four groups displayed a high abundance

of bacteria from the *Lachnospiraceae* family, which was largely attributed to higher

levels of bacteria from the *Blautia* genus (see Supplementary Table 1). The sausage

group was the only group with measureable *alistipes*, which was the 3[rd] most

abundant species 11.7 (5.5)%.

The most abundant bacteria at the family level in all four dietary groups is *Clostridiales*,

the next most abundant in the control, sausage, and pork groups is *Bacteroidales*, whilst

Erysipelotrichales is the second most abundant in the frankfurter group.

Discussion

This study found that a modest inclusion of a sodium-nitrite containing pork

frankfurter in the diet of APC^{min} mice significantly increased the number of intestinal

tumours present. A similar inclusion of either nitrite-free pork sausage or minced

pork did not increase tumour numbers. However, the numbers of intestinal foci

(ACF and MDF) did not differ between any groups. Frankfurter consumption was

also associated with higher levels of oxidative stress, as indicated by serum MDA

and urinary DHN-MA, than any other dietary group. Unbiased PCA analysis of

serum metabolomics data showed a distinct clustering of the frankfurter group

which distinguished this diet from all others. This metabolic shift resulted from

vastly more serum metabolites being significantly altered in Frankfurter-fed mice

than any other group. When hierarchical clustering was performed on a shortlist of

the 50 most affect metabolites across all groups it was demonstrated that the

metabolic status of mice fed minced pork and those fed pork sausage were most

similar to one another. It also illustrated the widerspread metabolic divergance of

frankfurter-fed mice. Of the many metabolites profiled trimethylamine N-oxide

(TMAO) stood out, since it has previously has linked to CRC development [9], and

here it was uniquely elevated in the frankfurter group.

Intestinal microbiota play a significant role in TMAO production and

precursor molecules of TMAO found in red meat, eggs, milk and fish. However, the

increases in circulating TMAO here most likely result from alterations in the gut

microbiome. It is clear that all dietary interventions in this study impacted the

composition of faecal microbiota to some extent. However, the reduced microbial

diversity (Shannon index) of the frankfurter group suggested that this diet resulted

in gut dysbiosis. The most remarkable observation was the sheer dominance of *D.*

newyorkenesis in the frankfurter group. The levels of this organism in all other groups

fell well below threshold cut-off values. *D. newyorkenesis* is a poorly characterised

organism and we questioned whether *D. newyorkenesis* is a driver of TMAO

formation which is subsequently detected in the blood circulation. The genome of

this recently identified bactierial species [10] has was sequenced recently (but is not yet

fully annotated). We utilized the makeblastdb application and produced a

searchable BLAST databases from FASTA files obtained from D. newyorkenesis and

carried out pairwise alignments to search for gene sequences encoding enzmyes

involved in TMA-synthesis pathways (including choline TMA-lyase (cutC) and

carnitine oxygenase (cntA)) [11] but we did not find any significant alignments, which

would suggest that this organism is not involved in TMAO production.

Nonetheless, at the genus level *Dubosiella* is positively associated with obesity in

mice [12] and at the 'family' level, *Erysipeltrchaceae*, is positively associated with CRC in

murine models and humans [13,14]. It is not however clear if predominance of

Erysipeltrchaceae is causative of CRC or if colonisation occurs in response to CRC. The

Clostridiales family, although present in all four groups, was noticeably less prevalent

in the frankfurter group. It may be that *Erysipeltrchaceae* to some extent displaced

Clostridiales, but another explanation could be the known ability of nitrites to prevent

the growth of organisms from the *clostridium* genus.

An in-depth literature search reveals that ten pre-clinical studies have thusfar

evaluated the effect of nitrite consumption on the development of CRC. The

deficiency of most of these studies is that cancer was chemically induced and not

spontaneous. None-the-less, five studies reported nitrite exposure to increase cancer

risk [15-19]. Three studies reported that nitrite treatment made no difference [20-22], and

one study reported nitrite to actually be beneficial [23]. Our study has several

advantages over previous studies in this area. Firstly, it is the only study to provide

a nitrite-containing meat matrix to $Apc^{Min+/+}$ mice, which could be described as a

'gold-standard' model of spontaneous CRC development. This minimises the known

unpredictability of cancer development (especially in location and timing) of

chemically-induced CRC models. Secondly, all previous pre-clinical studies linking

nitrite to CRC development, provided a minimium of 50% processed meat in the

diet. Not only are such levels unrealistic for humans, they disproportionately alter

the overall nutritional profile of the diet. Our study has detected a measurable effect

with just 15% dietary inclusion, and although this still represents a relatively high

intake of processed meat it clearly demonstrates that lower dietary quantities can

exacerbate the disease. Thirdly, this study has investigated other relatively

unexplored mechanisms through which diets impacts CRC. The application of both

metabolome and microbiome data makes this study particularly unique. Lipid

peroxidation (i.e. the oxidative destruction of membrane lipids and production of

compounds impairing cells and tissues) was evident and this is in keeping with

other studies [16,18]. The aforementioned rise in TMAO in the frankfurter group could

also be linked linked to oxidative stress [24]. TMAO is generated from the metabolism

of a number of nutrients including carnitine and choline which are present in pork

meat. Increased TMAO in the circulation apparently induces superoxide production,

a reactive oxygen species linked to oxidative stress [24]. TMAO's production of reactive

oxygen species has been confirmed in cells *in vitro* [25]. The association between

TMAO, oxidative stress and cancer still needs to be verified, but mounting evidence

suggests that TMAO could play a role in chronic inflammation as a cause of

carcinogenesis in CRC [26]. Furthermore, *in silico* studies suggest that TMAO

metabolism and CRC share many similar genetic pathways [27].

In sharp contrast to the frankfurter group, the minced pork and pork sausage

group did not display elevated serum TMAO. The TMAO responses did not mirror

those of common TMAO precusors. Choline levels were the same in all groups. As

would be expected carnitine (and several acylcarnitines) were high in all three pork

diet interventions. Another TMAO precursor betaine (found typically in plants but

not in meat) did not differ in any groups either. Therefore, in these experiments the

supply of precursors does not appear to be important factor in TMAO production,

and the suspicion would be that altered microbiota are more pivotal. A limitation of

this study is the fact that the microbial profiling undertaken involved the analysis of

16S rRNA amplicon rather than a whole-metagenomics approach, thus it was not

possible to evaluate whether there was an abundance of bacterial species with genes

encoding the enzymatic machinery involved in TMAO production.

An obvious advantage of the present study design is that it reflects real-world

human food choices. To this end it directly compared commercially available pork

products on an equivalent gram-for-gram basis. The unfortunate consequence of this

approach is that the nutritional properties of the pelleted diets cannot be precisely-

matched, or be isocaloric, which is a potential limitation. Another limitation is that is

was not possible to record food intake, and we cannot pinpoint the underlying

reasons for the higher bodyweights of mice in the frankfurter group. It is possible

that mice found this pelleted diet more palatable and consumed more, but there are

also differences in the levels of fibre, and the types of fat in the diets which could

have contributed also to this effect. It is not known if raised bodyweight exacerbated

CRC pathology in this group, but this is very questionable given that the pork

sausage group had the lowest body weight, but also the second highest number of

tumours overall. There were metabolic differences evident in the frankfurter group,

for instance, the significantly elevated diglycerides and triglycerides, which were

likely explained by the higher saturated fat content of this diet. The decreased

phosphatidylcholine and lysophosphatidylcholine (lysoPC) levels may also be a

consequence of this, but equally, similar observations have been noted in CRC

patients [28]. Increased lysoPC metabolism in tumour cells *in vitro* is one explanation

for a decline in their blood concentrations.

The results from this study clearly show that not all processed meats carry the same

risk of CRC development, and that the consumption of nitrite-containing processed

meat exacerbates CRC pathology in a genetically predisposed murine model.

However, higher dietary fat and increased body weight cannot completely be ruled

out as potential confounding factors. The metabolomic and gut microbiome data

demonstrate the systemic alterations that can occur as result of simple dietary

choices. These also provide important mechanistic data. Further dose-response

studies should be performed, perhaps using humanised or colon-specific CRC

mouse models, and employing frankfurter as a model food matrix. Such work is

required to determine the levels at which nitrite inclusion becomes harmful.

Methods

Animal models

Forty female $Apc^{Min/+}$ mice were purchased from Charles River and delivered to Axis

Bioservices (Coleraine, Northern Ireland) where the study was conducted. All mice

were four weeks old on delivery. The mice were all fed an ad lib maintenance diet of

AIN76 for a two week acclimitization period (Figure 1a). Mice were caged in groups

of 3 or 4 in a 12 hour light, 12 hour dark cycle. Mice had unlimited access to their

respective diets and water for the duration of the study which was 12 weeks, with

the exception of 12 hours prior to blood collection, when they were fasted. Body

weight was measured every 3–4 days. We attempted to measure food intake in a

similar manner, but since diets were required (for welfare reasons) to be placed in

dishes in home cages this led to extremely erroneous data and was subsequently

abandoned. Mice were randomly assigned to one of four treatment groups,

described below. Studies in mice underwent Ethical Review and were performed in

accordance with the UK Animals (Scientific Procedures) Act 1986.

Diets

Each treatment group received one of four diets for the duration of the study, the

dietary groups included; Control, which consisted of 100% AIN76 (Altromin,

Germany). Sausage meat (15%)/AIN76 (85%), Frankfurter (15%)/AIN76 (85%),

unprocessed pork (15%)/AIN76 (85%). Sausage meat was manufactured and

provided by Finnebrogue Artisan (Downpatrick, UK) as was the unprocessed pork.

Nitrite-containing Frankfurter was manufactured by Herta (Illkirch, France) and

purchased from a local supermarket. All meat was freeze dried by European Freeze

Dry (UK), and powderized. Altromin then incorporated this into pellet composed of

15% meat powder and 85% AIN76. The nutritional properties of each diet were

analysed after pelleting (Supplementary Table 2).

Biological samples

Blood samples were collected from the tails of all animals at 3 time points, week 0,

week 3, week 6, a final cardiac puncture blood was taken at the termination of the

study (week 8). Urine was collected from individual animals at baseline and week 8.

Faecal samples at the same timepoints but were collected from each cage.

Tumor scoring in the gastrointestinal tract

Animals were sacrificed using CO_2 asphyxiation and their organs harvested. As

described previously[29], the intestinal tract was removed washed in phosphate

buffered saline and sectioned into the duodenum, jejunum, ileum, and colon. The

sections were opened longitudinally and fixed in 10% formalin, colons were stained

for 5 minutes in 0.05% methylene blue solution (Sigma), and small intestine were

stained for 48 hours in 0.05% methylene blue. Using x25 magnification all tumors

present were counted by one researcher in a blinded manner. The minimum size for

accepting and counting a tumor was 0.5mm diameter.

Identification of aberrant crypt foci and mucin depleted foci

Aberrant crypt foci (ACF) were measured in the colon by a well described method [30].

The colons were washed in saline, and stained with 0.2% methylene blue for 5 mins.

They were placed on a glass slide and visualised using a magnification of 32x on a

light microscope. Mucin depleted foci (MDF) were also measured in the colon, the

sections were washed in saline, and stained with 1% high-iron diamine alcian blue

procedure for 30 mins. All scoring was completed by one researcher and verified by

a second researcher in a blinded manner.

Lipid peroxidation

Serum samples from every time point were analysed for malondialdehyde (MDA)

using ELISA methodologies (Abcam, Cambridge UK). In brief, thiobarbituric acid,

serum and standards were added to 96-well microtiter plate and incubated at 95^{o}C

for 60 mins and cooled on ice. An aliquot from each well was added to the

colorimetric plate and measured at OD532 nm using a plate reader. Urine samples

from baseline and week 8 were analysed for 1,4-dihydroxynonane mercapturic acid

(DHN-MA) using a Bertin Bioreagent enzyme immunoassay (Wolverhampton UK).

Briefly, buffer, standards and urine were added to each well, the detection antibody

was then added, and the plate was incubated overnight at 4^{o}C. Ellman's reagent was

added, and the absorbance was measured at 414nm.

Metabolite profiling

The MxP Quant 500 kit (Biocrates, Life science AG, Innsbruck, Austria) was used to

quantify amino acids, acylcarnitines, bile acids, biogenic amines, carboxylic acids,

ceramides, cholesterol esters, diacylglycerols, fatty acids, glycerophospholipids,

glycosylceramides, sphingolipids, and triacylglycerols in serum samples from

baseline and week 8. Serum (10μL) samples were analysed using a triple-quadrupole

mass spectrometer (Xevo TQ-S, Waters Corporation, Milford, CT, USA). Samples

were dispensed into a 96-well plate, dried at room temperature for 30 min and then

50 μL of phenylisothiocyanate (PITC) was added. After a further 20 min incubation

at room temperature samples were dried under nitrogen for 1 h. Ammonium acetate

(3 mM) was added, and the plate was shaken at room temperature for 30 min and

centrifuged for 2 min at 500 g. Each well then had 150 μL transferred from it to a 96

deep well plate along with and equal volume of dH_2O. A 96-well plate containing

490 μL of FIA solvent is shaken along with 10 μL of extract at room temperature for

5 mins. From each well 5 μL of sample Samples were injected through the MxP

Quant 500 kit column system for amino acids and biogenic amines analysis. Other

metabolites were semi-quantified using the same mass spectrometer without column

separation by the flow injection analysis (FIA). Metabolite concentrations were

calculated using the Analyst/MetIDQ software and expressed as μM in serum.

Microbial profiling

A QIAamp DNA stool mini kit (Qiagen, Manchester, UK) was used to extract DNA

from faecal samples. All reagents, unless otherwise stated were provided by Qiagen.

Faecal samples were weighed (220mg) and added to a 2ml Eppendorf tube along

with 1.4 ml of ASL buffer, which was vortexed and heated for 5 mins at 95°C. The

contents were added to a 2 ml screw capped tube containing 0.5 g of both 0.1 mm

and 0.5 mm zirconium beads. The tube was then added to a FastPrep 24 g5 bead

beater (MP biomedicals, Eschwege, Germany) for 5 x 20 s bursts. The suspension

was subsequently spun at 14,000 g for 3 mins. The supernatant (1.2 ml) was collected

and transferred into a QIAamp silica-gel membrane tube and spun for 3 min, and

DNA extracted in accordance with the manufacturer's instructions. Amplicon

sequencing was carried out on an Illumina MiSeq. Variable regions V4-V5 of the

bacterial 16S ribosomal RNA gene were amplified from extracted DNA using PCR conditions

and custom primers as described in the Microbiome Helper protocol [31]. The forward (515FB =

GTGYCAGCMGCCGCGGTAA) and reverse (926R = CCGYCAATTYMTTTRAGTTT)

primers used Nextera Illumina index tags and sequencing adapters fused to the 16S sequences.

Each sample was amplified with a different combination of index barcodes to allow for sample

identification after multiplex sequencing. After amplification, paired-end 300 + 300 bp v3

sequencing was performed for all samples on an Illumina MiSeq. Raw sequencing reads

were trimmed as previously described [32].

Analysis of 16S sequencing data

Analysis of 16S sequencing data was carried out on the Hardiman Laboratory

GRANDE server. Primer sequences were removed from the sequencing reads using

cutadapt (v 1.14) [33], and primer-trimmed files were imported into QIIME2 (v.

2019.10.0) [34]. Forward and reverse paired-end reads were joined using VSEARCH (v

2.9.0) [35] and inputted to Deblur [36] to correct reads and obtain amplicon sequence

variants (ASVs). Taxonomy was assigned to ASVs using the SILVA rRNA gene

database [37] and the "feature-classifier" option in QIIME2, sklearn. Rare sequences

were removed from the feature table. Any sequences below an arbitrary cut-off

frequency of 100 were also removed. Estimates of alpha-diversity (Observed ASVs,

Shannon Diversity), beta-diversity (weighted UniFrac), and relative abundance of

ASVs were obtained using QIIME2 (v 2019.10.0). Differential abundance tests were

done using the Analysis of compositions of microbiomes (ANCOM)plugin [38].

Statistical analysis

All data were tested for normality using Kologorov-Smirnoff and where appropriate

data was transformed. Group comparisons for tumour numbers, ACF and MDF

were performed by a One-way ANOVA with LSD post-hoc test (two-tailed; SPSS

version 21). Lipid peroixdation markers DHN-MA and MDA at week 8 were

compared between groups whilst controlling for baseline concentrations by an

ANCOVA. Metabolomic data were cube-root transformed, underwent mean-

centered scaling and compared using univariate (ANOVA), multivariate and

hierarchial clustering tools within MetaboAnalyst 4.0 (http://metaboanalyst.ca).

Data Availability Statement

Raw mass spectrometry data is available on the Metabolomics Workbench
(https://www.metabolomicsworkbench.org/) under study ST002313,
[http://dx.doi.org/doi: 10.21228/M81M72]. The 16S rRNA Seq data have been
submitted to the European Nucleotide Archive (ENA), accession number
PRJEB56215.

Acknowledgements

Studies investigating links between meat consumption and CRC are in-part
supported by Agri-Food QUEST, a membership-based, industry-led Innovation
Centre for agri-food business in Northern Ireland. Agri-Food QUEST is supported
by funding from Invest Northern Ireland (InvestNI) but also were partially
supported by commercial funding from: Finnebrogue Artisan, Karro Food group,

465 and Cranswick. These industrial partners were not involved design or writing of the
466 manuscript, the analysis/interpretation of the data, or the decision to publish.

Competing Interests

469 W.C., X.P., G.H., J.M., A.H, and B.D.G declare no competing interests. C.T.E.
470 declares that he is an editor for npj Science of Food, but was not involved in the
471 journal's review of, or decisions related to, this manuscript.

Author Contributions

473 W.C. organised, managed and conducted laboratory work. X.P. conducted
474 metabolomic studies. G.H., J.M. and A.H. performed metagomic sequencing
475 analysis. W.C. and B.D.G. wrote the manuscript. B.D.G, G.H. and C.T.E edited the
476 manuscript. All authors approved the final text of the manuscript.

References

References

1   Sung, H. *et al.* Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and
Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* **71**, 209-249 (2021).
https://doi.org:10.3322/caac.21660

2   Jeter, J. M., Kohlmann, W. & Gruber, S. B. Genetics of colorectal cancer. *Oncology (Williston*
Park) **20**, 269-276; discussion 285-266, 288-269 (2006).

3   Bouvard, V. *et al.* Carcinogenicity of consumption of red and processed meat. *Lancet Oncol*
16, 1599-1600 (2015).

4   Crowe, W., Elliott, C. T. & Green, B. D. A Review of the In Vivo Evidence Investigating the
Role of Nitrite Exposure from Processed Meat Consumption in the Development of
Colorectal Cancer. *Nutrients* **11** (2019). https://doi.org:10.3390/nu11112673

5   Jakszyn, P. *et al.* Endogenous versus exogenous exposure to N-nitroso compounds and
gastric cancer risk in the European Prospective Investigation into Cancer and Nutrition (EPIC-
EURGAST) study. *Carcinogenesis* **27**, 1497-1501 (2006).
https://doi.org:10.1093/carcin/bgl019

6   Pothuraju, R. *et al.* Mucins, gut microbiota, and postbiotics role in colorectal cancer. *Gut*
Microbes **13**, 1974795 (2021). https://doi.org:10.1080/19490976.2021.1974795

7   Castellarin, M. *et al.* Fusobacterium nucleatum infection is prevalent in human colorectal
carcinoma. *Genome Res* **22**, 299-306 (2012). https://doi.org:10.1101/gr.126516.111

8   Kostic, A. D. *et al.* Genomic analysis identifies association of Fusobacterium with colorectal
carcinoma. *Genome Res* **22**, 292-298 (2012). https://doi.org:10.1101/gr.126573.111

9   Bae, S. *et al.* Plasma choline metabolites and colorectal cancer risk in the Women's Health
Initiative Observational Study. *Cancer Res* **74**, 7442-7452 (2014).
https://doi.org:10.1158/0008-5472.CAN-14-1835

10   Cox, L. M. *et al.* Corrigendum: Description of two novel members of the family
Erysipelotrichaceae: Ileibacterium valens gen. nov., sp. nov. and Dubosiella newyorkensis,
gen. nov., sp. nov., from the murine intestine, and emendation to the description of
Faecalibacterium rodentium. *Int J Syst Evol Microbiol* **67**, 4289 (2017).
https://doi.org:10.1099/ijsem.0.002339

11   Rath, S., Heidrich, B., Pieper, D. H. & Vital, M. Uncovering the trimethylamine-producing
bacteria of the human gut microbiota. *Microbiome* **5**, 54 (2017).
https://doi.org:10.1186/s40168-017-0271-9

12   Wang, W. *et al.* Optimal Dietary Ferulic Acid for Suppressing the Obesity-Related Disorders in
Leptin-Deficient Obese C57BL/6J -ob/ob Mice. *J Agric Food Chem* **67**, 4250-4258 (2019).
https://doi.org:10.1021/acs.jafc.8b06760

13   Chen, W., Liu, F., Ling, Z., Tong, X. & Xiang, C. Human intestinal lumen and mucosa-
associated microbiota in patients with colorectal cancer. *PLoS One* **7**, e39743 (2012).
https://doi.org:10.1371/journal.pone.0039743

14   Zhu, Q. *et al.* Analysis of the intestinal lumen microbiota in an animal model of colorectal
cancer. *PLoS One* **9**, e90849 (2014). https://doi.org:10.1371/journal.pone.0090849

15   Santarelli, R. L. *et al.* Calcium inhibits promotion by hot dog of 1,2-dimethylhydrazine-
induced mucin-depleted foci in rat colon. *Int J Cancer* **133**, 2533-2541 (2013).
https://doi.org:10.1002/ijc.28286

16   Santarelli, R. L. *et al.* Meat processing and colon carcinogenesis: cooked, nitrite-treated, and
oxidized high-heme cured meat promotes mucin-depleted foci in rats. *Cancer Prev Res*
(Phila) **3**, 852-864 (2010). https://doi.org:10.1158/1940-6207.Capr-09-0160

17   Pierre, F. H. *et al.* Calcium and α-tocopherol suppress cured-meat promotion of chemically
induced colon carcinogenesis in rats and reduce associated biomarkers in human volunteers.
Am J Clin Nutr **98**, 1255-1262 (2013). https://doi.org:10.3945/ajcn.113.061069

18 Pierre, F. H. *et al.* Freeze-dried ham promotes azoxymethane-induced mucin-depleted foci
and aberrant crypt foci in rat colon. *Nutr Cancer* **62**, 567-573 (2010).
https://doi.org:10.1080/01635580903532408
19 Bastide, N. M. *et al.* Red Wine and Pomegranate Extracts Suppress Cured Meat Promotion of
Colonic Mucin-Depleted Foci in Carcinogen-Induced Rats. *Nutr Cancer* **69**, 289-298 (2017).
https://doi.org:10.1080/01635581.2017.1263745
20 Chenni, F. Z. *et al.* Heme-induced biomarkers associated with red meat promotion of colon
cancer are not modulated by the intake of nitrite. *Nutr Cancer* **65**, 227-233 (2013).
https://doi.org:10.1080/01635581.2013.749291
21 Parnaud, G., Peiffer, G., Taché, S. & Corpet, D. E. Effect of meat (beef, chicken, and bacon)
on rat colon carcinogenesis. *Nutr Cancer* **32**, 165-173 (1998).
https://doi.org:10.1080/01635589809514736
22 Bastide, N. M. *et al.* A central role for heme iron in colon carcinogenesis associated with red
meat intake. *Cancer Res* **75**, 870-879 (2015). https://doi.org:10.1158/0008-5472.CAN-14-
2554
23 Iscovich, J. M. *et al.* Colon cancer in Argentina. I: Risk from intake of dietary items. *Int J
Cancer* **51**, 851-857 (1992). https://doi.org:10.1002/ijc.2910510603
24 Li, T., Chen, Y., Gua, C. & Li, X. Elevated Circulating Trimethylamine N-Oxide Levels
Contribute to Endothelial Dysfunction in Aged Rats through Vascular Inflammation and
Oxidative Stress. *Front Physiol* **8**, 350 (2017). https://doi.org:10.3389/fphys.2017.00350
25 Yue, C. *et al.* Trimethylamine N-oxide prime NLRP3 inflammasome via inhibiting ATG16L1-
induced autophagy in colonic epithelial cells. *Biochem Biophys Res Commun* **490**, 541-551
(2017). https://doi.org:10.1016/j.bbrc.2017.06.075
26 Chan, C. W. H. *et al.* Trimethylamine-N-oxide as One Hypothetical Link for the Relationship
between Intestinal Microbiota and Cancer - Where We Are and Where Shall We Go? *J
Cancer* **10**, 5874-5882 (2019). https://doi.org:10.7150/jca.31737
27 Xu, R., Wang, Q. & Li, L. A genome-wide systems analysis reveals strong link between
colorectal cancer and trimethylamine N-oxide (TMAO), a gut microbial metabolite of dietary
meat and fat. *BMC Genomics* **16 Suppl 7**, S4 (2015). https://doi.org:10.1186/1471-2164-16-
s7-s4
28 Zhao, Z. *et al.* Plasma lysophosphatidylcholine levels: potential biomarkers for colorectal
cancer. *J Clin Oncol* **25**, 2696-2701 (2007). https://doi.org:10.1200/jco.2006.08.5571
29 Reima, H., Saar, H., Innos, K. & Soplepmann, J. Methylene blue intra-arterial staining of
resected colorectal cancer specimens improves accuracy of nodal staging: A randomized
controlled trial. *Eur J Surg Oncol* **42**, 1642-1646 (2016).
https://doi.org:10.1016/j.ejso.2016.06.001
30 Bird, R. P. Observation and quantification of aberrant crypts in the murine colon treated with
a colon carcinogen: preliminary findings. *Cancer Lett* **37**, 147-151 (1987).
https://doi.org:10.1016/0304-3835(87)90157-1
31 Comeau, A. M., Douglas, G. M. & Langille, M. G. Microbiome Helper: a Custom and
Streamlined Workflow for Microbiome Research. *mSystems* **2** (2017).
https://doi.org:10.1128/mSystems.00127-16
32 Huse, S. M., Huber, J. A., Morrison, H. G., Sogin, M. L. & Welch, D. M. Accuracy and quality of
massively parallel DNA pyrosequencing. *Genome Biol* **8**, R143 (2007).
https://doi.org:10.1186/gb-2007-8-7-r143
33 Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17**, 10-12 (2011). https://doi.org:https://doi.org/10.14806/ej.17.1.200.
34 Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data science
using QIIME 2. *Nat Biotechnol* **37**, 852-857 (2019). https://doi.org:10.1038/s41587-019-
0209-9

35   Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. VSEARCH: a versatile open source
tool for metagenomics. *PeerJ* **4**, e2584 (2016). https://doi.org/10.7717/peerj.2584
36   Amir, A. *et al.* Deblur Rapidly Resolves Single-Nucleotide Community Sequence Patterns.
mSystems **2** (2017). https://doi.org/10.1128/mSystems.00191-16
37   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing
and web-based tools. *Nucleic Acids Res* **41**, D590-596 (2013).
https://doi.org/10.1093/nar/gks1219
38   Mandal, S. *et al.* Analysis of composition of microbiomes: a novel method for studying
microbial composition. *Microb Ecol Health Dis* **26**, 27663 (2015).
https://doi.org/10.3402/mehd.v26.27663

Figure Legends

Figure 1: APC^{Min} mice consuming 15% frankfurter (85% chow) for 8 weeks have
significantly more gastrointestinal tumours, higher levels of lipid peroxidation,
and greater body weight than APC^{Min} mice consuming a control diet (100% chow).
a) after acclimatisation mice were fed one of four chow diets which were either
Control (n=9; standard chow (red)), or chow with a 15% incorporation of pork mince
(n=9; blue), pork sausage (n=10; light blue), or nitrite-containing frankfurter sausage
(n=10; green). **b)** numbers of tumours did not differ between any groups in any of
the specific regions of the GI tract. However, compared with control signifcantly
more tumours were displayed by the frankfurter group alone. **c)** terminal serum
MDA concentrations of mice consuming frankfurter were elevated compared to that
of control terminal samples, after controlling for baseline using an ANCOVA. Urine
DHN-MA concentrations were compared between groups at the same timepoint
using an ANOVA. Levels in mice consuming frankfurter supplemented chow had
higher levels at 28 and 56 days. **d)** bodyweights of frankfurter-fed mice were
consistently higher than control mice. Error bars shown represent s.e.m.

Figure 2 Serum metabolite profiles of frankfurter-fed mice deviate most from
control, whereas those fed pork or sausage align more closely. Serum collected at
56 days under went metabolomic profiling: **a)** shows scores plots of unsupervised
PCA based on the first three principle components (PCs) for Control (n=9; standard
chow (red)), or chow with a 15% incorporation of pork mince (n=9; blue), pork
sausage (n=10; light blue), or nitrite-containing frankfurter sausage (n=10; green).
Metabolomic data explained 89.4% of the total variation (76.9%, 8.0% and 4.5% for
PC1, PC2 and PC3, respectively). There was a tendency for profiles from the
frankfurter group to separate along the PC1 axis. Profiles from control, pork and
sausages groups clustered more closely with variation more restricted to PC2. **b)** is a
Venn diagram showing the number of the 599 metabolites measured which
significantly differed in group comparisons with control. c) heat map with
hierarchical clustering based on the top 50 metabolites ranked by FDR –corrected P-

value (ANOVA). Each row represents a metabolite and each column a sample. Red (high) and blue (low) indicates relative concentrations. Dendrograms indicating similarity are shown at top (samples) on left side (metabolites). Serum metabolite measurements readily clustered mice by diet with pork and nitrite-free sausage being most similar to one another. Frankfurter consumption was characterised by significantly higher levels of certain species of diglycerides (DG), triglycerides (TG) and acylcarnitines (C), and by lower levels of certain phosphatidylcholines (PC/LysoPC) and cholesterol esters (CE). Other notable observations included the differing bile acid responses of taurocholic acid (TCA) and cholic acid (CA). The profound variability of trimethylamine N-oxide (TMAO) responses in frankfurters was also very evident. Control samples are shown in red, minced pork in dark blue, nitrite-free sausage in light blue, and frankfurter (nitrite-containing) in green.

Figure 3 Serum TMAO (trimethylamine N-oxide) is higher and more variable in mice receiving frankfurter diet, yet TMAO precursor metabolites are not similarly affected. Serum levels of TMAO (at 56 days), typically formed from choline, betaine and carnitines were significantly higher in mice receiving frankfurter-containing chow. Levels of choline and betaine did not differ between groups. Mice consuming any meat-containing chow had higher levels of carnitine, acetylcarnitine, and propionylcarnitine. Control samples are shown in red (n=9), minced pork in dark blue (n=10), nitrite-free sausage in light blue (n=9), and frankfurter (nitrite-containing) in green (n=10). Data shown are violin plots containing individual samples values with the median and quartile values represented as dotted lines.

Figure 4 Faecal microbiome profiles differ most in frankfurter-fed mice with reduced micribial diversity and a profound shift in abundance of taxa between feeding groups. Faecal samples were collected at 56 days from each mouse cage for metagenomic sequencing providing cage replicates (n=3) for each treatment. **a)** shows scores plots of unsupervised PCA based on the first three principle components (PCs). Species level microbiome data explained 73.4% of the total variation (42.9%, 15.5% and 15.0% for PC1, PC2 and PC3, respectively). **b)** the Shannon diversity index as calculated by Qiime 2 indicates a reduction in microbial diversity in the frankfurter-fed animals. Individual values are shown along with means (dotted lines), and the error bars shown represent s.e.m. **c)** Relative abundance of microbial taxa is shown at the level of the phylum, the family, and the species. The taxa differed most profoundly in the frankfurter-fed animals. Bacterial species with an abundance of <1% were aggregated and referred to as 'All others' (white bars in figure). At the species level the dominance of *Dubiosiella Newyorkenesis* in the frankfurter group was a particularly striking observation. Control samples are shown in red, minced pork in dark blue, nitrite-free sausage in light blue, and frankfurter (nitrite-containing) in green. uncult. – uncultured bacteria; ambig. – ambiguous taxa.

Table 1. Numbers of aberrant crypt foci and mucin depleted foci did not significantly differ between any groups

	Control	Pork	Sausage	Frankfurter
ACF	102.44 (10.86)	115.11 (17.03)	101.80 (10.71)	101.90 (15.18)
MDF	6.11 (2.76)	5.98 (2.67)	6.5 (2.59)	5.08 (3.12)

ACF- aberrant crypt foci; MDF- mucin depleted foci. Mean values were compared between groups by analysis of variance, alpha value of less than 0.05 was accepted as statistically significant. Values in parenthesis are standard deviations for the corresponding mean.

(a) Study plan

(b) Gastrointestinal tumours

(c) Oxidative stress

(d) Body weight

(a) Perturbed metabolite profiles

(b) no. of metabolites affected

(c) Clustering based on metabolite profiles

(a) Altered faecal microbiota

(b) Reduced microbial diversity

(c) Profound changes in microbial composition

Phylum

Frankfurter group
Pork group
Sausage group
Control group

0   25   50 (%)   75   100

■ Firmicutes ■ Bacteroidetes ■ Verrucomicrobia ■ Actinobacteria

Family

Frankfurter group
Pork group
Sausage group
Control group

0   25   50 (%)   75   100

■ Clostridiales ■ Bacteroidales ■ Verrucomicrobiales ■ Erysipelotrichales
■ Lactobacillales ■ Coriobacteriales ■ Bifidobacteriales ■ Betaproteopbacteriales
■ Bacillales

Species

Frankfurter group
Pork group
Sausage group
Control group

0   25   50   75   100

■ D. newyorkensis ■ Thetaiotaomicron ■ Lachnospiraceae
■ Blautia ■ Muribaculaceae uncult. ■ Akkermansia uncult.
■ Romboutsia uncult. ■ Clostridium sensu stricto uncult. ■ Ruminiclostridium uncult.
■ Ruminiclostridium_9 uncult. ■ Lachnospiraceae_uncult. ■ L. johnsonii
■ Ruminiclostridium_9 ■ Alistipes ■ Erysipelatoclostridium ambig.
■ Ruminococcaceae uncult. ■ Coprostanoligenes ■ Lachnoclostridium uncult.
■ Parasutterella uncult. □ All others