Novel diversity panel facilitates genomic dissection of lodging in tef (*Eragrostis tef*)

Shiran Ben-Zeev[1], Timo Hellwig[1,2,*], Muluken Demeile[1,3], Vered Barak[1] Sasha Vorobyova[4], Sariel Hübner[4], Yehoshua Saranga[1,**]

[1] Robert H. Smith Faculty of Agriculture, Food and Environment, The Hebrew University of Jerusalem, Israel

[2] Volcani Center, Agricultural Research Organization, Israel

[3] Ethiopian Institute of Agricultural Research, Ethiopia

[4] Galilee Research Institute (MIGAL) and Tel-Hai College, Israel

* Current affiliation: Institute for Plant Genetics, Heinrich Heine University Düsseldorf, Germany

** Corresponding: Y. Saranga, shuki.saranga@mail.huji.ac.il

Email addresses:

S. Ben-Zeev, shiran.benzeev@mail.huji.ac.il; T Helwig, t.hellwig@posteo.de; M. Demeile, muluken.alemu@mail.huji.ac.il; V. Barak, vered1503@gmail.com; S. Voroboyova, sashi398@gmail.com; S. Hübner, sarielh@migal.org.il; Y. Saranga, shuki.saranga@mail.huji.ac.il

Submission date:

Manuscript content: 2 tables, 7 figures

Supplementary materials: 2 tables, 2 figures

Word count: 3702

Highlights

Here we report on the assembly and genomic characterization of a tef diversity panel (TDP-300). This unique resource was used for GWAS to identify genomic regions associated with plant lodging.

Abstract

Underutilized species that are not widely cultivated (known as orphan crops) present opportunities to increase crop diversity and food security. Tef [*Eragrostis tef* (Zucc.) Trotter] is known for its high-quality grain and forage. Root-borne lodging is a devastating problem in tef cultivation, leading to large economic losses and limiting its widespread adoption. The aim of this study was to identify genomic regions that are associated with tef lodging. A diversity panel comprised of 297 lines (TDP-300) was assembled, genotyped, and phenotyped across 4 field environments. Genome-wide association study identified 29 genomic regions associated with lodging; in all cases with a minor allele conferring reduced lodging. The eleven most prominent sites were located in or near genes, 5 of them with a putative role, of which 3 were to be associated with root development. The identification of lodging-related sites in the current study may advance understanding of the mechanisms underlying tef lodging and crop improvement. To the best of our knowledge, this is the first diversity panel developed for tef. This platform is expected to be useful for detecting genomic regions and markers associated with traits of interest in tef, as well as for breeding and molecular investigations.

Key words

24   diversity panel, GBS, GWAS, lodging, tef, teff

Abbreviations

GWAS – Genome wide association study

LI – lodging index

SNP – Single nucleotide polymorphism

TDP-300 – tef diversity panel 300

VCF – variant calling file

WAE – weeks after emergence

WL - water-limited treatment

WW - well-watered treatment

Introduction

Sustaining agricultural productivity and food security is becoming ever more challenging for many reasons, including climate change (Rosenzweig *et al.*, 2014), lack of diversity (Renard and Tilman, 2019; Tadele, 2019), diminishing soil and water resources (Wuepper *et al.*, 2020), increasing human population (Tilman *et al.*, 2011) and changing global consumer demands (Khoury *et al.*, 2014). These dramatic changes are affecting the world's major crops - maize, wheat and rice (Zhang *et al.*, 2021). A viable strategy to enhance agricultural productivity and food security is to increase crop diversity (Huang *et al.*, 2002), with a particular emphasis on resilience to challenging environmental conditions such as waterlogging, drought and salinity (Smith *et al.*, 1989; Renard and Tilman, 2019; Girija *et al.*, 2021). The diversification of crops grown for human and animal consumption is important ecologically as well as nutritionally (Khoury *et al.*, 2014). The importance of crop diversity is even more salient in light of the current climate change (Kamenya *et al.*, 2021).

Underutilized species that are not widely cultivated (known as orphan crops) present opportunities to increase crop diversity. The reasons for their minor importance in global agriculture have been debated (Tadele, 2019; Allaby, 2021; Milla and Osborne, 2021), but there is consensus on their potential to improve food production and security. Some information on growing conditions, genetics and ecology is available for a number of orphan crops (Mabhaudhi *et al.*, 2019). Combining the existing information with genomics and genetics could help convert them into large-scale crops, thus benefiting from both their diversity and their resilience (Varshney *et al.*, 2012; Dawson *et al.*, 2019). An important tool for the introduction of a new crop into modern agriculture, and to support its breeding and development, is the genomic database (Brook *et al.*, 2008). Genomic insights into the control of various traits have been proposed as a target for orphan crops (Varshney et al. 2012). Genome-wide association studies (GWAS) have long been used to expose the genetic basis of traits (Ikegawa, 2012), and the use of genomic data derived from genotyping-by-sequencing (GBS) assays for GWAS is an accepted method (Bellucci *et al.*, 2015). Recently, GWAS have been applied to several African orphan crops: fonio millet (*Digitaria exilis*) (Abrouk *et al.*, 2020), finger millet (*Eleusine coracana*) (Puranik *et al.*, 2020) and pigeon pea (*Cajanus cajan*) (Zhao *et al.*, 2020). The key role of orphan crops in areas that are unsuitable for common major crops suggests their tremendous potential for introduction into challenging environments. In addition, orphan crops contain genes that may help develop similar resilience in major crops (Jamnadass *et al.*, 2020; Kamenya *et al.*, 2021).

Tef [*Eragrostis tef* (Zucc.) Trotter] is an allotetraploid C4 (2n = 20) cereal crop with a genome size

of 622 Mbp (VanBuren *et al.*, 2020), mostly grown in Ethiopia where it was presumably

domesticated (D'Andrea, 2008). Tef grains are gluten-free, and contain high amounts of nutritious

compounds and all nine essential amino acids (Reda, 2014; Shumoy *et al.*, 2018; Kebede, 2021);

it is therefore commonly referred to as a "superfood". In addition, tef is an excellent forage grass

(Ditsch, 2007; Matthew Davidson, 2018). Tef grows in a wide range of environments, from sea

level to over 2000 m elevation, with seasonal precipitation of 200 to over 2000 mm, and from

drought-prone areas and saline soils to water logging conditions (Ketema, 1983*a*, 1993; Assefa

et al., 2013). These characteristics have led to an interest in adopting tef in countries outside the

Horn of Africa (Miller, 2008; Norberg *et al.*, 2009; Brown, 2017; Ben-Zeev *et al.*, 2018).

Adapting a traditional crop, such as tef, to modern, highly mechanized cultivation requires

reevaluation and calibration of agronomic practices in tandem with genomic studies and breeding.

The introduction of tef as a new crop to irrigated Mediterranean conditions presents many

challenges, at each step of its growth, some of which have been targeted in our recent studies

(Ben-Zeev *et al.*, 2020*a,b*). The relatively minor attention paid to tef in research, as well as the

lack of a reference genome and diversity panel, has hindered studies on the genomic basis of

key traits in this crop plant (Girija *et al.*, 2021). Nevertheless, some progress has been made in

these aspects over the last few decades. AFLP (Bai *et al.*, 1999), RFLP (Zhang *et al.*, 2001) SSR

(Yu et al. 2006, 2007; Zeid et al. 2011) markers were used to map QTLs for various traits onto

linkage groups. Two tef reference genomes (Cannarozzi *et al.*, 2014; VanBuren *et al.*, 2020) were

recently published, the latter identifying exceptional subgenome divergence. A most recent study

used the published reference genome to identify genomic regions associated with tef grain

nutritional traits (Ereful *et al.*, 2022). These studies are opening new avenues for further

exploration of tef's genomic potential. Recently published reviews have called for genomic studies

specifically in tef, showcasing their contribution to other cereals (Girija *et al.*, 2021; Numan *et al.*,

2021).

Plant lodging, defined as the permanent displacement of the culm from its vertical position, is the

most devastating problem in tef cultivation. Lodging can be divided into stem lodging, in which

the stem bends or breaks, and root lodging, in which the angle between the stem and the soil is

a changed due to root disanchoring(Berry *et al.*, 2004), the latter being the prime cause of tef

lodging(Ben-Zeev *et al.*, 2020*b*). Various studies estimated that tef lodging cause 30–50% yield

losses (Ketema, 1997; Van Delden *et al.*, 2010; Zeid *et al.*, 2010), reduce grain and feed quality

(Ketema, 1983*b;* Van Delden *et al.*, 2012; Reda, 2014) and lead to large economic penalties. The

damage caused by lodging is exacerbated in modern mechanized farming due to the difficulty in

harvesting lodging plants. Irrigation, required for tef cultivation under semiarid and arid conditions

(Ben-Zeev *et al.*, 2018) and mostly applied via sprinklers or pivots, further exacerbates lodging

due to increased canopy weight while the soil is lubricated and the plant's anchorage loosened.

Lodging often occurs on a plot or field level - meaning that a single plant or group of plants which

start lodging have a literal "domino" effect.

Lodging is a dynamic and detrimental phenomenon (Berry *et al.*, 2000, 2004; Van Delden *et al.*,

2010; Blösch *et al.*, 2020), which is most likely to be affected by various characteristics and genes

at different developmental stages. Like most other traits, it is affected by environment (E),

management (M), genotype (G) and their interaction (G x E x M). The effects of select

management practices on lodging have been reported (Paff and Asseng 2018 and references

therein). Semi-dwarfism, as a basis for reduced lodging, has been targeted via mutagenesis (Zhu

et al. 2012; Jost et al. 2015), as well as by multiparental (Abraha et al. 2017) and biparental

populations (Bekana and Assefa 2021). In the current paper, we report on the assembly and

genomic characterization of a tef diversity panel (TDP-300) consisting of 297 lines, as a new

resource for tef genomics and breeding. This unique resource was used to identify genomic

regions associated with plant lodging, possibly the most devastating and challenging tef

phenotype, thus demonstrating the utility of TDP-300 in paving the way to improve lodging

resistance in tef.

Materials and Methods

Diversity panel assembly and genotyping

A set of ca. 400 tef accessions, collected by the USDA in Ethiopia during the 1950s and 60s, and stored since the early 1990s in the Israeli Gene Bank (Agricultural Research Organization, Volcani Center) was propagated and stabilized via single-plant selection (Ben-Zeev *et al.*, 2018). DNA was isolated from leaf tissue at the seedling stage using a Nucleospin II kit (Mecherey Nagel, Germany). Sequencing was conducted at the Australian Genome Research Facility (AGRF), using a ddRAD-based library-preparation protocol. DNA was digested using restriction enzymes Pstl and Nlall, ligated to adaptors, amplified, and sequenced. Reads were tested for quality, aligned to the tef reference genome (VanBuren *et al.*, 2020), and variants were called using the Stacks V2.3d pipeline (Catchen *et al.*, 2011).

In the subsequent filtering steps, only biallelic SNPs were retained. Genotype quality was set to a minimum of 15. A minimum of 5 reads was set to call a genotype, where heterozygous genotypes with read depth (DP) < 30 were converted to missing data. Minimum and maximum mean depth over individuals were set to 5 and 20, respectively, based on the depth distribution (Fig. S1a). The lines were highly homozygous, owing to tef's self-pollinating nature and the single-plant-selection origin of the samples. Due to the high homozygosity and the abundance of rare alleles, minor allele frequency (MAF) was set to >1%. Sites with >30% missing data were removed. Individual lines with >1% heterozygous loci and/or >30% missing genotypes were excluded from the data set. Individuals were filtered based on identity by state (IBS) using four internal controls (lines doubled in sequencing). The final variant calling file (VCF) included 28,837 polymorphic sites and 297 lines, hereafter termed tef diversity panel 300 (TDP-300).

Exploration of population structure

Linkage disequilibrium (LD) pruning was conducted with Plink (Purcell *et al.*, 2007). Windows of 20 kbp were shifted by 5 variants at each step and variants in one window were removed if they exhibited $r^2 \geq 0.5$. Population stratification was explored using sparse non-negative matrix factorization (sNMF) with the LD-pruned SNP set algorithm as implemented in the R package 'LEA' (Frichot and François, 2015). Because the regularization parameter (α) can have a large influence on the outcome of the analysis' incomparably small data sets (Frichot *et al.*, 2014), the

optimum α was determined by testing a range of α values. The α value with the minimum mean

cross-entropy error was used for the final analysis. The final analysis was performed with the

following parameters: α = 260, K ranging from 1 through 30, 10 independent repetitions, 5,000

iterations, and a tolerance error of 10^{-6}. Cross-entropy was calculated with 10% masked

genotypes. Model-free approaches such as sNMF do not depend on the explicit assumption of

Hardy–Weinberg equilibrium and therefore, they are appropriate for both inbreeding and

outbreeding species (Frichot *et al.*, 2014).

A neighbor-joining tree was created based on Prevosti's distance (Prevosti *et al.*, 1975) and 100

bootstrap values to estimate the node support. Principal coordinate analysis (PCoA) was

performed using GENPOFAD genetic distances (Joly *et al.*, 2015; R Core Team, 2020), R core

functions and the 'poppr' package (Kamvar *et al.*, 2015).

The clustering tendency of the genotype matrix was estimated using the approach of Lawson and

Jurs (1990). We clustered the LD-pruned genotype matrix using the following clustering

algorithms: k-means, partition around medoids (PAM), clustering large applications (CLARA) and

hierarchical clustering. In each algorithm, we allowed 2 through 30 clusters. Cluster quality was

assessed using connectivity (Handl *et al.*, 2005), Dunn index (Dunn, 1974), and silhouette width

(Rousseeuw, 1987).

Experimental layout and phenotyping

The TDP-300 was grown in the field in 2 consecutive years under well-watered (WW) and water-

limited (WL) conditions. In 2020, the experiment was conducted at the J. Margulies Experimental

Farm of the Hebrew University of Jerusalem in Rehovot (31.906° N, 34.799° E), Israel, using an

augmented experimental design with 17 check lines replicated in three blocks. In 2021, the

experiment was conducted at the Kvutzat Shiller farm (31.879° N, 34.777° E), Israel, using a

factorial (lines x treatment) split-plot design (irrigation in main plots, lines in subplots) with three

replicates. Soil types at the experimental locations were sandy loam and clay, respectively. Both

experiments were sown on 22 April of the respective years. Experimental plots in both years

consisted of six 2-m long and 15-cm spaced rows, sown at a seed rate equivalent to 6 kg ha^{-1}

based on our previous study and the common practice in Israel (Ben-Zeev et al. 2020b).

Experiments were sprinkler-irrigated until 2 weeks after emergence (WAE), then drip-irrigated.

Due to technical constraints, a total of 216 and 252 TDP-300 lines were tested in the 2020 and

2021 experiment, respectively; of these, a complete data set across all four environments (2 years

x 2 treatments) was available for 202 lines. Mineral nutrition and weed control were applied

according to commercial farming practices, and no other pesticides were required.

Lodging was assessed visually by two independent surveyors, once a week from 8 to 10 WAE,

using a scoring method adapted from Caldicott and Nuttall (1979) and recently described in Ben-

Zeev et al. (2020b). Lodging was scored on a 10-level severity scale (0 being an upright, non-

lodging plant and 9 being a fully lodging plant), and lodging prevalence was assessed (percentage

of the entire plot area). For each plot, the severity was multiplied by the prevalence to calculate

the lodging index (LI).

Data analyses and heritability estimates

All data were analyzed using JMP software (Version 15, SAS Institute Inc., Cary, NC, 1989–2021)

and R 4.1.3 (R Core Team 2020). Line performance under each of the four environments: 2020

water-limited (20WL), 2020 well-watered (20WW), 2021 water-limited (21WL), and 2021 well-

watered (21WW), was calculated according to the experimental designs employed in each year.

Owing to the augmented experimental design in 2020, data of individual plots were adjusted for

block effects within each treatment, based on the 17 check lines replicated in each block. For the

2021 experiments, least square (LS) means were calculated using a full factorial model including

E x G interaction and blocks. In addition, LS means were calculated jointly for all four

environments using a model consisting of environment and genotype factors.

Broad sense heritability (h^2) estimates of LI in each week were calculated using two approaches.

For the replicated 2021 experiment, h^2 was calculated using variance components estimated

based on ANOVA:

$$H_b^2 = \sigma^2{}_g / (\sigma^2{}_g + \frac{\sigma^2{}_{gXe}}{e}) \qquad [1]$$

where $\sigma^2{}_g$ = [(MS $_{(G)}$ - MS $_{(G \times E)}$/e], $\sigma^2{}_{gXe}$ = [(MS $_{(G)}$ + MS $_{(G \times E)}$/e], e is the number of environments

and MS is the mean square.

Because ANOVA was not applicable for the augmented (non-replicated) 2020 experiment, and

based on tef's extreme self-pollination nature (Ketema, 1983*a*), h^2 estimates of LI in each week

were calculated using a linear regression coefficient (Cahaner and Hillel, 1980) between the two

treatments within the same year. Similarly, h^2 was calculated across years (2020 vs. 2021) for

each treatment as well as for means of each year.

Genome-wide association

Genotypic and phenotypic data were combined to perform a genome-wide association analysis

using a mixed linear model including the full VCF file and an IBS kinship matrix. Due to the low

clustering tendency of the population (see Results section), principal components were not

included in the final model.

The threshold for significance was set based on the Bonferroni correction (0.05/n = 1.7 x 10^{-6},

where n is the number of markers) and the Benjamini-Hochberg method implemented in the R

package 'RAINBOWR' (Bonferroni, 1936; Benjamini and Hochberg, 1995; Hamazaki and Iwata,

2020; R Core Team, 2020). Manhattan plots and Q-Q plots were also produced using R. The

association test was conducted using TASSEL (Bradbury *et al.*, 2007). The mixed linear model,

which included an IBS kinship matrix was:

$$y = Z\alpha + X\beta + W\gamma + e \hspace{3cm} [2]$$

where y is the phenotype value, α is the value of the fixed effect other than SNP, β is the vector

of the SNP effect, γ is the vector of the kinship effect, and e is the vector of residual effects.

Results

Sequencing quality

The final genotypic file included 28,837 SNPs, representing an average density of 46 SNPs/Mbp. The LD-pruned data set consisted of 9,268 SNPs. Nevertheless, use of the unpruned data was preferred to enable the detection of all significant SNPs. Density of SNPs varied along the genome, but no large regions devoid of SNPs were observed (Fig. 1). Missing data per site and per sample were at a low level, with ~10% at the third quartile (Fig. S1a). Most sites (27,550) were completely homozygous, and only 146 sites displayed >10% heterozygous genotypes (Fig. S1b). A quarter of all loci displayed MAF < 0.05 indicating an excess of rare alleles (Fig. S1c). The nucleotide diversity (π) exhibited a dichotomous distribution, skewed toward its possible extreme values of 0 and 0.5; the left skew is in line with the high abundance of rare alleles, whereas the right skew could be the result of (almost) fixed alleles between the two subpopulations (Fig. S1d).

Population structure

Estimated clustering tendency was 0.66, indicating nearly random data distribution. The cross-entropy criterion of sNMF was equivocal (Fig. S2). Possible candidates for the optimum number of K were 8, 14, and 20. The optimum number of clusters using classical clustering algorithms (k-means, PAM, CLARA, hierarchical) depended on the method employed for cluster-quality assessment. Connectivity, Dunn index, and silhouette width suggested 2, 2, and 30 clusters, respectively. Based on the two former clustering methods (out of the three), two subpopulations seemed to best describe the data. Due to the rather piecemeal data available regarding collection sites and environmental backgrounds for the TDP-300 lines, we could not attribute the clustering to any underlying parameter. Yet, in line with the low clustering tendency value, a clear-cut subpopulation structure was not observed, and the optimum number of clusters could not be unambiguously determined. Consequently, almost all samples exhibited at least some degree of admixture regardless of the number of clusters, and admixture increased with the growing number of subpopulations (Fig. 2). PCoA results corroborated the structure found using clustering algorithms (Fig. 3).

Lodging index distribution

Wide ranges of LI values were observed within each environment across various sampling times (Fig. 4). In both years, lodging onset was recorded at 8 WAE, as reflected by the distribution of LI values at this time point: low LI values for most lines while a few lines exhibited severe lodging, and these shifted to higher values in subsequent weeks. The natural trend of lodging escalation was evident from the increase in average values in any of the four environments along the presented 3-week data. As a result, LI did not distribute normally in any environment or at any sampling time and various transformations failed to restore normality. Therefore, LI data recorded at 8 WAE, which were heavily skewed toward lower values, were not used for heritability or GWAS analyses.

Despite being drip-irrigated, where water does not impose physical force on the crop's canopy, greater LI values were recorded in the WW vs. WL treatment, with an up to 5-fold difference at 8 WAE in the 2021 season (Fig. 5). Notably, the WL treatment resulted in later lodging onset than the WW treatment in both years.

Heritability of lodging

ANOVA-based h^2 estimates for LI (Eq. 1) using the 2021 data set were 0.27–0.30 (Table 1). Correlation-based h^2, calculated for the 2020 experiments, resulted in higher values (0.49–0.55), while between-year estimates were intermediate (0.26–0.49) (all correlation coefficients were significant, $P < 0.0001$). Owing to the self-pollinating nature of tef, all loci are expected to be homozygous and with no dominant effect. Therefore, correlation-based h^2 is a good proxy for the additive variance.

Marker-trait association for lodging index

A GWAS was conducted for eight LI data sets separately, each representing a specific year (2020, 2021), treatment (WW, WL) and time (9 and 10 WAE) (Figs. 6, 7). Manhattan plots for each data set are presented with Bonferroni-corrected threshold and Benjamini-Hochberg threshold (applicable only when the former threshold was met).

A two-step approach was employed to select the most important sites associated with LI. First,
we screened for significant sites (meeting the Bonferroni-corrected threshold, 1.7 x 10^{-6}) or a less
stringent suggestive threshold ($P < 1$ x 10^{-4}) determined based on lowest P-values indicated by
the Benjamini-Hochberg correction. This process resulted in a list of 29 sites distributed across
10 of the 20 tef chromosomes (Table S1). The number of lines carrying the minor alleles varied
between 3 and 22 across the 29 significant or suggestive sites.

To minimize the potential of false positives, we then selected sites that were significant
(Bonferroni threshold) in at least one data set and suggestive in at least one other data set. This
process resulted in 11 selected sites considered of prime interest. These 11 sites outlined four
groups of lines which shared the same minor allele(s) at common site(s) (Table 2). Group 1
consisted of four TDP-300 lines with the same minor alleles at six common sites (on
chromosomes 1A, 1B, 3A and 7A). Groups 2 and 3 consisted of seven and eight lines,
respectively, having the same minor allele at a single site each (chromosome 2B and 1A,
respectively). Importantly, there was some overlap between lines belonging to groups 1–3. Group
4 consisted of 21 lines with the same minor allele at three common sites (all three on chromosome
7A). The lines in group 4 were not present in groups 1 and 2.

A t-test was used to test the effects of selected sites on LI recorded at 8-10 WAE, averaged (LS
means) across all four environments (Table 2). The phenotypic differences between the two allelic
groups were significant in most cases ($P < 0.05$), thus validating the effect of the selected sites.
The difference between lines carrying the minor vs. major alleles ranged between 100 and 280 LI
units, nearly one-third of the entire LI range.

Candidate genes in and near the selected sites

The tef genome browser (genomevolution.org/coge/GenomeInfo.pl?gid=50954) was used to
search for candidate genes located at or closest to (up to 200 kbp up or downstream) the selected
genomic loci associated with LI. Out of the 11 selected sites, 4 were located in genes (all 4 with
a putative function) and 7 were located near genes (1 with a putative function) (Table S2).

Discussion

TDP-300 – a new tool for tef genomics and breeding

Being a minor crop with narrow geographic distribution, availability of genomic infrastructure, genetic markers, and a mapping database is limited for tef (Tadele, 2019). Hence advanced genome-based, marker-assisted breeding of tef has remained mostly out of reach thus far (Bekana and Assefa, 2021; Numan *et al.*, 2021). Genomic sites related to lodging have scarcely been reported in tef, although a few studies pre-dating the published tef genomes (Cannarozzi *et al.*, 2014; VanBuren *et al.*, 2020) reported on genetic markers associated with lodging (Yu *et al.*, 2007; Zeid *et al.*, 2012). The difficulty involved in crossing tef (Berhe, 1975; Zeid *et al.*, 2010; Habteab *et al.*, 2015) and the deficiency of reference genomes (Adnew *et al.*, 2005; Tadele, 2019; Girija *et al.*, 2021) may have hindered the development of biparental populations and diversity panels, respectively, both being essential for genetic mapping. Hence, the TDP-300 reported in the current study may become a useful tool for genomic dissection of various traits of interest in tef and subsequently, their advancement via marker-assisted breeding and their investigation in other germplasms.

Structure analysis

To test the clustering tendency, Lawson and Jurs (1990) compared the observed structure with the structure of "white noise". In the absence of any clusters, the expected value was 0.5, whereas a value of 0.75 indicated a clustering tendency at the 90% confidence level (Han *et al.*, 2011). In the TDP-300, the estimated tendency was 0.66, suggesting very low clustering. In line with this, the clustering approaches produced ambiguous results and did not indicate a single optimum number of clusters that would be favorable for GWAS. We opted to use two clusters (subpopulations) as the optimum number of subpopulations to apply the coloring schemes in the figures (Figs. 2, 3), because two independent approaches pointed to this cluster number. Yet, the varying results indicated that the collection cannot be described by any clear-cut subpopulation structure. Hence, for GWAS, we assumed no population structure in the TDP-300.

Distribution and heritability of lodging

Lodging is a complicated phenomenon resulting from plant (shoot and root) traits (Berry *et al.*,
2004), environmental (soil properties and moisture) factors (Pinthus, 1974; Berry *et al.*, 2000),
and management practices (irrigation, sowing depth and density) (Ben-Zeev *et al.*, 2020*a,b*). The
nature of tef lodging as a dynamic and largely irremediable phenomenon is clearly depicted in the
LI distributions at 8–10 WAE across the four studied environments (Figs. 4, 5). Due to the rather
uniform onset of lodging and the rarity of lodging-resistant lines, LI distribution was heavily skewed
toward lower values at 8 WAE, wider at 9 WAE, and moved toward the high end of LI values at
10 WAE.

Heritability estimates ranged between 0.26 and 0.55 (Table 1) at different times (WAE), under
different environments, and with different calculation methods. In line with our results, previous
studies on tef lodging have reported h^2 values of 0.16 to 0.55 (Chanyalew, 2010; Ayalneh *et al.*,
2011; Bekana and Assefa, 2021). In the current study, lower values (0.27–0.30) were obtained
from the ANOVA-based calculation, and the highest values from a between-environment
correlation in the 2020 season. Since correlation-based h^2 tends to be overestimated, it is the
authors' opinion that h^2 values of 0.30–0.45 for tef lodging are probably realistic. These estimates,
considered low to intermediate, reflect the small genetic variation and large effects of environment
and management on tef lodging.

SNP markers associated with lodging

Owing to the nature of lodging in tef and the rarity (or near absence) of non-lodging lines, the LI
did not distribute normally, particularly at 8 WAE. Transformations failed to restore normality, and
therefore LI8 was excluded and the original data sets of LI9 and LI10 were used for GWAS, albeit
with a careful consideration of the results. Our two-step site screening resulted in 11 selected
sites which outlined four groups of lines sharing the same minor allele(s) at common site(s) (Table
2). Groups 1 and 4 shared 6 and 3 common sites, respectively (Table 2). The current data and
panel do not allow distinguishing between the effect of the various sites associated with each
group, and their apparent phenotypic effect may result from either a single site (with all other sites
being neutral), closely linked sites or a few independent sites. Previous mapping studies of tef
lodging (Yu *et al.*, 2007; Zeid *et al.*, 2011) pre-dated the published reference genomes, and
therefore we could not compare the previously published sites with those identified in the current
study.

Due to the lodging-assessment method, LI ranged between 0 and 900. A somewhat wider
distribution resulted from the block adjustment of the 2020 augmented design experiment (Fig. 4)
or the LS mean calculated for the 2021 replicated experiment (Fig. 5). The effects of SNP sites
on LS mean LIs of the four combined environments confirmed the effects of the selected sites
and demonstrated the magnitude of the genetic effect on LI (Table 2). Various sites seem to affect
lodging at different times during plant development, suggesting their involvement in different plant
traits and/or developmental processes, such as crown root number and diameter, culm diameter
and number of tillers, as suggested in our previous publication (Ben-Zeev et al. 2020b). A number
of published studies have targeted reduced plant heightt as a means of reducing lodging in tef
(Zhu et al. 2012; Jost et al. 2015; Jifar et al. 2017; Bekana and Assefa 2021). Mapping plant
height and panicle length assessed under the same four environments reported here identified a
number of sites associated with these traits (data not shown). However, no overlap was found
between sites associated with lodging and those associated with plant height or panicle length.
While plant height determines the bending moment applied by the canopy, root-borne lodging
depends also on the diameter of the root-soil cone, the bending strength of the crown and the
shear strength of the soil (Shah *et al.*, 2019). Indeed, 3 of the 5 candidate genes with a putative
function were reported to be involved in root development(Beisson *et al.*, 2007; Wu *et al.*, 2016;
Lee *et al.*, 2019; Su *et al.*, 2019).

In all 29 significant or suggestive sites (Table S1), the minor allele was associated with reduced
lodging in line with the rarity of lodging resistance in tef. Of these, the selected 11 sites, which
exhibited an effect on lodging across two data sets and were confirmed in the combined four-
environment data, can be readily utilized for selecting lines with improved lodging resistance.

Concluding remarks

The sites identified in this study were found to reduce the LI by up to 280 units (Table 2), which
may play a vital role in reducing lodging damage in traditional as well as modern tef cultivation,
thus improving farmers' livelihood and food security. Combining several alleles associated with
lodging resistance may result in a greater reduction of LI. The identification of lodging-related sites
in the current study, as well as in future studies, can help understand the mechanisms underlying
tef lodging and provide a basis to outline the genetic markers for a non-lodging tef ideotype.

This study presents the first tef diversity panel (TDP-300) subjected to modern genotyping technique (GBS-based SNP markers). Phenotyping of TDP-300 under four environments in the field revealed wide diversity in a number of phenotypic traits, most of which will be presented in our follow-up publication. Lodging was used in the current study to examine the utility of the panel for genomic mapping of complex traits. The success in identifying genomic regions associated with LI demonstrates the potential of TDP-300 for further tef research.

Seeds of TDP-300, as well as our VCF, will soon be available for distribution to allow further research of this important and intriguing orphan crop. We invite the scientific community to utilize this resource to promote tef research, which in turn will support broader tef cultivation and production (Tadele, 2019; Kebede, 2021; Numan *et al.*, 2021), crop diversification and food security.

Acknowledgements

This study was supported by The Israel Ministry of Agriculture and Rural Development, Chief Scientist Foundation (grant no. 12-02-0008) and The Israel Innovation Authority, Challenge Program (grant no. 73546). The authors wish to acknowledge the use of the services and facilities of AGRF. We thank the Netafim irrigation company for providing the irrigation system for the 2020 season. We are grateful to Noa Kirby, Philip Wagali, Ido Zeltzer, Neta Levinson, Ori Harash, Ariel Joseph, May Shoshan, Eran Dagan, Emma Braslavsky, Eric and Rotem Ben-Zeev who enabled performance of this study with their hard work and insights. S.B.-Z. is indebted to the Robert H. Smith Foundation for a doctoral fellowship award. M. D. is The Heinrich Bonnenberg Scholarship Awardee. Y.S. is the incumbent of the Haim Gvati Chair in Agriculture.

Author Contributions

Shiran Ben-Zeev and Yehoshua Saranga assembled the diversity panel, conceptualized and designed the study. Shiran Ben-Zeev Muluken Demeile and Vered Barak collected the data, propagated the plants and extracted the DNA. Timo Hellwig conducted the population structure and genomic data analysis and supported the association study. Sasha Vorobyova and Sariel Hübner analyzed the raw genomic data and provided insight into the population structure and the association study. Shiran Ben-Zeev and Yehoshua Saranga analyzed the phenotypic and

genomic data and wrote the first draft of the manuscript. All authors read and approved the final

manuscript.

Conflict of interest

431   The author declare no conflict of interest.

Data availability statement

434   Genomic data and seeds are readily available for the scientific community. VCF files will be made

435   available on our web site, sequence data as well as seeds of the TDP300 will be provided upon

436   request.

References

Abraha MT, Hussein S, Laing M, Assefa K. 2017. Early generation genetic variation and heritability of yield and related traits among tef populations. Journal of Crop Science and Biotechnology **20**, 379–386.

Abrouk M, Ahmed HI, Cubry P, *et al.* 2020. Fonio millet genome unlocks African orphan crop diversity for agriculture in a changing climate. Nature Communications **11**.

Adnew T, Ketema S, Tefera H, Sridhara H. 2005. Genetic diversity in tef [Eragrostis tef (Zucc.) Trotter] germplasm. Genetic Resources and Crop Evolution **52**, 891–902.

Allaby R. 2021. First come, first served for ancient crops. Nature Plants **7**, 542–543.

Assefa K, Chanyalew S, Tadele Z. 2013. *Achievements and prospects of tef improvement*.

Ayalneh T, Habtamu Z, Amsalu A. 2011. Genetic Variability, Heritability and Genetic Advance in Tef (Eragrostis tef (Zucc.) Trotter) Lines at Sinana and Adaba. International Journal of Plant Breeding and Genetics **6**, 40–46.

Bai G, Tefera H, Ayele M, Nguyen HT. 1999. A genetic linkage map of tef [Eragrostis tef (Zucc.) Trotter] based on amplified fragment length polymorphism. Theoretical and Applied Genetics **99**, 599–604.

Beisson F, Li Y, Bonaventure G, Pollard M, Ohlrogge JB. 2007. The Acyltransferase GPAT5 Is Required for the Synthesis of Suberin in Seed Coat and Root of Arabidopsis. **19**, 351–368.

Bekana G, Assefa K. 2021. Genetic Variation , Heritability and Genetic Advance Among Semi-Dwarf Tef [Eragrostis tef (Zucc .) Trotter] Recombinant Inbred Lines with Emphasis to Lodging. **9**, 92–104.

Bellucci A, Torp AM, Bruun S, Magid J, Andersen SB, Rasmussen SK. 2015. Association Mapping in Scandinavian Winter Wheat for Yield, Plant Height, and Traits Important for Second-Generation Bioethanol Production. Frontiers in Plant Science **6**, 1–12.

Ben-Zeev S, Bimro J, Barak V, Saranga Y. 2018. Phenotypic diversity and heritability in

Eragrostis tef under irrigated Mediterranean conditions. Israel Journal of Plant Sciences **65**, 222–231.

Ben-Zeev S, Kerzner S, Rabinovitz O, Saranga Y. 2020*a*. Optimizing Sowing Depth of Tef for Irrigated Mediterranean Conditions: From Laboratory to Field Studies. Agronomy **10**, 16.

Ben-Zeev S, Rabinovitz O, Orlov-Levin V, Chen A, Graff N, Goldwasser Y, Saranga Y. 2020*b*. Less Is More: Lower Sowing Rate of Irrigated Tef (Eragrostis tef) Alters Plant Morphology and Reduces Lodging. Agronomy **10**, 570.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal statistical society: series B (Methodological) **57**, 289–300.

Berhe T. 1975. A breakthrough in tef-breeding technique. Information Bulletin on the Near East Cereal Improvement and Production Project (FAO).

Berry PM, Griffin JM, Sylvester-Bradley R, Scott RK, Spink JH, Baker CJ, Clare RW. 2000. Controlling plant form through husbandry to minimise lodging in wheat. Field Crops Research **67**, 59–81.

Berry PM, Sterling M, Spink JH, Baker CJ, Sylvester-Bradley R, Mooney SJ, Tams AR, Ennos AR. 2004. Understanding and Reducing Lodging in Cereals. Advances in Agronomy **84**, 217–271.

Blösch R, Plaza-Wüthrich S, Barbier de Reuille P, Weichert A, Routier-Kierzkowska A-L, Cannarozzi G, Robinson S, Tadele Z. 2020. Panicle Angle is an Important Factor in Tef Lodging Tolerance. Frontiers in Plant Science **11**, 1–12.

Bonferroni C. 1936. Teoria statistica delle classi e calcolo delle probabilita. Pubblicazioni del R Istituto Superiore di Scienze Economiche e Commericiali di Firenze **8**, 3–62.

Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES. 2007. TASSEL: software for association mapping of complex traits in diverse samples. Bioinformatics **23**, 2633–2635.

Brook AJ, Woodcock BA, Sinka M, Vanbergen AJ. 2008. Experimental verification of suction sampler capture efficiency in grasslands of differing vegetation height and structure. Journal of

Applied Ecology **45**, 1357–1363.

Brown R. 2017. *Using Teff as a Summer Cover Crop Using Teff as a Summer Cover Crop*.

Cahaner A, Hillel J. 1980. Estimating heritability and genetic correlation between traits from generations F2 and F 3 of self-fertilizing species: a comparison of three methods. TAG. Theoretical and applied genetics. Theoretische und angewandte Genetik **58**, 33–8.

Caldicott JJB, Nuttall AM. 1979. A method for the assessment of lodging in cereal crops. Journal of the National Institute of Agricultural Botany **15**, 88–91.

Cannarozzi G, Plaza-Wüthrich S, Esfeld K, *et al.* **2014. Genome and transcriptome sequencing identifies breeding targets in the orphan crop tef (Eragrostis tef). BMC Genomics **15, 581.

Catchen JM, Amores A, Hohenlohe P, Cresko W, Postlethwait JH. 2011. Stacks: building and genotyping loci de novo from short-read sequences. G3: Genes| genomes| genetics **1**, 171–182.

Chanyalew S. 2010. Genetic analyses of agronomic traits of tef (Eragrostis tef) genotypes. Research Journal of Agriculture and Biological Sciences **6**, 912–916.

D'Andrea AC. 2008. T'ef (Eragrostis tef) in ancient agricultural systems of highland Ethiopia. Economic Botany **62**, 547–566.

Dawson IK, Powell W, Hendre P, Bančič J, Hickey JM, Kindt R, Hoad S, Hale I, Jamnadass R. 2019. The role of genetics in mainstreaming the production of new and orphan crops to diversify food systems and support human nutrition. New Phytologist **224**, 37–54.

Van Delden SH, Vos J, Ennos AR, Stomph TJ. 2010. Analysing lodging of the panicle bearing cereal teff (Eragrostis tef). New Phytologist **186**, 696–707.

Van Delden SH, Vos J, Stomph TJ, Brouwer G, Struik PC, Alaunyte I, Stojceska V, Plunkett A, Ainsworth P, Derbyshire E. 2012. Improving the quality of nutrient-rich Teff (Eragrostis tef) breads by combination of enzymes in straight dough and sourdough breadmaking. European Journal of Agronomy **37**, 105–114.

Ditsch DC. 2007. *Options for Summer Grazing*.

Dunn JC. 1974. Well-separated clusters and optimal fuzzy partitions. Journal of cybernetics **4**, 95–104.

Ereful NC, Jones H, Fradgley N, Boyd L, Cherie HA, Milner MJ. 2022. Whole-Genome Sequencing and Analysis of Nutritional Traits in a Core Set of Ethiopian Teff (Eragrostis tef) Varieties.

Frichot E, François O. 2015. LEA: An R package for landscape and ecological association studies. Methods in Ecology and Evolution **6**, 925–929.

Frichot E, Mathieu F, Trouillon T, Bouchard G, François O. 2014. Fast and efficient estimation of individual ancestry coefficients. Genetics **196**, 973–983.

Girija A, Jifar H, Jones C, Yadav R, Doonan J, Mur LAJ. 2021. Tef: a tiny grain with enormous potential. Trends in Plant Science **27**, 220–223.

Habteab G, Richard B, Shimelis H, Mark L. 2015. The efficacy of four gametocides for induction of pollen sterility in Eragrostis tef (Zucc.) Trotter. African Journal of Biotechnology **14**, 774–780.

Hamazaki K, Iwata H. 2020. RAINBOW: Haplotype-based genome-wide association study using a novel SNP-set method. PLoS computational biology **16**, e1007663.

Han J, Kamber M, Pei J. 2011. Data Mining: Concepts and. Techniques (3rd ed), Morgan Kauffman.

Handl J, Knowles J, Kell DB. 2005. Computational cluster validation in post-genomic data analysis. Bioinformatics **21**, 3201–3212.

Huang J, Pray C, Rozelle S. 2002. Enhancing the crops to feed the poor. Nature **418**, 678–684.

Ikegawa S. 2012. A short history of the genome-wide association study: where we were and where we are going. Genomics & informatics **10**, 220.

Jamnadass R, Mumm RH, Hale I, *et al.* 2020. Enhancing African orphan crops with genomics. Nature Genetics **52**, 356–360.

Jifar H, Tesfaye K, Assefa K, Chanyalew S, Tadele Z. 2017. Semi-dwarf tef lines for high seed yield and lodging tolerance in Central Ethiopia. African Crop Science Journal **25**, 419–439.

Joly S, Bryant D, Lockhart PJ. 2015. Flexible methods for estimating genetic distances from single nucleotide polymorphisms. Methods in Ecology and Evolution **6**, 938–948.

Jost M, Esfeld K, Burian A, Cannarozzi G, Chanyalew S, Kuhlemeier C, Assefa K, Tadele Z. 2015. Semi-dwarfism and lodging tolerance in tef (Eragrostis tef) is linked to a mutation in the alpha-Tubulin 1 gene. Journal of Experimental Botany **66**, 933–944.

Kamenya SN, Mikwa EO, Song B, Odeny DA. 2021. Genetics and breeding for climate change in Orphan crops. Theoretical and Applied Genetics, 1–29.

Kamvar ZN, Brooks JC, Grunwald NJ. 2015. Novel R tools for analysis of genome-wide population genetic data with emphasis on clonality. Front Genet. 2015; 6: 208.

Kebede W. 2021. Quality traits ; measurement and possible breeding method to improve the quality traits of tef [Eragrostis tef (Zucc .) Trotter]. International Journal of Novel Research in Life Sciences **8**, 29–40.

Ketema S. 1983*a*. Studies of Lodging, Floral Biology and Breeding Techniques in Tef (Eragrostis Tef (Zucc.) Trotter).

Ketema S. 1983*b*. Studies of Lodging Floral Biology and Bredding Techniques in Tef (Eragrostis tef).

Ketema S. 1993. *Tef (Eragrostis Tef): Breeding, Genetic Resources, Agronomy, Utilization and Role in Ethiopian Agriculture.* Adis Abab, Ethopia: Institute of Agricultural Research.

Ketema S. 1997. *Eragrostis tef (Zucc.) Trotter: Promoting the conservation and use of underutilized and neglected crops.* Rome: Institute of Plant Genetics and Crop Plant Research, Gatersleben/International Plant Genetic Resources Institute.

Khoury CK, Bjorkman AD, Dempewolf H, Ramirez-Villegas J, Guarino L, Jarvis A, Rieseberg LH, Struik PC. 2014. Increasing homogeneity in global food supplies and the implications for food security. Proceedings of the National Academy of Sciences of the United States of America **111**, 4001–4006.

Lawson RG, Jurs PC. 1990. New index for clustering tendency and its application to chemical problems. Journal of chemical information and computer sciences **30**, 36–41.

Lee K, Park SJ, Han JH, Jeon Y, Pai H, Kang H. 2019. A chloroplast-targeted pentatricopeptide repeat protein PPR287 is crucial for chloroplast function and Arabidopsis development. **244**, 1–10.

Mabhaudhi T, Chimonyo VGP, Hlahla S, Massawe F, Mayes S, Nhamo L, Modi AT. 2019. Prospects of orphan crops in climate change. Planta **250**, 695–708.

Matthew Davidson J. 2018. Evaluating teff grass as a summer forage.

Milla R, Osborne CP. 2021. Crop origins explain variation in global agricultural relevance. Nature Plants **7**, 598–607.

Miller D. 2008. *Teff As an Alternative Summer Forage*.

Negash W. 2017. Effect of Row Spacing on Yield and Yield Components of Teff [Eragrostis tef (Zucc .) Trotter] Varieties in Gonji Kolela District , North Western Ethiopia. Journal of Biology, Agriculture and Healthcare **7**, 35–43.

Norberg S, Roseberg R, Charlton B, Shock C. 2009. Teff: A new warm-season annual grass for Oregon. OSU Extension Catalog **1**, 2008–2009.

Numan M, Khan AL, Asaf S, Salehin M, Beyene G, Tadele Z, Ligaba-Osena A. 2021. From traditional breeding to genome editing for boosting productivity of the ancient grain tef [eragrostis tef (Zucc.) trotter]. Plants **10**.

Paff K, Asseng S. 2018. A review of tef physiology for developing a tef crop model. European Journal of Agronomy **94**, 54–66.

Pinthus MJ. 1974. Lodging in Wheat, Barley, and Oats: The Phenomenon, its Causes, and Preventive Measures. Advances in Agronomy **25**, 209–263.

Prevosti A, Ocana J, Alonso G. 1975. Distances between populations of Drosophila subobscura, based on chromosome arrangement frequencies. Theoretical and Applied Genetics **45**, 231–241.

Puranik S, Sahu PP, Beynon S, Srivastava RK, Sehgal D, Ojulong H, Yadav R. 2020. Genome-wide association mapping and comparative genomics identifies genomic regions governing grain nutritional traits in finger millet (Eleusine coracana L. Gaertn.). Plants People Planet **2**, 649–662.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, De Bakker PIW, Daly MJ. 2007. PLINK: a tool set for whole-genome association and population-based linkage analyses. The American journal of human genetics **81**, 559–575.

R Core Team RC. 2020. R: A Language and Environment for Statistical Computing.

Reda A. 2014. Achieving food security in Ethiopia by promoting productivity of future world food tef: a review. Advances in Plants & Agriculture Research **2**.

Renard D, Tilman D. 2019. National food production stabilized by crop diversity. Nature **571**, 257–260.

Rosenzweig C, Elliott J, Deryng D, *et al.* 2014. Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison. Proceedings of the National Academy of Sciences of the United States of America **111**, 3268–3273.

Rousseeuw PJ. 1987. Silhouettes: a graphical aid to the interpretation and validation of cluster analysis. Journal of computational and applied mathematics **20**, 53–65.

Shah L, Yahya M, Mehar S, *et al.* 2019. Improving Lodging Resistance : Using Wheat and Rice as Classical Examples.

Shumoy H, Pattyn S, Raes K. 2018. Tef protein: Solubility characterization, in-vitro digestibility and its suitability as a gluten free ingredient. LWT - Food Science and Technology **89**, 697–703.

Smith RL, Hoveland CS, Hanna WW. 1989. Water Stress and Temperature in Relation to Seed Germination of Pearl Millet and Sorghum. Agronomy Journal **81**, 303–305.

Su HG, Li B, Song XY, Ma J, Chen J, Zhou Y Bin, Chen M, Min DH, Xu ZS, Ma YZ. 2019. Genome-wide analysis of the DYW subgroup PPR gene family and identification of gmPPR4 responses to drought stress. International Journal of Molecular Sciences **20**, 1–21.

Tadele Z. 2019. Orphan crops: their importance and the urgency of improvement. Planta **250**,

677–694.

Tilman D, Balzer C, Hill J, Befort BL. 2011. Global food demand and the sustainable intensi fi cation of agriculture. PNAS | **108**.

VanBuren R, Man Wai C, Wang X, *et al.* 2020. Exceptional subgenome stability and functional divergence in the allotetraploid Ethiopian cereal teff. Nature Communications **11**, 884.

Varshney RK, Ribaut JM, Buckler ES, Tuberosa R, Rafalski JA, Langridge P. 2012. Can genomics boost productivity of orphan crops? Nature Biotechnology **30**, 1172–1176.

Wu Y, Xun Q, Guo Y, Zhang J, Cheng K, Shi T, He K, Hou S, Gou X, Li J. 2016. Genome-Wide Expression Pattern Analyses of the Arabidopsis Leucine-Rich Repeat Receptor-Like Kinases. Molecular Plant **9**, 289–300.

Wuepper D, Borrelli P, Finger R. 2020. Countries and the global rate of soil erosion. Nature Sustainability **3**, 51–55.

Yu J-K, Graznak E, Breseghello F, Tefera H, Sorrells ME. 2007. QTL mapping of agronomic traits in tef [Eragrostis tef (Zucc) Trotter]. BMC plant biology **7**, 2–13.

Yu JK, Sun Q, Rota ML, Edwards H, Tefera H, Sorrells ME. 2006. Expressed sequence tag analysis in tef (Eragrostis tef (Zucc) Trotter). Genome **49**, 365–372.

Zeid M, Assefa K, Haddis A, Chanyalew S, Sorrells ME. 2012. Genetic diversity in tef (Eragrostis tef) germplasm using SSR markers. Field Crops Research **127**, 64–70.

Zeid M, Belay G, Mulkey S, Poland J, Sorrells ME. 2011. QTL mapping for yield and lodging resistance in an enhanced SSR-based map for tef. Theoretical and Applied Genetics **122**, 77–93.

Zeid M, Yu J-K, Assefa K, Tefera H, Sorrells ME, Belay G. 2010. Breeding tef [Eragrostis tef (Zucc.) trotter]: conventional and molecular approaches. Plant Breeding **130**, 1–9.

Zhang D, Ayele M, Tefera H, Nguyen HT. 2001. RFLP linkage map of the Ethiopian cereal tef [Eragrostis tef (Zucc) Trotter]. Theoretical and Applied Genetics **102**, 957–964.

Zhang Y, Niu H, Yu Q. 2021. Impacts of climate change and increasing carbon dioxide levels

on yield changes of major crops in suitable planting areas in China by the 2050s. Ecological Indicators **125**, 107588.

Zhao J, Bayer PE, Ruperao P, Saxena RK, Khan AW, Golicz AA, Nguyen HT, Batley J, Edwards D, Varshney RK. 2020. Trait associations in the pangenome of pigeon pea (Cajanus cajan). Plant Biotechnology Journal **18**, 1946–1954.

Zhu Q, Smith SM, Ayele M, Yang L, Jogi A, Chaluvadi SR, Bennetzen JL. 2012. High-throughput discovery of mutations in tef semi-dwarfing genes by next-generation sequencing analysis. Genetics **192**, 819–829.

Tables and figures

Table 1. ANOVA-based and correlation-based broad sense heritability (h^2) estimates of lodging indices at 9 and 10 weeks after emergence (LI9 and LI10, respectively) and water-limited (WL), well-watered (WW) or averages across environments (LS means).

Data sets used	h^2 estimates	
	LI9	LI10
ANOVA-based estimate		
2021 (two environments)	0.30	0.27
Correlation-based estimate		
2020WL vs. 2020WW	0.49	0.55
2020WL vs. 2021WL	0.44	0.48
2020WW vs. 2021WW	0.26	0.36
LS means 2020 vs. LS means 2021	0.44	0.49

Table 2. Selected genomic sites associated with lodging indices (LI) in two or more data sets (see Table S1) divided into four groups carrying a minor allele at the common sites of interest. Presented are SNP ratio in the TDP-300 lines tested across four environments (out of 202), mean LI of the various allelic groups and *P*-values of t-test for LS (significant *P-values* in bold) means for all four environments combined. Lines carrying minor allele are listed; those tested across all environments are underlined.

Groups of lines	Genomic sites	SNP ratio, minor/major in TDP-300 lines tested across 4 environ-ments	Mean phenotypic effect of minor vs. major alleles and t-test probability across 4 combined environments			Lines carrying minor allele at the common sites
			LI8	LI9	LI10	
1	1A:20268178 1A:21091933 1B:22065152 1B:22075858 3A:8334174 7A:17140288	4/293 (2/200)	1; 139 *P* = 0.09	101; 181 *P* = 0.31	386; 646 *P* = 0.014	RTC146, RTC172, RTC177, RTC195
2	2B:17571095	7/290 (4/198)	1; 141 *P* = 0.013	155; 180 *P* = 0.65	445; 647 *P* = 0.007	RTC146, RTC161, RTC167, RTC172, RTC195, RTC236, RTC274
3	1A:31534966	8/281 (5/197)	22; 141 *P* = 0.021	138; 181 *P* = 0.39	486; 647 *P* = 0.017	RTC146, RTC167, RTC172, RTC177, RTC180b, RTC195, RTC239, RTC276
4	7A:25753459 7A:25872958 7A:25888367	21/276 (12/190)	44; 144 *P* = 0.003	62; 187 *P* = 0.0001	432; 656 *P* < 0.0001	RTC4, RTC79b, RTC111, RTC155a, RTC157, RTC192, RTC211, RTC233, RTC254, RTC262, RTC264, RTC265, RTC266, TRC267, RTC279a, RTC286, RTC302a, RTC302b, RTC307, RTC320, RTC325b

Figure 1. Distribution of SNP density over chromosomes. Outer circle (green): filtered variant set. Inner circle (red): LD-pruned variant set.

Figure 2. Neighbor-joining tree and sNMF bar plots with the number of subpopulations, K, ranging from 2 to 8. Only bootstrap values >70% are shown. Sample labels in the neighbor-joining tree are colored according to sNMF subpopulation assignment, with K = 2 if they had at least 60% subpopulation fraction of one subpopulation.

Figure 3. Genomic Principal coordinate analysis (PcoA) showing the relationship between the two subpopulations. Principal component (PC) 1 and 2 cumulatively explained ~15.5% of the overall genetic variation

Figure 4. Distributions of lodging index values at 8, 9 and 10 weeks after emergence (LI8, 9, 10, respectively) under well-watered (WW) and water-limited (WL) treatments in the 2020 season. Mean values are indicated for each distribution.

Figure 5. Distributions of lodging index values at 8, 9 and 10 weeks after emergence (LI8, 9, 10, respectively) under well-watered (WW) and water-limited (WL) treatments in the 2021 season. Mean values are indicated for each distribution.

Figure 6. Manhattan and Q-Q plots of the association of 28,837 SNP with lodging index at 9 and 10 weeks after emergence (LI9 and LI10, respectively) under well-watered (WW) and water-limited (WL) treatments in the 2020 season. Red and green dotted lines (respectively) represent the significance threshold based on the Bonferroni correction threshold and Benjamini-Hochberg correction threshold (where applicable).

Figure 7. Manhattan and Q-Q plots of the association of 28,837 SNP with lodging index at 9 and 10 weeks after emergence (LI9 and LI10, respectively) under well-watered (WW) and water-limited (WL) treatments in the 2021 season. Red and green dotted lines (respectively) represent the significance threshold based on the Bonferroni correction threshold and Benjamini-Hochberg correction threshold (where applicable).

Supplementary material

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
missing data per site	0	0.03704	0.06061	0.07962	0.10774	0.31313
missing data per sample	0.01179	0.04064	0.06343	0.07962	0.10421	0.27864
observed heterozygousity	0	0	0	0.00164	0	0.26407
minor allele frequency	0.01010	0.04437	0.13402	0.17569	0.28723	0.50000
nucleotide diversity	0.02003	0.08495	0.23252	0.24806	0.41019	0.50113

Figure S1. Quality assessment of the final SNP set described by missing data per site and sample, observed heterozygosity, minor allele frequency and nucleotide diversity (π). Min., minimum; 1st Qu., first quartile; 3rd Qu., third quartile; Max., maximum.

Figure S2. Cross-entropy criterion of sNMF averaged over replications for K ranging from 0 to 30.

Table S1. Significant sites (meeting the Bonferroni threshold, 1.7 x 10^{-6}), and suggestive threshold (p < 1 x 10^{-4}).

Sites	SNP (minor/major)	SNP ratio (minor/major)	20WW_ LI9	20WW_ LI10	20WL_ LI9	20WL_ LI10	21WW_ LI9	21WW_ LI10	21WL_ LI9	21WL_ LI10
1A_20268178	A/G	4/293	N.S.	N.S.	N.S.	N.S.	0.00000156	1.56E-08	N.S.	N.S.
1A_21091933	C/T	4/293	N.S.	N.S.	N.S.	N.S.	0.00000156	1.56E-08	N.S.	N.S.
1A_31534966	C/A	8/289	N.S.	N.S.	N.S.	N.S.	0.000092	0.000000462	N.S.	N.S.
1A_35589206	C/G	6/291	N.S.	N.S.	N.S.	N.S.	N.S.	0.00000735	N.S.	N.S.
1A_35747247	A/G	6/291	N.S.	N.S.	N.S.	N.S.	N.S.	0.00000735	N.S.	N.S.
1B_22065152	A/G	4/293	N.S.	N.S.	N.S.	N.S.	0.00000156	1.56E-08	N.S.	N.S.
1B_22075858	A/G	4/293	N.S.	N.S.	N.S.	N.S.	0.00000156	1.56E-08	N.S.	N.S.
1B_4782451	T/C	3/294	N.S.	N.S.	N.S.	N.S.	N.S.	0.00000106	N.S.	N.S.
2B_17571095	G/A	7/290	N.S.	N.S.	N.S.	N.S.	0.000000825	0.0000248	N.S.	N.S.
2B_500639	A/G	6/291	N.S.	N.S.	N.S.	N.S.	N.S.	0.00000415	N.S.	N.S.
3A_6416217	G/A	5/292	N.S.	N.S.	N.S.	N.S.	N.S.	3.41E-08	N.S.	N.S.
3A_6430248	T/G	8/289	N.S.	N.S.	N.S.	N.S.	N.S.	0.00000153	N.S.	N.S.
3A_6459194	G/A	5/292	N.S.	N.S.	N.S.	N.S.	N.S.	0.000000468	N.S.	N.S.
3A_8334174	A/G	5/292	N.S.	N.S.	N.S.	N.S.	0.00000156	1.56E-08	N.S.	N.S.
3B_19818793	A/G	5/292	N.S.	N.S.	N.S.	N.S.	N.S.	3.41E-08	N.S.	N.S.
4A_20600242	T/C	5/292	N.S.	N.S.	N.S.	N.S.	N.S.	3.41E-08	N.S.	N.S.
4B_28815304	C/T	8/289	N.S.	N.S.	N.S.	N.S.	0.00000303	N.S.	N.S.	N.S.
6B_18856981	T/C	8/289	N.S.	0.00000807	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.
7A_17140288	G/A	4/293	N.S.	N.S.	N.S.	N.S.	0.00000156	1.56E-08	N.S.	N.S.
7A_1854867	T/C	3/294	N.S.	N.S.	N.S.	N.S.	N.S.	0.00000106	N.S.	N.S.
7A_25753459	A/G	21/276	N.S.	N.S.	N.S.	0.000000444	N.S.	N.S.	N.S.	0.00000104
7A_25872958	A/C	21/276	N.S.	N.S.	N.S.	0.000000444	N.S.	N.S.	N.S.	0.00000104
7A_25888367	C/T	22/275	N.S.	N.S.	N.S.	0.000000148	N.S.	N.S.	N.S.	0.0000023
7A_2607424	G/A	5/292	N.S.	N.S.	N.S.	N.S.	N.S.	0.00000424	N.S.	N.S.
7A_3229656	A/G	3/294	N.S.	N.S.	N.S.	N.S.	N.S.	0.00000106	N.S.	N.S.
10B_11041396	A/G	5/292	N.S.	N.S.	N.S.	N.S.	N.S.	0.00000665	N.S.	N.S.
10B_13932694	A/G	5/292	N.S.	N.S.	N.S.	N.S.	N.S.	3.41E-08	N.S.	N.S.
10B_4601363	T/C	5/292	N.S.	N.S.	N.S.	N.S.	N.S.	0.000000466	N.S.	N.S.
10B_9081256	C/T	7/290	N.S.	N.S.	N.S.	N.S.	N.S.	0.000000273	N.S.	N.S.

Table S2. Candidate genes located at (in bold) or near (200 kbp up or downstream) the selected genomic sites associated with LI.

Genomic site	Genes at site
1A:20268178	Uncharacterized
1A:21091933	Uncharacterized
1B:22065152	**Setaria italica acyl transferase 1 ***
1B:22075858	Uncharacterized
3A:8334174	Uncharacterized
7A:17140288	Uncharacterized
2B:17571095	**Panicum virgatum LRR receptor-like serine/threonine-protein kinase RGI3 ****
1A:31534966	**Panicum virgatum cilia- and flagella-associated protein 58-like**
7A:25753459	**Setaria viridis glyoxylate/hydroxypyruvate reductase HPR3-like**
7A:25872958	Uncharacterized
7A:25888367	Setaria viridis pentatricopeptide repeat-containing protein At3g61360 ***

* Acyltransferase was associated with suberin synthesis in roots and seed coats (Beisson *et al.*, 2007)

** A gene family expressed in various parts of the roots (Wu *et al.*, 2016)

*** PPR proteins were involved in root and shoot development and abiotic stress (Lee *et al.*, 2019; Su *et al.*, 2019)