Title

The DC1 domain protein BINUCLEATE POLLEN is required for pollen development in

Arabidopsis thaliana

Running head

BNP is required for male gametogenesis

Corresponding author

D.F. Fiol. Instituto de investigaciones Biológicas IIB-CONICET- Universidad Nacional

de Mar del Plata, Funes 3250, Mar del Plata, Buenos Aires, 7600, Argentina. Tel +54

2234753030. Fax +54 22344724143.

E-mail: diefiol@mdp.edu.ar

Subject area: Growth and development

Number of black and white figures: 0

Number of colour figures: 7

Number of tables: 5

Number of supplementary figures: 9

Number of supplementary tables: 2

Title

The DC1 domain protein BINUCLEATE POLLEN is required for pollen development in

Arabidopsis thaliana

Running head

BNP is required for male gametogenesis

Author's names and affiliations

Leonardo A. Arias[1], Sebastián D'Ippolito[1], Jésica Frik[1], Natalia L. Amigo, Fernanda

Marchetti, Claudia A. Casalongué, Gabriela C. Pagnussat, Diego F. Fiol[*]

Instituto de investigaciones Biológicas IIB-CONICET- Universidad Nacional de Mar del

Plata, Mar del Plata, Buenos Aires, 7600, Argentina.

[1] these authors contributed equally to this work.

[*] Corresponding author: Diego Fernando Fiol

E-mail: diefiol@mdp.edu.ar. Fax +54 22344724143

Abstract

Development of the male gametophyte is a tightly regulated process that requires precise control of cell division and gene expression. A relevant aspect to understand the events underlying pollen development regulation constitutes the identification and characterization of the genes required for this process. In this work we showed that the DC1 domain protein BINUCLEATE POLLEN (BNP) is essential for pollen development and germination. Pollen grains carrying the defective *BNP* allele failed to complete mitosis II and are impaired in pollen germination. By yeast two-hybrid analysis and bimolecular fluorescence complementation assays, we identified a set of BNP-interacting proteins. Among confirmed interactors we found NAC family transcriptional regulators Vascular Plant One-Zinc Finger 1 (VOZ1) and VOZ2. VOZ1 localization changes during pollen development, moving to the vegetative nucleus at the tricellular stage. We observed that this relocalization requires BNP, as in the absence of BNP in pollen from *bnp/BNP* plants, VOZ1 nuclear localization is impaired. As *voz1voz2* double mutants showed the same developmental defect observed in *bnp* pollen grains, we propose that BNP requirement to complete microgametogenesis could be linked to its interaction with VOZ1/2 proteins. BNP could have a role of scaffold protein, recruiting VOZ1/2 to the endosomal system into assemblies that are required for their further translocation to the nucleus, where they act as transcriptional regulators.

Keywords: *Arabidopsis thaliana,* Divergent C1 domain, microgametogenesis*,* Pollen Mitosis II, endosomal system, Vascular Plant One-Zinc Finger.

1. Introduction

The development of the male gametophyte of flowering plants (pollen grain) takes place inside the anthers and involves two sequential phases: microsporogenesis and microgametogenesis. Microsporogenesis begins when a diploid microsporocyte undergoes meiotic division to form a tetrad of haploid microspores (Twell, 2011). During microgametogenesis a large vacuole is formed inside the released microspores and positions the nucleus in the periphery of the cell. This stage is followed by an asymmetric division or Pollen Mitosis I (PMI) that yields a large vegetative cell and a small germ cell. The germ cell is subsequently engulfed within the cytoplasm of the vegetative cell. The germ cell then undergoes a second division, Pollen Mitosis II (PMII), which results in twin sperm cells and yields the mature tricellular pollen. In species such as *Arabidopsis thaliana*, PMII takes place inside the pollen grain prior to anthesis. Although transcriptional studies have provide significant advances in the understanding of the molecular mechanisms controlling male gametophyte development, the picture is still far from complete (reviewed in (Hafidh et al., 2016). Studies on isolated mutants have also proved to be of great importance to identify novel genes and cellular mechanisms involved in this process (Borg et al., 2009).

Divergent C1 (DC1) domains are cysteine/histidine-rich zinc finger modules found exclusively in the plant kingdom. DC1 domains resemble the C1 domains present in protein kinase C (PKC) and in over a dozen of animal proteins (Colon-Gonzalez and Kazanietz, 2006). However, C1 domains are not frequent in plants, as they were only identified in diacylglycerol kinases (Escobar-Sepúlveda et al., 2017). C1 domains have the ability to bind the secondary messenger diacylglycerol (DAG) but also participate in protein-protein interactions and in the targeting of proteins to the membrane (Colon-Gonzalez and Kazanietz, 2006). Sequence and structure similarities between DC1 and C1 domains suggest that some of the roles played by C1 domains in animal proteins could be fulfilled in plants by DC1 domains (D'Ippólito et al., 2017).

The Arabidopsis genome encodes for over 140 uncharacterized proteins harboring DC1 domains (D'Ippólito et al., 2017). Reports regarding genes coding for DC1 domain proteins were limited to transcriptional responses. In Arabidopsis, *At5g17960* was described as responsive to hormones and stress treatments (Bhaskar et al., 2015) and *ULI3 (At5g59920)* was related to UV-B responses (Suesslin and Frohnmeyer, 2003). In other plant species, DC1 domain proteins have been also associated with stress and hormone responses (Shinya et al., 2007; Li et al., 2010; Hwang et al., 2014; Gao et al., 2016).

The first functional characterization of a DC1 domain protein, Vacuoleless Gametophytes (VLG), was recently reported (D'Ippólito et al., 2017). VLG localizes to

prevacuolar compartments/multivesicular bodies and it was shown to be essential for the development of the female and male gametophytes. A role for VLG was proposed during vacuole biogenesis, likely through the interaction with the VAMP protein PVA12 and the GDSL-motif lipase LTL1 (D'Ippólito et al., 2017).

In this work we studied one of the closest VLG homologs, encoded by *At2g44370* and identified that the DC1 domain protein BINUCLEATE POLLEN (BNP) plays a relevant role during the male gametophyte development in Arabidopsis. In addition, we demonstrated that BNP is required for pollen germination and early sporophytic developmental stages. To functionally characterize BNP we searched for protein interactors and identified transcriptional regulators VOZ1 and VOZ2 as BNP binding proteins and we showed that VOZ proteins co-localize to BNP, supporting a physiological interaction. We showed that VOZ1 localizes in the cytoplasm at early stages of pollen development, but moves to the vegetative nucleus at the tricellular stage. We observed that this relocalization requires BNP, as in the absence of BNP in pollen from *bnp/BNP* plants, VOZ1 nuclear localization is impaired. As *voz1voz2* mutants also exhibit pollen developmental defects (Celesnik et al., 2013), we propose that the interaction between BNP and VOZ proteins outside the nucleus might impact further on VOZ1/2 activities, resulting in functional consequences for gene expression. Thus, BNP could act as a scaffold protein recruiting VOZ1/2 and likely other proteins to the endosomal system into assemblies that might be required to accomplish VOZ1/2 translocation to the nucleus.

2. Results
BNP codes for a DC1 domain containing protein
At2g44370, here named *BINUCLEATE POLLEN* (*BNP*), is a single exon gene that encodes for a 250-amino acid long protein with three DC1 domains (Fig. 1B). Phylogenetic analysis of Arabidopsis DC1 domain protein family grouped BNP into the most divergent cluster, with At5g40590 and VLG being the closest homologs to BNP (Fig. 1C). Multiple sequence alignments of selected protein sequences of this cluster and DC1 domain proteins from *Capsicum annuum*, tobacco, wheat and cotton showed amino acid conservation mostly restricted to the residues that define the DC1 signature motif, which might be involved in Zn^{2+} coordination and folding of the domain (Fig. 1D). On the other hand, amino acids corresponding to loop regions in the folded domain - which might define its binding capabilities (Rahman and Das, 2015)- showed very low similarity, suggesting a diversity of interactors for the aligned proteins.

bnp is a gametophytic-acting mutation

Two Arabidopsis lines with T-DNA insertions in *BNP* (SALK_114889 –here named *bnp-*
1 and GK-008E01 -here named *bnp-2*) were studied (Fig. 1A). After backcrossing to
the Col-0 ecotype, the progeny of self-pollinated plants was genotyped. No
homozygous mutant plants were recovered in the offspring from *bnp-1/BNP* or *bnp-*
2/BNP self-pollinated plants. The ratio of hemizygous to WT plants in the progeny of
self-fertilized *bnp-1/BNP* plants was 1.44 (*n* = 183). For *bnp-2/BNP* plants, segregation
analysis for sulfadiazine resistance showed a ratio of resistant to sensitive plants of
1.31 (*n* = 2634). The proportion of plants harboring the transgene recovered in the
progeny of self-fertilized plants was lower than the Mendelian segregation ratio
expected for diploid sporophytic (3:1) or embryo lethal (2:1) mutants suggesting a
gametophytic defect. Furthermore, no obvious sporophytic phenotypes were observed
in the hemizygous plants, suggesting that *bnp* is a recessive mutation (Fig. S1). The
siliques of *bnp/BNP* plants showed a small percentage of aborted seeds, not different
to WT plants (Fig. S1).

As no homozygous plants were recovered and siliques from hemizygous plants
looked normal, the *BNP* mutation could be affecting pollen development, germination
or early development. To test this hypothesis, we performed reciprocal crosses
between *bnp/BNP* and WT plants and calculated the transmission efficiency of the
mutant allele (TE = resistant/WT offspring x 100) (Table 1). Only 15.5% of
microgametophytes carrying *bnp-1* and 22.0 % of the microgametophytes carrying *bnp-*
2 transmitted the insertion to the next generation (p<0.0001, Chi square test),
indicating that *bnp* affects male gametophyte development or function (Table 1). No
significant defects were observed for transmission through the female gametophyte
(90.8% and 95.9% for *bnp-1* and *bnp-2*, respectively, Table 1, p-values 0.51 and 0.66
respectively, Chi-square test).

Although the insertions in *BNP* compromised the transmission of the mutation
through the male gametophyte, a fraction (about 15-22%) of pollen grains are still be
able to transmit the insertion to the next generation. However, we did not recover
homozygous mutant plants in the self-progeny of hemizygous plants. Thus, we
investigated whether embryogenesis or seed germination were affected by the
mutation analyzing pistils and seeds from *bnp/BNP* plants. No obvious embryo defects
were observed in siliques from *bnp/BNP* plants, as neither developing siliques nor seed
sets showed differences from WT siliques (Fig. S1). However, seeds from *bnp/BNP*
self-pollinated plants showed reduced germination rates when compared to WT plants
(Table 2). Further analysis of the germinated seeds showed that a fraction of the
seedlings arrested soon after germination. Values of seedling lethality obtained were of
0.9% for WT (*n* = 440), 3.4% (*n* = 754) for seeds from *bnp-1/BNP* self-pollinated plants

and 4.5% (*n* = 973) for *bnp-2/BNP* self-pollinated plants (Table 2). PCR genotyping of
arrested seedlings showed that 75% (*n* = 12) corresponded to *bnp/bnp* homozygous
individuals.

Altogether these results suggest that the mutation might cause impairment in
male gametogenesis or function, reduced seed germination and seedling lethality,
which combined, explain the absence of homozygous mutant plants in the progeny of
selfed hemizygous plants.

183 Pollen development in *bnp* mutants is arrested at the binuclear stage
Viability of mature pollen was analyzed in *bnp-1/BNP* and *bnp-2/BNP* by means of
Alexander staining and FDA staining. No differences were observed compared to
pollen obtained from WT plants, indicating that the *bnp* mutation does not affect pollen
viability (Fig. S2).

We later analyzed nuclear composition of mature pollen using DAPI staining.
Normal mature pollen contains three nuclei: a vegetative nucleus and two sperm nuclei
generated by the mitotic division of the germ cell. Mature pollen from WT plants
showed 99.2% (*n* = 772) of trinucleate grains. On the other hand, *bnp/BNP* plants
presented only about 57-59% of trinucleate mature pollen grains (56.9% (*n* = 1123) in
bnp-1/BNP and 59.0% (*n* = 981) in *bnp-2/BNP*) (Table 3). The rest were found in a
binucleate stage, suggesting developmental arrest.

A genetic complementation assay confirmed that the insertions in *BNP* are
causing the abnormalities observed during pollen development. *bnp-1/BNP*
hemizygous plants transformed with a *ProBNP:BNP:GFP* translational fusion construct
showed trinucleate mature pollen grains at similar levels of WT plants (Fig. S3). This
result was verified in T3 plants of five independent complemented lines.

In order to better characterize the defect observed during pollen development,
bnp-1/BNP plants were crossed to *qrt1* homozygous plants, which are impaired in a
pectin methylesterase responsible for the separation of microspores tetrads (Francis et
al., 2006). F2 plants that were hemizygous for the insertion in *BNP* and homozygous
for *qrt1* (*bnp/BNP qrt1/qrt1*) were further analyzed. Mature tetrads were collected and
stained with DAPI. Consistently with our previous results, most tetrads (99.1%) of *bnp-*
1/BNP qrt1/qrt1 plants contained at least one microspore and, in most cases, two
microspores with an abnormal number of nuclei (Fig. 2A), indicating a failure to
complete PMII in about 46% of the pollen grains analyzed.

We next assessed germination of pollen grains in the tetrads. As shown in Fig.
2, germination rates were 37.7% in pollen collected from *bnp-1/BNP qrt1/qrt1* plants
and 68.2% in pollen grains from *BNP/BNP qrt1/qrt1* plants (Fig. 2F). This indicates a

reduction of 44.7% in the germination rate in pollen from *bnp-1/BNP qrt1/qrt1* plants
compared to pollen grain tetrads from *BNP/BNP qrt1/qrt1* plants. As the reduction in
the rate of germination observed in the hemizygous background *bnp-1/BNP* was less
than 50%, this suggests that although severely affected, a small fraction of *bnp* pollen
grains are still able to germinate.

Analysis of pollen grain germination in *bnp-2/BNP* plants showed comparable
values, as we calculated a decrease in germination rate of about 36% compared to
pollen grains from WT plants. Germination rates of 43% ($n = 787$) and 67% ($n = 443$)
were recorded for pollen grains collected from *bnp-2/BNP* and WT plants respectively.

These results showed that *BNP* is required to complete pollen development,
specifically to undergo PMII. Arrested pollen grains were viable, but germination was
impaired, which explains the reduced transmission of *bnp* through the male
gametophyte (Table 1).

BNP is expressed in pollen and young sporophytic tissues
Expression of *BNP* was studied in Arabidopsis pollen expressing the *BNP-GFP* fusion
driven by the *BNP* promoter, which was proved to be functional, as it complemented
the phenotype in *bnp/BNP* plants (Fig. S3). BNP-GFP was detected in cytoplasmic
speckles following a punctuate pattern at early stages of pollen development
(microspore tetrad and released microspore) and in discrete bigger compartments at
bicellular and tricellular stages (Fig. 3A and 3H). The same localization pattern was
observed in the pollen tube (PT) of germinating pollen grains (Fig. 3I-J).

To gain further knowledge of BNP roles *in planta*, a reporter construction
expressing the *GUS* gene under the control of *BNP* promoter was used to analyze *BNP*
expression. GUS activity accumulated at early stages of seedling development, mainly
in vascular tissues and developing leaves. It was also observed associated with
anthers at very early stages of floral development, indicating that expression of *BNP*
can be detected before meiosis (Fig. 3K-S). Expression data observed were
comparable to transcriptional data available in the efp Browser Data Analysis Tool
(https://bar.utoronto.ca/eplant/) (Fig. S4). These results also indicated that BNP could
be functional not only during pollen development but also at additional developmental
stages, such as sporogenesis, germination and early stages of sporophyte
development.

BNP interacts with transcription factors VOZ1 and VOZ2
To gain insight into possible functions of BNP we carried out a yeast-two-hybrid screen
to identify potential BNP protein interactors. Two closely related NAC family

transcription factors, VOZ1 and VOZ2, were identified among interactors (all interacting
clones are presented in Table S1). Interestingly, VOZ1 and VOZ2 were reported to
bind a pollen-specific promoter element (Mitsuda et al., 2004). BNP was also retrieved
in the Y2H screening as a self-interactor (Table S1). To confirm BNP-VOZ1, BNP-
VOZ2 and BNP-BNP interactions, we performed Bimolecular Fluorescence
complementation (BiFC) assays.

We made constructs to express BNP and each of the candidate proteins fused
either to the N- or the C-terminal of yellow fluorescent protein (YFP) fragments to
generate BNP-N-YFP and VOZ1-C-YFP, VOZ2-C-YFP or BNP-C-YFP fusion
constructs. The fusion constructs under control of the *CaMV 35S* promoter were used
to transiently transform *N. benthamiana* leaves and BiFC signals were analyzed using
confocal microscopy (Fig. 4). YFP signal was detected with VOZ1, VOZ2 and with BNP
indicating the interaction of the pairs BNP- VOZ1, BNP-VOZ2 and BNP-BNP (Fig. 4).
Control *N. benthamiana* leaves transfected either with BNP-N-YFP, VOZ1-C-YFP,
VOZ2-C-YFP or BNP-C-YFP did not show any BiFC signal (Fig. 4). An additional
control experiment using VLG (as BNP closest homolog, sharing 72.0% identity) did
not show an interaction with VOZ1, VOZ2 and BNP indicating the specificity of the
observed interactions (Fig. S5).

BNP mainly localizes to Rha1 positive vesicles and co-localizes with VOZ1 and
VOZ2
We used the lipophilic dye FM4-64 to gain insight on the subcellular localization of
BNP. After internalization, this dye is sequentially distributed to different organelle
membranes, following the endocytic pathway (Bolte et al., 2004). We followed FM4-64
in roots of plants expressing the fusion protein BNP-GFP at different time points. We
observed no co-localization of internalized FM4-64 and BNP-GFP at early time points
of incubation (10-30 min) discarding localization of BNP-GFP to early endosomes or to
components of the *trans*-Golgi Network (Fig. S6 A-E). However, high levels of co-
localization were observed at about 120 min of incubation, supporting a localization in
late endosomes (Fig. S6 F-J). In agreement with that, BNP-GFP did not co-localize
with Brefeldin bodies caused by the addition of the endosomal recycling inhibitor
Brefeldin A to FM4-64 treated roots (Fig. S6 K-O). In addition, treatment with the PI3K
inhibitor LY294002, which was reported to cause swelling and association of late
endosomes (Takáč et al, 2013) caused a significant increase in the area of BNP
compartments (Fig. S7).

To assess BNP localization, we used *N. benthamiana* to transiently express
BNP-GFP together with a late endosome/prevacuolar compartments/multivesicular

bodies marker, the Rab5 GTPase Rha1 fused to RFP (Foresti et al., 2010) or with a

Golgi apparatus marker, the rat sialyltransferase St fused to RFP (Wee et al., 1998).

Representative images shown in Fig. 5 and Pearson and Spearman correlation (PSC)

values indicated that BNP-GFP co-localized with Rha1-RFP (Fig. 5A-E) and did not co-

localize with St-RFP (Fig. 5F-J), suggesting that BNP is mainly confined to late

endosome/prevacuolar compartments/multivesicular bodies associated vesicles.

To further confirm that BNP and VOZ proteins are indeed localized in the same

subcellular compartment, we performed a co-localization study. *N. benthamiana* plants

were transiently co-transfected with constructions to express BNP-GFP and either

VOZ1-RFP or VOZ2-RFP. In these transfected cells BNP-GFP and VOZ1-RFP or

VOZ2-RFP can be found separately, however, a significant shared pixel population

indicates partial co-localization, confirmed by moderate correlation scores for Pearson

and Spearman tests (Fig. 5K-T).

BNP is required for nuclear localization of VOZ1 in pollen

We next analyzed VOZ1 localization in pollen at anthesis from *BNP/BNP* and

hemizygous *bnp-1/BNP* plants (Fig. 6). We observed that in mature trinucleated pollen

VOZ1 localizes to the cytoplasm and to the vegetative nucleus, however, in mature but

binucleated pollen from *bnp-1/BNP* plants, VOZ1 is localized only in the cytoplasm

(Fig. 6). This result suggested that the absence of a functional BNP could be affecting

the localization of VOZ to the nucleus

To test this, we transformed *BNP/BNP* and hemizygous *bnp-1/BNP* plants, both

in the *qrt1* background, with *ProUb10:VOZ1-RFP* and analyzed VOZ1-RFP intracellular

localization during pollen development (Fig. 7). The introduction of the mutation in the

qrt1 background allows to compare and analyze any difference in VOZ1 localization

that might arise from an impairment in BNP, as pollen tetrads of the *bnp-1/BNP*

genotype are constituted by two mutant (binucleate) and two WT (trinucleate) pollen

cells.

At earlier developmental stages (microspore and bicellular pollen) VOZ1 was

detected at the cytoplasm with very low signal intensities, showing no differences

between *qrt1/qrt1*, *BNP/BNP, VOZ1-RFP* and *qrt1/qrt1*, *bnp-1/BNP, VOZ1-RFP*

genotypes (Fig. 7A-H). In mature WT pollen, however, VOZ1-RFP was detected at

higher levels and its localization appeared distributed between the cytoplasm and the

nuclei (Fig. 7I-L). Most of the pollen tetrads (91.5%) from *qrt1/qrt1*, *BNP/BNP, VOZ1-*

RFP plants presented VOZ1 nuclear localization in the four cells of the tetrad (Table 4).

However, most of the tetrads from the *qrt1/qrt1*, *bnp-1/BNP,* VOZ1-RFP genotype

presented VOZ1 nuclear localization only in one or two pollen cells (85.2%, Table 4).

Nuclear composition of mature pollen from plants expressing VOZ1-RFP (*qrt1/qrt1,*
BNP/BNP, VOZ1-RFP and *qrt1/qrt1, bnp-1/BNP, VOZ1-RFP*) shown in Table 5
presented comparable values to the results showed in Fig. 2A for *qrt1/qrt1, BNP/BNP*
and *qrt1/qrt1, bnp-1/BNP* respectively.

Next, we analyzed if the lack of a functional BNP could also affect the levels of
VOZ1 in pollen. We quantified VOZ1-RFP by measuring red fluorescence intensity in
mature pollen cells of *qrt1/qrt1*, *BNP/BNP,* VOZ1-RFP and *qrt1/qrt1*, *bnp-2/BNP,*
VOZ1-RFP plants (Fig. S8). Interestingly, lower red fluorescence levels were recorded
in *qrt1/qrt1, bnp-2/BNP,* VOZ1-RFP plants suggesting that BNP could have a role in
the promotion of VOZ1 stabilization. Altogether, these results strongly suggest that
BNP is involved in VOZ1 localization, as its nuclear localization relies on the presence
of a functional BNP.

We then analyzed BNP localization during pollen development using WT plants
harboring stable *BNP-GFP* and *VOZ1-RFP* constructs. We observed that both, VOZ1
and BNP, localized in the cytoplasm at early developmental stages. Contrasting VOZ1,
no nuclear localization was recorded for BNP (Fig. S9). Partial co-localization was
detected in areas in the cytosol were BNP and VOZ1 might interact at microspore and
bicellular stages. This was not observed in mature pollen, where VOZ1 show both
nuclear and cytoplasmic localization. As BNP remained in the cytosol at every
developmental stage, forming speckles during binucleate and mature pollen stages,
any role related to VOZ proteins translocation/function would be performed outside the
nucleus and before pollen maturation.

3. Discussion
In this study we identified *BNP*, a gene coding for a DC1 domain protein that is
required for pollen development and germination. Analysis of hemizygous *bnp/BNP*
plants showed that transmission through the male gametophyte is severely impaired;
about half of pollen grains from *bnp/BNP* plants resulted arrested by the binucleate
stage and their germination was significantly reduced.

By means of Y2H and BiFC experiments, we identified and confirmed VOZ1
and VOZ2 as BNP interacting proteins. VOZ1 and VOZ2 are transcription factors that
belong to the NAC transcription factor family, and were reported as involved in the
control of flowering time (Celesnik et al., 2013; Yasui and Kohchi, 2014), either through
the binding of PhyB (Yasui et al., 2012), Flowering Locus C (Mimida et al., 2011) or
Constans (Koguchi et al., 2017; Kumar et al., 2018). VOZ proteins are also involved in
biotic and abiotic stress responses (Neill et al., 2002; Nakai et al., 2013; Song et al.,
2018; Schwarzenbacher et al., 2020). VOZ1 and VOZ2 were proved to be functionally

redundant, as single mutants for each gene showed normal phenotypes, but *voz1voz2*

double mutants displayed phenotypic defects (Yasui et al., 2012; Celesnik et al., 2013;

Nakai et al., 2013; Kumar et al., 2018). Interestingly, *voz1voz2* plants presented the

same pollen developmental defect observed in *bnp/BNP* plants, as a fraction of their

pollen grains contained only two nuclei –the vegetative nucleus and a single

reproductive nucleus (Celesnik et al., 2013). This fact supports a physiological role for

the interaction between BNP and VOZ proteins during pollen development.

BNP was localized mainly in Rha1 positive vesicles, which are defined as late

endosomes/ prevacuolar compartments/ multivesicular bodies. VOZ proteins are

functional in the nucleus, but they were initially identified in cytoplasmic speckles and a

cytoplasmic-nuclear translocation was proposed to modulate VOZ activity as

transcriptional regulators (Yasui et al., 2012; Koguchi et al., 2017). VOZ1 and VOZ2 do

not contain neither a transmembrane region nor localization signals such as NLS or

NES (Jensen et al., 2010). This suggests that VOZ1 and VOZ2 nuclear/cytoplasmic

localization must rely on another mechanism. Endocytic proteins have been previously

shown to participate in the spatiotemporal regulation of transcription factors (Palfy et

al., 2012; Wu et al., 2012; Serrano et al., 2016). They may facilitate the assembly of

multi-molecular complexes through protein–protein interaction domains and act as

regulators (Pyrzynska et al., 2009). As can be concluded from the analysis showed in

Fig. 6, 7 and Table 3 a similar situation could be taking place between the pairs BNP-

VOZ1, where VOZ1 is required to interact with BNP to reach the nuclear import

machinery.

The involvement of the endomembrane system in nuclear translocation has

been previously reported, as several membrane receptors undergo endocytosis and

are transported to the nucleus via endosomal trafficking (Mukherjee et al., 2006, Wang

et al., 2010; Wang et al., 2020). However the mechanisms of the process are still under

investigation and several mechanisms are proposed to allow the cargo reach the

nuclear import machinery, including retrograde movement of vesicles through the Golgi

and ER, the release of the cargo in the cytoplasm or the exposing of a NLS that is

recognized by importin- β (Ek-Ramos et al., 2014).

Implications of VOZ1/2 nuclear translocation may be related to VOZ1/2 role as

a transcriptional regulator, with such regulation occurring in the nucleus (Yasui et al.,

2012). Interestingly, VOZ proteins nuclear localization was rarely detected, and the

addition of NLS signals was usually required to place it the nucleus (Yasui et al., 2012;

Schwarzenbacher et al., 2020) However it was reported that degradation of VOZ1/2

takes also place in the nucleus, facilitated by the E3 ubiquitin ligase BRUTUS (Selote

et al., 2018). Thus, BNP role could be involved in the function and/or in the degradation
of VOZ1/2.

Our results also indicated that BNP could be essential not only during pollen
development but also at additional developmental stages, such as germination and
early stages of sporophyte development. In agreement with this, *BNP* expression was
detected very early during seedling development, in particular in vascular tissues (Fig.
3). The expression pattern of BNP also suggests that it could play roles in other
sporophytic tissues, either through the interaction with VOZ TFs, that are also
ubiquitously expressed through the plant (Yasui et al., 2012), or through the interaction
with other proteins. Potential BNP interacting proteins identified by Y2H experiments
(Table S1) provide a promising list of candidates that might be involved with BNP
function in other plant tissues.

Only one additional DC1 domain protein, VLG, has been characterized so far
and was likewise proved to be required for pollen development. However, *VLG*
deficiency caused microgametogenesis arrest at PMI, an earlier stage, yielding
unviable uninucleate pollen grains (D'Ippólito et al., 2017). On the other hand, VLG was
also essential for female gametogenesis (D'Ippólito et al., 2017).

Notably, the Arabidopsis genome encodes for over 140 uncharacterized
proteins harboring DC1 domains. Multiple sequence alignments of DC1 domain
proteins from different species showed amino acid conservation restricted to the
residues that define the DC1 signature motif, but divergence in the amino acids
corresponding to the folded domain -which might define binding capabilities (Rahman
and Das, 2015)- suggesting a diversity of interactors. Thus, DC1 domain proteins
emerge as a novel family of highly specialized modular scaffold proteins, capable to
promote the recruiting of binding targets and affect their localization and functionality.

Altogether, our results support a role for BNP acting as a scaffold protein
recruiting VOZ1 and VOZ2 to the endomembrane system. This recruitment appears to
facilitate VOZ1 -and likely VOZ2- translocation to the nucleus, promoting their activity
as transcriptional regulators during pollen development. In addition, a role of BNP in
the regulation of the levels VOZ proteins could not be discarded, as lower levels of
these proteins were present in mature pollen lacking a functional BNP.

4. Materials and Methods

Plant materials and growth conditions

Arabidopsis (Columbia [Col-0] ecotype) lines SALK_114889 (*bnp-1,* with T-DNA
insertion located in chr2 TAIR10 18,323,075) and GK-008E01 (*bnp-2,* with T-DNA
insertion located in chr2 TAIR10 18,323,371) were obtained from ABRC (Ohio State

University, Ohio, USA) (Alonso et al., 2003). Both lines were backcrossed twice to the
WT Columbia prior to their use. The presence of a single insertion in *At2g44370* and all
genotypes was confirmed by PCR-based genotyping and segregation analysis. When
indicated seeds were sterilized in 20% (v/v) sodium hypochlorite, washed with sterile
water and plated on MS plates with 50 µg/mL kanamycin or 7.5 µg/mL sulfadiazine,
stratified at 4°C for 48 h and placed at 23°C, 16 h light, 8 h dark. Resistant seedlings
were then transferred onto soil and grown under the conditions described above.
440

Molecular characterization of insertional lines
For *bnp-1*, the left border–genomic sequence junction was determined by PCR in
plants showing kanamycin resistance using the T-DNA-specific primer LBb1 combined
with the genomic sequence specific primers bnp-1 RP and bnp-1 LP (Table S2). For
bnp-2, the left border–genomic sequence junction was determined by PCR in plants
showing sulfadiazine resistance using the T-DNA–specific primer GKGT8474
combined with the genomic sequence specific primers bnp-2 RP and bnp-2 LP (Table
S2).
449

Segregation Analysis
For self-fertilization analysis, the progeny of self-pollinated hemizygous plants was
germinated on selective growth medium containing 50 µg/mL kanamycin for *bnp-1* or
7.5 µ/mL sulfadiazine for *bnp-2*, and the ratio of resistant to sensitive plants was
scored. Reciprocal crosses were performed as described previously
(Pagnussat et al., 2005).
456

Generation of vectors for plant transformation
Genomic DNA was extracted from rosette leaves or whole seedlings as described
(León et al., 2007). To generate *ProBNP:BNP-GFP* translational fusion the region
including *BNP* ORF and putative promoter (672 upstream the ATG codon) was
amplified by PCR using the primer combination listed in Table S2. The amplicon was
cloned using pENTR™ Directional TOPO® Cloning Kit (Invitrogen) using Gateway®
technology and the sequence was verified. The resulting plasmid *pENTR-ProBNP-*
BNP was subjected to the LR reaction using the destination vector pMDC107 (Curtis
and Grossniklaus, 2003). The *ProBNP:GUS* construct was generated amplifying the
BNP putative promoter region (672 upstream the ATG codon) by PCR using the primer
combination listed in Table S2. The amplicon was cloned into pENTR™/TOPO
(Invitrogen) and its sequence was verified. The resulting plasmid *pENTR-ProBNP* was
subjected to the LR reaction using the destination vector pMDC162 (Curtis and

Grossniklaus, 2003). For *Pro35S:BNP-GFP* construct, *BNP* ORF was PCR amplified

using as template clone DKLAT2G44370 obtained from ABRC (Ohio State University,

Ohio, USA) using the primer combination listed in Table S2. The amplicon was cloned

into pENTR^{TM}/TOPO (Invitrogen) and its sequence was verified. The resulting plasmid

pENTR-BNP was subjected to the LR reaction using the destination vector pMDC83

(Curtis and Grossniklaus, 2003). For *ProUb10:VOZ1-RFP* and *ProUb10:VOZ2-*

RFP constructions, respective ORFs were PCR amplified using primers listed in Table

S2, cloned in pENTR^{TM}/TOPO (Invitrogen) and sequence-verified. The resulting

plasmids *pENTR-VOZ1* and *pENTR-VOZ2* were subjected to the LR reaction using the

destination vector pUBN-RFP-DEST (Grefen et al., 2010).

Obtaining *bnp/BNP* plants in the *qrt1* background

To obtain *bnp* mutants in *qrt1* background, *bnp-1/BNP* was crossed with *qrt1*

homozygous plants. Kanamycin resistant plants from the progeny were selected and

allowed to self-fertilize. Homozygous *qrt1* plants harboring *bnp* mutations were

selected for further analysis (Johnson-Brousseau and McCormick, 2004).

In a similar way, *qrt1/qrt1-bnp-2/BNP*-VOZ1-RFP plants were obtained crossing *bnp-*

2/BNP plants expressing VOZ1-RFP with *qrt1* homozygous plants.

Transformation of *Agrobacterium tumefaciens* and Arabidopsis

Vectors were introduced into Agrobacterium strain GV3101 by electroporation.

Arabidopsis WT plants were transformed using the floral dip method (Clough and Bent,

1998). Transformant plants were selected based on their ability to survive in MS

medium with 15 mg. L^{-1} hygromycin. Resistant (green seedlings with true leaves) were

then transferred to soil and grown under the conditions described above.

Morphological and histological analyses

Pollen grain viability was assessed using Alexander's staining (Alexander, 1969) and

fluorescein diacetate (FDA) staining (Heslop-Harrison and Heslop-Harrison, 1970). For

pollen development analysis, successive buds in the same inflorescence were

collected (Lalanne and Twell, 2002), fixed in ethanol:acetic acid 3:1 until discoloration

and stored at room temperature. Anthers were washed, dissected and mounted on

microscopy slides with DAPI staining solution. For pollen tube germination, pollen

grains were incubated overnight in a dark moisture chamber at 22°C in 0.01% boric

acid, 5mM CaCl$_2$, 5mM KCl, 1 mM MgSO$_4$, 10% sucrose pH 7.5 and 1.5% agarose, as

described previously (Boavida and McCormick, 2007). For GUS staining, tissues from

ProBNP:GUS carrying plants were collected and incubated in GUS staining solution as

already described (Pagnussat et al., 2007). Gametophytes were observed on a Zeiss
Axio Imager A2 microscope using DIC optics. Seedlings were observed on a Nikon
SMZ800 Stereomicroscope and images were captured with a Canon DS126431
camera.
511

Yeast two-hybrid screen
The Y2H screening was performed by Hybrigenics Services S.A.S. (Paris, France). A
DNA fragment encoding Arabidopsis At2g44370 (amino acids 1-250) was PCR-
amplified and cloned into pB66 as C-terminal fusion to Gal4 DNA-binding domain
(Gal4-At2g44370) and used as a bait to screen Universal Arabidopsis Normalized
library containing 3.2 million of independent clones in pGADT7-RecAB vector. pB66
derive from the original pAS2ΔΔ (Fromont-Racine et al., 1997). For the Gal4 bait
construct, 35 million clones were screened using a mating approach with YHGX13
(Y187 ade2-101::loxP-kanMX-loxP, matα) and L40ΔGal4 (mata) yeast strains as
previously described (Fromont-Racine et al., 1997). 158 His+ colonies were selected
on a medium lacking tryptophan, leucine and histidine. The prey fragments of the
positive clones were amplified by PCR and sequenced at their 5' and 3' junctions. The
resulting sequences were used to identify the corresponding interacting proteins in the
GenBank database (NCBI) using a fully automated procedure, and in total, 23 different
clones were identified (Table S1). A confidence score (PBS, for Predicted Biological
Score) was attributed to each interaction as previously described (Formstecher et al.,
2005).
529

Co-localization experiments in *Nicotiana benthamiana*
A. tumefaciens strain GV3101 carrying *Pro35S:BNP-GFP* was co-infiltrated with *A.*
tumefaciens cells carrying either the late endosome marker *RFP-Rha1* or the Golgi
apparatus marker *St-RFP*, in leaf epidermal cells of *N. benthamiana* plants. Infiltrated
leaves were analyzed 48-72 h after infiltration on a confocal microscope (Nikon Eclipse
C1 Plus) using EZ-C1 3.80 imaging software and Ti-Control. Pearson and Spearman
correlation (PSC) plugin for ImageJ was used to calculated correlation coefficients from
a minimum of 400 individual regions of interest corresponding with punctate structures
were manually selected from at least 10 independent cell images (French et al., 2008).
539

Co-localization experiments in Arabidopsis
Arabidopsis root cells expressing a *35S:BNP-GFP* construction were used to analyze
the co-localization between BNP and the endocytic tracer FM4-64. Roots were
incubated for the indicated times with 6 μM FM4-64 solution, washed twice, and then

incubated in liquid medium or with the indicated inhibitors. For Brefeldin A treatments,
incubation was performed in 50 µM Brefeldin A solution for 15 min. For LY294002
treatments, incubation was performed in 100µM LY294002 solution for 120 min.
Confocal microscopy images were obtained at the indicated times with a confocal
microscope (Nikon Eclipse C1 Plus) using EZ-C1 3.80 imaging software and Ti-
Control. Pearson and Spearman correlation (PSC) plugin for ImageJ was used to
calculated correlation coefficients from a minimum of 400 individual regions of interest
from at least 10 independent cell images (French et al., 2008).

Bimolecular fluorescence complementation (BiFC) analysis
The cDNA sequences of At2g44370 (*BNP*), At1g28520 (*VOZ1*), At2g42400 (*VOZ2*)
and At2g17740 (*VLG*) were PCR amplified using the primers listed in Table S2. The
PCR products were cloned into pENTR^{TM}/TOPO (Invitrogen), sequenced and
recombined through BP reaction into BiFC destination plasmids pUBN-YN and pUBN-
YC (Grefen et al., 2010). The binary plasmids were then transformed into *A.*
tumefaciens strain GV3101 by electroporation. Split nYFP- and cYFP-tagged protein
pairs and the gene-silencing suppressor p19 were co-expressed in *N. benthamiana*
leaves by *A. tumefaciens* -mediated inoculation. Plant leaves were examined 48-72 h
post-infiltration with a confocal microscope (Nikon Eclipse C1 Plus) using a Super Fluor
40.0x/1.30/0.22 Oil objective, numerical aperture 1.300. Acquisition was performed
using EZ-C1 3.80 imaging software and Ti-Control. All pictures were acquired with the
same settings.

Quantification of VOZ1-RFP in mature pollen
High quality images of mature pollen grains from the genotypes *qrt1/qrt1, BNP/BNP,*
VOZ1-RFP and *qrt1/qrt1, bnp-1/BNP, VOZ1-RFP* plants were acquired and red
fluorescence was quantified. Red fluorescence background of mature pollen grains
from plants not expressing VOZ1-RFP was subtracted. Fluorescence was expressed in
arbitrary units relative to the calculated cell area of each pollen grain. Data points were
plotted on a violin plot using GraphPad, median (blue line) and quartiles (black lines)
are indicated. Box: Mean values ± SD and number of data points (n) is indicated. A
significant difference of the means was indicated using unpaired t-test.

Bioinformatics and phylogenetic analysis
Sequences from the DC1 domain family were identified using the Basic Local
Alignment Search Tool (BLAST) from NCBI (National Center for Biotechnology

Information) using initially BNP as a reference and retrieved sequences in iterative
searches. Protein sequences alignments were performed using MEGA7 (version
7.0.14) (Kumar et al., 2016). Phylogenetic trees were constructed using the neighbor-
joining method and the default settings of MEGA7 (version 7.0.14) (Kumar et al., 2016).
The evolutionary distances were computed using the Poisson correction method
(Zuckerkandl and Pauling, 1965) and are in the units of the number of amino acid
substitutions per site. The trees were drawn using FigTree (version 1.4.3)
(http://tree.bio.ed.ac.uk/). Graphic display of identities was visualized using Geneious
(version 9.1.4) (http://www.geneious.com) based on an identity matrix (Kearse et al.,
2012).
590

591 **5. Data availability**

592 All newly generated sequences are available for peer-review and registered in public
593 repositories. Sequence data from this article can be found in the Arabidopsis Genome
594 Initiative database under accession numbers: BNP (At2g44370), VOZ1 (At1g28520),
595 VOZ2 (At2g42400) and VLG (At2g17740).
596

6. Funding

This work was supported by grants to DFF from Agencia Nacional de Promoción
Cientifica y Técnica Argentina (PICT2013-1524 and PICT2017-0232), Consejo
Nacional de Investigaciones Cientificas y Técnicas (CONICET, PIP-0200), Comisión
de Investigaciones Cientificas (CIC-PBA) and Universidad Nacional de Mar del Plata; a
grant to GCP from the Howard Hughes Medical Institute (IECS award 55007430).
603

7. Acknowledgements

We would like to thank Michael Blatt and Christopher Grefen (Glasgow University, UK)
for providing pUB-Destination vectors, Jurgen Denecke (University of Leeds, UK) for
providing the constructs ST-RFP and Rha-RFP, Nicolas Setzes for assistance with
statistical analysis and Daniela Villamonte for technical assistance with confocal
microscopy. L.A.A., J.F. and F.M. are fellows of CONICET; N.L.A. is a fellow of
ANPCyT; D.F.F., G.C.P., C.A.C. and S.D. are researchers from CONICET.
611

8. Author contributions

D.F.F. conceived the original screening and research plans; D.F.F., G.C.P. and C.A.C.
supervised the experiments; L.A.A., S.D., J.F., N.L.A and F.M. performed most of the
experiments; D.F.F., G.C.P and L.A.A. designed the experiments and analyzed the
data; D.F.F. conceived the project and wrote the article with contributions of all the

authors; D.F.F. agrees to serve as the author responsible for contact and ensures
communication
619

620 **9. Conflicts of interest**

621 The authors declare no conflicts of interest.
622

10. References

Alexander MP (1969) Differential staining of aborted and nonaborted pollen. Stain
Technol. 44, 117-122

Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK,
Zimmerman J, Barajas P, Cheuk R, al e (2003) Genome-wide insertional
mutagenesis of *Arabidopsis thaliana*. Science 301, 653–657

Bhaskar RV, Mohanty B, Verma V, Wijaya E, Kumar PP (2015) A hormone-responsive
C1-domain-containing protein At5g17960 mediates stress response in
Arabidopsis thaliana. PLoS One 10, e0115418

Boavida LC, McCormick S (2007) Temperature as a determinant factor for increased
and reproducible in vitro pollen germination in *Arabidopsis thaliana*. Plant J. 52,
570-582

Bolte S, Talbot C, Boutte Y, Catrice O, Read ND, Satiat-Jeunemaitre B (2004) FM-dyes
as experimental probes for dissecting vesicle trafficking in living plant cells. J.
Microsc. 214, 159-173

Borg M, Brownfield L, Twell D (2009) Male gametophyte development: A molecular
perspective. J. Exp. Bot. 60, 1465-1478

Celesnik H, Ali GS, Robison FM, Reddy AS (2013) *Arabidopsis thaliana* VOZ
(VASCULAR PLANT ONE-ZINC FINGER) transcription factors are required for
proper regulation of flowering time. Biol. Open 2, 424-431

Clough SJ, Bent AF (1998) Floral dip: A simplified method for Agrobacterium-mediated
transformation of *Arabidopsis thaliana*. Plant J. 16, 735-743

Colon-Gonzalez F, Kazanietz MG (2006) C1 domains exposed: From diacylglycerol
binding to protein-protein interactions. Biochim. Biophys. Acta 1761, 827-837

Curtis MD, Grossniklaus U (2003) A gateway cloning vector set for high-throughput
functional analysis of genes in planta. Plant Physiol. 133, 462–469.

D'Ippólito S, Arias LA, Casalongué CA, Pagnussat GC, Fiol DF (2017) The DC1-
domain protein VACUOLELESS GAMETOPHYTES is essential for female and
male gametophyte development in Arabidopsis. Plant J. 90, 261-275

Ek-Ramos MJ, Avila J, Nelson Dittrich AC, Su D, Gray JW, Devarenne TP (2014) The
tomato cell death suppressor Adi3 is restricted to the endosomal system in

response to the *Pseudomonas syringae* effector protein AvrPto. PLOS One 9,
e110807

Escobar-Sepúlveda HF, Trejo-Téllez LI, Pérez-Rodríguez P, Hidalgo-Contreras JV,
Gómez-Merino FC (2017) Diacylglycerol kinases are widespread in higher plants
and display inducible gene expression in response to beneficial elements, metal,
and metalloid ions. Front. Plant Sci. 8, 129

Foresti O, Gershlick DC, Bottanelli F, Hummel E, Hawes C, Denecke J (2010) A
recycling-defective vacuolar sorting receptor reveals an intermediate
compartment situated between prevacuoles and vacuoles in tobacco. Plant Cell
22, 3992-4008

Formstecher E, Aresta S, Collura V, Hamburger A, Meil A, Trehin A, Reverdy C, Betin
V, Maire S, Brun C, Jacq B, Arpin M, Bellaiche Y, Bellusci S, Benaroch P,
Bornens M, Chanet R, Chavrier P, Delattre O, Doye V, Fehon R, Faye G, Galli T,
Girault JA, Goud B, de Gunzburg J, Johannes L, Junier MP, Mirouse V,
Mukherjee A, Papadopoulo D, Perez F, Plessis A, Rosse C, Saule S, Stoppa-
Lyonnet D, Vincent A, White M, Legrain P, Wojcik J, Camonis J, Daviet L (2005)
Protein interaction mapping: A Drosophila case study. Genome Res. 15, 376-384

Francis KE, Lam SY, Copenhaver GP (2006) Separation of Arabidopsis pollen tetrads
is regulated by quartet1, a pectin methylesterase gene. Plant Physiol. 142, 1004-
1013

French AP, Mills S, Swarup R, Bennett MJ, Pridmore TP (2008) Colocalization of
fluorescent markers in confocal microscope images of plant cells. Nature Protoc.
3, 619-628

Fromont-Racine M, Rain JC, Legrain P (1997) Toward a functional analysis of the
yeast genome through exhaustive two-hybrid screens. Nat. Genet. 16, 277-282

Gao S, Yang L, Zeng HQ, Zhou ZS, Yang ZM, Li H, Sun D, Xie F, Zhang B (2016) A
cotton miRNA is involved in regulation of plant response to salt stress. Sci.
Reports 6, 19736

Grefen C, Donald N, Hashimoto K, Kudla J, Schumacher K, Blatt MR (2010) A
ubiquitin-10 promoter-based vector set for fluorescent protein tagging facilitates
temporal stability and native protein distribution in transient and stable expression
studies. Plant J. 64, 355-365

Hafidh S, Fila J, Honys D (2016) Male gametophyte development and function in
angiosperms: A general concept. Plant Reprod. 29, 31-51

Heslop-Harrison J, Heslop-Harrison Y (1970) Evaluation of pollen viability by
enzymatically induced fluorescence; intracellular hydrolysis of fluorescein
diacetate. Stain Technol. 45, 115-120

Hwang IS, Choi DS, Kim NH, Kim DS, Hwang BK (2014) The pepper cysteine/histidine-
rich DC1 domain protein CaDC1 binds both RNA and DNA and is required for
plant cell death and defense response. New Phytol. 201, 518-530

Jensen Michael K, Kjaersgaard T, Nielsen Michael M, Galberg P, Petersen K, O'Shea
C, Skriver K (2010) The *Arabidopsis thaliana* NAC transcription factor family:
Structure–function relationships and determinants of ANAC019 stress signalling.
Biochem J. 426, 183-196

Johnson-Brousseau SA, McCormick S (2004) A compendium of methods useful for
characterizing Arabidopsis pollen mutants and gametophytically-expressed
genes. Plant J. 39, 761-775

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S,
Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A
(2012) Geneious basic: An integrated and extendable desktop software platform
for the organization and analysis of sequence data. Bioinformatics 28, 1647-1649

Koguchi M, Yamasaki K, Hirano T, Sato MH (2017) VASCULAR PLANT ONE-ZINC-
FINGER PROTEIN 2 is localized both to the nucleus and stress granules under
heat stress in Arabidopsis. Plant Signal. Behav. 12, e1295907

Kumar S, Choudhary P, Gupta M, Nath U (2018) VASCULAR PLANT ONE-ZINC
FINGER1 (VOZ1) and VOZ2 interact with CONSTANS and promote
photoperiodic flowering transition. Plant Physiol. 176, 2917-2930

Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular evolutionary genetics
analysis version 7.0 for bigger datasets. Mol. Biol. Evol. 33, 1870-1874

Lalanne E, Twell D (2002) Genetic control of male germ unit organization in
Arabidopsis. Plant Physiol. 129, 865-875

León G, Holuigue L, Jordana X (2007) Mitochondrial complex ii is essential for
gametophyte development in Arabidopsis. Plant Physiol. 143, 1534-1546

Li C, Lv J, Zhao X, Ai X, Zhu X, Wang M, Zhao S, Xia G (2010) TaCHP: A wheat zinc
finger protein gene down-regulated by abscisic acid and salinity stress plays a
positive role in stress tolerance. Plant Physiol. 154, 211-221

Mimida N, Kidou S, Iwanami H, Moriya S, Abe K, Voogd C, Varkonyi-Gasic E, Kotoda
N (2011) Apple Flowering Locus T proteins interact with transcription factors
implicated in cell growth and organ development. Tree Physiol. 31, 555-566

Mitsuda N, Hisabori T, Takeyasu K, Sato MH (2004) VOZ; isolation and
characterization of novel vascular plant transcription factors with a one-zinc finger
from *Arabidopsis thaliana*. Plant Cell Physiol. 45, 845-854

Mukherjee S, Tessema M, Wandinger-Ness A (2006) Vesicular trafficking of tyrosine
kinase receptors and associated proteins in the regulation of signaling and
vascular function. Circul. Res. 98, 743-756

Nakai Y, Nakahira Y, Sumida H, Takebayashi K, Nagasawa Y, Yamasaki K, Akiyama
M, Ohme-Takagi M, Fujiwara S, Shiina T, Mitsuda N, Fukusaki E, Kubo Y, Sato
MH (2013) Vascular plant one-zinc-finger protein 1/2 transcription factors
regulate abiotic and biotic stress responses in Arabidopsis. Plant J. 73, 761-775

Neill SO, Gould KS, Kilmartin PA, Mitchell KA, Markham KR (2002) Antioxidant
capacities of green and cyanic leaves in the sun species, *Quintinia serrata.*
Funct. Plant Biol. 29, 1437-1443

Pagnussat GC, Yu HJ, Ngo QA, Rajani S, Mayalagu S, Johnson CS, Capron A, Xie LF,
Ye D, Sundaresan V (2005) Genetic and molecular identification of genes
required for female gametophyte development and function in Arabidopsis.
Development 132, 603-614

Pagnussat GC, Yu HJ, Sundaresan V (2007) Cell-fate switch of synergid to egg cell in
Arabidopsis *eostre* mutant embryo sacs arises from misexpression of the BEL1-
like homeodomain gene BLH1. Plant Cell 19, 3578-3592

Palfy M, Remenyi A, Korcsmaros T (2012) Endosomal crosstalk: Meeting points for
signaling pathways. Trends Cell Biol. 22, 447-456

Pyrzynska B, Pilecka I, Miaczynska M (2009) Endocytic proteins in the regulation of
nuclear signaling, transcription and tumorigenesis. Mol. Oncol. 3, 321-338

Rahman GM, Das J (2015) Modeling studies on the structural determinants for the
dag/phorbol ester binding to C1 domain. J. Biomol Struct Dynam. 33, 219-232

Schwarzenbacher RE, Wardell G, Stassen J, Guest E, Zhang P, Luna E, Ton J (2020)
The IBI1 receptor of β-aminobutyric acid interacts with VOZ transcription factors
to regulate abscisic acid signaling and callose-associated defense. Mol. Plant 13,
1455-1469

Selote D, Matthiadis A, Gillikin JW, Sato MH, Long TA (2018) The E3 ligase brutus
facilitates degradation of VOZ1/2 transcription factors. Plant Cell Environ. 41,
2463-2474

Serrano I, Buscaill P, Audran C, Pouzet C, Jauneau A, Rivas S (2016) A non canonical
subtilase attenuates the transcriptional activation of defence responses in
Arabidopsis thaliana. eLife 5,

Shinya T, Galis I, Narisawa T, Sasaki M, Fukuda H, Matsuoka H, Saito M, Matsuoka K
(2007) Comprehensive analysis of glucan elicitor-regulated gene expression in
tobacco BY-2 cells reveals a novel MYB transcription factor involved in the
regulation of phenylpropanoid metabolism. Plant Cell Physiol. 48, 1404-1413

Song C, Lee J, Kim T, Hong JC, Lim CO (2018) VOZ1, a transcriptional repressor of
DREB2C, mediates heat stress responses in Arabidopsis. Planta 247, 1439-1448
Suesslin C, Frohnmeyer H (2003) An Arabidopsis mutant defective in UV-B light-
mediated responses. Plant J. 33, 591-601
Takáč T, Pechan T, Šamajová O, Šamaj J (2013) Vesicular trafficking and stress
response coupled to PI3K inhibition by LY294002 as revealed by proteomic and
cell biological analysis. J Proteome Res 12, 4435-4448
Twell D (2011) Male gametogenesis and germline specification in flowering plants.
Sex. Plant Reprod. 24, 149-160
Wang X, Xu M, Gao C, Zeng Y, Cui Y, Shen W, Jiang L (2020) The roles of
endomembrane trafficking in plant abiotic stress responses. J. Integr. Plant Biol.
62, 55-69
Wang Y-N, Wang H, Yamaguchi H, Lee H-J, Lee H-H, Hung M-C (2010) COPI-
mediated retrograde trafficking from the Golgi to the ER regulates EGFR nuclear
transport. Biochem. Biophys. Res. Comm, 399, 498-504
Wee EG-T, Sherrier DJ, Prime TA, Dupree P (1998) Targeting of active
sialyltransferase to the plant Golgi apparatus. Plant Cell 10, 1759-1768
Wu P, Wee P, Jiang J, Chen X, Wang Z (2012) Differential regulation of transcription
factors by location-specific egf receptor signaling via a spatio-temporal interplay
of erk activation. PloS One 7, e41354
Yasui Y, Kohchi T (2014) VASCULAR PLANT ONE-ZINC FINGER1 and VOZ2 repress
the FLOWERING LOCUS C clade members to control flowering time in
Arabidopsis. Biosci. Biotechnol. Biochem. 78, 1850-1855
Yasui Y, Mukougawa K, Uemoto M, Yokofuji A, Suzuri R, Nishitani A, Kohchi T (2012)
The phytochrome-interacting VASCULAR PLANT ONE-ZINC FINGER1 and
VOZ2 redundantly regulate flowering in Arabidopsis. Plant Cell 24, 3248-3263
Zuckerkandl E, Pauling L (1965) Evolutionary divergence and convergence in proteins.
Evolv. Genes Prot. 97, 97-166

Figure Legends

Figure 1. BNP codes for a DC1 domain containing protein. (A) Schematic view of a
region of Arabidopsis chromosome 2 (from 18,323,538 to 18,321,537 bp -TAIR10)
showing BNP genomic location and bnp-1 and bnp-2 insertion sites. (B) BNP amino
acid sequence with boxed DC1 domains. (C) Phylogenic tree of 140 DC1 domain
containing proteins in Arabidopsis. Bootstrap test results (1000 replicates) of the major

nodes are indicated. IDs of sequences grouped in clusters I to VIII are listed. (D)

Alignment of BNP with its closest Arabidopsis homologs and CaDC1 (AEI52549),

TaCHP (ACU80555), NtDC1a (BAF80452) and NtDC1b (BAF80453). Phylogenic tree

with bootstrap test results (1000 replicates) for the alignment of the complete 18

sequences is shown. Amino acid region represented for each sequence is indicated.

DC1 domains are framed, signature residues and binding loop regions are indicated

(C=Cys, H=His, — = loop). Higher intensity in grey scale denotes higher sequence

similarity.

Figure 2. Pollen development and germination is impaired in *bnp-1* mutants. (A)

Quantification of phenotypes observed in *BNP/BNP* and *bnp-1/BNP* plants in *qrt1/qrt1*

background. (B-E) Nuclear configuration types revealed by DAPI (4',6-Diamidino-2-

phenylindole) staining. Bars = 10 μm. (F) Quantification of pollen tube germination in

tetrads from *BNP/BNP* and *bnp-1/BNP* plants in *qrt1/qrt1* background. (G-K) Pollen

tube germination in tetrads. Bars = 20 μm.

Figure 3. *BNP* is expressed in pollen and young sporophytic tissues. (A-D)

Confocal images showing BNP-GFP through microgametogenesis. (E-H) Overlay

between DIC and GFP fluorescence. (A and E) Microspore tetrad, (B and F) released

microspore, (C and G) bicellular stage and (D and H) tricellular stage. (I) Image

showing a projection along the z axis of all planes showing GFP fluorescence. (J)

Overlay of DIC and GFP fluorescence showing a germinating pollen grain. (K-S) GUS

activity in Arabidopsis transgenic plants carrying the *pBNP:GUS* construct. (K)

Seedling 24 h after imbibition, (L) 48 h after imbibition, (M) 6 days post germination and

(N) 12 days post germination. (O) Root with lateral root primordium, (P) primary root

with root hairs, (Q) third leaf 8 days after imbibition (R) third leaf 14 days after

imbibition. (S) GUS signal was detected inside anthers at floral developmental stage

10. Three independent transgenic lines were used for this study. Pictures are

representative of the results obtained in all analyzed lines. Bars = 10 μm.

Figure 4. BNP interacts with itself, VOZ1 and VOZ2. Bimolecular fluorescence

complementation (BiFC) analysis of the interaction of BNP with itself, VOZ1 and VOZ2

in *N. benthamiana* leaves. Representative confocal microscopy images showing that

BNP interacts with (A–C) itself, (D-F) VOZ1 and (G-I) VOZ2. Reconstituted YFP

fluorescence (YFP) or DIC images of *N. benthamiana* leaves are shown in epidermal

cells co-infiltrated with *Agrobacterium tumefaciens* harboring the indicated constructs.

Control *N. benthamiana* leaves transfected with (J-L) BNP-N-YFP, (M-O) BNP-C-YFP,

(P-R) VOZ1-C-YFP, (S-U) VOZ2-C-YFP do not show any BiFC signal. The last row (C,

F, I, L, O, R, U) correspond to the magnification of regions of interest (insets) indicated

in the first row. Bars = 20 μm.

Figure 5. BNP co-localizes with VOZ1 and VOZ2 to Rha1 positive vesicles. (A, F,

K and P) Localization of BNP-GFP transiently expressed in *N. benthamiana* epidermal

cells. (B) Fluorescent marker for late endosomes/prevacuolar compartments/

multivesicular bodies, RFP-Rha1. (C) Merged image of BNP-GFP and RFP-Rha1. (D)

Shows an overlay with the corresponding bright-field image. (E, J, O and T) Pearson

and Spearman correlation (PSC) test calculated using a plugin for ImageJ; r values

indicate the level of co-localization ranging from +1 for perfect co-localization to −1 for

negative correlation. (G) Fluorescent marker for Golgi apparatus, St-RFP. (H) Merged

image of BNP-GFP and St-RFP. (I) Shows an overlay with the corresponding bright-

field image. (L) Localization of RFP-VOZ1. (M) Merged image of BNP-GFP and RFP-

VOZ1. (N) Shows the corresponding bright field image of the cells showed in (M). (Q)

Localization of RFP-VOZ2. (R) Merged image of BNP-GFP and RFP-VOZ2. (S) Bright

field image of the cells showed in (R). Bars = 20 µm.

Figure 6. VOZ1 localization in mature pollen from WT and *bnp-1/BNP* plants.

Mature pollen from (A- E) BNP/BNP, VOZ1-RFP and (F – J) bnp-1/BNP, VOZ1-RFP

plants was analized. Representative z-stack images of confocal micrographs displaying

(A, F) DAPI nuclear staining, (B, G) VOZ1-RFP fluorescence and (E, J) bright-field

images. (C, H) Merged DAPI and VOZ1-RFP fluorescence images and (D, I) merged

DAPI, VOZ1-RFP fluorescence and bright-field images Yellow arrows indicate the

vegetative nucleus and white arrows indicate the generative nucleus. Bars = 10 µm.

Figure 7. Nuclear localization of VOZ1 is affected in *bnp* mutants.

Pollen tetrads from *qrt1/qrt1, BNP/BNP, VOZ1-RFP* (A, B, E, F, I, J) and *qrt1/qrt1, bnp-*

1/BNP, VOZ1-RFP (C, D, G, H, K, L) plants were analyzed at microspore (A-D),

binucleate (E-H) and mature tricellular (I-L) stages. Representative z-stack images of

confocal micrographs displaying VOZ1-RFP fluorescence (B, D, F, H, J, L) and bright

field images (A, C, E, G, I, K) are shown. Arrowheads indicate nuclear localized VOZ1.

Bars = 10 µm.

Table 1. Transmission efficiency of *bnp* alleles. Transmission efficiency of the *bnp-1* and *bnp-2* alleles in reciprocal crosses between mutant and WT plants.

Parental genotypes		Genotype of progeny† Number of individuals		Transmission efficiency of:	p-value*
Female	Male	WT	*bnp-1/BNP*	*bnp-1*	
bnp-1/BNP	WT	98	89	♀ 90.8%	0.51
WT	*bnp-1/BNP*	181	28	♂ 15.5%	<0.0001
		WT	*bnp-2/ BNP*	*bnp-2*	
bnp-2/BNP	WT	217	208	♀ 95.9%	0.66
WT	*bnp-2/BNP*	254	56	♂ 22%	<0.0001

†No homozygous plants were detected; * Chi-square test for a 1:1 segregation hypothesis (degrees of freedom = 1).

Table 2. Viability of the progeny in *bnp* mutants. Seed germination and seedling lethality of the progeny of self-pollinated *bnp-1/BNP*, *bnp-2/BNP* and WT plants.

Parental genotype	% viable progeny	% unviable progeny		p-value*
		not germinated	seedling lethality	
WT	98.6 (435/441)	0.2 (1/441)	1.1 (5/441)	
bnp-1/BNP	94.8 (715/754)	1.7 (13/754)	3.4 (26/754)	<0.001
bnp-2/BNP	91.4 (688/753)	2.8 (21/753)	5.8 (44/753)	<0.00001

In parenthesis, number of individuals observed over the total analyzed. * Chi-square test for a hypothesis of no relationship between the genotype of the parents and the viability of the progeny (degree of freedom = 1).

Table 3. Nuclear composition of mature pollen in *bnp* mutants. Mature pollen from *bnp-1/BNP*, *bnp-2/BNP* and WT plants was stained using DAPI.

Genotype	% trinucleate pollen	p-value*
WT	99.2 (766/772)	
bnp-1/BNP	56.9 (639/1123)	<0.0001
bnp-2/BNP	59.0 (579/981)	<0.0001

In parenthesis, number of individuals observed over the total analyzed. * The statistical significance was determined by Student's *t*-test.

Table 4. VOZ1 localization in mature pollen in *bnp* mutants expressing VOZ1. Mature pollen tetrads from *qrt1/qrt1, BNP/BNP, VOZ1-RFP* and *qrt1/qrt1, bnp-1/BNP, VOZ1-RFP* plants.

Genotype	Observed VOZ1 localization				
	Four nuclear	Three nuclear, one cytoplasmic	Two nuclear, two cytoplasmic	One nuclear, three cytoplasmic	Four cytoplasmic
qrt1/qrt1, BNP/BNP, VOZ1-RFP	91.5 % (65/71)	7.0 % (5/71)	1.4 % (1/71)	0 % (0/71)	0 % (0/71)
qrt1/qrt1, bnp-1/BNP, VOZ1-RFP	1.3 % (1/76)	11.8 % (9/76)	57.8% (44/76)	27.6% (21/76)	1.3 % (1/76)

In parenthesis, number of individuals observed over the total analyzed.

Table 5. Nuclear composition of mature pollen in *bnp* mutants expressing VOZ1. Mature pollen tetrads from *qrt1/qrt1, BNP/BNP, VOZ1-RFP* and *qrt1/qrt1, bnp-1/BNP, VOZ1-RFP* plants.

Genotype	Observed nuclear phenotype			
	Four trinucleate	Three trinucleate, one binucleate	Two trinucleate, two binucleate	Two binucleate, one tetranucleate
qrt1/qrt1, BNP/BNP, VOZ1-RFP	97.3 % (71/73)	2.7 % (2/73)	0 % (0/73)	0 % (0/73)
qrt1/qrt1, bnp-1/BNP, VOZ1-RFP	3.6 % (4/110)	18.2 % (20/110)	77.3 % (85/110)	0.9 % (1/110)

In parenthesis, number of individuals observed over the total analyzed.

A

At2g44380>   *bnp-1* *bnp-2*   At2g44370

ATG   TAA   At2g44360>

200 bp

B

MNW MAARKPSVR**H**PS**H**NHPLRSHKAQVEEEII**C**SG**C**DLDLIGAAFK**C**TKSE**C**DYFL**H**KS**C**FELPRETR**H**KA**H**PDHPLILLYSPPYESTYT**C**DA**C**GEYGSGFTYN**C**SI**C**QYDV**H**VG**C**VSMPESVER

EG**H**A**H**PLTLLYRSPYQNGLIFN**C**DV**C**QETVPDNLWSYY**C**KE**C**DYGT**H**LHS**C**AVEEEEEAEEEEEPKRGGGSARGNTKSGGNKGGRGSAASELAAMLEAQREMEKMQIELHMEMQRAKIAKKARKACLKLİ

1   122

123   250

C

D

A

Genotype	Observed nuclear phenotype			
	Four trinucleate (image B)	Three trinucleate, one binucleate (image C)	Two trinucleate, two binucleate (image D)	Two trinucleate, one tetranucleate (image E)
BNP/BNP	96.9 % (187/193)	3.1 % (6/193)	0 % (0/193)	0 % (0/193)
bnp-1/BNP	0.9 % (2/229)	11.8 % (27/229)	83.4 % (191/229)	3.9 % (9/229)

F

Genotype	Pollen germination rate	Observed tetrad phenotype				
		Four germinated (image G)	Three germinated (image H)	Two germinated (image I)	One germinated (image J)	none germinated (image K)
BNP/BNP	68.2 % (573/840)	35.2 % (74/210)	35.6 % (75/210)	10.5 % (22/210)	5.7 % (12/210)	12.8 % (27/210)
bnp-1/BNP	37.7 % (291/772)	0% (0/193)	8.8 % (17/193)	46.6% (90/193)	31.1 % (60/193)	13.4 % (26/193)

| | *qrt1/qrt1, BNP/BNP, VOZ1-RFP* | *qrt1/qrt1, bnp-1/BNP, VOZ1-RFP* |