Title

The DC1 domain protein BINUCLEATE POLLEN is required for pollen development in *Arabidopsis thaliana*

Running title

BNP is required for male gametogenesis

Author's names and affiliations

Leonardo A. Arias[1], Sebastián D'Ippolito[1], Jésica Frik[1], Natalia L. Amigo, Claudia A. Casalongué, Gabriela C. Pagnussat, Diego F. Fiol[*]

Instituto de investigaciones Biológicas IIB-CONICET- Universidad Nacional de Mar del Plata, Mar del Plata, Buenos Aires, 7600, Argentina.

[1] these authors contributed equally to this work.

[*] Corresponding author: Diego Fernando Fiol

E-mail: diefiol@mdp.edu.ar. Fax +54 22344724143

Corresponding author

D.F. Fiol. Instituto de investigaciones Biológicas IIB-CONICET- Universidad Nacional de Mar del Plata, Funes 3250, Mar del Plata, Buenos Aires, 7600, Argentina. Tel +54 2234753030. Fax +54 22344724143.

E-mail: diefiol@mdp.edu.ar

ABSTRACT

Development of the male gametophyte is a tightly regulated process that requires
precise control of cell division and gene expression. A relevant aspect to understand
the events underlying pollen development regulation constitutes the identification and
characterization of the genes required for this process. In this work we showed that the
DC1 domain protein BINUCLEATE POLLEN (BNP) is essential for pollen development
and germination. Pollen grains carrying the defective *BNP* allele failed to complete
mitosis II and are impaired in pollen germination. By yeast two hybrid screening and
bimolecular fluorescence complementation assays, we identified a set of BNP-
interacting proteins. Among confirmed interactors we found NAC family transcriptional
regulators Vascular plant One-Zinc finger 1 (VOZ1) and VOZ2. VOZ1 localization
changes during pollen development, moving to the vegetative nucleus at the tricellular
stage. We observed that this relocalization requires BNP, as in their absence in pollen
from *bnp/BNP* plants, VOZ1 nuclear localization resulted impaired. As *voz1voz2* double
mutants showed the same developmental defect observed in *bnp* pollen grains*,* we
propose that BNP requirement to complete microgametogenesis could be linked to its
interaction with VOZ1/2 proteins. BNP could have a role of scaffold protein, recruiting
VOZ1/2 to the endosomal system into assemblies that are required for their further
translocation to the nucleus, where they act as transcriptional regulators.

Keywords: *Arabidopsis thaliana,* Divergent C1 domain, microgametogenesis*,* Pollen
Mitosis II, endosomal system, Vascular Plant One-Zinc Finger.

INTRODUCTION

The development of the male gametophyte of flowering plants (pollen grain) takes place inside the anthers and involves two sequential phases: microsporogenesis and microgametogenesis. Microsporogenesis begins when a diploid microsporocyte undergoes meiotic division to form a tetrad of haploid microspores (Twell, 2011). During microgametogenesis a large vacuole is formed inside the released microspores and positions the nucleus in the periphery of the cell. This stage is followed by an asymmetric division or Pollen Mitosis I (PMI) that yields a large vegetative cell and a small germ cell. The germ cell is subsequently engulfed within the cytoplasm of the vegetative cell. The germ cell then undergoes a second division, Pollen Mitosis II (PMII), which results in twin sperm cells and yields the mature tricellular pollen. In species such as *Arabidopsis thaliana*, PMII takes place inside the pollen grain prior to anthesis. Although transcriptional studies have provide significant advances in the understanding of the molecular mechanisms controlling male gametophyte development, the picture is still far from complete (reviewed in (Hafidh et al., 2016). Studies on isolated mutants have also proved to be of great importance to identify novel genes and cellular mechanisms involved in this process (Borg et al., 2009).

Divergent C1 (DC1) domains are cysteine/histidine-rich zinc finger modules found exclusively in the plant Kingdom. DC1 domains resemble the C1 domains present in PKC and in over a dozen of animal proteins (Colon-Gonzalez and Kazanietz, 2006). However, C1 domains are not frequent in plants, as they were only identified in diacylglycerol kinases (Escobar-Sepúlveda et al., 2017). C1 domains have the ability to bind the secondary messenger diacylglycerol (DAG) but also participate in protein-protein interactions and in the targeting of proteins to the membrane (Colon-Gonzalez and Kazanietz, 2006). Sequence and structure similarities between DC1 and C1 domains suggest that some of the roles played by C1 domains in animal proteins could be fulfilled in plants by DC1 domains (D'Ippólito et al., 2017).

The Arabidopsis genome encodes for over 140 uncharacterized proteins harboring DC1 domains. Reports regarding genes coding for DC1 domain proteins were limited to transcriptional responses. In Arabidopsis, *At5g17960* was described as responsive to hormones and stress treatments (Bhaskar et al., 2015) and *ULI3 (At5g59920)* was related to UV-B responses (Suesslin and Frohnmeyer, 2003). In other plant species, DC1 domain proteins have been also associated with stress and hormone responses (Shinya et al., 2007; Li et al., 2010; Hwang et al., 2014; Gao et al., 2016).

The first functional characterization of a DC1 domain protein, Vacuoleless Gametophytes (VLG), was recently reported (D'Ippólito et al., 2017). VLG localizes to

prevacuolar compartments/multivesicular bodies and it was proved to be essential for the development of the female and male gametophytes. A role for VLG was proposed during vacuole biogenesis, likely through the interaction with the VAMP protein PVA12 and the GDSL-motif lipase LTL1 (D'Ippólito et al., 2017).

In this work we identified the DC1 domain protein BINUCLEATE POLLEN (BNP) as an essential protein for male gametophyte development in Arabidopsis. In addition, we demonstrated that BNP is required for pollen germination and early sporophytic developmental stages. To functionally characterize BNP we searched for protein interactors and identified transcriptional regulators VOZ1 and VOZ2 as BNP binding proteins and we showed that VOZ proteins co-localize to BNP, supporting a physiological interaction. We showed that VOZ1 localizes in the cytoplasm at early stages of pollen development, but moves to the vegetative nucleus at the tricellular stage. We observed that this relocalization requires BNP, as in their absence in pollen from *bnp/BNP* plants, VOZ1 nuclear localization resulted impaired. As *voz1voz2* mutants also exhibit pollen developmental defects (Celesnik et al., 2013), we propose that the interaction between BNP and VOZ proteins outside the nucleus might impact further on VOZ1/2 activities, resulting in functional consequences for gene expression. Thus, BNP could act as a scaffold protein recruiting VOZ1/2 and likely other proteins to the endosomal system into assemblies that might be required to accomplish VOZ1/2 translocation to the nucleus.

RESULTS

BNP codes for a DC1 domain containing protein

At2g44370, here named *BINUCLEATE POLLEN* (*BNP*), is a single exon gene that encodes for a 250-amino acid long protein with three DC1 domains (Fig. 1B). Phylogenetic analysis of Arabidopsis DC1 domain protein family grouped BNP into the most divergent cluster, being At5g40590 and VLG its closest homologs (Fig. 1C). Multiple sequence alignments of selected protein sequences of this cluster and DC1 domain proteins from *C. annuum*, tobacco, wheat and cotton showed amino acid conservation mostly restricted to the residues that define the DC1 signature motif, which might be involved in Zn^2 coordination and folding of the domain (Fig. 1D). On the other hand, amino acids corresponding to loop regions in the folded domain -which might define its binding capabilities (Rahman and Das, 2015)- showed very low similarity, suggesting a diversity of interactors for the aligned proteins.

***bnp* is a gametophytic mutation**

Two Arabidopsis lines with T-DNA insertions in *BNP* (SALK_114889 –here named *bnp-*
1 and GK-008E01 -here named *bnp-2*) were studied (Fig. 1A). After backcrossing to
the Col-0 ecotype, the progeny of self-crossed plants was genotyped. No homozygous
mutant plants were recovered in the offspring from *bnp-1/BNP* or *bnp-2/BNP* self-
pollinated plants. The ratio of hemizygous to WT plants in the progeny of self-fertilized
bnp-1/BNP plants was 1.44 (*n* = 183). For *bnp-2/BNP* plants, segregation analysis for
sulfadiazine resistance showed a ratio of resistant to sensitive plants of 1.31 (*n* =
2634). The proportion of plants harboring the transgene recovered in the progeny of
self-fertilized plants was lower than the Mendelian segregation ratio expected for
diploid sporophytic (3:1) or embryo lethal (2:1) mutants suggesting a gametophytic
defect. Furthermore, no obvious sporophytic phenotypes were observed in the
hemizygous plants, suggesting that *bnp* is a recessive mutation (Fig. S1). The siliques
of *bnp/BNP* plants showed a small percentage of aborted seeds, not different to WT
plants (Fig. S1).

As no homozygous plants were recovered and siliques from hemizygous plants
looked normal, the *BNP* mutation could be affecting pollen development, germination
or early development. To test this hypothesis, we performed reciprocal crosses
between *bnp/BNP* and WT plants and calculated the transmission efficiency of the
mutant allele (TE = resistant/WT offspring x 100) (Table 1). Only 15.5% of
microgametophytes carrying *bnp-1* and 22.0 % of the microgametophytes carrying *bnp-*
2 transmitted the insertion to the next generation (p<0.0001, Chi square test),
indicating that *bnp* affects male gametophyte development or function (Table 1). No
significant defects were observed for transmission through the female gametophyte
(90.8% and 95.9% for *bnp-1* and *bnp-2*, respectively, table 1, p-values 0.51 and 0.66
respectively, Chi-square test).

Although the insertions in *BNP* compromised the transmission of the mutation
through the male gametophyte, a fraction (about 15-22%) of pollen grains might still be
able to transmit the insertion to the next generation. However, we did not recover
homozygous mutant plants in the self-progeny of hemizygous plants. Thus, we
investigated whether embryogenesis or seed germination were affected by the
mutation analyzing pistils and seeds from *bnp/BNP* plants. No obvious embryo defects
were observed in siliques from *bnp/BNP* plants, as neither developing siliques nor seed
sets showed differences from WT siliques (Fig. S1). However, seeds from *bnp/BNP*
self-pollinated plants showed reduced germination rates when compared to WT plants
(94.8% for *bnp-1/BNP* (*n* = 754), 93.3% for *bnp-2/BNP* (*n* = 973) and 98.9% for WT (*n*
= 440)). Further analysis of the germinated seeds showed that a fraction of the
seedlings arrested soon after germination. Values of seedling lethality obtained were of

0.9% for WT (*n* = 440), 3.4% (*n* = 754) for seeds from *bnp-1/BNP* self-pollinated plants

and 4.5% (*n* = 973) for *bnp-2/BNP* self-pollinated plants (Table 2). PCR genotyping of

arrested seedlings showed that 75% (*n* = 12) corresponded to *bnp/bnp* homozygous

individuals.

Altogether these results suggest that the mutation might cause impairment in

male gametogenesis or function, reduced seed germination and seedling lethality,

which combined, explain the absence of homozygous mutant plants in the progeny of

selfed hemizygous plants.

Pollen development in *bnp* mutants is arrested at the binuclear stage

Viability of mature pollen was analyzed in *bnp-1/BNP* and *bnp-2/BNP* by means of

Alexander staining and FDA staining. No differences were observed compared to

pollen obtained from WT plants, indicating that the *bnp* mutation does not affect pollen

viability (Fig. S2).

We later analyzed nuclear composition of mature pollen using DAPI staining.

Normal mature pollen contains three nuclei: a vegetative nucleus and two sperm nuclei

generated by the mitotic division of the generative nucleus. Mature pollen from WT

plants showed 99.2% (*n* = 772) of trinucleate grains. On the other hand, *bnp/BNP*

plants presented only about 57-59% of trinucleate mature pollen grains (56.9% (*n* =

1123) in *bnp-1/BNP* and 59.0% (*n* = 981) in *bnp-2/BNP*). The rest were found in a

binucleate stage, suggesting developmental arrest.

A genetic complementation assay confirmed that the insertions in *BNP* are

causing the abnormalities observed during pollen development. *bnp-1/BNP*

hemizygous plants transformed with a *ProBNP:BNP:GFP* translational fusion construct

showed trinucleate mature pollen grains at similar levels of WT plants (Fig. S3). This

result was verified in T3 plants of five independent complemented lines.

In order to better characterize the defect observed during pollen development,

bnp-1/BNP plants were crossed to *qrt1* homozygous plants, which are impaired in a

pectin methylesterase responsible for the separation of microspores tetrads (Francis et

al., 2006). F2 plants that were hemizygous for the insertion in *BNP* and homozygous

for *qrt1* (*bnp/BNP qrt1/qrt1*) were further analyzed. Mature tetrads were collected and

stained with DAPI. Consistently with our previous results, most tetrads (99.1%) of *bnp-*

1/BNP qrt1/qrt1 plants contained at least one microspore and, in most cases, two

microspores with an abnormal number of nuclei (Fig. 2A), indicating a failure to

complete PMII in about 46% of the pollen grains analyzed.

We next assessed germination of pollen grains in the tetrads. As shown in Fig.

2, germination rates were 37.7% in pollen collected from *bnp-1/BNP qrt1/qrt1* plants

and 68.2% in pollen grains from *BNP/BNP qrt1/qrt1* plants (Fig. 2F). This indicates a reduction of 44.7% in the germination rate in pollen from *bnp-1/BNP qrt1/qrt1* plants compared to pollen grain tetrads from *BNP/BNP qrt1/qrt1* plants. As the reduction in the rate of germination observed in the hemizygous background *bnp-1/BNP* was less than 50%, this suggests that although severely affected, a small fraction of *bnp* pollen grains is still able to germinate.

Analysis of pollen grain germination in *bnp-2/BNP* plants showed comparable values, as we calculated a decrease in germination rate of about 36% compared to pollen grains from WT plants. Germination rates of 43% ($n = 787$) and 67% ($n = 443$) were recorded for pollen grains collected from *bnp-2/BNP* and WT plants respectively.

These results showed that *BNP* is required to complete pollen development, specifically to undergo PMII. Arrested pollen grains were viable, but germination was impaired, which explains the reduced transmission of *bnp* through the male gametophyte (Table 1).

BNP is expressed in pollen and young sporophytic tissues

Expression of *BNP* was studied in Arabidopsis plants expressing the *BNP-GFP* fusion driven by the *BNP* promoter, which was proved to be functional, as it complemented the phenotype in *bnp/BNP* plants (Fig. S3). BNP-GFP was detected in cytoplasmic speckles following a punctuate pattern at early stages of pollen development (microspore tetrad and released microspore) and in discrete bigger compartments at bicellular and tricellular stages (Fig. 3A and 3H). The same localization pattern was observed in the pollen tube (PT) of germinating pollen grains (Fig. 3I-J).

To gain further knowledge of BNP roles *in planta*, a reporter construction expressing the *GUS* gene under the control of *BNP* promoter was used to analyze *BNP* expression. GUS activity accumulated at early stages of seedling development, mainly in vascular tissues and developing leaves. It was also observed associated to anthers at very early stages of floral development, indicating that expression of *BNP* can be detected before meiosis (Fig. 3K-S). Expression data observed were comparable to transcriptional data available in the efp Browser Data Analysis Tool (https://bar.utoronto.ca/eplant/) (Fig. S4). These results also indicated that BNP could be functional not only during pollen development but also at additional developmental stages, such as sporogenesis, germination and early stages of sporophyte development.

BNP interacts with transcription factors VOZ1 and VOZ2

To gain insight into possible functions of BNP we carried out a yeast-two-hybrid screening to identify potential BNP protein interactors. Two closely related NAC family transcription factors, VOZ1 and VOZ2, were identified among interactors (a selection of the identified clones is presented in Table S1). Interestingly, VOZ1 and VOZ2 were reported to bind a pollen-specific promoter element (Mitsuda et al., 2004). BNP was also retrieved in the Y2H screening as a self-interactor (Table S1). To confirm BNP-VOZ1, BNP-VOZ2 and BNP-BNP interactions, we performed Bimolecular Fluorescence complementation (BiFC) assays.

We made constructs to express BNP and each of the candidate proteins fused either to the N- or the C-terminal of yellow fluorescent protein (YFP) fragments to generate BNP-N-YFP and VOZ1-C-YFP, VOZ2-C-YFP or BNP-C-YFP fusion constructs. The fusion constructs under control of the CaMV 35S promoter were used to transiently transform *N. benthamiana* leaves and BiFC signals were analyzed using confocal microscopy (Fig. 4). YFP signal was detected with VOZ1, VOZ2 and with BNP indicating the interaction of the pairs BNP- VOZ1, BNP-VOZ2 and BNP-BNP (Fig. 4). Control *N. benthamiana* leaves transfected either with BNP-N-YFP, VOZ1-C-YFP, VOZ2-C-YFP or BNP-C-YFP did not show any BiFC signal (Fig. 4). An additional control experiment using VLG (as BNP closest homolog, sharing 72.0% identity) showed negative results with VOZ1, VOZ2 and BNP indicating the specificity of the observed interactions (Fig. S5).

BNP mainly localizes to Rha1 positive vesicles and co-localizes with VOZ1 and VOZ2

We used the lipophilic dye FM4-64 to gain insight on the subcellular localization of BNP. After internalization, this dye is sequentially distributed to different organelle membranes, following the endocytic pathway (Bolte et al., 2004). We followed FM4-64 in roots of plants expressing the fusion protein BNP-GFP at different time points. We observed no co-localization of internalized FM4-64 and BNP-GFP after 30 min to 2 h of incubation (Fig. S6), discarding localization of BNP-GFP to early endosomes or to components of the *trans*-Golgi Network. In agreement with that, BNP-GFP did not co-localize with Brefeldin bodies caused by the addition of the endosomal recycling inhibitor Brefeldin A to FM4-64 treated roots (Fig. S6).

To assess BNP localization, we used *N. benthamiana* to transiently express BNP-GFP together with a late endosome/prevacuolar compartments/multivesicular bodies marker, the Rab5 GTPase Rha1 fused to RFP (Foresti et al., 2010) or with a Golgi apparatus marker, the rat sialyltransferase St fused to RFP (Wee et al., 1998).

Representative images shown in Fig. 5 and Pearson and Spearman correlation (PSC) values indicated that BNP-GFP co-localized with Rha1-RFP (Fig. 5A-E) and did not co-localize with St-RFP (Fig. 5F-J), suggesting that BNP is mainly confined to late endosome/prevacuolar compartments/multivesicular bodies associated vesicles.

To further confirm that BNP and VOZ proteins are indeed localized in the same subcellular compartment, we performed a co-localization study. *N. benthamiana* plants were transiently co-transfected with constructs to express BNP-GFP and either VOZ1-RFP or VOZ2-RFP. In these transfected cells, a typical punctuate expression was detected for BNP-GFP and high levels of co-localization were observed with both, VOZ1-RFP and VOZ2-RFP (Fig. 5K-T).

BNP is required for nuclear localization of VOZ1 in pollen

We next analyzed VOZ1 localization in mature pollen from *BNP/BNP* and hemizygous *bnp-1/BNP* plants (Fig. 6). We observed that in mature trinucleated pollen VOZ1 localizes to the cytoplasm and to the vegetative nucleus, however, in mature but binucleated pollen from *bnp-1/BNP* plants, VOZ1 is localized only in the cytoplasm (Fig. 6). This result suggested that the absence of a functional BNP could be affecting the localization of VOZ to the nucleus

To test this, we transformed *BNP/BNP* and hemizygous *bnp-1/BNP* plants, both in the *qrt1* background, with *ProUb10:VOZ1-RFP* and analyzed VOZ1-RFP intracellular localization during pollen development (Fig. 7). The introduction of the mutation in the *qrt1* background allows to compare and analyze any difference in VOZ1 localization that might arise from an impairment in BNP, as pollen tetrads of the *bnp-1/BNP* genotype are constituted by two mutant (binucleate) and two WT (trinucleate) pollen cells.

At earlier developmental stages (microspore and bicellular pollen) VOZ1 was detected at the cytoplasm with very low signal intensities, showing no differences between *qrt1/qrt1*, *BNP/BNP, VOZ1-RFP* and *qrt1/qrt1*, *bnp-1/BNP*, *VOZ1-RFP* genotypes (Fig. 7A-H). In mature WT pollen, however, VOZ1-RFP was detected at higher levels and its localization appeared distributed between the cytoplasm and the nuclei. Most of the pollen tetrads (91.5%) from *qrt1/qrt1*, *BNP/BNP, VOZ1-RFP* plants presented VOZ1 nuclear localization in the four cells of the tetrad (Fig. 7M). However, most of the tetrads from the *qrt1/qrt1*, *bnp-1/BNP,* VOZ1-RFP genotype presented VOZ1 nuclear localization only in one or two pollen cells (85.2%, Fig. 7M).

Next, we analyzed if the lack of a functional BNP could also affect the levels of VOZ1 in pollen. We quantified VOZ1-RFP by measuring red fluorescence intensity in mature pollen cells of *qrt1/qrt1*, *BNP/BNP,* VOZ1-RFP and *qrt1/qrt1, bnp-2/BNP,*

303 VOZ1-RFP plants (Fig. S7). Interestingly, lower red fluorescence levels were recorded
304 in *qrt1/qrt1, bnp-2/BNP,* VOZ1-RFP plants suggesting that BNP could have a role in
305 the promotion of VOZ1 stabilization.
306
Altogether, these results strongly suggest that BNP is involved in VOZ1
localization, as its nuclear localization relies on the presence of a functional BNP.
We later analyzed BNP localization during pollen development using WT plants
harboring stable *BNP-GFP* and *VOZ1-RFP* constructs. We observed that both, VOZ1
and BNP, localized in the cytoplasm at early developmental stages. Contrasting VOZ1,
no nuclear localization was recorded for BNP (Fig. S8). Partial co-localization was
detected in areas in the cytosol were BNP and VOZ1 might interact at microspore and
bicellular stages. This was not observed in mature pollen, where VOZ1 show both
nuclear and cytoplasmic localization. As BNP remained in the cytosol at every
developmental stage, forming speckles during binucleate and mature pollen stages,
any role related to VOZ proteins translocation/function would be carried out outside the
nucleus and before pollen maturation.
319

DISCUSSION
In this study we identified *BNP*, a gene coding for a DC1 domain protein that is
required for pollen development and germination. Analysis of hemyzigous *bnp/BNP*
plants showed that transmission through the male gametophyte is severely impaired;
about half of pollen grains from *bnp/BNP* plants resulted arrested at PMII stage and
their germination was significantly reduced.
By means of Y2H and BiFC experiments, we identified and confirmed VOZ1
and VOZ2 as BNP interacting proteins. VOZ1 and VOZ2 are transcription factors that
belong to the NAC transcription factor family, and were reported as involved in the
control of flowering time (Celesnik et al., 2013; Yasui and Kohchi, 2014), either through
the binding of PhyB (Yasui et al., 2012), Flowering Locus C (Mimida et al., 2011) or
Constans (Koguchi et al., 2017; Kumar et al., 2018). VOZ proteins were also related to
biotic and abiotic stress responses (Neill et al., 2002; Nakai et al., 2013; Song et al.,
2018; Schwarzenbacher et al., 2020). VOZ1 and VOZ2 were proved to be functionally
redundant, as single mutants for each gene showed normal phenotypes, but *voz1voz2*
double mutants displayed phenotypic defects (Yasui et al., 2012; Celesnik et al., 2013;
Nakai et al., 2013; Kumar et al., 2018). Interestingly, these plants presented the same
pollen developmental defect observed in *bnp/BNP* plants, as a fraction of *voz1voz2*
pollen grains contained only two nuclei –the vegetative nucleus and a single

reproductive nucleus (Celesnik et al., 2013). This fact supports a physiological role for

the interaction between BNP and VOZ proteins during pollen development.

BNP was localized mainly in Rha1 positive vesicles, which are defined as late

endosomes/ prevacuolar compartments/ multivesicular bodies. VOZ proteins are

functional in the nucleus, but they were initially identified in cytoplasmic speckles and a

cytoplasmic-nuclear translocation was proposed to modulate VOZ activity as

transcriptional regulators (Yasui et al., 2012; Koguchi et al., 2017). VOZ1 and VOZ2 do

not contain neither a transmembrane region nor localization signals such as NLS or

NES (Jensen et al., 2010). This suggests that VOZ1 and VOZ2 nuclear/cytoplasmic

localization must rely on another mechanism. Endocytic proteins have been previously

shown to participate in the spatiotemporal regulation of transcription factors (Palfy et

al., 2012; Wu et al., 2012; Serrano et al., 2016). They may facilitate the assembly of

multi-molecular complexes through protein–protein interaction domains and act as

regulators (Pyrzynska et al., 2009). As can be concluded from the analysis showed in

Fig. 6, 7 and Table 3 a similar situation could be taking place between the pairs BNP-

VOZ1, where VOZ1 is required to interact with BNP to reach the nuclear import

machinery.

The involvement of the endomembrane system in nuclear translocation has

been previously reported, as several membrane receptors undergo endocytosis and

are transported to the nucleus via endosomal trafficking (Mukherjee et al., 2006)(Wang

et al., 2010; Wang et al., 2020) However the mechanisms of the process are still under

investigation and several mechanisms are proposed to allow the cargo reach the

nuclear import machinery, including retrograde movement of vesicles through the Golgi

and ER, the release of the cargo in the cytoplasm or the exposing of a NLS that is

recognized by importin- β (Ek-Ramos et al., 2014).

Implications of VOZ1/2 nuclear translocation may be related to VOZ1/2 role as

a transcriptional regulator, as it was proposed to be exerted in the nucleus (Yasui et al.,

2012). Interestingly, VOZ proteins nuclear localization was rarely detected, and the

addition of NLS signals was usually required to place it the nucleus (Yasui et al., 2012;

Schwarzenbacher et al., 2020) However it was reported that degradation of VOZ1/2

takes also place in the nucleus, facilitated by the E3 ubiquitin ligase BRUTUS (Selote

et al., 2018). Thus, BNP role could be involved in the function and/or in the degradation

of VOZ1/2.

Our results also indicated that BNP could be essential not only during pollen

development but also at additional developmental stages, such as germination and

early stages of sporophyte development. In agreement with this, *BNP* expression was

detected very early during seedling development, in particular in vascular tissues (Fig.

3). The expression pattern of BNP also suggests that it could play roles in other
sporophytic tissues, either through the interaction with VOZ TFs, that are also
ubiquitously expressed through the plant (Yasui et al., 2012), or through the interaction
with other proteins. Potential BNP interacting proteins identified by Y2H experiments
(Supplemental Table S2) provide a promising list of candidates that might be involved
with BNP function in other plant tissues.

Only one additional DC1 domain protein, VLG, has been characterized so far
and was likewise proved to be required for pollen development. However, *VLG*
deficiency caused microgametogenesis arrest at PMI, an earlier stage, yielding
unviable uninucleate pollen grains (D'Ippólito et al., 2017). On the other hand, VLG was
also essential for female gametogenesis (D'Ippólito et al., 2017).

Notably, the Arabidopsis genome encodes for over 140 uncharacterized
proteins harboring DC1 domains. Multiple sequence alignments of DC1 domain
proteins from different species showed amino acid conservation restricted to the
residues that define the DC1 signature motif, but divergence in the amino acids
corresponding to the folded domain -which might define binding capabilities (Rahman
and Das, 2015)- suggesting a diversity of interactors. Thus, DC1 domain proteins
emerge as a novel family of highly specialized modular scaffold proteins, capable to
promote the recruiting of binding targets and affect their localization and functionality.

Altogether, our results support a role for BNP acting as a scaffold protein
recruiting VOZ1 and VOZ2 to the endomembrane system. This recruitment appears to
facilitate VOZ1 -and likely VOZ2- translocation to the nucleus, promoting their activity
as transcriptional regulators during pollen development. In addition, a role of BNP in
the regulation of the levels VOZ proteins could not be discarded, as lower levels of
these proteins were present in mature pollen lacking a functional BNP.

MATERIALS AND METHODS
Plant materials and growth conditions
Arabidopsis (Columbia [Col-0] ecotype) lines SALK_114889 (*bnp-1,* with T-DNA
insertion located in chr2 TAIR10 18,323,075) and GK-008E01 (*bnp-2,* with T-DNA
insertion located in chr2 TAIR10 18,323,371) were obtained from ABRC (Ohio State
University, Ohio, USA) (Alonso et al., 2003). Both lines were backcrossed twice to the
WT Columbia prior to their use. Single insertion in At2g44370 after backcrosses and all
genotypes were confirmed by PCR-based genotyping and segregation analysis. When
indicated seeds were sterilized in 20% (v/v) sodium hypochlorite, washed with sterile
water and plated on MS plates with 50 µg/mL kanamycin or 7.5 µg/mL sulfadiazine,

stratified at 4°C for 48 h and placed at 23°C, 16 h light, 8 h dark. Resistant seedlings

were then transferred onto soil and grown under the conditions described above.

Molecular characterization of insertional lines

For *bnp-1*, the left border–genomic sequence junction was determined by PCR in

plants showing kanamycin resistance using the T-DNA-specific primer LBb1 combined

with the genomic sequence specific primers bnp-1 RP and bnp-1 LP (Table S1). For

bnp-2, the left border–genomic sequence junction was determined by PCR in plants

showing sulfadiazine resistance using the T-DNA–specific primer GKGT8474

combined with the genomic sequence specific primers bnp-2 RP and bnp-2 LP (Table

S1).

Segregation Analysis

For self-cross analysis, the progeny of self-pollinated hemizygous plants was

germinated on selective growth medium containing 50 µg/mL kanamycin for *bnp-1* or

7.5 µ/mL sulfadiazine for *bnp-2*, and the ratio of resistant to sensitive plants was

scored. Reciprocal crosses were performed as described previously

(Pagnussat et al., 2005).

Constructions

Genomic DNA was extracted from rosette leaves or whole seedlings as described

(León et al., 2007). To generate *ProBNP:BNP-GFP* translational fusion the region

including *BNP* ORF and putative promoter (672 upstream the ATG codon) was

amplified by PCR using the primer combination listed in Table S1. The amplicon was

cloned using pENTR™ Directional TOPO® Cloning Kit (Invitrogen) using Gateway®

technology and the sequence was verified. The resulting plasmid *pENTR-ProBNP-*

BNP was subjected to the LR reaction using the destination vector pMDC107 (Curtis

and Grossniklaus, 2003). The *ProBNP:GUS* construct was generated amplifying the

BNP putative promoter region (672 upstream the ATG codon) by PCR using the primer

combination listed in Table S1. The amplicon was cloned into pENTR™/TOPO

(Invitrogen) and its sequence was verified. The resulting plasmid *pENTR-ProBNP* was

subjected to the LR reaction using the destination vector pMDC162 (Curtis and

Grossniklaus, 2003). For *Pro35S:BNP-GFP* construct, *BNP* ORF was PCR amplified

using as template clone DKLAT2G44370 obtained from ABRC (Ohio State University,

Ohio, USA) using the primer combination listed in Table S1. The amplicon was cloned

into pENTR™/TOPO (Invitrogen) and its sequence was verified. The resulting plasmid

pENTR-BNP was subjected to the LR reaction using the destination vector pMDC83

(Curtis and Grossniklaus, 2003). For *ProUb10:VOZ1-RFP* and *ProUb10:VOZ2-*
RFP constructions, respective ORFs were PCR amplified using primers listed in Table
S1, cloned in pENTR^TM/TOPO (Invitrogen) and sequence-verified. The resulting
plasmids *pENTR-VOZ1* and *pENTR-VOZ2* were subjected to the LR reaction using the
destination vector pUBN-RFP-DEST (Grefen et al., 2010).
454

Obtaining *bnp/BNP* plants in the *qrt1* background

To obtain *bnp* mutants in *qrt1* background, *bnp-1/BNP* was crossed with *qrt1*
homozygous plants. Kan resistant plants from the progeny were selected and allowed
to self-fertilize. Homozygous *qrt1* plants harboring *bnp* mutations were selected for
further analysis (Johnson-Brousseau and McCormick, 2004).
In a similar way, *qrt1/qrt1-bnp-2/BNP*-VOZ1-RFP plants were obtained crossing *bnp-*
2/BNP plants expressing VOZ1-RFP with *qrt1* homozygous plants.
462

Transformation of *Agrobacterium tumefaciens* and Arabidopsis

464 Vectors were introduced into Agrobacterium strain GV3101 by electroporation.
465 Arabidopsis WT plants were transformed using the floral dip method (Clough and Bent,
466 1998). Transformant plants were selected based on their ability to survive in MS
467 medium with 15 mg. L^{-1} hygromicyn. Resistant (green seedlings with true leaves) were
468 then transferred to soil and grown under the conditions described above.
469

Morphological and histological analyses

Pollen grains viability was assessed using Alexander's staining (Alexander, 1969) and
fluorescein diacetate (FDA) staining (Heslop-Harrison and Heslop-Harrison, 1970). For
pollen development analysis, successive buds in the same inflorescence were
collected (Lalanne and Twell, 2002), fixed in ethanol:acetic acid 3:1 until discoloration
and stored at room temperature. Anthers were washed, dissected and mounted on
microscopy slides with DAPI staining solution. For pollen tube germination, pollen
grains were incubated overnight in a dark moisture chamber at 22°C in 0.01% boric
acid, 5mM CaCl$_2$, 5mM KCl, 1 mM MgSO$_4$, 10% sucrose pH 7.5 and 1.5% agarose, as
described previously (Boavida and McCormick, 2007). For GUS staining, tissues from
ProBNP:GUS carrying plants were collected and incubated in GUS staining solution as
already described (Pagnussat et al., 2007). Gametophytes were observed on a Zeiss
Axio Imager A2 microscope using DIC optics. Seedlings were observed on a Nikon
SMZ800 Stereomicroscope and images were captured with a Canon DS126431
camera.
485

Co-localization experiments in *Nicotiana benthamiana*

A. tumefaciens strain GV3101 carrying *Pro35S:BNP-GFP* was co-infiltrated with *A. tumefaciens* cells carrying either the late endosome marker *RFP-Rha1* or the Golgi apparatus marker *St-RFP*, in leaf epidermal cells of *N. benthamiana* plants. Infiltrated leaves were analyzed 48-72 h after infiltration on a confocal microscope (Nikon Eclipse C1 Plus) using EZ-C1 3.80 imaging software and Ti-Control. Pearson and Spearman correlation (PSC) plugin for ImageJ was used to calculated correlation coefficients from a minimum of 400 individual punctae from at least 10 independent cell images (French et al., 2008).

Yeast two-hybrid screen

The Y2H screening was performed by Hybrigenics Services S.A.S. (Paris, France). A DNA fragment encoding Arabidopsis At2g44370 (amino acids 1-250) was PCR-amplified and cloned into pB66 as C-terminal fusion to Gal4 DNA-binding domain (Gal4-At2g44370) and used as a bait to screen Universal Arabidopsis Normalized library containing 3.2 million of independent clones in pGADT7-RecAB vector. pB66 derive from the original pAS2ΔΔ (Fromont-Racine et al., 1997). For the Gal4 bait construct, 35 million clones were screened using a mating approach with YHGX13 (Y187 ade2-101::loxP-kanMX-loxP, mat□) and L40□Gal4 (mata) yeast strains as previously described (Fromont-Racine et al., 1997). 158 His+ colonies were selected on a medium lacking tryptophan, leucine and histidine. The prey fragments of the positive clones were amplified by PCR and sequenced at their 5' and 3' junctions. The resulting sequences were used to identify the corresponding interacting proteins in the GenBank database (NCBI) using a fully automated procedure. A confidence score (PBS, for Predicted Biological Score) was attributed to each interaction as previously described (Formstecher et al., 2005).

Bimolecular fluorescence complementation (BiFC) analysis

The cDNA sequences of At2g44370 (*BNP*), At1g28520 (*VOZ1*), At2g42400 (*VOZ2*) and At2g17740 (*VLG*) were PCR amplified using the primers listed in Table S1. The PCR products were cloned into pENTR™/TOPO (Invitrogen), sequenced and recombined through BP reaction into BiFC destination plasmids pUBN-YN and pUBN-YC (Grefen et al., 2010). The binary plasmids were then transformed into *A. tumefaciens* strain GV3101 by electroporation. Split nYFP- and cYFP-tagged protein pairs and the gene-silencing suppressor p19 were co-expressed in *N. benthamiana* leaves by *A. tumefaciens* -mediated inoculation. Plant leaves were examined 48-72 h post-infiltration with a confocal microscope (Nikon Eclipse C1 Plus) using a Super Fluor

523 40.0x/1.30/0.22 Oil objective, numerical aperture 1.300. Acquisition was performed
524 using EZ-C1 3.80 imaging software and Ti-Control. All pictures were acquired with the
525 same settings.
526

Quantification of VOZ1-RFP in mature pollen
High quality images of mature pollen grains from the genotypes *qrt1/qrt1, BNP/BNP,*
VOZ1-RFP and *qrt1/qrt1, bnp-1/BNP, VOZ1-RFP* plants were acquired and red
fluorescence was quantified. Red fluorescence background of mature pollen grains
from plants not expressing VOZ1-RFP was subtracted. Fluorescence was expressed in
arbitrary units relative to the calculated cell area of each pollen grain. Data points were
plotted on a violin plot using GraphPad, median (blue line) and quartiles (black lines)
are indicated. Box: Mean values ± SD and number of data points (n) is indicated. A
significant difference of the means was indicated using unpaired t-test.
536

Bioinformatics and phylogenetic analysis
Sequences from the DC1 domain family were identified using the Basic Local
Alignment Search Tool (BLAST) from NCBI (National Center for Biotechnology
Information) using initially BNP as a reference and retrieved sequences in iterative
searches. Protein sequences alignments were performed using MEGA7 (version
7.0.14) (Kumar et al., 2016). Phylogenetic trees were constructed using the neighbor-
joining method and the default settings of MEGA7 (version 7.0.14) (Kumar et al., 2016).
The evolutionary distances were computed using the Poisson correction method
(Zuckerkandl and Pauling, 1965) and are in the units of the number of amino acid
substitutions per site. The trees were drawn using FigTree (version 1.4.3)
(http://tree.bio.ed.ac.uk/). Graphic display of identities was visualized using Geneious
(version 9.1.4) (http://www.geneious.com) based on an identity matrix (Kearse et al.,
2012).
Accession Numbers
Sequence data from this article can be found in the Arabidopsis Genome Initiative
database under accession numbers: BNP (At2g44370), VOZ1 (At1g28520), VOZ2
(At2g42400) and VLG (At2g17740).
554

555 **ACKNOWLEDGEMENTS**
556 We would like to thank Michael Blatt and Christopher Grefen (Glasgow University, UK)
557 for providing pUB-Destination vectors, Jurgen Denecke (University of Leeds, UK) for
558 providing the constructs ST-RFP and Rha-RFP, Nicolas Setzes for assistance with

statistical analysis and Daniela Villamonte for technical assistance with confocal microscopy. This work was supported by grants to DFF from Agencia Nacional de Promoción Cientifica y Técnica Argentina (PICT2013-1524 and PICT2017-0232), Consejo Nacional de Investigaciones Cientificas y Técnicas (CONICET, PIP-0200), Comisión de Investigaciones Cientificas (CIC-PBA) and Universidad Nacional de Mar del Plata; a grant to GCP from the Howard Hughes Medical Institute (IECS award 55007430). LAA and JF are fellows of CONICET; NLA is a fellow of ANPCyT; DFF, GCP, CAC and SD are researchers from CONICET.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

D.F.F. conceived the original screening and research plans; D.F.F., G.C.P. and C.A.C. supervised the experiments; L.A.A., S.D., J.F. and N.L.A. performed most of the experiments; D.F.F., G.C.P and L.A.A. designed the experiments and analyzed the data; D.F.F. conceived the project and wrote the article with contributions of all the authors; D.F.F. agrees to serve as the author responsible for contact and ensures communication

REFERENCES

Alexander MP (1969) Differential staining of aborted and nonaborted pollen. Stain Technol. **44**: 117-122

Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Cheuk R, al e (2003) Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. Science **301**: 653–657

Bhaskar RV, Mohanty B, Verma V, Wijaya E, Kumar PP (2015) A hormone-responsive C1-domain-containing protein At5g17960 mediates stress response in *Arabidopsis thaliana*. PLoS One **10**: e0115418

Boavida LC, McCormick S (2007) Temperature as a determinant factor for increased and reproducible in vitro pollen germination in arabidopsis thaliana. Plant J. **52**: 570-582

Bolte S, Talbot C, Boutte Y, Catrice O, Read ND, Satiat-Jeunemaitre B (2004) FM-dyes as experimental probes for dissecting vesicle trafficking in living plant cells. J. Microsc. **214**: 159-173

Borg M, Brownfield L, Twell D (2009) Male gametophyte development: A molecular perspective. J. Exp. Bot. **60**: 1465-1478

Celesnik H, Ali GS, Robison FM, Reddy AS (2013) *Arabidopsis thaliana* VOZ (vascular plant one-zinc finger) transcription factors are required for proper regulation of flowering time. Biol. Open **2**: 424-431

Clough SJ, Bent AF (1998) Floral dip: A simplified method for Agrobacterium-mediated transformation of *Arabidopsis thaliana*. Plant J. **16**: 735-743

Colon-Gonzalez F, Kazanietz MG (2006) C1 domains exposed: From diacylglycerol binding to protein-protein interactions. Biochim. Biophys. Acta **1761**: 827-837

Curtis MD, Grossniklaus U (2003) A gateway cloning vector set for high-throughput functional analysis of genes in planta. Plant Physiol. **133**: 462–469.

D'Ippólito S, Arias LA, Casalongué CA, Pagnussat GC, Fiol DF (2017) The dc1-domain protein vacuoleless gametophytes is essential for female and male gametophyte development in arabidopsis. Plant J. **90**: 261-275

Ek-Ramos MJ, Avila J, Nelson Dittrich AC, Su D, Gray JW, Devarenne TP (2014) The tomato cell death suppressor ADI3 is restricted to the endosomal system in response to the pseudomonas syringae effector protein avrpto. PLOS One **9**: e110807

Escobar-Sepúlveda HF, Trejo-Téllez LI, Pérez-Rodríguez P, Hidalgo-Contreras JV, Gómez-Merino FC (2017) Diacylglycerol kinases are widespread in higher plants and display inducible gene expression in response to beneficial elements, metal, and metalloid ions. Front. Plant Sci. **8**: 129

Foresti O, Gershlick DC, Bottanelli F, Hummel E, Hawes C, Denecke J (2010) A recycling-defective vacuolar sorting receptor reveals an intermediate compartment situated between prevacuoles and vacuoles in tobacco. Plant Cell **22**: 3992-4008

Formstecher E, Aresta S, Collura V, Hamburger A, Meil A, Trehin A, Reverdy C, Betin V, Maire S, Brun C, Jacq B, Arpin M, Bellaiche Y, Bellusci S, Benaroch P, Bornens M, Chanet R, Chavrier P, Delattre O, Doye V, Fehon R, Faye G, Galli T, Girault JA, Goud B, de Gunzburg J, Johannes L, Junier MP, Mirouse V, Mukherjee A, Papadopoulo D, Perez F, Plessis A, Rosse C, Saule S, Stoppa-Lyonnet D, Vincent A, White M, Legrain P, Wojcik J, Camonis J, Daviet L (2005) Protein interaction mapping: A Drosophila case study. Genome Res. **15**: 376-384

Francis KE, Lam SY, Copenhaver GP (2006) Separation of arabidopsis pollen tetrads is regulated by quartet1, a pectin methylesterase gene. Plant Physiol. **142**: 1004-1013

French AP, Mills S, Swarup R, Bennett MJ, Pridmore TP (2008) Colocalization of
fluorescent markers in confocal microscope images of plant cells. Nature Protoc.
3: 619-628

Fromont-Racine M, Rain JC, Legrain P (1997) Toward a functional analysis of the
yeast genome through exhaustive two-hybrid screens. Nat. Genet. **16**: 277-282

Gao S, Yang L, Zeng HQ, Zhou ZS, Yang ZM, Li H, Sun D, Xie F, Zhang B (2016) A
cotton miRNA is involved in regulation of plant response to salt stress. Sci.
Reports **6**: 19736

Grefen C, Donald N, Hashimoto K, Kudla J, Schumacher K, Blatt MR (2010) A
ubiquitin-10 promoter-based vector set for fluorescent protein tagging facilitates
temporal stability and native protein distribution in transient and stable expression
studies. Plant J. **64**: 355-365

Hafidh S, Fila J, Honys D (2016) Male gametophyte development and function in
angiosperms: A general concept. Plant Reprod. **29**: 31-51

Heslop-Harrison J, Heslop-Harrison Y (1970) Evaluation of pollen viability by
enzymatically induced fluorescence; intracellular hydrolysis of fluorescein
diacetate. Stain Technol. **45**: 115-120

Hwang IS, Choi DS, Kim NH, Kim DS, Hwang BK (2014) The pepper
cysteine/histidine-rich DC1 domain protein CaDC1 binds both RNA and DNA and
is required for plant cell death and defense response. New Phytol. **201**: 518-530

Jensen Michael K, Kjaersgaard T, Nielsen Michael M, Galberg P, Petersen K,
O'Shea C, Skriver K (2010) The *Arabidopsis thaliana* NAC transcription factor
family: Structure–function relationships and determinants of ANAC019 stress
signalling. Biochem J. **426**: 183-196

Johnson-Brousseau SA, McCormick S (2004) A compendium of methods useful for
characterizing Arabidopsis pollen mutants and gametophytically-expressed
genes. Plant J. **39**: 761-775

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S,
Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P,
Drummond A (2012) Geneious basic: An integrated and extendable desktop
software platform for the organization and analysis of sequence data.
Bioinformatics **28**: 1647-1649

Koguchi M, Yamasaki K, Hirano T, Sato MH (2017) Vascular plant one-zinc-finger
protein 2 is localized both to the nucleus and stress granules under heat stress in
arabidopsis. Plant Signal. Behav. **12**: e1295907

Kumar S, Choudhary P, Gupta M, Nath U (2018) Vascular plant one-zinc finger1 (VOZ1) and VOZ2 interact with constans and promote photoperiodic flowering transition. Plant Physiol. **176**: 2917-2930

Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. Mol. Biol. Evol. **33**: 1870-1874

Lalanne E, Twell D (2002) Genetic control of male germ unit organization in arabidopsis. Plant Physiol. **129**: 865-875

León G, Holuigue L, Jordana X (2007) Mitochondrial complex ii is essential for gametophyte development in Arabidopsis. Plant Physiol. **143**: 1534-1546

Li C, Lv J, Zhao X, Ai X, Zhu X, Wang M, Zhao S, Xia G (2010) TaCHP: A wheat zinc finger protein gene down-regulated by abscisic acid and salinity stress plays a positive role in stress tolerance. Plant Physiol. **154**: 211-221

Mimida N, Kidou S, Iwanami H, Moriya S, Abe K, Voogd C, Varkonyi-Gasic E, Kotoda N (2011) Apple Flowering Locus T proteins interact with transcription factors implicated in cell growth and organ development. Tree Physiol. **31**: 555-566

Mitsuda N, Hisabori T, Takeyasu K, Sato MH (2004) VOZ; isolation and characterization of novel vascular plant transcription factors with a one-zinc finger from *Arabidopsis thaliana*. Plant Cell Physiol. **45**: 845-854

Mukherjee S, Tessema M, Wandinger-Ness A (2006) Vesicular trafficking of tyrosine kinase receptors and associated proteins in the regulation of signaling and vascular function. Circul. Res. **98**: 743-756

Nakai Y, Nakahira Y, Sumida H, Takebayashi K, Nagasawa Y, Yamasaki K, Akiyama M, Ohme-Takagi M, Fujiwara S, Shiina T, Mitsuda N, Fukusaki E, Kubo Y, Sato MH (2013) Vascular plant one-zinc-finger protein 1/2 transcription factors regulate abiotic and biotic stress responses in Arabidopsis. Plant J. **73**: 761-775

Neill SO, Gould KS, Kilmartin PA, Mitchell KA, Markham KR (2002) Antioxidant capacities of green and cyanic leaves in the sun species, *Quintinia serrata*. Funct. Plant Biol. **29**: 1437-1443

Pagnussat GC, Yu HJ, Ngo QA, Rajani S, Mayalagu S, Johnson CS, Capron A, Xie LF, Ye D, Sundaresan V (2005) Genetic and molecular identification of genes required for female gametophyte development and function in Arabidopsis. Development **132**: 603-614

Pagnussat GC, Yu HJ, Sundaresan V (2007) Cell-fate switch of synergid to egg cell in Arabidopsis *eostre* mutant embryo sacs arises from misexpression of the BEL1-like homeodomain gene BLH1. Plant Cell **19**: 3578-3592

Palfy M, Remenyi A, Korcsmaros T (2012) Endosomal crosstalk: Meeting points for
signaling pathways. Trends Cell Biol. **22**: 447-456

Pyrzynska B, Pilecka I, Miaczynska M (2009) Endocytic proteins in the regulation of
nuclear signaling, transcription and tumorigenesis. Mol. Oncol. **3**: 321-338

Rahman GM, Das J (2015) Modeling studies on the structural determinants for the
dag/phorbol ester binding to C1 domain. J. Biomol Struct Dynam. **33**: 219-232

Schwarzenbacher RE, Wardell G, Stassen J, Guest E, Zhang P, Luna E, Ton J
(2020) The IBI1 receptor of β-aminobutyric acid interacts with VOZ transcription
factors to regulate abscisic acid signaling and callose-associated defense. Mol.
Plant **13**: 1455-1469

Selote D, Matthiadis A, Gillikin JW, Sato MH, Long TA (2018) The E3 ligase brutus
facilitates degradation of VOZ1/2 transcription factors. Plant Cell Environ. **41**:
2463-2474

Serrano I, Buscaill P, Audran C, Pouzet C, Jauneau A, Rivas S (2016) A non
canonical subtilase attenuates the transcriptional activation of defence responses
in *Arabidopsis thaliana*. eLife **5**:

Shinya T, Galis I, Narisawa T, Sasaki M, Fukuda H, Matsuoka H, Saito M,
Matsuoka K (2007) Comprehensive analysis of glucan elicitor-regulated gene
expression in tobacco BY-2 cells reveals a novel MYB transcription factor
involved in the regulation of phenylpropanoid metabolism. Plant Cell Physiol. **48**:
1404-1413

Song C, Lee J, Kim T, Hong JC, Lim CO (2018) VOZ1, a transcriptional repressor of
DREB2C, mediates heat stress responses in Arabidopsis. Planta **247**: 1439-1448

Suesslin C, Frohnmeyer H (2003) An Arabidopsis mutant defective in UV-B light-
mediated responses. Plant J. **33**: 591-601

Twell D (2011) Male gametogenesis and germline specification in flowering plants.
Sex. Plant Reprod. **24**: 149-160

Wang X, Xu M, Gao C, Zeng Y, Cui Y, Shen W, Jiang L (2020) The roles of
endomembrane trafficking in plant abiotic stress responses. J. Integr. Plant Biol.
62: 55-69

Wang Y-N, Wang H, Yamaguchi H, Lee H-J, Lee H-H, Hung M-C (2010) Copi-
mediated retrograde trafficking from the golgi to the er regulates egfr nuclear
transport. Biochem. Biophys. Res. Comm, **399**: 498-504

Wee EG-T, Sherrier DJ, Prime TA, Dupree P (1998) Targeting of active
sialyltransferase to the plant Golgi apparatus. Plant Cell **10**: 1759-1768

Wu P, Wee P, Jiang J, Chen X, Wang Z (2012) Differential regulation of transcription
factors by location-specific egf receptor signaling via a spatio-temporal interplay
of erk activation. PloS One **7**: e41354

Yasui Y, Kohchi T (2014) Vascular plant one-zinc finger1 and VOZ2 repress the
flowering locus C clade members to control flowering time in Arabidopsis. Biosci.
Biotechnol. Biochem. **78**: 1850-1855

Yasui Y, Mukougawa K, Uemoto M, Yokofuji A, Suzuri R, Nishitani A, Kohchi T
(2012) The phytochrome-interacting vascular plant one-zinc finger1 and voz2
redundantly regulate flowering in arabidopsis. Plant Cell **24**: 3248-3263

Zuckerkandl E, Pauling L (1965) Evolutionary divergence and convergence in
proteins. Evolv. Genes Prot. **97**: 97-166

SUPPORTING INFORMATION

Supplemental Figure S1. *bnp/BNP* hemizygous plants showed normal sporophytic
phenotypes and seed sets*.*

Supplemental Figure S2. *bnp*/*BNP* hemizygous plants showed no differences in the
viability of mature pollen.

Supplemental Figure S3. Complementation of *bnp-1/BNP* plants with *ProBNP:BNP-*
GFP.

Supplemental Figure S4. *BNP* expression in Arabidopsis

Supplemental Figure S5. BNP-related protein VLG does not interacts with VOZ1,
VOZ2 or BNP.

Supplemental Figure S6. Analysis of co-localization of BNP and the endocytic tracer
FM4-64

Supplemental Figure S7. Quantification of VOZ1-RFP red fluorescence in mature
pollen of *qrt1/qrt1, BNP/BNP, VOZ1-RFP* and *qrt1/qrt1, bnp-1/BNP, VOZ1-RFP* plants

Supplemental Figure S8. BPN and VOZ1 localization during pollen development

Supplemental Table S1. Selected BNP interacting protein identified by a yeast two-
hybrid screening.

Supplemental Table S2. Primers used in the study

Figure Legends

Figure 1. BNP codes for a DC1 domain containing protein. (A) Schematic view of a region of Arabidopsis chromosome 2 (from 18,323,538 to 18,321,537 bp -TAIR10) showing BNP genomic location and bnp-1 and bnp-2 insertion sites. (B) BNP amino acid sequence with boxed DC1 domains. (C) Phylogenic tree of 140 DC1 domain containing proteins in Arabidopsis. Bootstrap test results (1000 replicates) of the major nodes are indicated. IDs of sequences grouped in clusters I to VIII are listed. (D) Alignment of BNP with its closest Arabidopsis homologs and CaDC1 (AEI52549), TaCHP (ACU80555), NtDC1a (BAF80452) and NtDC1b (BAF80453). Phylogenic tree with bootstrap test results (1000 replicates) for the alignment of the complete 18 sequences is shown. Amino acid region represented for each sequence is indicated. DC1 domains are framed, signature residues and binding loop regions are indicated (C=Cys, H=His, — = loop). Higher intensity in grey scale denotes higher sequence similarity.

Figure 2. Pollen development and germination is impaired in *bnp-1* mutants. (A) Quantification of phenotypes observed in *BNP/BNP* and *bnp-1/BNP* plants in *qrt1/qrt1* background. (B-E) Nuclear configuration types revealed by DAPI (4',6-Diamidino-2-phenylindole) staining. Bars = 10 µm. (F) Quantification of pollen tube germination in tetrads from *BNP/BNP* and *bnp-1/BNP* plants in *qrt1/qrt1* background. (G-K) Pollen tube germination in tetrads. Bars = 20 µm.

Figure 3. *BNP* is expressed in pollen and young sporophytic tissues. (A-D) Confocal images showing BNP-GFP through microgametogenesis. (E-H) Overlay between DIC and GFP fluorescence. (A and E) Microspore tetrad, (B and F) released microspore, (C and G) bicellular stage and (D and H) tricellular stage. (I) Image showing a projection along the z axis of all planes showing GFP fluorescence. (J) Overlay of DIC and GFP fluorescence showing a germinating pollen grain. (K-S) GUS activity in Arabidopsis transgenic plants carrying the *pBNP:GUS* construct. (K) Seedling 24 h after imbibition, (L) 48 h after imbibition, (M) 6 days post germination and (N) 12 days post germination. (O) Root with lateral root primordium, (P) primary root with root hairs, (Q) third leaf 8 days after imbibition (R) third leaf 14 days after imbibition. (S) GUS signal was detected inside anthers at floral developmental stage 10. Three independent transgenic lines were used for this study. Pictures are representative of the results obtained in all analyzed lines. Bars = 10 µm.

Figure 4. BNP interacts with itself, VOZ1 and VOZ2. Bimolecular fluorescence complementation (BiFC) analysis of the interaction of BNP with itself, VOZ1 and VOZ2 in *Nicotiana benthamiana* leaves. Representative confocal microscopy images showing that BNP interacts with (A–C) itself, (D-F) VOZ1 and (G-I) VOZ2. Reconstituted YFP fluorescence (YFP) or DIC images of *N. benthamiana* leaves are shown in epidermal

cells co-infiltrated with *Agrobacterium tumefaciens* harboring the indicated constructs.
Control *N. benthamiana* leaves transfected with (J-L) BNP-N-YFP, (M-O) BNP-C-YFP,
(P-R) VOZ1-C-YFP, (S-U) VOZ2-C-YFP do not show any BiFC signal. The last row (C,
F, I, L, O, R, U) correspond to the magnification of regions of interest (insets) indicated
in the first row. Bars = 20 μm.

Figure 5. BNP co-localizes with VOZ1 and VOZ2 to Rha1 positive vesicles. (A, F,
K and P) Localization of BNP-GFP transiently expressed in *Nicotiana benthamiana*
epidermal cells. (B) Fluorescent marker for late endosomes/prevacuolar compartments/
multivesicular bodies, RFP-Rha1. (C) Merged image of BNP-GFP and RFP-Rha1.
(D) Shows an overlay with the corresponding bright-field image. (E, J, O and T)
Pearson and Spearman correlation (PSC) test calculated using a plugin for ImageJ; r
values indicate the level of co-localization ranging from +1 for perfect co-localization to
−1 for negative correlation. (G) Fluorescent marker for Golgi apparatus, St-RFP. (H)
Merged image of BNP-GFP and St-RFP. (I) Shows an overlay with the
corresponding bright-field image. (L) Localization of RFP-VOZ1. (M) Merged image of
BNP-GFP and RFP-VOZ1. (N) Shows the corresponding bright field image of the cells
showed in (M). (Q) Localization of RFP-VOZ2. (R) Merged image of BNP-GFP and
RFP-VOZ2. (S) Bright field image of the cells showed in (R). Bars = 20 μm.

Figure 6. VOZ1 localization in mature pollen from WT and *bnp-1/BNP* plants.
Mature pollen from (A- E) BNP/BNP, VOZ1-RFP and (F – J) bnp-1/BNP, VOZ1-RFP
plants was analized. Representative z-stack images of confocal micrographs displaying
(A, F) DAPI nuclear staining, (B, G) VOZ1-RFP fluorescence and (E, J) bright-field
images. (C, H) Merged DAPI and VOZ1-RFP fluorescence images and (D, I) merged
DAPI, VOZ1-RFP fluorescence and bright-field images Yellow arrows indicate the
vegetative nucleus and white arrows indicate the generative nucleus. Bars = 10 μm.

Figure 7. Nuclear localization of VOZ1 is affected in *bnp* mutants.
Pollen tetrads from *qrt1/qrt1, BNP/BNP, VOZ1-RFP* (A, B, E, F, I, J) and *qrt1/qrt1, bnp-*
1/BNP, VOZ1-RFP (C, D, G, H, K, L) plants were analyzed at microspore (A-D),
binucleate (E-H) and mature tricellular (I-L) stages. Representative z-stack images of
confocal micrographs displaying VOZ1-RFP fluorescence (B, D, F, H, J, L) and bright
field images (A, C, E, G, I, K) are shown. Arrowheads indicate nuclear localized VOZ1.
Bars = 10 μm.
Mature pollen tetrads from four *qrt1/qrt1, BNP/BNP, VOZ1-RFP* and three *qrt1/qrt1,*
bnp-1/BNP, VOZ1-RFP plants were analyzed by fluorescent microscopy and VOZ1
localization (M) and nuclear configuration (N) were scored. Observed number and total
pollen tetrads are indicated in parenthesis.

Table 1. Transmission efficiency of *bnp* alleles. Transmission efficiency of the *bnp-1* and *bnp-2* alleles in reciprocal crosses between mutant and WT plants.

Parental genotypes		Genotype of progeny† Number of individuals		Transmission efficiency of:	p-value*
Female	Male	WT	*bnp-1/BNP*	*bnp-1*	
bnp-1/BNP	WT	98	89	♀ 90.8%	0.51
WT	*bnp-1/BNP*	181	28	♂ 15.5%	<0.0001
		WT	*bnp-2/ BNP*	*bnp-2*	
bnp-2/BNP	WT	217	208	♀ 95.9%	0.66
WT	*bnp-2/BNP*	254	56	♂ 22%	<0.0001

†No homozygous plants were detected; * Chi-square test for a 1:1 segregation hypothesis (degrees of freedom = 1).

Table 2. Viability of the progeny in *bnp* mutants. Seed germination and seedling lethality of the progeny of self-pollinated *bnp-1/BNP*, *bnp-2/BNP* and WT plants.

Parental genotype	% viable progeny	% unviable progeny		p-value*
		not germinated	seedling lethality	
WT	98.86 (435/440)	0.23 (435/440)	1.13 (435/440)	
bnp-1/BNP	94.83 (715/754)	1.72 (13/754)	3.45 (26/754)	<0.001
bnp-2/BNP	93.32 (908/753)	2.15 (21/753)	4.52 (44/753)	<0.00001

In parenthesis, number of individuals observed over the total analyzed. * Chi-square test for a hypothesis of no relationship between the genotype of the parents and the viability of the progeny (degree of freedom = 1).

Figure 1. BNP codes for a DC1 domain containing protein. (A) Schematic view of a region of Arabidopsis chromosome 2 (from 18,323,538 to 18,321,537 bp -TAIR10) showing BNP genomic location and bnp-1 and bnp-2 insertion sites. (B) BNP amino acid sequence with boxed DC1 domains. (C) Phylogenic tree of 140 DC1 domain containing proteins in Arabidopsis. Bootstrap test results (1000 replicates) of the major nodes are indicated. IDs of sequences grouped in clusters I to VIII are listed. (D) Alignment of BNP with its closest Arabidopsis homologs and CaDC1 (AEI52549), TaCHP (ACU80555), NtDC1a (BAF80452) and NtDC1b (BAF80453). Phylogenic tree with bootstrap test results (1000 replicates) for the alignment of the complete 18 sequences is shown. Amino acid region represented for each sequence is indicated. DC1 domains are framed, signature residues and binding loop regions are indicated (C=Cys, H=His, — = loop). Higher intensity in grey scale denotes higher sequence similarity.

A

Genotype	Observed nuclear phenotype			
	Four trinucleate (image B)	Three trinucleate, one binucleate (image C)	Two trinucleate, two binucleate (image D)	Two trinucleate, one tetranucleate (image E)
BNP/BNP	96.9 % (187/193)	3.1 % (6/193)	0 % (0/193)	0 % (0/193)
bnp-1/BNP	0.9 % (2/229)	11.8 % (27/229)	83.4 % (191/229)	3.9 % (9/229)

F

Genotype	Pollen germination rate	Observed tetrad phenotype				
		Four germinated (image G)	Three germinated (image H)	Two germinated (image I)	One germinated (image J)	none germinated (image K)
BNP/BNP	68.2 % (573/840)	35.2 % (74/210)	35.6 % (75/210)	10.5 % (22/210)	5.7 % (12/210)	12.8 % (27/210)
bnp-1/BNP	37.7 % (291/772)	0% (0/193)	8.8 % (17/193)	46.6% (90/193)	31.1 % (60/193)	13.4 % (26/193)

Figure 2. Pollen development and germination is impaired in bnp-1 mutants. (A) Quantification of phenotypes observed in BNP/BNP and bnp-1/BNP plants in qrt1/qrt1 background. (B-E) Nuclear configuration types revealed by DAPI (4',6-Diamidino-2-phenylindole) staining. Bars = 10 µm. (F) Quantification of pollen tube germination in tetrads from BNP/BNP and bnp-1/BNP plants in qrt1/qrt1 background. (G-K) Pollen tube germination in tetrads. Bars = 20 µm

Figure 3. BNP is expressed in pollen and young sporophytic tissues. (A-D) Confocal images showing BNP-GFP through microgametogenesis. (E-H) Overlay between DIC and GFP fluorescence. (A and E) Microspore tetrad, (B and F) released microspore, (C and G) bicellular stage and (D and H) tricellular stage. (I) Image showing a projection along the z axis of all planes showing GFP fluorescence. (J) Overlay of DIC and GFP fluorescence showing a germinating pollen grain. (K-S) GUS activity in Arabidopsis transgenic plants carrying the pBNP:GUS construct. (K) Seedling 24 h after imbibition, (L) 48 h after imbibition, (M) 6 days post germination and (N) 12 days post germination. (O) Root with lateral root primordium, (P) primary root with root hairs, (Q) third leaf 8 days after imbibition (R) third leaf 14 days after imbibition. (S) GUS signal was detected inside anthers at floral developmental stage 10. Three independent transgenic lines were used for this study. Pictures are representative of the results obtained in all analyzed lines. Bars = 10 µm.

Figure 4. BNP interacts with itself, VOZ1 and VOZ2. Bimolecular fluorescence complementation (BiFC) analysis of the interaction of BNP with itself, VOZ1 and VOZ2 in Nicotiana benthamiana leaves. Representative confocal microscopy images showing that BNP interacts with (A–C) itself, (D-F) VOZ1 and (G-I) VOZ2. Reconstituted YFP fluorescence (YFP) or DIC images of N. benthamiana leaves are shown in epidermal cells co-infiltrated with Agrobacterium tumefaciens harboring the indicated constructs. Control N. benthamiana leaves transfected with (J-L) BNP-N-YFP, (M-O) BNP-C-YFP, (P-R) VOZ1-C-YFP, (S-U) VOZ2-C-YFP do not show any BiFC signal. The last row (C, F, I, L, O, R, U) correspond to the magnification of regions of interest (insets) indicated in the first row. Bars = 20 µm.

Figure 5. BNP co-localizes with VOZ1 and VOZ2 to Rha1 positive vesicles. (A, F, K and P) Localization of BNP-GFP transiently expressed in Nicotiana benthamiana epidermal cells. (B) Fluorescent marker for late endosomes/prevacuolar compartments/ multivesicular bodies, RFP-Rha1. (C) Merged image of BNP-GFP and RFP-Rha1. (D) Shows an overlay with the corresponding bright-field image. (E, J, O and T) Pearson and Spearman correlation (PSC) test calculated using a plugin for ImageJ; r values indicate the level of co-localization ranging from +1 for perfect co-localization to −1 for negative correlation. (G) Fluorescent marker for Golgi apparatus, St-RFP. (H) Merged image of BNP-GFP and St-RFP. (I) Shows an overlay with the corresponding bright-field image. (L) Localization of RFP-VOZ1. (M) Merged image of BNP-GFP and RFP-VOZ1. (N) Shows the corresponding bright field image of the cells showed in (M). (Q) Localization of RFP-VOZ2. (R) Merged image of BNP-GFP and RFP-VOZ2. (S) Bright field image of the cells showed in (R). Bars = 20 μm.

Figure 6. VOZ1 localization in mature pollen from WT and bnp-1/BNP plants.
Mature pollen from (A- E) BNP/BNP, VOZ1-RFP and (F – J) bnp-1/BNP, VOZ1-RFP
plants was analized. Representative z-stack images of confocal micrographs displaying
(A, F) DAPI nuclear staining, (B, G) VOZ1-RFP fluorescence and (E, J) bright-field
images. (C, H) Merged DAPI and VOZ1-RFP fluorescence images and (D, I) merged
DAPI, VOZ1-RFP fluorescence and bright-field images Yellow arrows indicate the
vegetative nucleus and white arrows indicate the generative nucleus. Bars = 10 µm.

M

Genotype	Observed VOZ1 localization				
	Four nuclear	Three nuclear, one cytoplasmic	Two nuclear, two cytoplasmic	One nuclear, three cytoplasmic	Four cytoplasmic
qrt1/qrt1, BNP/BNP, VOZ1-RFP	91.5 % (65/71)	7.0 % (5/71)	1.4 % (1/71)	0 % (0/71)	0 % (0/71)
qrt1/qrt1, bnp-1/BNP, VOZ1-RFP	1.3 % (1/76)	11.8 % (9/76)	57.8% (44/76)	27.6% (21/76)	1.3 % (1/76)

N

Genotype	Observed nuclear phenotype			
	Four trinucleate	Three trinucleate, one binucleate	Two trinucleate, two binucleate	Two binucleate, one tetranucleate
qrt1/qrt1, BNP/BNP, VOZ1-RFP	97.3 % (71/73)	2.7 % (2/73)	0 % (0/73)	0 % (0/73)
qrt1/qrt1, bnp-1/BNP, VOZ1-RFP	3.6 % (4/110)	18.2 % (20/110)	77.3 % (85/110)	0.9 % (1/110)

Figure 7. Nuclear localization of VOZ1 is affected in bnp mutants.

Pollen tetrads from qrt1/qrt1, BNP/BNP, VOZ1-RFP (A, B, E, F, I, J) and qrt1/qrt1, bnp-1/BNP, VOZ1-RFP (C, D, G, H, K, L) plants were analyzed at microspore (A-D), binucleate (E-H) and mature tricellular (I-L) stages. Representative z-stack images of confocal micrographs displaying VOZ1-RFP fluorescence (B, D, F, H, J, L) and bright field images (A, C, E, G, I, K) are shown. Arrowheads indicate nuclear localized VOZ1. Bars = 10 µm.

Mature pollen tetrads from four qrt1/qrt1, BNP/BNP, VOZ1-RFP and three qrt1/qrt1, bnp-1/BNP, VOZ1-RFP plants were analyzed by fluorescent microscopy and VOZ1 localization (M) and nuclear configuration (N) were scored. Observed number and total pollen tetrads are indicated in parenthesis.