High neutralizing antibody levels against SARS-CoV-2 Omicron BA.1 and BA.2 after UB-612 vaccine booster

Farshad Guirakhoo[1]*, Shixia Wang[1], Chang Yi Wang[2], Hui-Kai Kuo[2], Wen-Jiun Peng[2],

Hope Liu[2], Lixia Wang[1], Marina Johnson[3], Adam Hunt[3], Mei Mei Hu[1], Thomas P. Monath[1],

Alexander Rumyantsev[1], David Goldblatt[3]

[1]Vaxxinity Inc., Dallas, TX, USA.

[2]United Biomedical Inc Asia; Taipei, Taiwan.

[3]Great Ormond Street Institute of Child Health, University College London; London, UK.

*Corresponding author:

Farshad Guirakhoo, PhD

Phone: +41 076-476-2153

Email: farshad@vaxxinity.com

Short title: UB-612 boosted neutralizing antibodies against Omicron

Abstract

The highly transmissible Omicron variant has caused high rates of breakthrough infections among vaccinated and convalescent individuals. Here, we demonstrate that a booster dose of UB-612 vaccine candidate delivered 7-9 months after primary vaccination increases neutralizing antibody levels by 131-, 61- and 49-fold against ancestral SARS-CoV-2, Omicron BA.1 and BA.2 variants, respectively. Based on the RBD protein binding antibody responses, we estimated a ~95% efficacy against symptomatic COVID-19 caused by the ancestral strain after a UB-612 booster. Our results support UB-612 vaccine as a potent booster against current and emerging SARS-CoV-2 variants.

Keywords: SARS-CoV-2; COVID-19; Omicron; variant; vaccine; booster; spike protein; receptor binding domain; antibody; neutralizing antibody.

Introduction

In November 2021, the Omicron (B.1.1.529) Variant of Concern (VOC) was first reported in South Africa and quickly spread, becoming the dominant SARS-CoV-2 variant worldwide. Omicron's high transmissibility and potential for immune system evasion, as suggested by its ability to infect and be transmitted by previously infected and vaccinated individuals, predicted a transmission advantage over the Delta variant and the displacement of the latter as dominant variant [1]. The Omicron variant has 3 major sublineages: BA.1, BA.2, and BA.3. While BA.1 caused most of the cases globally throughout November 2021 and February 2022, BA.2 is now the predominant SARS-CoV-2 variant[2].

The Omicron variant has over 50 new amino acid substitutions with >15 in the receptor-binding domain (RBD) of Spike (S) protein [3]. Although BA.2 shares many mutations with BA.1, these 2 sublineages differ by dozens of amino acid substitutions, especially at the S protein (**Fig. 1A**), some of which could be responsible for the rapid surge in BA.2 cases.

Given that over 90% of neutralizing antibodies are present in plasma of convalescent individuals and up to 99% of neutralizing antibodies elicited by vaccination with the mRNA-1273 vaccine are directed to RBD [4], these mutations could be largely responsible for Omicron's ability to evade neutralizing antibodies induced by the approved COVID-19 vaccines [5]. Multiple studies have shown a 20- to 30-fold reduction in neutralizing antibody activity against Omicron in the sera of primary vaccine recipients, compared with the ancestral SARS-CoV-2 or D614G viruses [5–7]. The emergence of new variants, in addition to rapidly waning immunity over time, has raised concerns about breakthrough infections in vaccinated individuals, and highlights the need for booster doses. Homologous or heterologous booster vaccines, all based on full-length S protein, restored protective neutralizing antibodies to levels achieved by

the primary immunization; however, these titers were 7.1-fold lower against Omicron BA.1 than

the ancestral strain, suggesting a continued risk of breakthrough infections in vaccinated

individuals over time [5].

In contrast to most of approved COVID-19 vaccines, which encode the full-length S

protein, the UB-612 vaccine candidate is composed of Wuhan-Hu S1-RBD-sFc fusion protein

and is enriched with 5 peptides representing Sarbecovirus-conserved Th and CTL epitopes on the

S2 subunit, Membrane (M), and Nucleocapsid (N) proteins [8]. A favorable safety and

tolerability profile for UB-612 was demonstrated in ~4000 participants in a Phase 1 trial and a

Phase 2 trial conducted in Taiwan [9]. In both trials, the UB-612 vaccine was found to have a

favorable safety profile and low reactogenicity. Two immunizations with UB-612 led to a >90%

seroconversion rate of neutralizing antibodies in vaccine recipients. In these same studies, UB-

612 was shown to elicit long-lasting neutralizing antibody titers similar to levels detected in

convalescent patients [10], that were cross-reactive against Delta and Omicron variants [9].

The objectives of this study were to evaluate the magnitude of neutralizing antibodies

elicited by a third dose (booster) of UB-612 against Omicron and their reactivity to recombinant

S and RBD protein antigens across various variants.

Methods

93   After receiving a 2-dose primary vaccine series or a booster given 7-9 months after the

second dose, sera from 15 participants (who consented to be in this study) in the Phase 1 trial

(UB-612, 100-µg dose), were tested in a SARS-CoV-2 live virus neutralization test (VNT) at

Vismederi, Siena, Italy (a Coalition for Epidemic Preparedness Innovations (CEPI) testing

laboratory for COVID-19 vaccines) **(Table S1)**. Previously, to establish an International

Reference Standard for anti−SARS-CoV-2 antibody detection, the VNT used in our analysis and

performed by Vismederi, was compared with other VNTs and found to be the most stringent

assay, resulting in a lower geometric mean titer (GMT) than other plaque reduction-, foci

reduction-, cytopathic effect (CPE)-, or pseudotyped virus-based neutralization assays [11]. The

detailed materials and methods for clinical trial and sample information, neutralization assays,

immunological assays and statistical analyses are described in the Supplemental Materials.

 Results

 A booster dose of UB-612 significantly increased the levels of neutralizing antibodies

 against the ancestral strain and Omicron variants

Two doses of UB-612 showed modest neutralizing activity against the authentic wild-

type SARS-CoV-2 ancestral strain (Victoria/1/2020) (GMT VNT_{50} of 47.0), and very low

activity against Omicron's BA.1 and BA.2 sublineages (GMT VNT_{50} of 10-11) (**Fig. 1B**) (n=15).

Similarly, a 2-dose immunization with mRNA vaccines has been shown to result in low levels of

Omicron neutralizing antibody responses: (i) mRNA-1273 on day 21 after immunization

stimulated a GMT $pVNT_{50}$ of 14, and (ii) BNT162b2 on day 28 after immunization lead to a

GMT $pVNT_{50}$ of 7 [6,7].

A booster dose of UB-612 delivered at 7-9 months after the primary series increased

neutralizing antibody titers against the ancestral strain, as well as Omicron BA.1 and BA.2

variants, to GMT VNT_{50} of 6,159, 670, and 485, respectively, which constitutes 131-, 61-, and

49-fold higher GMTs than those achieved after 2 doses (**Fig. 1B**). The estimated decrease in

neutralization titers against Omicron BA.1 and BA.2 in 15 sera obtained 2 weeks after UB-612

booster, was 9.2- and 12.7-fold, respectively, compared with the ancestral Victoria strain.

Previously, only a 5.5-fold decrease against BA.1 was reported when 20 sera from UB-612

vaccinated participants were tested in a pseudovirus-based neutralization assay (GMT $pVNT_{50}$ of

12,778 against the Wuhan strain, vs. 2,325 against the BA.1 strain) [9]. These data support the

breadth of UB-612−elicited neutralizing antibodies across multiple SARS-CoV-2 variants after

the booster, a differentiation property of UB-612 primarily attributed to its subunit protein RBD

antigenic component [10].

Expansion of antibody breadth following UB-612 booster

We evaluated reactivity of UB-612−elicited antibodies to S and RBD proteins, using 15 sera from the Phase 1 trial (V123) participants and 84 randomly selected sera from Phase 2 trial participants immunized with UB-612 (**Tables S1-S2**). These sera were tested in 2 ELISA-based assays for IgG direct binding to recombinant S and RBD proteins, and inhibition of recombinant S and RBD protein binding to the human angiotensin-converting enzyme 2 (hACE2) receptor. A third dose of UB-612 booster immunization stimulated broadly reactive IgG antibodies, effectively binding to RBDs of 14 divergent SARS-CoV-2 variants, including Alpha, Beta, Gamma, Delta, and Omicron (**Fig. 1C** and **Fig. S1**).

Compared with the second UB-612 dose, IgG binding titers against Omicron's RBD after the third dose increased by over 40-fold, and the titers against RBDs of other SARS-CoV-2 variants were also increased 30- to 50-fold after the booster dose. When the IgG titer ratio (in binding antibody units (BAU)/mL) of several variants was compared to the ancestral Wuhan strain, the normalized RBD antibody-binding responses to the tested variants were found to be similar after 2 or 3 doses: Alpha (0.98-fold), Beta (2.44-fold), Delta (1.33-fold), Gamma (1.77-fold), and Omicron (3.3-fold) after 2 doses; and Alpha (0.91-fold), Beta (1.8-fold), Delta (1.4-fold), Gamma (1.55-fold), and Omicron (3.7-fold) after 3 doses.

Similar to RBD binding, the results of the S-protein binding antibody responses (S:ACE2- and RBD:ACE2-blocking antibody titers), confirmed the extent of stability in ratios of parental to variant IgG antibodies stimulated by 2 or 3 doses of UB-612, despite an up to 60-fold increase in titers against different variants after the booster dose (**Fig. S2, Fig. S3**).

Vaccine efficacy prediction

We compared the level of UB-612−elicited IgG antibodies with data previously reported

for several authorized vaccines determined in equivalent S- and RBD-binding assays [12]. After

a 2-dose primary immunization series, the GMTs of UB-612−elicited IgG antibodies were 69

and 127 (BAU/mL) against the Wuhan S protein, and 235 and 494 (BAU/mL) against the RBD

antigen in sera from Phase 1 and Phase 2 participants, respectively (**Fig. S4**). These IgG

responses were comparable to those observed in individuals after the primary immunization with

adenovirus vectored vaccines (1-dose Ad26.COV2.S or 2-dose ChAdOx1-S) but were lower than

the response observed after 2 immunizations with mRNA vaccines. The additional booster dose

with UB-612 increased levels of both S- and RBD-protein binding IgG antibodies in the Phase 1

participants by more than 16- and 13-fold, and increased antibody GMTs to 2,138 and 6,767

(BAU/mL), respectively, matching those achieved by 2 immunizations with the mRNA vaccines.

We further utilized a vaccine efficacy prediction model based on the RBD activity of IgG

antibodies to the ancestral strain, extending previous models based on neutralizing antibodies

[13] or S protein−binding activities [12]. According to this model, the predicted vaccine efficacy

of UB-612 against symptomatic disease caused by the prototype strain after 2 doses is ~72%

(235 BAU/mL) with sera from 15 Phase 1 participants, ~82% (494 BAU/mL) with sera from 84

randomly selected Phase 2 participants, and ~95% after the booster dose (6,767 BAU/mL), with

sera from 15 Phase 1 participants (**Fig. 2 and Fig. S4**).

Discussions

It is difficult to compare neutralization activities of different vaccine platforms against

variants because there are currently no international standard reagents available for variants, and

each manufacturer uses a different assay, including live virus, pseudovirus-based, or chimeric

recombinant virus, with different endpoint readouts, such as colorimetric or CPE-based

evaluations. Moreover, each stock virus may contain different ratios of noninfectious to

infectious particles and different virus concentrations may be used for neutralizing sera from

vaccinees, both of which could influence the resulting titers. Keeping these caveats in mind, we

observed GMT levels after the UB-612 booster dose comparable to those elicited by a booster

dose of mRNA vaccine BNT162b2 [7] and mRNA-1273-50 μg [6]. After the booster dose (third

vaccination), the live virus neutralizing antibody GMTs against the ancestral strain and Omicron

BA.1 were 763 and 106 for BNT162b2 (7.2-fold loss) [7], and 2,423 and 850 for mRNA-1273-

50 μg (2.8-fold loss) [6]. The pseudovirus neutralization GMT titers were 6,539, 1,066, and 776

against the ancestral strain WA1/2020, Omicron BA.1, and BA.2, respectively, at 14 days after

the third dose of BNT162b2 (a 6.1- and 8.4-fold loss against BA.1 and BA.2, respectively,

compared with the ancestral strain). The homologous booster of mRNA-1273, BNT162b2, or

Ad26.COV.S S-based vaccines dramatically increased neutralizing antibodies to Omicron (20- to

30-fold) compared with the modest increase reported for the ancestral strain (1- to 4-fold), likely

due to a higher baseline titer for the ancestral strain compared with the variants.

Vaccination with UB-612 elicited highly cross-reactive IgG and neutralizing antibodies

to Omicron variants (with a 49- to 61-fold increase in VNT_{50}) and the ratio of binding antibodies

to ancestral strain/Omicron and other variants remained stable after the second and booster

immunizations. It has been demonstrated that the third dose of mRNA vaccines containing the

full-length S protein of SARS-CoV-2 can recall the persisting memory B cells clones that have

been affinity-matured after a long interval between the doses, as well as to develop new clones of

memory B cells to produce neutralizing antibodies to conserved RBD regions, enhancing the

breadth of cross-variant neutralization [14]. We believe that the UB-612 vaccine, which contains

only the RBD region of the S protein, may be more effective in recalling persisting memory B-

cell clones against the conserved epitopes to protect against variant-induced serious disease.

Another potential advantage of the UB-612 vaccine is the inclusion of Th/CTL peptides from S2,

M and N proteins that are highly conserved in Omicron variants (data not shown). There is

growing evidence supportive of importance of vaccine- elicited CD8+ T cells in protection

against Omicron [15].

In summary, a booster dose of UB-612 elicited robust S- and RBD-specific binding and

virus neutralizing antibodies against multiple SARS-CoV-2 variants, including Omicron BA.1

and BA.2. The magnitude and extent of reactivity of the neutralizing antibody responses after

UB-612 booster match those reported for the authorized vaccines, including BNT162b2 and

mRNA-1273. Additionally, UB-612 has been shown to stimulate T-cell responses against

conserved S2, N, and M peptides [9,10] and may provide long-lasting antibody responses [11]

that would further differentiate UB-612 from many authorized vaccines. As SARS-CoV-2

continues to evolve, several strategies are being explored to effectively prevent COVID-19

caused by newly emerging SARS-CoV-2 variants, including monovalent variant antigen

matching, multivalent, or universal vaccine approaches. Our results indicate that UB-612 could

offer an alternative strategy for a booster vaccine, eliciting long lasting neutralizing antibody

responses with extensive activity across currently circulating and potentially future SARS-CoV-2

variants.

Funding

This work was supported by Vaxxinity Inc.

Acknowledgments

We thank many colleagues at UBIAsia, UBP, and Vaxxinity who designed, developed, and produced the UB-612 vaccine candidate and performed the Phase 1 and 2 clinical trials in Taiwan. We also thank the Phase 1 (NCT04545749) and Phase 2 (NCT04773067) clinical trial participants, from whom the post-vaccination sera were obtained. We are also grateful to the Vismederi team for their work on live virus-neutralizing antibody assays.

Conflict of Interests

FG, SW, LW, MMH, TM, and AR are employees of Vaxxinity Inc, Dallas, TX, USA.

CYW, WJP, HKK, and HL are employees of United Biomedical Inc Asia, Taipei, Taiwan.

Author contributions

Conceived and conceptualized, wrote the manuscript: FG

Analyzed data, interpreted data, wrote the manuscript: SW

Received consent from the patients for the samples to be used for this study and organized shipment of their sera for binding and neutralization assays and reviewed the manuscript: CYW, WJP, HL, HKK

Assisted in planning of the study: MMH, TPM

Wrote and reviewed the manuscript: AR

Analyzed data: LW

Performed experiments and interpreted data: DG, MG, AH

Previous presentation of data

Part of data was previously presented at the World Vaccine Congress, April 17-20, 2022; and

was published in BioRxiv.

References

1. Viana R, Moyo S, Amoako DG, et al. Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in southern Africa. Nature **2022**; 603:679-686.

2. World Health Organization. Weekly epidemiological update on COVID-19 - 22 March 2022. Available from: https://www.who.int/publications/m/item/weekly-epidemiological-update-on-covid-19---22-march-2022. Accessed 24 April 2022.

3. Rahimi F, Abadi ATB. Is Omicron the last SARS-CoV-2 Variant of Concern? Arch Med Res **2022**; 53:336-338

4. Kleanthous H, Silverman JM, Makar KW, Yoon I-K, Jackson N, Vaughn DW. Scientific rationale for developing potent RBD-based vaccines targeting COVID-19. npj vaccines **2021**; 6:128

5. Pérez-Then E, Lucas C, Monteiro VS, et al. Neutralizing antibodies against the SARS-CoV-2 Delta and Omicron variants following heterologous CoronaVac plus BNT162b2 booster vaccination. Nat Med **2022**; 28:481–485

6. Pajon R, Doria-Rose NA, Shen X, et al. SARS-CoV-2 Omicron Variant Neutralization after mRNA-1273 Booster Vaccination. N Engl J Med **2022**; 386:1088-1091

7. Muik A, Lui BG, Wallisch A-K, et al. Neutralization of SARS-CoV-2 Omicron by BNT162b2 mRNA vaccine-elicited human sera. Science **2022**; 375;678-680

8. Guirakhoo F, Kuo L, Peng J, et al. A Novel SARS-CoV-2 Multitope Protein/Peptide Vaccine Candidate is Highly Immunogenic and Prevents Lung Infection in an AAV hACE2 Mouse Model and non-human primates. BioRxiv 399154 [Preprint] October 7, 2021 [cited Feb 13, 2022]; Available from: https://doi.org/10.1101/2020.11.30.399154

271   9.   Wang CY, Hwang K-P, Kuo H-K, et al. A multitope SARS-COV-2 vaccine provides long-

272   lasting B cell and T cell immunity against Delta and Omicron variants. J Clin Invest

273   [Online ahead of print] **2022**; Available from: https://www.jci.org/articles/view/157707

274   10.   Yi C, Hwang K-P, Kuo H-K, et al. Potent Anti-Delta Effect By a Booster Third-Dose of

275   UB-612, a Precision-Designed SARS-CoV-2 Multitope Protein-Peptide Research Square

276   [Preprint] October 18, 2021 [cited Feb 14, 2022]; Available from:

277   https://assets.researchsquare.com/files/rs-944205/v1/520a1fa9-da02-4683-bd0e-

278   a7068f0bf05c.pdf?c=1636030266

279   11.   Mattiuzzo G, Bentley EM, Hassall M, et al. Establishment of the WHO International

280   Standard and Reference Panel for anti-SARS-CoV-2 antibody on behalf of the ISARIC4C

281   Investigators. World Health Organization Expert Committee on Biological Standardization

282   **2020** [Online] Nov 18, 2020 [cited Feb 14, 2022] Available from

283   https://www.who.int/publications/m/item/WHO-BS-2020.2403

284   12.   Goldblatt D, Fiore-Gartland A, Johnson M, et al. Towards a population-based threshold of

285   protection for COVID-19 vaccines. Vaccine **2022**; 40:306–315.

286   13.   Cromer D, Steain M, Reynaldi A, et al. Neutralising antibody titres as predictors of

287   protection against SARS-CoV-2 variants and the impact of boosting: a meta-analysis. The

288   Lancet Microbe **2022**; 3:e52–e61.

289   14.   Muecksch F, Wang Z, Cho A, et al. Increased memory B cell potency and breadth after a

290   SARS-CoV-2 mRNA boost. Nature 2022; in press (https://doi.org/10.1038/s41586-022-

291   04778-y)

292   15.   Chandrashekar A, Yu J, McMahan K, et al. Vaccine protection against the SARS-CoV-2

293   Omicron variant in macaques. Cell 2022; 185:1549-1555.e11

Figure 1. SARS-CoV-2 Omicron BA.1 and BA.2 amino acid substitutions and UB-612

vaccine-induced antibody responses. Panel **A** shows the amino acid substitutions in Omicron's

BA.1 and BA.2 sublineage spike protein. The upper part is the S protein diagram, and the lower

part shows the substitutions. The ".", "-" and "+" represent sequence identical, deletion, and

insertion in Omicron BA.1 and BA.2 compared with the US-WA1/2020 virus, respectively.

Panel **B** shows the GMT VNT$_{50}$ neutralizing antibody titers against SARS-CoV-2 ancestral strain

Victoria/1/2020 (VIC01/2020) and Omicron (B.1.1.529) variant sublineages BA.1 and BA.2 in

sera from Phase 1 trial (V123) participants (n=15). The sera were collected at 28 days after 2

doses and at 14 days after the booster dose with UB-612 (100 µg). Data expressed in the

reciprocal dilutions for each serum sample and GMT (95% CI) are plotted. Panel C shows the

RBD-specific IgG binding titers against SARS-CoV-2 major VOCs in sera collected 28 days

after 2 doses and 14 days after 3 doses with UB-612 (100 µg) from Phase 1 trial participants

(n=15). The loss of antibody binding to the RBD of variants compared with the original RBD

(ancestral strain) remains stable between 2 and 3 doses of UB-612 vaccine, despite an increase in

310   levels of binding antibodies to RBD. GMT, geometric mean titers; VNT, virus neutralization

311   test.

Figure 2. Estimated UB-612 efficacy after 2 and 3 doses of UB-612 vaccine.

A model bridging vaccine-induced RBD IgG response to vaccine efficacy against symptomatic

COVID-19 caused by ancestral Wuhan [5]. Estimated efficacy of UB-612 after 2 doses is ~72%

(CI, 70%-80%) based on RBD binding IgG antibodies from 15 participants (Phase 1) (GMT 235

BAU/mL, 95% CI, 158-350, ~82% (CI, 80%-85%) based on RBD binding IgG antibodies (GMT

494 BAU/mL, 95% CI, 337-725, shown in this graph), and ~95% (93%-97%) after a booster

vaccination (GMT 6767 (95% CI, 4142-11,057). GMT, geometric mean titers.