Hemispheric lateralization in older adults who habitually play darts: a cross-sectional study using functional near-infrared spectroscopy.

Koki Toyofuku, ktoyofuku@mis.doshisha.ac.jp, Graduate School of Life and Medical Sciences, Doshisha University, Kyoto, Japan.

Satoru Hiwa[*]**,** shiwa@mail.doshisha.ac.jp, Department of Biomedical Sciences and Informatics, Doshisha University, Kyoto, Japan. ORCID: 0000-0001-6547-626X.

Kensuke Tanioka, ktanioka@mail.doshisha.ac.jp, Department of Biomedical Sciences and Informatics, Doshisha University, Kyoto, Japan. ORCID: 0000-0001-9621-5871.

Tomoyuki Hiroyasu, tomo@is.doshisha.ac.jp, Department of Biomedical Sciences and Informatics, Doshisha University, Kyoto, Japan. ORCID: 0000-0003-0730-7905.

Masaki Takeda, mtakeda@mail.doshisha.ac.jp, Faculty of Health and Sports Science, Doshisha University, Kyoto, Japan.

* Correspondence: Satoru Hiwa.

Abstract
Exercise training integrating physical and cognitive activities is gaining attention because of its potential benefits for brain health. This study focuses on exercise training using a dart game called Wellness Darts. Wellness Darts is a sport involving throwing darts and walking to pull them out of the board, memorizing the score, and subtracting it from the total score, thus allowing the performance of two tasks simultaneously: exercise and calculation. It is expected to maintain and improve cognitive function, and whether the continual dart training affects brain function is of great interest. Before conducting the longitudinal study revealing its effect on brain function, we aimed to cross-sectionally confirm the difference in hemispheric lateralization between expert and non-expert players. Functional near-infrared spectroscopy (fNIRS) was used to measure brain activity for three groups: an expert older group who practiced darts continually, a non-expert older control group, and a non-expert younger control group. Their brain activity patterns were quantified by lateralization index (LI) and compared between groups. The results showed that the younger and the expert older groups had significantly higher LI values than the non-expert older group and did not differ between the expert older and the younger groups. Our results suggest that the Wellness Darts possibly promote hemispheric lateralization.

Keywords: exercise training, darts, functional near-infrared spectroscopy, cognitive function, hemispheric lateralization

1   Introduction

Cognitive decline is one of the most crucial issues to be addressed in the aging population. Various interventions have been proposed to maintain and improve cognitive function in older adults, including physical activities and cognitive exercises. In this context, exercise and sports that integrate physical and cognitive activities are gaining attention because of their potential benefits for brain health.

Wu et al. found that Mind-Body exercise, particularly tai chi and dance, improves global cognition, cognitive flexibility, working memory, verbal fluency, and learning in cognitively impaired and non-disabled older adults [1]. Furthermore, Babaei et al. suggested that frequent moderate aerobic activity can be associated with improved neurocognitive performance for older adults [2]. Gheysen et al. stated that combined physical and cognitive activity programs should be promoted to prevent and treat cognitive decline in older adults [3]. Similarly, Rieker et al. found that performing cognitive and physical exercise simultaneously and interactive training produced the most significant gains in executive functions, speed, and global cognition, and the most significant improvements in physical functions [4]. Also, physical activity associated with exercise training has been shown to cause substantial changes in brain health and cognitive performance, potentially affecting a wide range of cognitive functions, including memory, attention, and executive function [5]. Additionally, Liu et al. suggested exercise-induced enhancement of executive function in young individuals, providing practical evidence to encourage regular exercise to maintain cognitive function and promote brain health [6]. These previous studies imply that exercise training, especially the combined physical and cognitive activity program, is one of the most effective interventions for brain health.

This study focused on exercise training using a darts game called Wellness Darts. Darts is one of the sports that seek fun and gameplay by subtracting the score obtained by the player from the points they have and reducing it to zero. Wellness Darts also has the unique feature of calculating scores by rote [7]. "Rote," in this case, refers to the process of quickly mentally calculating the score of a dart thrown by a player on a dartboard and subtracting it from the points the player has. Wellness Darts is a combined physical and cognitive activity program that involves not only throwing darts but also walking to pull them out of the board, memorizing the score, and subtracting it from the total score, thus allowing the performance of two tasks at the same time: exercise and calculation [7]. Simultaneous performance of the motor and cognitive tasks, each effective in improving cognitive function as a single task, is more effective in increasing PFC activity and improving cognitive function than performing either task alone [8]. Takeda et al. conducted cognitive tests before, one month, and two months after an experiment on a Wellness Darts intervention group and a control group of older adults [7]. The results showed that continuing Wellness Darts increased older adults' short-term memory test (STM test) scores. However, the impact of continual practice of Wellness Darts on brain function is still being determined. To develop Wellness darts as a combined physical and cognitive activity program, it is essential to understand their potential effects on brain function.

This study investigates the cognitive status and brain function of participants who practiced Wellness Darts in older adults. Their characteristics of brain function and the performance of cognitive tests will be compared cross-sectionally among older adults who are proficient with darts, older adults who are inexperienced with darts, and younger adults who are inexperienced with darts. The Hemispheric Asymmetry Reduction in OLDer adults (HAROLD) model proposed by Cabeza et al. asserts that the degree of Frontal lobe (FL) lateralization seen in youth decreases with age [9, 10, 11,12, 13, 14, 15, 16]. In addition, Erickson et al. reported that when cognitive training was conducted on healthy older adults, lateralization was observed with improved cognitive performance,

bringing the brain activation pattern closer to that of younger adults [12]. That is why increased
lateralization can be used as a measure of cognitive improvement. On the other hand, brain regions
such as the inferior frontal gyrus (IFG), superior frontal gyrus (SFG), middle frontal gyrus (MFG)
[17], and DLPFC have been reported to be activated during mental calculation [18]. Furthermore, the
inferior parietal lobule (IPL) in patients with cognitive decline, such as MCI and Alzheimer's disease,
is reported to be less activated [19].

Based on these findings, we proposed the following three hypotheses: (1) older adults who practice
Wellness Darts (referred to as the 'expert older group') will show a level of hemispheric
lateralization similar to that of younger people who had never played Wellness Darts before (referred
to as 'younger group' as a control), with activation of the dorsal superior frontal gyrus (SFGdor) and
MFG, which are known to be activated during mental arithmetic. (2) There is no difference in the
level of IPL activation between the expert older group and the younger group. (3) There is a positive
correlation between the level of hemispheric lateralization in brain activity and the number of days of
Wellness Dart experience in the expert older group. These three hypotheses were tested by measuring
brain activity using functional near-infrared spectroscopy (fNIRS), a non-invasive functional brain
imaging system robust to body movement. The third group, the older adults who had never played
Wellness Darts (referred to as the 'non-expert older group'), was used as the older control to confirm
that hypotheses (1) and (2) were valid only for the expert older group.

2   Materials and Methods

2.1   Participants

Twenty-one healthy younger adults (13 males, seven females; mean age: 22.5 ± 0.8 years) who had
never played Wellness Darts, 21 healthy older adults (8 males, 13 females; mean age: 75.1 ± 4.3
years) who continually played Wellness Darts, and 21 healthy older adults (4 males, 17 females;
mean age: 74.1 ± 4.5 years) who had never played Wellness Darts participated in the experiment.
These three participant groups were referred to as the younger group, the expert older group, and the
non-expert older group. The participants for the expert older group were recruited from Wellness
Darts players in the Kyotanabe-Doshisha Sports Club (KDSC), one of the community sporting clubs
in Kyoto, Japan. The participants for the non-expert group were also recruited based on the
acquaintanceship of the members of the expert older group. Only healthy participants over 60 could
participate in this study for these two groups. For the younger group, the healthy undergraduate and
graduate school students aged over 20 were recruited from Doshisha University, Kyoto, Japan. The
sample size for each group was determined with a significance level of 0.05 and a power of 0.8 to
detect the effect size of Cohen's $d = 0.79$ reported in a previous study comparing the degree of
hemispheric lateralization in older adults using fNIRS [**Error! Reference source not found.**]
(G*Power was used for the calculation).

2.2   Wellness Darts task

Wellness Darts is a darts game developed by the Japan Wellness Darts Association. The standard
game of Wellness Darts is the "Zero One Game 251". In the game, each player is assigned a score of
251 and throws darts until the score reaches zero, subtracting the score of each dart thrown from the
holding points. The first player to get a score of exactly 0 wins. Each player can throw darts three
times in their turn (which is defined as one set) and is required to make their score exactly zero on the
third throw for either of five sets. In addition, each player must subtract the points earned from their

score by themselves. We assumed that these mental arithmetic operations, the planning of throws to
reach zero as quickly as possible, and the exercise associated with throwing darts could be effective
in improving cognitive function because the Simultaneous performance of the motor and cognitive
tasks said to be effective in increasing PFC activity and improving cognitive function [8]. However,
since the official rules of Wellness Darts place a physical burden on the participants due to the long
duration of the experiment, in this study, the initial score of the subjects was set at 128 points, and
each subject played the game for a maximum of five sets. Furthermore, the game was terminated
even if the score did not reach zero in the fifth set.

2.3   Procedure

Participants played up to five sets of Wellness Darts with a maximum of 128 holding points while
brain activity and head acceleration were simultaneously recorded. As described below, the
accelerations were used to remove noise due to head movements. In each of the three throws in one
set of the Wellness Darts game, the participants were instructed by the recorded voice prompts and
the beep cue from the experimental computer on when to hold the dart, throw, and start the
calculation; the participants performed each action following the instructions.

During the planning block set at the beginning of each throwing session, participants were asked to
determine where to hit (designate scoring points) within 5 s for the first throw and within 13 s for the
remaining two throws. At the beginning of the second and third throwing sessions, they were
provided an extra 8 s in the planning blocks to tally the points from the previous throws. Of note, we
excluded the beginning 8 s. Instead, we applied the later 5 s for the subsequent analyses because we
assumed that the first throw did not involve the confirmation of the hit point and employed different
cognitive processes than those used in determining where to hit. The planning block was followed by
dart-throwing within 3 s after the beep cue.

Following dart throwing, the participants were instructed to move their gaze away from the dartboard
to the fixation point and not to think of anything in particular. This waiting block consisted of a
random duration between 28 - 32 s to reduce the anticipatory response. After completing the three
throwing sessions, the participants commenced the calculation block, where they were instructed to
subtract the scores, they received during the previous throws from the total score and write the
remaining points on the scoring sheet. Then, they were asked to press a button to proceed to the 20 s
rest block. During the rest block, they stayed the same as in the waiting block. This procedure was
repeated for five sets. The overall procedure is illustrated in Figure 1.

All participants were informed of the methods and risks of the experiment and signed a written
informed consent form. This study was conducted in accordance with the Research Ethics Committee
of Doshisha University (Kyoto, Japan) (approval code: 20014). Before the experiment, participants
were asked to complete a questionnaire, a dart-throwing performance test, and five cognitive tests:
Simple Cognitive (SC) test, Computational Ability Test, Short-Term Memory (STM) Test, Trail
Making Peg Test (TMPT), and CogEvo. Participants then performed one set of dart-throwing before
being fitted with the fNIRS device to understand the experimental procedure.

2.4   fNIRS data acquisition

Changes in the concentration of oxyhemoglobin (HbO), deoxyhemoglobin (HbR), and total
hemoglobin (HbT) in the participants were measured using NIRSport2 (NIRx Medical Technologies,
LLC). The sampling frequency was 4.36 Hz, and the wavelengths of the near-infrared light were 760
and 850 nm, respectively [21]. As shown in Figure 2 (b), 16 sources and 14 detectors were placed on

the frontal, parietal, and temporal regions of the head. A pair of sources and detectors adjacent at 30-
mm intervals constituted one standard channel, and 44 standard channels were placed. In addition, a
short-distance detector bundle (NIRx Medical Technologies, LLC) was established to eliminate
extra-cerebral blood flow alterations [22, 23]. The signals obtained from the 16 short-distance
channels were used to detect extracerebral signals and to remove the effect of skin blood flow in the
signals measured by the standard channels. To remove noise from body motion, a six-axis
accelerometer (NIRx Medical Technologies, LLC) was used to measure head acceleration at a
sampling frequency of 4.36 Hz, the same frequency as the change in hemoglobin concentration [24].
The accelerometer was placed at the top of the head, as shown in Figure 2 (b). Presentation software
(Neurobehavioral System Inc.) was used to control the presentation and timing of the recorded voice
prompts and the beep cue during the experiment, and the signal measurement and execution of the
experimental task were synchronized. The 3D coordinates of the light source and detection probe
were measured using a 3D magnetic space digitizer (Fastrak, Polhemus, USA) to estimate the exact
measurement position of the brain area for each individual. The measured 3D coordinates were
entered into the 'register2polhemus' function, and the 'depthmap' function implemented in the NIRS
toolbox [25], and the distance from each standard channel to the brain surface was calculated. On the
basis of this distance, the regions of the brain defined by the automated anatomical labeling atlas [26]
corresponding to each standard channel were identified.

2.5　Cognitive tests

Five cognitive tests were conducted to identify differences in cognitive function between the three
participant groups. As the current study is a cross-sectional comparison, it is impossible to show
whether the continued practice of Wellness Darts in older adults affects cognitive function; however,
it was preliminarily investigated to confirm its possibility. That is why the statistical tests were
conducted between two groups: the expert older and the non-expert older groups.

2.5.1 Simple Cognitive test

The Simple Cognitive (SC) test was used to verify that the participant's cognitive function had not
declined [27]. The SC test is a 50-point cognitive function test that can identify early cognitive
decline that cannot be detected by the mini-mental state examination (MMSE), which is often used to
examine dementia. A score of fewer than 20 points on the SC test is considered suspicious of
cognitive decline. In this study, only participants with a score of 20 or higher were considered to
have no cognitive decline and were used in subsequent analyses.

2.5.2 Computational ability test

The participants' computational abilities were measured by administering a test consisting of 30
subtraction problems between two digits, between three and two digits, and between three digits.
Differences in mean scores on the computational ability test between the expert older group and the
non-expert older group were t-tested ($p < .05$).

2.5.3 STM

To evaluate the short-term memory of the participants, the Short-Term Memory (STM) Test was
conducted on a computer [7]. After the instruction "Please memorize the numbers in order" was
displayed on the monitor, nine disordered numbers were displayed on the screen one by one per
second. The participants were then instructed to "write the numbers in order," and they wrote the
memorized numbers on a piece of paper during a 20-s time frame. The difference in mean STM test
scores between the expert older group and the non-expert older group was t-tested ($p < .05$).

2.5.4 CogEvo

CogEvo was used to test the cognitive abilities of the participants; CogEvo is a computer-assisted
cognitive function test that assesses disorientation, attention, memory, planning, and spatial
awareness [28]. CogEvo can reportedly capture mild changes in cognitive function and is a simple
and convenient ICT tool to assess cognitive changes over time in middle-aged and older adults, as
well as during the preclinical stages of dementia [28, 29]. CogEvo has 12 different tasks; in this
experiment, we selected flashlight, visual search, number step, and just fit as tasks related to
Wellness Darts from the disorientation, attention, memory, planning, and spatial awareness
components of CogEvo, respectively. The difference in mean CogEvo scores between the expert
older group and the non-expert older group was t-tested ($p < .05$).

2.5.5 TMPT

The Trail making peg test (TMPT) was conducted to assess participants' cognitive functioning,
particularly attentional function and finger dexterity [30]. The TMPT is a peg-movement test that
reflects finger dexterity combined with the trail-making test that measures visual attention and
executive function [31]. In the TMPT, participants quickly insert 25 cylindrical wooden sticks, called
pegs, into numbered holes in numerical order. The board containing the pegs and the board
containing the holes were separated from each other. The difference in mean completion time was t-
tested ($p < .05$).

2.6   Dart throwing performance test

To confirm that the dart-throwing skill of the expert older group was higher than that of the control
groups, the dart-throwing performances of the three groups were tested. In this test, a target score of
65 points was set and participants were required to throw the darts 10 times and tally their scores to
match the target score. The differences in total dart performance test scores between the expert and
younger groups and between the expert and non-expert older groups were t-tested ($p < .05$).

2.7   Questionnaires

Participants were asked to respond to questions regarding their dominant hand, the number of days
per week they practiced Wellness Darts, and the number of years they had been practicing. From
these responses, the total number of days of experience for each participant was estimated.

2.8   fNIRS data processing

2.8.1 Preprocessing

Previous studies have concluded that HbO in fNIRS is more strongly related to blood oxygenation
level-dependent (BOLD) signals measured by functional magnetic resonance imaging (fMRI) than
HbR [32, 33, 34]. The fMRI has better spatial resolution than fNIRS and has been used in many
cognitive neuroscience studies. In this study, we used HbO to analyze brain activity. The fNIRS data
were preprocessed using the NIRS toolbox [26] to remove physiological noise such as skin blood
flow, heart rate, and pulse rate. Optical intensity data acquired from the fNIRS system were
converted to optical density (OD) data and then to Hb concentration data based on the modified Beer-
Lambert law [22]. A bandpass filter (0.008-0.09 Hz) was applied to remove noise such as low-
frequency drift and heartbeat [34, 35].

2.8.2 Activation analysis

The level of task-induced brain activation for each individual was estimated using the iteratively
reweighted least-squares model (AR-IRLS) with short separation (SS) and acceleration, a general
linear model (GLM)-based analysis method [24, 36]. In GLM, task-related brain activation was
modeled by the hemodynamic response function (HRF). The measured brain activity, i.e., HbO
signal, was regressed on the ideal task-related activity calculated from the canonical HRF and a
design matrix indicating the type of experimental task, and the partial regression coefficients were
estimated by IRLS, a least-squares method. The AR-IRLS allows brain activation analysis to
consider the effects of arbitrary confounding factors on brain activity, such as the SS signal measured
with a short-distance channel, blood pressure, respiration, and acceleration. In this study, we
introduced the SS signal and acceleration data as physiological regressors of GLM to reduce the
effects of skin blood flow and body motion on the HbO signal [24, 37].

2.8.3 Hemispheric lateralization analysis

The level of hemispheric lateralization was calculated from the lateralization index (LI) shown in
Equation 1.

$$LI = \frac{|AL - AR|}{|AL| + |AR|} \qquad (1)$$

where AL is the sum of β values in the regions of interest (ROIs) in the left hemisphere and AR is the
sum of those in the right hemisphere [38, 39]. We used absolute values for the LI values to allow for
the fact that the dominant hemisphere differs depending on the dominant hand.

2.9 Statistical analysis

2.9.1 Activation analysis

For individual-level analysis, the following two contrasts were created for the HbO data of the
younger group, the expert older group, and the non-expert older group: "rest vs. calculation" and "rest
vs. planning". The β-values for each experimental condition and each channel were calculated. The
statistical significance of the task-related brain activity was then verified by testing the null
hypothesis that the estimated β coefficient was not significantly different from zero (*t*-test, $p < .05$).
Note that multiple comparisons were corrected using the Benjamini-Hochberg method [37]. In
addition, to test the hypothesis that there is no difference in the level of IPL activation between older
adults who continually practiced Wellness Darts and younger adults who were not experienced in
Wellness Darts, β values of IPL were compared between the three groups (*t*-test, $p < .016$, Bonferroni
corrected).

Next, for the group-level analysis, we constructed the most consistent contrast images of group-level
brain activity using simultaneously the β values obtained from the individual-level analysis of all
participants, based on [25]. To estimate significant group-level activity, we used participants as
random effects and conditions (planning, calculation, and rest blocks) as fixed effects. Finally, all
participants were considered as a sample drawn from the population and a one-sample t-test was used
to test if the population mean was greater than zero with respect to the *t*-value. The significance level
was set at $p < .05$ (false discovery rate corrected at peak level), and the brain regions activated during
the calculation and planning blocks were estimated.

2.9.2 Hemispheric lateralization analysis

The LI values were used to compare the degree of hemispheric lateralization among the three groups
of participants (*t*-test, $p < .016$, Bonferroni corrected). In addition, we tested whether there was a
positive correlation between the number of days of Wellness Dart experience and LI values in the
expert older group (Spearmans' rank correlation test, $p < .05$).

3   Results

3.1   Descriptive statistics on three groups

The descriptive statistics on three groups, including age, days of practicing wellness darts, cognitive
test scores, and scores of dart-throwing performance tests, are summarized in Table 1. The following
subsections describe the details of the results including statistical tests.

3.2   fNIRS data

3.2.1 Hemispheric lateralization analysis

We hypothesized that the LI values of the younger group and the expert older group would be larger
than those of the non-expert older group. LI values were calculated from the respective β values of
the two task blocks (planning and calculation) for the three participant groups. The LI values in the
planning and calculation blocks of the younger group were 0.45 ± 0.34 and 0.49 ± 0.35, respectively.
The LI values in the planning and calculation blocks of the expert older group were 0.53 ± 0.38 and
0.61 ± 0.35, respectively. Finally, for the non-expert older group, they were 0.23 ± 0.18 and 0.50 ± 0.38
in the planning and calculation blocks, respectively. The box plots of the LI values in the planning
and calculation blocks for the three groups of participants are shown in Figure 3 (d) and Figure 4 (d),
respectively. Furthermore, the β maps of the participants who showed the closest values to the
median in the LI values for each group of participants are shown in Figure 3 (a)-(c) and Figure 4 (a)-
(c).

There were no significant differences between the LI values of the younger group and the expert
older group in the planning block ($t = 0.6730$, $p = 0.504$, $d = 0.213$). On the other hand, in the
planning block, the LI values of the expert older group were significantly greater than those of the
non-expert older group ($t = 3.188$, $p = 0.003$, $d = 1.005$). Furthermore, the LI values of the younger
group were significantly greater than those of the non-expert older group in the planning block ($t =$
2.58, $p = 0.014$, $d = 0.815$).

In the calculation block, there were no differences in LI values between the younger group and the
expert older group ($t = 1.009$, $p = 0.319$, $d = 0.319$). Similarly, there were no differences in LI
between the expert older group and the non-expert older group ($t = 0.828$, $p = 0.413$, $d = 0.262$).
There were also no differences in LI values between the younger group and the non-expert older
group ($t = 0.144$, $p = 0.886$, $d = 0.046$).

We also hypothesized that the number of days of Wellness Dart experience in the expert older group
would be positively correlated with LI values. The results of an Spearman's rank correlation test
between the number of days of Wellness Dart experience and LI values showed a negative significant
correlation for the planning block ($\rho = -0.439$, $p = 0.046$, Figure 5 (a)) but no significant correlation
for the calculation block ($\rho = -0.037$, $p = 0.874$, Figure 5 (b)).

To investigate the Wellness Dart experience of the expert older group in terms of skill, the mean
darts performance test score (target score was 65 points; the closer the total score of 10 dart throws
was to 65 points, the better the skill) was compared between the three participant groups. The mean
scores for the younger group, the expert older group, and the non-expert older group were 39.3±13.4,
49.9±5.5, and 31.9±13.7, respectively. There were no participants who exceeded the 65 points. As
shown in Figure 6, the expert older group showed better dart performance compared to the younger
group ($t = 3.247$, $p = 0.003$, $d = 1.022$). Similarly, the expert older group showed better performance
than the non-expert older group ($t = 5.400$, $p = 0.00001$, $d = 1.700$).

Furthermore, inspired by the results of no significant 'positive' correlation between the number of
days of Wellness Dart experience and LI values, we assumed that dart proficiency could be assessed
not only by the number of days of experience but also by dart-throwing performance. Therefore, a
positive correlation between the total dart performance test score and the degree of hemispheric
lateralization of brain activity in the expert older group was also tested (Spearman's rank correlation
test, $p < .05$). It should be noted that this is an exploratory analysis based on the result shown in
Figure 5. However, no significant correlation between the total dart performance test score and the LI
value was confirmed (planning block: $\rho = 0.324$, $p = 0.152$, Figure 7 (a); calculation block: $\rho = -0.225$, $p = 0.327$, Figure 7 (b)).
344

3.2.2 Activation analysis

The activation analysis shown in Figure 8 indicates that in the planning block, there was no
difference between the left and right IPL (IPL.L: $t = 0.738$, $p = 0.466$, $d = 0.234$. IPL.R: $t = 0.134$, p
$= 0.894$, $d = 0.043$). Similarly, there was no difference between the expert older group and the non-
expert older group (IPL.L: $t = 0.042$, $p = 0.966$, $d = 0.014$; IPL.R: $t = 0.821$, $p = 0.419$, $d = 0.257$).
Furthermore, there was no difference between the younger group and the non-expert older group
(IPL.L: $t = 0.711$, $p = 0.481$, $d = 0.226$. IPL.R: $t = 0.967$, $p = 0.343$, $d = 0.312$).

In the calculation block, there was no difference between the left and right IPL (IPL.L and IPL.R) of
the younger group and the expert older group (IPL.L: $t = 1.211$, $p = 0.236$, $d = 0.381$; IPL.R: $t =$
0.206, $p = 0.838$, $d = 0.068$). Similarly, there was no difference between the expert older group and
the non-expert older group (IPL.L: $t = 1.625$, $p = 0.115$, $d = 0.513$; IPL.R: $t = 1.642$, $p = 0.109$, $d =$
0.523). There was no difference between the younger group and the non-expert older group (IPL.L: t
$= 2.176$, $p = 0.036$, $d = 0.688$. IPL.R: $t = 1.844$, $p = 0.074$, $d = 0.621$).

Tables 2 to 7 summarize the results of individual-level analysis and show the regions of interest
where the p-values are less than 0.05. Tables 8 to 10 summarize the results of group-level analysis
and show only the ROIs. There was significant activation in the ROIs in the individual-level analysis
in the expert older group but no significant activation in the group-level analysis.

3.3 Cognitive tests

3.3.1 Computational ability test

364 The mean computational test scores for the younger group, the expert older group, and the non-expert
365 older group were 24.0 ± 3.8, 20.5 ± 3.8, and 17.1 ± 5.5, respectively. As shown in Figure 9, the
366 expert older group had higher scores on the computational skills test than the non-expert older group
367 ($t = 2.275$, $p = 0.029$, $d = 0.719$).

3.3.2 STM test

369 The mean STM test scores of the younger group, the expert older group, and the non-expert older
370 group were 74.6 ± 13.6, 57.1 ± 9.5, and 49.4 ± 9.9, respectively. As shown in Figure 10, the expert
371 older group showed higher STM test scores than the non-expert older group ($t = 2.475$, $p = 0.018$, $d = 0.792$).
372 0.792).

3.3.3 CogEvo

374 The scores of the five CogEvo tasks for each group of participants are summarized in Figure 11. In
375 the expert older and non-expert older groups, there were no significant differences in the CogEvo
376 scores (disorientation: $t = 1.892$, $p = 0.067$, $d = 0.598$, visual search: $t = 1.435$, $p = 0.159$, $d = 0.454$;
377 flashlight: $t = 1.386$, $p = 0.174$, $d = 0.438$; number step: $t = 1.722$, $p = 0.093$, $d = 0.550$; just fit: $t = 0.625$, $p = 0.534$, $d = 0.198$).
378 0.625, $p = 0.534$, $d = 0.198$).

3.3.4 TMPT

380 The mean TMPT completion times for the younger, expert older, and non-expert older groups were
381 41.1 ± 4.5, 58.1 ± 8.5, and 58.9 ± 9.2, respectively. There was no significant difference in TMPT
382 completion time between the older and non-expert older groups ($t = 0.272$, $p = 0.787$, $d = 0.086$).

4   Discussion

4.1   Hypothesis 1

As seen in the results of the individual-level analysis in Tables 2-7, we were able to confirm the
activation of SFGdor and MFG in the three participant groups. However, there was no significant
activation of these regions in the group-level analysis. This may be because the mean activation level
was estimated to be lower after averaging the results of the group as the individual activated areas
varied even in the same region due to the distinct differences in the arrangement of measurement
channels in the fNIRS device.

On the other hand, Figure 3 (d) shows that the LI values of the younger group and the expert older
group were significantly higher than those of the non-expert older group in the planning block. In
addition, there was no difference in LI values between the younger group and the expert older group.
This indicates that the brain activation patterns of participants in the younger group and the expert
older group are hemispherically lateralized during the planning of dart-throwing and while
determining where to hit. The β-map in Figure 3 (a)-(c) also shows that the activation levels in the
younger and expert older groups were higher in one hemisphere, while the non-expert older group
showed no hemispheric difference. In the non-expert older group, there is no difference between the
hemispheres, and bilateral activation levels of SFGdor and MFG are high. This result is consistent
with a previous study in which left-right hemispherical asymmetry increased when cognitive training
was performed in healthy older adults [12]. In summary, although SFGdor and MFG were not
significantly activated in the expert older group in group-level analysis, Hypothesis 1 was established
during the planning of dart throwing, suggesting that continual Wellness Dart training promotes
hemispheric lateralization in brain activity in older adults.

4.2   Hypothesis 2

Figure 8 indicates that there were no significant differences in the β-values of the left and right IPL
between the younger group and the expert older group. Further, there was no significant difference in
the β-values of the left and right IPL between the younger and non-expert older groups, and between

the expert and non-expert older groups. However, in the three participant groups in the present study,
the IPL activation was comparable, suggesting that there was no difference in IPL activation between
the older groups. The previous study to which the hypothesis refers involved healthy subjects and
patients with MCI and Alzheimer's disease [19]. The participants in the present study were only
healthy adults, suggesting that the difference in cognitive function between the older groups and the
younger group did not affect IPL activation.

4.3   Hypothesis 3

As shown in Figure 5, there was no positive correlation between the number of days of Wellness Dart
experience and the LI value of the expert older group; therefore, hypothesis 3 was not supported. On
the other hand, in Figure 6, the dart performance of the expert older group was higher than that of the
other two groups. In addition, Figure 7 implies a positive correlation between dart performance and
LI value in the planning block, although it is not statistically significant. The total score of our dart
performance test indicates the accuracy of dart throwing, which is improved by the number of days of
experience. Therefore, the results of Figures 6 and 7 may contribute to the validity of hypothesis 3.
Also, since the current results suggested weak correlations, the correlations should be tested again
with a larger sample size. These issues would be investigated in future longitudinal controlled studies
of the Wellness Darts intervention that measure dart performance levels to verify the increase in LI
values promoted by Wellness Dart training.

4.4   Cognitive tests

428 In addition to hypotheses testing, the results of three cognitive function tests to determine the impact
429 of Wellness Darts on cognitive function are also discussed.

4.4.1 Computational ability test

The score of the expert older group was significantly higher than that of the non-expert older group.
In other words, the expert older group had better computational ability than the non-expert older
group. According to previous studies, cognitive function, particularly for high-load tasks, declines as
the brain deteriorates with age [40]. Therefore, the results of the computational ability test in this
study suggest that continual practice of Wellness Darts can reduce the decline of calculation ability
due to aging.

4.4.2 STM test

438 The STM test scores of the expert older group were higher than that of the non-expert older group,
439 and the difference was significant. Fandakova et al. [41] have shown that short-term memory
440 capacity declines with age. Therefore, our results indicate that constant training with Wellness Darts
441 can reduce the age-related decline in short-term memory capacity.

4.4.3 CogEvo

No significant differences were found in the scores between the expert older group and the non-
expert older group on the five CogEvo tasks. Although CogEvo has been reported to be able to
significantly distinguish between Alzheimer's disease, MCI, and cognitively normal groups of older
adults [28, 29], our results showed that the expert older group and the non-expert older group did not
differ significantly in their scores on the five CogEvo tasks. In fact, our results indicated that the
expert older and non-expert older groups had similar cognitive levels. In other words, the older
groups in our experiment had normal levels of cognitive function or the differences in cognitive

function were at levels that were undetectable by CogEvo. Another explanation could be that the
older groups were unfamiliar with the operation of CogEvo as it was conducted using a tablet device.

4.4.4 TMPT

As seen in Figure 12, there was no significant difference in the completion time of TMPT between
the expert older and the non-expert older groups. TMPT requires skill from the participant as a
component of physical function. We expected that participants engaged in ongoing Wellness Darts
exercises would demonstrate improved skillfulness; however, in this cross-sectional study, no
differences were observed between the expert older and non-expert older groups. A previous
longitudinal study that divided older adults into a dual-task (DT) exercise group and a control group
observed a significant difference in TMPT before and after intervention in the DT exercise group, but
no significant difference was identified in TMPT between the two groups [42]. Due to the pilot
nature of the current study based on a cross-sectional comparison, a possible future direction could be
to investigate whether the Wellness Darts intervention contributes to improving the TMPT
completion time in longitudinal randomized controlled trials.

4.5   Strengths of the current study, limitations, and future direction

Based on our findings, only hypothesis 1 was supported in the planning of dart throwing and in
determining where to hit. Due to the pilot nature of this study, the sample size was limited, and the
effects of several unseen parameters such as education level, socio-economic factors, body mass
index, and smoking were not controlled and investigated. These limitations make the current study
insufficient to demonstrate that Wellness Darts truly affected brain lateralization. However, despite
these limitations, the current study revealed that older adults with Wellness Dart experience have a
higher tendency toward hemispheric lateralization than those with no experience. The current results
are valuable in motivating us to proceed to future longitudinal randomized controlled trials to explore
the effects of continual Wellness Dart training on hemispheric lateralization with a larger sample size.

5   Conclusions

Wellness Darts is a cognitively challenging sport that combines exercise and calculation. Here we
aimed to cross-sectionally confirm the difference in hemispheric lateralization between expert and
non-expert players as a pilot study prior to the longitudinal randomized controlled trials. The results
showed that the younger and the expert older groups had significantly higher LI values than the non-
expert older group and that these values did not differ between the expert older and the younger
groups. These results suggest that the Wellness Darts possibly promote hemispheric lateralization.
481

6   Declarations

6.1   Ethics approval and consent to participate

Studies involving human participants were reviewed and approved by the Research Ethics
Committee of Doshisha University, Kyoto, Japan (approval code: 20014), and conducted in
accordance with the Declaration of Helsinki. All participants were informed about the experimental
method as well as the risks and signed written informed consent forms to participate in the study.

6.2   Consent for publication

he participants provided their written informed consent prior to enrolment in the study.

6.3   Availability of data and materials

The datasets generated during and/or analyzed during the current study are available from the
corresponding author upon reasonable request.

6.4   Competing interests

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as potential conflicts of interest.

6.5   Funding

This work was supported by JSPS KAKENHI, Grant Number JP20K11403.

6.6   Authors' contributions

KoT, SH, TH, and MT contributed to the conception and design of the study. KoT and SH wrote the
manuscript. KoT, SH, and MT conducted experiments. KoT performed data analysis. SH, KeT, TH,
and MT advised on the data analysis and experimental design. All authors contributed to the
manuscript revision and read and approved the submitted version.

6.7   Acknowledgments

We are deeply grateful to the members of the Kyotanabe Doshisha Sports Club (KDSC) for their
cooperation in the experiment. We would like to thank Editage (www.editage.com) for English
language editing.

7   References

1.   Chunxiao Wu, Qu Yi, Xiaoyan Zheng, Shaoyang Cui, Bin Chen, Liming Lu, Chunzhi Tang.
Effects of Mind-Body Exercises on Cognitive Function in Older Adults: A Meta-Analysis.
Journal of the American Geriatrics Society, 67(4):749-758, April 2019. doi: 10.1111/jgs.15714.
Epub 2018 Dec 18. PMID: 30565212.
2.   Parvin Babaei and Helya Bolouki Azari. Exercise Training Improves Memory Performance in
Older Adults: A Narrative Review of Evidence and Possible Mechanisms. Frontiers in Human
Neuroscience, 15:771553, 2022. doi: 10.3389/fnhum.2021.771553.
3.   Freja Gheysen, Louise Poppe, Ann DeSmet, Stephan Swinnen, Greet Cardon, Ilse De
Bourdeaudhuij, Sebastien Chastin and Wim Fias. Physical activity to improve cognition in older
adults: can physical activity programs enriched with cognitive challenges enhance the effects? A
systematic review and meta-analysis. International Journal of Behavioral Nutrition and Physical
Activity, 15(1):63, July 4, 2018. doi: 10.1186/s12966-018-0697-x. PMID: 29973193; PMCID:
PMC6032764.
4.   Jennifer A. Rieker, José M. Reales, Mónica Muiños, Soledad Ballesteros. The Effects of
Combined Cognitive-Physical Interventions on Cognitive Functioning in Healthy Older Adults:
A Systematic Review and Multilevel Meta-Analysis. Frontiers in Human Neuroscience.
16:838968, doi: 10.3389/fnhum.2022.838968.
5.   Neva J Kirk-Sanchez and Ellen L McGough. Physical exercise and cognitive performance in the
elderly: current perspectives. Clinical Interventions in Aging, 9:51-62, 2014.
6.   Liu J, Min L, Liu R, et al. The effect of exercise on cerebral blood flow and executive function
among young adults: a double-blinded randomized controlled trial. Scientific Reports, 13:8269,
2023. https://doi.org/10.1038/s41598-023-33063-9.
7.   Masaki Takeda, Nao Yasuda, Satsuki Ito, Masato Abe, Yohei Takamichi, Kazuma Yanagisawa,
and Zsolt Radak. Effects of habitual darts training on cognitive function in elderly people. The
Harris Science Review of Doshisha University, 58(2):53–60, 2017. doi:
10.14988/pa.2017.0000015587.
8.   Eric Hermand, Bertrand Tapie, Olivier Dupuy, Sarah Fraser, Maxence Compagnat, Jean Yves
Salle, Jean Christophe Daviet, and Anaick Perrochon. Prefrontal cortex activation during dual
task with increasing cognitive load in subacute stroke patients: A pilot study. Frontiers in Aging
Neuroscience, 11:160, 2019.
9.   Roberto Cabeza. Hemispheric asymmetry reduction in older adults: the harold model.
Psychology and aging, 17(1):85, 2002.
10. Roberto Cabeza, Nicole D Anderson, Sylvain Houle, Jennifer A Mangels, and Lars Nyberg.
Age-related differences in neural activity during item and temporal-order memory retrieval: a
positron emission tomography study. Journal of Cognitive Neuroscience, 12(1):197–206, 2000.
11. Laura D Müller, Anne Guhn, Julia BM Zeller, Stefanie C Biehl, Thomas Dresler, Tim Hahn,
Andreas J Fallgatter, Thomas Polak, Jürgen Deckert, and Martin J Herrmann. Neural correlates
of a standardized version of the trail making test in young and elderly adults: a functional near-
infrared spectroscopy study. Neuropsychologia, 56:271-9, 2014.
12. Kirk I Erickson, Stanley J Colcombe, Ruchika Wadhwa, Louis Bherer, Matthew S Peterson,
Paige E Scalf, Jennifer S Kim, Maritza Alvarado, and Arthur F Kramer. Training-induced
plasticity in older adults: effects of training on hemispheric asymmetry. Neurobiology of Aging,
28(2):272-83, 2007.
13. Roberto Cabeza, Nicole D Anderson, Jill K Locantore, and Anthony R McIntosh. Aging
gracefully: compensatory brain activity in high-performing older adults. NeuroImage,
17(3):1394–1402, 2002.

14. Patricia A Reuter-Lorenz, John Jonides, Edward E Smith, Alan Hartley, Andrea Miller, Christina Marshuetz, and Robert A Koeppe. Age differences in the frontal lateralization of verbal and spatial working memory revealed by pet. Journal of cognitive neuroscience, 12(1):174–187, 2000.

15. Leonore Blum, David Rosenbaum, Benjamin Röben, Katja Dehnen, Walter Maetzler, Ulrike Suenkel, Andreas J Fallgatter, Ann-Christine Ehlis, and Florian G Metzger. Age-related deterioration of performance and increase of cortex activity comparing time-versus item-controlled fnirs measurement. Scientific Reports, 11(1):1–13, 2021.

16. Anouk Vermeij, Arenda HEA van Beek, Babette LR Reijs, Jurgen AHR Claassen, and Roy PC Kessels. An exploratory study of the effects of spatial working-memory load on prefrontal activation in low- and high-performing elderly. Frontiers in aging neuroscience, 6:303, 2014.

17. Martin Verner, Martin J Herrmann, Stefan J Troche, Claudia M Roebers, and Thomas H Rammsayer. Cortical oxygen consumption in mental arithmetic as a function of task difficulty: a near-infrared spectroscopy approach. Frontiers in human neuroscience, 7:217, 2013.

18. Gert Pfurtscheller, Günther Bauernfeind, Selina Christin Wriessnegger, and Christa Neuper. Focal frontal (de)oxyhemoglobin responses during simple arithmetic. International Journal of Psychophysiology, 76(3):186–192, 2010.

19. Matthias L Schroeter, Timo Stein, Nina Maslowski, and Jane Neumann. Neural correlates of alzheimer's disease and mild cognitive impairment: a systematic and quantitative meta-analysis involving 1351 patients. NeuroImage, 47(4):1196–1206, 2009.

20. Michael K Yeung, Sophia L. Sze, Jean Woo, Timothy Kwok, David H. K. Shum, Ruby Yu, and Agnes S. Chan. Altered Frontal Lateralization Underlies the Category Fluency Deficits in Older Adults with Mild Cognitive Impairment: A Near-Infrared Spectroscopy Study. Frontiers in Aging Neuroscience, 8:59, 2016.

21. Vanessa J Pereira, Karl G Linden, and Howard S Weinberg. Evaluation of uv irradiation for photolytic and oxidative degradation of pharmaceutical compounds in water. Water Research, 41(19):4413–4423, 2007.

22. Theodore J Huppert, Solomon G Diamond, Maria A Franceschini, and David A Boas. Homer: a review of time-series analysis methods for near-infrared spectroscopy of the brain. Applied optics, 48(10):D280–D298, 2009.

23. Ilias Tachtsidis and Felix Scholkmann. False positives and false negatives in functional near-infrared spectroscopy: issues, challenges, and the way forward. Neurophotonics, 3(3):031405, 2016.

24. Alexander von Lühmann, Xinge Li, Klaus-Robert Müller, David A Boas, and Meryem A Yücel. Improved physiological noise regression in fnirs: A multimodal extension of the general linear model using temporally embedded canonical correlation analysis. NeuroImage, 208:116472, 2020.

25. Hendrik Santosa, Xuetong Zhai, Frank Fishburn, and Theodore Huppert. The nirs brain analyzir toolbox. Algorithms, 11(5):73, 2018.

26. Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., Joliot, M. (2002). Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. NeuroImage, 15(1), 273–289.

27. Nagata, H., Yamamoto, Y. and Sakaguchi, R. (2010), P2-123: Our newly developed Simple Cognitive test can be also used by computer or paper. Alzheimer's & Dementia, 6: S349-S349. https://doi.org/10.1016/j.jalz.2010.05.1170

28. Sadanobu Ichii, Takumi Nakamura, Takeshi Kawarabayashi, Masamitsu Takatama, Tetsuya Ohgami, Kazushige Ihara, and Mikio Shoji. Cogevo, a cognitive function balancer, is a sensitive and easy psychiatric test battery for age-related cognitive decline. Geriatrics & Gerontology International, 20(3):248–255, 2020.

29. Hajime Takechi and Hiroshi Yoshino. Usefulness of cogevo, a computerized cognitive
assessment and training tool, for distinguishing patients with mild alzheimer's disease and mild
cognitive impairment from cognitively normal older people. Geriatrics & Gerontology
International, 21(2):192–196, 2021.

30. Jieun Yoon, Hiroko Isoda, and Tomohiro Okura. Evaluation of beneficial effect of a dual-task
exercise based on japanese transitional games in older adults: a pilot study. Aging (Albany NY),
12(19):18957, 2020.

31. Jordi Llinàs-Reglà, Joan Vilalta-Franch, Secundino López-Pousa, Laia Calvó-Perxas, David
Torrents Rodas, Josep Garre-Olmo. The Trail Making Test. 24(2):183-196, 2017.

32. Gary Strangman, Joseph P Culver, John H Thompson, and David A Boas. A quantitative
comparison of simultaneous bold fmri and nirs recordings during functional brain activation.
NeuroImage, 17(2):719–731, 2002.

33. Xu Cui, Signe Bray, Daniel M Bryant, Gary H Glover, and Allan L Reiss. A quantitative
comparison of nirs and fmri across multiple cognitive tasks. NeuroImage, 54(4):2808–
2821,2011.

34. Theodore J Huppert, Richard D Hoge, Solomon Gilbert Diamond, Maria Angela Franceschini,
and David A Boas. A temporal comparison of bold, asl, and nirs hemodynamic responses to
motor stimuli in adult humans. NeuroImage, 29(2):368–382, 2006.

35. Shuntaro Sasai, Fumitaka Homae, Hama Watanabe, Akihiro T Sasaki, Hiroki C Tanabe,
Norihiro Sadato, and Gentaro Taga. A nirs–fmri study of resting state network.NeuroImage,
63(1):179–193, 2012.

36. Congcong Huo, Gongcheng Xu, Wenhao Li, Hui Xie, Tengyu Zhang, Ying Liu, and Zengyong
Li. A review on functional near-infrared spectroscopy and application in stroke rehabilitation.
Medicine in Novel Technology and Devices, page 100064, 2021.

37. Hendrik Santosa, Xuetong Zhai, Frank Fishburn, Patrick J Sparto, and Theodore J Huppert.
Quantitative comparison of correction techniques for removing systemic physiological signal in
functional near-infrared spectroscopy studies. Neurophotonics, 7(3):035009,2020.

38. Mohamed L Seghier. Laterality index in functional mri: methodological issues. Magnetic
resonance imaging, 26(5):594–601, 2008.

39. Eiju Watanabe, Atsushi Maki, Fumio Kawaguchi, Kaoru Takashiro, Yuichi Yamashita, Hideaki
Koizumi, and Yoshiaki Mayanagi. Non-invasive assessment of language dominance with near-
infrared spectroscopic mapping. Neuroscience letters, 256(1):49–52, 1998.

40. Kirstin L Bierre, Samuel JE Lucas, Hayley Guiney, James D Cotter, and Liana Machado.
Cognitive difficulty intensifies age-related changes in anterior frontal hemodynamics: novel
evidence from near-infrared spectroscopy. Journals of Gerontology Series A: Biomedical
Sciences and Medical Sciences, 72(2):181–188, 2017.

41. Yana Fandakova, Myriam C Sander, Markus Werkle-Bergner and Yee Lee Shing. Age
differences in short-term memory binding are related to working memory performance across
the lifespan. Psychology and Aging, 29(1):140–149, 2014.

42. Nathalie Fournet, Jean-Luc Roulin, Fanny Vallet, Marine Beaudoin, Stefan Agrigoroaei, Adeline
Paignon, Cécile Dantzer, and Olivier Desrichard. Evaluating short-term and working memory in
older adults: French normative data. Aging & mental health, 16(7):922–930, 2012.

8   Figure legends

Figure 1. Overview of the experiment. A. The experimental procedure. B. The experimental setup.

Figure 2. Probe displacement of fNIRS. (a) Monte-Carlo simulation results over the frontal cortex,
temporal cortex, and parietal cortex. Red dots represent emitters, blue dots represent detectors, and
yellow lines represent measurement channels, respectively (created using Homer 2 AtlasViewer; v2.8,
p2.1: https://www.nitrc.org/frs/shownotes.php?release_id=3956). The color bar represents the spatial
sensitivity of fNIRS measurements. (b) The two-dimensional fNIRS montage using the International
10-20 measurement system as reference is presented.

Figure 3. βmaps and LI values in the planning block. (a)-(c) βmaps of participants with LI values
close to the median in younger, expert older, and non-expert older groups are shown, respectively. (d)
Distribution of LI values for the three participant groups. *p < .016. βmaps were visualized using the
xjView toolbox and BrainNet viewer.

Figure 4. βmaps and LI values in the calculation block. (a)-(c) βmaps of participants with LI values
close to the median in the younger, expert older, and non-expert older groups are shown, respectively.
(d) Distribution of LI values for the three participant groups. *p < .016. βmaps were visualized using
the xjView toolbox and BrainNet viewer.

Figure 5. Correlation between the number of days of Wellness Dart experience and LI values in the
expert older group. (a) planning block. (b) calculation block.

Figure 6. Dart performance level in the younger, expert older, and non-expert older group. *p<.05.

Figure 7. Correlation between darts performance level and LI value in the expert older group. (a)
planning block. (b) calculation block.

Figure 8. β values of IPL in the younger, expert older, and non-expert older group.

Figure 9. Computational ability test scores for the younger, expert older, and non-expert older group.
*p <.05.

Figure 10. STM test scores for the younger, expert older, and non-expert older group. *p <.05.

Figure 11. Scores on five different CogEvo tasks for the three groups of participants. (a) Orientation.
(b) Visual search. (c) Flashlight. (d) Number step. (e) Just Fit.

Figure 12. TMPT completion time for the younger, expert older, and non-expert older group.

9   Tables

Table 1. Descriptive statistics on the three groups.

Characteristics		Expert older group	Nonexpert older group (older control)	Younger group (nonexpert younger control)
age (years)		75.1±4.3	74.1±4.5	22.5±0.8
Number of days of Wellness Darts experience (days)		170.7±145.6	0.0	0.0
Cognitive tests	SC test	41.1±7.8	44.5±5.8	48.8±2.5
	Computational ability test	20.5±3.8	17.1±5.5	24.0±3.8
	STM test	57.1±9.5	49.4±9.9	74.6±13.6
	TMPT	58.1±8.5	58.9±9.2	41.1±4.5
	CogEvo: disorientation	361.8±52.7	278.2±74.5	341.5±68.8
	CogEvo: visual search	177.9±39.0	161.3±33.8	282.0±41.3
	CogEvo: flashlight	364.3±146.9	308.1±106.4	604.3±169.2
	CogEvo: number step	197.4±40.8	173.5±45.9	280.9±42.8
	CogEvo: just fit	332.2±104.9	342.4±98.7	442.1＋78.2
Dart-throwing performance		49.9±5.5	31.9±13.7	39.3±13.4

680   **Table 2.** Brain regions that showed task-related activation for each individual in the planning block
681   (younger group). Only regions of interest that showed $p < 0.05$ (FDR corrected at peak level) are
682   shown.

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	Participant	Anatomical Region	t-value	p-value (FDR-corrected)
	IPL.L	3.765	5.76E-04		MFG.L	5.449	2.54E-07
1	MFG.L	4.408	4.06E-05	11	MFG.R	8.413	8.76E-16
	MFG.R	10.33	5.57E-23		SFGdor.L	6.024	1.12E-08

Hemispheric lateralization in dart players

	SFGdor.L	7.434	1.48E-12		SFGdor.R	8.779	5.39E-17
	SFGdor.R	9.248	9.70E-19		MFG.L	4.957	1.77E-06
2	IPL.R	-4.224	3.05E-04	12	MFG.R	5.836	2.56E-08
	MFG.L	2.498	0.038		SFGdor.L	6.686	1.67E-10
	MFG.R	-2.941	0.012		SFGdor.R	5.824	2.63E-08
	SFGdor.R	-2.978	0.012		IPL.L	-2.443	0.035
3	IPL.R	3.342	0.002	13	MFG.L	5.219	1.93E-06
	MFG.R	6.801	1.32E-10		SFGdor.L	5.193	2.03E-06
	SFGdor.R	5.896	2.37E-08		MFG.R	7.688	3.71E-13
4	IPL.L	-3.846	1.84E-04	14	SFGdor.R	6.225	2.75E-09
	IPL.R	-4.344	2.73E-05		IPL.L	-3.952	2.21E-04
	MFG.L	-6.759	7.67E-11	15	MFG.R	8.127	8.13E-15
	MFG.R	-2.298	0.028		SFGdor.R	9.337	7.87E-19
	SFGdor,L	-4.55	1.07E-05		IPL.L	-6.443	3.36E-09
	SFGdor.R	-2.273	0.029	16	MFG.R	2.473	0.026
5	MFG.L	8.83	9.44E-18	17	IPL.L	-4.459	9.81E-05

	MFG.R	6.576	2.80E-10		SFGdor.L	−2.842	0.02
	SFGdor.L	19.18	2.87E-77		IPL.L	−9.742	1.12E-20
	SFGdor.R	7.335	1.45E-12		MFG.L	−3.147	0.005
6	IPL.R	6.645	3.50E-10	18	MFG.R	−3.923	3.48E-04
	SFGdor.L	−4.996	2.78E-06		SFGdor.L	−5.72	7.17E-08
	SFGdor.R	2.399	0.028		SFGdor.R	−4.389	5.23E-05
	IPL.L	2.771	0.008		IPL.L	6.362	7.05E-10
	IPL.R	4.191	6.82E-05		MFG.L	7.276	2.11E-12
7	MFG.R	4.289	4.64E-05	19	MFG.R	7.561	3.03E-13
	SFGdor.L	4.583	1.38E-05		SFGdor.L	6.035	4.98E-09
	SFGdor.R	4.365	3.38E-05		SFGdor.R	10.65	3.69E-24
	IPL.L	4.411	1.56E-05	20	MFG.R	4.538	1.88E-05
	IPL.R	4.257	3.09E-05		SFGdor.R	4.636	1.26E-05
8	MFG.L	5.811	1.70E-08		IPL.L	2.346	0.033
	MFG.R	5.317	2.31E-07	21	MFG.L	3.582	9.94E-04
	SFGdor.L	9.745	5.83E-21		MFG.R	12.54	9.28E-34
	SFGdor.R	6.694	7.43E-11		SFGdor.L	4.791	5.92E-06

	IPL.L	3.432	0.002		SFGdor.R	12.01	3.62E-31
10	MFG.L	3.113	0.004				
	MFG.R	4.145	1.42E-04				
	SFGdor.R	3.909	3.05E-04				

683 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
684 gyrus; R: right; L: Left.

685 **Table 3.** Brain regions that showed task-related activation for each individual in the calculation block
686 (younger group). Only regions of interest that showed p < 0.05 (FDR corrected at peak level) are
687 shown.

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	Participant	Anatomical Region	t-value	p-value (FDR-corrected)
1	MFG.R	7.509	2.55E-12		IPL.L	−3.342	0.002
	SFGdor.L	4.194	2.04E-04		MFG.L	5.89	1.75E-08
	SFGdor.R	5.533	7.67E-05	11	MFG.R	6.951	2.72E-11
2	IPL.R	−8.217	1.51E-14		SFGdor.L	6.464	6.57E-10
	MFG.R	−3.969	8.20E-04		SFGdor.R	8.092	8.43E-15
	SFGdor.R	−5.505	7.29E-07	13	IPL.L	−2.444	0.035
3	IPL.L	−	8.59E-07		MFG.L	5.21	1.93E-06

		5.429				9	
	MFG.R	5.85	8.88E-08		SFGdor.L	5.193	2.03E-06
	SFGdor.R	2.523	0.023		MFG.L	3.061	0.004
4	IPL.L	-8.516	7.35E-17	14	MFG.R	7.079	9.36E-12
	IPL.R	-9.993	1.64E-22		SFGdor.L	3.61	6.08E-04
	MFG.L	-2.591	0.011		SFGdor.R	6.778	6.56E-11
	MFG.R	-2.234	0.029	15	MFG.R	7.157	1.92E-11
	SFGdor.L	-5.707	2.23E-08		SFGdor.R	8.688	2.59E-16
	SFGdor.R	-4.508	9.39E-06	16	IPL.L	-6.433	2.05E-09
5	IPL.L	-3.004	0.007		IPL.L	-6.99	4.20E-11
	MFG.L	6.701	1.05E-10	17	MFG.L	-7.918	5.44E-14
	MFG.R	2.79	0.013		SFGdor.L	-8.635	2.21E-16
	SFGdor.L	15.949	2.31E-54		SFGdor.R	-3.074	0.006

Hemispheric lateralization in dart players

	SFGdor.R	2.956	0.008		IPL.L	−4.03	1.73E-04
6	IPL.R	3.981	2.01E-04		MFG.L	−5.082	2.31E-06
	MFG.R	3.87	2.90E-04	18	MFG.R	−3.547	9.51E-04
	SFGdor.L	−5.067	2.42E-06		SFGdor.L	−6.342	2.48E-09
	SFGdor.R	4.977	3.63E-06		SFGdor.R	−4.291	6.13E-05
7	IPL.L	3.011	0.009		IPL.L	3.566	6.62E-04
	IPL.R	4.247	1.77E-04		IPL.R	−2.674	0.011
	IPL.L	3.933	1.30E-04	19	MFG.L	4.345	3.13E-05
8	IPL.R	3.329	0.001		MFG.R	7.278	1.86E-12
	MFG.L	5.935	7.59E-09		SFGdor.L	4.855	3.25E-06
	MFG.R	2.072	0.044		SFGdor.R	11.48	2.56E-28
	SFGdor.L	8.248	2.22E-15	20	IPL.L	−5.517	6.64E-08

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	participant	Anatomical Region	t-value	p-value (FDR-corrected)
	SFGdor.R	2.922	0.004		MFG.L	−2.113	0.042
	IPL.L	−3.998	1.99E-04		MFG.R	4.332	2.16E-05
9	IPL.R	−5.626	1.14E-07		SFGdor.R	3.732	2.57E-04
	MFG.L	−3.412	0.001		MFG.L	3.828	3.12E-04
				21	MFG.R	14.12	6.09E-42
					SFGdor.L	3.692	4.94E-04
					SFGdor.R	12.36	3.38E-33

688 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
689 gyrus; R: right; L: Left.

690 **Table 4.** Brain regions that showed task-related activation in the planning block for each individual
691 (expert older group). Only regions of interest with p < 0.05 (FDR corrected at peak level) are shown.

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	participant	Anatomical Region	t-value	p-value (FDR-corrected)
1	IPL.R	4.418	4.97E-05	12	MFG.L	−5.516	1.33E-06
	MFG.R	−4.478	3.88E-05		MFG.R	−4.304	3.30E-04

Hemispheric lateralization in dart players

Group	Region	Value	p	Group	Region	Value	p
2	IPL.L	5.142	1.98E-06	13	MFG.L	6.548	1.97E-10
	IPL.R	4.167	1.24E-04		MFG.R	3.212	0.002
	MFG.L	-4.8	9.13E-06	14	MFG.L	5.044	3.19E-06
	SFGdor.L	-3.54	0.001		MFG.R	6.031	2.05E-08
3	MFG.L	-7.369	1.65E-12		SFGdor.L	4.537	2.84E-05
	SFGdor.L	-4.309	4.51E-05		SFGdor.R	2.859	0.009
4	IPL.R	-5.123	1.03E-06	15	IPL.L	6.276	3.28E-09
	MFG.L	4.82	4.32E-06		IPL.R	7.845	1.55E-13
5	IPL.R	2.806	0.01		MFG.L	4.419	3.04E-05
	SFGdor.L	-3.32	0.002		MFG.R	5.777	6.03E-08
6	MFG.L	-2.802	0.014		SFGdor.R	4.734	8.11E-06
7	MFG.L	4.296	1.29E-04	16	IPL.R	-4.795	2.22E-05
	SFGdor.L	3.35	0.003	17	IPL.L	-4.80	5.20E-06

Participant	Anatomical	t-	p-value	Participant	Anatomical	t-	p-value
		2				6	
8	IPL.L	-4.035	2.56E-04		MFG.L	-2.944	0.006
8	SFGdor.R	-2.635	0.02		MFG.L	-2.989	0.007
9	IPL.L	3.995	4.11E-04	19	MFG.R	-5.129	1.74E-06
9	SFGdor.L	3.754	9.62E-04	19	SFGdor.L	-4.712	1.02E-05
10	IPL.L	-3.172	0.007	19	SFGdor.R	-2.744	0.012
10	IPL.R	7.868	2.43E-13	20	MFG.L	2.756	0.023
10	MFG.R	2.39	0.049	20	SFGdor.L	3.671	0.001
11	MFG.R	7.11	3.42E-11	21	MFG.R	3.168	0.005
11	SFGdor.R	4.948	1.11E-05	21	SFGdor.R	5.267	1.52E-05

692 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
693 gyrus; R: right; L: Left.

694 **Table 5.** Brain regions that showed task-related activation for each individual in the calculation block
695 (expert older group). Only regions of interest with $p < 0.05$ (FDR corrected at peak level) are shown.

Participa	Anatomica	t-	p-value	Participa	Anatomica	t-	p-value

Hemispheric lateralization in dart players

nt	l Region	value	(FDR-corrected)	nt	l Region	value	(FDR-corrected)
1	IPL.R	4.418	4.97E-05	13	MFG.L	5.359	2.13E-07
	MFG.R	-4.478	3.88E-05		MFG.R	5.988	6.77E-09
2	IPL.L	5.142	1.98E-06		SFGdor.L	3.457	9.19E-04
	IPL.R	4.167	1.24E-04		SFGdor.R	3.344	0.001
	MFG.L	-4.8	9.13E-06	14	MFG.R	3.831	4.78E-04
	SFGdor.L	-3.54	0.001	15	IPL.L	8.888	1.50E-17
3	MFG.L	-7.369	1.65E-12		IPL.R	7.401	1.02E-12
	SFGdor.L	-4.309	4.51E-05		MFG.L	6.731	8.57E-11
4	IPL.R	-5.123	1.03E-06		MFG.R	8.754	3.76E-17
	MFG.L	4.82	4.32E-06		SFGdor.L	5.877	1.52E-08
5	IPL.R	2.806	0.01		SFGdor.R	7.612	2.69E-13
	SFGdor.L	-3.32	0.002	16	MFG.R	8.742	5.85E-17
6	MFG.L	-2.802	0.014		SFGdor.L	4.577	2.61E-05
7	MFG.L	4.296	1.29E-04		SFGdor.R	9.739	1.51E-20
	SFGdor.L	3.352	0.003	17	IPL.L	-3.792	3.28E-04
8	IPL.L	-4.035	2.56E-04		MFG.R	4.375	3.15E-05
	SFGdor.R	-2.635	0.02	18	IPL.L	-2.839	0.012

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	Participant	Anatomical Region	t-value	p-value (FDR-corrected)
9	IPL.L	3.995	4.11E-04		MFG.L	4.933	6.41E-06
	SFGdor.L	3.754	9.62E-04		SFGdor.L	3.433	0.002
10	IPL.L	-3.172	0.007	19	SFGdor.R	3.007	0.013
	IPL.R	7.868	2.43E-13		MFG.L	-11.83	5.34E-31
	MFG.R	2.39	0.049	20	MFG.R	-14.5	1.92E-45
11	MFG.R	7.11	3.42E-11		SFGdor.L	-68.23	0
	SFGdor.R	4.948	1.11E-05		SFGdor.R	-93.95	0
12	MFG.L	-5.55	5.50E-07		IPL.R	3.885	2.64E-04
	MFG.R	-4.926	1.23E-05		MFG.L	-4.797	6.94E-06
				21	MFG.R	4.554	2.01E-05
					SFGdor.L	21.789	4.15E-98
					SFGdor.R	7.121	1.20E-11

696 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
697 gyrus; R: right; L: Left.

698 **Table 6.** Brain regions that showed task-related activation for each individual in the planning block
699 (non-expert older group). Only regions of interest with $p < 0.05$ are shown.

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	Participant	Anatomical Region	t-value	p-value (FDR-corrected)

Hemispheric lateralization in dart players

Group	Region	Value	P	Group	Region	Value	P
1	IPL.L	4.003	3.30E-04	13	IPL.L	-6.647	1.57E-10
	MFG.R	2.857	0.013		IPL.R	-8.842	2.49E-17
2	IPL.L	-4.04	9.84E-04		MFG.L	6.663	1.48E-10
3	MFG.L	-4.888	1.43E-05		MFG.R	3.744	3.67E-04
	MFG.R	-4.201	1.22E-04		SFGdor.R	5.344	3.00E-07
	SFGdor.L	-4.201	1.22E-04	14	IPL.L	2.801	0.009
	SFGdor.R	-4.027	2.30E-04		IPL.R	2.918	0.006
4	MFG.L	-5.972	2.00E-08		MFG.L	-1.914	0.078
	MFG.R	-2.455	0.028		SFGdor.L	-2.504	0.019
	SFGdor.L	-5.021	2.74E-06	15	IPL.L	4.378	4.65E-05
	SFGdor.R	-3.152	0.004		MFG.R	-3.081	0.005
5	MFG.L	4.23	2.98E-04	16	IPL.L	-3.582	0.004

	Region	Value	p		Region	Value	p
	MFG.R	3.643	0.002		MFG.L	-3.388	0.006
	SFGdor.R	5.05	1.69E-05		SFGdor.R	-2.628	0.031
7	MFG.R	-3.466	0.002		IPL.L	-3.349	0.002
7	SFGdor.L	-4.202	2.30E-04		MFG.L	-2.394	0.027
8	MFG.L	4.709	3.04E-05	17	SFGdor.L	-9.055	2.14E-18
8	MFG.R	2.838	0.014		SFGdor.R	-7.301	1.48E-12
8	SFGdor.R	3.386	0.003		IPL.L	14.66	4.98E-46
9	IPL.L	6.602	2.53E-09		MFG.L	5.508	1.02E-07
9	SFGdor.L	-3.184	0.008	18	SFGdor.L	4.667	7.56E-06
10	IPL.L	2.37	0.04		SFGdor.R	5.935	8.93E-09
10	MFG.L	7.529	8.86E-13		IPL.L	8.061	2.25E-14
10	MFG.R	5.415	3.04E-07	19	SFGdor.L	3.538	8.87E-04

	SFGdor.L	7.37	2.32E-12		SFGdor.R	3.536	8.87E-04
	SFGdor.R	6.141	5.18E-09		IPL.L	-2.806	0.018
11	MFG.L	-4.217	1.30E-04	20	IPL.R	-3.373	0.004
	SFGdor.L	-3.748	6.50E-04		MFG.L	-8.005	7.56E-14
	SFGdor.R	-2.585	0.023		SFGdor.L	-2.431	0.046
				21	MFG.L	-2.649	0.024

700 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
701 gyrus; R: right; L: Left.

702 **Table 7.** Brain regions that showed task-related activation for each individual in the calculation block
703 (non-expert older group). Only regions of interest with p < 0.05 are shown.

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	Participant	Anatomical Region	t-value	p-value (FDR-corrected)
1	IPL.L	3.241	0.003	13	IPL.L	-8.63	1.57E-16
	SFGdor.L	-2.994	0.006		IPL.R	-8.138	4.42E-15

Hemispheric lateralization in dart players

Group	Region	Value	p-value	Group	Region	Value	p-value
2	IPL.L	-7.816	7.98E-13	15	MFG.L	5.221	5.11E-07
	MFG.L	-3.72	0.002		MFG.R	3.503	9.32E-04
3	IPL.L	-2.901	0.008		SFGdor.R	5.609	6.61E-08
	IPL.R	-2.726	0.013		IPL.R	-2.658	0.025
	MFG.L	-5.086	1.88E-06		MFG.L	-4.417	8.75E-05
	SFGdor.L	-2.479	0.024	16	IPL.L	-4.386	4.12E-05
	SFGdor.R	8.724	4.59E-17		IPL.R	-3.726	5.33E-04
4	MFG.L	-3.32	0.01		MFG.L	-6.352	2.58E-09
5	IPL.R	2.419	0.032		MFG.R	-4.675	1.22E-05
	MFG.L	7.139	3.06E-11		SFGdor.L	-2.931	0.007
	MFG.R	6.862	1.73E-10		SFGdor.R	-6.661	5.64E-10
	SFGdor.L	3.649	9.31E-04	17	IPL.L	-	7.40E-08

Hemispheric lateralization in dart players

							5.71	
	SFGdor.R	8.793	1.29E-16			SFGdor.L	−4.989	2.74E-06
7	SFGdor.L	−2.798	0.011			SFGdor.R	−2.72	0.014
8	MFG.L	10.47	7.82E-24			IPL.L	8.34	3.71E-16
	MFG.R	5.262	7.12E-07			MFG.R	−2.519	0.017
	SFGdor.R	5.782	5.24E-08		18	SFGdor.L	7.127	3.42E-12
9	IPL.L	10.99	1.15E-25			SFGdor.R	5.032	9.95E-07
	SFGdor.L	−3.241	0.003			IPL.L	7.062	2.39E-11
10	MFG.L	4.3	6.68E-05		19	MFG.R	−3.826	3.31E-04
	MFG.R	2.867	0.011			IPL.L	3.576	0.001
	SFGdor.L	5.681	9.16E-08			IPL.R	4.542	2.56E-05
11	MFG.L	−4.033	2.33E-04		20	MFG.L	9.534	1.23E-19
	SFGdor.L	−2.915	0.009			MFG.R	3.608	0.001
	SFGdor.R	−2.952	0.008			SFGdor.L	4.643	1.73E-05

	MFG.R	-2.837	0.013		SFGdor.R	5.79	9.68E-08
12	SFGdor.R	-2.926	0.01		MFG.L	-3.354	0.003
				21	MFG.R	-2.275	0.044
					SFGdor.R	-2.226	0.049

704 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
705 gyrus; R: right; L: Left.

706 **Table 8.** Task-related activation in the region of interest revealed by group-level analysis (younger
707 group).

Condition	Anatomical Region	t-value	p-value (FDR-corrected)
planning	IPL.L	0.15	0.909
	MFG.L	1.931	0.186
	MFG.R	3.321	0.022
	SFGdor.L	3.155	0.022
	SFGdor.R	3.822	0.012
calculation	IPL.L	0.139	0.454
	MFG.L	3.321	0.022
	MFG.R	3.321	0.022

	SFGdor.L	2.866	0.052
	SFGdor.R	5.847	5.72E-05

708 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
709 gyrus; R: right; L: Left.

710 **Table 9.** Task-related activation in the region of interest revealed by group-level analysis (expert
711 older group).

Condition	Anatomical Region	t-value	p-value (FDR-corrected)
planning	IPL.L	0.127	0.988
	IPL.R	1.706	0.548
	MFG.L	-0.163	0.988
	MFG.R	1.238	0.709
	SFGdor.L	0.343	0.988
	SFGdor.R	1.04	0.817
calculation	IPL.L	0.545	0.723
	IPL.R	0.367	0.82
	MFG.L	0.999	0.495
	MFG.R	1.542	0.289
	SFGdor.L	1.115	0.272
	SFGdor.R	0.989	0.495

712 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
713 gyrus; R: right; L: Left.

714 **Table 10.** Task-related activation in the region of interest revealed by group-level analysis (non-
715 expert older group).

Condition	Anatomical Region	t-value	p-value (FDR-corrected)
planning	IPL.L	2.282	0.385
	MFG.L	0.446	0.943
	MFG.R	-0.928	0.794
	SFGdor.L	-0.331	0.943
	SFGdor.R	-0.684	0.904
calculation	IPL.L	1.234	0.525
	MFG.L	0.309	0.876
	MFG.R	-1.118	0.54
	SFGdor.L	-0.496	0.803
	SFGdor.R	-0.469	0.815

716 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
717 gyrus; R: right; L: Left.

(a)

Recorded voice prompts from the experimental computer

Beep cue to start throwing a dart

(b)

Dartboard

Fixation point

Button response device

fNIRS system

Scoring sheet

(a)

Log₁₀ sensitivity (mm⁻¹)

-2.0 -1.8 -1.6 -1.4 -1.2 -1.0 -0.8 -0.6 -0.4 -0.2

(b)

Front

L   R

Back

Detector

Source

Short Distance Detector

30mm

8mm

Accelerometer

● Source
● Detector
● Short distance detector
● Accelerometer

(a) HbO: younger group
SFGdor.R   SFGdor.R
MFG.R   MFG.R
HbO β-value
-200   200

(b) HbO: expert older group
SFGdor.R   SFGdor.R
MFG.R   MFG.R
HbO β-value
-300   300

(c) HbO: non-expert older group
SFGdor.R   SFGdor.R
MFG.R   MFG.R
HbO β-value
-500   500

(d)
LI value
2.00
1.50
1.00
0.50
0.00
-0.50

younger group   expert older group   non-expert older group

(a) HbO: younger group

(b) HbO: expert older group

(c) HbO: non-expert older group

(d)

(a) planning
(b) calculation

(a) planning

(b) calculation