Hemispheric lateralization in older adults who habitually play darts: a cross-sectional study using functional near-infrared spectroscopy.

Koki Toyofuku, ktoyofuku@mis.doshisha.ac.jp, Graduate School of Life and Medical Sciences, Doshisha University, Kyoto, Japan.

Satoru Hiwa[*]**,** shiwa@mail.doshisha.ac.jp, Department of Biomedical Sciences and Informatics, Doshisha University, Kyoto, Japan. ORCID: 0000-0001-6547-626X.

Kensuke Tanioka, ktanioka@mail.doshisha.ac.jp, Department of Biomedical Sciences and Informatics, Doshisha University, Kyoto, Japan. ORCID: 0000-0001-9621-5871.

Tomoyuki Hiroyasu, tomo@is.doshisha.ac.jp, Department of Biomedical Sciences and Informatics, Doshisha University, Kyoto, Japan. ORCID: 0000-0003-0730-7905.

Masaki Takeda, mtakeda@mail.doshisha.ac.jp, Faculty of Health and Sports Science, Doshisha University, Kyoto, Japan.

* Correspondence: Satoru Hiwa.

Abstract

Purpose: Exercise training is one of the most effective ways of preventing dementia. This study focuses on exercise training using a dart game called Wellness Darts. Wellness Darts is a sport involving not only throwing darts but also walking to pull them out of the board, memorizing the score, and subtracting it from the total score, thus allowing the performance of two tasks simultaneously: exercise and calculation. It is expected to prevent dementia, and of great interest, whether the continual dart training affects brain function. Before conducting the longitudinal study revealing the Wellness Darts-intervention effect on brain function, here we aimed to cross-sectionally confirm the difference in hemispheric lateralization between expert and non-expert players.

Methods: Functional near-infrared spectroscopy (fNIRS) was used to measure brain activity for three groups: an expert older group who practiced darts continually, a non-expert older control group, and a non-expert younger control group. Their brain activity patterns were quantified by lateralization index (LI) and compared between groups.

Results: The results showed that the younger and the expert older groups had significantly higher LI values than the non-expert older group and that these values did not differ between the expert older and the younger groups.

Conclusion: Our results suggest that the Wellness Darts possibly promote hemispheric lateralization.

Keywords: exercise training, darts, functional near-infrared spectroscopy, cognitive function, hemispheric lateralization

Data availability statement: The datasets generated during and/or analysed during the current study
are available from the corresponding author on reasonable request.

Funding statement: This work was supported by JSPS KAKENHI, Grant Number JP20K11403.

Conflict of interest disclosure: The authors declare that the research was conducted in the absence
of any commercial or financial relationships that could be construed as potential conflicts of interest.

Ethics statement: Studies involving human participants were reviewed and approved by the
Research Ethics Committee of Doshisha University, Kyoto, Japan (approval code: 20014).

Patient consent statement: All participants were informed about the experimental method as well as
the risks and signed written informed consent forms to participate in the study.

Author contribution statement: KoT, SH, TH, and MT contributed to conception and design of the
study. KoT and SH wrote the manuscript. KoT, SH, and MT conducted experiments. KoT performed
data analysis. SH, KeT, TH, and MT advised on the data analysis and experimental design. All
authors contributed to manuscript revision, read, and approved the submitted version.

Acknowledgments: We are deeply grateful to the members of the Kyotanabe Doshisha Sports Club
(KDSC) for their cooperation in the experiment. We would like to thank Editage (www.editage.com)
for English language editing.

1   Introduction

Dementia is a general term for symptoms in which cognitive function declines more than expected
due to biological aging. Also, mild cognitive impairment (MCI) is an intermediate between normal
cognitive aging and dementia (Lissek and Suchan, 2021). In the MCI stage, the patients do not
interfere with their everyday activities. That is why early intervention at the MCI stage is crucial to
prevent dementia.

Prevention of dementia requires, among other things, the management of hypertension, i.e., the
identification and control of risk factors for stroke and cardiovascular disease (Hanon et al., 2004;
Román, 2005). However, although these measures can prevent hypertension-related dementia, they
are ineffective in preventing dementia caused by other factors. Physical inactivity, diabetes mellitus,
smoking, and depressive symptoms are also associated with the risk of developing Alzheimer's
disease, the most common form of dementia. Among these, physical inactivity is one of the most
preventable risk factors for Alzheimer's disease and higher levels of physical activity have been
shown to reduce the risk of developing Alzheimer's disease (Meng et al., 2020). Therefore,
promoting physical activity to prevent dementia has received much attention. Physical activity
associated with exercise training has been shown to cause significant changes in brain health and
cognitive performance, with potential effects on a wide range of cognitive functions, including
memory, attention, and executive function (McGough et al., 2014). It has also been reported that
executive functioning improves after low-intensity exercise, with increased activity in the
dorsolateral prefrontal cortex (DLPFC) and frontal pole, brain regions associated with executive
function (Byun et al., 2014).

This study focused on exercise training using a darts game called Wellness Darts. Darts is one of the
sports that seek fun and gameplay by subtracting the score obtained by the player from the points he
or she has and reducing it to zero. Wellness Darts also has the unique feature of calculating scores by
rote (Takeda et al., 2017); patients with MCI and Alzheimer's disease are said to perform poorly on
calculation tasks, especially subtraction and multiplication, and show this tendency from the MCI
stage onwards (Martin., 2003). Wellness Darts is a cognitively challenging sport that involves not
only throwing darts, but also walking to pull them out of the board, memorizing the score, and
subtracting it from the total score, thus allowing the performance of two tasks at the same time:
exercise and calculation (Takeda et al., 2017). Simultaneous performance of the motor and cognitive
tasks, each effective in improving cognitive function as a single task, is more effective in increasing
PFC activity and improving cognitive function than performing either task alone (Hermand et al.,
2019). Takeda et al. conducted cognitive tests before, one month, and two months after an
experiment on a Wellness Darts intervention group and a control group, consisting of older adults
(Takeda et al., 2017). The results showed that the continuation of Wellness Darts increased older
adults' short-term memory test (STM test) scores. However, the impact of continual practice of
Wellness Darts on brain function is unclear.

This study aims to investigate the current cognitive status and brain function of participants who
practiced Wellness Darts in older adults. In particular, their characteristics of brain function and the
performance of cognitive tests will be compared cross-sectionally among older adults who are
proficient with darts, older adults who are inexperienced with darts, and younger adults who are
inexperienced with darts. The Hemispheric Asymmetry Reduction in OLDer adults (HAROLD)
model proposed by Cabeza et al. asserts that the degree of Frontal lobe (FL) lateralization seen in
youth decreases with age (Cabeza et al., 2002; Cabeza et al., 2000; Müller et al., 2014; Erickson et
al., 2007; Cabeza et al., 2002; Reuter-Lorenz et al., 2000; Blum et al., 2021; Vermeij et al., 2014). In

addition, Erickson et al. reported that when cognitive training was conducted on healthy older adults,
an increase in lateralization was observed together with the improvement of cognitive performance,
bringing the brain activation pattern closer to that of younger adults (Erickson et al., 2007). That is
why increased lateralization can be potentially used as a measure of cognitive improvement. On the
other hand, brain regions such as the inferior frontal gyrus (IFG), superior frontal gyrus (SFG),
middle frontal gyrus (MFG), (Verner et al., 2013), and DLPFC have been reported to be activated
during mental calculation (Pfurtscheller et al., 2010). Furthermore, the inferior parietal lobule (IPL)
in patients with cognitive decline such as MCI and Alzheimer's disease is reported to be less
activated (Schroeter et al., 2009).

Based on these findings, we proposed the following three hypotheses: (1) older adults who practice
Wellness Darts (referred to as the 'expert older group') will show a level of hemispheric
lateralization similar to that of younger people who had never played Wellness Darts before (referred
to as 'younger group' as a control), with activation of the dorsal superior frontal gyrus (SFGdor) and
MFG, which are known to be activated during mental arithmetic. (2) There is no difference in the
level of IPL activation between the expert older group and the younger group. (3) There is a positive
correlation between the level of hemispheric lateralization in brain activity and the number of days of
Wellness Dart experience in the expert older group. These three hypotheses were tested by measuring
brain activity using functional near-infrared spectroscopy (fNIRS), a non-invasive functional brain
imaging system robust to body movement. The third group, the older adults who had never played
Wellness Darts (referred to as the 'non-expert older group'), was used as the older control to confirm
that hypotheses (1) and (2) were valid only for the expert older group.

2   Materials and Methods

2.1   Participants

Twenty-one healthy younger adults (13 males, 7 females; mean age: 22.5 ± 0.8 years) who had never
played Wellness Darts, 21 healthy older adults (8 males, 13 females; mean age: 75.1 ± 4.3 years) who
continually played Wellness Darts, and 21 healthy older adults (4 males, 17 females; mean age: 74.1
± 4.5 years) who had never played Wellness Darts participated in the experiment. These three
participant groups were referred to as the younger group, the expert older group, and the non-expert
older group, respectively. The participants for the expert older group were recruited from Wellness
Darts players in the Kyotanabe-Doshisha Sports Club (KDSC), one of the community sporting clubs
in Kyoto, Japan. The participants for the non-expert group were also recruited based on the
acquaintanceship of the member of the expert older group. Only healthy participants aged over 60
could participate in this study for these two groups. For the younger group, the healthy undergraduate
and graduate school students aged over 20 were recruited from Doshisha University, Kyoto, Japan.
The sample size for each group was determined with a significance level of 0.05 and a power of 0.8
to detect the effect size of Cohen's $d = 0.79$ reported in a previous study comparing the degree of
hemispheric lateralization in the elderly using fNIRS (Yeung et al., 2020) (G*Power was used for the
calculation).

2.2   Wellness Darts task

Wellness Darts is a darts game developed by the Japan Wellness Darts Association. The standard
game of Wellness Darts is the "Zero One Game 251". In the game, each player is assigned a score of
251 and throws darts until the score reaches zero, subtracting the score of each dart thrown from the
holding points. The first player to reach a score of exactly 0 wins. Each player can throw darts three
times in their turn (which is defined as one set) and is required to make his/her score exactly zero on

the third throw for either of five sets. In addition, each player must subtract the number of points
earned from their score by themselves. We assumed that these mental arithmetic operations, the
planning of throws to reach zero as quickly as possible, and the exercise associated with throwing
darts could be effective in improving cognitive function because the Simultaneous performance of
the motor and cognitive tasks said to be effective in increasing PFC activity and improving cognitive
function (Hermand et al., 2019). However, since the official rules of Wellness Darts place a physical
burden on the participants due to the long duration of the experiment, in this study, the initial score of
the subjects was set at 128 points, and each subject played the game for a maximum of five sets.
Furthermore, the game was terminated even if the score did not reach zero in the fifth set.

152 2.3 Procedure

Participants played up to five sets of Wellness Darts with a maximum of 128 holding points, while
brain activity and head acceleration were simultaneously recorded. The accelerations were used to
remove noise due to head movements, as described below. In each of the three throws in one set of
the Wellness Darts game, the participants were instructed by the recorded voice prompts and the beep
cue from the experimental computer on when to hold the dart, throw, and start the calculation; the
participants performed each action following the instructions.

During the planning block that was set at the beginning of each throwing session, participants were
asked to determine in their mind where to hit (designate scoring points) within 5 s for the first throw
and within 13 s for the remaining two throws. At the beginning of the second and third throwing
sessions, they were provided an extra 8 s in the planning blocks to tally the points from the previous
throws. Of note, we excluded the beginning 8 s and instead applied the later 5 s for the subsequent
analyses because we assumed that the first throw did not involve the confirmation of the hit point and
employed different cognitive processes than those used in determining where to hit. The planning
block was followed by dart-throwing within 3 s after the beep cue.

Following dart throwing, the participants were instructed to move their gaze away from the dartboard
to the fixation point and not to think of anything in particular. This waiting block consisted of
random duration between 28 - 32 s to reduce the anticipatory response. After completing the three
throwing sessions, the participants commenced the calculation block where they were instructed to
subtract the scores they received during the previous throws from the total score and write the
remaining points on the scoring sheet. Then, they were asked to press a button to proceed to the 20 s
rest block. During the rest block, they stayed the same as in the waiting block. This procedure was
repeated for five sets. The overall procedure is illustrated in Figure 1.

All participants were informed of the methods and risks of the experiment and signed a written
informed consent form. This study was conducted in accordance with the Research Ethics Committee
of Doshisha University (Kyoto, Japan) (approval code: 20014). Before the experiment, participants
were asked to complete a questionnaire, a dart-throwing performance test, and five cognitive tests:
Simple Cognitive (SC) test, Computational Ability Test, Short-Term Memory (STM) Test, Trail
Making Peg Test (TMPT), and CogEvo. Participants then performed one set of dart-throwing before
being fitted with the fNIRS device in order to understand the experimental procedure.

182 2.4 fNIRS data acquisition

Changes in the concentration of oxy-hemoglobin (HbO), deoxy-hemoglobin (HbR), and total-
hemoglobin (HbT) in the participants were measured using NIRSport2 (NIRx medical technologies,
LLC). The sampling frequency was 4.36 Hz, and the wavelengths of the near-infrared light were 760

and 850 nm, respectively (Pereira et al., 2007). As shown in Figure 2 (b), 16 sources and 14 detectors
were placed on the frontal, parietal, and temporal regions of the head. A pair of sources and detectors
adjacent to each other at 30-mm intervals constituted one standard channel, and 44 standard channels
were placed. In addition to these, a short-distance detector bundle (NIRx medical technologies, LLC)
was placed to eliminate extra-cerebral blood flow alterations (Huppert et al., 2009; Tachtsidis et al.,
2016). The signals obtained from the 16 short-distance channels were used to detect extracerebral
signals and to remove the effect of skin blood flow in the signals measured by the standard channels.
To remove noise from body motion, a six-axis accelerometer (NIRx medical technologies, LLC) was
used to measure head acceleration at a sampling frequency of 4.36 Hz, the same frequency as the
change in hemoglobin concentration (Lu☐hmann et al., 2020). The accelerometer was placed at the
top of the head as shown in Figure 2 (b). Presentation software (Neurobehavioral system Inc.) was
used to control the presentation and timing of the recorded voice prompts and the beep cue during the
experiment, and the signal measurement and execution of the experimental task were synchronized.
The 3D coordinates of the light source and detection probe were measured using a 3D magnetic
space digitizer (Fastrak, Polhemus, USA) to estimate the exact measurement position of the brain
area for each individual. The measured 3D coordinates were entered into the 'register2polhemus'
function and the 'depthmap' function implemented in the NIRS toolbox (Santosa et al., 2018), and the
distance from each standard channel to the brain surface was calculated. On the basis of this distance,
the regions of the brain defined by the automated anatomical labeling atlas (Tzourio-Mazoyer et al.,
2002) corresponding to each standard channel were identified.

2.5   Cognitive tests

Five cognitive tests were conducted to identify differences in cognitive function between the three
participant groups. As the current study is a cross-sectional comparison, it is impossible to show
whether the continued practice of Wellness Darts in older adults affects cognitive function; however,
it was preliminarily investigated to confirm its possibility. That is why the statistical tests were
conducted between two groups: the expert older and the non-expert older groups.

2.5.1 Simple Cognitive test

The Simple Cognitive (SC) test was used to verify that the participant's cognitive function had not
declined (Nagata et al., 2010). The SC test is a 50-point cognitive function test that can identify early
cognitive decline that cannot be detected by the mini-mental state examination (MMSE), which is
often used to examine dementia. A score of fewer than 20 points on the SC test is considered
suspicious of cognitive decline. In this study, only participants with a score of 20 or higher were
considered to have no cognitive decline and were used in subsequent analyses.

2.5.2 Computational ability test

The participants' computational abilities were measured by administering a test consisting of 30
subtraction problems between two digits, between three and two digits, and between three digits.
Differences in mean scores on the computational ability test between the expert older group and the
non-expert older group were t-tested ($p < .05$).

2.5.3 STM

To evaluate the short-term memory of the participants, the Short-Term Memory (STM) Test was
conducted on a computer (Takeda et al., 2017). After the instruction "Please memorize the numbers
in order" was displayed on the monitor, nine disordered numbers were displayed on the screen one by
one per second. The participants were then instructed to "write the numbers in order," and they wrote

229 the memorized numbers on a piece of paper during a 20-s time frame. The difference in mean STM
230 test scores between the expert older group and the non-expert older group was *t*-tested ($p < .05$).

2.5.4 CogEvo

232 CogEvo was used to test the cognitive abilities of the participants; CogEvo is a computer-assisted
233 cognitive function test that assesses disorientation, attention, memory, planning, and spatial
234 awareness (Ichii et al., 2020). CogEvo can reportedly capture mild changes in cognitive function and
235 is a simple and convenient ICT tool to assess cognitive changes over time in middle-aged and older
236 adults, as well as during the preclinical stages of dementia (Ichii et al., 2020; Takechi et al., 2021).
237 CogEvo has 12 different tasks; in this experiment, we selected flashlight, visual search, number step,
238 and just fit as tasks related to Wellness Darts from the disorientation, attention, memory, planning,
239 and spatial awareness components of CogEvo, respectively. The difference in mean CogEvo scores
240 between the expert older group and the non-expert older group was *t*-tested ($p < .05$).

2.5.5 TMPT

242 The Trail making peg test (TMPT) was conducted to assess participants' cognitive functioning,
243 particularly attentional function and finger dexterity (Yoon et al., 2020). The TMPT is a peg-
244 movement test that reflects finger dexterity combined with the trail-making test that measures visual
245 attention and executive function (Llinàs-Reglà et al., 2017). In the TMPT, participants quickly insert
246 25 cylindrical wooden sticks, called pegs, into numbered holes in numerical order. The board
247 containing the pegs and the board containing the holes were separated from each other. The
248 difference in mean completion time was *t*-tested ($p < .05$).

2.6 Dart throwing performance test

250 To confirm that the dart-throwing skill of the expert older group was higher than that of the control
251 groups, the dart-throwing performances of the three groups were tested. In this test, a target score of
252 65 points was set and participants were required to throw the darts 10 times and tally their scores to
253 match the target score. The differences in total dart performance test scores between the expert and
254 younger groups and between the expert and non-expert older groups were t-tested ($p < .05$).

2.7 Questionnaires

256 Participants were asked to respond to questions regarding their dominant hand, the number of days
257 per week they practiced Wellness Darts, and the number of years they had been practicing. From
258 these responses, the total number of days of experience for each participant was estimated.

2.8 fNIRS data processing

2.8.1 Preprocessing

261 Previous studies have concluded that HbO in fNIRS is more strongly related to blood oxygenation
262 level-dependent (BOLD) signals measured by functional magnetic resonance imaging (fMRI) than
263 HbR (Strangmen et al., 2002; Cui et al., 2011; Huppert et al., 2006). The fMRI has better spatial
264 resolution than fNIRS and has been used in many cognitive neuroscience studies. In this study, we
265 used HbO to analyze brain activity. The fNIRS data were preprocessed using the NIRS toolbox
266 (Santosa et al., 2018) to remove physiological noise such as skin blood flow, heart rate, and pulse
267 rate. Optical intensity data acquired from the fNIRS system were converted to optical density (OD)
268 data and then to Hb concentration data based on the modified Beer-Lambert law (Huppert et al.,

269   2009). A bandpass filter (0.008-0.09 Hz) was applied to remove noise such as low-frequency drift
270   and heartbeat (Huppert et al., 2006; Sasai et al., 2012).

2.8.2 Activation analysis

The level of task-induced brain activation for each individual was estimated using the iteratively
reweighted least-squares model (AR-IRLS) with short separation (SS) and acceleration, a general
linear model (GLM)-based analysis method (Lu□hmann et al., 2020; Huo et al., 2021). In GLM,
task-related brain activation was modeled by the hemodynamic response function (HRF). The
measured brain activity, i.e., HbO signal, was regressed on the ideal task-related activity calculated
from the canonical HRF and a design matrix indicating the type of experimental task, and the partial
regression coefficients were estimated by IRLS, a least-squares method. The AR-IRLS allows brain
activation analysis to consider the effects of arbitrary confounding factors such as the SS signal
measured with a short-distance channel, blood pressure, respiration, and acceleration on brain activity.
In this study, we introduced the SS signal and acceleration data as physiological regressors of GLM
to reduce the effects of skin blood flow and body motion on the HbO signal (Lu□hmann et al., 2020;
Santosa et al., 2020).

2.8.3 Hemispheric lateralization analysis

The level of hemispheric lateralization was calculated from the lateralization index (LI) shown in
Equation 1.

$$LI = \frac{|AL - AR|}{|AL| + |AR|} \qquad (1)$$

where AL is the sum of β values in the regions of interest (ROIs) in the left hemisphere and AR is the
sum of those in the right hemisphere (Seghier et al., 2008, Watanabe et al., 1998). We used absolute
values for the LI values to allow for the fact that the dominant hemisphere differs depending on the
dominant hand.

2.9   Statistical analysis

2.9.1 Activation analysis

For individual-level analysis, the following two contrasts were created for the HbO data of the
younger group, the expert older group, and the non-expert older group: "rest vs. calculation" and "rest
vs. planning". The β-values for each experimental condition and each channel were calculated. The
statistical significance of the task-related brain activity was then verified by testing the null
hypothesis that the estimated β coefficient was not significantly different from zero (*t*-test, $p < .05$).
Note that multiple comparisons were corrected using the Benjamini-Hochberg method (Santosa et al.,
2020) . In addition, to test the hypothesis that there is no difference in the level of IPL activation
between older adults who continually practiced Wellness Darts and younger adults who were not
experienced in Wellness Darts, β values of IPL were compared between the three groups (*t*-test, *p*
$< .016$, Bonferroni corrected).

Next, for the group-level analysis, we constructed the most consistent contrast images of group-level
brain activity using simultaneously the β values obtained from the individual-level analysis of all
participants, based on Santosa et al. (2018). To estimate significant group-level activity, we used
participants as random effects and conditions (planning, calculation, and rest blocks) as fixed effects.
Finally, all participants were considered as a sample drawn from the population and a one-sample t-
test was used to test if the population mean was greater than zero with respect to the *t*-value. The

significance level was set at $p < .05$ (false discovery rate corrected at peak level), and the brain
regions activated during the calculation and planning blocks were estimated.

2.9.2 Hemispheric lateralization analysis

The LI values were used to compare the degree of hemispheric lateralization among the three groups
of participants (t-test, $p < .016$, Bonferroni corrected). In addition, we tested whether there was a
positive correlation between the number of days of Wellness Dart experience and LI values in the
expert older group (Spearmans' rank correlation test, $p < .05$).

3   Results

3.1   Descriptive statistics on three groups

The descriptive statistics on three groups, including age, days of practicing wellness darts, cognitive
test scores, and scores of dart-throwing performance tests, are summarized in Table 1. The following
subsections describe the details of the results including statistical tests.

3.2   fNIRS data

3.2.1 Hemispheric lateralization analysis

We hypothesized that the LI values of the younger group and the expert older group would be larger
than those of the non-expert older group. LI values were calculated from the respective β values of
the two task blocks (planning and calculation) for the three participant groups. The LI values in the
planning and calculation blocks of the younger group were 0.45 ± 0.34 and 0.49 ± 0.35, respectively.
The LI values in the planning and calculation blocks of the expert older group were 0.53 ± 0.38 and
0.61 ± 0.35, respectively. Finally, for the non-expert older group, they were 0.23 ± 0.18 and 0.50 ± 0.38
in the planning and calculation blocks, respectively. The box plots of the LI values in the planning
and calculation blocks for the three groups of participants are shown in Figure 3 (d) and Figure 4 (d),
respectively. Furthermore, the β maps of the participants who showed the closest values to the
median in the LI values for each group of participants are shown in Figure 3 (a)-(c) and Figure 4 (a)-
(c).

There were no significant differences between the LI values of the younger group and the expert
older group in the planning block ($t = 0.6730$, $p = 0.504$, $d = 0.213$). On the other hand, in the
planning block, the LI values of the expert older group were significantly greater than those of the
non-expert older group ($t = 3.188$, $p = 0.003$, $d = 1.005$). Furthermore, the LI values of the younger
group were significantly greater than those of the non-expert older group in the planning block ($t =$
2.58, $p = 0.014$, $d = 0.815$).

In the calculation block, there were no differences in LI values between the younger group and the
expert older group ($t = 1.009$, $p = 0.319$, $d = 0.319$). Similarly, there were no differences in LI
between the expert older group and the non-expert older group ($t = 0.828$, $p = 0.413$, $d = 0.262$).
There were also no differences in LI values between the younger group and the non-expert older
group ($t = 0.144$, $p = 0.886$, $d = 0.046$).

We also hypothesized that the number of days of Wellness Dart experience in the expert older group
would be positively correlated with LI values. The results of an Spearman's rank correlation test

between the number of days of Wellness Dart experience and LI values showed a negative significant
correlation for the planning block (ρ = -0.439, p = 0.046, Figure 5 (a)) but no significant correlation
for the calculation block (ρ = -0.037, p = 0.874, Figure 5 (b)).

To investigate the Wellness Dart experience of the expert older group in terms of skill, the mean
darts performance test score (target score was 65 points; the closer the total score of 10 dart throws
was to 65 points, the better the skill) was compared between the three participant groups. The mean
scores for the younger group, the expert older group, and the non-expert older group were 39.3±13.4,
49.9±5.5, and 31.9±13.7, respectively. There were no participants who exceeded the 65 points. As
shown in Figure 6, the expert older group showed better dart performance compared to the younger
group (t = 3.247, p = 0.003, d = 1.022). Similarly, the expert older group showed better performance
than the non-expert older group (t = 5.400, p = 0.00001, d = 1.700).

Furthermore, inspired by the results of no significant 'positive' correlation between the number of
days of Wellness Dart experience and LI values, we assumed that dart proficiency could be assessed
not only by the number of days of experience but also by dart-throwing performance. Therefore, a
positive correlation between the total dart performance test score and the degree of hemispheric
lateralization of brain activity in the expert older group was also tested (Spearman's rank correlation
test, p < .05). It should be noted that this is an exploratory analysis based on the result shown in
Figure 5. However, no significant correlation between the total dart performance test score and the LI
value was confirmed (planning block: ρ = 0.324, p = 0.152, Figure 7 (a); calculation block: ρ = -
0.225, p = 0.327, Figure 7 (b)).

3.2.2 Activation analysis

The activation analysis shown in Figure 8 indicates that in the planning block, there was no
difference between the left and right IPL (IPL.L: t = 0.738, p = 0.466, d = 0.234. IPL.R: t = 0.134, p
= 0.894, d = 0.043). Similarly, there was no difference between the expert older group and the non-
expert older group (IPL.L: t = 0.042, p = 0.966, d = 0.014; IPL.R: t = 0.821, p = 0.419, d = 0.257).
Furthermore, there was no difference between the younger group and the non-expert older group
(IPL.L: t = 0.711, p = 0.481, d = 0.226. IPL.R: t = 0.967, p = 0.343, d = 0.312).

In the calculation block, there was no difference between the left and right IPL (IPL.L and IPL.R) of
the younger group and the expert older group (IPL.L: t = 1.211, p = 0.236, d = 0.381; IPL.R: t =
0.206, p = 0.838, d = 0.068). Similarly, there was no difference between the expert older group and
the non-expert older group (IPL.L: t = 1.625, p = 0.115, d = 0.513; IPL.R: t = 1.642, p = 0.109, d =
0.523). There was no difference between the younger group and the non-expert older group (IPL.L: t
= 2.176, p = 0.036, d = 0.688. IPL.R: t = 1.844, p = 0.074, d = 0.621).

Tables 2 to 7 summarize the results of individual-level analysis and show the regions of interest
where the p-values are less than 0.05. Tables 8 to 10 summarize the results of group-level analysis
and show only the ROIs. There was significant activation in the ROIs in the individual-level analysis
in the expert older group but no significant activation in the group-level analysis.

3.3   Cognitive tests

3.3.1 Computational ability test

The mean computational test scores for the younger group, the expert older group, and the non-expert
older group were 24.0 ± 3.8, 20.5 ± 3.8, and 17.1 ± 5.5, respectively. As shown in Figure 9, the

expert older group had higher scores on the computational skills test than the non-expert older group
($t = 2.275$, $p = 0.029$, $d = 0.719$).

3.3.2 STM test

393   The mean STM test scores of the younger group, the expert older group, and the non-expert older
394   group were 74.6 ± 13.6, 57.1 ± 9.5, and 49.4 ± 9.9, respectively. As shown in Figure 10, the expert
395   older group showed higher STM test scores than the non-expert older group ($t = 2.475$, $p = 0.018$, $d =$
396   0.792).

3.3.3 CogEvo

398   The scores of the five CogEvo tasks for each group of participants are summarized in Figure 11. In
399   the expert older and non-expert older groups, there were no significant differences in the CogEvo
400   scores (disorientation: $t = 1.892$, $p = 0.067$, $d = 0.598$, visual search: $t = 1.435$, $p = 0.159$, $d = 0.454$;
401   flashlight: $t = 1.386$, $p = 0.174$, $d = 0.438$; number step: $t = 1.722$, $p = 0.093$, $d = 0.550$; just fit: $t =$
402   0.625, $p = 0.534$, $d = 0.198$).

3.3.4 TMPT

404   The mean TMPT completion times for the younger, expert older, and non-expert older groups were
405   41.1 ± 4.5, 58.1 ± 8.5, and 58.9 ± 9.2, respectively. There was no significant difference in TMPT
406   completion time between the older and non-expert older groups ($t = 0.272$, $p = 0.787$, $d = 0.086$).

4   Discussion

4.1   Hypothesis 1

As seen in the results of the individual-level analysis in Tables 2-7, we were able to confirm the
activation of SFGdor and MFG in the three participant groups. However, there was no significant
activation of these regions in the group-level analysis. This may be because the mean activation level
was estimated to be lower after averaging the results of the group as the individual activated areas
varied even in the same region due to the distinct differences in the arrangement of measurement
channels in the fNIRS device.

On the other hand, Figure 3 (d) shows that the LI values of the younger group and the expert older
group were significantly higher than those of the non-expert older group in the planning block. In
addition, there was no difference in LI values between the younger group and the expert older group.
This indicates that the brain activation patterns of participants in the younger group and the expert
older group are hemispherically lateralized during the planning of dart-throwing and while
determining where to hit. The β-map in Figure 3 (a)-(c) also shows that the activation levels in the
younger and expert older groups were higher in one hemisphere, while the non-expert older group
showed no hemispheric difference. In the non-expert older group, there is no difference between the
hemispheres, and bilateral activation levels of SFGdor and MFG are high. This result is consistent
with a previous study in which left-right hemispherical asymmetry increased when cognitive training
was performed in healthy older adults (Erickson et al., 2007). In summary, although SFGdor and
MFG were not significantly activated in the expert older group in group-level analysis, Hypothesis 1
was established during the planning of dart throwing, suggesting that continual Wellness Dart
training promotes hemispheric lateralization in brain activity in older adults.

4.2   Hypothesis 2

Figure 8 indicates that there were no significant differences in the β-values of the left and right IPL
between the younger group and the expert older group. Further, there was no significant difference in
the β-values of the left and right IPL between the younger and non-expert older groups, and between
the expert and non-expert older groups. However, in the three participant groups in the present study,
the IPL activation was comparable, suggesting that there was no difference in IPL activation between
the older groups. The previous study to which the hypothesis refers involved healthy subjects and
patients with MCI and Alzheimer's disease (Schroeter et al., 2009). The participants in the present
study were only healthy adults, suggesting that the difference in cognitive function between the older
groups and the younger group did not affect IPL activation.

4.3   Hypothesis 3

As shown in Figure 5, there was no positive correlation between the number of days of Wellness Dart
experience and the LI value of the expert older group; therefore, hypothesis 3 was not confirmed. On
the other hand, in Figure 6, the dart performance of the expert older group was higher than that of the
other two groups. In addition, Figure 7 implies a positive correlation between dart performance and
LI value in the planning block, although it is not statistically significant. The total score of our dart
performance test indicates the accuracy of dart throwing, which is improved by the number of days of
experience. Therefore, the results of Figures 6 and 7 may contribute to the validity of hypothesis 3.
Also, since the current results suggested weak correlations, the correlations should be tested again
with a larger sample size. These issues would be investigated in future longitudinal controlled studies
of the Wellness Darts intervention that measure dart performance levels to verify the increase in LI
values promoted by Wellness Dart training.

4.4   Cognitive tests

In addition to hypotheses testing, the results of three cognitive function tests to determine the impact
of Wellness Darts on cognitive function are also discussed.

4.4.1 Computational ability test

The score of the expert older group was significantly higher than that of the non-expert older group.
In other words, the expert older group had better computational ability than the non-expert older
group. According to previous studies, cognitive function, particularly for high-load tasks, declines as
the brain deteriorates with age (Bierre et al., 2017). Therefore, the results of the computational ability
test in this study suggest that continual practice of Wellness Darts can reduce the decline of
calculation ability due to aging.

4.4.2 STM test

The STM test scores of the expert older group were higher than that of the non-expert older group,
and the difference was significant. Fandakova et al. (2014) have shown that short-term memory
capacity declines with age. Therefore, our results indicate that constant training with Wellness Darts
can reduce the age-related decline in short-term memory capacity.

4.4.3 CogEvo

No significant differences were found in the scores between the expert older group and the non-
expert older group on the five CogEvo tasks. Although CogEvo has been reported to be able to
significantly distinguish between Alzheimer's disease, MCI, and cognitively normal groups of older
adults (Ichii et al., 2020; Takechi et al., 2021), our results showed that the expert older group and the
non-expert older group did not differ significantly in their scores on the five CogEvo tasks. In fact,

our results indicated that the expert older and non-expert older groups had similar cognitive levels. In
other words, the older groups in our experiment had normal levels of cognitive function or the
differences in cognitive function were at levels that were undetectable by CogEvo. Another
explanation could be that the older groups were unfamiliar with the operation of CogEvo as it was
conducted using a tablet device.

4.4.4 TMPT

As seen in Figure 12, there was no significant difference in the completion time of TMPT between
the expert older and the non-expert older groups. TMPT requires skill from the participant as a
component of physical function. We expected that participants engaged in ongoing Wellness Darts
exercises would demonstrate improved skillfulness; however, in this cross-sectional study, no
differences were observed between the expert older and non-expert older groups. A previous
longitudinal study that divided older adults into a dual-task (DT) exercise group and a control group
observed a significant difference in TMPT before and after intervention in the DT exercise group, but
no significant difference was identified in TMPT between the two groups (Yoon et al., 2020). Due to
the pilot nature of the current study based on a cross-sectional comparison, a possible future direction
could be to investigate whether the Wellness Darts intervention contributes to improving the TMPT
completion time in longitudinal randomized controlled trials.

4.5 Strengths of the current study, limitations, and future direction

Based on our findings, only hypothesis 1 was confirmed in the planning of dart throwing and in
determining where to hit. Due to the pilot nature of this study, the sample size was limited, and the
effects of several unseen parameters such as education level, socio-economic factors, body mass
index, and smoking were not controlled and investigated. These limitations make the current study
insufficient to demonstrate that Wellness Darts truly affected brain lateralization. However, despite
these limitations, the current study revealed that older adults with Wellness Dart experience have a
higher tendency toward hemispheric lateralization than those with no experience. The current results
are valuable in motivating us to proceed to future longitudinal randomized controlled trials to explore
the effects of continual Wellness Dart training on hemispheric lateralization with a larger sample size.

5 Conclusions

Wellness Darts is a cognitively challenging sport that combines exercise and calculation. Here we
aimed to cross-sectionally confirm the difference in hemispheric lateralization between expert and
non-expert players as a pilot study prior to the longitudinal randomized controlled trials. The results
showed that the younger and the expert older groups had significantly higher LI values than the non-
expert older group and that these values did not differ between the expert older and the younger
groups. These results suggest that the Wellness Darts possibly promote hemispheric lateralization.

6 References

Lissek V, Suchan B. Preventing dementia? Interventional approaches in mild cognitive impairment.
Neurosci Biobehav Rev. 2021 Mar;122:143-164. doi: 10.1016/j.neubiorev.2020.12.022. Epub 2021
Jan 10. PMID: 33440197.

Hanon O and Forette F. Prevention of dementia: lessons from syst-eur and progress. Journal of the
Neurological Sciences, 226(1):71–74, 2004.

Román G.C. Vascular dementia prevention:a risk factor analysis. Cerebrovasc Diseases, 20(2):91–
100, 2005.

Qing Meng, Muh-Shi Lin, and I-Shiang Tzeng. Relationship between exercise and alzheimer's
disease: A narrative literature review. Frontiers in Neuroscience, 14, 2020.

McGough EL Kirk-Sanchez NJ. Physical exercise and cognitive performance in the elderly: current
perspectives. Clinical Interventions in Aging, 9:51–62, 2014.

Kyeongho Byun, Kazuki Hyodo, Kazuya Suwabe, Genta Ochi, Yosuke Sakairi, Morimasa Kato,
Ippeita Dan, and Hideaki Soya. Positive effect of acute mild exercise on executive function via
arousal-related prefrontal activations: An fnirs study. NeuroImage, 98(Supplement C):336–345, 2014.

Martin RC, Annis SM, Darling LZ, Wadley V, Harrell L, and Marson DC. Loss of calculation
abilities in patients with mild and moderate alzheimer disease. Archives of Neurology, 60(11):1585–
1589, 2003.

Eric Hermand, Bertrand Tapie, Olivier Dupuy, Sarah Fraser, Maxence Compagnat, Jean Yves Salle,
Jean Christophe Daviet, and Anaick Perrochon. Prefrontal cortex activation during dual task with
increasing cognitive load in subacute stroke patients: A pilot study. Frontiers in Aging Neuroscience,
11:160, 2019.

Masaki Takeda, Nao Yasuda, Satsuki Ito, Masato Abe, Yohei Takamichi, Kazuma Yanagisawa, and
Zsolt Radak. Effects of habitual darts training on cognitive function in elderly people. The Harris
Science Review of Doshisha University, 58(2):53–60, 2017.

Roberto Cabeza. Hemispheric asymmetry reduction in older adults: the harold model. Psychology
and aging, 17(1):85, 2002.

Roberto Cabeza, Nicole D Anderson, Sylvain Houle, Jennifer A Mangels, and Lars Nyberg. Age-
related differences in neural activity during item and temporal-order memory retrieval: a positron
emission tomography study. Journal of Cognitive Neuroscience, 12(1):197–206, 2000.

Laura D Müller, Anne Guhn, Julia BM Zeller, Stefanie C Biehl, Thomas Dresler, Tim Hahn, Andreas
J Fallgatter, Thomas Polak, Jürgen Deckert, and Martin J Herrmann. Neural correlates of a
standardized version of the trail making test in young and elderly adults: a functional near-infrared
spectroscopy study. Neuropsychologia, 56:271–279,2014.

Kirk I Erickson, Stanley J Colcombe, Ruchika Wadhwa, Louis Bherer, Matthew S Peterson, Paige E
Scalf, Jennifer S Kim, Maritza Alvarado, and Arthur F Kramer. Training-induced plasticity in older
adults: effects of training on hemispheric asymmetry. Neurobiology of aging, 28(2):272–283, 2007.

Roberto Cabeza, Nicole D Anderson, Jill K Locantore, and Anthony R McIntosh. Aging gracefully:
compensatory brain activity in high-performing older adults. NeuroImage, 17(3):1394–1402, 2002.

Patricia A Reuter-Lorenz, John Jonides, Edward E Smith, Alan Hartley, Andrea Miller, Christina
Marshuetz, and Robert A Koeppe. Age differences in the frontal lateralization of verbal and spatial
working memory revealed by pet. Journal of cognitive neuroscience, 12(1):174–187, 2000.

Leonore Blum, David Rosenbaum, Benjamin Röben, Katja Dehnen, Walter Maetzler, Ulrike Suenkel,
Andreas J Fallgatter, Ann-Christine Ehlis, and Florian G Metzger. Age-related deterioration of
performance and increase of cortex activity comparing time-versus item-controlled fnirs
measurement. Scientific Reports, 11(1):1–13, 2021.

Anouk Vermeij, Arenda HEA van Beek, Babette LR Reijs, Jurgen AHR Claassen, and Roy PC
Kessels. An exploratory study of the effects of spatial working-memory load on prefrontal activation
in low- and high-performing elderly. Frontiers in aging neuroscience, 6:303, 2014.

Martin Verner, Martin J Herrmann, Stefan J Troche, Claudia M Roebers, and Thomas H Rammsayer.
Cortical oxygen consumption in mental arithmetic as a function of task difficulty: a near-infrared
spectroscopy approach. Frontiers in human neuroscience, 7:217, 2013.

Gert Pfurtscheller, Günther Bauernfeind, Selina Christin Wriessnegger, and Christa Neuper. Focal
frontal (de)oxyhemoglobin responses during simple arithmetic. International Journal of
Psychophysiology, 76(3):186–192, 2010.

Matthias L Schroeter, Timo Stein, Nina Maslowski, and Jane Neumann. Neural correlates of
alzheimer's disease and mild cognitive impairment: a systematic and quantitative meta-analysis
involving 1351 patients. NeuroImage, 47(4):1196–1206, 2009.

Michael K Yeung and Agnes S Chan. Functional near-infrared spectroscopy reveals decreased
resting oxygenation levels and task-related oxygenation changes in mild cognitive impairment and
dementia: A systematic review. Journal of psychiatric research, 124:58– 76, 2020.

Vanessa J Pereira, Karl G Linden, and Howard S Weinberg. Evaluation of uv irradiation for
photolytic and oxidative degradation of pharmaceutical compounds in water. Water Research,
41(19):4413–4423, 2007.

Theodore J Huppert, Solomon G Diamond, Maria A Franceschini, and David A Boas. Homer: a
review of time-series analysis methods for near-infrared spectroscopy of the brain. Applied optics,
48(10):D280–D298, 2009.

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., … Joliot,
M. (2002). Automated anatomical labeling of activations in SPM using a macroscopic anatomical
parcellation of the MNI MRI single-subject brain. NeuroImage, 15(1), 273–289.

Ilias Tachtsidis and Felix Scholkmann. False positives and false negatives in functional near-infrared
spectroscopy: issues, challenges, and the way forward. Neurophotonics, 3(3):031405, 2016.

Alexander von Lühmann, Xinge Li, Klaus-Robert Müller, David A Boas, and Meryem A Yücel.
Improved physiological noise regression in fnirs: A multimodal extension of the general linear model
using temporally embedded canonical correlation analysis. NeuroImage, 208:116472, 2020.

Hendrik Santosa, Xuetong Zhai, Frank Fishburn, and Theodore Huppert. The nirs brain analyzir
toolbox. Algorithms, 11(5):73, 2018.

Nagata, H., Yamamoto, Y. and Sakaguchi, R. (2010), P2-123: Our newly developed Simple
Cognitive test can be also used by computer or paper. Alzheimer's & Dementia, 6: S349-S349.
https://doi.org/10.1016/j.jalz.2010.05.1170

Sadanobu Ichii, Takumi Nakamura, Takeshi Kawarabayashi, Masamitsu Takatama, Tetsuya Ohgami,
Kazushige Ihara, and Mikio Shoji. Cogevo, a cognitive function balancer, is a sensitive and easy
psychiatric test battery for age-related cognitive decline. Geriatrics & Gerontology International,
20(3):248–255, 2020.

Hajime Takechi and Hiroshi Yoshino. Usefulness of cogevo, a computerized cognitive assessment
and training tool, for distinguishing patients with mild alzheimer's disease and mild cognitive
impairment from cognitively normal older people. Geriatrics & Gerontology International,
21(2):192–196, 2021.

Jieun Yoon, Hiroko Isoda, and Tomohiro Okura. Evaluation of beneficial effect of a dual-task
exercise based on japanese transitional games in older adults: a pilot study. Aging (Albany NY),
12(19):18957, 2020.

Gary Strangman, Joseph P Culver, John H Thompson, and David A Boas. A quantitative comparison
of simultaneous bold fmri and nirs recordings during functional brain activation. NeuroImage,
17(2):719–731, 2002.

Xu Cui, Signe Bray, Daniel M Bryant, Gary H Glover, and Allan L Reiss. A quantitative comparison
of nirs and fmri across multiple cognitive tasks. NeuroImage, 54(4):2808–2821,2011.

Theodore J Huppert, Richard D Hoge, Solomon Gilbert Diamond, Maria Angela Franceschini, and
David A Boas. A temporal comparison of bold, asl, and nirs hemodynamic responses to motor
stimuli in adult humans. NeuroImage, 29(2):368–382, 2006.

Shuntaro Sasai, Fumitaka Homae, Hama Watanabe, Akihiro T Sasaki, Hiroki C Tanabe, Norihiro
Sadato, and Gentaro Taga. A nirs–fmri study of resting state network.NeuroImage, 63(1):179–193,
2012.

Congcong Huo, Gongcheng Xu, Wenhao Li, Hui Xie, Tengyu Zhang, Ying Liu, and Zengyong Li. A
review on functional near-infrared spectroscopy and application in stroke rehabilitation. Medicine in
Novel Technology and Devices, page 100064, 2021.

Hendrik Santosa, Xuetong Zhai, Frank Fishburn, Patrick J Sparto, and Theodore J Huppert.
Quantitative comparison of correction techniques for removing systemic physiological signal in
functional near-infrared spectroscopy studies. Neurophotonics, 7(3):035009,2020.

Ana Carolina De Campos, Theresa Sukal-Moulton, Theodore Huppert, Katharine Alter, and Diane L
Damiano. Brain activation patterns underlying upper limb bilateral motor coordination in unilateral
cerebral palsy: an fnirs study. Developmental Medicine & Child Neurology, 62(5):625–632, 2020.

Mohamed L Seghier. Laterality index in functional mri: methodological issues. Magnetic resonance
imaging, 26(5):594–601, 2008.

Eiju Watanabe, Atsushi Maki, Fumio Kawaguchi, Kaoru Takashiro, Yuichi Yamashita, Hideaki
Koizumi, and Yoshiaki Mayanagi. Non-invasive assessment of language dominance with near-
infrared spectroscopic mapping. Neuroscience letters, 256(1):49–52, 1998.

Kirstin L Bierre, Samuel JE Lucas, Hayley Guiney, James D Cotter, and Liana Machado. Cognitive
difficulty intensifies age-related changes in anterior frontal hemodynamics: novel evidence from

near-infrared spectroscopy. Journals of Gerontology Series A: Biomedical Sciences and Medical
Sciences, 72(2):181–188, 2017.

Nathalie Fournet, Jean-Luc Roulin, Fanny Vallet, Marine Beaudoin, Stefan Agrigoroaei, Adeline
Paignon, Cécile Dantzer, and Olivier Desrichard. Evaluating short-term and working memory in
older adults: French normative data. Aging & mental health, 16(7):922–930, 2012.

7   Figure legends

Figure 1. Overview of the experiment. A. The experimental procedure. B. The experimental setup.

Figure 2. Probe displacement of fNIRS. (a) Monte-Carlo simulation results over the frontal cortex,
temporal cortex, and parietal cortex. Red dots represent emitters, blue dots represent detectors, and
yellow lines represent measurement channels, respectively (created using Homer 2 AtlasViewer; v2.8,
p2.1: https://www.nitrc.org/frs/shownotes.php?release_id=3956). The color bar represents the spatial
sensitivity of fNIRS measurements. (b) The two-dimensional fNIRS montage using the International
10-20 measurement system as reference is presented.

Figure 3. βmaps and LI values in the planning block. (a)-(c) βmaps of participants with LI values
close to the median in younger, expert older, and non-expert older groups are shown, respectively. (d)
Distribution of LI values for the three participant groups. *p < .016. βmaps were visualized using the
xjView toolbox and BrainNet viewer.

Figure 4. βmaps and LI values in the calculation block. (a)-(c) βmaps of participants with LI values
close to the median in the younger, expert older, and non-expert older groups are shown, respectively.
(d) Distribution of LI values for the three participant groups. *p < .016. βmaps were visualized using
the xjView toolbox and BrainNet viewer.

Figure 5. Correlation between the number of days of Wellness Dart experience and LI values in the
expert older group. (a) planning block. (b) calculation block.

Figure 6. Dart performance level in the younger, expert older, and non-expert older group. *p<.05.

Figure 7. Correlation between darts performance level and LI value in the expert older group. (a)
planning block. (b) calculation block.

Figure 8. β values of IPL in the younger, expert older, and non-expert older group.

Figure 9. Computational ability test scores for the younger, expert older, and non-expert older group.
*p <.05.

Figure 10. STM test scores for the younger, expert older, and non-expert older group. *p <.05.

Figure 11. Scores on five different CogEvo tasks for the three groups of participants. (a) Orientation.
(b) Visual search. (c) Flashlight. (d) Number step. (e) Just Fit.

Figure 12. TMPT completion time for the younger, expert older, and non-expert older group.

8   Tables

Table 1. Descriptive statistics on the three groups.

Characteristics		Expert older group	Nonexpert older group (older control)	Younger group (nonexpert younger control)
age (years)		75.1±4.3	74.1±4.5	22.5±0.8
Number of days of Wellness Darts experience (days)		170.7±145.6	0.0	0.0
Cognitive tests	SC test	41.1±7.8	44.5±5.8	48.8±2.5
	Computational ability test	20.5±3.8	17.1±5.5	24.0±3.8
	STM test	57.1±9.5	49.4±9.9	74.6±13.6
	TMPT	58.1±8.5	58.9±9.2	41.1±4.5
	CogEvo: disorientation	361.8±52.7	278.2±74.5	341.5±68.8
	CogEvo: visual search	177.9±39.0	161.3±33.8	282.0±41.3
	CogEvo: flashlight	364.3±146.9	308.1±106.4	604.3±169.2
	CogEvo: number step	197.4±40.8	173.5±45.9	280.9±42.8
	CogEvo: just fit	332.2±104.9	342.4±98.7	442.1＋78.2
Dart-throwing performance		49.9±5.5	31.9±13.7	39.3±13.4

663   **Table 2.** Brain regions that showed task-related activation for each individual in the planning block
664   (younger group). Only regions of interest that showed $p < 0.05$ (FDR corrected at peak level) are
665   shown.

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	Participant	Anatomical Region	t-value	p-value (FDR-corrected)
	IPL.L	3.765	5.76E-04		MFG.L	5.449	2.54E-07
1	MFG.L	4.408	4.06E-05	11	MFG.R	8.413	8.76E-16
	MFG.R	10.33	5.57E-23		SFGdor.L	6.024	1.12E-08

Hemispheric lateralization in dart players

	SFGdor.L	7.434	1.48E-12		SFGdor.R	8.779	5.39E-17
	SFGdor.R	9.248	9.70E-19		MFG.L	4.957	1.77E-06
2	IPL.R	−4.224	3.05E-04	12	MFG.R	5.836	2.56E-08
	MFG.L	2.498	0.038		SFGdor.L	6.686	1.67E-10
	MFG.R	−2.941	0.012		SFGdor.R	5.824	2.63E-08
	SFGdor.R	−2.978	0.012	13	IPL.L	−2.443	0.035
3	IPL.R	3.342	0.002		MFG.L	5.219	1.93E-06
	MFG.R	6.801	1.32E-10		SFGdor.L	5.193	2.03E-06
	SFGdor.R	5.896	2.37E-08	14	MFG.R	7.688	3.71E-13
4	IPL.L	−3.846	1.84E-04		SFGdor.R	6.225	2.75E-09
	IPL.R	−4.344	2.73E-05	15	IPL.L	−3.952	2.21E-04
	MFG.L	−6.759	7.67E-11		MFG.R	8.127	8.13E-15
	MFG.R	−2.298	0.028		SFGdor.R	9.337	7.87E-19
	SFGdor,L	−4.55	1.07E-05	16	IPL.L	−6.443	3.36E-09
	SFGdor.R	−2.273	0.029		MFG.R	2.473	0.026
5	MFG.L	8.83	9.44E-18	17	IPL.L	−4.459	9.81E-05

	MFG.R	6.576	2.80E-10		SFGdor.L	−2.842	0.02
	SFGdor.L	19.18	2.87E-77		IPL.L	−9.742	1.12E-20
	SFGdor.R	7.335	1.45E-12		MFG.L	−3.147	0.005
	IPL.R	6.645	3.50E-10	18	MFG.R	−3.923	3.48E-04
6	SFGdor.L	−4.996	2.78E-06		SFGdor.L	−5.72	7.17E-08
	SFGdor.R	2.399	0.028		SFGdor.R	−4.389	5.23E-05
	IPL.L	2.771	0.008		IPL.L	6.362	7.05E-10
	IPL.R	4.191	6.82E-05		MFG.L	7.276	2.11E-12
7	MFG.R	4.289	4.64E-05	19	MFG.R	7.561	3.03E-13
	SFGdor.L	4.583	1.38E-05		SFGdor.L	6.035	4.98E-09
	SFGdor.R	4.365	3.38E-05		SFGdor.R	10.65	3.69E-24
	IPL.L	4.411	1.56E-05	20	MFG.R	4.538	1.88E-05
	IPL.R	4.257	3.09E-05		SFGdor.R	4.636	1.26E-05
8	MFG.L	5.811	1.70E-08		IPL.L	2.346	0.033
	MFG.R	5.317	2.31E-07		MFG.L	3.582	9.94E-04
	SFGdor.L	9.745	5.83E-21	21	MFG.R	12.54	9.28E-34
	SFGdor.R	6.694	7.43E-11		SFGdor.L	4.791	5.92E-06

	IPL.L	3.432	0.002	SFGdor.R	12.01	3.62E-31
10	MFG.L	3.113	0.004			
	MFG.R	4.145	1.42E-04			
	SFGdor.R	3.909	3.05E-04			

666 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
667 gyrus; R: right; L: Left.

668 **Table 3.** Brain regions that showed task-related activation for each individual in the calculation block
669 (younger group). Only regions of interest that showed p < 0.05 (FDR corrected at peak level) are
670 shown.

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	Participant	Anatomical Region	t-value	p-value (FDR-corrected)
1	MFG.R	7.509	2.55E-12		IPL.L	−3.342	0.002
	SFGdor.L	4.194	2.04E-04		MFG.L	5.89	1.75E-08
	SFGdor.R	5.533	7.67E-05	11	MFG.R	6.951	2.72E-11
2	IPL.R	−8.217	1.51E-14		SFGdor.L	6.464	6.57E-10
	MFG.R	−3.969	8.20E-04		SFGdor.R	8.092	8.43E-15
	SFGdor.R	−5.505	7.29E-07	13	IPL.L	−2.444	0.035
3	IPL.L	−	8.59E-07		MFG.L	5.21	1.93E-06

		5.429				9	
	MFG.R	5.85	8.88E-08		SFGdor.L	5.193	2.03E-06
	SFGdor.R	2.523	0.023		MFG.L	3.061	0.004
4	IPL.L	−8.516	7.35E-17	14	MFG.R	7.079	9.36E-12
	IPL.R	−9.993	1.64E-22		SFGdor.L	3.61	6.08E-04
	MFG.L	−2.591	0.011		SFGdor.R	6.778	6.56E-11
	MFG.R	−2.234	0.029	15	MFG.R	7.157	1.92E-11
	SFGdor.L	−5.707	2.23E-08		SFGdor.R	8.688	2.59E-16
	SFGdor.R	−4.508	9.39E-06	16	IPL.L	−6.433	2.05E-09
5	IPL.L	−3.004	0.007		IPL.L	−6.99	4.20E-11
	MFG.L	6.701	1.05E-10		MFG.L	−7.918	5.44E-14
	MFG.R	2.79	0.013	17	SFGdor.L	−8.635	2.21E-16
	SFGdor.L	15.949	2.31E-54		SFGdor.R	−3.074	0.006

Hemispheric lateralization in dart players

Group	Region	value	p		Group	Region	value	p
	SFGdor.R	2.956	0.008			IPL.L	−4.03	1.73E-04
	IPL.R	3.981	2.01E-04			MFG.L	−5.082	2.31E-06
6	MFG.R	3.87	2.90E-04		18	MFG.R	−3.547	9.51E-04
	SFGdor.L	−5.067	2.42E-06			SFGdor.L	−6.342	2.48E-09
	SFGdor.R	4.977	3.63E-06			SFGdor.R	−4.291	6.13E-05
7	IPL.L	3.011	0.009			IPL.L	3.566	6.62E-04
	IPL.R	4.247	1.77E-04			IPL.R	−2.674	0.011
	IPL.L	3.933	1.30E-04		19	MFG.L	4.345	3.13E-05
	IPL.R	3.329	0.001			MFG.R	7.278	1.86E-12
8	MFG.L	5.935	7.59E-09			SFGdor.L	4.855	3.25E-06
	MFG.R	2.072	0.044			SFGdor.R	11.48	2.56E-28
	SFGdor.L	8.248	2.22E-15		20	IPL.L	−5.517	6.64E-08

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	participant	Anatomical Region	t-value	p-value (FDR-corrected)
	SFGdor.R	2.922	0.004		MFG.L	2.113	0.042
9	IPL.L	−3.998	1.99E-04		MFG.R	4.332	2.16E-05
	IPL.R	−5.626	1.14E-07		SFGdor.R	3.732	2.57E-04
	MFG.L	−3.412	0.001		MFG.L	3.828	3.12E-04
				21	MFG.R	14.12	6.09E-42
					SFGdor.L	3.692	4.94E-04
					SFGdor.R	12.36	3.38E-33

671 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
672 gyrus; R: right; L: Left.

673 **Table 4.** Brain regions that showed task-related activation in the planning block for each individual
674 (expert older group). Only regions of interest with p < 0.05 (FDR corrected at peak level) are shown.

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	participant	Anatomical Region	t-value	p-value (FDR-corrected)
1	IPL.R	4.418	4.97E-05	12	MFG.L	−5.516	1.33E-06
	MFG.R	−4.478	3.88E-05		MFG.R	−4.304	3.30E-04

Hemispheric lateralization in dart players

2	IPL.L	5.142	1.98E-06	13	MFG.L	6.548	1.97E-10
	IPL.R	4.167	1.24E-04		MFG.R	3.212	0.002
	MFG.L	-4.8	9.13E-06	14	MFG.L	5.044	3.19E-06
	SFGdor.L	-3.54	0.001		MFG.R	6.031	2.05E-08
3	MFG.L	-7.369	1.65E-12		SFGdor.L	4.537	2.84E-05
	SFGdor.L	-4.309	4.51E-05		SFGdor.R	2.859	0.009
4	IPL.R	-5.123	1.03E-06	15	IPL.L	6.276	3.28E-09
	MFG.L	4.82	4.32E-06		IPL.R	7.845	1.55E-13
5	IPL.R	2.806	0.01		MFG.L	4.419	3.04E-05
	SFGdor.L	-3.32	0.002		MFG.R	5.777	6.03E-08
6	MFG.L	-2.802	0.014		SFGdor.R	4.734	8.11E-06
7	MFG.L	4.296	1.29E-04	16	IPL.R	-4.795	2.22E-05
	SFGdor.L	3.35	0.003	17	IPL.L	-4.80	5.20E-06

		2				6	
8	IPL.L	−4.035	2.56E−04		MFG.L	−2.944	0.006
8	SFGdor.R	−2.635	0.02		MFG.L	−2.989	0.007
9	IPL.L	3.995	4.11E−04	19	MFG.R	−5.129	1.74E−06
9	SFGdor.L	3.754	9.62E−04	19	SFGdor.L	−4.712	1.02E−05
10	IPL.L	−3.172	0.007		SFGdor.R	−2.744	0.012
10	IPL.R	7.868	2.43E−13	20	MFG.L	2.756	0.023
10	MFG.R	2.39	0.049	20	SFGdor.L	3.671	0.001
11	MFG.R	7.11	3.42E−11	21	MFG.R	3.168	0.005
11	SFGdor.R	4.948	1.11E−05	21	SFGdor.R	5.267	1.52E−05

Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal gyrus; R: right; L: Left.

Table 5. Brain regions that showed task-related activation for each individual in the calculation block (expert older group). Only regions of interest with $p < 0.05$ (FDR corrected at peak level) are shown.

Participa	Anatomica	t−	p-value	Participa	Anatomica	t−	p-value

Hemispheric lateralization in dart players

nt	l Region	value	(FDR-corrected)	nt	l Region	value	(FDR-corrected)
1	IPL.R	4.418	4.97E-05		MFG.L	5.359	2.13E-07
1	MFG.R	-4.478	3.88E-05	13	MFG.R	5.988	6.77E-09
2	IPL.L	5.142	1.98E-06	13	SFGdor.L	3.457	9.19E-04
2	IPL.R	4.167	1.24E-04		SFGdor.R	3.344	0.001
2	MFG.L	-4.8	9.13E-06	14	MFG.R	3.831	4.78E-04
2	SFGdor.L	-3.54	0.001		IPL.L	8.888	1.50E-17
3	MFG.L	-7.369	1.65E-12		IPL.R	7.401	1.02E-12
3	SFGdor.L	-4.309	4.51E-05	15	MFG.L	6.731	8.57E-11
4	IPL.R	-5.123	1.03E-06	15	MFG.R	8.754	3.76E-17
4	MFG.L	4.82	4.32E-06		SFGdor.L	5.877	1.52E-08
5	IPL.R	2.806	0.01		SFGdor.R	7.612	2.69E-13
5	SFGdor.L	-3.32	0.002		MFG.R	8.742	5.85E-17
6	MFG.L	-2.802	0.014	16	SFGdor.L	4.577	2.61E-05
7	MFG.L	4.296	1.29E-04		SFGdor.R	9.739	1.51E-20
7	SFGdor.L	3.352	0.003	17	IPL.L	-3.792	3.28E-04
8	IPL.L	-4.035	2.56E-04		MFG.R	4.375	3.15E-05
8	SFGdor.R	-2.635	0.02	18	IPL.L	-2.839	0.012

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	Participant	Anatomical Region	t-value	p-value (FDR-corrected)
9	IPL.L	3.995	4.11E-04		MFG.L	4.933	6.41E-06
	SFGdor.L	3.754	9.62E-04		SFGdor.L	3.433	0.002
10	IPL.L	-3.172	0.007	19	SFGdor.R	3.007	0.013
	IPL.R	7.868	2.43E-13	20	MFG.L	-11.83	5.34E-31
	MFG.R	2.39	0.049		MFG.R	-14.5	1.92E-45
11	MFG.R	7.11	3.42E-11		SFGdor.L	-68.23	0
	SFGdor.R	4.948	1.11E-05		SFGdor.R	-93.95	0
12	MFG.L	-5.55	5.50E-07	21	IPL.R	3.885	2.64E-04
	MFG.R	-4.926	1.23E-05		MFG.L	-4.797	6.94E-06
					MFG.R	4.554	2.01E-05
					SFGdor.L	-21.789	4.15E-98
					SFGdor.R	7.121	1.20E-11

679 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
680 gyrus; R: right; L: Left.

681 **Table 6.** Brain regions that showed task-related activation for each individual in the planning block
682 (non-expert older group). Only regions of interest with $p < 0.05$ are shown.

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	Participant	Anatomical Region	t-value	p-value (FDR-corrected)

Hemispheric lateralization in dart players

	Region	Value	p		Region	Value	p
1	IPL.L	4.003	3.30E-04	13	IPL.L	6.647	1.57E-10
	MFG.R	2.857	0.013		IPL.R	-8.842	2.49E-17
2	IPL.L	-4.04	9.84E-04		MFG.L	6.663	1.48E-10
3	MFG.L	-4.888	1.43E-05		MFG.R	3.744	3.67E-04
	MFG.R	-4.201	1.22E-04		SFGdor.R	5.344	3.00E-07
	SFGdor.L	-4.201	1.22E-04	14	IPL.L	2.801	0.009
	SFGdor.R	-4.027	2.30E-04		IPL.R	2.918	0.006
4	MFG.L	-5.972	2.00E-08		MFG.L	-1.914	0.078
	MFG.R	-2.455	0.028		SFGdor.L	-2.504	0.019
	SFGdor.L	-5.021	2.74E-06	15	IPL.L	4.378	4.65E-05
	SFGdor.R	-3.152	0.004		MFG.R	-3.081	0.005
5	MFG.L	4.23	2.98E-04	16	IPL.L	-3.582	0.004

Hemispheric lateralization in dart players

Group	Region	Value	p	Group	Region	Value	p
	MFG.R	3.643	0.002		MFG.L	-3.388	0.006
	SFGdor.R	5.05	1.69E-05		SFGdor.R	-2.628	0.031
7	MFG.R	-3.466	0.002		IPL.L	-3.349	0.002
7	SFGdor.L	-4.202	2.30E-04		MFG.L	-2.394	0.027
8	MFG.L	4.709	3.04E-05	17	SFGdor.L	-9.055	2.14E-18
8	MFG.R	2.838	0.014		SFGdor.R	-7.301	1.48E-12
8	SFGdor.R	3.386	0.003		IPL.L	14.66	4.98E-46
9	IPL.L	6.602	2.53E-09	18	MFG.L	5.508	1.02E-07
9	SFGdor.L	-3.184	0.008		SFGdor.L	4.667	7.56E-06
	IPL.L	2.37	0.04		SFGdor.R	5.935	8.93E-09
10	MFG.L	7.529	8.86E-13	19	IPL.L	8.061	2.25E-14
	MFG.R	5.415	3.04E-07		SFGdor.L	3.538	8.87E-04

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	Participant	Anatomical Region	t-value	p-value (FDR-corrected)
	SFGdor.L	7.37	2.32E-12		SFGdor.R	3.536	8.87E-04
	SFGdor.R	6.141	5.18E-09		IPL.L	-2.806	0.018
11	MFG.L	-4.217	1.30E-04	20	IPL.R	-3.373	0.004
	SFGdor.L	-3.748	6.50E-04		MFG.L	-8.005	7.56E-14
	SFGdor.R	-2.585	0.023		SFGdor.L	-2.431	0.046
				21	MFG.L	-2.649	0.024

683 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
684 gyrus; R: right; L: Left.

685 **Table 7.** Brain regions that showed task-related activation for each individual in the calculation block
686 (non-expert older group). Only regions of interest with $p < 0.05$ are shown.

Participant	Anatomical Region	t-value	p-value (FDR-corrected)	Participant	Anatomical Region	t-value	p-value (FDR-corrected)
1	IPL.L	3.241	0.003	13	IPL.L	-8.63	1.57E-16
	SFGdor.L	-2.994	0.006		IPL.R	-8.138	4.42E-15

Hemispheric lateralization in dart players

Group	Region	Value	p-value	Group	Region	Value	p-value
2	IPL.L	−7.816	7.98E-13	15	MFG.L	5.221	5.11E-07
2	MFG.L	−3.72	0.002		MFG.R	3.503	9.32E-04
3	IPL.L	−2.901	0.008		SFGdor.R	5.609	6.61E-08
	IPL.R	−2.726	0.013		IPL.R	−2.658	0.025
	MFG.L	−5.086	1.88E-06		MFG.L	−4.417	8.75E-05
	SFGdor.L	−2.479	0.024		IPL.L	−4.386	4.12E-05
	SFGdor.R	8.724	4.59E-17		IPL.R	−3.726	5.33E-04
4	MFG.L	−3.32	0.01	16	MFG.L	−6.352	2.58E-09
	IPL.R	2.419	0.032		MFG.R	−4.675	1.22E-05
5	MFG.L	7.139	3.06E-11		SFGdor.L	−2.931	0.007
	MFG.R	6.862	1.73E-10		SFGdor.R	−6.661	5.64E-10
	SFGdor.L	3.649	9.31E-04	17	IPL.L	−	7.40E-08

						5.71	
	SFGdor.R	8.793	1.29E-16		SFGdor.L	-4.989	2.74E-06
7	SFGdor.L	-2.798	0.011		SFGdor.R	-2.72	0.014
8	MFG.L	10.47	7.82E-24		IPL.L	8.34	3.71E-16
	MFG.R	5.262	7.12E-07		MFG.R	-2.519	0.017
	SFGdor.R	5.782	5.24E-08	18	SFGdor.L	7.127	3.42E-12
9	IPL.L	10.99	1.15E-25		SFGdor.R	5.032	9.95E-07
	SFGdor.L	-3.241	0.003		IPL.L	7.062	2.39E-11
10	MFG.L	4.3	6.68E-05	19	MFG.R	-3.826	3.31E-04
	MFG.R	2.867	0.011		IPL.L	3.576	0.001
	SFGdor.L	5.681	9.16E-08		IPL.R	4.542	2.56E-05
11	MFG.L	-4.033	2.33E-04	20	MFG.L	9.534	1.23E-19
	SFGdor.L	-2.915	0.009		MFG.R	3.608	0.001
	SFGdor.R	-2.952	0.008		SFGdor.L	4.643	1.73E-05

12	MFG.R	-2.837	0.013		SFGdor.R	5.79	9.68E-08
	SFGdor.R	-2.926	0.01		MFG.L	-3.354	0.003
				21	MFG.R	-2.275	0.044
					SFGdor.R	-2.226	0.049

687 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
688 gyrus; R: right; L: Left.

689 **Table 8.** Task-related activation in the region of interest revealed by group-level analysis (younger
690 group).

Condition	Anatomical Region	t-value	p-value (FDR-corrected)
planning	IPL.L	0.15	0.909
	MFG.L	1.931	0.186
	MFG.R	3.321	0.022
	SFGdor.L	3.155	0.022
	SFGdor.R	3.822	0.012
calculation	IPL.L	0.139	0.454
	MFG.L	3.321	0.022
	MFG.R	3.321	0.022

	SFGdor.L	2.866	0.052
	SFGdor.R	5.847	5.72E-05

691 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
692 gyrus; R: right; L: Left.

693 **Table 9.** Task-related activation in the region of interest revealed by group-level analysis (expert
694 older group).

Condition	Anatomical Region	t-value	p-value (FDR-corrected)
planning	IPL.L	0.127	0.988
	IPL.R	1.706	0.548
	MFG.L	-0.163	0.988
	MFG.R	1.238	0.709
	SFGdor.L	0.343	0.988
	SFGdor.R	1.04	0.817
calculation	IPL.L	0.545	0.723
	IPL.R	0.367	0.82
	MFG.L	0.999	0.495
	MFG.R	1.542	0.289
	SFGdor.L	1.115	0.272
	SFGdor.R	0.989	0.495

695 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
696 gyrus; R: right; L: Left.

697 **Table 10.** Task-related activation in the region of interest revealed by group-level analysis (non-
698 expert older group).

Condition	Anatomical Region	t-value	p-value (FDR-corrected)
planning	IPL.L	2.282	0.385
	MFG.L	0.446	0.943
	MFG.R	-0.928	0.794
	SFGdor.L	-0.331	0.943
	SFGdor.R	-0.684	0.904
calculation	IPL.L	1.234	0.525
	MFG.L	0.309	0.876
	MFG.R	-1.118	0.54
	SFGdor.L	-0.496	0.803
	SFGdor.R	-0.469	0.815

699 Abbreviations: IPL: Parietal Inf lobule; MFG: middle frontal gyrus; SFGdor: dorsal superior frontal
700 gyrus; R: right; L: Left.

(a)

Recorded voice prompts from the experimental computer

Beep cue to start throwing a dart

(b)

Dartboard

Fixation point

Button response device

fNIRS system

Scoring sheet

(a)

(b)

Front

L ● ● R

Back

Detector

Source

30mm

Short Distance Detector

8mm

Accelerometer

A

● Source
● Detector
● Short distance detector
● Accelerometer

Log₁₀ sensitivity (mm⁻¹)
-2.0 -1.8 -1.6 -1.4 -1.2 -1.0 -0.8 -0.6 -0.4 -0.2

(a) HbO: younger group

SFGdor.R   SFGdor.R
MFG.R   MFG.R

HbO β-value

−200   200

(b) HbO: expert older group

SFGdor.R   SFGdor.R
MFG.R   MFG.R

HbO β-value

−300   300

(c) HbO: non-expert older group

SFGdor.R   SFGdor.R
MFG.R   MFG.R

HbO β-value

−500   500

(d)

younger group   expert older group   non-expert older group

(a) planning

ρ = -0.439
p = 0.046

(b) calculation

ρ = -0.037
p = 0.874

(a) planning

(b) calculation