Disruption of the plastid-hosted iron-sulfur cluster biogenesis pathway in *Toxoplasma gondii* has pleiotropic effects irreversibly impacting parasite viability

Eléa A. Renaud[1‡], Sarah Pamukcu[1‡], Aude Cerutti[1], Laurence Berry[1], Catherine Lemaire-Vieille[2], Yoshiki Yamaryo-Botté[2], Cyrille Y. Botté[2], and Sébastien Besteiro[1*]

[1] LPHI, Univ Montpellier, CNRS, UMR5235, Montpellier, France

[2] ApicoLipid Team, Centre National de la Recherche Scientifique, Institute for Advanced Biosciences, Institut National de la Santé et de la Recherche Médicale, UMR5309, U1209, Université Grenoble Alpes, Grenoble, France

[‡] These authors contributed equally to this work; author order was determined randomly

[*] For correspondence: Sébastien Besteiro, sebastien.besteiro@inserm.fr

Running title: Importance of the Toxoplasma SUF iron-sulfur cluster pathway

Keywords: Toxoplasma gondii, iron metabolism, iron-sulfur protein, parasite metabolism, apicoplast, lipid

Abstract

Like many other apicomplexan parasites, *Toxoplasma gondii* contains a plastid harbouring key metabolic pathways, including the SUF pathway that is involved in the biosynthesis of iron-sulfur clusters. These cofactors are key for a variety of proteins involved in important metabolic reactions, potentially including plastidic pathways for the synthesis of isoprenoid and fatty acids. It was shown previously that impairing the NFS2 cysteine desulfurase, involved in the first step of the SUF pathway, leads to an irreversible killing of intracellular parasites. However, the metabolic impact of disrupting the pathway remained unexplored. We have generated another mutant of the pathway, deficient for the SUFC ATPase, and we have investigated in details the phenotypic consequences of TgNFS2 and TgSUFC depletion on parasite homeostasis. Our analysis confirms that Toxoplasma SUF mutants are severely and irreversibly impacted in growth: cell division and membrane homeostasis are particularly affected. Lipidomic analysis suggests a defect in apicoplast-generated fatty acids, along with a simultaneous increase in scavenging of host-derived lipids. However, addition of exogenous lipids did not allow full restauration of growth, suggesting other more important cellular functions were impacted in addition to fatty acid synthesis. For instance, we have shown that the SUF pathway is also key for generating isoprenoid-derived precursors necessary for the proper targeting of GPI-anchored proteins as well as for the parasite gliding motility. Thus, plastid-generated iron-sulfur clusters support the functions of proteins involved in several vital downstream cellular pathways, which implies the SUF machinery may be explored for discovering new potential anti-Toxoplasma targets.

Introduction

Apicomplexan parasites are some of the most prevalent and morbidity-causing pathogens worldwide. Noticeably, they comprise *Plasmodium* species that can naturally infect humans and cause the deadly malaria in tropical and subtropical areas of the world (1). Although less lethal, another apicomplexan parasite called *Toxoplasma gondii* can cause serious illness in animals, including humans, and has a widespread host range and geographical distribution (2). These protists are obligate intracellular parasites that rely to a large extent on their host cells for nutrient acquisition and for protection from the immune system. Through their evolutionary history, *Plasmodium* and *Toxoplasma* have inherited a plastid from a secondary endosymbiosis event involving the engulfment of a red alga whose photosynthetic capability previously originated from the acquisition of a cyanobacterium (3). Although the ability to perform photosynthesis has been lost during evolution when the ancestors of Apicomplexa became parasitic (4), the plastid has retained critical metabolic functions. For instance it hosts pathways for the synthesis of heme (together with the mitochondrion), fatty acids (FASII), isoprenoid precursors (through the so-called non-mevalonate pathway), and iron-sulfur (Fe-S) clusters (5, 6). Because of its origin and its metabolic importance, the apicoplast is particularly attractive to look for potential drug targets (7).

As some of the earliest catalytic cofactors on earth (8), Fe-S clusters are found in all kingdoms of life, associated with proteins involved in a number of key cellular functions like the synthesis of metabolites, the replication and repair of DNA, the biogenesis of ribosomes and the modification of tRNAs (9). The biosynthesis of Fe-S clusters necessitates a complex machinery for assembling ferrous (Fe^{2+}) or ferric (Fe^{3+}) iron and sulfide (S^{2-}) ions, and delivering the resulting Fe-S cluster to target client proteins (10). In eukaryotes Fe-S proteins are present in various subcellular compartments like the cytosol and the nucleus, but also organelles of endosymbiotic origin like mitochondria or plastids, and thus require compartment-specific biogenesis systems. The three main eukaryotic Fe-S synthesis pathways comprise the ISC (iron-sulfur cluster) machinery, hosted by the mitochondrion, the cytosolic Fe-S protein assembly (CIA) machinery, important not only for the generation of cytosolic, but also of nuclear Fe-S proteins, and the SUF (sulfur formation) pathway that is found in plastids (9).

Like in plants and algae, apicoplast-containing apicomplexan parasites seem to express the machinery corresponding to the three eukaryotic pathways. For instance, in *T. gondii* recent investigations have shown that the CIA, ISC and SUF pathways are all essential for parasite fitness (11, 12). From a biochemical point of view, mitochondrial and plastidic Fe-S cluster biosynthesis pathways follow a similar general pattern: cysteine desulfurases produce sulfur from L-cysteine, then scaffold proteins provide a molecular platform allowing assembly of iron and sulfur into a cluster, and finally carrier proteins deliver the cluster to target apoproteins. Importantly, targeting the *T. gondii* mitochondrial ISC pathway through disruption of scaffold protein ISU1 was shown to lead to a reversible growth arrest and to trigger differentiation into a stress-resistant form; while on the other hand, targeting the plastidic SUF pathway by inactivating NFS2 function led to an irreversible lethal phenotype (12). Like for many apicoplast-hosted pathways, enzymes belonging to the SUF machinery are essentially absent from the mammalian host and as such they may be seen as good potential drug targets. This has sparked considerable interest for the SUF pathway in *Plasmodium*, which has been shown to be important for the viability of several developmental stages of the parasite (13–17).

To better understand the contribution of the SUF pathway to *T. gondii* viability, we have generated a conditional mutant for the scaffold protein TgSUFC and conducted a thorough phenotypic

characterization of this mutant, together with the TgNFS2 mutant we have previously generated (12).
Our results confirm that inactivating the plastid-hosted SUF pathway in *T. gondii* leads to irreversible
and marked effects on membrane homeostasis, impacting the division process and parasite viability.
We show that these effects are likely due to impairment in the function of several key plastidic Fe-S
proteins, which have pleiotropic downstream metabolic consequences for the parasite.

Results

A Toxoplasma SUFC homolog in the apicoplast

Searching for homologs of the plant SUF system in the ToxoDB.org database, we have previously
shown an overall good conservation for the plastidic Fe-S cluster synthesis pathway (12). Among the
candidates for members of the SUF machinery, we have identified a potential *T. gondii* homolog of
SUFC, member of a Fe-S cluster scaffold complex comprising SUFC, SUFB and SUFD (Fig. 1A). This
complex is also present in prokaryotes, where it was first characterized (18): it was shown that
bacterial SufC is an ATP-binding cassette (ABC)-like ATPase component essential for proper Fe-S
cluster assembly (19). Alignment of the amino acid sequences of the *T. gondii* SUFC candidate (entry
TGGT1_225800 in the ToxoDB.org database (20)) and its *Escherichia coli* counterpart showed a good
overall conservation (56% of identity), particularly in the motifs that are characteristic of ABC
ATPAses (Fig. 1B). The *T. gondii* protein presents a N-terminal extension when compared with *E. coli*
SufC, which may contain a transit peptide for targeting to the apicoplast. Accordingly, it was
predicted with high probability to be a plastid-localized protein by the Deeploc 1.0
(http://www.cbs.dtu.dk/services/DeepLoc-1.0/) algorithm, although the exact position of the N-
terminal transit peptide sequence could not be defined. Data from global mapping of *T. gondii*
proteins subcellular location by hyperLOPIT spatial proteomics (21) also suggested an apicoplast
localization for TGGT1_225800. To assess whether this protein is a real functional homolog, we first
performed complementation assays of an *E. coli* SufC mutant, for which growth is slowed, especially
when limiting iron availability with a specific chelator (22). We could show that expressing the
predicted functional domain of TGGT1_225800 restored bacterial growth (Fig. 1C), even in the
presence of the iron chelator, confirming this protein (hereafter named TgSUFC) is functional.

In order to detect TgSUFC expression and assess its sub-cellular localization in the tachyzoite stage
(the fast-replicating stage associated with acute toxoplasmosis (2)), we epitope-tagged the native
protein. This was performed by inserting a sequence coding for a C-terminal triple hemagglutinin
(HA) epitope tag at the endogenous *TgSUFC* locus by homologous recombination (Figure S1). It was
achieved in the TATi ΔKu80 cell line, favoring homologous recombination and allowing
transactivation of a Tet operator-modified promoter that we subsequently used for generating a
conditional mutant in this background (23–25). Immunoblot analysis with an anti-HA antibody
revealed two products, likely corresponding to the immature and mature forms (resulting from the
cleavage of the transit peptide upon import into the organelle) of TgSUFC (Fig. 2A).
Immunofluorescence assay (IFA) with the anti-HA antibody and co-staining with an apicoplast marker
confirmed that TgSUFC localizes to this organelle in *T. gondii* tachyzoites (Fig. 2B).

Depletion of TgSUFC blocks parasite growth

Next, we generated a conditional TgSUFC mutant cell line in the TgSUFC-HA-expressing TATi ΔKu80
background. Replacement of the endogenous promoter by an inducible-Tet07SAG4 promoter was

achieved through a single homologous recombination at the locus of interest, yielding the cKD
TgSUFC-HA cell line (Fig. S2). In this cell line, the addition of anhydrotetracycline (ATc) can repress
transcription through a Tet-Off system (26). Initial phenotypic characterization was performed on
two independent clones, which were found to behave similarly and thus only one was analysed
further. It should be noted that the promoter replacement resulted in a slightly higher expression of
TgSUFC, but did not change the maturation profile of the protein (Fig. 2A). Down-regulation of
TgSUFC was assessed by growing the parasites in the presence of ATc. Immunoblot and IFA analyses
showed a decrease of TgSUFC to almost undetectable levels after as early as one day of ATc
treatment (Fig. 2A and B). We also generated a complemented cell line constitutively expressing an
additional TY-tagged (27) copy of TgSUFC from the *uracil phosphoribosyltransferase* (*UPRT*) locus,
driven by a *tubulin* promoter (Fig. S3). This cell line, named cKD TgSUFC-HA comp, was found by
immunoblot (Fig. 2C) and IFA (Fig. 2D), to be stably expressing TgSUFC while the HA-tagged copy was
down-regulated in the presence of ATc.

We first evaluated the consequences of TgSUFC depletion on parasite fitness *in vitro* by performing a
plaque assay, which determines the capacity of the mutant and complemented parasites to produce
lysis plaques on a host cells monolayer in absence or continuous presence of ATc for 7 days (Fig. 3).
Depletion of TgSUFC prevented plaque formation, which was restored in the complemented cell lines
(Fig. 3A and B). Our previous analysis of another SUF pathway mutant (TgNFS2, (12)) suggested that
the impact on the pathway leads to irreversible death of the parasites, so we sought to verify this by
removing the ATc after 7 days of incubation and monitoring plaque formation. We confirmed that
depleting TgSUFC was irreversibly impacting parasite viability, as ATc removal did not lead to the
appearance of plaques (Fig. 3C). We next assessed whether this defect in the lytic cycle is due to a
replication problem. Mutant and control cell lines were preincubated in absence or presence of ATc
for 48 hours and released mechanically, before infecting new host cells and were then grown for an
additional 24 hours in ATc prior to parasite counting. We noted that the incubation with ATc led to
an accumulation of vacuoles with fewer TgSUFC mutant parasites, but that it was not the case in the
complemented cell lines (Fig. 3D). Overall, our data show that depleting TgSUFC leads to an
irreversible impact on parasite growth, as previously described for other SUF mutant TgNFS2 (12).

SUF pathway mutants display important membrane defects during cell division

T. gondii tachyzoites divide by a process called endodyogeny, whereby two daughter cells will
assemble inside a mother cell (28). Among the structures which are essential as a scaffold for
daughter cell formation is the inner membrane complex (IMC), a system of flattened vesicles
underlying the plasma membrane and is supported by a cytoskeletal network. The IMC also supports
anchorage for the glideosome, the protein complex powering parasite motility (29). As for several
other cellular structures, there is a combination of *de novo* assembly and recycling of maternal
material during IMC formation in daughter cells (30). To get more precise insights into the impact of
the impairment of the SUF pathway on parasite division, we incubated the TgNFS2 and TgSUFC
mutant parasites with ATc for up to 2 days and stained them for IMC protein IMC3 to detect growing
daughter cells (Fig. 4A). IMC3 is an early marker of daughter cell budding (31), which is usually
synchronized within the same vacuole. However, after two days of ATc treatment an increasing
portion of the vacuoles showed a lack of synchronicity for daughter cell budding for both mutant cell
lines, although the effect was more pronounced for the TgSUFC mutant (Fig. 4A and B).

Then, we used electron microscopy to get a subcellular view of the consequences of TgNFS2 and
TgSUFC depletion on the cell division process. Strikingly, in parasites grown in the continuous

presence of ATc for three days, we observed cytokinesis completion defects. As budding daughter
cells emerge, they normally incorporate plasma membrane material that is partly recycled from the
mother, leaving only a basal residual body. Here, in both TgNFS2 and TgSUFC mutant cell lines
daughter cells remained tethered through patches of plasma membrane (Fig. 4C). However, most
membrane-bound subcellular compartments seemed to have an otherwise normal morphology.
Hence, this highlighted an early and important defect in plasma membrane synthesis and/or
segregation during daughter cell budding. We previously observed major cell division defects after
long term (five days or more) continuous incubation of cKD TgNFS2-HA parasites with ATc (12). When
assessing the cKD TgSUFC-HA parasites in the same conditions, co-staining with apicoplast and IMC
markers revealed similar defects, including organelle segregation problems and an abnormal
membranous structures (Fig. 4D).

Depletion of TgSUFC has an impact on the apicoplast

Computational prediction of the Fe-S proteome combined with hyperLOPIT localization data suggests
there is a limited number of apicoplast proteins potentially containing Fe-S clusters (12). However,
these candidates are supposedly very important for the parasite. They include: IspG and IspH, two
oxidoreductases involved in isoprenoid synthesis (32); LipA, a lipoyl synthase important for the
function of the pyruvate dehydrogenase (PDH) complex (33); MiaB, which is likely a tRNA
modification enzyme (34); as well as of course proteins that are directly involved in Fe-S synthesis,
and the plastidic ferredoxin (Fd) that is an important electron donor that regulates several
apicoplast-localized pathways (35–37). Dataset from a CRISPR-based genome-wide screen suggests
that most of these candidates are important for fitness *in vitro* (38).

We first assessed whether depletion of TgSUFC led to a partial apicoplast loss, as was previously
shown for TgNFS2 (12). Although slowed down in growth, some parasites eventually egressed during
the course of the experiments and were used to reinvade host cells, and were kept for a total of 5
days in the presence of ATc (Fig. 5A). This is reminiscent to the so-called "delayed death" effect,
observed when inhibiting apicoplast metabolism, that often results in slow-kill kinetics (39).
Quantification of the apicoplast marker TgCPN60 showed a progressive loss of this protein (Fig. 5A
and B). As this could reflect a specific impact on this protein marker rather than a general loss of the
organelle, we also stained the parasites with fluorescent streptavidin (mainly detects the biotinylated
apicoplast protein acetyl-CoA carboxylase (40)), confirming a similar loss of signal after 5 days of
incubation with ATc (Fig. S4). This suggests TgSUFC depletion leads to a progressive but late loss of
the apicoplast. Of note, this effect on the organelle is less marked than when TgNFS2 is depleted
(12).

One of the apicoplast-localized Fe-S proteins whose activity can be assessed is LipA, which is
responsible for the lipoylation of a single apicoplast target protein, the E2 subunit of the PDH (41).
We performed an immunoblot analysis with an anti-lipoic acid on protein extracts from cKD TgSUFC-
HA parasites kept in the presence of ATc for up to 4 days (Fig. 5 C and D). We noticed a progressive
decrease in lipoylated PDH-E2 to almost no signal after 3 days of ATc incubation. Using an antibody
that we specifically raised against the E2 subunit of the PDH, we verified this was not due to a
decrease in global levels of this particular protein. Overall, this is comparable to what we previously
described upon depletion of TgNFS2 (12). Thus, disrupting the SUF pathway has direct consequences
on Fe-S proteins-dependent metabolic pathways hosted by the apicoplast, and prolonged depletion
of SUF proteins can even lead to partial loss of the organelle.

Impact of SUF pathway disruption on fatty acid metabolism

The PDH complex generates acetyl-CoA, which is the first step of the FASII system in the apicoplast. This pathway generates fatty acid (FA) precursors that are subsequently elongated in the endoplasmic reticulum (ER) (42), to generate for instance essential membrane building blocks. We first wanted to evaluate the impact of the perturbation of the SUF pathway on parasite lipids. Total lipids from the TgNFS2 and TgSUFC mutant cell lines were determined by gas chromatography-mass spectrometry (GC-MS). For the NFS2 mutant we noted small but significant decrease in abundance of shorter FAs (C12-C17), suggesting de novo FA synthesis is affected (Fig. 6A), however this was not as clear for the SUFC mutant (Fig. 6B).

While FASII was shown previously to be important for tachyzoite fitness (41), recent investigations have shown that tachyzoites are able to scavenge FA precursors from their host cells and at least partly compensate for lack of de novo synthesis (43, 44). To investigate whether the SUFC mutant may also be affected in apicoplast FA synthesis but may have developed a compensatory mechanism, we thus sought to assess the impact of TgSUFC depletion on $de\ novo$-synthesized versus scavenged lipids by stable isotope precursor labeling with ^{13}C glucose combined with mass spectrometry-based lipidomic analyses (43, 45). These analyses highlighted an increase in scavenging from the host cells is happening in absence of TgSUFC, as shown for the C16:0 FA (Fig. 6C). This reflects most likely a mechanism for compensating the lack of $de\ novo$-made FAs by increasing scavenging host-derived FAs.

The ability of tachyzoites to survive lack of de novo lipid synthesis is highly dependent on the availability of exogenous lipid precursors. Others have shown that FASII mutants could be rescued by the addition of palmitic (C16:0) or myristic (C14:0) acid for instance (42, 44, 46). We thus tried to compensate the impact of TgNFS2 and TgSUFC depletion on the parasite lipid homeostasis by supplementing the growth medium with these FAs and performing plaque assays. Palmitic acid supplementation partially restored growth of the TgNFS2 mutant (Fig. 7A and B), while with myristic acid only after long term incubation (up to two weeks) these mutant parasites started growing back (Fig. 7C and D). In contrast, depletion of TgSUFC was not efficiently compensated by exogenous fatty supplementation (Fig. 7). As our lipidomic analysis has shown that depletion of TgNFS2 affects de novo lipid production (Fig. 6A), so it makes sense that providing exogenous lipids can improve fitness of the mutant parasites. On the other hand, TgSUFC mutant parasites are likely able to scavenge lipids from the host as a compensation for an altered de novo synthesis (Fig. 6C), and medium supplementation with more exogenous lipids does not seem to further improve their fitness.

This highlights differences between the two SUF pathway mutants with regards to adaptation to the perturbation of lipid homeostasis, which may be explained by different kinetics in depletion of the respective proteins, or a different impact on global apicoplast function. In any case, our data confirms apicoplast-based lipid production is affected in SUF pathway mutants, and while these parasites establish compensatory mechanisms by scavenging exogenous lipid precursors, they do not allow them recovering to full fitness. This suggests that perturbation of apicoplast FA synthesis is not the only, and perhaps not even the primary, effect of SUF pathway disruption impacting growth.

SUF pathway disruption also has an impact on isoprenoid-dependent pathways

The other main apicoplast-localized biosynthetic pathway potentially affected by disruption of the SUF machinery is isoprenoid synthesis, through the two Fe-S-containing proteins IspG and IspH that are needed for the synthesis of the five carbon precursor isopentenyl pyrophosphate (IPP) and its

isomer dimethylallyl pyrophosphate (DMAPP) (32). Synthesis of these isoprenoid building blocks is
the only essential metabolic function of the apicoplast in the asexual intraerythrocytic stages of
Plasmodium, where the loss of the organelle can be simply compensated by supplementation with
exogenous IPP (47). *Plasmodium* SUF mutants survive when cultured in the presence of IPP,
confirming the essential role of the Fe-S synthesizing pathway in this parasite is likely for isoprenoid
synthesis (13). Isoprenoid synthesis is also vital for *T. gondii* tachyzoites (48), however IPP
supplementation to the culture medium is not possible because, unlike for *Plasmodium*, the highly
charged IPP does not efficiently reach the parasite cytoplasm. Tachyzoites can nevertheless scavenge
some isoprenoids precursors from their host cells (49). Apicoplast-derived isoprenoid precursors are
mostly known for their involvement in important posttranslational protein modifications like
prenylation, glycosylphosphatidylinositol (GPI) anchoring, as well glycosylation, in addition to the
synthesis of quinones and several antioxidant molecules (32).

In *Plasmodium* it is believed that prenylated proteins that regulate vesicle trafficking are key in the
delayed death phenotype caused by apicoplast loss (50). As geranylgeraniol (GGOH), an isoprenoid
precursor for protein farnesylation/prenylation, was successfully used to at least partially
complement deficiencies in apicoplast isoprenoid production (49, 51), we tried to supplement the
culture medium with GGOH and perform plaque assays with the SUF mutants we generated.
However, we did not detect any restoration of growth (Fig. S5A). In contrast to *Plasmodium* (52, 53),
the prenylome of *T. gondii* is still largely uncharacterized, but using an anti-farnesyl antibody, we did
not detect obvious alterations in the general profile of prenylated proteins upon depletion of TgNFS2
or TgSUFC (Fig. S5B). Although apicoplast-generated IPP and DMAPP are also necessary for
synthesizing the farnesyl diphosphate used for protein prenylation in *T. gondii* tachyzoites, it is thus
possible that these parasites can initially scavenge host-derived isoprenoids to compensate for a
deficient *de novo* production (49). In any case, altogether our results suggest that defects in protein
farnesylation/prenylation may not be one of the primary consequences of SUF pathway disruption in
T. gondii.

We next sought to investigate the potential impact of TgNFS2 or TgSUFC depletion on GPI anchoring.
The SAG-related sequence (SRS) family comprising proteins related to SAG1, the first characterized *T.*
gondii surface antigen, is arguably the best characterized family of GPI-anchored proteins in the
parasite (54). We pre-incubated mutant parasites for three days with ATc, and allowed them to
invade and grow into host cells for an extra day in the presence of ATc before using specific
antibodies to detect GPI-anchored SAG1 and SAG3. We could see obvious signs of mislocalization for
these two proteins that, instead of keeping a homogenous peripheral distribution, they were often
seen accumulated at the apex or base of the parasites, or found within the parasitophorous vacuole
space (Fig. 8A and B). Interestingly, while the SAG1 protein appears to be distributed differently by
IFA, our previous immunoblot analyses suggest there is no drastic change in the total amount of
protein upon TgNFS2 or TgSUFC depletion ((12) and Fig. 2A). It was previously shown that the
deletion of SAG1's GPI anchor leads to a constitutive secretion of this protein to the parasitophorous
vacuole space (55). It should be noted that, in contrast to the distribution of GPI-anchored SAG
proteins, in these experimental conditions the overall structure of the IMC appeared unaffected (Fig.
8A and B). Hence, our data suggest that disruption of the SUF pathway perturbs GPI anchor
formation.

Another important posttranslational modification depending on isoprenoid-containing dolichol is
glysosylation. Several key proteins of the glideosome complex are supposedly glycosylated, and as a
consequence glycosylation inhibition has been reported to impact parasite motility (56, 57). We thus
performed gliding motility assays on the SUF mutants. Typically, this is done by monitoring the

shedding of SAG1, that leaves trails when tachyzoites glide on solid substrates. Perhaps as a
consequence of SUF pathway disruption on SAG1 targeting, trails they were clearly less abundant
upon depletion of TgNFS2 and TgSUFC (Fig. 8C and D). We could nevertheless detect and measure
trails, whose mean length provides an estimate of overall motility rates, and they were found to be
significantly smaller in absence of TgNFS2 or TgSUFC (Fig. 8E). These results suggest that protein
glycosylation is also affected upon disruption of the SUF pathway. Overall, our findings indicate that
the depletion of proteins of the SUF pathway leads to defects in the synthesis of isoprenoids
precursors with consequences on the post-translational modification and targeting of proteins of the
peripheral membrane system.

Discussion

Beside components of the SUF Fe-S cluster synthesis machinery, the apicoplast harbors only a small
number of putative Fe-S proteins (Fig. 9, (12)). Yet, they are all presumably important for parasite
fitness as suggested by their negative scores with a CRISPR-based genome-wide screen (Fig. 9, (38)).
However, not all are expected to be absolutely essential *in vitro*, tRNA modifying enzyme MiaB for
instance, has only a moderately low fitness score and has been shown recently in to be dispensable
for *Plasmodium* intraerythrocytic stages (37). Similarly, the LipA lipoyl synthase essential for the
function of the E2 subunit of the PDH complex, which is in turn crucial for generating the acetyl-CoA
necessary for de novo FA synthesis in the apicoplast, seems dispensable for *Plasmodium* blood stages
(37), for which the FASII system is not essential (58). However, given the potentially greater
importance of FASII in *T. gondii* tachyzoites (41), this is something we investigated further. We
demonstrated disruption of the SUF pathway impaired LipA function and PDH-E2 lipoylation (Fig. 5)
and, likely as a consequence of this and FASII perturbation, general production of myristic and
palmitic acid in the parasites (Fig. 6). However, the growth defect of SUF mutants could only be
partially complemented by FA supplementation for the TgNFS2 mutant, and not at all for the TgSUFC
mutant (Fig. 7), suggesting perturbation of FASII is not the primary cause of death for these mutants.
While it is undoubtedly a metabolic pathway that plays a central role for parasite fitness, the view on
the essentiality of FASII in *T. gondii* tachyzoites has recently evolved. There is now published
evidence that parasites can survive *in vitro* even when FASII enzymes are inactivated (44, 59) and
there is clearly flexibility in the adaptation to lipid sources (60). As tachyzoites can readily scavenge
and incorporate FAs from exogenous sources into their own range of lipids (43, 44, 61), in the end
the essentiality of the FASII pathway depends largely on the culture conditions.

The other main metabolic pathway that depends directly on Fe-S proteins in the apicoplast is for the
synthesis of isoprenoid precursors. When their isoprenoid production is inhibited, *T. gondii*
tachyzoites can scavenge some of these precursors from the host cell, leading to a delayed death
effect (49, 62), but that cannot fully compensate for a lack of de novo synthesis. Isoprenoids are a
large and diverse class of lipids whose cellular functions in Apicomplexa still remain to be extensively
characterised, but they include synthesis of vitamins and cofactors (ubiquinone), and are also
involved in important posttranslational modifications of proteins, like prenylation, GPI-anchoring and
glycosylation (Fig. 9, (32)). Ubiquinone is a central molecule in the mitochondrial electron transport
chain (ETC): its quinone head is the functional group for the transfer of electrons, whereas the
isoprenoid tail primarily serves for anchoring to the inner mitochondrial membrane. The
mitochondrial ETC is a validated drug target in Apicomplexa, for which complex III inhibitor
atovaquone has been used in therapeutic strategies (63). However, while *T. gondii* mitochondrial ETC
mutants are severely impaired in growth (64–67), it seems that genetic or pharmaceutical
inactivation (with atovaquone) is reversible ((12, 66)), and may lead to stage conversion and a

metabolic dormant state rather than complete death of the parasites. On the contrary, the viability
of the SUF mutants is irreversibly affected (Fig. 3C and (12)). So although the impact SUF protein
depletion on the isoprenoid pathway is likely to lead to a deficiency in ubiquinone synthesis, which in
turn would contribute to a decrease in parasite fitness, this is probably not the main reason for the
irreversible death of the parasites.

In *Plasmodium* blood stages, where isoprenoid synthesis is the only essential pathway hosted by the
apicoplast, disrupting the prenylation of Rab GTPAses, which are involved in vesicular trafficking,
contributes to delayed death (50). In *T. gondii*, interestingly, perturbation of Rab function can lead to
intracellular accumulation and patchy surface distribution of SAG1 proteins, and results in defects in
the delivery of new membrane required for completing daughter cell segregation at the end of
cytokinesis (68, 69). This bears some similarity with some phenotypes we have observed in the SUF
mutants, yet while we cannot completely exclude Rab prenylation is perturbed in the SUF mutants,
we did not identify any particular prenylation problem in the parasites, and we failed to complement
their growth defect with prenylation precursor GGOH (Fig. S5). We thus investigated other important
isoprenoid-dependent protein modifications. For instance, upon sugar addition dolichol can be used
for the formation of GPI anchors, or act as a donor for protein glycosylation. Our phenotypic analysis
revealed that the targeting of GPI-anchored surface proteins and the gliding motility of the parasites,
which relies on glycosylated proteins, were clearly affected upon disruption of the SUF pathway (Fig.
8). This confirms the importance of the apicoplast Fe-S cluster synthesis machinery for isoprenoid
metabolism. In *Plasmodium*, interfering with apicoplast-hosted isoprenoid production affects the
morphology of the organelle (70) but depletion of IspG and IspH does not lead to loss of the
apicoplast (37), while interfering with the SUF pathway does (13). We also observe a late impact on
the organelle (Fig. 5), that suggest the phenotypic consequences of SUF proteins in *T. gondii* are
indeed multifactorial and would extend beyond the simple disruption of the isoprenoid pathway.

At the cellular level, one of the most visible consequence of long term depletion of SUF proteins is
the membrane defects in the late stages of cytokinesis (Fig. 4). Interestingly, treatment of tachyzoites
with the FASII inhibitor triclosan or inactivating the FASII component acyl carrier protein was shown
to lead to severe problems in cytokinesis completion, with tethered daughter cells resembling the
phenotype we have described here (46). A similar phenotype was also observed in the mutant for
TgATS2, an apicoplast-located acyltransferase (43). This points to a central role for the apicoplast to
provide specific precursors for membrane biogenesis during cytokinesis, and to a SUF-dependent
FASII function is important for the homeostasis of the parasite plasma membrane. It should however
be noted that some isoprenoid-dependent cellular mediators may also contribute to plasma
membrane synthesis during cell division. For instance, disruption of Rab-controlled vesicular
trafficking, leads to very similar phenotypes of incomplete cytokinesis, with tachyzoites still fused
along their lateral surface (68, 69). Glycosylated IMC proteins associated with gliding motility are also
important for IMC formation and the cell division process (71). It is also possible that yet unidentified
GPI-anchored *T. gondii* proteins may be involved in plasma membrane formation or recycling: GPI
synthesis is essential for *T. gondii* survival (72), but the function of individual GPI-anchored proteins
remain largely overlooked. The detrimental effects of SUF depletion on plasma membrane
homeostasis may thus manifest through both FASII and isoprenoid perturbation. Moreover, as some
isoprenoid-dependent modifications are also linked to FA acid synthesis, like GPI anchors of *T. gondii*
surface proteins that also necessitate phospholipid moieties (73), a simultaneous impact on the two
pathways may enhance the phenotypic output.

One key apicoplast-located Fe-S protein is Fd, which has a central role in the function of Fe-S-
dependent apicoplast enzymes: it is potentially providing electrons to other apicoplast Fe-S enzymes

like MiaB, IspG, IspH and LipA. The role of Fd has recently been investigated in Apicomplexa. In
Plasmodium blood stage, the loss of parasite viability upon Fd depletion was likely mostly due to the
importance of Fd for the isoprenoid synthesis pathway (37), which is the only essential apicoplast-
located pathway in this developmental stage. Fd is equally essential for *T. gondii*, tachyzoite survival,
where it was shown to impact both FASII and isoprenoid synthesis (35), in a similar fashion to our
SUF mutants. Whether the vital importance of Fe-S cluster synthesis and associated apicoplast redox
metabolism is solely through its key role in isoprenoid synthesis is thus less clear in *T. gondii* than in
Plasmodium. As *T. gondii* tachyzoites grow in host cell types that can potentially provide them with
more resources, the ability to scavenge exogenous metabolites creates a complex situation whereby
metabolic pathways like FASII may be only essential in certain particular conditions. The nutrient-rich
in vitro culture systems may also mask some important contributions. In any case, because of their
upstream role in cellular functions important for parasite fitness, Fd and SUF mutants clearly have
pleiotropic effects. More importantly, we have confirmed here that disrupting the SUF machinery
leads to an irreversible death of the tachyzoites, which is not the case, for example, when the
mitochondrial Fe-S cluster machinery is inactivated (12). For all these reasons, and also because of its
absence from mammalian hosts of the parasite, the SUF pathway has a strong potential identifying
novel drug targets.

Experimental procedures

Parasites and cells culture. Tachyzoites of the TATi ΔKu80 *T. gondii* strain (25), as well as derived
transgenic parasites generated in this study, were maintained by serial passage in monolayers of
human foreskin fibroblast (HFF, American Type Culture Collection, CRL 1634) grown in Dulbecco's
modified Eagle medium (Gibco), supplemented with 5% decomplemented fetal bovine serum, 2-mM
L-glutamine and a cocktail of penicillin-streptomycin at 100 μg/ml.

Bioinformatic analyses. Sequence alignment was performed using the MUltiple Sequence
Comparison by Log-Expectation (MUSCLE) algorithm of the Geneious 6.1.8 software suite
(http://www.geneious.com). Transit peptide and localization prediction was done with the Deeploc
1.0 (http://www.cbs.dtu.dk/services/DeepLoc-1.0/) algorithm.

Heterologous expression in *E. coli*. Construct for designing the recombinant protein used for *E. coli*
complementation was defined by aligning the amino acid sequences of TgSUFC with its *E. coli*
counterparts. A 894 bp fragment corresponding to amino acids 220-518, was amplified by
polymerase chain reaction (PCR) from *T. gondii* cDNA using primers ML4200/ML4010 (sequences of
the primers used in this study are found in Table S1). The fragment was cloned into the pUC19
plasmid (Thermo Fisher Scientific) using the HindIII/BamHI restriction sites. The SufC *E. coli* mutant
from the Keio collection (obtained from the The *Coli* Genetic Stock Center at the University of Yale:
stain number JW1672-1), was transformed with the plasmid expressing the TgSUFC recombinant
protein and selected with ampicillin. For growth assays (22), overnight stationary phase cultures
were adjusted to the same starting OD_{600} of 0.6 in salt-supplemented M9 minimal media containing
0.4% glucose and varying amounts of the 2,2⊡-Bipyridyl iron chelator (Sigma-Aldrich). Growth was
monitored through OD_{600} measurement after 7, 14 and 24 hours at 37°C in a shaking incubator.

Generation of the HA-tagged TgSUFC cell line.

A CRISPR-based strategy was used. Using the pLIC-HA$_3$-CAT plasmid as a template, a PCR was performed with the KOD DNA polymerase (Novagen) to amplify the tag and the resistance gene expression cassette with primers ML3980/ML3981, that also carry 30⬚bp homology with the 3⬚ end of the corresponding genes. A specific single-guide RNA (sgRNA) was generated to introduce a double-stranded break at the 3⬚ of the *TgSUFC* gene, using primers ML3952/ML3953, and the protospacer sequences were introduced in the Cas9-expressing pU6-Universal plasmid (Addgene, ref #52694) (38). The TATi ΔKu80 cell line was transfected and transgenic parasites were selected with chloramphenicol and cloned by serial limiting dilution.

Generation of TgSUFC conditional knock-down and complemented cell lines. The conditional knock-down cell line for *TgSUFC* was generated based on the Tet-Off system using the DHFR-TetO7Sag4 plasmid (74) using a CRISPR-based strategy. Using the DHFR-TetO7Sag4 plasmid as a template, a PCR was performed with the KOD DNA polymerase (Novagen) to amplify the promoter and the resistance gene expression cassette with primers ML4107/ML4108 that also carry 30⬚bp homology with the 5⬚ end of the *TgSUFC* gene. A specific single-guide RNA (sgRNA) was generated to introduce a double-stranded break at the 5⬚ of the *TgSUFC* locus. Primers used to generate the guide were ML4109/ML4110 and the protospacer sequences were introduced in the pU6-Universal plasmid (Addgene ref#52694) (38). The TgSUFC-HA cell line was transfected with the donor sequence and the Cas9/guide RNA-expressing plasmid and transgenic parasites were selected with pyrimethamine and cloned by serial limiting dilution.

The cKD TgSUFC-HA cell line was complemented by the addition of an extra copy of the *TgSUFC* gene put under the dependence of a tubulin promoter at the *uracil phosphoribosyltransferase* (*UPRT*) locus. *TgSUFC* (1,557 bp) whole cDNA sequence was amplified by reverse transcription (RT)-PCR with primers ML4815/ML4816. They were then cloned downstream of the *tubulin* promoter sequence of the pUPRT-TUB-Ty vector (25) to yield the pUPRT-TgSUFC plasmid. This plasmid was then linearized prior to transfection of the mutant cell line. The recombination efficiency was increased by co-transfecting with the Cas9-expressing pU6-UPRT plasmids generated by integrating *UPRT*-specific protospacer sequences (with primers ML2087/ML2088 for the 3' and primers ML3445/ML3446 for the 5') which were designed to allow a double-strand break at the *UPRT* locus. Transgenic parasites were selected using 5-fluorodeoxyuridine and cloned by serial limiting dilution to yield the cKD TgSUFC-HA comp cell line.

Anti-TgPDH-E2 antibody production. A polyclonal antibody was raised in rabbit against a peptide (ISLIQAKGLSLISASSSPA) specific of TgPDH-E2 by the Proteogenix company. The peptide was conjugated to Keyhole limpet haemocyanin carrier protein prior to immunization and the whole serum was affinity-purified against the peptide for increased specificity.

Immunoblot analysis. Protein extracts from 10^7 freshly egressed tachyzoites were prepared in Laemmli sample buffer, separated by SDS-PAGE and transferred onto nitrocellulose membrane using the BioRad Mini-Transblot system according to the manufacturer's instructions. Rat monoclonal antibody (clone 3F10, Roche) was used to detect HA-tagged proteins. Other primary antibodies used were mouse monoclonal anti-TY tag (27), rabbit anti-lipoic acid antibody (ab58724, Abcam), mouse anti-SAG1 (75), rabbit anti-CPN60 (76), mouse anti-actin (77), and rabbit anti-farnesyl polyclonal antibody (PA1-12554, Life Technologies).

Immunofluorescence microscopy. For immunofluorescence assays (IFA), intracellular tachyzoites grown on coverslips containing HFF monolayers, were either fixed for 20 min with 4% (w/v) paraformaldehyde in PBS and permeabilized for 10 min with 0.3% Triton X-100 in PBS or fixed for 5 min in cold methanol (for SAG labelling). Slides/coverslips were subsequently blocked with 0.1% (w/v) BSA in PBS. Primary antibodies used (at 1/1,000, unless specified) were rat anti-HA tag (clone 3F10, Roche), mouse anti-TY tag (27), rabbit anti-CPN60 (76), rabbit anti-IMC3 (78), mouse anti-SAG1 (75), and mouse anti-SAG3 (79). Staining of DNA was performed on fixed cells by incubating them for 5 min in a 1 µg/ml 4,6-diamidino-2-phenylindole (DAPI) solution. All images were acquired at the Montpellier RIO imaging facility from a Zeiss AXIO Imager Z2 epifluorescence microscope driven by the ZEN software v2.3 (Zeiss). Z-stack acquisition and maximal intensity projection was performed to quantify apicoplast loss. Adjustments for brightness and contrast were applied uniformly on the entire image.

Electron Microscopy. Parasites were pretreated for three days with ATc, and then used to infect HFF monolayers and grown for an extra 24 hours in ATc. They were fixed with 2.5% glutaraldehyde in cacodylate buffer 0.1 M pH7.4. Coverslips were then processed using a Pelco Biowave pro+ (Ted Pella). Briefly, samples were postfixed in 1% OsO_4 and 2% uranyl acetate, dehydrated in acetonitrile series and embedded in Epon 118 using the following parameters: Glutaraldehyde (150 W ON/OFF/ON 1-min cycles); two buffer washes (40 s 150 W); OsO_4 (150 W ON/OFF/ON/OFF/ON 1-min cycles); two water washes (40 s 150 W); uranyl acetate (100 W ON/OFF/ON 1-min cycles); dehydration (40 s 150 W); resin infiltration (350 W 3-min cycles). Fixation and infiltration steps were performed under vacuum. Polymerization was performed at 60°C for 48 hr. Ultrathin sections at 70 nM were cut with a Leica UC7 ultramicrotome, counterstained with uranyl acetate and lead citrate and observed in a Jeol 1200 EXII transmission electron microscope from the MEA Montpellier Electron Microscopy Platform. All chemicals were from Electron Microscopy Sciences, and solvents were from Sigma.

Plaque assay. Confluent monolayers of HFFs were infected with freshly egressed parasites, which were left to grow for 7 days in the absence or presence of ATc (unless stated). For some experiments, the medium was supplemented with 50 µM palmitic acid (P0500, Sigma-Aldrich), 50 µM myristic acid (70082, Sigma-Aldrich), or 20 µM geranylgeranyol (G3278, Sigma-Aldrich). They were then fixed with 4% v/v paraformaldehyde (PFA) and plaques were revealed by staining with a 0.1% crystal violet solution (V5265, Sigma-Aldrich).

Gliding assay. 10^7 freshly egressed parasites were resuspended in 300 µl of motility buffer (Ringer's solution: 155 mM NaCl, 3 mM KCl, 2 mM $CaCl_2$, 1 mM $MgCl_2$, 3 mM NaH_2PO_4, 10 mM HEPES, 10 mM glucose). 100 µl were deposited on poly-L-lysine coated microscope slides (J2800AMNZ, Thermo Scientific), in a well delineated with a hydrophobic pen (PAP Pen, Kisker Biotech). Parasites were left to glide for 15 minutes in an incubator at 37°C, then the suspension was carefully removed and parasites were fixed with 4% (w/v) paraformaldehyde in PBS. Immunostaining was performed with an anti-SAG1 antibody (75) as described above, but without permeabilization. Trail deposition images were acquired with a 63x objective on a Zeiss AXIO Imager Z2 epifluorescence microscope and processed with ImageJ v. 1.53f51, using the NeuronJ plugin as described previously (35).

Lipidomic analyses. Parasite lipidomic analyses were conducted as previously described (43, 80). Briefly, the parasites were grown for 72 h in +/- ATc conditions within a confluent monolayer of HFF in flasks (175 cm^2). At each time point, parasites were harvested as intracellular tachyzoites (1×10^7 cell equivalents per replicate) after syringe filtration with 3-µm pore size membrane. These parasites were metabolically quenched by rapid chilling in a dry ice-ethanol slurry bath and then centrifuged down at 4°C. The parasite pellet was washed with ice-cold PBS thrice, before transferring the final pellet to a microcentrifuge tube. Then total lipids were extracted in chloroform/methanol/water (1:3:1, v/v/v) containing PC (C13:0/C13:0), 10 nmol and C21:0 (10 nmol) as internal standards for extraction. Polar and apolar metabolites were separated by phase partitioning by adding chloroform and water to give the ratio of chloroform/methanol/water as 2:1:0.8 (v/v/v). For lipid analysis, the organic phase was dried under N$_2$ gas and dissolved in 1-butanol to obtain 1µl butanol/10^7 parasites.

Total lipid analysis – The extracted total lipid sample was then added with 1 nmol pentadecanoic acid (C15:0) as internal standard as stated before using Trimethylsulfonium hydroxide for total FA content. Resultant FA methyl esters (FAMEs) were analyzed by GC-MS as previously described (45). All FAMEs were identified by comparison of retention time and mass spectra from GC-MS with authentic chemical standards. The concentration of FAMEs was quantified after initial normalization to different internal standards and finally to parasite number.

Stable isotope metabolic labeling experiment.

Tracking host-derived FAs – (monitoring parasite scavenging capacities)

The HFF cells were grown (1×10^8 cell equivalents per replicate) to confluency in the presence of stable isotope U-^{13}C-glucose at a final concentration of 800 µM added to a glucose-free DMEM. These ^{13}C-pre labelled HFF were then infected with *Tg*LIPIN-ikD parasites in the presence of normal-glucose containing DMEM under +/-ATc (0.5 µg/ml). The host HFF and parasites were metabolically quenched separately, and their lipid content was quantified by GC-MS as described above. ^{13}C incorporation to each FA was calculated as the percent of the metabolite pool containing one or more ^{13}C atoms after correction for natural abundance and the amount of ^{13}C-carbon source in the culture medium. The degree of the incorporation of ^{13}C into FAs (%carbon incorporation) was determined by the mass isotopomer distribution of each FAMEs. Mass isotopomer distribution was obtained from the shift in isotopic mass dependent on the amount of ^{12}C carbons compared to the integration of ^{13}C carbon atoms. The total abundance of ^{13}C-labelled FAs was obtained by calculating the concentration of all isotopomers of ^{13}C-labelled FAMEs and finally normalizing to authentic internal standards and parasite number. The total abundance of ^{13}C-labelled FAs was analyzed initially for HFF to check labelling of the metabolites (described previously). Later, the same was calculated for parasites to confirm direct uptake of ^{13}C-labelled fatty acids from the host.

Data availability

All data are contained within the manuscript. Material described is available upon request from the corresponding author.

Supporting information

This article contains supporting information.

Acknowledgments

We are grateful to B. Striepen, L. Sheiner, D. Soldati-Favre, M.J. Gubbels, V. Carruthers, and J.F.
Dubremetz for sharing antibodies, strains and plasmids. We wish to thank K. Semenovskaya for
technical help with some constructs. We also thank the Montpellier Rio Imaging facility for providing
access to their microscopes, as well as the electron microscopy imaging facility of the University of
Montpellier.

Funding and additional information

AC was supported by a fellowship from the Fondation pour la Recherche Médicale (Equipe FRM
EQ20170336725). SB and CYB acknowledge support from the Labex Parafrap (ANR-11-LABX-0024),
the Agence Nationale de la Recherche (ANR-21-CE44-0010 to CYB and SB, and ANR-19-CE15-0023 to
SB). The funders had no role in study design, data collection and analysis, decision to publish, or
preparation of the manuscript.

Conflict of interest

The authors declare that they have no conflicts of interest with the contents of this article.

References

1. White, N. J., Pukrittayakamee, S., Hien, T. T., Faiz, M. A., Mokuolu, O. A., and Dondorp, A. M.
(2014) Malaria. *Lancet*. **383**, 723–735
2. Sanchez, S. G., and Besteiro, S. (2021) The pathogenicity and virulence of Toxoplasma gondii.
Virulence. **12**, 3095–3114
3. Janouskovec, J., Horak, A., Obornik, M., Lukes, J., and Keeling, P. J. (2010) A common red algal
origin of the apicomplexan, dinoflagellate, and heterokont plastids. *Proceedings of the National*
Academy of Sciences. **107**, 10949–10954
4. van Dooren, G. G., and Striepen, B. (2013) The algal past and parasite present of the apicoplast.
Annu. Rev. Microbiol. **67**, 271–289
5. van Dooren, G. G., and Hapuarachchi, S. V. (2017) The dark side of the chloroplast: biogenesis,
metabolism and membrane biology of the apicoplast. in *Advances in Botanical Research*, pp.
145–185, Elsevier, **84**, 145–185
6. Seeber, F., and Soldati-Favre, D. (2010) Metabolic pathways in the apicoplast of apicomplexa. in
International Review of Cell and Molecular Biology, pp. 161–228, **281**, 161–228
7. Biddau, M., and Sheiner, L. (2019) Targeting the apicoplast in malaria. *Biochem Soc Trans*. **47**,
973–983
8. Imlay, J. A. (2006) Iron-sulphur clusters and the problem with oxygen. *Mol Microbiol*. **59**, 1073–
1082
9. Lill, R. (2009) Function and biogenesis of iron–sulphur proteins. *Nature*. **460**, 831–838

10.   Braymer, J. J., Freibert, S. A., Rakwalska-Bange, M., and Lill, R. (2021) Mechanistic concepts of
614   iron-sulfur protein biogenesis in Biology. *Biochimica et Biophysica Acta (BBA) - Molecular Cell*
615   *Research*. **1868**, 118863

616   11.   Aw, Y. T. V., Seidi, A., Hayward, J. A., Lee, J., Victor Makota, F., Rug, M., and van Dooren, G. G.
617   (2020) A key cytosolic iron-sulfur cluster synthesis protein localises to the mitochondrion of
618   Toxoplasma gondii. *Mol Microbiol*. 10.1111/mmi.14651

619   12.   Pamukcu, S., Cerutti, A., Bordat, Y., Hem, S., Rofidal, V., and Besteiro, S. (2021) Differential
620   contribution of two organelles of endosymbiotic origin to iron-sulfur cluster synthesis and
621   overall fitness in Toxoplasma. *PLoS Pathog*. **17**, e1010096

622   13.   Gisselberg, J. E., Dellibovi-Ragheb, T. A., Matthews, K. A., Bosch, G., and Prigge, S. T. (2013) The
623   Suf iron-sulfur cluster synthesis pathway is required for apicoplast maintenance in malaria
624   parasites. *PLoS Pathog*. **9**, e1003655

625   14.   Haussig, J. M., Matuschewski, K., and Kooij, T. W. A. (2014) Identification of vital and
626   dispensable sulfur utilization factors in the Plasmodium apicoplast. *PLoS ONE*. **9**, e89718

627   15.   Kumar, B., Chaubey, S., Shah, P., Tanveer, A., Charan, M., Siddiqi, M. I., and Habib, S. (2011)
628   Interaction between sulphur mobilisation proteins SufB and SufC: Evidence for an iron–sulphur
629   cluster biogenesis pathway in the apicoplast of Plasmodium falciparum. *International Journal*
630   *for Parasitology*. **41**, 991–999

631   16.   Charan, M., Singh, N., Kumar, B., Srivastava, K., Siddiqi, M. I., and Habib, S. (2014) Sulfur
632   mobilization for Fe-S cluster assembly by the essential SUF pathway in the Plasmodium
633   falciparum apicoplast and its Inhibition. *Antimicrob. Agents Chemother*. **58**, 3389–3398

634   17.   Charan, M., Choudhary, H. H., Singh, N., Sadik, M., Siddiqi, M. I., Mishra, S., and Habib, S. (2017)
635   [Fe-S] cluster assembly in the apicoplast and its indispensability in mosquito stages of the
636   malaria parasite. *FEBS J*. **284**, 2629–2648

637   18.   Takahashi, Y., and Tokumoto, U. (2002) A third bacterial system for the assembly of Iron-Sulfur
638   clusters with homologs in archaea and plastids. *J. Biol. Chem*. **277**, 28380–28383

639   19.   Hirabayashi, K., Yuda, E., Tanaka, N., Katayama, S., Iwasaki, K., Matsumoto, T., Kurisu, G.,
640   Outten, F. W., Fukuyama, K., Takahashi, Y., and Wada, K. (2015) Functional dynamics revealed
641   by the structure of the SufBCD complex, a novel ATP-binding cassette (ABC) protein that serves
642   as a scaffold for Iron-Sulfur cluster biogenesis. *Journal of Biological Chemistry*. **290**, 29717–
643   29731

644   20.   Harb, O. S., and Roos, D. S. (2020) ToxoDB: functional genomics resource for Toxoplasma and
645   related organisms. *Methods Mol Biol*. **2071**, 27–47

646   21.   Barylyuk, K., Koreny, L., Ke, H., Butterworth, S., Crook, O. M., Lassadi, I., Gupta, V., Tromer, E.,
647   Mourier, T., Stevens, T. J., Breckels, L. M., Pain, A., Lilley, K. S., and Waller, R. F. (2020) A
648   comprehensive subcellular atlas of the Toxoplasma proteome via hyperLOPIT provides spatial
649   context for protein functions. *Cell Host & Microbe*. 10.1016/j.chom.2020.09.011

650   22.   Outten, F. W., Djaman, O., and Storz, G. (2004) A suf operon requirement for Fe-S cluster
651   assembly during iron starvation in Escherichia coli: suf operon role during iron starvation.
652   *Molecular Microbiology*. **52**, 861–872

653   23.   Fox, B. A., Ristuccia, J. G., Gigley, J. P., and Bzik, D. J. (2009) Efficient gene replacements in
654   Toxoplasma gondii strains deficient for nonhomologous end joining. *Eukaryotic Cell*. **8**, 520–529

655   24.   Huynh, M.-H., and Carruthers, V. B. (2009) Tagging of endogenous genes in a Toxoplasma
656   gondii strain lacking Ku80. *Eukaryot Cell*. **8**, 530–539

657   25.   Sheiner, L., Demerly, J. L., Poulsen, N., Beatty, W. L., Lucas, O., Behnke, M. S., White, M. W., and
658   Striepen, B. (2011) A Systematic Screen to Discover and Analyze Apicoplast Proteins Identifies a
659   Conserved and Essential Protein Import Factor. *PLoS Pathog*. 10.1371/journal.ppat.1002392

660   26.   Meissner, M., Brecht, S., Bujard, H., and Soldati, D. (2001) Modulation of myosin A expression
661   by a newly established tetracycline repressor-based inducible system in Toxoplasma gondii.
662   *Nucleic Acids Res*. **29**, E115

663   27.   Bastin, P., Bagherzadeh, A., Matthews, K. R., and Gull, K. (1996) A novel epitope tag system to
664   study protein targeting and organelle biogenesis in Trypanosoma brucei. *Molecular and*
665   *Biochemical Parasitology*. **77**, 235–239

666   28.   Francia, M. E., and Striepen, B. (2014) Cell division in apicomplexan parasites. *Nat. Rev.*
667   *Microbiol.* **12**, 125–136

668   29.   Harding, C. R., and Meissner, M. (2014) The inner membrane complex through development of
669   Toxoplasma gondii and Plasmodium. *Cell Microbiol.* **16**, 632–641

670   30.   Ouologuem, D. T., and Roos, D. S. (2014) Dynamics of the Toxoplasma gondii inner membrane
671   complex. *J Cell Sci.* **127**, 3320–3330

672   31.   Gubbels, M.-J., Wieffer, M., and Striepen, B. (2004) Fluorescent protein tagging in Toxoplasma
673   gondii: identification of a novel inner membrane complex component conserved among
674   Apicomplexa. *Molecular and Biochemical Parasitology*. **137**, 99–110

675   32.   Imlay, L., and Odom, A. R. (2014) Isoprenoid metabolism in apicomplexan parasites. *Curr Clin*
676   *Microbiol Rep*. **1**, 37–50

677   33.   Thomsen-Zieger, N., Schachtner, J., and Seeber, F. (2003) Apicomplexan parasites contain a
678   single lipoic acid synthase located in the plastid. *FEBS Lett*. **547**, 80–86

679   34.   Pierrel, F., Douki, T., Fontecave, M., and Atta, M. (2004) MiaB protein is a bifunctional radical-S-
680   adenosylmethionine enzyme involved in thiolation and methylation of tRNA. *J Biol Chem*. **279**,
681   47555–47563

682   35.   Henkel, S., Frohnecke, N., Maus, D., McConville, M. J., Laue, M., Blume, M., and Seeber, F.
683   (2022) Toxoplasma gondii apicoplast-resident ferredoxin is an essential electron transfer
684   protein for the MEP isoprenoid-biosynthetic pathway. *J Biol Chem*. **298**, 101468

685   36.   Frohnecke, N., Klein, S., and Seeber, F. (2015) Protein–protein interaction studies provide
686   evidence for electron transfer from ferredoxin to lipoic acid synthase in *Toxoplasma gondii*.
687   *FEBS Letters*. **589**, 31–36

688   37.   Swift, R. P., Rajaram, K., Elahi, R., Liu, H. B., and Prigge, S. T. (2022) Roles of ferredoxin-
689   dependent proteins in the apicoplast of Plasmodium falciparum parasites. *mBio*.
690   10.1128/mbio.03023-21

691   38.   Sidik, S. M., Huet, D., Ganesan, S. M., Huynh, M.-H., Wang, T., Nasamu, A. S., Thiru, P., Saeij, J.
692   P. J., Carruthers, V. B., Niles, J. C., and Lourido, S. (2016) A Genome-wide CRISPR Screen in
693   Toxoplasma Identifies Essential Apicomplexan Genes. *Cell*. **166**, 1423-1435.e12

694   39.   Kennedy, K., Crisafulli, E. M., and Ralph, S. A. (2019) Delayed death by plastid inhibition in
695   apicomplexan parasites. *Trends in Parasitology*. **35**, 747–759

696   40.   Jelenska, J., Crawford, M. J., Harb, O. S., Zuther, E., Haselkorn, R., Roos, D. S., and Gornicki, P.
697   (2001) Subcellular localization of acetyl-CoA carboxylase in the apicomplexan parasite
698   Toxoplasma gondii. *Proceedings of the National Academy of Sciences*. **98**, 2723–2728

699   41.   Mazumdar, J., H Wilson, E., Masek, K., A Hunter, C., and Striepen, B. (2006) Apicoplast fatty acid
700   synthesis is essential for organelle biogenesis and parasite survival in Toxoplasma gondii. *Proc.*
701   *Natl. Acad. Sci. U.S.A.* **103**, 13192–13197

702   42.   Ramakrishnan, S., Docampo, M. D., Macrae, J. I., Pujol, F. M., Brooks, C. F., van Dooren, G. G.,
703   Hiltunen, J. K., Kastaniotis, A. J., McConville, M. J., and Striepen, B. (2012) Apicoplast and
704   endoplasmic reticulum cooperate in fatty acid biosynthesis in apicomplexan parasite
705   Toxoplasma gondii. *J. Biol. Chem.* **287**, 4957–4971

706   43.   Amiar, S., Katris, N. J., Berry, L., Dass, S., Duley, S., Arnold, C.-S., Shears, M. J., Brunet, C.,
707   Touquet, B., McFadden, G. I., Yamaryo-Botté, Y., and Botté, C. Y. (2020) Division and adaptation
708   to host environment of apicomplexan parasites depend on apicoplast lipid metabolic plasticity
709   and host organelle remodeling. *Cell Reports*. **30**, 3778-3792.e9

710   44.   Liang, X., Cui, J., Yang, X., Xia, N., Li, Y., Zhao, J., Gupta, N., and Shen, B. (2020) Acquisition of
711   exogenous fatty acids renders apicoplast-based biosynthesis dispensable in tachyzoites of
712   Toxoplasma. *J Biol Chem*. **295**, 7743–7752

45. Dubois, D., Fernandes, S., Amiar, S., Dass, S., Katris, N. J., Botté, C. Y., and Yamaryo-Botté, Y.
(2018) Toxoplasma gondii acetyl-CoA synthetase is involved in fatty acid elongation (of long
fatty acid chains) during tachyzoite life stages. *J. Lipid Res.* **59**, 994–1004
46. Martins-Duarte, É. S., Carias, M., Vommaro, R., Surolia, N., and de Souza, W. (2016) Apicoplast
fatty acid synthesis is essential for pellicle formation at the end of cytokinesis in Toxoplasma
gondii. *J Cell Sci.* **129**, 3320–3331
47. Yeh, E., and DeRisi, J. L. (2011) Chemical rescue of malaria parasites lacking an apicoplast
defines organelle function in blood-stage Plasmodium falciparum. *PLoS Biol.* **9**, e1001138
48. Nair, S. C., Brooks, C. F., Goodman, C. D., Sturm, A., Strurm, A., McFadden, G. I., Sundriyal, S.,
Anglin, J. L., Song, Y., Moreno, S. N. J., and Striepen, B. (2011) Apicoplast isoprenoid precursor
synthesis and the molecular basis of fosmidomycin resistance in Toxoplasma gondii. *J. Exp.*
Med. **208**, 1547–1559
49. Li, Z.-H., Ramakrishnan, S., Striepen, B., and Moreno, S. N. J. (2013) Toxoplasma gondii relies on
both host and parasite isoprenoids and can be rendered sensitive to Atorvastatin. *PLoS Pathog.*
9, e1003665
50. Kennedy, K., Cobbold, S. A., Hanssen, E., Birnbaum, J., Spillman, N. J., McHugh, E., Brown, H.,
Tilley, L., Spielmann, T., McConville, M. J., and Ralph, S. A. (2019) Delayed death in the malaria
parasite Plasmodium falciparum is caused by disruption of prenylation-dependent intracellular
trafficking. *PLoS Biol.* **17**, e3000376
51. Li, Z.-H., King, T. P., Ayong, L., Asady, B., Cai, X., Rahman, T., Vella, S. A., Coppens, I., Patel, S.,
and Moreno, S. N. J. (2021) A plastid two-pore channel essential for inter-organelle
communication and growth of Toxoplasma gondii. *Nat Commun.* **12**, 5802
52. Suazo, K. F., Schaber, C., Palsuledesai, C. C., Odom John, A. R., and Distefano, M. D. (2016)
Global proteomic analysis of prenylated proteins in Plasmodium falciparum using an alkyne-
modified isoprenoid analogue. *Sci Rep.* **6**, 38615
53. Gisselberg, J. E., Zhang, L., Elias, J. E., and Yeh, E. (2017) The prenylated proteome of
Plasmodium falciparum reveals pathogen-specific prenylation activity and drug mechanism-of-
action. *Mol Cell Proteomics.* **16**, S54–S64
54. Jung, C., Lee, C. Y.-F., and Grigg, M. E. (2004) The SRS superfamily of Toxoplasma surface
proteins. *International Journal for Parasitology.* **34**, 285–296
55. Striepen, B., Yingxin He, C., Matrajt, M., Soldati, D., and Roos, D. S. (1998) Expression, selection,
and organellar targeting of the green fluorescent protein in Toxoplasma gondii. *Molecular and*
Biochemical Parasitology. **92**, 325–338
56. Fauquenoy, S., Morelle, W., Hovasse, A., Bednarczyk, A., Slomianny, C., Schaeffer, C., Van
Dorsselaer, A., and Tomavo, S. (2008) Proteomics and glycomics analyses of N-glycosylated
structures involved in Toxoplasma gondii-host cell interactions. *Molecular & Cellular*
Proteomics. **7**, 891–910
57. Luk, F. C. Y., Johnson, T. M., and Beckers, C. J. (2008) N-linked glycosylation of proteins in the
protozoan parasite Toxoplasma gondii. *Mol Biochem Parasitol.* **157**, 169–178
58. Vaughan, A. M., O'Neill, M. T., Tarun, A. S., Camargo, N., Phuong, T. M., Aly, A. S. I., Cowman, A.
F., and Kappe, S. H. I. (2009) Type II fatty acid synthesis is essential only for malaria parasite late
liver stage development. *Cellular Microbiology.* **11**, 506–520
59. Xu, X.-P., Elsheikha, H. M., Liu, W.-G., Zhang, Z.-W., Sun, L.-X., Liang, Q.-L., Song, M.-X., and Zhu,
X.-Q. (2021) The role of type II fatty acid synthesis enzymes FabZ, ODSCI, and ODSCII in the
pathogenesis of Toxoplasma gondii infection. *Front Microbiol.* **12**, 703059
60. Shunmugam, S., Arnold, C.-S., Dass, S., Katris, N. J., and Botté, C. Y. (2022) The flexibility of
Apicomplexa parasites in lipid metabolism. *PLoS Pathog.* **18**, e1010313
61. Krishnan, A., Kloehn, J., Lunghi, M., Chiappino-Pepe, A., Waldman, B. S., Nicolas, D., Varesio, E.,
Hehl, A., Lourido, S., Hatzimanikatis, V., and Soldati-Favre, D. (2020) Functional and
computational genomics reveal unprecedented flexibility in stage-specific Toxoplasma
metabolism. *Cell Host & Microbe.* 10.1016/j.chom.2020.01.002

62. Amberg-Johnson, K., and Yeh, E. (2019) Host cell metabolism contributes to delayed-death
kinetics of apicoplast inhibitors in Toxoplasma gondii. *Antimicrob Agents Chemother.* **63**,
e01646-18

63. Baggish, A. L., and Hill, D. R. (2002) Antiparasitic agent atovaquone. *Antimicrob Agents*
Chemother. **46**, 1163–1173

64. Huet, D., Rajendran, E., van Dooren, G. G., and Lourido, S. (2018) Identification of cryptic
subunits from an apicomplexan ATP synthase. *eLife.* **7**, e38097

65. Seidi, A., Muellner-Wong, L. S., Rajendran, E., Tjhin, E. T., Dagley, L. F., Aw, V. Y., Faou, P.,
Webb, A. I., Tonkin, C. J., and van Dooren, G. G. (2018) Elucidating the mitochondrial proteome
of Toxoplasma gondii reveals the presence of a divergent cytochrome c oxidase. *Elife.*
10.7554/eLife.38131

66. Hayward, J. A., Rajendran, E., Zwahlen, S. M., Faou, P., and van Dooren, G. G. (2021) Divergent
features of the coenzyme Q:cytochrome c oxidoreductase complex in Toxoplasma gondii
parasites. *PLoS Pathog.* **17**, e1009211

67. Maclean, A. E., Bridges, H. R., Silva, M. F., Ding, S., Ovciarikova, J., Hirst, J., and Sheiner, L.
(2021) Complexome profile of Toxoplasma gondii mitochondria identifies divergent subunits of
respiratory chain complexes including new subunits of cytochrome bc1 complex. *PLoS Pathog.*
17, e1009301

68. Agop-Nersesian, C., Naissant, B., Ben Rached, F., Rauch, M., Kretzschmar, A., Thiberge, S.,
Menard, R., Ferguson, D. J. P., Meissner, M., and Langsley, G. (2009) Rab11A-controlled
assembly of the inner membrane complex is required for completion of apicomplexan
cytokinesis. *PLoS Pathog.* **5**, e1000270

69. Venugopal, K., Chehade, S., Werkmeister, E., Barois, N., Periz, J., Lafont, F., Tardieux, I., Khalife,
J., Langsley, G., Meissner, M., and Marion, S. (2020) Rab11A regulates dense granule transport
and secretion during Toxoplasma gondii invasion of host cells and parasite replication. *PLoS*
Pathog. **16**, e1008106

70. Okada, M., Rajaram, K., Swift, R. P., Mixon, A., Maschek, J. A., Prigge, S. T., and Sigala, P. A.
(2021) *Critical role for isoprenoids in apicoplast biogenesis by malaria parasites*, Microbiology,
10.1101/2021.08.19.456988

71. Harding, C. R., Egarter, S., Gow, M., Jiménez-Ruiz, E., Ferguson, D. J. P., and Meissner, M. (2016)
Gliding associated proteins play essential roles during the formation of the inner membrane
complex of Toxoplasma gondii. *PLoS Pathog.* **12**, e1005403

72. Wichroski, M. J., and Ward, G. E. (2003) Biosynthesis of glycosylphosphatidylinositol Is essential
to the survival of the protozoan parasite Toxoplasma gondii. *Eukaryot Cell.* **2**, 1132–1136

73. Tomavo, S., Schwarz, R. T., and Dubremetz, J. F. (1989) Evidence for glycosyl-
phosphatidylinositol anchoring of Toxoplasma gondii major surface antigens. *Mol. Cell. Biol.* **9**,
4576–4580

74. Morlon-Guyot, J., Berry, L., Chen, C.-T., Gubbels, M.-J., Lebrun, M., and Daher, W. (2014) The
Toxoplasma gondii calcium-dependent protein kinase 7 is involved in early steps of parasite
division and is crucial for parasite survival. *Cell. Microbiol.* **16**, 95–114

75. Couvreur, G., Sadak, A., Fortier, B., and Dubremetz, J. F. (1988) Surface antigens of Toxoplasma
gondii. *Parasitology.* **97 (Pt 1)**, 1–10

76. Agrawal, S., van Dooren, G. G., Beatty, W. L., and Striepen, B. (2009) Genetic evidence that an
endosymbiont-derived endoplasmic reticulum-associated protein degradation (ERAD) system
functions in import of apicoplast proteins. *J. Biol. Chem.* **284**, 33683–33691

77. Herm-Gotz, A. (2002) Toxoplasma gondii myosin A and its light chain: a fast, single-headed,
plus-end-directed motor. *The EMBO Journal.* **21**, 2149–2158

78. Anderson-White, B. R., Ivey, F. D., Cheng, K., Szatanek, T., Lorestani, A., Beckers, C. J., Ferguson,
D. J. P., Sahoo, N., and Gubbels, M.-J. (2011) A family of intermediate filament-like proteins is
sequentially assembled into the cytoskeleton of Toxoplasma gondii. *Cell. Microbiol.* **13**, 18–31

79.   Cesbron-Delauw, M. F., Tomavo, S., Beauchamps, P., Fourmaux, M. P., Camus, D., Capron, A.,
and Dubremetz, J. F. (1994) Similarities between the primary structures of two distinct major
surface proteins of Toxoplasma gondii. *J Biol Chem*. **269**, 16217–16222
80.   Dass, S., Shunmugam, S., Berry, L., Arnold, C.-S., Katris, N. J., Duley, S., Pierrel, F., Cesbron-
Delauw, M.-F., Yamaryo-Botté, Y., and Botté, C. Y. (2021) Toxoplasma LIPIN is essential in
channeling host lipid fluxes through membrane biogenesis and lipid storage. *Nat Commun*. **12**,
2813

Abbreviations

ABC (ATP-binding cassette), ATc (anhydrotetracycline), cKD (conditional knock-down), DMAPP
(dimethylallyl diphosphate), ER (endoplasmic reticulum), ETC (electron transport chain), FA (fatty
acid), FAME (fatty acid methyl ester), FASII (type II fatty acid synthase), FBS (fetal bovine serum), Fd
(ferredoxin), Fe-S (iron-sulfur), GC-MS (gas chromatography-mass spectrometry), GGOH
(geranylgeraniol), GPI (glycosylphosphatidylinositol), HA (hemagglutinin), HFF (human foreskin
fibroblasts), IFA (immunofluorescence assay), IMC (inner membrane complex), IPP (isopentenyl
diphosphate), ISC (iron–sulfur cluster), LipA (lipoic acid synthase A), LOPIT (localization of organelle
proteins by isotope tagging), PDH (pyruvate dehydrogenase), surface antigen (SAG), SUF (sulfur
utilization factor), TATi (tetracycline-inducible transactivator), UPRT (uracil
phosphoribosyltransferase)

Figure legends

Figure 1. The *Toxoplasma gondii* functional homolog of SufC. A) Schematic representation of the
molecular machinery for Fe-S cluster synthesis in the apicoplast of *T. gondii*. B) Alignment of the
predicted amino acid sequence of TgSUFC and its homolog from *Escherischia coli*. Motifs that are
potentially important for ATPase activity are outlined in blue. C) Functional complementation of
bacterial mutants for SufC. Growth of 'wild-type' (WT) *E. coli* K12 parental strain, the SufC mutant
strain and the mutant strain complemented ('comp') by the *T. gondii* homolog, was assessed by
monitoring the optical density at 600 nm in the presence or not of an iron chelator (2,2'-bipyridyl,
'chel'). Values are mean from n = 3 independent experiments ±SEM. * denotes $p \leq 0.05$, Student's t-
test, when comparing values obtained in the absence of chelator for the mutant cell line versus the
complemented one.

**Figure 2. Generation of conditional knock-down and complemented cell lines for the apicoplast-
localized TgSUFC.** A) Immunoblot analysis with anti-HA antibody shows precursor (p) and mature (m)
forms of C-terminally HA-tagged TgSUFC and efficient down-regulation of the protein after 24 hours
of incubation with ATc. Anti-SAG1 antibody was used as a loading control. B) HA-tagged TgSUFC
(green) localizes to the apicoplast (labelled with marker TgCPN60, red) and is efficiently down-
regulated upon addition of ATc for 24 hours. Scale bar represents 5 µm. DNA was labelled with DAPI.

Parasite shape is outlined. C) Immunoblot analysis of the conditional TgSUFC knock-down cell line
expressing a TY-tagged version of the protein shows similar processing profile and stable expression
after 48 hours of ATc addition. D) Immunofluorescence assay confirms co-localization of the
regulatable HA-tagged TgSUFC (green) and the TY-tagged additional copy (red), whose expression is
retained after 24 hours of incubation with ATc. Scale bar represents 5 µm. DNA was labelled with
DAPI. Parasite shape is outlined.

Figure 3. Depletion of TgSUFC affects in vitro growth of the tachyzoites irresversibly. A) Plaque
assays were carried out by infecting HFF monolayers with the TgSUFC2-HA conditional knock-down
and complemented cell lines. They were grown for 7 days ± ATc. B) Measurements of lysis plaque
areas highlight a significant defect in the lytic cycle when TgSUFC is depleted. Values are means of n =
3 experiments ± SEM. Mean value of a TATi ΔKu80 control grown in the presence of Atc (not shown
on the left) was set to 100% as a reference. **** denotes $p \leq 0.0001$, Student's t-test. Scale bar =
2mm. C) Plaque assays for the TgSUFC mutant was performed as described in A), but ATc was
washed out after 7 days (7d+ATc 7d-ATc) or not (14d+ATc), and parasites were left to grow for an
extra 7 days. No plaque was observed upon Atc removal. Shown are images from one representative
out of three independent experiments. Scale bar = 1mm. D) TgSUFC mutant and complemented cell
lines, as well as their parental cell line and the TATi ΔKu80 control, were grown in HFF in the
presence or absence of ATc for 48 hours, and subsequently allowed to invade and grow in new HFF
cells for an extra 24 hours in the presence of ATc. Parasites per vacuole were then counted. Values
are means ± SEM from n = 3 independent experiments for which 200 vacuoles were counted for each
condition.

Figure 4. Depletion of TgNFS2 or TgSUFC leads to membrane defects during cell division. A) TgNFS2-
HA and TgSUFC-HA conditional knock-down parasites as well as a TATi ΔKu80 control were grown in
the presence of ATc for up to 2 days and were stained with an anti-TgIMC3 antibody (in red, to
outline parasites and internal buds - top). Scale bar represents 5 µm. DNA was labelled with DAPI
(blue). B) The percentage of vacuoles presenting asynchronous division described in B) has been
quantified and is represented as a means of n = 3 experiments ± SEM (bottom). ** denotes $p \leq 0.01$,
Student's t-test. C) Electron microscopy analysis of TgNFS2-HA and TgSUFC-HA conditional mutants
grown for 4 days in the presence of ATc shows default in plasma membrane separation during
parasite division, as displayed on insets representing magnifications of selected parts of the
respective left image. D) cKD TgSUFC-HA parasites that were grown in the presence of ATc for 5 days
were co-stained with anti-TgIMC3 to outline the inner membrane complex and anti-TgCPN60 (an
apicoplast marker), which highlighted abnormal membrane structures and organelle segregation
problems. Scale bar represents 5 µm. DNA was labelled with DAPI.

Figure 5. TgSUFC depletion impacts apicoplast-hosted Fe-S pathways. A) cKD TgSUFC-HA parasites
were kept in the presence of ATc for up to five days and the aspect of the apicoplast was evaluated
by microscopic observation using the specific CPN60 marker. After 3 days, parasites egressed and
were used to reinvade new host cells for subsequent timepoints. Scale bar represents 5 µm. DNA was
labelled with DAPI. DIC: differential interference contrast. B) Using the labelling described in A),
apicoplast loss in vacuoles was monitored after two to five days of incubation with ATc. Data are
mean values from n = 3 independent experiments ±SEM. ns, not significant; * $p \leq 0.05$, **** $p \leq$
0.0001, Student's t-test. C) A decrease in the lipoylation of the E2 subunit of pyruvate dehydrogenase

(TgPDH-E2), which depends on the apicoplast-hosted Fe-S-containing lipoyl synthase LipA, was
observed by immunoblot using an anti-lipoic acid antibody on cell extracts from cKD TgSUFC-HA
parasites kept with ATc for an increasing period of time. A polyclonal antibody raised against PDH-E2
was used as a control for global abundance of the protein and for apicoplast integrity. TgACT1 was
used as a loading control. D) Decrease of lipoylated TgPDH-E2 was quantified by band densitometry
and normalized with the internal loading control. Data represented are mean ±SEM of n = 3
independent experiments. ns, not significant, *≤ 0.05, ** p ≤ 0.01, *** p ≤ 0.001, **** p ≤ 0.0001,
ANOVA comparison.

Figure 6. Lipidomic analysis of the consequences of TgNFS2 and TgSUFC depletion. Ratio of total
lipid content for conditional knock-down cell lines for TgNFS2 (A) and TgSUFC (B) in the presence or
in the absence of ATc. (C) Stable isotope labelling combined to GCMS analyses allows the
quantification of host FA flux towards the parasite. [13]C incorporation levels of C16:0 illustrates a
significant increase of host scavenged fatty acids for the TgSUFC mutant. Data are mean from n=4
independent experiments ±SEM, * p ≤ 0.05 Student's t-test

Figure 7. Exogenous supplementation with short chain fatty acids only partially restore fitness of
cKD TgNFS2 and cKD TgSUFC mutant parasites *in vitro*. Plaque assays were performed as described
in Figure 3A, in the absence or presence of 50 μM of palmitic (C16:0; A, B) or myristic (C14:0; C, D)
acid. Plaque sizes were measured and area was expressed as a percentage of the value obtained
after 7 days of growth in the absence of ATc. are mean values from n = 3 independent experiments
±SEM. * p ≤ 0.05, ** p ≤ 0.01, Student's t-test. Palmitic acid allow partial restoration of plaques with
the cKD TgNFS2 mutant and to a lesser extent with the cKD TgSUFC mutant after 7 days of growth;
myristic acid partially restored plaques but only for the cKD TgNFS2 mutant and after two weeks of
incubation.

Figure 8. The depletion of TgNFS2 or TgSUFC leads to mislocalization of GPI-anchored surface
antigens and impacts gliding motility. A) TgNFS2 and TgSUFC conditional mutants were grown for
three days in the presence or absence of ATc and allowed to invade host cells for another 24 hours in
the presence or absence of ATc. Parasites were then co-stained for inner membrane complex marker
IMC3 (red) together with GPI-anchored protein SAG1 or SAG3 (green). As shown on insets
representing selected parts of the images, the depletion of TgNFS2 or TgSUFC leads to the
accumulation of SAGs in the vacuolar space (white arrowhead) or concentration at the apex or base
of the parasite (red arrowhead). Scale bar represents 10 μm. DNA was labelled with DAPI. B)
Quantification of the abnormal distribution of SAG labellings in vacuoles containing more than two
parasites. Data are mean values from n = 3 independent experiments ±SEM. **** p ≤ 0.0001,
Student's t-test. C) Representative views of a gliding assay showing lower abundance of SAG1 trails
upon TgNFS2 or TgSUFC depletion (inverted grayscale images). Scale bar represents 10 μm. D)
Quantification of the trail to parasite ratio on at least ten randomly selected fields. Data are mean
values from n = 3 independent experiments ±SEM. ns, not significant; * p ≤ 0.05, ** p ≤ 0.01,
Student's t-test. E) Individual measurements of SAG1 trail lengths. Horizontal lines represent mean
values from n = 3 independent experiments ±SEM. ns, not significant; **** p ≤ 0.0001, Student's t-
test.

Figure 9. Schematic representation of th main celllular pathways that depend on apicoplast Fe-S
proteins. For selected apicoplast-located Fe-S proteins squared red numbers represent CRISPR
fitness score of the corresponding gene (genes that contribute to *in vitro* parasite fitness are
represented by negative scores; values below −2.5 highlight increasing likelihood of being essential).
Fd: ferredoxin; IspG/IspH: oxidoreductases catalysing the last two steps of IPP/DMAPP synthesis;
LipA: lipoyl synthase; PDH-E2: E2 subunit of the pyruvate dehydrogenase complex; CoA: coenzyme A;
IPP: isopentenyl diphosphate; DMAPP: dimethylallyl diphosphate; FASII: fatty acid synthesis type II;
GPP: Geranyl diphosphate; FPP: farnesyl diphosphate; GGPP: Geranylgeranyl diphosphate; GPI:
Glycosylphosphatidylinisotol; dol-P: dolichol phosphate.

Supporting Information

This article contains supporting information (Figures S1-S5, Table S1)

A

SUF pathway

B

```
                    1              10              20              30              40              50              60
TgSUFCTGGT1_225800  M A S L P V F L V E L D R W K H H V W L R R L R C L L V S P F V S L S V L P V L L P F L F N F V T S P D I V P S S A L T
EcSufC EGT68402

                    70              80              90             100             110             120
TgSUFCTGGT1_225800  L K S T S A F L S S P A S R F S P P L S P L S C V S S R P S L S P S S S A S S A P S S P S S A S L G V R P H F I S V P A
EcSufC EGT68402

                   130             140             150             160             170             180
TgSUFCTGGT1_225800  A S G T E L A A S L S S T T S S F I G S S S S T R P A L N R D S A C R Q T V S F P A R L S P T L F P L S P D T E D G R G
EcSufC EGT68402

                   190             200             210             220             230             240
TgSUFCTGGT1_225800  S F S D L Q Q L R D S R E E R F A A L K S M R E Q E T P R F Q A E D W T P D Q P L L E I K D L R V E A E E D G Q E F L K
EcSufC EGT68402                                                               M F S I K D L H V S V E D - - K A I L R

                   250             260             270             280             290             300
                                            Walker A
TgSUFCTGGT1_225800  G V N A T I M P G E V H A T V G R N G S G K S T L S K V L A G Y P S Y K V T A G E V R Y K G L D L T E L P I D N R G L A
EcSufC EGT68402      G L S L A V R P G E V H A I M G P N G S G K S T L S A T L A G R E D Y E V T G G I V E F K G K D L L A L S P E D R A G E

                   310             320             330             340             350             360
                              Q loop
TgSUFCTGGT1_225800  G L P L A F Q Y P I E L P L V S N L E P L R V A F N E R R K W K E P E V V S Y E E R E L V E G R L K E V G L D P S F L
EcSufC EGT68402      G I F M A F Q Y P V E L P G V S N Q F F L Q T A L N A V R S Y R G Q E T L D R F D F Q D L M E E K I A L L K M P E D L L

                   370             380             390             400             410             420
                           ABC signature              Walker B   D-loop
TgSUFCTGGT1_225800  D R P L N Y G F S G G E K K R N E L L Q M L V L D P E L V M L D E I D S G L D V D S F N I T A N A I K R F S K R K G K S
EcSufC EGT68402      T R S V N V G F S G G E K K R N D I L Q M A V L E P E L C I L D E S D S G L D I D A L K V V A D G V N S L R D G K - R S

                   430             440             450             460             470             480
                           H motif
TgSUFCTGGT1_225800  P L V T T H Y K K L L D V L Q P H K I H V M H A G K I V L S G S M D L A G Q I E A E G F Q A L V G A A A E G E E E E E R
EcSufC EGT68402      F I I V T H Y Q R I L D Y I K E D Y V H V L Y Q G R L V K S G D F T T L V K Q L E Q Q G Y G W Y T - - - - - - D Q Q

                   490             500             510    518
TgSUFCTGGT1_225800  D R R E D E G E R E E H E G E D E A E S L L S P S A A G G G R R N L D R L L
EcSufC EGT68402
```

C

A

cKD TgSUFC-HA | cKD TgSUFC-HA comp

no ATc

+ ATc

2 mm

B

Plaque area (% of control)

ATc: + − + − +

TATi ΔKu80 | cKD TgSUFC-HA | cKD TgSUFC-HA comp

**** ****

C

cKD TgSUFC-HA

1 mm

7d +ATc
7d -ATc | 14d +ATc

D

number of vacuoles

ATc: + + − + − +

TATi ΔKu80 | TgSUFC-HA | cKD TgSUFC-HA | cKD TgSUFC-HA comp

parasites per vacuole

A

	1d +ATc	2d +ATc
TATi ΔKu80		
cKD TgNFS2-HA		
cKD TgSUFC-HA		

B

Vacuoles with asynchronous division

- ☐ 1d+ATc
- ☐ 2d+ATc

TATi ΔKu80 — cKD TgNFS2-HA — cKD TgSUFC-HA

**

C

cKD TgNFS2-HA

5 μm — 500 nm

cKD TgSUFC-HA

5 μm — 500 nm

D

IMC

DAPI

Apicoplast

Merge