Synapses in oligodendrocyte precursor cells are dynamic and contribute to Ca^{2+}

activity

Jiaxing Li[1, *], Tania Miramontes[1], Tim Czopka[2], Kelly Monk[1, *]

1 Vollum Institute, Oregon Health & Science University, Portland, OR, USA

2 Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK

* Corresponding authors

Cell-cell interactions are critical to central nervous system (CNS) structure and function. Direct

neuron-neuron interactions occur via synapses, but only one type of neuron-glia synapse is

known to exist: those formed between neurons (pre-) and oligodendrocyte precursor cells

(OPCs) (post-)[1,2]. These synapses are thought to bridge neuronal activity to OPCs, which are

critical for developmental myelination and adult motor learning[3]. However, the composition

and assembly of neuron-OPC synapses and their *in vivo* functions remain largely unclear.

Here, using zebrafish to label postsynaptic organizers PSD-95 and Gephyrin in OPCs, we found

that synapses in OPCs possess dynamic assembly properties and are developmentally

regulated. We also found that synapses and voltage-gated Ca^{2+} channels (VGCCs) play

important roles in driving intrinsic OPC Ca^{2+} activity that occurs mainly in microdomains,

while evoked synaptic release has a limited role that diminishes through development. Finally,

disrupting a key synaptic gene, *neuroligin3b,* specifically in OPCs impairs OPC differentiation

and subsequent myelination. Our results establish an *in vivo* model of dynamic neuron-OPC

synapses mediating intrinsic OPC Ca^{2+} activity and impacting OPC differentiation, pointing to

plastic neuron-OPC interactions and synapse-dependent and

-independent Ca^{2+} pathways in OPCs.

Neuron-OPC synapse assembly and disassembly is mysterious. This is largely due to lack of

labeling methods, especially *in vivo*. The composition of these synapses is also largely unknown,

with the exception of some neurotransmitter and synaptic receptors (*e.g.*, glutamatergic and

GABAergic) that have been revealed by pharmacology and electrophysiology[3]. OPCs exhibit

heterogeneity[4,5], raising the question of whether synapse assembly differs among OPCs. Unlike

canonical postsynaptic cells (*e.g.*, neurons and muscle cells), OPCs migrate and constantly

remodel their processes, and many OPCs differentiate and transform these processes into

myelin sheaths[3]. Therefore, OPCs may employ special mechanisms of synapse assembly to

accommodate their unique biology, a notion further supported by transcriptomic studies

suggesting different synaptic gene expression in OPCs from neurons[6,7]. These transcriptomic

studies, however, do indicate the presence of PSD-95 (a major MAGUK family member) and

Gephyrin in OPCs (Extended Data Fig. 1)[6], which are abundant postsynaptic scaffold proteins at

excitatory and inhibitory neuron-neuron synapses, respectively[8], making them promising

candidates for neuron-OPC synapse labeling.

Previous *ex vivo* rodent studies linked axon stimulation and neuron-OPC synapses to

Ca^{2+} rise in oligodendrocyte lineage cell somas and proximal processes[9–11]. While these studies

demonstrate the capacity of neuron-OPC synapses to induce OPC Ca^{2+} activity, it remains

unknown if these synapses contribute to generating intrinsic OPC Ca^{2+} activity *in vivo*.

Furthermore, recent *in vivo* zebrafish studies revealed intrinsic Ca^{2+} events in microdomains of

OPC processes[4,12], but the role of synapses in generating these events is largely unknown. Here

we leveraged zebrafish to study synapse assembly mechanisms in OPCs *in vivo* and to address

the role of synapses in OPC biology and intrinsic Ca^{2+} activity.

Postsynaptic structures (PSSs) in OPCs are dynamic, assemble at preferential locations, and

many lack presynaptic structure alignment.

We generated PSD-95-GFP and GFP-Gephyrin constructs and drove their expression

with a pan-oligodendrocyte lineage promoter (*sox10*, see Methods). At 3 days post fertilization

(dpf) in the spinal cord, we found both markers present in punctate and cytosolic forms in OPCs

(Fig. 1a, see methods for OPC identification), consistent with patterns observed in neurons[13,14],

with OPC puncta size resembling that of neurons (Fig. 1a-b and Extended Data Fig. 2a). We

detected 29 ± 4.4 (N=18) PSD-95-GFP and 27 ± 3.5 (N=14) GFP-Gephyrin puncta per OPC.

Through immunohistochemistry (IHC), we verified the presence of MAGUK (and PSD-95) and

Gephyrin puncta in spinal cord OPCs (Extended Data Fig. 3). We found similar Gephyrin puncta

number between *in vivo* imaging and IHC, and higher PSD-95 puncta number by *in vivo* imaging

than by IHC (Extended Data Fig. 3h), possibly due to increased synaptogenesis with PSD-95

overexpression[15]. Despite the overexpression caveat, the IHC results verified our *in vivo*

methods for labeling PSS puncta in OPCs. Through time-lapse imaging, we found that the

number of PSS puncta per cell remains stable in a neuron, but fluctuates in an OPC over 30

minutes of observation (Fig. 1c and Supplementary Videos 1-4). The puncta of PSD-95-GFP and

GFP-Gephyrin in OPCs are considerably shorter-lived than in neurons (Fig. 1d). Many puncta in

neurons last more than 10,000 seconds; conversely, we found no such stable puncta in OPCs

(Fig. 1e). While some PSS puncta in OPCs appear or disappear as processes remodel (Extended

Data Fig. 2b and Supplemental Video 3), others do when processes are relatively stable

(Extended Data Fig. 2c-d and Supplemental Video 1). Some PSS puncta in OPCs appear

preferentially at certain locations, verified by frequency maps (Fig. 1f-g). Further analysis

revealed that PSS puncta in these "hotspots" repeatedly assemble and disassemble (10-12

"hotspots" per cell, Fig. 1h and Extended Data Fig. 2c-d). Notably, we observed this

phenomenon in not only dynamic processes but also stable processes (Extended Data Fig. 2c-d),

suggesting that the repeatedly assembled and disassembled "hotspots" are not simply caused

by process instability.

Next, we tested whether these dynamic PSS puncta form complete synapses by

examining their alignment with presynaptic structures (pan-presynaptic marker Synapsin 1/2[16]).

We found nearly a 50% alignment rate for MAGUK and Gephyrin in OPCs and a reduced

colocalization ratio between pre- and post-synaptic staining compared to neurons (Extended

Data Fig. 4). This suggests that nearly 50% of PSSs found in OPCs are part of a functional

synapse unit while the rest lack synaptic partners. Similar non-aligned PSSs have been reported

in developing neurons and are considered to be immature[17,18]. From these data, we conclude

that PSSs in OPCs are more dynamic than neurons, assemble and disassemble frequently, and

that many are likely immature.

PSSs number in OPCs are developmentally regulated, with few regional and subgroup

differences.

Given previously reported OPC heterogeneity[4,5], we wondered if PSSs in OPCs are

regulated spatially, temporally, and/or between subgroups. Using IHC, we found that MAGUK

puncta numbers are similar across CNS regions, while Gephyrin puncta number is significantly

higher in the forebrain and midbrain compared to the spinal cord (Extended Data Fig. 5a),

suggesting a preference of OPCs to form synapses with inhibitory neurons in the brain. Two

subgroups of OPCs have been reported in zebrafish spinal cord that exhibit distinct

transcriptional profiles and myelination potential[4]. Despite these differences, both subgroups

possess similar PSS numbers (Extended Data Fig. 5b-c).

114   We found that both MAGUK (and PSD-95) and Gephyrin puncta numbers in OPCs

increase as animals develop using *in vivo* imaging and IHC (Fig. 2a-b). Time-lapse imaging of

OPCs revealed that coincident with differentiation, PSS number is reduced (Fig. 2c-d), which

could account for reduced synaptic inputs previously reported at this stage[19,20]. In cases where

we observed differentiating oligodendrocytes that possess both processes and myelin sheaths,

puncta were almost exclusively in processes (Fig. 2e), suggesting preferential localization of

PSSs to processes rather than nascent myelin sheaths. These results show that PSS number in

OPCs increases as animals develop and decreases upon OPC differentiation.

Synapses and VGCCs contribute to generating intrinsic OPC Ca^{2+} activity in vivo.

To investigate the role of synapses in intrinsic OPC Ca^{2+} activity *in vivo*, we employed a plasma-

membrane-targeting GCaMP6s (See Methods). At single cell resolution, we found that the

majority (99%) of Ca^{2+} events in OPCs occur in microdomains on processes (Fig. 3a): we

recorded 624 local events (Supplementary Video 5) and 6 somal events (Supplementary Video 6)

from 180 cells, 107 fish, 30 hours imaging period. To test if intrinsic OPC Ca^{2+} activity is

mediated by synapses, we first examined the proximity between Ca^{2+} events and synapses,

visualized by a presynaptic marker, synaptophysin-TagRFP[4]. Through 3D analysis, we found that

34 ± 9% of Ca^{2+} microdomains are in close proximity (<0.1 μm) to synaptophysin-TagRFP puncta

at 5 dpf (Fig. 3b-e). In some cases, we observed Ca^{2+} events initiate from a "puncta"-like region

in close proximity to synapses (Extended Data Fig. 6a-c). Next, we blocked synaptic release by

expressing tetanus toxin light chain (TeTxLc)-tdtomato[21] in axons (Fig. 3f). Compared to OPCs

neighboring few TeTxLc-expressing axons (< 4), OPCs neighboring many TeTxLc-expressing

axons (> 4) showed significantly reduced frequency of Ca^{2+} activity, peak amplitude, basal signal

intensity, and microdomain length (Fig. 3g and Extended Data Fig. 6d). Together, these results

establish an *in vivo* role for synapses in generating intrinsic OPC Ca^{2+} activity that mainly occurs

in microdomains.

We next asked what receptors or channels at synapses mediate intrinsic OPC Ca^{2+}

activity. We tested NMDA receptors, CP-AMPA receptors, and VGCCs, as suggested by previous

ex vivo studies[9–11], by administering their antagonists (AP5, IEM-1460, and cadmium,

respectively) via ventricle injection as described previously[12] (Fig. 3h). We confirmed successful

drug delivery: injections induced paralysis by 3 minutes post-injection (AP5, 15/19; IEM-1460,

14/14; cadmium, 9/10), and dextran co-injected with antagonist (dextran MW3000, larger than

the antagonists) diffused and encompassed OPCs prior to imaging in all cases (45/45) (Extended

Data Fig. 6e). We found that AP5 and IEM-1460 did not inhibit the overall Ca^{2+} activity (Fig. 3i

and Extended Data Fig. 6f). In contrast, cadmium administration significantly reduced

microdomain number per OPC (4.4 ± 1.0 to 2.0 ± 0.5) as well as the frequency of Ca^{2+} activity

per microdomain (1.8 ± 0.2 to 0.9 ± 0.2 x 10^{-3} Hz) (Fig. 3i, Extended Data Fig. 6f, Supplementary

Video 7-8). The overall 50-55% reduction in both analyses suggests that VGCCs, but not NMDA

or CP-AMPA receptors, play a critical role in generating intrinsic OPC Ca^{2+} activity *in vivo*.

Neuronal activity and evoked synaptic release play a limited role in intrinsic OPC Ca^{2+} activity

during early development.

Expanding from a previous study[12], we examined the role of neuronal activity and evoked

synaptic release in driving intrinsic OPC Ca^{2+} activity at two developmental stages (3 dpf, when

OPCs initiate differentiation and 5 dpf, when many OPCs complete differentiation[22]). Similar to

previous results with 3 dpf larvae[12], upon ventricle injection (Fig. 3h) of tetrodotoxin (TTX,

sodium channel blocker), OPCs exhibited reductions in the frequency of Ca^{2+} activity (Extended

Data Fig. 7a and Supplementary Video 9). In addition, we observed reduced microdomain

number per OPC (3.0 ± 0.4 *vs.* 2.2 ± 0.3) and peak amplitude (2.1 ± 0.2 *vs.* 1.6 ± 0.1 fold)

(Extended Data Fig. 7a). The overall 25-33% reduction in these analyses suggests a partial

requirement of neuronal activity and evoked synaptic release for intrinsic OPC Ca^{2+} activity.

However, we found no such requirement at 5 dpf (Extended Data Fig. 7a). We excluded the

possibility of ineffective TTX delivery in three ways. We observed larval paralysis within 3

minutes post-injection (30/30), a complete block of motor neuron Ca^{2+} activity (Extended Data

Fig. 7b-c), and dextran co-injected with TTX (dextran MW3000, larger than TTX) surrounded the

imaged OPCs (120/120) (Extended Data Fig. 7d). Therefore, our results point to a limited role of

neuronal activity and evoked synaptic release in driving intrinsic OPC Ca^{2+} activity, which

diminishes over development. These results also suggest that the role of synapses we observed

(Fig. 3) were likely mediated by spontaneous synaptic release.

Synaptic component nlgn3b is important for oligodendrocyte differentiation and myelination.

The role of neuron-OPC synapses in development is unclear due to diverse and

sometimes conflicting results of individual receptor disruption[23–26]. To overcome this, we

targeted *neuroligin3* (*nlgn3*), which localizes to both excitatory and inhibitory synapses,

regulates neuronal synapse structure and function[27], and is enriched in OPCs (Extended Data Fig.

1)[6]. We developed and verified a cell-specific CRISPR/Cas9-mediated gene disruption system

(Extended Data Fig. 8) and targeted the duplicated *nlgn3* genes (*nlgn3a* and *nlgn3b*) in zebrafish.

We observed an increased OPC to myelinating oligodendrocyte ratio in *nlgn3b*-targeted cells at

3 dpf (ctrl 3:61, *nlgn3a* 0:19, *nlgn3b* 9:25) and at 5 dpf (ctrl 1:66, *nlgn3a* 2:56, *nlgn3b* 18:47,

Extended Data Fig. 8e), suggesting that loss of *nlgn3* disrupts proper OPC differentiation. At 5

dpf, almost all myelinating oligodendrocytes in control animals possess only myelin sheaths. In

contrast, several *nlgn3b*-disrupted oligodendrocytes possess both myelin sheaths and

cytoplasmic processes with abnormal trapezoidal structures (ctrl: 0 out of 67, *nlgn3a*: 0 out of

58, *nlgn3b*: 3 out of 65, Fig. 4a and extended Data Fig. 8f), indicating developmental defects

when Neuroligin3b is lost. Additionally, disruption of *nlgn3b* leads to reduced total length of

myelin sheaths per oligodendrocyte at 3 dpf (Fig. 4b), which was largely resolved by 5 dpf (Fig.

4c), suggesting a temporary delay in myelin sheath formation. Together, these results suggest a

role for Neuroligin3 specifically and neuron-OPC synapses more broadly in oligodendrocyte

differentiation and myelination.

Discussion

The landmark discovery of neuron-OPC synapses has inspired several studies to examine

their properties with electrophysiology and EM[3], yet we still lack basic understanding of their

assembly and function. Here, we identified PSSs composed of MAGUK (PSD-95) and Gephyrin in

zebrafish OPCs and revealed unique synapse features: dynamic assembly, lack of consistent

synaptic partners, and frequent assembly at specific locations. Interestingly, these features

have also been reported in some neurons, although to a lesser extent[17,18,28,29]. These synapse

features may enable flexible neuron-OPC synapse formation in tune with neuronal inputs.

Development influences synapse formation in OPCs: PSS number doubles within a few days

during animal development and decreases upon OPC differentiation and myelin sheath

formation. Meanwhile, OPC development seems to require proper synapse formation, revealed

by oligodendrocyte-specific disruption of *nlgn3b*. Together, these findings pinpoint a

bidirectional and mutually influencing relationship between dynamic synapse assembly and

OPC development (Fig. 4d) and a newly discovered developmental role of *nlgn3b*, previously

shown to promote glioma growth[30].

Several previous studies investigated synaptic mechanisms underlying Ca^{2+} elevation in

OPCs[9–11], yet the role of synapses in driving intrinsic OPC Ca^{2+} activity *in vivo* and in

microdomains was never explored. Our work suggests the requirement of synapses in

generating intrinsic OPC Ca^{2+} activity that mainly occur in microdomains. We further showed

that Ca^{2+} influx is likely mediated by VGCCs, but not CP-AMPA or NMDA receptors. This is in

contrast with a previous *ex vivo* study[9] and could be explained by the different mechanisms

underlying somal *vs.* microdomain Ca^{2+} events. To our surprise, neuronal activity and evoked

synaptic release contribute moderately to intrinsic OPC Ca^{2+} activity at 3 dpf but not 5 dpf.

Considering that OPC differentiation initiates at 3 dpf, it suggests an OPC sensitivity to neuronal

activity through Ca^{2+} signaling at this critical developmental stage. Overall, our results establish

an *in vivo* model of intrinsic OPC Ca^{2+} activity: synapses, likely through spontaneous release,

play an important role with evoked release and neuronal activity contributing during early

development.

References

1. Bergles, D. E., Roberts, J. D. B., Somogyl, P. & Jahr, C. E. Glutamatergic synapses on

oligodendrocyte precursor cells in the hippocampus. *Nature* **405**, 187–191 (2000).

2. Lin, S. C. & Bergles, D. E. Synaptic signaling between GABAergic interneurons and

oligodendrocyte precursor cells in the hippocampus. *Nat. Neurosci.* **7**, 24–32 (2004).

3. Bergles, D. E. & Richardson, W. D. Oligodendrocyte development and plasticity. *Cold*

Spring Harb. Perspect. Biol. **8**, a020453 (2016).

4. Marisca, R. *et al.* Functionally distinct subgroups of oligodendrocyte precursor cells

integrate neural activity and execute myelin formation. *Nat. Neurosci.* **23**, 363–374

(2020).

5. Spitzer, S. O. *et al.* Oligodendrocyte Progenitor Cells Become Regionally Diverse and

Heterogeneous with Age. *Neuron* **101**, 459-471.e5 (2019).

6. Zhang, Y. *et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons,

and vascular cells of the cerebral cortex. *J. Neurosci.* **34**, 11929–11947 (2014).

7. Marques, S. *et al.* Oligodendrocyte heterogeneity in the mouse juvenile and adult central

nervous system. *Science (80-.).* **352**, 1326–1329 (2016).

8. Sheng, M. & Kim, E. The postsynaptic organization of synapses. *Cold Spring Harb.*

Perspect. Biol. **3**, a005678–a005678 (2011).

9.   Barron, T. & Kim, J. H. Neuronal input triggers Ca2+ influx through AMPA receptors and

voltage-gated Ca2+ channels in oligodendrocytes. *Glia* **67**, 1922–1932 (2019).

10.   Sun, W., Matthews, E. A., Nicolas, V., Schoch, S. & Dietrich, D. NG2 glial cells integrate

synaptic input in global and dendritic calcium signals. *Elife* **5**, (2016).

11.   Hamilton, N., Vayro, S., Wigley, R. & Butt, A. M. Axons and astrocytes release ATP and

glutamate to evoke calcium signals in NG2-glia. *Glia* **58**, 66–79 (2010).

12.   Krasnow, A. M., Ford, M. C., Valdivia, L. E., Wilson, S. W. & Attwell, D. Regulation of

developing myelin sheath elongation by oligodendrocyte calcium transients in vivo. *Nat.*

Neurosci. **21**, 24–30 (2018).

13.   Yoshii, A. & Constantine-Paton, M. BDNF induces transport of PSD-95 to dendrites

through PI3K-AKT signaling after NMDA receptor activation. *Nat. Neurosci.* **10**, 702–711

(2007).

14.   Meier, J. & Grantyn, R. A Gephyrin-Related Mechanism Restraining Glycine Receptor

Anchoring at GABAergic Synapses. *J. Neurosci.* **24**, 1398–1405 (2004).

15.   Nikonenko, I. *et al.* PSD-95 promotes synaptogenesis and multiinnervated spine

formation through nitric oxide signaling. *J. Cell Biol.* **183**, 1115–1127 (2008).

16.   Easley-Neal, C., Fierro, J., Buchanan, J. A. & Washbourne, P. Late Recruitment of Synapsin

to Nascent Synapses Is Regulated by Cdk5. *Cell Rep.* **3**, 1199–1212 (2013).

17.   Gerrow, K. *et al.* A preformed complex of postsynaptic proteins is involved in excitatory

synapse development. *Neuron* **49**, 547–562 (2006).

18.   Panzer, J. A. *et al.* Neuromuscular synaptogenesis in wild-type and mutant zebrafish. *Dev.*

Biol. **285**, 340–357 (2005).

19.   Kukley, M., Nishiyama, A. & Dietrich, D. The fate of synaptic input to NG2 glial cells:

Neurons specifically downregulate transmitter release onto differentiating

oligodendroglial cells. *J. Neurosci.* **30**, 8320–8331 (2010).

20.   De Biase, L. M., Nishiyama, A. & Bergles, D. E. Excitability and synaptic communication

within the oligodendrocyte lineage. *J. Neurosci.* **30**, 3600–3611 (2010).

21.   Koudelka, S. *et al.* Individual Neuronal Subtypes Exhibit Diversity in CNS Myelination

Mediated by Synaptic Vesicle Release. *Curr. Biol.* **26**, 1447–1455 (2016).

22.   Almeida, R. & Lyons, D. Oligodendrocyte development in the absence of their target

axons in Vivo. *PLoS One* **11**, (2016).

23.   Balia, M., Benamer, N. & Angulo, M. C. A specific GABAergic synapse onto

oligodendrocyte precursors does not regulate cortical oligodendrogenesis. *Glia* **65**, 1821–

1832 (2017).

24.   de Biase, L. M. *et al.* NMDA receptor signaling in oligodendrocyte progenitors is not

required for oligodendrogenesis and myelination. *J. Neurosci.* **31**, 12650–12662 (2011).

25.   Zonouzi, M. *et al.* GABAergic regulation of cerebellar NG2 cell development is altered in

perinatal white matter injury. *Nat. Neurosci.* **18**, 674–682 (2015).

26.   Lundgaard, I. *et al.* Neuregulin and BDNF Induce a Switch to NMDA Receptor-Dependent

Myelination by Oligodendrocytes. *PLoS Biol.* **11**, (2013).

27.   Krueger-Burg, D., Papadopoulos, T. & Brose, N. Organizers of inhibitory synapses come of

age. *Curr. Opin. Neurobiol.* **45**, 66–77 (2017).

28.   Wierenga, C. J., Becker, N. & Bonhoeffer, T. GABAergic synapses are formed without the

involvement of dendritic protrusions. *Nat. Neurosci.* **11**, 1044–1052 (2008).

29.   Villa, K. L. *et al.* Inhibitory Synapses Are Repeatedly Assembled and Removed at

Persistent Sites In Vivo. *Neuron* **89**, 756–769 (2016).

30.   Venkatesh, H. S. *et al.* Neuronal activity promotes glioma growth through neuroligin-3

secretion. *Cell* **161**, 803–816 (2015).

Methods

Zebrafish lines and husbandry

We used the following existing zebrafish lines and strains: *Tg(sox10:Kalta4)*[31], *Tg(olig1:Kalta4)*[4],

Tg(huc:synaptophysin-tagRFP)[4], *Tg(olig2:dsRed)*, *Tg(sox10:GFP-caax)*, *nacre*, and AB. The

following lines were newly generated for this study: *Tg(10xUAS:myrGCaMP6s)*,

Tg(olig1:myrGCaMP6s), *Tg(10xUAS:myrGFP)*, *Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:ctrl-*

sgRNA1;U6:ctrl-sgRNA2), *Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:nlgn3a^{ex1}-*

sgRNA;U6:nlgn3a^{ex4}-sgRNA), *Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:nlgn3b^{ex1}-*

sgRNA;U6:nlgn3b^{ex3}-sgRNA). All experimental fish were crossed into the *nacre* background for

optimal optics without pigmentation. The Kalta4/UAS system allows sparse labeling and single

cell analysis. The genotypes of fish used for each experiment were indicated in figure legends or

Supplementary Table 3. All animals were kept at 28° C with a 14 h/10 h light/dark cycle.

Larvae and juveniles were nurtured with rotifer suspension and dry food (Gemma 75 and 150,

respectively). Adults were fed with a combination of brine shrimp and dry food (Gemma 300).

All zebrafish experiments and procedures were performed in compliance with institutional

ethical regulations for animal testing and research at Oregon Health and Science University

(OHSU). Experiments were approved by the Institutional Animal Care and Use Committee of

OHSU. Animal and cell number were indicated in figure legends or Supplementary Table 2.

Tissue sectioning and immunohistochemistry

To prepare sagittal and transverse sections, zebrafish larvae at indicated stages were first

anesthetized with tricaine and then fixed in 4% PFA, 0.1% Triton at 4° C overnight. Heads and

tails were removed to enable better penetration of fixative. The fish were then washed in PBS,

0.1% Triton for 5 minutes and passed in 10%, 20%, and 30% sucrose in PBS, followed by OCT

embedding and cryosectioning of 20-25 μm slices. The sections were brought to room

temperature for 45 min and rehydrated in PBS, 0.2% Triton for 1 hour. 3% BSA, 5% NGS, 0.2%

Triton in PBS was used as blocking buffer. Primary antibodies: chicken anti-GFP (Invitrogen

A10262), 1:1000, mouse anti-Pan-MAGUK (NeuroMab K28/86) 1:500, mouse anti-Gephyrin

(Synaptic Systems 147011) 1:1000, rabbit anti-Synapsin 1/2 (Synaptic Systems 106002) 1:1000,

mouse anti PSD-95 6G6-1C9 (MAB1596) 1:1000. Secondary antibodies: A488 anti-chicken, A594

anti-rabbit, A647 anti-mouse IgGI, A647 anti-mouse (Thermo Fisher). The stained samples were

mounted in prolong Glass (Invitrogen P36980) for at least a day before imaging.

Plasmid construction

In zebrafish, *gephyrin* is duplicated (*gephyrina* and *gephyrinb*), and there are multiple isoforms

for both genes. All isoforms are almost identical in protein sequence except for the inclusion or

exclusion of C3 and C4 cassettes. We amplified several isoforms from cDNA and decided to

follow up *gephyrinb* isoform X6 (XM_021468341) because it encodes a protein that is 96%

identical and 98% similar to the rat P1 isoform (NP_074056.2)[32] that has been used to track

synapses previously, and, *gephyrina* isoform X11 (XM_017351650.2) because it contains a C3

cassette previously reported to be present in glia cells[33].

Total RNA was extracted from 3 dpf zebrafish AB larvae (50 fish) using RNAeasy Plus Kit

(Qiagen 74134). cDNA was synthesized using SuperScript IV Reverse Transcriptase (Invitrogen).

2 sets of primers (Supplementary Table 1) were used to clone the full length *gephyrina* and

gephyrinb using high fidelity Q5 DNA polymerase, followed by A-addition and TA cloning into

pCRII-TOPO backbones. Using Gibson assembly (NEB E5510S), we added EGFP to the 5' end and

a linker GGGGSGGGGSGGGGS in between, and integrated it into pCS 2.1 backbones with an

AVRII digestion site. The 5' end was chosen based on previous imaging studies[32]. In transient

injections, GFP-Gephyrina and GFP-Gephyrinb exhibit similar expression profiles; therefore, we

used GFP-Gephyrinb for our studies and refer to the GFP-3xGGGGS-Gephyrinb as GFP-

Gephyrinb throughout. Using Gateway BP reaction (Invitrogen 11786100), we generated

pME_ATG-GFP-Gephyrinb-TAA. Using AVRII and KPNI-HF, we cut GFP-Gephyrinb (without an

ATG) from the pCS 2.1 backbone and inserted it in frame into p3E_p2A to generate p3E_p2A-

GFP-Gephyrinb-TGA-polyA.

PSD-95/dlg4b transcript NM_214728.1 encodes a protein that is 76% identical and 83%

similar to rat *Dlg4* (NP_062567.1), and was thus chosen for use in our studies. We added GFP to

the 3'end because N-terminal palmitoylation of PSD-95 is necessary for its synaptic localization,

and receptor clustering[34,35]. It was cloned from cDNA with primers (Supplementary Table 1)

similar to *gephyrin* as described above. We refer to PSD-95-3xGGGGS-GFP as PSD-95-GFP. We

generated pME_ATG-PSD-95-GFP-TAA and p3E_p2A-PSD-95-GFP-TGA-polyA.

pME_myrGFP and pME_myrTagRFP-T were generated using PCR and BP reaction, using

Fyn myristoylation and palmitoylation domain. pGP-CMV-GCaMP6s (Addgene 40753) was used

as a template and 3 primers were used in a Q5 PCR (15 cycles at 71° C and 20 cycles at 72° C) to

amplify GCaMP6s and add Kozak sequence, myr (Fyn myristoylation and palmitoylation

domain), attB1, and attB2 sequences. This PCR product was used to generate

pME_myrGCaMP6s-TAA using BP reaction.

We obtained multiple plasmids from Tol2 kits: pDestTol2pA2, pDONR221, p5E_10xUAS,

p3E_polyA, p3E_p2A[36], p5E_sox10$_{7.2kb}$[31], pDestTol2pACryGFP (pDestTol2GE),

pDestTol2pACrymCherry (pDestTol2RE).

Using Gateway LR reactions (Invitrogen 11791020), we generated the following

constructs: PA2-10xUAS:myrEGFP-PolyA, GE-10xUAS:myrGCaMP6s-TAA-polyA, PA2-

olig1:myrGCaMP6s-TAA-polyA, GE-10xUAS:GFP-Gephyrinb-TAA-polyA, RE-10xUAS:GFP-

Gephyrina-TAA-polyA, GE-10xUAS:PSD-95-GFP-TAA-polyA, PA2-sox10:myrTagRFP-T-P2A-GFP-

Gephyrinb-polyA, PA2-sox10:myrTagRFP-T-P2A-PSD95-GFP-polyA. We used Mnx1:Gal4[37] to

drive myrGCaMP6s expression in motor neurons. We used Huc-Kalta4[31], UAS:TeTxLc-

tdtomato[21] to drive TeTxLc expression in neurons. All constructs generated in this study were

fully sequenced (except *sox10* and *olig1* promoters).

Cell specific CRISPR/Cas9 plasmid construction and CRISPR/Cas9-mediated gene disruption

We used 5xUAS-hsp70promoter-Cas9-T2A-GFP, U6:sgRNA1;U6:sgRNA2[38] (Addgene 74009) as a

backbone and performed the following modifications: replacement of the 5xUAS-hsp70

promoter with the 10xUAS-E1b-bactin-Kozak promoter because we found the original promoter

expressed in many non-specific cell types when injected into *Tg(sox10:Kalta4)*; replacement of

T2A-GFP with myrmScarletI-p2A and added to the 5'end of Cas9; addition of polyA to the 3' end

of Cas9. Four sets of primers and 2 previously made plasmids (Tol2-10xUAS-E1b-mScarletI and

Tol2-10xUAS:cas9-P2A-TagRFPTcaax) were used to generate 4 segments. These segments were

integrated into the backbone by NEBuilder HiFi assembly (NEB E2621S). The new construct is

referred to as 10xUAS:myrmScarletI-p2A-Cas9, U6:sgRNA1;U6:sgRNA2.

CHOPCHOP and Crisprscan were used to identify 2 sgRNA sequences for each gene of

interest: *gephyrina* exon 2, GGTCATGAACGAGATCTTTGAGG; *gephyrinb* exon 7,

GGAGAGCCGGCAGGATGAACTGG; *myrf* exon 3, GACCCAACGGCGGTGCACCCGGG; *myrf* exon 4,

GCTGGAGTCCGGAGGAGACTCGG; *nlgn3a* exon 1, GGTGTATGATGCTTGTCCGGGGG; *nlgn3a* exon

4, GCCTTGTGTACTTAACCGGCTGG; *nlgn3b* exon 1, GATGCGGGTCGCTGTTGCCACGG; *nlgn3b*

exon 3, GGTAGTGTGCTGGCCAGCTACGG. The cutting efficiencies of these sgRNAs were verified

(>95%) via enzyme digestion of the targeted locus PCR product (CRISPRtest primers,

Supplementary Table 1). F0 larvae with *gephyrina* and *gephyrinb* sgRNA injected together with

1 ng Cas9 protein were used to examine the specificity of the Gephyrin antibody. BsaI and

BsmBI were used to insert the sgRNA into 10xUAS:myrmScarletI-p2A-Cas9,

U6:sgRNA1;U6:sgRNA2. The plasmids were injected with Tol2 transposase mRNA to generate

transgenic fish. *Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:myrf^{exon3};U6:myrf^{exon4})*,

Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:nlgn3a^{exon1};U6:nlgn3a^{exon4}), and

Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:nlgn3b^{exon1};U6:nlgn3b^{ex34}) larvae with sparsely labeled

oligodendrocyte were examined for morphology and sheath characteristics.

Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:sgRNA1;U6:sgRNA2) larvae were used as controls.

Sparse labeling and generation of transgenic lines

Fertilized zebrafish eggs at the 1-cell stage were microinjected with 1 nL of an injection

solution containing 20 ng/μL plasmid DNA, 25 ng/μL Tol2 transposase mRNA, 0.02% phenol

red and 0.2M KCl. Injected F0 animals were either used for single-cell analysis or raised to

adulthood to generate transgenic lines. To generate *Tg(10xUAS:myrGFP)* and

Tg(10xUAS:myrGCaMP6s), adult F0 animals were outcrossed with *Tg(sox10:Kalta4)* zebrafish,

and F1 progenies were screened for germline transmission of the fluorescent reporters. F1

animals were further outcrossed and maintained in the *nacre* background to establish stable

transgenic lines with single copy insertion.

Mounting larvae for in vivo imaging

For all live imaging experiments except those utilizing myrGCaMP6s, larvae were anesthetized

with 0.16 mg/mL (600 μM) tricaine in embryo medium. The fish were mounted in 0.8% low

melting agarose (Sigma A9414) on a coverslip for morphology and sheath examination or in 1.5%

low melting agarose in a dish for time lapse imaging (hours). For myrGCaMP6s imaging, larvae

were immobilized with 0.5 mg/ml non-depolarizing acetylcholine receptor antagonist

mivacurium chloride (Abcam) and mounted in 1.5% low melting agarose in a dish.

Confocal microscopy

All imaging was performed on an upright Zeiss LSM 980 confocal with Airyscan 2 in 4Y fast

mode. For fixed tissues, a 63x/1.4 NA oil objective was used, and for live imaging, a 20x/1.0 NA

water objective was used. All images were taken in Zeiss software. We used the following lasers:

488 nm for EGFP, myrGCaMP6s, and Alexa488; 561 nm for TagRFP, mScarletI, and Alexa594;

639 nm for Alexa647. Fixed tissue was imaged with 0.15 μm z spacing with 1.6x to 2.5x zoom.

For *in vivo* analyses, PSD-95-GFP and GFP-Gephyrin were imaged with 2 μm z-spacing at an

interval of 5 min (for 30 min to 1 hour) or 20 min (for longer than 1 hour) at a zoom between

3.7x-10x depending on OPC cell size. myrGCaMP6s was imaged with 2 μm z-spacing at 0.5 Hz (4

planes scanned) for 10 min or 20 min (drug injection experiments) as previously described[4,12].

Images acquired at 0.5 Hz were analyzed along with images acquired at 10 Hz with a single

plane. No major differences were found on the Ca^{2+} spike characteristics, but 0.5 Hz acquisition

allows inclusion of more stacks and increases the number of events detected. Therefore, 0.5 Hz

was used throughout the study.

Drug delivery through ventricle injection

For myrGCaMP6s experiments involving drug delivery, we injected drugs via ventricle injection

as described previously[12]. Briefly, following 10 min GCaMP6s imaging, a 1 nL mixture containing

0.2M KCL, 0.5% Dextran Texas Red 3000 MW (Thermo), and molecular biology grade H_2O

(control), TTX (0.5 mM), IEM-1460 (20 mM), AP5 (30 mM), or cadmium (100 mM) (all drugs

were kind gifts from Henrique von Gersdorff's lab) were pressure injected into the hindbrain

ventricle using an ASI MPPI-3. Between 15 and 35 min after injection, Dextran was imaged,

followed by another 10 min of GCaMP6s imaging. Dextran surrounding the imaged OPCs was

used to verify the drug delivery. TTX, IEM1460, AP5, and cadmium all caused fish paralysis,

further verifying efficient delivery of active drugs. After imaging, fish were removed from

mounting media. Larvae injected with control solution recovered full swimming activity within 3

minutes, and drug-injected paralyzed larvae maintained normal heartbeats for at least 3 hours.

Quantification and image analysis

OPC labeling during development

We used *sox10* and *olig1* to label OPCs. *sox10* drives expression in both OPCs and myelinating

oligodendrocytes while *olig1* labels only OPCs. OPCs and myelinating oligodendrocytes can be

distinguished by cell morphology: OPCs only possess processes and myelinating

oligodendrocytes possess sheaths. Animal ages and corresponding OPC activity: 55-60 hpf

(hours post-fertilization, when OPCs are born, and migrate), 3 dpf (when OPCs initiate

differentiation), 5-14 dpf (when many OPCs complete differentiation.)

Alignment rate and colocalization ratio

Images of fixed tissue were analyzed in Imaris. "Surface" function was used to create a cell

surface and "Spot" function was used to generate $Spot_{PSS\ puncta}$. Then the cell surface was used

as a mask to generate $Spot_{PSS\ puncta\ in\ OPCs}$, which was further verified by crosschecking its overlap

with $Spot_{PSS\ puncta}$. $Spot_{PSS\ puncta\ in\ OPCs}$ was analyzed for puncta number or density. To examine the

alignment of MAGUK (Gephyrin) and Synapsin, we used "Spot" function to generate $Spot_{Synapsin}$

$_{puncta}$. By determining the proximity $Spot_{PSS\ puncta\ in\ OPCs}$ to $Spot_{Synapsin\ puncta}$ using MATLAB-enabled

"Colocalize Spots" function (<0.1 μm), we could identify PSSs in OPCs that align with Synapsin

and calculate the alignment rate. To calculate the colocalization ratio of MAGUK (Gephyrin)-

Synapsin, we used the Imaris "Coloc" function to generate a new M&S coloc channel based on

the overlapping pixels between MAGUK (Gephyrin) and Synapsin channel above thresholds.

"Surface" function was used to generate M&S coloc surface of the whole section, MAGUK

(Gephyrin) (M) surface of the whole section, and OPC surface. OPC surface was used as a mask

to generate masked M&S coloc channel and MAGUK (Gephyrin) channel. These masked

channels were then used to generate M&S coloc surface in OPCs and M surface in OPCs." The

colocalization ratio in OPCs and neurons was calculated as follows:

$$Colocalization\ ratio_{OPCs} = \frac{Volume_{M\&S\ coloc\ surface\ in\ OPCs}}{Volume_{M\ surface\ in\ OPCs}};$$

$$Colocalization\ ratio_{neurons} =$$

$$\frac{Volume_{M\&S\ coloc\ surface\ of\ the\ whole\ section} - Volume_{M\&S\ coloc\ surface\ in\ OPCs}}{Volume_{M\ surface\ of\ the\ whole\ section} - Volume_{M\ Surface\ in\ OPCs}}.$$

PSS puncta identification and analysis

Live Images of PSD-95-GFP and GFP-Gephyrin were processed in ImageJ and synaptic puncta

were identified and tracked with Trackmate[39]. The images were first maximum projected and

stabilized using "Image Stabilizer." In Trackmate, parameters were set as follows: LOG mode,

diameter 0.75 μm, Quality>100, contrast>0.19 and signal noise >0.35, LAP linking with 0.5 μm

distance and 2 max gap frames allowed were applied. The identified puncta at each frame were

then manually examined to remove poor quality tracks. Neighboring puncta may sometimes be

incorrectly linked between timelapses; therefore, manual examinations were done to fix these

link errors. The data output was analyzed in MATLAB with custom scripts to calculate puncta

number per cell at each timepoint, puncta duration and diameter. Heatmaps of puncta over

time were generated using the output data with Heatscatter

(https://www.mathworks.com/matlabcentral/fileexchange/47165-heatscatter-plot-for-

variables-x-and-y). To generate figures of mean and standard deviation as shade, we used a

modified script (https://www.mathworks.com/matlabcentral/fileexchange/29534-stdshade).

To analyze the puncta number during OPC differentiation, time points when OPC shows first

sheath (differentiation) and when myelinating oligodendrocyte's last process disappeared and

all process transformed into sheaths (maturation) were noted for each cell. In MATLAB, this

information is used to divide the puncta number into before and after differentiation or

maturation for each cell. The traces were generated after the data was realigned to set the

timepoint of differentiation or maturation as time 0.

Ca^{2+} events analysis

myrGCaMP6s images of z-stacks were maximum projected and stabilized using "Image

Stabilizer." The images were assigned with random names using "blindr"

(https://github.com/U8NWXD/blindr) before analysis and reassigned with the original names

after analysis. The regions of Ca^{2+} events (ROI) were identified by "GECIquant"[40] first and then

manually examined frame by frame. The intensity and area of each ROI were exported and

processed in MATLAB through custom scripts. Briefly, $\Delta F/F_{average}$ ($\Delta F=F- F_{average}$) was calculated

to generate traces and to identify intensity peaks using "findpeaks" in MATLAB with filtering

parameters (3 standard deviations above average, prominence larger than 40% of the baseline,

and peak area larger than 2.5). Rare elevated Ca^{2+} signal for extended periods of time (longer

than 50 seconds) were excluded. Peaks were manually verified against the image. Several Ca^{2+}

characteristics were measured: peak amplitude, peak duration, Ca^{2+} microdomain number, Ca^{2+}

microdomain area, and frequency of Ca^{2+} activity. A region with multiple OPCs was analyzed for

TeTxLc experiment, while single OPC was analyzed for the rest of experiments. To examine the

proximity of synaptophysin-TagRFP to Ca^{2+} microdomain, Ca^{2+} microdomains were identified as

above, and their locations were first compared to synaptophysin-TagRFP manually plane by

plane and then verified by Imaris. In Imaris, using "Spot" function and "Surface" function, we

generated synaptophysin-TagRFP puncta and OPC surface, with which we identified the puncta

that were close to OPC surfaces using a MATLAB-enabled "Spot-to-Surface" function (0.1 µm

threshold). These puncta were manually checked against Ca^{2+} microdomains identified through

morphological landmarks in Imaris.

Oligodendrocyte and myelin sheath analysis

To analyze myelin sheath length and number, we used ImageJ ROI manager to trace all sheaths

of sparsely labeled oligodendrocytes between segments 1 and 15 in spinal cord. The number of

oligodendrocytes analyzed per fish was limited to 3 to ensure even weight among individual fish.

Sheath traces and the corresponding images were blinded using "Blind analysis encrypter." The

sheath length and number were then analyzed in MATLAB with custom scripts and matched

with the original genotype following analysis.

Statistics

Data are shown as mean ± SEM., with individual data points shown as circles. We conducted

Shapiro-Wilk's test and Levene's test to verify data normality and homoscedasticity,

respectively, before performing two-sided *t*-test for unpaired groups, paired *t*-test for paired

groups, and one-way ANOVA and Tukey's test for more than 3 groups. If the data did not pass

normality tests, we used Kruskal-Wallis test for more than 3 groups. These analyses were

performed either in GraphPad Prism or MATLAB. The significance and statistical analyses

employed were indicated in figure legends. Animal and cell number used for each experiment,

t-values and df for *t*-tests, and *F* values and df for ANOVA were indicated in either figure

legends or Supplemental Table 2. Data collection was not performed blindly due to the

experimental conditions, but analyses were performed blindly as indicated. Sample size was

chosen based on standard deviation and effect (http://www.biomath.info/power/ttest.htm).

Sex is not yet determined in zebrafish at the ages employed in our study and so cannot be

considered as a biological variable.

Reporting summary

Further information on research design is available in the Nature Research Reporting Summary

linked to this manuscript.

Data availability

All research materials are available upon request. Source data are provided with this paper.

Code availability

All custom scripts are available upon request.

31.   Almeida, R. G. & Lyons, D. A. Intersectional gene expression in zebrafish using the split

KalTA4 system. *Zebrafish* **12**, 377–386 (2015).

32.   Fuhrmann, J. C. *et al.* Gephyrin interacts with dynein light chains 1 and 2, components of

motor protein complexes. *J. Neurosci.* **22**, 5393–5402 (2002).

33.   Smolinsky, B., Eichler, S. A., Buchmeier, S., Meier, J. C. & Schwarz, G. Splice-specific

functions of gephyrin in molybdenum cofactor biosynthesis. *J. Biol. Chem.* **283**, 17370–

17379 (2008).

34.   Craven, S. E., El-Husseini, A. E. & Bredt, D. S. Synaptic targeting of the postsynaptic

density protein PSD-95 mediated by lipid and protein motifs. *Neuron* **22**, 497–509 (1999).

35.   Topinka, J. R. & Bredt, D. S. N-terminal palmitoylation of PSD-95 regulates association

with cell membranes and interaction with K+ channel K(v)1.4. *Neuron* **20**, 125–134 (1998).

36.   Kwan, K. M. *et al.* The Tol2kit: A multisite gateway-based construction Kit for Tol2

transposon transgenesis constructs. *Dev. Dyn.* **236**, 3088–3099 (2007).

37.   Wyart, C. *et al.* Optogenetic dissection of a behavioural module in the vertebrate spinal

cord. *Nature* **461**, 407–410 (2009).

38.   Di Donato, V. *et al.* 2C-Cas9: A versatile tool for clonal analysis of gene function. *Genome*

Res. **26**, 681–692 (2016).

39.   Tinevez, J. Y. *et al.* TrackMate: An open and extensible platform for single-particle

tracking. *Methods* **115**, 80–90 (2017).

40.   Srinivasan, R. *et al.* Ca2+ signaling in astrocytes from Ip3r2 -/- mice in brain slices and

during startle responses in vivo. *Nat. Neurosci.* **18**, 708–717 (2015).

Acknowledgments

This work was supported by an NMSS postdoctoral fellowship and a Warren Alpert

Distinguished Scholar Award (JL) and NIH/NINDS grant 1R21NS120650 (KRM). We thank Dwight

Bergles and members of the Monk lab for discussions and critical reading of the manuscript. We

are indebted to Austin Forbes, Tia Perry, and Grace Halsell-Vore for fish care. We thank Rafael

Almeida, Paul Brehm, Marc Freeman, Henrique von Gersdorff, Eric Gouaux, David Lyons, Alex

Nechiporuk, and Teresa Nicolson for reagents and suggestions as well as Fernanda Coelho,

Stefanie Kaech Petrie and OHSU microscopy core staff for feedback and assistance in imaging.

We thank Lori Vaskalis for graphic design.

Author Contributions

J.L. and K.R.M. conceived the project. J.L. carried out experiments and data analyses. T.M.

generated constructs for *nlgn3a* and *nlg3b* disruption. T.C. provided key resources prior to their

publication. J.L. and K.R.M wrote the manuscript, and all authors edited the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information is available for this paper.

Correspondence and requests for materials should be addressed to Jiaxing Li or Kelly Monk.

Figure 1. PSD-95-GFP and GFP-Gephyrin imaging reveals dynamic synapses that assemble at specific locations in OPCs in vivo. a-b, Representative images of PSD-95-GFP and GFP-Gephyrin (magenta) in spinal cord (**a**) OPCs and (**b**) neurons at 3 dpf. Cell membranes indicated with cyan. **c**, Quantification of puncta number per cell within 30 minutes for PSD-95-GFP and GFP-Gephyrin in neurons and OPCs, normalized to the average. Black line, average; shaded area, sd. **d-e**, Quantification of PSD-95-GFP (green) and GFP-Gephyrin (magenta) (**d**) puncta duration in log scale and (**e**) % of stable puncta that lasts over 10,000 seconds in neurons and OPCs at 3 dpf. From left, n = 13, 3, 5 and 7 cells, from 6, 3, 4 and 4 fish. **f,g**, Heatmaps of puncta appearance frequency over time for (**f**) PSD-95-GFP and (**g**) GFP-Gephyrin. 30% of max frequency is used as coloring threshold: below threshold, grey; above threshold, a shade of magenta corresponding to the frequency. Loci with high puncta appearance frequency (> 3 sd above average) are marked with arrowheads. The numbers with green arrowheads indicate puncta shown in time lapse images in Extended Data Fig. 2c-d. Bottom left, snapshots of OPCs with puncta marked with magenta circles. **h**, Quantification of loci with high puncta appearance frequency in OPCs. From left, n = 3 and 7 cells from 3 and 5 fish. All data are represented as mean ± SEM; unpaired *t*-test; Scale bar, 5 μm.

Figure 2. PSSs in OPCs are developmentally regulated. **a**, Quantification of PSD-95-GFP (green) and GFP-Gephyrin (magenta) puncta number per cell in the spinal cord at 55 hpf and 3 dpf by *in vivo* imaging. **b**, Quantification of MAGUK (green) and Gephyrin (magenta) puncta number per cell in spinal cord at different ages by IHC. **c-d**, Quantification of (**c**) PSD-95-GFP and (**d**) GFP-Gephyrin puncta number in OPCs before and after the last process disappearance, which is set as time zero. Left, traces show the average (black line) and the sd (shade) with individual traces (grey line). Right, paired comparison before and after time zero. **c**, n= 12 cells from 8 fish; **d**, n= 12 cells from 8 fish. **e**, Representative images of PSD-95-GFP and GFP-Gephyrin (magenta) in differentiating oligodendrocytes (cyan) at 55-60 hpf. All data are represented as mean ± SEM; N.S., not significant; (**a**) unpaired *t*-test, (**b**) one-way ANOVA followed by Tukey's test, and (**c-d**) ratio-paired *t*-test; scale bar 5 μm.

Figure 3. Synapses and VGCCs, but not NMDA or CP-AMPA receptors, contribute to generating intrinsic OPC Ca^{2+} activity in vivo. **a**, Representative images of plasma membrane-targeting myrGCaMP6s in an OPC (grey) showing activity in microdomains (cyan). **b-c**, Representative images of (**b**) an OPC (grey) visualized by myrGCaMP6s showing (**c**) proximity between an microdomain with Ca^{2+} activity (cyan) and presynaptic synaptophysin-tagRFP (magenta). **d**, 3-D reconstruction showing synaptophysin-tagRFP puncta (magenta) that are within 0.1 μm from the OPC surface (arrowheads). **e**, Quantification of the percentage of Ca^{2+} microdomains that are associated with synaptophysin-tagRFP (n = 7 cells from 7 fish). **f**, Representative images of OPCs neighboring few (<4) and many (≥4) axons expressing *TeTxLc-tdtomato*. **g**, Quantification of frequency of Ca^{2+} activity and peak amplitude in OPCs that neighbor few (<4) or many (≥4) TeTxLc-expressing axons at 5 dpf. From left, n = 14 and 26 cells from 8 and 15 fish. **h**, Schematic of hindbrain ventricle injection. **i**, Quantification of Ca^{2+} microdomain number per OPC and frequency of Ca^{2+} activity per microdomain in OPCs before (blue) and after (magenta) injection of 1 nL of AP5 (30 mM), Cadmium (100 mM), or IEM1460 (20 mM) solution at 5 dpf. Paired data indicated with grey line. From left, n = 13, 16 and 13 cells from 7, 7 and 7 fish. All data are represented as mean ± SEM; N.S., not significant; (**g**) unpaired *t*-test or (**i**) paired *t*-test; scale bar 5 μm.

Figure 4. Synaptic component nlgn3b is important for oligodendrocyte differentiation and myelination.
a, Morphology of control (ctrl), and *nlgn3b*-targeted myelinating oligodendrocytes at 5 dpf. Top, 3D-reconstructed single oligodendrocyte with sheath (colored) and non-sheath (white). Bottom, single plane with sheath (black arrowheads) and processes (magenta arrowheads). **b-c**, Quantification of total length of myelin sheaths of myelinating OLs at (**b**) 3 dpf and (**c**) 5 dpf in ctrl, *nlgn3a*-, and *nlgn3b*-targeted fish. From left, 3 dpf, n = 61, 19 and 25 cells from 18, 14 and 18 fish; 5 dpf, n = 66, 56 and 47 cells from 24, 36 and 30 fish. All data are represented as mean ± SEM; N.S., not significant; (**b,c**) Kruskal-Wallis test; scale bar 5 μm. **d**, Schematic model: **d'**, synapses containing MAGUK (green) and Gephyrin (red) form between neurons and OPCs; PSSs in OPCs (green with black border) increase in number through development and decrease upon differentiation, while disruption of synaptic gene *nlgn3b* impairs OPC differentiation; **d''**, PSSs in OPCs are dynamic with frequent assembly, and many of them do not align with presynaptic active zone (magenta); **d'''**, Neuron-OPC synapses are in proximity to some Ca^{2+} microdomains (blue), and, together with VGCCs (Black), contribute to generating intrinsic OPC Ca^{2+} activity. Evoked synaptic release contributes to intrinsic OPC Ca^{2+} activity at 3 dpf but not 5 dpf.

646 ***Extended Data Figure 1. PSD-95 (dlg4), PSD-93 (dlg2), gephyrin, and nlgn3 are expressed in OPCs.***
647 Summary of RNA-seq expression reported of mammalian *Dlg4 (MAGUK)*, *Dlg2 (MAGUK)*, *Gephyrin*, and
648 *Nlgn3* in different CNS cell types[26].

Extended Data Figure 2. in vivo analysis of PSSs. a, Quantification of PSD-95-GFP (green) and GFP-Gephyrin (magenta) puncta diameter in neurons and OPCs at 2-3 dpf in the spinal cord. From left, n = 10, 11, 15 and 29 cells from at least 8 fish. **b**, Representative images of GFP-Gephyrin (magenta) as an OPC remodels its processes (cyan) at 3 dpf. 3 enlarged regions were shown at different time points (minutes) and puncta were indicated with arrowheads. **c-d**, Representative time-lapse images of (**c**) PSD-95-GFP and (**d**) GFP-Gephyrin within OPC processes. The numbers indicate the corresponding puncta in **Fig. 1f**,**g**. Puncta 1 and 3 are in stable processes; puncta 2 and 4 are in dynamic processes. The dashed lines outline the process, and arrowheads indicate puncta appearance. Time unit is in minutes. All data are represented as mean ± SEM; (**a**) unpaired *t*-test. Scale bar 5 μm.

Extended Data Figure 3. Postsynaptic organizers, MAGUK, PSD-95, and Gephyrin, are present in
zebrafish spinal cord OPCs. **a**, Schematic fish model showing sagittal planes. **b-c**, Two representative
single-sagittal-plane IHC images showing (**b**) MAGUK and (**c**) Gephyrin within OPCs in *Tg(sox10:GFP-caax)*
spinal cord at 3 dpf. OPCs identified by GFP and shown in cyan. Arrowheads indicate PSS puncta. The
color was inverted to optimize visualization. **d**, Schematic fish model showing the transverse planes. **e-g**,
Representative images of (**e**) MAGUK, (**f**) PSD-95 and (**g**) Gephyrin staining in
Tg(olig1:Kalta4,10xUAS:myrGFP) spinal cord at 3 dpf. From left, spinal cord sections, enlarged images,
and 3D-reconstructed images to reveal OPC surface and PSS puncta (magenta sphere). **h-i**,
Quantification of MAGUK and Gephyrin (**g**) puncta number in OPCs and (**h**) puncta density in OPCs. **h**, n=
= 20 and 24 cells from 9 and 10 fish; **i**, n = 24 and 30 cells from 9 and 10 fish. All data are represented as
mean ± SEM; unpaired *t*-test; scale bar 5 μm.

Extended Data Figure 4. Many but not all PSS puncta in OPCs align with presynaptic structures. a-b, Representative single-sagittal-plane IHC images showing (**a**) MAGUK and (**b**) Gephyrin with presynaptic marker Synapsin in *Tg(sox10:GFP-caax)* spinal cord at 3 dpf. From left, merged channels, MAGUK or Gephyrin channel, and Synapsin channels. Black arrowheads indicate the alignment of postsynaptic puncta with presynaptic Synapsin. Magenta arrowheads point to postsynaptic puncta in OPCs that do not align with presynaptic Synapsin. **c-h**, Representative transverse IHC images of Synapsin and (**c-e**) MAGUK or (**f-h**) Gephyrin staining in *Tg(olig1:Kalta4,10xUAS:myrGFP)* spinal cord OPCs at 3 dpf. From left, (**c,f**) spinal cord sections and zoomed-in 3D reconstructed images to reveal PSS puncta (green sphere) (**d,g**) that align with presynaptic puncta, and (**e,h**) that do not. The arrowheads indicate the PSS puncta. **i**, The % of MAGUK and gephyrin puncta on OPC processes that align with presynaptic Synapsin in forebrain & midbrain, hindbrain, and spinal cord at 3 dpf. From left, n= 32, 57, 38, 33, 16 and 39 cells from at least 10 fish. **j**, Quantification of MAGUK (green) and Gephyrin (magenta) colocalization ratio with Synapsin in OPCs, normalized to the colocalization ratio in neurons at 2 dpf and 14 dpf. From left, n = 9, 7, 9 and 7 fish. All data are represented as mean ± SEM; N.S., not significant; (**i**) one-way ANOVA followed by Tukey's test or (**j**) paired *t*-test; Scale bar 5 μm.

Extended Data Figure 5. PSS numbers in OPCs show few regional and no subgroup differences. a,
Quantification of MAGUK (green) and Gephyrin (magenta) puncta number per OPC at 3 dpf across the
CNS regions indicated by the schematic: forebrain & midbrain, hindbrain, and spinal cord. From left, n =
34, 59, 43, 34, 18 and 40 cells from at least 10 fish. **b**, Schematic model showing the neuron-rich (NR)
and axon & synapse-rich (AR) regions and two representative images showing OPCs with soma localized
to NR (top) and AR (bottom) regions in *Tg(olig1:Kalta4,10xUAS:myrGFP)* spinal cord. **c**, Quantification of
MAGUK (green) and Gephyrin (magenta) puncta number per cell between OPCs with their soma
localized to NR and AR regions at 2-3 dpf. From left, n = 14, 5, 15 and 10 cells from at least 5 fish. All data
are represented as mean ± SEM; N.S., not significant; (**a**) one-way ANOVA followed by Tukey's test and
(**c**) unpaired *t*-test; scale bar 5 μm.

Extended Data Figure 6. Synapses contribute to generating intrinsic OPC Ca^{2+} activity in vivo. a, An OPC
visualized by myrGCaMP6s. **b,** In the enlarged region from (**a**), two Ca^{2+} spikes initiate as puncta and
then spread to form microdomains, indicated by blue arrowheads. The time is in seconds. **c,** The two
microdomains from (**b**) were in proximity (within 0.1 μm) with presynaptic synaptophysin-tagRFP
(magenta), indicated by white arrowheads. **d,** Quantification of basal intensity and Feret's diameter in
OPCs that neighbor few (< 4) or many (≥ 4) TeTxLc-expressing axons at 5 dpf. From left, n = 14 and 26
cells from 8 and 15 fish. **e,** Representative single-plane images of individual spinal cord OPCs
immediately and 10 min after hindbrain ventricle injection of dextran MW3000 Texas Red (magenta) in
Tg(olig1:Kalta4,10xUAS:myrGFP) larva. **f,** Quantification of peak duration and area of Ca^{2+} microdomains,
frequency of Ca^{2+} activity and peak amplitude before (blue) and after (magenta) APV, Cadmium, or
IEM1460 injection at 5 dpf. Data in pairs indicated with grey lines. From left, n = 13, 16 and 13 cells from
7, 7 and 7 fish. All data are represented as mean ± SEM; N.S., not significant; (**d**) unpaired *t*-test and (**f**)
paired *t*-test; scale bar 5 μm.

Extended Data Figure 7. TTX has limited effects on intrinsic OPC Ca^{2+} activity at 3 dpf but not 5 dpf.
a, Quantification of Ca^{2+} microdomain number per OPC, frequency of Ca^{2+} activity per microdomains, peak amplitude, peak duration, and area of Ca^{2+} microdomain in OPCs before (blue) and after (magenta) injection of 1 nL of control or TTX (0.5 mM) solution. Paired data indicated with grey line. *Tg(sox10:Kalta4,10xUAS:myrGCaMP6s)* larvae at 3- and 5 dpf were analyzed. From left, n = 15 ,31, 13 and 17 cells from 7, 13, 6 and 9 fish. **b**, Representative images of primary motoneurons visualized by injecting *mnx1:gal4* into *Tg(10xUAS:myrGCaMP6s)*. **c**, Quantification of frequency of Ca^{2+} activity per cell before (blue) and after (magenta) TTX injection in motoneurons at 5 dpf. n = 7 and 6 cells from 5 and 5 fish. **d**, Representative image of an OPC surrounded by dextran MW3000 by the time of imaging after TTX injection. All data are represented as mean ± SEM; N.S., not significant; paired *t*-test; Scale bar 5 μm.

738
Extended Data Figure 8. A cell-specific knockdown approach effectively disrupts oligodendrocyte
development. **a**, Schematic of Tol2 plasmid (10xUAS:myrmScarletI-p2A-Cas9, U6:sgRNA1;U6:sgRNA2)
constructed and used to induce cell-specific Cas9 expression with membrane labeling and 2 separate
U6-driven sgRNAs. **b**, Schematic of larva with sparsely labeled oligodendrocytes resulting from
Tg(sox10:Kalta4) crossed with transgenic fish carrying the plasmid from (**a**). **c**, Representative images of
single myelinating oligodendrocyte in the spinal cord of *Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:ctrl-*
sgRNA1;U6:ctrl-sgRNA2) and *Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:myrf^{ex3}-sgRNA;U6:myrf^{ex4}-sgRNA)*
at 6 dpf. **d**, Quantification of myelin sheath length from (**c**). n = 24 and 22 cells from 15 fish. **e**,
Representative image of an OPC and a myelinating oligodendrocyte (OL). **f**, 4 single planes of a
trapezoid-shaped process in an otherwise myelinating OL in *nlgn3b*-targeted larvae at 5 dpf. Myelin
sheaths indicated by black arrowheads and processes by magenta arrowheads. All data are represented
as mean ± SEM; N.S., not significant; unpaired *t*-test; scale bar 5 μm.
751
752

Supplementary information

Supplementary video 1. Time-lapse images of PSD-95-GFP in an OPC.
Representative video of sox10:myrTagRFP-T-P2A-PSD95-GFP in an OPC from the spinal cord of a 3 dpf larva. GFP-Gephyrin (magenta) puncta are dynamic in location and size, while the OPC processes are relatively stable (cyan).

Supplementary video 2. Time-lapse images of PSD-95-GFP in a neuron.
Representative video of 10xUAS:PSD95-GFP in a neuron in the spinal cord of a *Tg(sox10:Kalta4)* larva at 3 dpf.

Supplementary video 3. Time-lapse images of GFP-Gephyrin in an OPC.
Representative video of sox10:myrTagRFP-T-P2A-GFP-Gephyrin in an OPC from the spinal cord of a 3 dpf larva. GFP-Gephyrin (magenta) puncta are dynamic in location and size, while the OPC migrates and remodels its processes (cyan).

Supplementary video 4. Time-lapse images of GFP-Gephyrin in a neuron.
Representative video of 10xUAS:GFP-Gephyrin in a neuron in the spinal cord of a *Tg(sox10:Kalta4)* larva at 3 dpf.

Supplementary video 5. Time-lapse images of myrGCaMP6s in an OPC.
Representative video of Ca^{2+} activity in an OPC in a *Tg(olig1:Kalta4,10xUAS:myrGCaMP6s)* larva at 5 dpf.

Supplementary video 6. Time-lapse images of myrGCaMP6s in an OPC reveal somal Ca^{2+} activity.
Video of Ca^{2+} activity in an OPC with multiple events including a single somal event in a *Tg(olig1:Kalta4,10xUAS:myrGCaMP6s)* larva at 5 dpf.

Supplementary video 7. Time-lapse images of myrGCaMP6s in an OPC before and after injection of control solution.
Representative 20 min-video of Ca^{2+} activity in an OPC before and after injection of control solution in a *Tg(olig1:Kalta4,10xUAS:myrGCaMP6s)* larva at 5 dpf. Yellow arrows indicate the Ca^{2+} events.

Supplementary video 8. Time-lapse images of myrGCaMP6s in an OPC before and after injection of cadmium.
Representative 20 min-video of Ca^{2+} activity in an OPC before and after injection of cadmium in a *Tg(olig1:Kalta4,10xUAS:myrGCaMP6s)* larva at 5 dpf. Yellow arrows indicate the Ca^{2+} events.

Supplementary video 9. Time-lapse images of myrGCaMP6s in an OPC before and after injection of TTX at 3 dpf.
Representative 20-minute video of Ca^{2+} activity in an OPC before and after injection of TTX in a *Tg(sox10:Kalta4,10xUAS:myrGCaMP6s)* larva at 3 dpf. Yellow arrows indicate the Ca^{2+} events.

Supplementary Table 1. A list of primers used in this study.
Supplementary Table 2. Statistical information.
Supplementary Table 3. Fish used for experiments.

sgRNA   -   +   -   +

Supplementary Figure 1. Selected sgRNAs are efficient cutting targeted genomic regions.
Representative gel images of digested PCR product of genomic regions targeted by the sgRNAs. Each
lane represents a single embryo at 1 dpf. The left 4 lanes are uninjected (-) and the right 4 lanes are
injected (+) with sgRNAs and Cas9 protein.