Synapses and Ca^{2+} activity in oligodendrocyte precursor cells predict where myelin

sheaths form

Jiaxing Li[1, *], Tania Miramontes[1], Tim Czopka[2], Kelly Monk[1, *]

1 Vollum Institute, Oregon Health & Science University, Portland, OR, USA

2 Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK

*Corresponding authors

SUMMARY

In the nervous system, only one type of neuron-glial synapse is known to exist: that between

neurons and oligodendrocyte precursor cells (OPCs). Neuron-OPC synapses are thought to

bridge neuronal activity to OPCs. However, their composition, assembly, downstream

signaling, and *in vivo* functions remain largely unclear. Here, we use zebrafish to address

these questions and identify postsynaptic molecules PSD-95 and Gephyrin in OPCs. They

increase during early development and decrease upon OPC differentiation. PSD-95 and

Gephyrin in OPCs are highly dynamic and frequently assemble at "hotspots." Gephyrin

"hotspots" and synapse-associated Ca^{2+} activity in OPCs predict where myelin sheaths form in

oligodendrocytes. Further analyses reveal that spontaneous synaptic release from neurons

mediates most OPC Ca^{2+} activity, while evoked synaptic release contributes only in early

development. Finally, disruption of the synaptic molecule Nlgn3 impairs OPC differentiation

and myelination. Together, we propose that neuron-OPC synapses are dynamically

assembled and predetermine myelination patterns through Ca^{2+} signaling.

INTRODUCTION

Oligodendrocyte precursor cells (OPCs) are a dynamic cell type in the central nervous system (CNS) with frequent migration, proliferation, process remodeling, and differentiation. OPCs and their interactions with neurons play important roles in CNS health and disease such as myelination and remyelination[1], neural network remodeling[2], and glioma progression[3]. Synapses form between neurons (presynaptic) and OPCs (postsynaptic) and represent the only type of glial synapse in the nervous system. The composition of neuron-OPC synapses is largely unknown, with the exception of some synaptic receptors (*e.g.*, glutamatergic and GABAergic) that were revealed by pharmacology and electrophysiology[4]. Neuron-OPC synapse assembly and regulation *in vivo* are mysterious, due to a paucity of tools to label these structures. Previous studies knocking down single synaptic receptor types found diverse and sometimes conflicting effects on OPC development and myelination: blocking NMDA receptors reduced or had no effect on myelination[5,6]; blocking GABA receptors enhanced or had no effect on OPC proliferation[7,8]. This is possibly due to redundancy, as OPCs possess multiple synaptic receptors[4]. Therefore, our understanding of the functions of neuron-OPC synapses is incomplete.

OPCs are hypothesized to sense the "activity" of neurons through neuron-OPC synapses. Several studies support this idea: neuronal activity can modulate OPC development and myelination[9-13]; *ex vivo* rodent studies using axonal stimulation observed synapse-dependent Ca^{2+} increases in oligodendrocyte lineage cells, mostly in somas[14-16]. However, the *in vivo* relationship between neuronal activity, neuron-OPC synapses, and OPC Ca^{2+} activity without exogenous stimulation is unknown. Furthermore, the role of synapses in forming microdomain (MD) Ca^{2+} events in OPC processes[17,18] is not addressed.

Here we identify postsynaptic scaffold proteins membrane-associated guanylate

kinases (MAGUKs), PSD-95 (a MAGUK member), and Gephyrin in zebrafish OPCs. The

postsynaptic puncta number in OPCs increases during early development, reduces upon OPC

differentiation, and exhibits small differences across the CNS regions but no difference between

OPC subgroups. *In vivo* imaging of PSD-95 and Gephyrin puncta in OPCs reveals that the puncta

are highly dynamic, resulting in extensive synaptic coverage. Although dynamic, postsynaptic

puncta repeatedly assemble at approximately 10 hotspots per OPC. Gephyrin hotspots at the

OPC stage predict where myelin sheaths form at the oligodendrocyte stage, in a synaptic

release-dependent manner. Further analysis reveals that Ca^{2+} activity is associated with neuron-

OPC synapses and similarly predicts where myelin sheaths form. The majority of *in vivo* OPC

Ca^{2+} events occur in MDs, require spontaneous synaptic release and, during early development,

require evoked synaptic release from neurons. Finally, we find that disruption of *neuroligin3*

(*nlgn3*), a gene important for synapse structure and function, impairs OPC development and

myelination. Collectively, these results suggest that dynamically assembled neuron-OPC

synapses predetermine myelination patterns through spontaneous and evoked synaptic Ca^{2+}

signaling in OPCs.

RESULTS

Postsynaptic organizers MAGUK/PSD-95 and Gephyrin are present in zebrafish spinal cord

OPCs

Transcript expression of *psd-95/dlg4* (often associated with Glutamate receptors) and

gephyrin (often associated with GABA and Glycine receptors) were previously reported in OPCs

(Extended Data Fig. 1a-b)[17,19]. We detected the presence of MAGUK/PSD-95 and Gephyrin

puncta in zebrafish spinal cord OPCs using immunohistochemistry (IHC) at 3 days post-

fertilization (dpf) (Fig. 1a-c and Extended Data Fig. 1c-f). We identified 16.6 ± 2.5 MAGUK

puncta and 29 ± 3.1 Gephyrin puncta per OPC (Fig. 1d), which were localized to OPC somas and

processes (Extended Data Fig. 1g). Gephyrin antibody specificity was verified in zebrafish with

gephyrin CRISPR/Cas9 knockdown (Extended Data Fig. 1h-i). To further validate these

observations, we used Fibronectin intrabodies generated by mRNA display (FingRs) that bind to

endogenous PSD-95 and Gephyrin *in vivo*[20]. By expressing FingR.GFP.PSD-95 and

FingR.GFP.Gephyrin in OPCs, we found FingR.GFP enrichment in somas and puncta in processes

(Fig. 1e), similar to previous reports in neurons[20]. There were 20.3 ± 2.4 FingR.GFP.PSD-95 and

21.1 ± 4.0 FingR.GFP.Gephyrin puncta per OPC, which was not significantly different than IHC

(Fig. 1d). These results identify MAGUK/PSD-95 and Gephyrin puncta in OPCs, extending

previous work that reported PSD-95 in myelin sheaths[21].

Given that OPCs are heterogeneous according to transcriptomic and electrophysiological

studies[22,23], we examined the number of postsynaptic puncta across developmental stages and

CNS regions using IHC. We observed an increasing trend through development: both MAGUK

and Gephyrin puncta numbers increased from 2 dpf (14.1 ± 3.1 MAGUK; 23.6 ± 2.8 Gephyrin) to

14 dpf (56 ± 14.2 MAGUK; 50.9 ± 8.9 Gephyrin) (Extended Data Fig. 2a). The postsynaptic

puncta numbers per cell were overall similar across CNS regions, except that Gephyrin puncta

numbers were higher in the forebrain and midbrain (20.2 ± 1.2) compared to the spinal cord

(13.8 ± 0.9) (Extended Data Fig. 2b), suggesting a preference of OPCs to form synapses with

inhibitory neurons in the brain. Within the zebrafish spinal cord, OPC subgroups were

previously reported to reside in different regions (Extended Data Fig. 2c) [17]. We found that the postsynaptic puncta numbers were similar between these OPC subgroups (Extended Data Fig. 2d).

Not all MAGUK and Gephyrin puncta in OPCs are associated with other synaptic molecules

Next, we tested whether these puncta align with presynaptic structures. Using the pan-presynaptic marker Synapsin (Fig. 2a-f), we identified MAGUK and Gephyrin puncta in OPCs that align with (Fig. 2b, e) or that do not align with (Fig. 2c, f) Synapsin in transverse sections. We found similar results in sagittal sections (Extended Data Fig. 3a-b). $46.1 \pm 7.4\%$ and $38.3 \pm 4.8\%$ of MAGUK and Gephyrin puncta in an OPC, respectively, align with Synapsin in the spinal cord, and a similar percentage of puncta in an OPC align with Synapsin in other CNS regions (Fig. 2g). To compare pre- and post-synaptic association at neuron-OPC synapses *vs.* neuron-neuron synapses, we calculated colocalization ratios between Synapsin and MAGUK or Gephyrin. Compared to neuron-neuron synapses (100%, normalized) from the same sections, MAGUK at neuron-OPC synapses showed $103 \pm 22\%$ (2 dpf) and $70 \pm 8\%$ (14 dpf) colocalization, while Gephyrin at neuron-OPC synapses showed $69 \pm 10\%$ (2 dpf) and $87 \pm 5\%$ (14 dpf) colocalization (Extended Data Fig. 3c). In addition, $42.1 \pm 4.8\%$ of Gephyrin puncta in an OPC align with Glycine receptors (Fig. 2h-i), and the colocalization ratio between Gephyrin and Glycine receptors at neuron-OPC synapses ($53.2 \pm 6.6\%$) was less than that at neuron-neuron synapses (100%, normalized) from the same sections (Extended Data Fig. 3d). These results suggest that compared to neurons, more MAGUK and Gephyrin puncta in OPCs are not associated with other synaptic molecules, and therefore may not represent fully functional synapses.

132 *In vivo* **imaging reveals downregulation of PSD-95-GFP and GFP-Gephyrin upon OPC**

133 **differentiation**

134   To understand synapse assembly and disassembly in OPCs, we conducted *in vivo* time-

135 lapse imaging. Due to the susceptibility of FingR.GFP to photobleaching, we generated PSD-95-

136 GFP and GFP-Gephyrin constructs (see Methods) and drove their expression with *sox10*, a pan-

137 oligodendrocyte lineage promoter. At 3 dpf in the spinal cord, we found both markers present

138 in punctate and cytosolic forms in OPCs (Fig. 3a), consistent with patterns observed in

139 neurons[24,25]. PSD-95-GFP and GFP-Gephyrin puncta size in OPCs resembled that of neurons (Fig.

140 3a-b and Extended Data Fig. 4a). There were 28.7 ± 3.9 PSD-95-GFP and 27.2 ± 3.5 GFP-

141 Gephyrin puncta per OPC (Fig. 1d), which was similar to IHC and FingR.GFP results. The slightly

142 higher PSD-95-GFP puncta number may be due to increased synaptogenesis caused by PSD-95

143 overexpression[26]. We assessed presynaptic alignment of our expression constructs *in vivo*, and

144 found that 32.8 ± 5.5% PSD-95-GFP and 43.2 ± 3.8% GFP-Gephyrin aligned with synaptophysin

145 (Extended Data Fig. 4b-c), similar to IHC. In addition, we observed an increase of PSD-95-GFP

146 and GFP-Gephyrin from 2 dpf (4.6 ± 1.1 PSD-95-GFP; 6.9 ± 1.0 GFP-Gephyrin) to 3 dpf (29.4 ±

147 4.4 PSD-95-GFP; 27.2 ± 3.5 GFP-Gephyrin) (Extended Data Fig. 4d). This increasing trend during

148 early development was comparable to IHC results (Extended Data Fig. 2a), suggesting that the

149 GFP-tagged PSD-95 and Gephyrin are regulated similarly as the endogenous proteins.

150   We wondered if postsynaptic puncta assembly and disassembly were regulated upon

151 OPC differentiation. To test this, we tracked individual OPCs over the course of their

152 development and measured the postsynaptic puncta number per cell every 20 minutes.

153 Because OPC differentiation takes several hours for all processes to either retract or transform

into nascent myelin sheaths, we used the disappearance of the last exploratory process as a

reference point. When we aligned the results from different cells, we found a decreasing trend

during differentiation; the postsynaptic puncta numbers reduced from 9.8 ± 2.5 PSD-95-GFP

and 16.5 ± 2.8 GFP-Gephyrin before the reference point to 3.9 ± 1.2 PSD-95-GFP and 9.1 ± 2.0

GFP-Gephyrin after the reference point (Fig. 3c-d). In cases where we observed differentiating

oligodendrocytes that possessed both processes and myelin sheaths, puncta were almost

exclusively in processes (Fig. 3e), suggesting preferential localization of synaptic puncta to

processes rather than nascent myelin sheaths. Altogether, these results show a downregulation

of postsynaptic puncta upon OPC differentiation, consistent with reduced synaptic inputs

reported from *ex vivo* electrophysiological studies in rodents[27,28].

PSD-95-GFP and GFP-Gephyrin puncta in OPCs are dynamic and repeatedly assemble at

hotspots

We next sought to analyze the dynamics of PSD-95 and Gephyrin in OPCs *in vivo* using

our validated expression system. The number of postsynaptic puncta per cell remained stable in

neurons, but fluctuated in OPCs over 30 minutes of observation (Fig. 4a and Supplementary

Videos 1-2). Individual puncta in neurons lasted 14,041 ± 1,187 s or about 234 min (PSD-95-GFP)

and 12,348 ± 3,461s or about 206 min (GFP-Gephyrin), while those in OPCs lasted only 968 ±

105s or about 16 min (PSD-95-GFP) and 271 ± 53s or about 5 min (GFP-Gephyrin) (Fig. 4b). 78 ±

4% (PSD-95-GFP) and 41 ± 12% (GFP-Gephyrin) of puncta in neurons lasted more than 10,000

seconds or 167 min; conversely, only 0.4 ± 0.4% (PSD-95-GFP) and 0 ± 0% (GFP-Gephyrin) of

puncta in OPCs did so (Fig. 4c). OPCs possess both remodeling and stable processes[17]; we found

puncta assembly and disassembly in both types of processes (Fig. 4f-g), suggesting that

postsynaptic puncta dynamics in OPCs were not simply caused by process instability. In addition,

we tracked GFP-Gephyrin up to 7 hours and found that puncta from the same OPC covered a

large space with a convex hull volume of $1.21 \pm 0.21 \times 10^4$ μm^3, which was 50 times the volume

of an OPC (242 ± 28 μm^3, Fig. 4d-e). Therefore, the observed dynamic assembly enables a large

synaptic coverage by OPCs.

While postsynaptic puncta in OPCs were dynamic and covered a large space, some

postsynaptic puncta repeatedly assembled and disassembled in the same regions of OPC

processes (Fig. 4f-g). This raises the question if specific locations are preferred for synaptic

assembly. To test this, we plotted PSD-95-GFP and GFP-Gephyrin puncta over time and overlaid

heatmaps of puncta frequencies within 1 μm x 1 μm x 4 μm volumes (Fig. 4h-i). We found

approximately 10 such volumes where frequencies of PSD-95-GFP and GFP-gephyrin puncta

were 3 standard deviations higher than the average (Fig. 4j), and refer to these volumes as

hotspots. The Gephyrin hotspots represented only 0.3% of total synaptic space coverage of an

OPC (Fig. 4e). These results suggest that some postsynaptic puncta in OPCs assemble

repeatedly and preferentially in the confined spaces of hotspots.

Gephyrin hotspots in OPCs predict future myelin sheath formation in a synaptic release-

mediated manner

In zebrafish spinal cord, a myelinating oligodendrocyte forms an average of 10 myelin

sheaths[29], similar to the hotspot number we observed per OPC. We wondered if these hotspots

in OPCs were associated with future myelin sheath formation (Fig. 5a-b). To test this, we

measured the distances from Gephyrin hotspot centers (centroids of 1 μm x 1 μm x 4 μm

volumes) in an OPC to boundaries of sheaths formed after the OPC differentiated and

myelinated axons (Fig. 5c-d). We used the first nascent sheath appearance (time 0) to separate

the OPC stage and the myelinating oligodendrocyte stage. It takes up to 5 hours for an

oligodendrocyte to complete differentiation[29]; therefore, we identified where sheaths formed

approximately 6 hours after time 0 (Fig. 5c-d). In comparison, we selected 5-30 representative

points outside hotspots (nonhotspot) in an OPC at the beginning of imaging (see Methods) and

measured their distances to future myelin sheaths. The hotspot-sheath distance was

significantly smaller than the nonhotspot-sheath distance (Fig. 5e). Further, we examined the

percentage of hotspots or nonhotspots that predicted where myelin sheaths formed. We chose

1 μm as a threshold because we analyzed previous EM datasets of 5 dpf zebrafish[30] and found

an average distance of 1.4 μm between 2 myelinated axons in the dorsal spinal cord. Within 1

μm, 29.6 ± 8.7% of hotspots in an OPC predicted future sheath locations, while only 5.9 ± 3.4%

of nonhotspots did (Fig. 5f). Next, we asked if the relationship between OPC hotspots and

future myelin sheaths was dependent on synaptic release. We blocked synaptic release by

injecting tetanus toxin (TeNT) into ventricles between 2-3 dpf (Fig. 5c). We verified the TeNT

delivery to OPCs in the CNS with Dextran injection (see Methods). TeNT administration caused a

right-shift in hotspot-sheath distance distribution (Fig. 5g). The hotspot-sheath distance was not

significantly smaller than the nonhotspot-sheath distance with TeNT treatment (Fig. 5h). More

importantly, TeNT administration abolished the predication of hotspots within 1 μm (Fig. 5i).

This was not due to changes in OPC convex hull volume, OPC hotspot number, or myelin sheath

number (Extended Data Fig. 5a-c). Together, these results suggest that GFP-Gephyrin hotspots

are localized in proximity to future myelin sheaths and can predicate their location in a synapse-

release dependent manner.

OPC Ca^{2+} activity is associated with synapses and predicts myelin sheath formation

Given the role of Ca^{2+} signaling in sheath remodeling[18,31], we examined the relationship

of Ca^{2+} signaling events with synapses and sheath formation *in vivo*. With a plasma membrane-

targeted GCaMP6s, we analyzed OPC Ca^{2+} activity at single-cell resolution *in vivo* (Fig. 6a-b). We

found that 99% of Ca^{2+} events in OPCs occurred in MDs on processes, and 97% of OPCs

exhibited only MD Ca^{2+} activity during our recordings (Fig. 6c, Supplemental Video 3-4). We

examined the proximity between Ca^{2+} events in MDs and synapses, visualized by a presynaptic

marker, synaptophysin-TagRFP[17]. Through 3D reconstruction, we found that 34 ± 9% of MD Ca^{2+}

events were in <0.1 μm proximity to synaptophysin-TagRFP puncta at 5 dpf (Fig. 6d-f). We also

observed that Ca^{2+} events initiated from a "puncta"-like region in <0.1 μm proximity to

synapses (Fig. 6g-h). Next, we performed time-lapse imaging to measure the distance from MD

Ca^{2+} events in an OPC to sheaths formed after the OPC differentiated and myelinated axons (Fig.

6i-j). In comparison, we selected 5-30 representative points outside MDs (nonMD) in an OPC at

the beginning of imaging and measured their distances to future myelin sheaths. The MD-

sheath distance was significantly smaller than the nonMD-sheath distance (Fig. 6k). Next, we

examined the percentage of MDs that predicted where myelin sheaths formed. Within 1 μm,

32.0 ± 9.4% of MDs in an OPC predicted future sheath locations, while only 4.3 ± 1.2% of

nonhotspots did (Fig. 6l). These measurements of MD-sheath are similar to those for GFP-

Gephyrin hotspots and sheaths, suggesting that OPC Ca^{2+} activity *in vivo*, mainly occurring in

MDs, along with Gephyrin hotspots, predetermines where a subset of myelin sheaths form (Fig.

6m).

Spontaneous synaptic release contributes to generating OPC Ca^{2+} activity *in vivo*

Given the close relationship between Ca^{2+} activity and synapses, we investigated the

role of synapses in OPC Ca^{2+} activity *in vivo*. First, we blocked synaptic release by sparsely

expressing tetanus toxin light chain (TeNTlc)-tdtomato[32] in axons at 5 dpf (Fig. 7a). Compared to

OPCs that neighbored few TeNTlc-expressing axons (< 4, few), those that neighbored many

TeNTlc-expressing axons (> 4, many) exhibited significantly reduced frequency of Ca^{2+} activity

(few 0.020 ± 0.002 vs. many 0.013 ± 0.002 Hz), peak amplitude (few 1.81 ± 0.12 vs. many $1.4 \pm$

0.08 fold), basal signal intensity (few 1.00 ± 0.10 vs. many 0.74 ± 0.06), and MD length (few 7.03

± 0.38 vs. many 6.00 ± 0.25 µm) (Fig. 7b-c and Extended Data Fig. 6a). Next, we delivered

tetanus toxin (TeNT) to the CNS through ventricle injection at 5 dpf and analyzed Ca^{2+} activity

before and after injection (Fig. 7d). To ensure that TeNT (MW 150,000) was delivered

successfully, we coinjected Dextran (MW 3,000 or 150,000) and verified that Dextran

surrounded OPCs before Ca^{2+} imaging (Fig. 7e and Extended Data Fig. 6b). While control

solution had no effect on MD event number (before 17.0 ± 2.6 vs. after 15.5 ± 2.4) or frequency

of Ca^{2+} activity (before 0.0015 ± 0.0001 Hz vs. after 0.0014 ± 0.0001 Hz), TeNT administration

led to nearly 45% reduction in MD event number (before 25.2 ± 2.3 vs. after 14.4 ± 1.5) and

frequency of Ca^{2+} activity (before 0.0019 ± 0.0001 Hz vs. after 0.0010 ± 0.0001 Hz) (Fig. 7f, g,

Extended Data Fig. 6c and Supplementary Video 5).

These effects could be due to reduced neuronal activity and evoked synaptic release by

TeNT. To our surprise, we found that blocking neuronal action potentials with the voltage-gated

Na^+ channel blocker tetrodotoxin (TTX) had no effect on the OPC Ca^{2+} activity at 5 dpf (MD

number per OPC: before 2.9 ± 0.6 vs. after 2.4 ± 0.5. Frequency: before 0.0014 ± 0.0002 Hz vs.

after 0.0012 ± 0.0002 Hz) (Fig. 7h and Extended Data Fig. 6d). We excluded the possibility of

ineffective TTX delivery in three ways: we observed larval paralysis within 3 minutes post-

injection (30/30), a complete block of motor neuron Ca^{2+} activity (Extended Data Fig. 6e-f), and

dextran co-injected with TTX (dextran MW3000, larger than TTX) surrounded the imaged OPCs

(120/120). Therefore, the reduced Ca^{2+} activity upon TeNT injection was most likely due to the

effects of TeNT on vesicular release at neuron-OPC synapses, rather than neuronal activity. We

did, however, observe reduced OPC Ca^{2+} activity upon TTX injection at the earlier timepoint of 3

dpf (MD number per OPC: before 3.2 ± 0.4 vs. after 2.4 ± 0.3. frequency: before 0.0014 ± 0.0001

Hz vs. after 0.0010 ± 0.0001 Hz), consistent with a previous report[18] and suggesting a

diminishing role of neuronal activity and evoked synaptic release over development. Altogether,

our results establish a role for spontaneous synaptic release in generating OPC Ca^{2+} activity *in*

vivo.

Disruption of *nlgn3b* in OPCs impairs OPC development and myelination.

276   Given how synaptic hotspots and Ca^{2+} activity predict myelin sheath formation, we next

sought to examine whether synaptic assembly was important for OPC development and

myelination. We targeted *neuroligin3* (*nlgn3*), which localizes to both excitatory and inhibitory

synapses, regulates neuronal synapse structure and function[33], and is enriched in OPCs

(Extended Data Fig. 1a-b)[17,19]. We developed and verified a cell-specific CRISPR/Cas9-mediated

gene disruption system (Extended Data Fig. 7a-e) and targeted the duplicated *nlgn3* genes

(*nlgn3a* and *nlgn3b*) in zebrafish. We observed a reduced oligodendrocyte percentage among

nlgn3b-targeted cells at 3 dpf and 5 dpf (Extended Data Fig. 7f-g), suggesting that loss of *nlgn3*

disrupts proper OPC differentiation. At 5 dpf, all labeled oligodendrocytes in control animals

possessed only myelin sheaths; however, 7.6 ± 1.8% of *nlgn3b*-disrupted oligodendrocytes

possessed both myelin sheaths and cytoplasmic processes with abnormal trapezoidal structures

(Fig. 8a and Extended Data Fig. 7h-i), indicating developmental defects when Nlgn3b was lost.

Additionally, disruption of *nlgn3b* impaired myelin sheath formation (Fig. 8b): at 3 dpf,

individual sheath length (ctrl 13.6 ± 0.6 µm *vs. nlgn3b* 11.4 ± 0.5 µm), sheath number (ctrl 9.9 ±

0.3 *vs. nlgn3b* 9.2 ± 0.3), and total length of all sheaths (ctrl 125.6 ± 4.8 µm *vs. nlgn3b* 98.0 ± 4.1

µm) per oligodendrocyte were reduced (Fig. 8c); at 5 dpf, sheath number (ctrl 11.2 ± 0.5 *vs.*

nlgn3b 9.1 ± 0.4) and total length of all sheaths (ctrl 368.8 ± 11.7 µm *vs. nlgn3b* 290.4 ± 10.3 µm)

per oligodendrocyte were reduced (Fig. 8d). Together, these results suggest a role for Nlgn3

specifically and synaptic assembly more broadly in oligodendrocyte differentiation and

myelination.

DISCUSSION

The landmark discovery of neuron-OPC synapses over 20 years ago has inspired a

number of studies to examine their properties with electrophysiology and EM [4], yet we still lack

basic understanding of their assembly and function. Here, we establish an *in vivo* model of

these synapses in zebrafish OPCs (Fig. 8e): they are composed of postsynaptic MAGUK (PSD-95)

and Gephyrin, and are developmentally regulated; they lack consistent synaptic partners and

exhibit dynamic synapse features; they cover large spatial volumes over time, but frequently

and repeatedly assemble at specific and confined "hotspots"; Gephyrin hotspots and synaptic

release-mediated Ca^{2+} activity predict myelination patterns; and disruption of the synaptic

molecule Nlgn3b impairs OPC development and myelination.

Previous studies suggest that neuronal activity and presynaptic release bias OL

myelination to "active" axons[10,12]. However, it is not clear what mechanisms mediate this

preference. Here, we identify a postsynaptic mechanism in OPCs that depends on presynaptic

release and predicts future myelination, which might partially explain how neuronal activity

regulates myelination. We observed approximately 30% predication rate with 1 μm threshold

by both GFP-Gephyrin hotspots and MD Ca^{2+} in OPCs, suggesting they may act in the same

pathway to determine where myelin sheaths form. One possible scenario is that presynaptic

release from neurons leads to synaptic activation in a nearby OPC process and subsequent Ca^{2+}

activity in MDs. After the OPC "senses" this neuronal location, it could enhance the synaptic

assembly in this location through Ca^{2+} signaling. Through such a positive-feedback mechanism,

"hotspots" form, while the pre- and post-synaptic interactions influence the cytoskeleton to

transform a process into a myelin sheath. The 30% prediction rate also reflects the likelihood of

other synaptic mechanisms at play, for example, PSD-95 hotspots or neuropeptides. During OPC

differentiation, we and others observed transient myelin sheaths that disappear rapidly after

formation[34]. In the future, it will be interesting to examine if hotspots and MD Ca^{2+} predict the

formation of transient as well as the stable sheaths. What type of presynaptic release is

involved in predetermining myelination patterns? Multiple neurotransmitters and

neuropeptides can regulate the development of oligodendrocyte lineage cells[5–8,35]. Therefore,

different types of neurons across CNS regions may employ one or a combination of

neurotransmitters and/or neuropeptides to influence myelination of their axons. Here, we

focus on Gephyrin-containing synapses and predict that inhibitory neurotransmitters, such as

glycine and GABA, may be involved in regulating the dynamic assembly and enrichment at

hotspots of Gephyrin. Meanwhile, how the presynaptic release sites apposed to OPC

postsynaptic sites are distributed in neurons is still not fully understood. A recent study

suggests presynaptic release sites are assembled in hotspots towards which myelin sheaths

grow[36]. Together with our data, this suggests that both pre- and post-synaptic sites of neuron-

OPC synapses are dynamically assembled at "hotspots" that predetermine where future myelin

sheaths form on axons.

Dynamic synapses lacking certain pre- or post-synaptic molecules such as PSD-95 and

Gephyrin puncta in OPCs have been reported in developing neurons and are considered to be

immature[37–40]. Therefore, many synaptic puncta in OPCs likely do not form functional and/or

mature synapses. These synapse features are in tune with an OPC's dynamic cell biological

properties in which proliferation, migration, and process remodeling must be coordinated

before myelination. Interestingly, transcripts and/or proteins of synaptic genes are detected

even after OPCs differentiate and myelinate axons, although often at reduced levels[17,19,21,22]. It

is possible that the synaptic machinery and mechanisms we identify in OPCs also play a role in

sheath elongation and shortening. The dynamically assembled postsynaptic components in

OPCs cover a rather large space compared to the volume of an OPC alone, allowing these cells

to effectively survey their environments. Our observation that synaptic puncta numbers reduce

upon OPC differentiation provides a structural basis for the previously reported reduced

synaptic functions during this transition[27,28]. These data, together our observations that

synaptic puncta in OPCs increase during early development, suggest that synaptic assembly in

OPCs is finely regulated by development.

Several studies have identified synaptic mechanisms underlying Ca^{2+} elevation in OPCs

under conditions of increased neuronal activity (*e.g.*, through axon stimulation) with a focus on

somal activity[14–16,41]. We find that OPC Ca^{2+} activity *in vivo* mainly occurs in MDs and requires

spontaneous synaptic release. Neuronal activity and evoked synaptic release play a role in OPC

Ca^{2+} activity at 3 dpf, but to our surprise not at 5 dpf. Considering that OPC differentiation

initiates at 3 dpf, this suggests that OPCs are sensitive to neuronal activity through Ca^{2+}

signaling at this critical stage of development. This sensitivity may be important for OPC

differentiation and myelination. In our Ca^{2+} assays, we observe "outlier" OPCs that behave

differently from the majority of OPCs. For example, some OPCs exhibit no change or even an

increase of Ca^{2+} activity upon TeNT treatment while most OPCs exhibit reduction; some OPCs

exhibit reduced Ca^{2+} activity while others exhibit no change of or increased Ca^{2+} activity upon

TTX treatment at 5 dpf. These "outlier" OPCs and activity could be due to Ca^{2+} generated from

synapse-independent mechanisms, such as ER-mediated Ca^{2+} release. In line with this, nearly 61%

of Ca^{2+} MDs do not associate with presynaptic markers, and nearly 55% residual Ca^{2+} activity

remains after TeNT treatment, suggesting synapse-independent Ca^{2+}-inducing mechanisms in

OPCs. Overall, we find that OPC Ca^{2+} activity *in vivo* is complex and might be mediated by

different mechanisms than OPCs in *ex vivo* assays. While previous studies used axonal

stimulation to study neuronal activity-induced OPC Ca^{2+} response, our study of *in vivo* OPC Ca^{2+}

activity reveals that neuronal activity does not contribute to Ca^{2+} activity at certain

developmental stages. This suggests that axon stimulation may drive extrinsic Ca^{2+} activity that

otherwise would not naturally occur, cautioning the usage of axon stimulation to study OPC

Ca^{2+} activity.

Nlgn3's function is best understood at synapses. Our finding that *nlgn3b* knockdown in

oligodendrocyte lineage cells impairs OPC differentiation and myelination suggests a potential

role of synapses in OPCs. This is consistent with our observations that GFP-Gephyrin hotspots

and Ca^{2+} MDs in OPCs predict where myelin sheaths form in oligodendrocytes. Given Nlgn3's

broad presence and function at both Glutamatergic, GABAergic and Glycinergic synapses, we

expected that this approach has a larger impact on OPC activity and influences more OPCs than

disruption of single synaptic receptors. In line with our findings, an *in vitro* study found that

incubating neuron-glia co-cultures with extracellular domains of Nlgn3 inhibits OPC

differentiation and axon wrapping[42]. However, we can't rule out a non-synaptic function of

Nlgn3 that could contribute to these phenotypes. In astrocytes, *Nlgn3*'s knockdown reduced

astrocyte neuropil infiltration volumes[43]. Because astrocytes do not form synapses with

neurons, this study suggests a non-synaptic role of Nlgn3 in astrocytes. Whether Nlgn3 plays a

non-synaptic role in OPCs and oligodendrocytes remains to be understood.

The signaling pathways downstream of synapse/Ca^{2+} MDs in OPCs also remain to be

elucidated. Local Ca^{2+} activity might alter the cytoskeleton via Ca^{2+}-dependent kinases [44] or

affect lipid raft domains to change local signaling[41]. Notably, focal adhesion kinases (FAKs) can

affect cell motility and interactions with other cells and are tightly connected to Ca^{2+} activity[45].

FAKs promotes glioma growth downstream of Nlgn3 in response to neuronal activity[3,46], and we

find that that *nlgn3* knockdown leads to impaired OPC differentiation and oligodendrocyte

myelination. Therefore, Nlgn3 and FAKs may act downstream of synapses and Ca^{2+} signaling to

regulate OPC behavior during development as well as glioma growth. Overall, our *in vivo* study

establishes a relationship between neuronal activity, synaptic assembly, and Ca^{2+} activity at

neuron-OPC synapses, and reveals their roles in predicting where myelin sheaths form in

oligodendrocytes. This model inspires future directions to further investigate the pre- and post-

synaptic mechanisms underlying neuron-OPC interactions that regulate OPC development and

shape myelination.

ACKNOWLEDGMENTS

This work was supported by NMSS postdoctoral fellowship FG-1907-34613 (JL), Warren Alpert

Distinguished Scholar Award (JL), and NIH/NINDS grant 1R21NS120650 (KRM). We thank

Dwight Bergles, Anusha Mishra, Alex Nechiporuk, and members of the Monk lab for discussions

and critical reading of the manuscript. We are indebted to Austin Forbes, Tia Perry, and Grace

Halsell-Vore for fish care. We thank Rafael Almeida, Paul Brehm, Marc Freeman, Henrique von

Gersdorff, Eric Gouaux, David Lyons, Lei Ma, and Teresa Nicolson for reagents and suggestions

as well as Fernanda Coelho, Stefanie Kaech Petrie, and the OHSU microscopy core staff for

feedback and assistance in imaging. We thank Lori Vaskalis for graphic design.

AUTHOR CONTRIBUTIONS

410 J.L. and K.R.M. conceived the project. J.L. carried out experiments and data analyses. T.M.

411 generated constructs for *nlgn3a* and *nlg3b* disruption. T.C. provided key resources prior to their

412 publication. J.L. and K.R.M wrote the manuscript, and all authors edited the manuscript.

DECLARATION OF INTERESTS

415 The authors declare no competing interests.

FIGURE TITLES AND LEGENDS

Fig. 1. Postsynaptic organizers MAGUK/PSD-95 and Gephyrin are present in zebrafish spinal cord OPCs
(a) Top, schematic fish model showing transverse section orientation; bottom, distribution of soma and axons & synapses in spinal cord transverse sections.
(b-c) Representative IHC images of (b) MAGUK and (c) Gephyrin in *Tg(olig1:Kalta4,10xUAS:myrGFP)* spinal cord at 3 dpf. From left, spinal cord sections, enlarged regions, and 3D-reconstructed images. In 3D-reconstructed images, only the postsynaptic puncta that fall within OPC process volume are shown (magenta sphere).
(d) Quantification of puncta number per OPC by IHC (MAGUK and Gephyrin), FingR.GFP (PSD-95 and Gephyrin), and GFP-tagged (PSD-95 and Gephyrin) in the spinal cord. From left, n = 20, 32, 18, 24, 15, and 14 cells from at least 10 fish; F[69] = 3.937, F[52] = 1.381.
(e) Representative images and enlarged regions of FingR.GFP.PSD-95 and FingR.GFP.Gephyrin driven by *olig1* promoter in OPCs. OPC somas are indicated with arrows.
All data are represented as mean ± SEM; (d) one-way ANOVA followed by Tukey's test; scale bars 5 μm.

Fig. 2. Not all MAGUK and Gephyrin puncta in OPCs are associated with other synaptic components
(a-f) Representative transverse IHC images of (a-c) MAGUK and (d-f) Gephyrin with Synapsin in *Tg(olig1:Kalta4,10xUAS:myrGFP)* spinal cord OPCs at 3 dpf. (a and d) Spinal cord sections; (b, c, e, f) enlarged regions with 3D reconstruction (left) and Synapsin alignment (right). The postsynaptic puncta that fall within OPC process volumes are shown in 3D-reconstructed images (green sphere) and are indicated with arrowheads in Synapsin alignment images. Postsynaptic puncta either align (b and e) with Synapsin or do not (c and f).
(g) The % of MAGUK and gephyrin puncta in an OPC that align with presynaptic Synapsin in different CNS regions at 3 dpf. From left, n= 32, 57, 38, 33, 16 and 39 cells from at least 10 fish; F[214] = 3.5240.
(h) Representative IHC images of Gephyrin and Glycine receptors (GlyR) in *Tg(olig1:Kalta4,10xUAS:myrGFP)* spinal cord OPCs at 3 dpf. OPC boundaries are indicated in green. Gephyrin puncta either align with GlyR (green arrowheads) or do not (magenta arrowheads).
(i) The % of Gephyrin puncta in an OPC that align with GlyR in spinal cord at 3 dpf. n = 12 cells from 7 fish.
All data are represented as mean ± SEM; N.S., not significant; (g) one-way ANOVA followed by Tukey's test; Scale bars 5 μm.

Fig. 3. *In vivo* imaging reveals downregulation of PSD-95-GFP and GFP-Gephyrin upon OPC differentiation

(a-b) Representative images of PSD-95-GFP and GFP-Gephyrin (magenta) in spinal cord (a) OPCs and (b) neurons at 3 dpf. Cell membranes indicated with cyan.

(c-d) Quantification of (c) PSD-95-GFP and (d) GFP-Gephyrin puncta number in OPCs before and after the last process disappearance, set as time zero, during differentiation. Left, individual traces (grey lines) with the average (black line) and the standard deviation (shade). Right, paired comparison before and after time zero. (c) n= 12 cells from 8 fish; t[11] = 3.922. (d) n= 12 cells from 8 fish; t[11] = 2.705.

(e) Representative images of PSD-95-GFP and GFP-Gephyrin (magenta) in differentiating oligodendrocytes (cyan) at 2-3 dpf. Nascent sheaths are indicated with black arrowheads, and processes are indicated with magenta arrowheads.

All data are represented as mean ± SEM; N.S., not significant; (c-d) ratio-paired *t*-test; scale bars 5 μm.

Fig. 4. PSD-95-GFP and GFP-Gephyrin puncta in OPCs are dynamic and repeatedly assemble at hotspots

(a) Normalized puncta number per cell of PSD-95-GFP and GFP-Gephyrin in neurons and OPCs within 30 minutes. Black line, average; shaded area, standard deviation. From left, n = 12, 3, 4, and 6 cells from 6, 10, 4, and 4 fish.

(b-c) Quantification of PSD-95-GFP (green) and GFP-Gephyrin (magenta) puncta duration in (b) log scale and (c) % of stable puncta that lasts over 10,000 seconds in neurons and OPCs at 3 dpf. (b) n = 13 and 3 cells from 6 and 3 fish; t[14] = 5.152 and t[9] = 4.82. (c) n = 5 and 7 cells from 4 and 4 fish; t[14] = 9.989 and t[10] = 4.159.

(d) Convex hull of GFP-Gephyrin puncta in an OPC over 7 hours. The OPC membrane in the beginning of imaging is shown in magenta.

(e) Quantification of convex hull volume of GFP-Gephyrin from (d), OPC volume and volume of GFP-Gephyrin hotspot space in an OPC. From left, n= 11, 23, and 18 cells from at least 10 fish; t[39] = 6.561 and t[32] = 8.480.

(f-g) Representative time-lapse images of (f) PSD-95-GFP and (g) GFP-Gephyrin within OPC processes. The numbers indicate the corresponding puncta in h and i. Puncta #1 and #3 are in stable processes; puncta #2 and #4 are in dynamic processes. The dashed lines outline the process, and arrowheads indicate puncta appearance. Time unit is in minutes.

(h-i) Heatmaps of puncta appearance frequency over time for (h) PSD-95-GFP and (i) GFP-Gephyrin. 30% of max frequency is used as coloring threshold: below threshold, grey; above threshold, a shade of magenta corresponding to the frequency. Loci with high puncta appearance frequency (> 3 standard deviations above average) are marked with arrowheads. The numbers with green arrowheads indicate puncta shown in time lapse images in Fig. 4f-g. Bottom left, images of OPCs with puncta marked with magenta circles.

(j) Quantification of loci with high puncta appearance frequency in OPCs. From left, n = 3 and 7 cells from 3 and 5 fish.

All data are represented as mean ± SEM; (b, c, and e) unpaired *t*-test; Scale bars, 5 μm.

Fig. 5. Gephyrin hotspots predict future myelin sheath formation in a synaptic release-mediated manner

(a-b) Representative images of GFP-Gephyrin (green) in an oligodendrocyte lineage cell (magenta) at 2 different stages: (a) OPC stage that is 8 hours prior to myelination and (b) oligodendrocyte stage that is 6 hours after myelination begins.

(c) Schematic showing the imaging protocol of individual OPCs that differentiate into myelinating oligodendrocytes. The first sheath appearance is designated as time 0. Images from up to 10 hours to 1 hour before time 0 are used to identify OPC hotspots, and images around 6 hours after time 0 are used to define sheath regions. TeNT is delivered through ventricle injection before imaging.

(d) Single-plane merged image of Gephyrin puncta frequency heatmap at OPC stage and sheath regions
at oligodendrocyte stage. Hotspots (magenta), hotspot centers (green), and sheaths (yellow) are
indicated. Hotspots that existed within where myelin sheaths form are indicated with arrowheads.
(e) Quantification of hotspot-sheath and nonhotspot-sheath distances. n = 8 cells from 7 fish.
(f) Percentage of hotspots or nonhotspots that predict where myelin sheaths form within 1 μm in an
OPC. n = 8 cells from 7 fish.
(g) Frequency distribution of hotspot-sheath distances in control- and TeNT-treated OPCs. F[13, 168] =
11.39.
(h) Quantification of hotspot-sheath and nonhotspot-sheath distances with TeNT treatment. n = 6 cells
from 6 fish.
(i) Percentage of hotspots or nonhotspots that predict where myelin sheaths form within 1 μm in an
OPC with TeNT treatment. n = 6 cells from 6 fish.
(e,f,h,i) Wilcoxon matched-pairs signed rank test; (g) Two-way ANOVA and Fisher's LSD test; scale bars 5
μm.
517
Fig. 6. OPC Ca^{2+} activity is associated with synapses and predicts myelin sheath formation
(a) Representative images of plasma membrane-targeted myrGCaMP6s in an OPC (grey) showing activity
in MDs (cyan).
(b) Individual traces plotting myrGCaMP6s intensity ($\Delta F/F_{average}$) in 5 MDs from Fig. 6A. Scale bar, 100%
and 50s.
(c) Left, the occurrences of Ca^{2+} activity in MDs vs. soma; right, the number of OPCs with only MD
activity vs. OPCs with both MD and somal activity. n = 180 cells from 107 fish.
(d) Representative images of a Ca^{2+} MD (cyan) in an OPC and presynaptic synaptophysin-tagRFP
(magenta).
(e) 3-D reconstruction from (d) showing synaptophysin-tagRFP puncta (magenta) that are within 0.1 μm
from the OPC surface (arrowheads).
(f) The percentage of Ca^{2+} MDs that are associated with synaptophysin-tagRFP. n = 7 cells from 7 fish.
(g) Two Ca^{2+} spikes initiate as puncta and then spread to form MDs, indicated by blue arrowheads. The
time is in seconds.
(h) The two MDs from (g) initiate in proximity (within 0.1 μm) to presynaptic synaptophysin-tagRFP
(magenta), indicated by white arrowheads.
(i) Ca^{2+} imaging protocol of individual OPCs that differentiate into oligodendrocytes. The first sheath
appearance is designated as time 0. Images up to 6 hours before time 0 are used to identify OPC Ca^{2+}
events in MDs, and images around 6 hours after time 0 are used to define sheath regions.
(j) Merged single-plane image of Ca^{2+} MDs at OPC stage and sheath regions at oligodendrocyte stage.
Ca^{2+} MDs (blue) and sheaths (yellow) are indicated.
(k) Quantification of MD-sheath and nonMD-sheath distances. n = 9 cells from 7 fish.
(l) Percentage of MDs or nonMDs that predict where myelin sheaths form within 1 μm in an OPC. n = 9
cells from 7 fish.
(m) Schematic showing GFP-Gephyrin hotspots and MDs in OPCs similarly predict where myelin sheaths
form in oligodendrocytes.
(k and l) Wilcoxon matched-pairs signed rank test; scale bars 5 μm.
545
Fig. 7. Spontaneous synaptic release contributes to generating OPC Ca^{2+} activity *in vivo*
(a) Schematic showing OPCs neighboring few or many TeNTlc-expressing axons in our sparse-labeling
approach.
(b) Representative images of OPCs neighboring few (<4) and many (≥4) axons expressing TeNTlc-
tdtomato.

(c) The frequency of Ca^{2+} activity and peak amplitude in OPCs that neighbor few (<4) and many ($\geq$4)
TeNTlc-expressing axons at 5 dpf. Few, n = 14 cells from 8 fish; many, n = 26 cells from 15 fish; t[37] =
2.420 and t[38] = 2.958.
(d) Experimental paradigm of reagent delivery and Ca^{2+} imaging.
(e) Representative single-plane images of individual spinal cord OPCs immediately and 10 min after
ventricle injection of Dextran[3000] Texas Red (magenta) in *Tg(olig1:Kalta4,10xUAS:myrGFP)* animals.
(f) Representative images of myrGCaMP6s (grey) and Ca^{2+} MDs (cyan) in control and TeNT treatment
condition.
(g) Ca^{2+} MD number and frequency of Ca^{2+} activity per MD in OPCs before (blue) and after (magenta)
injection of control or TeNT solution at 5 dpf. Paired data are indicated with grey lines. Control, n = 17
from 13 fish; TeNT, n = 26 cells from 18 fish; from left, t[16] = 1.303, t[512] = 0.5899, t[25] = 6.189 and
t[959] = 8.853.
(h) Ca^{2+} MD number and frequency of Ca^{2+} activity per MDs in OPCs before (blue) and after (magenta)
injection of control or TTX solution at 3 dpf and 5 dpf. Paired data are indicated with grey lines. From left,
n = 15 ,31, 13, and 17 cells from 7, 13, 6, and 9 fish; t[14] = 1.000, t[30] = 2.2010, t[12] = 0.1125, t[16] =
0.8552, t[66] = 0.6158, t[163] = 2.3440, t[81] = 0.5877 and t[89] = 0.5379.
All data are represented as mean ± SEM; N.S., not significant; (c) unpaired *t*-test or (g-h) paired *t*-test;
scale bars 5 μm.
569
Fig. 8. Disruption of *nlgn3b* in OPCs impairs development and myelination.
(a) Morphology of oligodendrocytes with control- (ctrl) and *nlgn3b*-targeted for cell-specific knockdown
at 5 dpf. Top, 3D-reconstructed single oligodendrocyte with sheath (colored) and non-sheath (gray).
Bottom, single plane with sheath (black arrowheads) and processes (magenta arrowheads).
(b) Representative images of myelin sheaths from oligodendrocytes with control- (ctrl) and *nlgn3b*-
targeted for cell-specific knockdown at 3 and 5 dpf.
(c-d) Quantification of myelin sheath average lengths, number, and total length per oligodendrocyte
from (b) at 3 dpf (c) and 5 dpf (d) in ctrl-, *nlgn3a*-, and *nlgn3b*-targeted fish. (c) n = 121, 30, and 170 cells
from 78, 26, and 97 fish. (d) n = 118, 80, and 152 cells from 80, 61, and 93 fish.
(e) *In vivo* model of neuron-OPC synapses: they are composed of MAGUK and Gephyrin and contribute
to OPC Ca^{2+} activity through spontaneous release; they exhibit immature features: lack of synaptic
components and fast assembly, which enable OPCs to survey large spatial volumes; synaptic assembly
hotspots and associated Ca^{2+} activity at the OPC stage predict where a subset of myelin sheaths form at
the oligodendrocyte stage; synaptic assembly is regulated through development while disruption of the
synaptic molecule *nlgn3b* impairs OPC development and myelination.
All data are represented as mean ± SEM; N.S., not significant; (c-d) Kruskal-Wallis test followed by
Dunn's multiple comparisons test; scale bar 5 μm.
587

EXTENDED DATA FIGURE TITLES AND LEGENDS

Extended Data Fig. 1. MAGUK/PSD-95, Gephyrin, and Nlgn3 are present in OPCs, related to Fig. 1
(a) RNA-seq expression of murine *Dlg4* (*PSD-95*), *Gephyrin*, and *Nlgn3* in different CNS cell types[19].
Microglia and Macrophage (Micro/Macro); Newly formed oligodendrocyte (NFO); Myelinating
oligodendrocyte (MOL).

(b) Single cell RNA-seq expression of zebrafish *dlg4* (*a/b*), *gephyrin* (*a/b*), and *nlgn3* (*a/b*)[17]. Vascular
lepotomeningeal cell (VLMC); Myelinating oligodendrocyte (MOL).
(c) Representative IHC images of PSD-95 in *Tg(olig1:Kalta4,10xUAS:myrGFP)* spinal cord at 3 dpf. From
left, spinal cord sections, enlarged regions, and 3D-reconstructed images to reveal OPC surface and
postsynaptic puncta within OPC processes (magenta sphere).
(d) Schematic fish model showing sagittal planes.
(E-F) Two representative single-sagittal-plane IHC images showing (e) MAGUK and (f) Gephyrin within
OPCs in *Tg(sox10:GFP-caax)* spinal cord at 3 dpf. OPCs identified by GFP and shown in cyan. Arrowheads
indicate synaptic puncta. The colors are inverted to optimize visualization.
(g) % of puncta localized to OPC soma and process. MAGUK, n = 18 cells from 8 fish; Gephyrin, n = 40
cells from 15 fish.
(h) IHC images of Gephyrin in the 5 dpf spinal cord in wild-type and F0 larvae with *gephyrin* knocked
down by sgRNA targeting *gephyrina*[ex2] and *gephyrinb*[ex7] and Cas9 protein.
(i) Quantification of Gephyrin intensity in (h). n = 5 and 9 fish for each group; t[12] = 7.322.
All data are represented as mean ± SEM; (i) unpaired *t*-test; scale bars 5 μm.
609
Extended Data Fig. 2. MAGUK and Gephyrin puncta numbers in OPCs increase during early
development and exhibit small differences across CNS regions, related to Fig. 1
(a) Quantification of MAGUK (green) and Gephyrin (magenta) puncta number per cell in the spinal cord
at different ages by IHC. From left, n = 10, 8, 8, 8, 7, 8, 8, and 7 cells from at least 6 fish; t[33] = 5.730
and t[30] = 7.040.
(b) Quantification of MAGUK (green) and Gephyrin (magenta) puncta number per OPC at 3 dpf across
different CNS regions. From left, n = 34, 59, 43, 34, 18, and 40 cells from at least 10 fish; F[227] = 13.10.
(c) Schematic model showing the neuron-rich (NR) and axon & synapse-rich (AR) regions and two
representative images showing OPCs with soma localized to NR (top) and AR (bottom) regions in
Tg(olig1:Kalta4,10xUAS:myrGFP) spinal cord.
(d) Quantification of MAGUK (green) and Gephyrin (magenta) puncta number per cell in OPCs localized
to NR or AR regions at 2-3 dpf. From left, n = 14, 5, 15 and 10 cells from at least 5 fish; t[17] = 0.3372 and
t[23] = 0.8770.
All data are represented as mean ± SEM; N.S., not significant; (a-b) one-way ANOVA followed by Tukey's
test and (d) unpaired *t*-test; scale bar 5 μm.
625
Extended Data Fig. 3. More MAGUK and Gephyrin puncta in OPCs are not associated with other
synaptic molecules compared to neurons, related to Fig. 2
(a-b) Representative single-sagittal-plane IHC images of (a) MAGUK and (b) Gephyrin with presynaptic
marker Synapsin in *Tg(sox10:GFP-caax)* spinal cord at 3 dpf. From left, merged channels, MAGUK or
Gephyrin channel, and Synapsin channel. Black arrowheads indicate the postsynaptic puncta in OPCs
that align with presynaptic Synapsin. Magenta arrowheads indicate the postsynaptic puncta in OPCs that
do not align with presynaptic Synapsin.
(c) Colocalization ratio of MAGUK (green) or Gephyrin (magenta) in OPCs with Synapsin, normalized to
neurons at 2 dpf and 14 dpf. From left, n = 9, 7, 9, and 7 OPCs from at least 5 fish; t[8] = 0.1426, t[6] =
3.6380, t[8] = 3.2880, t[6] = 2.5400.
(d) Colocalization ratio of Gephyrin with GlyR in OPCs, normalized to neurons at 3 dpf. n = 11 cells from
7 fish; t[10] = 7.049.
All data are represented as mean ± SEM; N.S., not significant; (c-d) paired *t*-test; Scale bars 5 μm.
639
640 **Extended Data Fig. 4. *In vivo* characterization of PSD-95-GFP and GFP-gephyrin, related to Fig. 3**

(a) Puncta diameter of PSD-95-GFP (green) and GFP-Gephyrin (magenta) in neurons and OPCs at 2-3 dpf in the spinal cord. From left, n = 10, 11, 15, and 29 cells from at least 8 fish; t[19] = 1.8200 and t[42] = 0.2447.

(b) Representative images of PSD-95-GFP or GFP-Gephyrin (green) and synaptophysin-TagRFP (magenta) in spinal cord OPCs at 3 dpf. From left, single plane images, enlarged regions, and Imaris "Spot"-processing to reveal pre- (magenta sphere) and post- (green sphere) synaptic puncta.

(c) The % of PSD-95-GFP and GFP-gephyrin puncta in an OPC that align with presynaptic synaptophysin-TagRFP in the spinal cord in (b). n = 13 and 20 cells from at least 10 fish.

(d) Quantification of PSD-95-GFP (green) and GFP-Gephyrin (magenta) puncta number per cell in the spinal cord at 2 dpf and 3 dpf. From left, n = 16, 18, 29, and 14 cells from at least 5 fish; t[32] = 5.238 and t[41] = 7.147.

All data are represented as mean ± SEM; (a, d) unpaired *t*-test. Scale bars 5 μm.

Extended Data Fig. 5. Gephyrin hotspots predict myelin sheath formation in a synaptic release-mediated manner, related to Fig. 5

(a-c) Quantification of (a) convex hull volume, (b) hotspot number, and (c) sheath number in control and TeNT treatment conditions. (a) n = 11 and 6 cells from 10 and 6 fish; (b) n = 18 and 6 cells from 16 and 6 fish; (c) n = 11 and 6 cells from 10 and 6 fish.

All data are represented as mean ± SEM; (a-c) Mann-Whitney test.

Extended Data Fig. 6. The role of synaptic release and neuronal activity in generating OPC Ca^{2+} activity, related to Fig. 7

(a) Basal GCaMP6s intensity and MD feret's diameter in OPCs that neighbor few (< 4) and many (≥ 4) TeTxLc-expressing axons at 5 dpf. Few, n = 14 cells from 8 fish; many, n= 26 cells from 15 fish; t[38] = 2.375 and t[38] = 2.351.

(b) Representative image of individual spinal cord OPCs 50 min after ventricle injection of Dextran^{150,000} Antonia Red (magenta) in *Tg(olig1:Kalta4,10xUAS:myrGcamp6s)* fish.

(c) Quantification of peak amplitude, peak duration, area of Ca^{2+} MD, frequency of Ca^{2+} activity per cell, basal intensity, and MD Feret's diameter in OPCs before (blue) and after (magenta) injection of control or TeNT solution at 5 dpf. Paired data are indicated with grey lines. Control, n = 17 cells from 13 fish; TeNT, n = 26 cells from 18 fish; from left, t[16] = 0.4575, t[25] = 1.158, t[16] = 0.3358, t[25] = 2.312, t[16] = 1.159, t[25] = 1.872, t[16] = 0.5831, t[25] = 5.077, t[16] = 1.039, t[27] = 4.028, t[16] = 1.434, and t[29] = 1.421.

(d) Quantification of Ca^{2+} MD number and frequency of Ca^{2+} activity per MD in OPCs before (blue) and after (magenta) injection of control or TTX solution at 3 dpf and 5 dpf. Paired data are indicated with grey lines. From left, n = 15 ,31, 13, and 17 cells from 7, 13, 6, and 9 fish; t[9] = 1.3490, t[23] = 2.2340, t[9] = 0.7957, t[10] = 1.0540, t[9] = 0.4645, t[23] = 0.8590, t[9] = 0.8009, t[10] = 0.2402, t[9] = 0.4605, t[23] = 0.7737, t[9] = 0.2900 and t[10] = 0.2361.

(e) Representative images of a primary motor neuron visualized by injecting *mnx1:gal4* into *Tg(10xUAS:myrGCaMP6s).*

(f) Frequency of Ca^{2+} activity per cell before (blue) and after (magenta) injection of control or TTX solution in motor neurons at 5 dpf. n = 7 and 6 cells from 5 and 5 fish; t[6] = 1.6220 and t[5] = 3.7970.

All data are represented as mean ± SEM; N.S., not significant; (a) unpaired *t*-test or (c-d, f) paired *t*-test; scale bars 5 μm.

Extended Data Fig. 7. Disruption of *nlgn3b* in OPCs impairs OPC development and myelination, related to Fig. 8

(a) Schematic of Tol2 plasmid (10xUAS:myrmScarletI-p2A-Cas9, U6:sgRNA1;U6:sgRNA2) constructed and used to induce cell-specific Cas9 expression with membrane labeling and 2 separate U6-driven sgRNAs.

(b) Representative gel images of digested PCR product of genomic regions targeted by the sgRNAs. Each lane represents a single embryo at 1 dpf. The left 4 lanes are uninjected (-) and the right 4 lanes are injected (+) with sgRNAs and Cas9 protein.

(c) Schematic of larva with sparsely labeled oligodendrocyte lineage cells resulting from *Tg(sox10:Kalta4)* crossed with transgenic fish carrying the plasmid from (a).

(d) Representative images of single oligodendrocytes in the spinal cord of *Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:ctrl-sgRNA1;U6:ctrl-sgRNA2)* and *Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:myrf^{ex3}-sgRNA;U6:myrf^{ex4}-sgRNA)* at 6 dpf. Knockdown of Myrf serves as a control for our targeting approach.

(e) Quantification of myelin sheath length in (d). n = 24 and 22 cells from 15 fish; t[44] = 2.247.

(f) Representative images of an OPC and an oligodendrocyte.

(g) Percentage of oligodendrocytes (in contrast to OPCs) in ctrl-, *nlgn3a*-, and *nlgn3b*-targeted fish at 3 dpf and 5 dpf. 3 dpf, n = 144, 82, and 152 cells from 78, 26, and 97 fish, F[6] = 15.90; 5 dpf, n = 119, 84, and 170 cells from 80, 61, and 93 fish, F[6] = 7.343.

(h) Percentage of oligodendrocytes that possess both myelin sheath and abnormal processes in ctrl-, *nlgn3a*-, and *nlgn3b*-targeted fish at 5 dpf. n = 119, 84, and 170 cells from 80, 61, and 93 fish, F[6] = 13.52.

(i) 4 single planes of a trapezoid-shaped process in an oligodendrocyte in *nlgn3b*-targeted larvae at 5 dpf. Myelin sheaths are indicated by black arrowheads and abnormal processes by magenta arrowheads.

All data are represented as mean ± SEM; N.S., not significant; (e) unpaired *t*-test and (g,h) one-way ANOVA followed by Tukey's test; scale bars 5 μm.

SUPPLEMENTARY VIDEOS WITH TITLES AND LEGENDS

Video 1. Time-lapse images of GFP-Gephyrin in an OPC
Representative video of sox10:myrTagRFP-T-P2A-GFP-Gephyrin in an OPC that eventually differentiates in the spinal cord of a 3 dpf larva.

Video 2. Time-lapse images of GFP-Gephyrin in a neuron
Representative video of 10xUAS:GFP-Gephyrin in a neuron in the spinal cord of a *Tg(sox10:Kalta4)* larva at 3 dpf.

Video 3. Time-lapse images of myrGCaMP6s in an OPC
Representative video of Ca^{2+} activity in an OPC in a *Tg(olig1:Kalta4,10xUAS:myrGCaMP6s)* larva at 5 dpf.

Video 4. Time-lapse images of myrGCaMP6s in an OPC with both MD and somal Ca^{2+} activity
Video of Ca^{2+} activity in an OPC with multiple events including a somal event in a *Tg(olig1:Kalta4,10xUAS:myrGCaMP6s)* larva at 5 dpf.

Video 5. Time-lapse images of myrGCaMP6s in an OPC before and after injection of TeNT at 5 dpf
Representative 10-minute video of Ca^{2+} activity in an OPC before (left) and after (right) injection of TeNT in a *Tg(olig1:Kalta4,10xUAS:myrGCaMP6s)* larva at 5 dpf.

SUPPLEMENTARY TABLES WITH TITLES AND LEGENDS

Table 1. A list of primers used in this study

733   Related to STARS Methods.

METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contacts, Jiaxing Li (lijia@ohsu.edu) or Kelly Monk (monk@ohsu.edu).

Materials availability

All reagents generated in this paper will be shared by the lead contacts upon request.

Data and code availability

All data reported in this paper will be shared by the lead contacts upon request.

All original code is available from the lead contacts upon request.

Any additional information required to reanalyze the data reported in this paper is available

from the lead contacts upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

We used the following existing zebrafish lines and strains: *Tg(sox10:Kalta4)*[47], *Tg(olig1:Kalta4)*[17],

Tg(huc:synaptophysin-tagRFP)[17], *Tg(olig2:dsRed)*, *Tg(sox10:GFP-caax)*, *nacre*, and AB. The

following lines were newly generated for this study: *Tg(10xUAS:myrGCaMP6s)*,

Tg(olig1:myrGCaMP6s), *Tg(10xUAS:myrGFP)*, *Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:ctrl-*

sgRNA1;U6:ctrl-sgRNA2), *Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:nlgn3a^{ex1}-*

sgRNA;U6:nlgn3a^{ex4}-sgRNA), *Tg(10xUAS:myrmScarletI-p2A-Cas9, U6:nlgn3b^{ex1}-*

sgRNA;U6:nlgn3b^{ex3}-sgRNA). All experimental fish were crossed into the *nacre* background for

optimal optics without pigmentation. The Kalta4/UAS system allowed sparse labeling and single

cell analysis. The genotypes of fish used for each experiment are indicated in figure legends.

Animals in the fish facility were maintained at 28° C with a 14 h/10 h light/dark cycle. Larvae

and juveniles were nurtured with rotifer suspension and dry food (Gemma 75 and 150,

respectively). Adults were fed with a combination of brine shrimp and dry food (Gemma 300).

Male and female fish of 3 to 12 months-old were mated; larval fish of 2 dpf to 14 dpf were used

for experiments (sex is not determined at these stages in zebrafish). All zebrafish experiments

and procedures were performed in compliance with institutional ethical regulations for animal

testing and research at Oregon Health and Science University (OHSU). Experiments were

approved by the Institutional Animal Care and Use Committee of OHSU. Animal and cell

numbers are indicated in figure legends.

METHOD DETAILS

Tissue sectioning and immunohistochemistry

To prepare sagittal and transverse sections, zebrafish larvae at indicated stages were first

anesthetized with tricaine and then fixed in 4% PFA, 0.1% Triton at 4° C overnight. Heads and

tails were removed to enable better penetration of fixative. The fish were then washed in PBS,

0.1% Triton for 5 minutes and passed in 10%, 20%, and 30% sucrose in PBS, followed by OCT

embedding and cryosectioning of 20-25 μm slices. The sections were brought to room

temperature for 45 minutes and rehydrated in PBS, 0.2% Triton for 1 hour. 3% BSA, 5% NGS, 0.2%

Triton in PBS was used as blocking buffer. Primary antibodies: chicken anti-GFP (Invitrogen

A10262), 1:1000, mouse anti-Pan-MAGUK (NeuroMab K28/86) 1:500, mouse anti-Gephyrin

(Synaptic Systems 147011) 1:1000, rabbit anti-Synapsin 1/2 (Synaptic Systems 106002) 1:1000,

mouse anti PSD-95 6G6-1C9 (MAB1596) 1:1000, Rabbit anti-Glycine receptor (Synaptic Systems

146008) 1:1000. Secondary antibodies: A488 anti-chicken, A594 anti-rabbit, A647 anti-mouse

IgGI, A647 anti-mouse (Thermo Fisher). The stained samples were mounted in prolong Glass

(Invitrogen P36980) for at least a day before imaging.

Plasmid construction

In zebrafish, *gephyrin* is duplicated (*gephyrina* and *gephyrinb*), and there are multiple isoforms

for both genes. All isoforms are almost identical in protein sequence except for the inclusion or

exclusion of C3 and C4 cassettes. We amplified several isoforms from cDNA and decided to

follow up two: *gephyrinb* isoform X6 (XM_021468341) because it encodes a protein that is 96%

identical and 98% similar to the rat P1 isoform (NP_074056.2)[48] that has been used to track

synapses previously; and, *gephyrina* isoform X11 (XM_017351650.2) because it contains a C3

cassette previously reported to be present in glia cells[49].

Total RNA was extracted from 3 dpf zebrafish AB larvae (50 fish) using RNAeasy Plus Kit

(Qiagen 74134). cDNA was synthesized using SuperScript IV Reverse Transcriptase (Invitrogen).

Two sets of primers (Supplementary Table 1) were used to clone the full length *gephyrina* and

gephyrinb using high fidelity Q5 DNA polymerase, followed by A-addition and TA cloning into

pCRII-TOPO backbones. Using Gibson assembly (NEB E5510S), we added EGFP to the 5' end and

a linker GGGGSGGGGSGGGGS in between, and integrated it into pCS 2.1 backbones with an

AVRII digestion site. The 5' end was chosen based on previous imaging studies[48]. In transient

injections, GFP-Gephyrina and GFP-Gephyrinb exhibited similar expression profiles; therefore,

we used GFP-Gephyrinb for our studies and refer to the GFP-3xGGGGS-Gephyrinb as GFP-

Gephyrinb throughout the manuscript. Using Gateway BP reaction (Invitrogen 11786100), we

generated pME_ATG-GFP-Gephyrinb-TAA. Using AVRII and KPNI-HF, we cut GFP-Gephyrinb

(without an ATG) from the pCS 2.1 backbone and inserted it in frame into p3E_p2A to generate

p3E_p2A-GFP-Gephyrinb-TGA-polyA.

PSD-95/dlg4b transcript NM_214728.1 encodes a protein that is 76% identical and 83%

similar to rat *Dlg4* (NP_062567.1) and was thus chosen for use in our studies. We added GFP to

the 3'end because N-terminal palmitoylation of PSD-95 is necessary for its synaptic localization,

and receptor clustering[50,51]. It was cloned from cDNA with primers (Supplementary Table 1)

similar to *gephyrin* as described above. We refer to PSD-95-3xGGGGS-GFP as PSD-95-GFP. We

generated pME_ATG-PSD-95-GFP-TAA and p3E_p2A-PSD-95-GFP-TGA-polyA constructs.

pCAG_FingR.GFP-Gephyrin-CCR5TC (Addgene 46296) and pCAG_FingR.GFP-PSD-95-

CCR5TC (Addgene 46295) were used to amplify the coding region with additional 5' zinc finger

binding sequence to regulate their expression[20], followed by BP reaction to generate

pME_zincfinger_FingR.GFP-PSD-95-CCR5TC and pME_zincfinger_FingR.GFP-Gephyrin-CCR5TC

constructs.

pME_myrGFP and pME_myrTagRFP-T were generated using PCR and BP reaction, using

Fyn myristoylation and palmitoylation domain. pGP-CMV-GCaMP6s (Addgene 40753) was used

as a template and 3 primers were used in a Q5 PCR (15 cycles at 71° C and 20 cycles at 72° C) to

amplify GCaMP6s and add a Kozak sequence, myr (Fyn myristoylation and palmitoylation

domain), attB1, and attB2 sequences. This PCR product was used to generate

pME_myrGCaMP6s-TAA using BP reaction.

We obtained multiple plasmids from Tol2 kits: pDestTol2pA2, pDONR221, p5E_10xUAS,

p3E_polyA, p3E_p2A[52], p5E_sox10$_{7.2kb}$[47], pDestTol2pACryGFP (pDestTol2GE),

pDestTol2pACrymCherry (pDestTol2RE).

Using Gateway LR reactions (Invitrogen 11791020), we generated the following

constructs: PA2-10xUAS:myrEGFP-PolyA, GE-10xUAS:myrGCaMP6s-TAA-polyA, PA2-

olig1:myrGCaMP6s-TAA-polyA, GE-10xUAS:GFP-Gephyrinb-TAA-polyA, RE-10xUAS:GFP-

Gephyrina-TAA-polyA, GE-10xUAS:PSD-95-GFP-TAA-polyA, PA2-sox10:myrTagRFP-T-P2A-GFP-

Gephyrinb-polyA, PA2-sox10:myrTagRFP-T-P2A-PSD95-GFP-polyA, GE-

10xUAS:zincfinger_FingR.GFP-PSD-95-CCR5TC, GE-10xUAS:zincfinger_FingR.GFP-Gephyrin-

CCR5TC. We used Mnx1:Gal4[53] to drive myrGCaMP6s expression in motor neurons. We used

Huc:Kalta4 [47], UAS:TeNTlc-tdtomato[32] to drive TeNTlc expression in neurons. All constructs

generated in this study were fully sequenced (except *sox10* and *olig1* promoters, which are well

established tools).

Cell specific CRISPR/Cas9 plasmid construction and CRISPR/Cas9-mediated gene disruption

We used 5xUAS:hsp70promoter-Cas9-T2A-GFP, U6:sgRNA1;U6:sgRNA2[54] (Addgene 74009) as a

backbone and performed the following modifications: replacement of the 5xUAS:hsp70

promoter with the 10xUAS:E1b-bactin-Kozak promoter because we found the original promoter

expressed in many non-specific cell types when injected into *Tg(sox10:Kalta4)*; replacement of

T2A-GFP with myrmScarletI-p2A and added to the 5'end of Cas9; addition of polyA to the 3' end

of Cas9. Four sets of primers and 2 previously made plasmids (Tol2-10xUAS:E1b-mScarletI and

Tol2-10xUAS:cas9-P2A-TagRFPTcaax) were used to generate 4 segments. These segments were

integrated into the backbone by NEBuilder HiFi assembly (NEB E2621S). The new construct is

referred to as 10xUAS:myrmScarletI-p2A-Cas9, U6:sgRNA1;U6:sgRNA2.

CHOPCHOP and Crisprscan were used to identify 2 sgRNA sequences for each gene of

interest: *gephyrina* exon 2, GGTCATGAACGAGATCTTTGAGG; *gephyrinb* exon 7,

GGAGAGCCGGCAGGATGAACTGG; *myrf* exon 3, GACCCAACGGCGGTGCACCCGGG; *myrf* exon 4,

GCTGGAGTCCGGAGGAGACTCGG; *nlgn3a* exon 1, GGTGTATGATGCTTGTCCGGGGG; *nlgn3a* exon

4, GCCTTGTGTACTTAACCGGCTGG; *nlgn3b* exon 1, GATGCGGGTCGCTGTTGCCACGG; *nlgn3b*

exon 3, GGTAGTGTGCTGGCCAGCTACGG. The cutting efficiencies of these sgRNAs were verified

(>95%) via enzyme digestion of the targeted locus PCR product (CRISPRtest primers, Fig. S3E).

F0 larvae with *gephyrina* and *gephyrinb* sgRNA injected together with 1 ng Cas9 protein were

used to examine the specificity of the Gephyrin antibody. BsaI and BsmBI were used to insert

the sgRNA into 10xUAS:myrmScarletI-p2A-Cas9, U6:sgRNA1;U6:sgRNA2. The plasmids were

injected with Tol2 transposase mRNA to generate transgenic fish. *Tg(10xUAS:myrmScarletI-*

p2A-Cas9, U6:myrf^{exon3};U6:myrf^{exon4}), Tg(10xUAS:myrmScarletI-p2A-Cas9,

U6:nlgn3a^{exon1};U6:nlgn3a^{exon4}), and Tg(10xUAS:myrmScarletI-p2A-Cas9,

U6:nlgn3b^{exon1};U6:nlgn3b^{ex34}) larvae with sparsely labeled oligodendrocyte were examined for

morphology and sheath characteristics. *Tg(10xUAS:myrmScarletI-p2A-Cas9,*

U6:sgRNA1;U6:sgRNA2) larvae were used as controls.

Sparse labeling and generation of transgenic lines

Fertilized zebrafish eggs at the 1-cell stage zebrafish zygotes were microinjected with 1 nL of

an injection solution containing 20 ng/µL plasmid DNA, 25 ng/µL Tol2 transposase mRNA, 0.02%

phenol red and 0.2M KCl. Injected F0 animals were either used for single-cell analysis or raised

to adulthood to generate transgenic lines. To generate stable lines, F1 progenies were screened

for germline transmission of the fluorescent reporters. F1 animals were further outcrossed and

maintained in the *nacre* background to establish stable transgenic lines with single copy

insertion.

Mounting larvae for *in vivo* imaging

For most live imaging experiments, larvae were anesthetized with 0.16 mg/mL (600 µM)

tricaine in embryo medium. The fish were mounted in 0.8% low melting agarose (Sigma A9414)

on a coverslip for morphology and sheath examination or in 1.5% low melting agarose in a dish

for time lapse imaging (hours). For Gephyrin hotspot and myrGCaMP6s imaging, larvae were

immobilized with 0.5 mg/ml non-depolarizing acetylcholine receptor antagonist mivacurium

chloride (Abcam) and mounted in 1.5% low melting agarose in a dish.

Confocal microscopy

All imaging was performed on an upright Zeiss LSM 980 confocal with Airyscan 2 in 4Y fast

mode. For fixed tissues, a 63x/1.4 NA oil objective was used, and for live imaging, a 20x/1.0 NA

water objective was used. All images were taken in Zeiss software. We used the following lasers:

488 nm for EGFP, myrGCaMP6s, and Alexa488; 561 nm for TagRFP, mScarletI, tdtomato, and

Alexa594; 639 nm for Alexa647. Fixed tissue was imaged with 0.15 µm z spacing with 1.6x to

2.5x zoom. For *in vivo* analyses, PSD-95-GFP and GFP-Gephyrin were imaged with 2 µm z-

spacing at an interval of 5 min (for 30 min to 1 hour) or 20 min (for longer than 1 hour) at a

zoom between 3.7x-10x depending on OPC cell size. myrGCaMP6s was imaged with 2 µm z-

spacing at 0.5 Hz (4 planes scanned) for 10 min or 20 min (drug injection experiments) as

previously described[17,18]. Images acquired at 0.5 Hz were analyzed along with images acquired

at 10 Hz with a single plane. No major differences were found on the Ca^{2+} spike characteristics,

but 0.5 Hz acquisition allowed inclusion of more stacks and increased the number of events

detected. Therefore, 0.5 Hz was used throughout the study. For tracking OPCs that differentiate

into myelinating oligodendrocyte, Ca^{2+} imaging was done repeatedly for 5 min at the beginning

of each hour.

Drug delivery through ventricle injection

For myrGCaMP6s experiments involving drug delivery, we injected drugs via ventricle injection

as described previously[18]. In TTX treatment, following 10 min GCaMP6s imaging, a 1 nL mixture

containing 0.2M KCL, 0.5% Dextran Texas Red 3000 MW (Thermo), and molecular biology grade

H_2O (control) or TTX (0.5 mM, kind gifts from Henrique von Gersdorff's lab) were pressure

injected into the hindbrain ventricle using an ASI MPPI-3. Between 15 and 35 min after injection,

Dextran was imaged, followed by another 10 min of GCaMP6s imaging. Dextran surrounding

the imaged OPCs was used to verify the drug delivery. TTX caused fish paralysis, further

verifying efficient delivery of active drugs. After imaging, fish were removed from mounting

media. Larvae injected with control solution recovered full swimming activity within 3 minutes,

and TTX-injected paralyzed larvae maintained normal heartbeats for at least 3 hours. In TeNT

treatment, an 8 nL mixture containing 0.2M KCL, 0.5% Dextran Antonia Red 150,000 MW

(Sigma), and TeNT (listlabs 190A, 0.5 µM) were pressure injected into the hindbrain ventricle.

Between 50 and 70 min after injection, Dextran was imaged, followed by another 10 min of

GCaMP6s imaging. TeNT-injected larvae showed reduced swimming activity.

QUANTIFICATION AND STATISTICAL ANLYSIS

OPC labeling during development

We used *sox10* and *olig1* to label OPCs. *sox10* drives expression in both OPCs and myelinating oligodendrocytes while *olig1* labels only OPCs. OPCs and myelinating oligodendrocytes could be distinguished by cell morphology: OPCs only possessed processes and myelinating oligodendrocytes possessed sheaths. Animal ages and corresponding OPC activity: 2 dpf (when OPCs are born, and migrate), 3 dpf (when many OPCs initiate differentiation), 5-14 dpf (when many OPCs complete differentiation.)

Alignment rate and colocalization ratio

Images of fixed tissue were analyzed in Imaris. "Surface" function was used to create a cell surface (Smooth and Surface Grain Size = 0.0850 μm; Diameter of Largest Sphere = 5 μm; Threshold between 900-1200; Number of Voxels above 100). "Spot" function was used to generate $Spot_{postsynaptic\ puncta}$ (Estimated XY Diameter = 0.35 μm; Estimated Z Diameter = 0.5 μm; Background subtraction; quality above 2500 to 2800). Then the cell surface was used as a mask to generate $Spot_{postsynaptic\ puncta\ in\ OPCs}$, which was further verified by crosschecking its overlap with $Spot_{postsynaptic\ puncta}$. $Spot_{postsynaptic\ puncta\ in\ OPCs}$ was analyzed for puncta number or density. To examine the alignment of MAGUK (Gephyrin) and Synapsin, we used "Spot" function to generate $Spot_{Synapsin\ puncta}$. By determining the proximity $Spot_{postsynaptic\ puncta\ in\ OPCs}$ to $Spot_{Synapsin\ puncta}$ using MATLAB-enabled "Colocalize Spots" function (<0.1 μm), we could identify postsynaptic puncta in OPCs that aligned with Synapsin and calculate the alignment rate. To calculate the colocalization ratio of MAGUK (Gephyrin)-Synapsin, we used the Imaris "Coloc" function to generate a new M&S coloc channel based on the overlapping pixels between MAGUK (Gephyrin) and Synapsin channel above thresholds. "Surface" function was used to

generate M&S coloc surface of the whole section, MAGUK (Gephyrin) (M) surface of the whole

section, and OPC surface. OPC surface was used as a mask to generate masked M&S coloc

channel and MAGUK (Gephyrin) channel. These masked channels were then used to generate

M&S coloc surface In OPCs and M surface in OPCs." The colocalization ratio in OPCs and

neurons was calculated as follows:

$$Colocalization\ ratio_{OPCs} = \frac{Volume_{M\&S\ coloc\ surface\ in\ OPCs}}{Volume_{M\ surface\ in\ OPCs}};$$

$$Colocalization\ ratio_{neurons} =$$

$$\frac{Volume_{M\&S\ coloc\ surface\ of\ the\ whole\ section} - Volume_{M\&S\ coloc\ surface\ in\ OPCs}}{Volume_{M\ surface\ of\ the\ whole\ section} - Volume_{M\ Surface\ in\ OPCs}}.$$

Postsynaptic puncta identification and analysis

Live Images of PSD-95-GFP and GFP-Gephyrin were processed in ImageJ, and synaptic puncta

were identified and tracked with Trackmate[55]. The images were first maximally projected and

stabilized using "Image Stabilizer." In Trackmate, parameters were set as follows: LOG mode,

diameter 0.75 µm, Quality>100, contrast>0.19 and signal noise >0.35, LAP linking with 0.5 µm

distance and 2 max gap frames allowed were applied. The identified puncta at each frame were

then manually examined to remove poor quality tracks. Neighboring puncta may sometimes be

incorrectly linked between timelapses; therefore, manual examinations were done to fix these

link errors. The data output was analyzed in MATLAB with custom scripts to calculate puncta

number per cell at each timepoint, puncta duration and diameter. To generate Fig.s of mean

and standard deviation as shade, we used a modified script

(https://www.mathworks.com/matlabcentral/fileexchange/29534-stdshade). To analyze the

puncta number during OPC differentiation, time points when OPC shows first sheath

(differentiation) and when myelinating oligodendrocyte's last process disappeared and all

process transformed into sheaths (maturation) were noted for each cell. In MATLAB, this

information is used to divide the puncta number into before and after differentiation or

maturation for each cell. The traces were generated after the data was realigned to set the

timepoint of differentiation or maturation as time 0.

Hotspot analysis

Heatmaps of puncta over time were generated using the output data with Heatscatter

(https://www.mathworks.com/matlabcentral/fileexchange/47165-heatscatter-plot-for-

variables-x-and-y). With a sampling volume of 1 x 1 x 4 µm, regions with frequency higher than

3 standard deviations were identified as hotspots using custom MATLAB scripts, which also

measured the distance between the center of hotspots and future sheaths. In time-lapse

imaging that encompassed both OPC and oligodendrocyte stages, we used the first nascent

sheath appearance (time 0) to separate OPC stage and oligodendrocyte stage. In some cases,

the first nascent sheath appeared between our observation timepoints or disappeared for half

an hour; therefore, we decided to confine the OPC stage up to one hour before time 0. The

distances between hotspots and myelin sheaths were measured using custom MATLAB scripts.

To choose 5-30 representative points outside hotspots in an OPC, we first used Trackmate to

identify 80-400 points along OPC processes (not soma) using the RFP channel. We removed the

points that are within 5 µm from the hotspot centers in the OPC and clustered the rest of points

into 5-30 groups (nonhotspots) using Lloyd's algorithm in MATLAB. Then we measured the

distances from each nonhotspot center to its closest myelin sheaths within the same OPC. To

examine how hotspots or nonhotspots predicted myelin sheaths, we analyzed the frequency

distribution of hotspot-sheath and nonhotspot-sheath distances, and calculated the percentage

of distances that were less than 1 μm.

Ca^{2+} events analysis

myrGCaMP6s images of z-stacks were maximum projected and stabilized using "Image

Stabilizer." The images were assigned with random names using "blindr"

(https://github.com/U8NWXD/blindr) before analysis and reassigned with the original names

after analysis. The regions of Ca^{2+} events (ROI) were identified by "GECIquant"[56] first and then

manually examined frame by frame. The intensity and area of each ROI were exported and

processed in MATLAB through custom scripts. Briefly, $\Delta F/F_{average}$ ($\Delta F=F- F_{average}$) was calculated

to generate traces and to identify intensity peaks using "findpeaks" in MATLAB with filtering

parameters (3 standard deviations above average, prominence larger than 40% of the baseline,

and peak area larger than 2.5). Rare elevated Ca^{2+} signal for extended periods of time (longer

than 50 seconds) were excluded. Peaks were manually verified against the image. Several Ca^{2+}

characteristics were measured: peak amplitude, peak duration, Ca^{2+} MD number, Ca^{2+} MD area,

and frequency of Ca^{2+} activity. A region with multiple OPCs was analyzed for TeNTlc and TeNT

experiment, while single OPCs were analyzed for the rest of experiments. To examine the

proximity of synaptophysin-TagRFP to Ca^{2+} MD, Ca^{2+} MDs were identified as above, and their

locations were first compared to synaptophysin-TagRFP manually plane by plane and then

verified by Imaris. In Imaris, using "Spot" and "Surface" functions, we generated synaptophysin-

TagRFP puncta and OPC surface, with which we identified the puncta that were close to OPC

surfaces using a MATLAB-enabled "Spot-to-Surface" function (0.1 μm threshold). These puncta

were manually checked against Ca^{2+} MDs identified through morphological landmarks in Imaris.

The distance between Ca^{2+} and future sheaths were measured in MATLAB. In TeNTlc

experiments, the number of axons expressing TeNTlc-tdtomato was counted in each image. If

the number was less than 4, the corresponding image was grouped as "neighboring few TeNTlc

axons"; otherwise, the image was grouped as "neighboring many TeNTlc axons". The distances

between Ca^{2+} MDs and future myelin sheaths were measured using custom MATLAB scripts. To

choose 5-30 representative points outside MDs in an OPC, we first used Trackmate to identify

80-400 points along OPC processes (not soma) using the RFP channel. We removed the points

that are within 5 μm from the MD centers in the OPC and clustered the rest of points into 5-30

groups (nonMDs) using Lloyd's algorithm in MATLAB. Then we measured the distances from

each nonMD center to its closest myelin sheaths within the same OPC. To examine how MDs or

nonMDs predicted myelin sheaths, we analyzed the frequency distribution of MD-sheath and

nonMD-sheath distances, and calculated the percentage of distances that were less than 1 μm.

1007

Oligodendrocyte and myelin sheath analysis

To analyze myelin sheath length and number, we used ImageJ ROI manager to trace all sheaths

of sparsely labeled oligodendrocytes between segments 1 and 15 in spinal cord. The number of

oligodendrocytes analyzed per fish was limited to 3 to ensure even weight among individual fish.

Sheath traces and the corresponding images were blinded using "File Name Encrypter" in

ImageJ. The sheath length and number were then analyzed in MATLAB with custom scripts and

matched with the original genotype following analysis. To calculate the oligodendrocyte %, in

each imaging experiment, the number of oligodendrocytes from one genotype was counted

and divided to the total number of cells (both oligodendrocytes and OPCs).

Statistics

Most data are shown as mean ± SEM., with individual data points shown as circles. We

conducted Shapiro-Wilk's test and Levene's test to verify data normality and homoscedasticity,

respectively, before performing two-sided *t*-test for unpaired groups, paired *t*-test for paired

groups, and one-way ANOVA and Tukey's test for more than 3 groups. If the data did not pass

normality tests, we used Kruskal-Wallis test for more than 3 groups and Mann-Whitney test for

2 groups. These analyses were performed either in GraphPad Prism or MATLAB. The

significance and statistical analyses employed were indicated in Fig. legends. Animal and cell

number used for each experiment, t-values and df for *t*-tests, and *F* values and df for ANOVA

were indicated in figure legends. Data collection was not performed blindly due to the

experimental conditions, but analyses were performed blindly as indicated. Sample size was

chosen based on standard deviation and effect (http://www.biomath.info/power/ttest.htm).

Sex is not yet determined in zebrafish at the ages employed in our study and so cannot be

considered as a biological variable.

Reporting summary

Further information on research design is available in the Nature Research Reporting Summary

linked to this manuscript.

Data availability

All research materials are available upon request. Source data are provided with this paper.

1039 **Code availability**

1040 All custom scripts are available upon request.

1041

REFERENCES

1. Franklin, R. J. M. & Ffrench-Constant, C. Remyelination in the CNS: From biology to

therapy. *Nature Reviews Neuroscience* vol. 9 839–855 (2008).

2. Xiao, Y., Petrucco, L., Hoodless, L. J., Portugues, R. & Czopka, T. Oligodendrocyte

precursor cells sculpt the visual system by regulating axonal remodeling. *Nat. Neurosci.*

25, 280–284 (2022).

3. Venkatesh, H. S. *et al.* Targeting neuronal activity-regulated neuroligin-3 dependency in

high-grade glioma. *Nature* **549**, 533–537 (2017).

4. Bergles, D. E. & Richardson, W. D. Oligodendrocyte development and plasticity. *Cold*

Spring Harb. Perspect. Biol. **8**, a020453 (2016).

5. Lundgaard, I. *et al.* Neuregulin and BDNF Induce a Switch to NMDA Receptor-Dependent

Myelination by Oligodendrocytes. *PLoS Biol.* **11**, (2013).

6. de Biase, L. M. *et al.* NMDA receptor signaling in oligodendrocyte progenitors is not

required for oligodendrogenesis and myelination. *J. Neurosci.* **31**, 12650–12662 (2011).

7. Zonouzi, M. *et al.* GABAergic regulation of cerebellar NG2 cell development is altered in

perinatal white matter injury. *Nat. Neurosci.* **18**, 674–682 (2015).

8. Balia, M., Benamer, N. & Angulo, M. C. A specific GABAergic synapse onto

oligodendrocyte precursors does not regulate cortical oligodendrogenesis. *Glia* **65**, 1821–

1832 (2017).

9.   Mitew, S. *et al.* Pharmacogenetic stimulation of neuronal activity increases myelination

1062   in an axon-specific manner. *Nat. Commun.* **9**, 306 (2018).

1063   10.   Hines, J. H., Ravanelli, A. M., Schwindt, R., Scott, E. K. & Appel, B. Neuronal activity biases

1064   axon selection for myelination in vivo. *Nat. Neurosci.* **18**, 683–689 (2015).

1065   11.   Gibson, E. M. *et al.* Neuronal activity promotes oligodendrogenesis and adaptive

1066   myelination in the mammalian brain. *Science (80-.).* **344**, (2014).

1067   12.   Mensch, S. *et al.* Synaptic vesicle release regulates myelin sheath number of individual

1068   oligodendrocytes in vivo. *Nat. Neurosci.* **18**, 628–630 (2015).

1069   13.   Nagy, B., Hovhannisyan, A., Barzan, R., Chen, T. J. & Kukley, M. Different patterns of

1070   neuronal activity trigger distinct responses of oligodendrocyte precursor cells in the

1071   corpus callosum. *PLoS Biol.* **15**, (2017).

1072   14.   Barron, T. & Kim, J. H. Neuronal input triggers Ca2+ influx through AMPA receptors and

1073   voltage-gated Ca2+ channels in oligodendrocytes. *Glia* **67**, 1922–1932 (2019).

1074   15.   Sun, W., Matthews, E. A., Nicolas, V., Schoch, S. & Dietrich, D. Ng2 glial cells integrate

1075   synaptic input in global and dendritic calcium signals. *Elife* **5**, (2016).

1076   16.   Hamilton, N., Vayro, S., Wigley, R. & Butt, A. M. Axons and astrocytes release ATP and

1077   glutamate to evoke calcium signals in NG2-glia. *Glia* **58**, 66–79 (2010).

1078   17.   Marisca, R. *et al.* Functionally distinct subgroups of oligodendrocyte precursor cells

1079   integrate neural activity and execute myelin formation. *Nat. Neurosci.* **23**, 363–374

1080   (2020).

1081   18.   Krasnow, A. M., Ford, M. C., Valdivia, L. E., Wilson, S. W. & Attwell, D. Regulation of

1082   developing myelin sheath elongation by oligodendrocyte calcium transients in vivo. *Nat.*

Neurosci. **21**, 24–30 (2018).

19.   Zhang, Y. *et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons,

and vascular cells of the cerebral cortex. *J. Neurosci.* **34**, 11929–11947 (2014).

20.   Gross, G. G. *et al.* Recombinant Probes for Visualizing Endogenous Synaptic Proteins in

Living Neurons. *Neuron* **78**, 971–985 (2013).

21.   Hughes, A. N. & Appel, B. Oligodendrocytes express synaptic proteins that modulate

myelin sheath formation. *Nat. Commun.* **10**, (2019).

22.   Marques, S. *et al.* Oligodendrocyte heterogeneity in the mouse juvenile and adult central

nervous system. *Science (80-.).* **352**, 1326–1329 (2016).

23.   Spitzer, S. O. *et al.* Oligodendrocyte Progenitor Cells Become Regionally Diverse and

Heterogeneous with Age. *Neuron* **101**, 459-471.e5 (2019).

24.   Yoshii, A. & Constantine-Paton, M. BDNF induces transport of PSD-95 to dendrites

through PI3K-AKT signaling after NMDA receptor activation. *Nat. Neurosci.* **10**, 702–711

(2007).

25.   Meier, J. & Grantyn, R. A Gephyrin-Related Mechanism Restraining Glycine Receptor

Anchoring at GABAergic Synapses. *J. Neurosci.* **24**, 1398–1405 (2004).

26.   Nikonenko, I. *et al.* PSD-95 promotes synaptogenesis and multiinnervated spine

formation through nitric oxide signaling. *J. Cell Biol.* **183**, 1115–1127 (2008).

27.   Kukley, M., Nishiyama, A. & Dietrich, D. The fate of synaptic input to NG2 glial cells:

Neurons specifically downregulate transmitter release onto differentiating

oligodendroglial cells. *J. Neurosci.* **30**, 8320–8331 (2010).

28.   De Biase, L. M., Nishiyama, A. & Bergles, D. E. Excitability and synaptic communication

within the oligodendrocyte lineage. *J. Neurosci.* **30**, 3600–3611 (2010).

29.   Czopka, T., ffrench-Constant, C. & Lyons, D. A. Individual oligodendrocytes have only a

few hours in which to generate new myelin sheaths invivo. *Dev. Cell* **25**, 599–609 (2013).

30.   Hildebrand, D. G. C. *et al.* Whole-brain serial-section electron microscopy in larval

zebrafish. *Nature* **545**, 345–349 (2017).

31.   Baraban, M., Koudelka, S. & Lyons, D. A. Ca 2+ activity signatures of myelin sheath

formation and growth in vivo. *Nat. Neurosci.* **21**, 19–25 (2018).

32.   Koudelka, S. *et al.* Individual Neuronal Subtypes Exhibit Diversity in CNS Myelination

Mediated by Synaptic Vesicle Release. *Curr. Biol.* **26**, 1447–1455 (2016).

33.   Krueger-Burg, D., Papadopoulos, T. & Brose, N. Organizers of inhibitory synapses come of

age. *Curr. Opin. Neurobiol.* **45**, 66–77 (2017).

34.   Almeida, A. R. & Macklin, W. B. Early myelination involves the dynamic and repetitive

ensheathment of axons which resolves through a low and consistent stabilization rate.

bioRxiv 2022.07.17.500324 (2022) doi:10.1101/2022.07.17.500324.

35.   Osso, L. A., Rankin, K. A. & Chan, J. R. Experience-dependent myelination following stress

is mediated by the neuropeptide dynorphin. *Neuron* **109**, 3619-3632.e5 (2021).

36.   Almeida, R. G. *et al.* Myelination induces axonal hotspots of synaptic vesicle fusion that

promote sheath growth. *Curr. Biol.* **31**, 3743-3754.e5 (2021).

37.   Gerrow, K. *et al.* A preformed complex of postsynaptic proteins is involved in excitatory

synapse development. *Neuron* **49**, 547–562 (2006).

38.   Panzer, J. A. *et al.* Neuromuscular synaptogenesis in wild-type and mutant zebrafish. *Dev.*

Biol. **285**, 340–357 (2005).

39.   Wierenga, C. J., Becker, N. & Bonhoeffer, T. GABAergic synapses are formed without the

1128   involvement of dendritic protrusions. *Nat. Neurosci.* **11**, 1044–1052 (2008).

1129   40.   Villa, K. L. *et al.* Inhibitory Synapses Are Repeatedly Assembled and Removed at

1130   Persistent Sites In Vivo. *Neuron* **89**, 756–769 (2016).

1131   41.   Wake, H., Lee, P. R. & Fields, R. D. Control of local protein synthesis and initial events in

1132   myelination by action potentials. *Science (80-.).* **333**, 1647–1651 (2011).

1133   42.   Proctor, D. T. *et al.* Axo-glial communication through neurexin-neuroligin signaling

1134   regulates myelination and oligodendrocyte differentiation. *Glia* **63**, 2023–2039 (2015).

1135   43.   Stogsdill, J. A. *et al.* Astrocytic neuroligins control astrocyte morphogenesis and

1136   synaptogenesis. *Nature* **551**, 192–197 (2017).

1137   44.   Waggener, C. T., Dupree, J. L., Elgersma, Y. & Fuss, B. CaMKIIβ regulates oligodendrocyte

1138   maturation and CNS myelination. *J. Neurosci.* **33**, 10453–10458 (2013).

1139   45.   Mitra, S. K., Hanson, D. A. & Schlaepfer, D. D. Focal adhesion kinase: In command and

1140   control of cell motility. *Nature Reviews Molecular Cell Biology* vol. 6 56–68 (2005).

1141   46.   Venkatesh, H. S. *et al.* Neuronal activity promotes glioma growth through neuroligin-3

1142   secretion. *Cell* **161**, 803–816 (2015).

1143   47.   Almeida, R. G. & Lyons, D. A. Intersectional gene expression in zebrafish using the split

1144   KalTA4 system. *Zebrafish* **12**, 377–386 (2015).

1145   48.   Fuhrmann, J. C. *et al.* Gephyrin interacts with dynein light chains 1 and 2, components of

1146   motor protein complexes. *J. Neurosci.* **22**, 5393–5402 (2002).

1147   49.   Smolinsky, B., Eichler, S. A., Buchmeier, S., Meier, J. C. & Schwarz, G. Splice-specific

1148   functions of gephyrin in molybdenum cofactor biosynthesis. *J. Biol. Chem.* **283**, 17370–

17379 (2008).

50.   Craven, S. E., El-Husseini, A. E. & Bredt, D. S. Synaptic targeting of the postsynaptic

density protein PSD-95 mediated by lipid and protein motifs. *Neuron* **22**, 497–509 (1999).

51.   Topinka, J. R. & Bredt, D. S. N-terminal palmitoylation of PSD-95 regulates association

with cell membranes and interaction with K+ channel K(v)1.4. *Neuron* **20**, 125–134 (1998).

52.   Kwan, K. M. *et al.* The Tol2kit: A multisite gateway-based construction Kit for Tol2

transposon transgenesis constructs. *Dev. Dyn.* **236**, 3088–3099 (2007).

53.   Wyart, C. *et al.* Optogenetic dissection of a behavioural module in the vertebrate spinal

cord. *Nature* **461**, 407–410 (2009).

54.   Di Donato, V. *et al.* 2C-Cas9: A versatile tool for clonal analysis of gene function. *Genome*

Res. **26**, 681–692 (2016).

55.   Tinevez, J. Y. *et al.* TrackMate: An open and extensible platform for single-particle

tracking. *Methods* **115**, 80–90 (2017).

56.   Srinivasan, R. *et al.* Ca2+ signaling in astrocytes from Ip3r2 -/- mice in brain slices and

during startle responses in vivo. *Nat. Neurosci.* **18**, 708–717 (2015).

1164

Figure 1

Figure 2

a OPC
PSD-95-GFP myrTagRFP-T
PSD-95-GFP
GFP-Gephyrin myrTagRFP-T
GFP-Gephyrin

b Interneuron
PSD-95-GFP myrTagRFP-T
PSD-95-GFP
GFP-Gephyrin myrTagRFP-T
GFP-Gephyrin

c PSD-95-GFP
Disappearance of last process
p=0.0024

d GFP-Gephyrin
Disappearance of last process
p=0.0205

e PSD-95-GFP myrTagRFP-T
GFP-Gephyrin myrTagRFP-T

Figure 3

Figure 4

a 8 hours prior to myelination
GFP-Gephyrin myrmScarlet-I

b 6 hours into myelination

c Ctrl or TeNT
First sheath appearance
Injection
-10   0   6   (hours)
Continuous imaging

d
Hotspots
Hotspot centers
Sheath
freq. max
30%

e
Distance to myelin sheaths (µm)
p=0.0078
Hotspot   Nonhotspot

f
% predicting myelin sheath location
p=0.0156
Hotspot   Nonhotspot

g
Ratio
p=0.0406   p=0.0038
Ctrl
TeNT
Hotspot-sheath distance (µm)

h
Distance to myelin sheaths (µm)
p=0.0649
TeNT Hotspot   TeNT Nonhotspot

i
% predicting myelin sheath location
p=0.6250
TeNT Hotspot   TeNT Nonhotspot

Figure 5

a myrGCaMP6s MD

b 100% | 50 s

c
Occurances
6
624
MD
Soma

OPCs
5
175
only MD activity
MD and somal activity

d myrGCaMP6s MD synaptophysin

e membrane synaptophysin

f % of MDs assocd. with synaptophysin

g 0" 2" 4" 6"

h myrGCaMP6s synaptophysin | synaptophysin

i Calcium imaging
0   6 (hours)
MDs identified | Sheaths identified

j MDs Sheath

k Distance to myelin sheaths (µm)
p=0.0039
MD   NonMD

l % predicting myelin sheath location
p=0.0156
MD   NonMD

m 1 µm

Figure 6

Figure 7

Figure 8