Task feedback suggests a post-perceptual component to serial dependence

Jacqueline M. Fulvio*[1], Bas Rokers[2], Jason Samaha[3]

[1]Psychology, University of Wisconsin-Madison, Madison, WI 53562 USA

[2]Psychology, New York University Abu Dhabi, Abu Dhabi, UAE. Dept. of Psychology

and Center for Neural Science, New York University, New York, NY 10003, USA.

[3]Department of Psychology, University of California, Santa Cruz, Santa Cruz, CA

95060, USA

*Correspondence: jacqueline.fulvio@wisc.edu

Funding: J.M.F. and B.R. were supported by funding provided by Facebook Reality and
Google Daydream. The funders had no role in study design, data collection and
analysis, decision to publish, or preparation of the manuscript.

Abstract (162 words)

Decisions across a range of perceptual tasks are biased toward past stimuli. Such serial
dependence is thought to be an adaptive low-level mechanism that promotes perceptual
stability across time. However, recent studies suggest post-perceptual mechanisms
may also contribute to serially-biased responses, calling into question a single locus of
serial dependence and the nature of integration of past and present sensory inputs. We
measured serial dependence in the context of a 3D motion perception task where
uncertainty in the sensory information varied substantially from trial to trial. We found
that serial dependence varied with stimulus properties that impact sensory uncertainty
on the current trial. Reduced stimulus contrast was associated with an increased bias
toward the previous trial's stimulus direction. Critically, performance feedback, which
reduced sensory uncertainty, abolished serial dependence. These results provide clear
evidence for a post-perceptual locus of serial dependence in 3D motion perception and

30 support the role of serial dependence as a response strategy in the face of substantial

31 sensory uncertainty.

Keywords

34 Serial dependence; 3D motion perception; sensory uncertainty; feedback

Introduction

In recent years, a number of perceptual phenomena have been explained by appealing

to the idea that perceptual reports are not uniquely determined by external stimulus

properties but also depend on internal states of the observer. For instance, the success

of Bayesian-observer models in accounting for biases in perception support the role of

internal states in perception (Knill & Richards, 1996; Weiss et al., 2002; Knill, 2007;

Girshick et al., 2011; Rokers et al., 2018). A recent phenomenon that illustrates the

contribution of internal state is serial dependence: the finding that recent stimulus

history can bias current perceptual reports (Fründ et al., 2014; Fischer & Whitney,

2014).

Serial dependence is thought to be adaptive. Because most features in the environment

are stable, especially over short time scales, the brain may incorporate recent

information into current estimates as a means of reducing noise in instantaneous

sensory signals. Here, we sought to understand the contribution of sequential effects to

estimating 3D motion direction and the extent to which such serial biases are impacted

by task feedback and sensory uncertainty.

Serial dependencies have been observed across a range of high- and low-level

perceptual dimensions, task structures, and sensory modalities (Burr & Cicchini, 2014;

Fischer & Whitney, 2014; Liberman et al., 2014; Cicchini et al., 2014; Xia et al., 2016;

Alais et al., 2017; Bliss et al., 2017; Cicchini et al., 2017; Cicchini et al., 2018; Kiyonaga

et al., 2017; van Bergen & Jehee, 2017; Clifford et al., 2018; Fornaciai & Park, 2018;

Liberman et al., 2018; Alexi et al., 2018; Manassi et al., 2018; Suárez-Pinilla et al.,

2018; Fritsche & de Lange, 2019; Van der Burg et al., 2019; Pascucci et al., 2019;

Samaha et al., 2019; Barbosa & Compte, 2020; Kim et al., 2020; Cicchini et al., 2021;

de Azevedo Neto & Bartels, 2021; Ceylan et al., 2021; Kim & Alais, 2021; Pascucci &

Plomp, 2021; Manassi & Whitney, 2022; Goettker & Stewart, 2022). It is clear that serial

dependence is not simply due to hysteresis in motor processes as the effect does not

depend on the need for a motor response (Fisher & Whitney, 2014; Cicchini et al., 2018;

Manassi et al., 2018; Fornaciai & Park, 2018). However, beyond that, serial

dependencies appear to take on many forms and arise from different sources.

Numerous perceptual bases for serial dependence have been described. Recent work

has identified retinal error signals as one source of serial dependence in oculomotor

behavior (Goettker & Stewart, 2022), isolating the effects to the earliest stages of visual

processing. Serial dependence has been demonstrated to be retinotopic for orientation

judgments for stimuli up to 22 degrees of visual angle apart (Collins, 2019), which led to

the speculation that serial dependence may also arise at later stages of visual

processing. The bias was thought to arise from the activation of inferior temporal

neurons with wide receptive fields by orientation-selective neurons in striate and

extrastriate cortex, which then feed back to lower visual areas., This feedback was

thought to elevate activation of neurons with similar orientation tuning, resulting in the

biasing the population response toward the recently encountered orientation on the next

trial. Using a reverse correlation technique with classification images, Murai & Whitney

(2021) demonstrated that serial dependence biases the priors ('perceptual templates')

used by the visual system to selectively weigh the visual input according to task

demands. Serial dependence has also been shown to operate on perception of features

and integrated objects (Collins, 2021), in the integration of dynamic optic flow and static

form information in estimates of heading direction (Wang et al., 2022), and to directly

alter appearance of perceptual stimuli (Collins, 2020; Cicchini et al., 2017; Cicchini, et

al., 2021; Manassi & Whitney, 2022). Other work using orientation judgment tasks has

shown attractive serial dependence when past and previous stimuli were close in

orientation (Cicchini et al., 2017).

However, using a similar task, attractive serial dependence has also been argued to

operate at the decision level (Fritsche et al., 2017; see also Pascucci, et al., 2019), thus

implicating post-perceptual contributions. Further evidence for post-perceptual

contributions comes from the demonstration of negative (i.e., repulsive) serial

dependence (Pascucci et al., 2019) or serial dependence disappearing altogether (Bae

& Luck, 2020) when a previous stimulus is encoded but not reported. Moreover, the

finding that attractive serial dependence is observed between stimuli that share no

overlap in low-level features, but which require the same type of response (Ceylan et

al., 2021) supports a post-perceptual component. Recent modeling work has also

suggested that serial dependence arises in the mechanisms responsible for the readout

of the encoded stimulus in visual cortex (Sheehan & Serences, 2022). Finally, serial

dependence has been shown in other cognitive domains including working memory

(Fritsche et al., 2017; Bliss et al., 2017; Barbosa et al., 2020) and confidence (Rahnev

et al, 2015; Samaha et al., 2019).

The computational principle that determines when and how previous and current inputs

are combined also does not appear to be singular. The sensory cue integration

literature suggests cues are combined proportional to their associated uncertainty

(Trommershäuser et al., 2011; van Bergen & Jehee, 2019). Some accounts of serial

dependence are consistent with the cue integration literature and have demonstrated

uncertainty-based weighting of previous trial and current trial inputs (Cicchini et al.,

2014; Fischer & Whitney, 2014; Cicchini et al., 2018; Samaha et al., 2019; Kim & Alais,

2021). In some cases, serial dependence has been shown to be strongest when past

and present trial inputs are similar (Fritsche, 2016; Cicchini et al., 2018; Fritsche & de

Lange, 2019; Lidström, 2019). Recent work have also suggested that uncertainty-

weighting is not a general principle in that the uncertainty associated with only the

current, but not the previous trial, may be incorporated (Ceylan et al., 2021; Gallagher &

Benton, 2022). More generally, it has been suggested that serial dependence reflects

changes in criteria, resulting from integration with perceptual expectations, as a distinct

mechanism from priming (Galluzzi et al., 2022). Finally, it is typically difficult to

numerically assess an observer's estimated reliability of both previous and current trial

information, making uncertainty-weighted predictions challenging to test.

In the current study, we investigated serial dependence in the context of 3D motion

perception. Although the perception of 3D motion is critical to navigate our environment

and interact with our surroundings, the perception of 3D motion in the laboratory is often

systematically biased (e.g., Fulvio et al., 2015; Harris & Dean, 2003; Harris & Drga,

2005; Lages, 2006; Rushton & Duke, 2007; Welchman et al., 2004; Welchman et al.,

2008). Specifically, 3D motion stimuli tend to be associated with highly variable, but

well-characterized, estimates of perceptual uncertainty. For example, observers

frequently misreport the motion-in-depth direction of an object such that when

approaching, it is reported as receding, and vice versa (Fulvio et al., 2015; Fulvio &

Rokers, 2017). Such errors are not simply the result of guessing, but rather arise in a

systematic and principled way. First, the frequency with which they occur increases with

increased uncertainty associated with the motion signal, as occurs, for example, when

the contrast of the stimulus is reduced (Fulvio et al., 2015). Second, these errors are

specific to motion-in-depth as they rarely occur for lateral (i.e., leftward and rightward)

object motion. Third, these phenomena can be explained by a Bayesian-observer model

that optimally weighs incoming sensory information with prior knowledge about object

motion in general (Rokers et al., 2018). Finally, the frequency of perceptual misreports

decreases with task feedback (Fulvio & Rokers, 2017).

Because of the considerable variability in sensory uncertainty of 3D motion stimuli, and

their lawful relationship to stimulus contrast and motion direction, they are an ideal

candidate to investigate response biases due to serial dependence. Specifically, we

have shown previously that report feedback reduces the motion-in-depth misreports and

improves overall performance (i.e., percentage of interceptions). Rapid learning occurs

early in the first block of training followed by a much slower improvement over time

(Fulvio & Rokers, 2017). Because the performance improvements occur much more

rapidly than would be expected based on perceptual learning, we believe the

performance improvements are due to the recruitment of the available visual cues in the

stimulus display rather than changes in low level sensory processing *per se* (see also

Fulvio, Ji, et al., 2020). That is, experience with the 3D display appears to overcome

'flatness' priors developed through use of other displays outside the laboratory, where

binocular depth cues are typically unavailable. Thus, if serial dependence is diminished

with task feedback, the results would support a post-perceptual component of serial

bias in this task. Thus, by exploiting the relationship between 3D motion stimuli and their

associated perceptual uncertainty, we aimed to gain insight into the computational

principles underlying serial dependence and identify the source of serial effects.

Methods

Participants

Data from 72 participants were used in the current analysis. All were recruited from the

University of Wisconsin - Madison undergraduate student population and were naive to

the purposes of the study. The study was approved by the University of Wisconsin -

Madison institutional review board. All participants gave informed consent and received

course credit for their participation. After initial data analysis, data from one participant

were excluded from further analysis due to excessive serial bias beyond three standard

deviations of the group mean. This resulted in a final set of data from 71 participants (35

females) upon which all results reported in the text are based.

Experimental design

All participants completed three blocks of a 3D motion-in-depth extrapolation task while

wearing a virtual reality head-mounted display. We designed the experiment to

resemble the classic video game Pong to make the task intuitive to participants. A major

difference between our experiments and the classic video game however, is that we

used virtual reality to present stimuli in three dimensions (primarily the x-z plane),

whereas classic Pong presents stimuli in 2D (the x-y plane). See Supplementary

Materials for a video illustrating the stimulus and experimental procedure. In all, 37

participants from the final set of 71 used in the current analyses completed the task

without feedback on the accuracy of their reports, 10 participants completed the task

with auditory feedback, and 24 participants completed the task with visual and auditory

feedback.

Visual stimuli

On each trial, the participant fixated the center of a gray circular aperture (7.5 deg

radius) within a 1/f noise-mapped planar surface rendered at a viewing distance of 45

cm. A white spherical target (ball, 0.25 deg radius) appeared at fixation and traveled

along a random trajectory in the x-z plane for 1 s before disappearing (see **Figure 1**).

The target was rendered under perspective projection, so that both monocular (looming)

and binocular (disparity) cues to motion-in-depth were present. During the response

phase, a rectangular block (paddle, 0.5 cm × 1 cm × 0.5 cm) textured with 1/f

random noise was presented.

Fig. 1. **Experimental paradigm.** Upper panel: Participants viewed a 3D scene inside a
virtual reality headset that consisted of a 1/f noise-mapped planar surface ("wall") with a
circular aperture. Lower panel: A small fixation patch with superimposed nonius lines
resided in the center of the aperture. The trial sequence consisted of a white circular
target of variable contrast ('low', 'mid', or 'high') appearing at fixation and traversing a
random trajectory along the x (lateral) and z (depth) planes for 1 second before
disappearing. (Note that for display purposes, the targets are shown separately at
eccentricity in the "Visual display elements" panel.) When the target disappeared,
participants adjusted the position of a 3D rectangular block ("paddle") to the location
along an invisible orbit (drawn as a dashed circle here for display purposes) that would
intercept the target had it continued along its trajectory. One group of participants did
not receive feedback and pressed a key to begin the next trial with a new target. A
second group of participants received auditory feedback on every trial, and a third group
received visual and auditory feedback on every trial. The visual feedback sequence
comprised the target reappearing at its last visible location and continuing along its

trajectory to the invisible orbit. An interception occurred when the target reached the

paddle; otherwise, the setting resulted in a miss. The auditory feedback consisted of an

auditory signal corresponding to a "hit" sounded upon interception, otherwise, a "miss"

sound was played.

Procedure

Participants were asked to report the motion direction of the white circular target, i.e. a

ball. To do so, participants adjusted the position of the paddle so that it would intercept

the target. Paddle position was restricted to an invisible circular orbit centered on the

fixation point (**Fig. 1**) and was initialized at a random position on each trial. In the no-

feedback group, participants locked in their responses and continued to the next trial

after a 1 second fixation interval. In the feedback group, participants instead saw the

target reappear at its last location and continue along its trajectory until reaching the

(invisible) circular orbit. If the paddle had been placed such that the target was

intercepted, a cowbell sound played, otherwise, a "swish" sound played. The next trial

then began after the 1-second fixation interval.

Within a block, white target stimuli were presented at three different luminance levels

defined in terms of their Weber contrast levels computed by

$$\frac{(L_{target} - L_{background})}{L_{background}}$$

Where L_{target} and $L_{background}$ correspond to the luminances of the target and background,

respectively. The three target luminance levels were 77.92 cd/m^2, 69.02 cd/m^2, 65.68

cd/m^2 and presented against a gray background (44.56 cd/m^2), which corresponded to

0.75 ('high'), 0.55 ('mid'), and 0.47 ('low') Weber contrast levels, respectively.

Henceforth we use the labels 'high', 'mid', and 'low' to refer to these contrast levels just

as a matter of convenience. The different target contrast levels were presented in a

random, counterbalanced order. Across blocks, three rendering conditions were tested

in a random order across participants: (i) 'fixed' in which the scene did not update

according to the participant's head motion; (ii) 'active' in which the scene updated

according to the participant's head motion; (iii) 'lagged' in which the scene updated

according to the participant's head motion, but with a random lag of 50 ms on each trial

(no-feedback group) or a random lag on each trial drawn from the uniform distribution [0

500 ms] (feedback group). We note that the three rendering conditions were included in

the study design to address questions beyond the scope of this investigation and do not

materially impact the serial dependence results reported here. These details are

described more extensively in a separate study investigating the role of response

feedback in the recruitment of visual cues in virtual reality (Fulvio & Rokers, 2017).

Participants in the no-feedback and auditory feedback groups completed 225 trials per

block for a total of 675 trials. Participants in the auditory + visual feedback group

completed 120 trials per block for a total of 360 trials.

Analysis of current trial performance

Response error on each trial was computed as the difference between presented and

reported motion direction. We computed the circular distance between the midpoint of

the participant's paddle setting and the endpoint of the presented target trajectory using

the *circ_dist.m* function of the Circular Statistics Toolbox for Matlab (Berens, 2009; **Fig.**

2a). Error distributions were plotted and summarized according to the mean standard

deviation of the signed errors, mean absolute error, and proportion of motion-in-depth

misreports for comparisons (i). across target contrast level conditions within each

feedback condition using one-factor repeated-measures ANOVA. In some cases, the

assumption of sphericity was violated, so a Greenhouse-Geisser correction was

included in the model; , and (ii). across feedback conditions using two-sample *t*-tests. In

some cases, a two-sample *F*-test for equal variances produced significant results,

indicating that the two samples violated the equal variance assumption. In those cases,

a Welch's two-sample *t*-test was used where indicated.

Serial dependence analysis

Serial dependence was quantified using a pipeline similar to that used by Samaha et al.

(2019). For each subject, the response error on each trial was computed as the angular

distance between the presented and reported motion trajectory. Following previous

serial dependence work (Bliss et al., 2017; Fritsche et al., 2017; Samaha et al., 2019),

high-error trials were omitted from analysis. In this task, high-error trials were defined as

those in which participants misreported the direction of the target's motion in depth (i.e.,

when participants reported an approaching target as receding and vice versa,

henceforth named "depth misreport trials" contrasted with "correct depth trials"). Such

depth misreport trials were more prevalent in the no-feedback condition compared to

the feedback conditions (38.3% vs. 18.1% (auditory) and 18.5% (auditory + visual),

respectively, on average across conditions). We emphasize that these depth misreports

were not simply random settings - the lateral (x) component of the target's motion was

almost always reported correctly on these trials. We also carried out the analyses with

depth misreport trials included and include them where appropriate, primarily in

Supplemental Materials. To remove any global directional biases for individual

participants, we subtracted their mean signed error across trials from each trial. On

average, these mean signed errors computed across all trials were small ($M_{no\text{-}feedback} =$

$0.353°$, $SD_{no\text{-}feedback} = 3.173°$; $M_{auditoryfeedback} = -0.179°$, $SD_{auditoryfeedback} = 2.622°$;

$M_{auditory+visualfeedback} = 0.165°$ $SD_{auditory+visualfeedback} = 3.391°$).

Model-based serial dependence was then quantified using the standard approach of

sorting each participant's demeaned errors according to the angular difference between

the presented target direction on the current and previous trials. We note that because

the presented target directions spanned the full 360° space, the angular difference scale

ranged from [-180°, 180°], twice the range of studies using orientation stimuli. The first

trial of each block was omitted because it did not have a previous trial. When trials were

removed due to response misreports as described above, the most immediately

preceding non-misreport trial served as the previous trial. This approach is reasonable

given previous work showing that serial dependence may extend to at least three trials

back (Fischer & Whitney, 2014).

The sorted errors for each participant were then smoothed by a 45-degree wide median

filter in 1 degree increments. Similar results were obtained with smaller filters. We

averaged the smoothed data within each group and fitted the averages of each group

with a derivative-of-Gaussian (DoG) function of the following form:

$$y = xawce^{-(wx)^2}$$

where x is the relative orientation of the target of the previous trial, a is the amplitude of

the peaks of the curve, w scales the width of the Gaussian derivative, and c is a

constant $\sqrt{2}/e^{-0.5}$, which scales the amplitude parameter to numerically match the

height of the curve in degrees. Fitting was achieved by minimizing the sum of squared

errors using the *lsqcurvefit* MATLAB routine, with amplitude a and width w parameters

free to vary between [-10 deg, 10 deg] and [0.001, 0.5], respectively.

Statistical significance of the group-level DoG fits was assessed through a bias-

corrected and accelerated (BCa) bootstrapping procedure. On each of 100,000

iterations, we sampled subjects with replacement and fit a DoG to the average of the

bootstrap sampled data. The value of the amplitude parameter was retained after each

iteration, resulting in a distribution of the amplitude parameter of our sample. We also

computed the jackknife distribution of amplitude parameters by computing all of the

leave-one-sample-out DoG fits. The confidence interval and p-value of the amplitudes of

each group fit when compared to zero (i.e., no serial dependence) were computed using

the *BCa_bootstrap* MATLAB function (Van Snellenberg, 2018) which utilized both the

bootstrapped and jackknife amplitude distributions.

Between-group comparisons of DoG fit amplitudes were carried out using permutation

testing. On each of 100,000 iterations, we shuffled the group labels (e.g., 'no feedback'

and 'auditory feedback') and then assigned each data set with one of the shuffled

labels. We then grouped the data by the shuffled labels and fit a DoG to the average of

each shuffled group. The value of the amplitude parameter was retained after each

iteration, resulting in a permutation distribution of the amplitude parameter. The p-value

of the original difference in amplitudes between the two groups was derived from the

permutation distribution as the proportion of the permutations greater than the original

difference.

Following Samaha et al. (2019), we additionally carried out a model-free analysis to

verify that the results were not specific to the model-fitting. For each participant, we

computed the median (signed) error across trials where the relative difference between

the current and previous stimulus fell within the 0-180 deg interval, and subtracted that

from the median error on trials within the -180-0 deg interval. Thus, positive and

negative values indicate an attractive or repulsive bias, respectively. Statistical testing

was performed using two-tailed *t*-tests, one-way ANOVAs, or a linear mixed effects

model as described in the main text.

Both model-based and model-free analyses were carried out multiple times on trial

sortings according to several questions addressed below. For the basic comparisons

between serial dependence on previous target direction, and between performance with

and without feedback, trials across all three blocks were used. For block-based

comparisons, serial dependence was estimated separately for each of the three blocks.

For target contrast-based comparisons, serial dependence was estimated for each of

the three target contrast levels, combining trials across blocks.

Results

Performance improves with both task feedback and target contrast

We first considered performance as a function of current task conditions. For each trial,

we computed the response error as the circular distance between the presented and

reported target trajectory (**Fig. 2a**). When feedback was not provided, mean absolute

error in motion direction report changed as a function of target contrast (low: M = 55.1

deg (SD = 11.8 deg); mid: M = 51.2 deg (SD = 12.6 deg); high: M = 44.7 deg (SD = 13.2

deg); $F_{(1.664, 59.905)} = 27.310$, $p < .001$) as did the proportion of motion-in-depth

misreports (F(1.699,61.156) = 14.172, p < .001) and the standard deviations of the

response errors (**Fig. 2b bar graph inset**).

When feedback was provided, mean absolute error in motion direction report was

significantly reduced when compared to the performance of the no-feedback group for

both feedback types (auditory: M = 28.5 deg (SD = 8.5 deg); t(45) = 5.549, p < .001;

auditory + visual: M = 24.9 deg (SD = 13.7 deg); t(59) = 7.702, p < .001) with no

difference between the two feedback conditions (t(32) = 0.7691, p .448) with the same

pattern observed in the standard deviations of the response errors (**Fig. 2c bar graph**

inset). Relatedly, feedback was associated with a reduction in motion-in-depth

misreports (38.3% (no feedback) vs. 18.1% (auditory feedback; t(45) = 7.056, p < .001)

and 18.5% (visual + auditory feedback; t(59) = 7.5584, p < .001), on average, with no

difference between the two feedback conditions t(32) = 0.1027, p = .919).

Mean absolute error in motion direction report for the auditory feedback group changed

as a function of target contrast (low: M = 36.8 deg (SD = 14.9 deg); mid: M = 30.2 deg

(SD = 13.2 deg); high: M = 27.8 deg (SD = 10.9 deg); F(2,18) = 11.115, p < .001) as did

the proportion of motion-in-depth misreports (F(2,18) = 10.828, p < .001) and the

standard deviations of the response errors (**Fig. 2d bar graph inset**). Mean absolute

error in motion direction report for the auditory + visual feedback group also changed as

a function of target contrast (low: M = 28.2 deg (SD = 17.7 deg); mid: M = 25.6 deg (SD

= 14.1 deg); high: M = 21.4 deg (SD = 13.2 deg); F(2,46) = 9.037, p < .001) as did the

proportion of motion-in-depth misreports (F(2,46) = 3.720, p = .032) and the standard

deviations of the response errors (**Fig. 2d bar graph inset**). In summary, response

errors in the judgment of 3D motion direction decreased within the same observer as

stimulus contrast increased. Similarly, response errors decreased when observers

received task feedback, compared to when observers did not receive feedback.

We next describe the results of the serial dependence analyses in turn. For

completeness, Table 1 at the end of the Results section summarizes all key findings.

Fig. 2. Behavioral performance on the current trial. (**a.**) Error in report on the current

trial is defined as the circular distance in degrees between the current trial target

direction and the current trial reported direction. (**b.**) Between-subjects error histograms

for the group that did not receive feedback for the high (light blue), mid (red), and low

(yellow orange) target contrast levels. Left inset depicts the standard deviation of the

errors for the three target contrast levels. (**c.**) Between-subjects error histograms for the

group that did not receive feedback (green), the group that received auditory feedback

(maroon) and the group that received both visual and auditory feedback (dark blue).

The inset depicts the standard deviation of the errors for the feedback groups. (**d.**)

Same format as (**b.**) for the auditory feedback group (left) and the auditory + visual

feedback group (right). Error bars correspond to +/- 1 *SEM*; ** corresponds to

significance at the Bonferroni-corrected alpha value = .0167.

Prior stimuli bias current responses

We next computed serial dependence in the 3D motion perception task in the absence of task feedback. In keeping with previous practice, we omitted high error trials (i.e., motion-in-depth misreport trials; but see **Figs. S1-S2** in *Supplementary Materials* for results including all trials). Consistent with previous studies, current responses were biased toward the previous trial's presented direction (serial dependence amplitude, a = 2.56°, 95% CI: [1.39°, 3.87°], $p < .001$) ; **Fig. 3a**). The significant serial dependence was confirmed with our model-free measure of serial bias (M = 3.01°; $t(36) = 3.3050$, $p = .002$; **Fig. 3b**). Indeed, the magnitude of bias is consistent with prior work using motion and orientation stimuli (Alais et al., 2017; Fritsche & de Lange, 2019; Samaha et al., 2019; Kim et al., 2020; Celyan et al., 2021; but see Fischer & Whitney, 2014 who reported biases >8° in several conditions).

Fig. 3. **Task feedback reduces serial dependence on previous target direction. (a.)**
Serial dependence curves and DoG fits to error data with high error trials omitted (i.e.,
on correct depth report trials following correct depth report trials), when no feedback
was provided (green), when auditory feedback was provided (red), and when auditory +
visual feedback was provided (blue), sorted according to the relative direction of the
current trial's target with respect to the previous trial's target direction. Shaded bands
represent +/-1 *SEM*. **corresponds to bootstrapped *p*-values < .001; n.s. = non-
significant; ☐*a* refers to the difference in amplitude parameter between no feedback and
auditory feedback groups (red font) and between no feedback and auditory + visual
feedback groups (blue font) with corresponding *p*-values derived from permutation
testing. **(b.)** Model-free serial bias with respect to the previous trial's target direction for
the three feedback conditions. Circular symbols correspond to individual subject biases;
black symbols correspond to group means. Error bars represent +/-1 *SEM*.
corresponds to *p*-values < .01; *corresponds to *p*-values < .05. **(c.) Same formats as

(a.) & **(b.)** split by block. One outlier with bias > 45° in the no feedback condition of

Block 1 was excluded from the plot.

Task feedback reduces serial dependence

The significant serial bias reported above extends the finding of serial dependence to

the 3D motion domain. To further examine the nature of serial dependence in this task,

we analyzed the impact of task feedback. In previous work, we have shown that

feedback is associated with an improved ability to intercept the target (Fulvio & Rokers,

2017). Furthermore, our previous work suggests that visual and auditory report

feedback improve performance by encouraging the appropriate recruitment of available

cues in estimating the motion-in-depth direction (Fulvio & Rokers, 2017; Fulvio, Ji, et al.,

2020). This, in turn, reduces the prevalence of motion-in-depth misreports. Our

interpretation of the mechanism through which feedback impacts performance in this

task is consistent with a post-perceptual locus of feedback-based effects - rather than

feedback impacting the quality of early sensory representations, feedback appears to

affect the integration of the sensory estimates based on the various cues available in

the stimulus. We therefore hypothesized that feedback would also be associated with a

reduction in serial dependence as cue recruitment provides more reliable sensory

signals.

Consistent with this expectation, task feedback eliminated serial dependence of the

current trial's report on the previous trial's target motion direction according to the

model-based analysis (auditory feedback: a = 0.12°, 95% CI: [-2.97°, 1.18°], p = .96;

auditory + visual feedback: a = 0.81°, 95% CI: [-0.39°, 1.45°], p = .23; **Fig. 3a**), with no

difference in serial dependence observed between the two feedback groups with the

model-based measure (Δa = 0.6898°, p = .173). However, the model-free analysis

suggests a small bias remained with feedback. For the auditory + visual feedback

group, this bias remained attractive toward the previous report (M = 1.26°; t(23) =

2.5225, p = .02), whereas the bias was repulsive from the previous report for the

auditory feedback group (M = -1.01°; t(9) = -2.3571, p = .043; **Fig. 3b**). The difference in

the model-free measure of serial bias for the two feedback groups was significant (t(32)

= 2.7509, p = .01).

Critically, however, both measures revealed a significant reduction in serial dependence

with feedback compared to performance when feedback was not provided (auditory

feedback: (model-based) $\Box a$ = 2.44°, p = .031; (model-free): Welch's two-sample

t(44.91) = 3.9952, p < .001; auditory + visual feedback: (model-based): $\Box a$ = 1.75°, p =

.019**;** (model-free): Welch's two-sample t(53.36) = 1.6845, p = .049).

Investigating how serial dependence changes across the three experimental blocks, we

found that dependence on previous target direction was already reduced compared to

reports without feedback by the end of the first block, with both auditory feedback

(model-based: a = 1.209°, 95% CI: [-1.5421°, 2.2492°], p = .578; model-free: M =

1.0292°, t(9) = 1.1174, p = 0.293) and auditory + visual feedback (model-based: a =

2.088°, 95% CI: [-1.128°, 3.838°], p = .213; model-free: M = 2.796°, t(23) = 2.306, p =

0.031) compared to the no-feedback group, (model-based: a = 3.576°, 95% CI: [1.820°,

5.796°], p = .009; model-free: M = 4.202°, t(36) = 2.672, p = 0.011). The difference

failed to reach significance by both measures for the auditory feedback and no feedback

group comparison (model-based: $\Box a$ = 2.555°, p = .101; model-free: Welch's two-

sample t(44.154) = 1.7408, p = .089; **Fig. 3c**). Likewise, the difference failed to reach

significance by both measures for the auditory + visual feedback and no feedback group

comparison (model-based: $\Box a$ = 1.488°, p = .148; model-free: Welch's two-sample

t(58.928) = 0.7079, p = .48).

The model-based measure indicated no detectable serial dependence in blocks 2 and 3

with auditory feedback, whereas the model-free measure indicated a significant

repulsive bias in block 2 (block 2: model-based: a = 0.0464°, 95% CI: [-7.1092°,

0.9545°], p = .459; model-free: M = -1.9897°, t(9) = 2.6201, p = 0.028; block 3: model-

based: a = -0.4366°, 95% CI: [-2.7976°, 1.7943°], p = .951; model-free: M = -1.2362°,

t(9) = 1.0245 , p = 0.33). This significance repulsive bias was significantly different from

the model-free estimate for both the no feedback and auditory + visual feedback

conditions ($p < .01$ for both comparisons). With auditory + visual feedback, serial

dependence was undetectable in blocks 2 and 3 by both measures (block 2: model-

based: $a = 1.241°$, 95% CI: [-1.108°, 3.194°], $p = .246$; model-free: $M = 1.421°$, $t(23) =$

1.470, $p = 0.155$; block 3: model-based: $a = 1.257°$, 95% CI: [-1.310°, 4.064°], $p = .512$;

model-free: $M = 0.917°$, $t(23) = 1.102$, $p = 0.28$).

Because our study design included additional manipulations, some of which were not

relevant to the current analyses, we fit a linear mixed effects model to the model-free

serial bias data that accounts for all manipulations. Main effects of (i) feedback (3 levels:

auditory, auditory + visual, & without), (ii) current trial target contrast (3 levels: low, mid,

& high), (iii) head-tracking condition (3 levels: fixed, active, & lagged), and (iv) block

order (3 levels: first, second, third) were included in the analysis as fixed effects.

Feedback and head-tracking condition were coded as categorical variables, and block

order was coded as an ordinal variable. Additionally, the interaction between block and

feedback was included. As random effects, the model included intercepts for subjects.

Visual inspection of residual plots did not reveal any obvious deviations from

homoscedasticity or normality. P-values were obtained by F-tests for each term in the

linear mixed-effects model. The main effect of current trial target contrast was significant

($F(1,627) = 25.321$, $p < .001$). The main effect of feedback remained marginally

significant when the feedback-block interaction was included in the model ($F(2,627) =$

2.844, $p = .056$), but was statistically significant when this term was left out ($F(2,631) =$

11.537, $p < .001$). There were no main effects of head-tracking condition ($F(2,631) =$

0.0996, $p = .905$) or block ,($F(2,631) = 0.616$, $p = .541$), and we did not find a block by

feedback interaction ($F(4,627) = 0.07716$, $p = .989$). The failure to find a significant

effect of block order is consistent with the notion that task feedback already affects

performance in an observer's first block of the task. It is worth noting that because there

was variability in the effect of block due to the differential effects of head-tracking

conditions on performance overall (see Fulvio & Rokers, 2017), specific effects of these

factors are likely under-powered and obscured in this data set. This may also be an

indirect source of the reduced effect of feedback when the interaction term was included

in the model. A more targeted investigation of block order by feedback effects may be of

interest for future research. We focus on one factor that directly impacts sensory

uncertainty - target contrast - below.

Current trial sensory uncertainty increases serial dependence

A sensory uncertainty account of serial dependence predicts that greater uncertainty in

the estimate of the current target's motion leads to greater reliance on recent

experience. And, conversely, that greater uncertainty in the previous trial estimate

should lead to a smaller impact of recent experience. Here, we tested both predictions

by quantifying serial dependence as a function of our manipulation of target contrast.

We first measured serial dependence as a function of current trial target contrast.

Increases in current trial target contrast were associated with reduced serial

dependence on the previous trial's target direction. In particular, serial dependence was

abolished on high target contrast trials, consistent with a sensory uncertainty account

(**Fig. 4a**). Without feedback, both reduced target contrast levels were associated with

significant serial dependence according to both serial dependence measures (model-

based: a_{high} = 0.174°, 95% CI: [-2.631°, 0.927°], p = .265; a_{mid} = 4.299°, 95% CI:

[2.633°, 6.048°], p < .001; a_{low} = 4.374°, 95% CI: [2.467°, 6.252°], p = .01; model-free:

low: M = 4.269°, $t(36)$ = 2.929, p = .006; mid: M = 3.467°, $t(36)$ = 2.644, p = .012). With

auditory + visual feedback, only the lowest target contrast level was associated with

significant serial dependence according to both measures (model-based: a_{high} = -0.976°,

95% CI: [-1.869°, 1.069°], p = .151; a_{mid} = 0.987°, 95% CI: [-1.138°, 2.171°], p = .569;

a_{low} = 3.279°, 95% CI: [1.696°, 5.309°], p < .001; model-free: low: M = 2.41°, $t(23)$ =

3.021, p = .006). With auditory feedback, none of the target contrast levels was

associated with significant serial dependence according to both measures (a_{high} =

1.438°, 95% CI: [-1.188°, 3.785°], p = .160; a_{mid} = -1.485°, 95% CI: [-2.875°, 0.717°], p =

.273; a_{low} = 1.367°, 95% CI: [-2.605°, 2.498°], p = .132).

Dependencies were significantly smaller when feedback was provided compared to

when it was not provided for the mid target contrast level only (permutation test *p*-values

≤.0037 for both mid target contrast level comparisons; *p* ≥ .0913 for all other

comparisons). Furthermore, analysis of the model-free serial bias measure revealed a

significant effect of current trial target contrast for the no feedback group ($F(1,109)$ =

6.147, *p* = .015) and for the auditory + visual feedback group ($F(1,70)$ = 5.319, *p* =

.024), but the effect failed to reach significance for the auditory feedback group ($F(1,28)$

= 3.703, *p* = .065).

Interestingly, the relationship with the previous trial's target contrast was less straight-

forward (**Fig. 4b**). A sensory uncertainty based explanation might predict that greater

target contrast on the previous trial would be associated with a more reliable estimate

on that trial, which should have greater impact on the current trial's estimate. With

auditory + visual feedback, the model-based measure indicated no reliance on the

previous target direction at any target contrast level ($a_{previoushigh}$ = 0.789°, 95% CI: [-

1.225°, 1.558°], *p* = .586; $a_{previousmid}$ = 0.911°, 95% CI: [-1.417°, 1.909°], *p* = .792;

$a_{previouslow}$ = 0.709°, 95% CI: [-1.379°, 1.663°], *p* = .902), whereas the model-free

measure revealed a small positive bias on trials following mid-level target contrast trials

only (M = 1.845°, $t(23)$ = 2.0859, *p* = .048). The model-based measure also indicated no

reliance on the previous target direction at any target contrast level for the auditory

feedback group ($a_{previoushigh}$ = 0.173°, 95% CI: [-2.905°, 1.917°], *p* = .965; $a_{previousmid}$ = -

1.124°, 95% CI: [-4.187°, 1.206°], *p* = .501; $a_{previouslow}$ = 1.610°, 95% CI: [-0.969°,

2.776°], *p* = .081), but the model-free measure revealed significant negative biases on

trials following mid- and high-level target contrast trials (mid: M = -1.872°, $t(9)$ = 4.218, *p*

= .002; high: M = -1.525°, $t(9)$ = -2.381, *p* = .041). However, in the no-feedback

condition, reliance on the previous trial's target direction was significant when the

previous trial's target contrast was mid or low, but not when it was high by both

measures (model-based: $a_{previoushigh}$ = 0.604°, 95% CI: [-3.275°, 2.084°], *p* = .89;

$a_{previousmid}$ = 3.769°, 95% CI: [2.268°, 5.283°], *p* < .001; $a_{previouslow}$ = 4.212°, 95% CI:

[2.598°, 5.885°], *p* < .001; model-free: low: M = 5.243°, $t(36)$ = 4.247, *p* < .001; mid: M =

4.703°, $t(36)$ = 4.187, *p* < .001**).**

Dependencies were significantly smaller when auditory + visual feedback was provided

compared to when it was not provided when the previous trial's target contrast was low

(permutation test p-value = .003) and trending in the same direction when auditory

feedback was provided (permutation test p-value = .061). When the previous trial's

target contrast was at the mid-level, serial dependence was significantly smaller for the

two groups in which feedback was provided compared to the no feedback condition

(permutation test $p \leq$ .0088 for both comparisons). No differences were observed for the

comparisons of serial dependence when the previous trial's target contrast level was

high (permutation test $p \geq$ .384 for both comparisons). This pattern of results suggests

that the reliability of the current trial's sensory estimate is a strong predictor of the

degree of serial dependence in this task, but the reliability of the previous trial sensory

reliability, less so (see also Gallagher & Benton, 2022).

In an additional analysis, we considered the *relative* contrast between the current and

the previous trial and found that dependence on the previous trial's target direction was

only significant for the no feedback group when the current trial's target contrast

matched the previous trial's target contrast according to both serial bias measures (a =

5.405°, 95% CI: [3.549°, 7.572°], $p <$ .001; all other p-values $\geq$ .2448; model-free: M =

6.085°, $t(36)$ = 4.127, $p <$ .001; **Fig. S3**). In summary, these results suggest that

sensory uncertainty as manipulated either through task feedback or stimulus contrast on

the current trial, strongly modulates serial dependence.

Fig. 4. Serial dependence is modulated by sensory uncertainty due to target contrast. (a.) Left: Serial dependence curves and DoG fits to error data with high error trials omitted (i.e., on correct depth report trials following correct depth report trials), when no feedback was provided (green), when auditory feedback was provided (red), and when auditory + visual feedback was provided (blue), sorted according to the relative direction of the current trial's target with respect to the previous trial's target direction and split out by current trial target contrast. Shaded bands represent +/-1 *SEM*. **corresponds to bootstrapped *p*-values < .001; n.s. = non-significant; *p*-values at the bottom of each graph refer to the difference in amplitude parameter between no feedback and auditory feedback groups (red font) and between no feedback and auditory + visual feedback groups (blue font) derived from permutation testing. Right: Model-free serial bias with respect to the previous trial's target direction for the three feedback conditions, split out by current trial target contrast. Circular symbols correspond to individual subject biases; black symbols correspond to group means.

636 Error bars represent +/-1 *SEM*. **corresponds to p-values < .01; *corresponds to p-

637 values < .05; n.s. = non-significant. **(b.)** Same formats as **(a.)** split according to previous

638 trial target contrast.

Analysis	Serial bias measure	
	DoG amplitude	Model-free
Overall	NF: 2.56° ** AF: 0.12° AV: 0.81° --------------- AF vs. NF *	NF: 3.01° ** AF: **-1.01° *** AV: 1.26° * --------------- AF vs. NF ** AV vs. NF * AF vs. AV *
By block (B1 / B2 / B3)	NF: 3.58° ** / 2.21° * / 2.58° * AF: 1.21° / 0.05° / **-0.44°** AV: 2.09° / 1.24° / 1.26° --------------- B3: AF vs. AV *	NF: 4.20° * / 2.90° ** / 2.20° * AF: 1.03° / **-1.99° *** / **-1.24°** AV: 2.80° * / 1.42° / 0.92° --------------- B2: AF vs. NF ** B2: AF vs. AV **
Target contrast - current trial (Low / Mid / High)	NF: 4.37° * / 4.30° ** / 0.17° AF: 1.37° / **-1.49°** / 1.44° AV: 3.28° ** / 0.99° / **-0.98°** --------------- Mid: AF vs. NF ** Mid: AV vs. NF **	NF: 4.27° * / 3.47° ** / 1.21° AF: **-1.18°** / **-1.90° *** / 0.88° AV: 2.41° ** / 1.14° / 0.04° --------------- Low: AF vs. NF * Low: AF vs. AV * Mid: AF vs. NF * Mid: AF vs. AV *
Target contrast (previous trial) (Low / Mid / High)	NF: 4.21° ** / 3.77° ** / 0.60° AF: 1.61° / **-1.12°** / 0.17° AV: 0.71° / 0.91° / 0.79° --------------- Mid: AF vs. NF ** Mid: AV vs. NF **	NF: 5.24° ** / 4.70° ** / **-0.70°** AF: 0.64° / **-1.87° *** / **-1.53° *** AV: 1.42° / 1.84° * / 0.85° --------------- Low: AV vs NF ** Mid: AF vs. NF ** Mid: AV vs. AF * High: AV vs. AF *
Target contrast (relative current and previous trial) (Curr lower / Curr higher / Curr same)	NF: 1.88° / 1.44° / 5.40° ** AF: **-0.002°** / 0.39° / 0.29° AV: 1.26° / **-0.79°** / 1.34° --------------- Curr same: AF vs. NF ** Curr same: AV vs. NF **	NF: 0.75° / 1.91° / 6.08° ** AF: **-1.76°** / **-0.28°** / **-0.84°** AV: 1.20° / **-0.02° *** / 2.20° --------------- Curr lower: AV vs AF * Curr same: AF vs. NF **

		Curr same: AV vs. NF *

Table 1. Summary of serial bias results. The first three lines of each cell report the bias
obtained in each of the three conditions. Additional lines report significant comparisons among
feedback conditions. *NF* = no feedback; *AF* = auditory feedback; *AV* = auditory + visual
feedback. Bolded values highlight repulsive biases. *corresponds to *p* < .05; **corresponds to *p*
< .01.

Discussion

Serial dependence – the impact of recent stimulus history on current perceptual reports

(Fründ et al., 2014; Fischer & Whitney, 2014) – is a pervasive finding in many

psychophysical domains. In the current study, we show that serial dependence occurs

in psychophysical reports of 3D motion direction. Critically, however, we showed that

serial dependence is not a given - reducing sensory uncertainty by increasing target

contrast or providing performance feedback on a trial-to-trial basis was associated with

near abolishment of serial dependence.

The target contrast-driven effects on serial dependence extend recent findings. In

particular, the sensory reliability of the current stimulus was a stronger and more

consistent predictor of serial dependence than the reliability of the previous stimulus.

Whereas increases in current trial target contrast were associated with reductions in

serial dependence for both feedback and no feedback groups, the effect of previous trial

target contrast was less clear. We note that one potential limitation of our design was

the use of above-detection threshold stimuli. Including lower contrast targets may have

revealed serial dependence effects in the feedback conditions as well. Nevertheless,

these results are consistent with other work showing that uncertainty in the current

sensory estimate drives serial dependence more than previous trial uncertainty (e.g.,

Pascucci et al., 2019; Ceylan et al., 2021; Gallagher & Benton, 2022). Our results also

support previous work showing that similarity between successive stimuli matters

(Fritsche, 2016; Cicchini et al., 2018; Fritsche & de Lange, 2019; Lidström, 2019).

By contrast, the feedback-driven attenuation of serial dependence is a novel finding. In

fact, our results appear to contradict a recent report of *increased* attractive serial bias

found in numerosity estimates with feedback (Fornaciai & Park, 2022). In that study,

feedback was provided to participants in the first eight experimental blocks, followed by

four blocks without feedback. Because serial dependence was smaller for the final four

no feedback blocks compared to the four blocks with feedback, and on par with serial

dependence observed in a previous study by the group in the absence of feedback

(Fornaciai & Park, 2020), the interpretation was that feedback encouraged greater

weight on the previous trial's stimulus, especially when the feedback on the previous

trial was 'correct'. However, with the fixed block order design, it is not clear whether the

effects were in fact feedback-based or merely experience-based. Indeed, in our study,

we showed that as experience evolved in the task, serial dependence declined – even

the no-feedback group's performance exhibited hints of a decline in serial dependence

over blocks. However, we emphasize that although feedback was associated with a

general improvement in behavioral performance, including a reduction in response

errors, those who learned to reduce response errors more over the course of the

experiment did not necessarily reduce serial dependence more. Because the

relationship between error variability and serial bias was not systematic, it does not

seem that the effect of feedback on sensory reliability is the sole driver of the effect of

feedback on serial dependence. Thus, taken together, both sets of results, those

reported here combined with the recent work by Fornaciai & Park (2022) support the

notion that serial dependence may originate at higher, post-perceptual levels, as also

suggested by other results across the literature (Fritsche et al., 2017; Pascucci et al.,

2019; Bae & Luck, 2020; Ceylan et al., 2021).

We note that in the auditory + visual feedback group, participants were presented again

with the stimulus as well as their response during the feedback stage. Thus, participants

in this group were provided with an opportunity to update their estimate of the motion

direction and target endpoint on that trial. Nevertheless, this did not encourage

continued dependence on the previous target direction beyond the first block. When the

visual component was removed from the feedback for the auditory feedback group,

serial dependence was, if anything, repulsive. The distinct nature of the feedback

signals in these two conditions may nevertheless have contributed to the subtle

differences in serial dependence between the two feedback groups. Taken together, our

results raise the possibility that serial dependence is subject to strategic control, driven

at least in part by error signals from feedback about the accuracy of one's previous

perceptual estimates.

In previous work, we interpreted improvements in accuracy of 3D motion-in-depth

judgments with feedback as the result of the inexperienced participants learning to

overcome "flatness" priors developed through extensive real-world use of 2D displays

and to compel recruitment of the available depth cues that they were otherwise

discounting (Fulvio & Rokers, 2017; Fulvio, Ji et al. 2020). Given that integration of

multiple cues enhances reliability of visual perception (Chang et al., 2020; Hillis et al.,

2004; Knill & Saunders, 2003; Murphy et al., 2013; Oruc et al., 2013; Preston et al.,

2009; Rideaux & Welchman, 2018; Welchman et al., 2005), increasing the number of

cues integrated in estimating the target's motion direction may be one way in which

feedback reduces serial dependence. (We note, however, that our design leaves open

the possibility that other factors may also have accounted for the effects of feedback,

given that adjustment responses are subject to some motor error and possible biases

towards/away from certain directions. Future work employing other paradigms such as

2AFC tasks can help clarify the impact of feedback on sensory estimation in this task.)

Nevertheless, the reduction in serial dependence in 3D motion reports observed in the

current data set suggests that serial dependence may be a strategy employed

specifically when the number or reliability of available sensory signals is low, supporting

variable serial dependence across task conditions and trials within the same task

context.

Feedback may also play a role in improving participants' understanding of the task

structure. Whereas real-world stimuli are often stable over various time scales such that

taking information from the recent past into consideration when making current

decisions could be adaptive (Fischer & Whitney, 2014; Kiyonaga et al., 2017; Braun et

al., 2018), psychophysical stimuli, such as those in the current study, are often

independent and (pseudo-)randomly presented by design. Serial dependence for

unrelated stimuli is clearly suboptimal from an experimental point of view, but until

participants are made aware of their mistakes, generalizing from experience provides a

way of overcoming uncertainty. Indeed, we showed previously that participants in the

no-feedback group exhibited a bias to report high contrast targets as approaching and

reduced contrast targets as receding (Fulvio & Rokers, 2017), consistent with a dimmer-

is-further-away response heuristic (Cooper & Norcia, 2014). In the current results, we

show that serial dependence in the reports of the no-feedback group was significantly

greater for reduced target contrast targets, thus supporting serial dependence as a

behavioral strategy to overcome sensory uncertainty.

In conclusion, our results provide novel evidence supporting the notion that serial

dependence can emerge as the result of a high level strategy employed to overcome

sensory uncertainty. Our results are in line with the idea that serial dependence may

reflect a flexible perceptual mechanism that is not limited to sensory components of

external stimuli, but instead exploits all available information (Fornaciai & Park, 2022),

including external feedback-based error signals.

Citation diversity statement

To promote transparency surrounding citation practice (Dworkin et al. 2019; Zurn et al.

2020) and mitigate biases leading to under-citation of work led by women relative to

other papers as demonstrated across several scientific domains (e.g., Dworkin et al.

2019; Maliniak et al. 2013; Caplar et al. 2017; Fulvio et al. 2021), we proactively aimed

to include references that reflect the diversity of the field and quantified the gender

breakdown of citations in this article according to the first names of the first and last

authors using the Gender Citation Balance Index web tool

(https://postlab.psych.wisc.edu/gcbialyzer/). This article contains 50.0% man/man,

12.5% man/woman, 26.6% woman/man and 10.9% woman/woman citations. For

comparison, proportions estimated from articles in five prominent neuroscience journals

(as reported in Dworkin et al. 2019) are 58.4% man/man, 9.4% man/woman, 25.5%

woman/man and 6.7% woman/woman. Note that the estimates may not always reflect

gender identity and do not account for intersex, non-binary, or transgender individuals.

References

Alais, D., Leung, J., & van der Burg, E. (2017). Linear summation of repulsive and

attractive serial dependencies: Orientation and motion dependencies sum in motion

perception. *Journal of Neuroscience, 37*, 4381-4390.

https://doi.org/10.1523/JNEUROSCI.4601-15.2017

Alexi, J., Cleary, D., Dommisse, K., Palermo, R., Kloth, N., Burr, D., & Bell, J. (2018).

Past visual experiences weigh in on body size estimation. *Scientific reports, 8*(1), 1-8.

https://doi.org/10.1038/s41598-017-18418-3

Bae, G. Y., & Luck, S. J. (2020). Serial dependence in vision: Merely encoding the

previous-trial target is not enough. *Psychonomic bulletin & review, 27*(2), 293-300.

https://doi.org/10.3758/s13423-019-01678-7

Barbosa, J., & Compte, A. (2020). Build-up of serial dependence in color working

memory. *Scientific reports, 10*(1), 1-7. https://doi.org/10.1038/s41598-020-67861-2

Barbosa, J., Stein, H., Martinez, R. L., Galan-Gadea, A., Li, S., Dalmau, J., ... Compte,

A. (2020). Interplay between persistent activity and activity-silent dynamics in the

prefrontal cortex underlies serial biases in working memory. *Nature Neuroscience,*

23(8), 1016–1024. https://doi.org/10.1038/s41593-020-0644-4

Berens, P. (2009). CircStat: A Matlab Toolbox for Circular Statistics, *Journal of*

Statistical Software, *(31)*10. http://www.jstatsoft.org/v31/i10

Bliss, D. P., Sun, J. J., & D'Esposito, M. (2017). Serial dependence is absent at the time

of perception but increases in visual working memory. *Scientific Reports, 7,* 1-13.

https://doi.org/10.1038/s41598-017-15199-7

Braun, A., Urai, A. E., & Donner, T. H. (2018). Adaptive history biases result from

confidence-weighted accumulation of past choices. *Journal of Neuroscience, 38(*10),

2418-2429. https://doi.org/10.1523/JNEUROSCI.2189-17.2017

Burr, D., & Cicchini, G. M. (2014). Vision: efficient adaptive coding. *Current Biology,*

24(22), R1096-R1098. https://doi.org/10.1016/j.cub.2014.10.002

Caplar, N., Tacchella, S., & Birrer, S. (2017). Quantitative evaluation of gender bias in

astronomical publications from citation counts. *Nature astronomy, 1*(6), 1-5.

https://doi.org/10.1038/s41550-017-0141

Ceylan, G., Herzog, M. H., & Pascucci, D. (2021). Serial dependence does not originate

from low-level visual processing. *Cognition, 212*, 104709, 1-7.

https://doi.org/10.1016/j.cognition.2021.104709

Chang, T. Y., Thompson, L., Doudlah, R., Kim, B., Sunkara, A., & Rosenberg, A.

(2020). Optimized but not maximized cue integration for 3D visual perception. *eNeuro*,

7(1), 1-18.

https://doi.org/10.1523/ENEURO.0411-19.2019

Cicchini, G. M., Anobile, G., & Burr, D. C. (2014). Compressive mapping of number to

space reflects dynamic encoding mechanisms, not static logarithmic transform.

Proceedings of the National Academy of Sciences, *111*(21), 7867-7872.

https://doi.org/10.1073/pnas.1402785111

Cicchini, G. M., Mikellidou, K., & Burr, D. (2017). Serial dependencies act directly on

perception. *Journal of vision, 17*(14):6, 1-9. https://doi.org/10.1167/17.14.6

Cicchini, G. M., Mikellidou, K., & Burr, D. (2018). The functional role of serial

dependence. *Proceedings of the Royal Society B: Biological Sciences, 285*(1890), 1-8.

https://doi.org/10.1098/rspb.2018.1722

Cicchini, G. M., Benedetto, A., & Burr, D. C. (2021). Perceptual history propagates

down to early levels of sensory analysis. *Current Biology*, *31*(6), 1245-1250.

https://doi.org/10.1016/j.cub.2020.12.004

Clifford, C. W., Watson, T. L., & White, D. (2018). Two sources of bias explain errors in

facial age estimation. *Royal Society open science, 5*(10), 1-10.

https://doi.org/10.1098/rsos.180841

Collins, T. (2019). The perceptual continuity field is retinotopic. *Scientific Reports,* 9,

18841, 1-6. https://doi.org/10.1038/s41598-019-55134-6

Collins, T. (2020). Serial dependence alters perceived object appearance. *Journal of*

Vision, *20*(13):9, 1-8. https://doi.org/10.1167/jov.20.13.9

Collins, T. (2021). Serial dependence occurs at the level of both features and integrated

object representations. *Journal of Experimental Psychology: General, 151*(8), 1821–

1832. https://doi.org/10.1037/xge0001159

Cooper, E. A., & Norcia, A. M. (2014). Perceived depth in natural images reflects

encoding of low-level luminance statistics. *Journal of neuroscience*, *34*(35), 11761-

11768. https://doi.org/10.1523/JNEUROSCI.1336-14.2014

de Azevedo Neto, R. M., & Bartels, A. (2021). Disrupting short-term memory

maintenance in premotor cortex affects serial dependence in visuomotor integration.

Journal of Neuroscience, 41(45), 9392-9402.

https://doi.org/10.1523/JNEUROSCI.0380-21.2021

Dworkin, J. D., Linn, K. A., Teich, E. G., Zurn, P., Shinohara, R. T., & Bassett, D. S.

(2020). The extent and drivers of gender imbalance in neuroscience reference lists.

Nature neuroscience, 23(8), 918-926. https://doi.org/10.1038/s41593-020-0658-y

Fischer, J., & Whitney, D. (2014). Serial dependence in visual perception. *Nature*

Neuroscience, 17, 738- 743. https://doi.org/10.1038/nn.3689

Fornaciai, M., & Park, J. (2018). Attractive serial dependence in the absence of an

explicit task. *Psychological Science, 29*, 437-446.

https://doi.org/10.1177/0956797617737385

Fornaciai, M., & Park, J. (2020). Attractive serial dependence between memorized

stimuli. *Cognition, 200*, 104250, 1-11. https://doi.org/10.1016/j.cognition.2020.104250

Fornaciai, M., & Park, J. (2022). The effect of abstract representation and response

feedback on serial dependence in numerosity perception. *Attention, Perception, &*

Psychophysics, 84, 1-15. https://doi.org/10.3758/s13414-022-02518-y

Fritsche, M. (2016). To Smooth or not to Smooth: Investigating the Role of Serial

Dependence in Stabilizing Visual Perception [Masters Dissertation, Radboud

University], https://theses.ubn.ru.nl/handle/123456789/3193

Fritsche, M., Mostert, P., & de Lange, F. P. (2017). Opposite effects of recent history on

perception and decision. *Current Biology, 27*, 590-595.

https://doi.org/10.1016/j.cub.2017.01.006

Fritsche, M., & de Lange, F. P. (2019). The role of feature-based attention in visual

serial dependence. *Journal of vision, 19*(13):21, 1-13. https://doi.org/10.1167/19.13.21

Fründ, I., Wichmann, F. A., & Macke, J. H. (2014). Quantifying the effect of intertrial

dependence on perceptual decisions. *Journal of Vision, 14*(7):9, 1-16,

https://doi.org/10.1167/14.7.9

944 Fulvio, J. M., Rosen, M. L., & Rokers, B. (2015). Sensory uncertainty leads to

945 systematic misperception of the direction of motion in depth. *Attention, Perception, &*

946 *Psychophysics*, *77*(5), 1685-1696. https://doi.org/10.3758/s13414-015-0881-x

948 Fulvio, J.M. & Rokers, B. (2017). Use of cues in virtual reality depends on visual

949 feedback. *Scientific reports, 7*(1), 1-13. https://doi.org/10.1038/s41598-017-16161-3

951 Fulvio, J.M., Ji, M., Thompson, L., Rosenberg, A. & Rokers, B. (2020). Cue-dependent

952 effects of VR experience on motion-in-depth sensitivity. *Plos One, 15*(3), e0229929, 1-

953 14. https://doi.org/10.1371/journal.pone.0229929

955 Fulvio, J. M., Akinnola, I., & Postle, B. R. (2021). Gender (im) balance in citation

956 practices in cognitive neuroscience. *Journal of Cognitive Neuroscience, 33*(1), 3-7.

957 https://doi.org/10.1162/jocn_a_01643

959 Gallagher, G. K., & Benton, C. P. (2022). Stimulus uncertainty predicts serial

960 dependence in orientation judgements. *Journal of Vision, 22*(1):6, 1-14.

961 https://doi.org/10.1167/jov.22.1.6

963 Galluzzi, F., Benedetto, A., Cicchini, G. M., & Burr, D. C. (2022). Visual priming and

964 serial dependence are mediated by separate mechanisms. *Journal of Vision, 22*(10):1,

965 1-12. https://doi.org/10.1167/jov.22.10.1

967 Girshick, A. R., Landy, M. S., & Simoncelli, E. P. (2011). Cardinal rules: visual

968 orientation perception reflects knowledge of environmental statistics. *Nature*

969 *neuroscience*, *14*(7), 926-932. https://doi.org/10.1038/nn.2831

971 Goettker, A., & Stewart, E. E. (2022). Serial dependence for oculomotor control

972 depends on early sensory signals. *Current Biology, 32*, 2956–2961.

973 https://doi.org/10.1016/j.cub.2022.05.011

Harris, J. M., & Dean, P. J. A. (2003). Accuracy and precision of binocular 3-D motion

perception. *Journal of Experimental Psychology. Human Perception and Performance,*

29(5), 869–881. https://doi.org/10.1037/0096-1523.29.5.869

Harris, J. M., & Drga, V. F. (2005). Using visual direction in three-dimensional motion

perception. *Nature Neuroscience, 8*(2), 229–233. https://doi.org/10.1038/nn1389

Hillis, J. M., Watt, S. J., Landy, M. S., & Banks, M. S. (2004). Slant from texture and

disparity cues: Optimal cue combination. *Journal of vision, 4*(12):1, 967-992.

https://doi.org/10.1167/4.12.1

Kim, S., Burr, D., Cicchini, G. M., & Alais, D. (2020). Serial dependence in perception

requires conscious awareness. *Current Biology, 30*(6), R257-R258.

https://doi.org/10.1016/j.cub.2020.02.008

Kim, S., & Alais, D. (2021). Individual differences in serial dependence manifest when

sensory uncertainty is high. *Vision Research, 188*, 274-282.

https://doi.org/10.1016/j.visres.2021.08.001

Kiyonaga, A., Scimeca, J. M., Bliss, D. P., & Whitney, D. (2017). Serial Dependence

across Perception, Attention, and Memory. *Trends in Cognitive Sciences, 21*(7), 493-

497. https://doi.org/10.1016/j.tics.2017.04.011

Knill, D. C. (2007). Robust cue integration: A Bayesian model and evidence from cue-

conflict studies with stereoscopic and figure cues to slant. *Journal of vision, 7*(7):5, 1-24.

https://doi.org/10.1167/7.7.5

1001

1002   Knill, D. C., & Richards, W. (Eds.). (1996). *Perception as Bayesian inference.*

1003   Cambridge University Press.

1004

Knill, D. C., & Saunders, J. A. (2003). Do humans optimally integrate stereo and texture information for judgments of surface slant?. *Vision research*, *43*(24), 2539-2558. https://doi.org/10.1016/S0042-6989(03)00458-9

Lages, M. (2006). Bayesian models of binocular 3-D motion perception. *Journal of Vision, 6*(4), 508–522. https://doi.org/10.1167/6.4.14

Liberman, A., Fischer, J., & Whitney, D. (2014). Serial dependence in the perception of faces. *Current Biology, 24*, 2569-2574. https://doi.org/10.1016/j.cub.2014.09.025

Liberman, A., Manassi, M., & Whitney, D. (2018). Serial dependence promotes the stability of perceived emotional expression depending on face similarity. *Attention, Perception, & Psychophysics, 80*(6), 1461-1473. https://doi.org/10.3758/s13414-018-1533-8

Lidström, A. (2019). Visual Uncertainty in Serial Dependence: Facing Noise [Masters Dissertation, Kristianstad University], http://urn.kb.se/resolve?urn=urn:nbn:se:hkr:diva-19356

Maliniak, D., Powers, R., & Walter, B. F. (2013). The gender citation gap in international relations. *International Organization, 67*(4), 889-922. https://doi.org/10.1017/S0020818313000209

Manassi, M., Liberman, A., Kosovicheva, A., Zhang, K., & Whitney, D. (2018). Serial dependence in position occurs at the time of perception. *Psychonomic bulletin & review, 25*(6), 2245-2253. https://doi.org/10.3758/s13423-018-1454-5

Manassi, M., & Whitney, D. (2022). Illusion of visual stability through active perceptual serial dependence. *Science advances, 8*(2), eabk2480, 1-8. https://doi.org/10.1126/sciadv.abk2480

1036 Murai, Y., & Whitney, D. (2021). Serial dependence revealed in history-dependent

1037 perceptual templates. *Current Biology*, *31*(14), 3185-3191.

1038 https://doi.org/10.1016/j.cub.2021.05.006

1039

1040 Murphy, A. P., Ban, H., & Welchman, A. E. (2013). Integration of texture and disparity

1041 cues to surface slant in dorsal visual cortex. *Journal of Neurophysiology*, *110*(1), 190-

1042 203. https://doi.org/10.1152/jn.01055.2012

1043

1044 Oruç, I., Maloney, L. T., & Landy, M. S. (2003). Weighted linear cue combination with

1045 possibly correlated error. *Vision research*, *43*(23), 2451-2468.

1046 https://doi.org/10.1016/S0042-6989(03)00435-8

1047

1048 Pascucci, D., Mancuso, G., Santandrea, E., Della Libera, C., Plomp, G., & Chelazzi, L.

1049 (2019). Laws of concatenated perception: Vision goes for novelty, decisions for

1050 perseverance. *PLoS biology*, *17*(3), e3000144.

1051 https://doi.org/10.1371/journal.pbio.3000144

1052

1053 Pascucci, D., & Plomp, G. (2021). Serial dependence and representational momentum

1054 in single-trial perceptual decisions. *Scientific Reports, 11*(1), 1-8.

1055 https://doi.org/10.1038/s41598-021-89432-9

1056

1057 Preston, T. J., Kourtzi, Z., & Welchman, A. E. (2009). Adaptive estimation of three-

1058 dimensional structure in the human brain. *Journal of Neuroscience*, *29*(6), 1688-1698.

1059 https://doi.org/10.1523/JNEUROSCI.5021-08.2009

1060

1061 Rahnev, D., Koizumi, A., McCurdy, L. Y., D'Esposito, M., & Lau, H. (2015). Confidence

1062 leak in perceptual decision making. *Psychological science*, *26*(11), 1664-1680.

1063 https://doi.org/10.1177/0956797615595037

1064

1065 Rideaux, R., & Welchman, A. E. (2018). Proscription supports robust perceptual

1066 integration by suppression in human visual cortex. *Nature Communications*, *9*(1), 1-14.

1067 https://doi.org/10.1038/s41467-018-03400-y

1068

1069 Rokers, B., Fulvio, J.M., Pillow, J., Cooper, E.A. (2018). Systematic misperceptions of

1070 3D motion explained by Bayesian inference. *Journal of Vision*, *18*(3), 1-23.

1071 https://doi.org/10.1167/jov.18.3.23

1072

1073 Rushton, S. K., & Duke, P. A. (2007). The use of direction and distance information in

1074 the perception of approach trajectory. *Vision Research, 47*(7), 899–912.

1075 https://doi.org/10.1016/j.visres.2006.11.019

1076

1077 Samaha, J., Switzky, M., & Postle, B. R. (2019). Confidence boosts serial dependence

1078 in orientation estimation. *Journal of Vision, 19*(4):25, 1-13.

1079 https://doi.org/10.1167/19.4.25

1080

1081 Sheehan, T. C., & Serences, J. T. (2022). Attractive serial dependence overcomes

1082 repulsive neuronal adaptation. *PLoS biology*, *20*(9), e3001711, 1-31.

1083 https://doi.org/10.1371/journal.pbio.3001711

1084

1085 Suárez-Pinilla, M., Seth, A. K., & Roseboom, W. (2018). Serial dependence in the

1086 perception of visual variance. *Journal of Vision, 18*(7):4, 1-24,

1087 https://doi.org/10.1167/18.7.4

1088

1089 Trommershäuser, J., Konrad K., & Landy, M.S. (eds), Sensory Cue Integration,

1090 *Computational Neuroscience Series* (2011; online edn, Oxford Academic, 20 Sept.

1091 2012). https://doi.org/10.1093/acprof:oso/9780195387247.001.0001

1092

1093 Welchman, A. E., Tuck, V. L., & Harris, J. M. (2004). Human observers are biased in

1094 judging the angular approach of a projectile. *Vision Research, 44*(17), 2027–2042.

1095 https://doi.org/10.1016/j.visres.2004.03.014

1096

1097 Welchman, A. E., Deubelius, A., Conrad, V., Bülthoff, H. H., & Kourtzi, Z. (2005). 3D

1098 shape perception from combined depth cues in human visual cortex. *Nature*

1099 *neuroscience, 8*(6), 820-827. https://doi.org/10.1038/nn1461

1100

1101 Welchman, A. E., Lam, J. M., & Bülthoff, H. H. (2008). Bayesian motion estimation

1102 accounts for a surprising bias in 3D vision. *Proceedings of the National Academy of*

1103 *Sciences of the United States of America, 105*(33), 12087–12092.

1104 https://doi.org/10.1073/pnas.0804378105

1105

1106 Xia, Y., Leib, A. Y., & Whitney, D. (2016). Serial dependence in the perception of

1107 attractiveness. *Journal of vision, 16*(15), 28-28. https://doi.org/10.1167/16.15.28

1108

1109 van der Burg, E., Rhodes, G., & Alais, D. (2019). Positive sequential dependency for

1110 face attractiveness perception. *Journal of vision, 19*(12):6, 1-16.

1111 https://doi.org/10.1167/19.12.6

1112

1113 van Bergen, R., & Jehee, J. (2017). Uncertainty in cortical stimulus representations

1114 predicts serial dependence effects in orientation perception. *Journal of Vision, 17*(10):

1115 590. https://doi.org/10.1167/17.10.590

1116

1117 van Bergen, R. S., & Jehee, J. F. (2019). Probabilistic representation in human visual

1118 cortex reflects uncertainty in serial decisions. *Journal of Neuroscience, 39*(41), 8164-

1119 8176. https://doi.org/10.1523/JNEUROSCI.3212-18.2019

1120

1121 van Snellenberg, J.X. (2018). BCa_bootstrap

1122 (https://www.mathworks.com/matlabcentral/fileexchange/69119), MATLAB Central File

1123 Exchange. Retrieved August, 2021.

1124

1125 Wang, X. Y., Gong, X. M., Sun, Q., & Li, X. (2022). Attractive effects of previous form

1126 information on heading estimation from optic flow occur at perceptual stage. *Journal of*

1127 *Vision*, *22*(12):18, 1-15. https://doi.org/10.1167/jov.22.12.18

1128

1129 Weiss, Y., Simoncelli, E. P., & Adelson, E. H. (2002). Motion illusions as optimal

1130 percepts. *Nature neuroscience*, *5*(6), 598-604. https://doi.org/10.1038/nn0602-858

1131

1132 Zurn, P., Bassett, D. S., & Rust, N. C. (2020). The citation diversity statement: a

1133 practice of transparency, a way of life. *Trends in Cognitive Sciences, 24(*9), 669-672.

1134 https://doi.org/10.1016/j.tics.2020.06.009

1135

1136

1137