Title: The effects of explicit and implicit information on modulation of corticospinal excitability during

hand-object interactions.

Running title: Effects of sensorimotor information on corticospinal excitability during grasping

Guy Rens[1,2], Marco Davare[3,4], Vonne van Polanen[2,5]

[1]The Brain and Mind Institute, University of Western Ontario, London, Ontario N6A 3K7, Canada.

[2]KU Leuven, Leuven Brain Institute, 3001 Leuven, Belgium

[3]Department of Clinical and Movement Neurosciences, UCL Queen Square Institute of Neurology,

University College London, London, United Kingdom

[4]Faculty of Life Sciences and Medicine, King's College London, London SE1 1UL, United Kingdom.

[5]Movement Control and Neuroplasticity Research Group, Department of Movement Sciences, Biomedical

Sciences group, KU Leuven, 3001 Leuven, Belgium

*Corresponding Author:

Guy Rens

The Brain and Mind Institute

University of Western Ontario

Ontario N6A 3K7, Canada

grens@uwo.ca

Funding statement: This work was funded by a Research Foundation Flanders (FWO) Odysseus Project

(Fonds Wetenschappelijk Onderzoek, Belgium: G/0C51/13N) awarded to MD and 12X7118N/Research

Foundation Flanders (FWO) awarded to VVP.

Acknowledgements: The authors would like to thank Immele Meeusen and Hannah De Buck for their help

in the data collection of experiment 1 and 2, respectively. We also would like to thank René Clerckx for

programming the force tracking task in experiment 1.

Conflict of interest: The authors declare to have no conflict of interest.

Abstract

Fingertip force scaling during hand-object interactions typically relies on visual information about the

object and sensorimotor memories from previous object interactions. Here, we investigated whether

contextual information, that is not explicitly linked to the intrinsic object properties (e.g., size or weight)

but is informative for motor control requirements, can mediate force scaling. For this, we relied on two

separate behavioral tasks during which we applied transcranial magnetic stimulation (TMS) to probe

corticospinal excitability (CSE), as a window onto the primary motor cortex role in controlling fingertip

forces. In experiment 1, participants performed a force tracking task, where we manipulated available

implicit and explicit visual information. That is, either the force target was fully visible, or only the force

error was displayed as a deviation from a horizontal line. We found that participants' performance was

better when the former condition, in which they had explicit access to predictive information. However,

we did not find differences in CSE modulation based on the type of visual information. On the other hand,

CSE was modulated by the change in muscle contraction, i.e., contraction vs. relaxation and fast vs. slow

changes. In sum, these findings indicate that CSE only reflects the ongoing motor command. In experiment

2, other participants performed a sequential object lifting task of visually identical objects that were

differently weighted, in a seemingly random order. Within this task, we hid short series of incrementally

increasing object weights. This allowed us to investigate whether participants would scale their forces for

specific object weights based on the previously lifted object (i.e., sensorimotor effect) or based on the

implicit information about the hidden series of incrementally increasing weights (i.e., extrapolation

beyond sensorimotor effects). Results showed that participants did not extrapolate fingertip forces based

on the hidden series but scaled their forces solely on the previously lifted object. Unsurprisingly, CSE was

not modulated differently when lifting series of random weights versus series of increasing weights.

Altogether, these results in two different grasping tasks suggest that CSE encodes ongoing motor

components but not sensorimotor cues that are hidden within contextual information.

Highlights

- Explicit visual cues and sensorimotor experience are key for object grasping
- Investigating influence of implicit, contextual information on object grasping
- Explicit but not implicit cues improve motor performance during object grasping
- Explicit but not implicit cues modulate corticospinal excitability

1. Introduction

Skilled hand object interactions require predictive planning of fingertip forces adjusted to the properties of the object (Gordon et al., 1993) as, for instance, a fragile object may be damaged when exerting too much force. Conventionally, it has been argued that individuals estimate object weight based on two mechanisms. First, by relying on visual cues and well-established "priors", such as the size-weight (Gordon et al., 1991) and density-weight relationships (Baugh et al., 2012), object weight can be estimated. Second, when visual cues are not indicative of object weight (e.g., a suitcase that might be empty or filled) then individuals rely on a "sensorimotor memory". This sensorimotor memory stores short-term information about object weight based on previous interactions with the same or similar objects (Baugh et al., 2012; Fu et al., 2010; Gordon et al., 1993; Johansson & Westling, 1988; Vonne van Polanen & Davare, 2015).

Interestingly, Cashaback et al. (2017) demonstrated that predictive object lifting not only relies on visual cues or sensorimotor memories, but also incorporates probabilistic information. In their study, participants minimized the potential prediction error across many lifts instead of optimizing their fingertip forces for the mostly anticipated object. In addition, Mawase & Karniel (2010) argued that an individual's sensorimotor system can use information the individual is not aware of: The authors showed that individuals predictively scale their fingertip forces to a non-existent heavier weight after lifting a series of increasing weights that were implicitly presented within larger sequences primarily consisting of random weights. Other studies on hand actions have supported the notion that motor control mechanisms can adapt their sensorimotor predictions towards environmental variability for minimizing the general prediction error across multiple trials (e.g., Hadjiosif & Smith, 2015; Körding et al., 2004; Landy et al., 2012; Zhang et al., 2015). Importantly, it has been argued that finetuning motor commands to environmental variability relies both on explicit (i.e., conscious strategies) and implicit (i.e., unaware/subconscious adaptation) learning mechanisms (Albert et al., 2022). Notably, studies teasing apart explicit and implicit components of sensorimotor learning typically rely on reaching, and not grasping (e.g., Wolpert & Flanagan, 2016).

While it seems likely that sensorimotor learning during grasping relies on similar mechanisms as reaching due to the interconnectedness of their respective neural pathways (Gerbella et al., 2017), performance errors are typically perceived differently between reaching and grasping. During reaching, performance errors can be perceived by a visual mismatch between the target position and the actual end-position. During grasping, lifting errors are only perceived through a haptic mismatch between the expected and the actual object weight (Castiello, 2005; Johansson & Westling, 1984).

Due to (i) these differences between grasping and reaching, (ii) most of the scientific literature

relying on reaching rather than grasping to understand implicit learning (for instance see Albert et al.,

2022; Heald et al., 2021) and (iii) grasping studies typically relying on explicit visual cues to indicate object

weight (e.g., Buckingham et al., 2014; Loh et al., 2010; Rens et al., 2021a), it is still unclear how the brain

encodes implicit information for grasping. While it is plausible that the cortical grasping network (Davare

et al., 2011; Gerbella et al., 2017) primarily relies on explicit visual cues and sensorimotor memories, it

could still benefit from implicit learning. For instance, when lifting many opaque objects with varying

weights (e.g., baggage handlers) it seems preferable to minimize the average prediction error than to try

to predict each object's weight (as in the study of Cashaback et al., 2017).

Importantly, the primary motor cortex (M1) plays a pivotal role in controlling hand-object

interactions. When disrupting M1 activity transiently with repetitive Transcranial Magnetic Stimulation

(rTMS), individuals fail to predictively scale their fingertip forces based on sensory feedback (Chouinard

et al., 2005; Parikh et al., 2020; Schabrun et al., 2008). In addition, Loh et al. (2010) showed that

corticospinal excitability (CSE), probed with TMS over M1 during lift planning, reflects the weight of the

previously lifted object but becomes reflective of a forthcoming weight when indicated by a visual cue. In

previous work from our group, we showed that CSE modulation can reflect an object's weight distribution

when indicated through action observation (Rens et al., 2021b). Arguably, these studies (Loh et al., 2010;

Rens et al., 2021b) and similar ones (e.g., Baugh et al., 2012; Fu et al., 2010; for a review see Johansson &

Flanagan, 2009) investigated how task requirements, indicated by explicit visual cues or sensorimotor

memories are represented in corticospinal outputs during hand-object interactions.

To our knowledge, it is not well understood how CSE encodes hand-object interactions when task

requirements are only implicitly present. For instance, in the study of Mawase & Karniel (2010), the

question arises whether CSE modulation would represent force predictions based on an implicit series of

increasing weights or the last lifted weight as in study of Loh et al. (2010). Noteworthy, using different

behavioral modalities, CSE has been shown to be modulated by implicit (probabilistic) information. For

instance, using a reaction time task, Van Elswijk et al. (2007) showed that CSE modulation increases with

increasing response signal expectancy. That is, the longer it takes for the cue to appear, the more likely it

is to appear, which increases CSE modulation. This finding was replicated by Dupin et al. (2019). Using a

similar reaction time paradigm, Bestmann et al. (2008) showed that CSE modulation increases in case of

low entropy (i.e., low uncertainty about sampling one trial from all possible trials) and low surprise (i.e.,

probability of the upcoming trial on itself). In another study, Senot et al. (2011) argued that, during lift

observation, CSE is modulated by both written labels (i.e., explicit cues) and movement kinematics (i.e.,

implicit cues) indicating object weight. Similar results were found in another action observation study by

Puglisi et al. (2017): CSE was modulated by explicit visual cues but also by implicit visual cues, i.e., in the

authors' words: "visual cues that were not the focus of attention".

In the present study, we aimed to extend on studies investigating how environmental uncertainty

affects grasping (see Cashaback et al., 2017; Hadjiosif & Smith, 2015; Körding et al., 2004; Landy et al.,

2012) and explore how the sensorimotor system can integrate implicit information for grasping. For this,

we used two separate experiments to investigate the integration of implicit information during

continuous force exertion by squeezing (experiment 1) and during discrete object lifting (experiment 2).

In addition to these behavioral tasks, we included TMS to probe CSE and investigate the underlying motor

cortical mechanisms.

For experiment 1 we used a force tracking task and manipulated the type of visual information

participants received about their ongoing motor performance. Participants were instructed to track a

sinusoidal force curve by exerting increasing and decreasing force at different rates of change (slope of

the curve) with their fingertips on force sensors. The type of the visual information was varied by

displaying either the actual sinusoidal curve or a horizontal line. Importantly, in the sinusoidal curve

condition, participants could rely on both predictive mechanisms, by observing the curve's slope, and

feedback-driven mechanisms, by observing their performance error (i.e., distance between the target

curve and the line representing their performance). If participants would perform the task perfectly, their

performance would overlap with the force target (sinusoidal curve). If they would not generate force at

all, their performance would show as a horizontal line. In this way, we argued that both prediction- and

feedback-related visual cues would be explicitly available. In contrast, in the horizontal line condition,

participants tracked the same sinusoidal curve although the actual force target was hidden from the

participants. Instead, they could only see their performance error (i.e., deviation from the sinusoidal

curve) as deviations from a horizontal line. Thus, in this condition, if participants would perform the

sinusoidal force tracking task perfectly, their performance would show as a horizontal line. Conversely, if

participants would not exert force at all, their performance would be represented as the inverse of the

sinusoidal target curve. As such, we argued that in this horizontal line condition, participants only had

explicit access to their performance errors: they could only implicitly deduct their location on the hidden

sinusoidal curve as (a) the force scaling requirements (curve slope) was not visible and (b) the performance

error was shown in an inverse manner. Finally, TMS was applied to probe CSE during different phases

(speed: fast or slow rate of change; target line: sinusoidal or horizontal; direction: up- or downscaling of

fingertip forces). We argued that by manipulating the type of visual information, we could investigate how

explicit and implicit information affects CSE modulation during visuomotor control of continuous hand-
object interactions.

Using a similar paradigm, Kimura et al. (2003) asked participants to contract and relax their elbow
in a sinusoidal manner. They found that CSE increased during the contraction compared to relaxation
phase, even though TMS was applied when the same amount of force was applied in both phases. In line
with Kimura et al. (2003), we hypothesised that, in both conditions, CSE would be larger during increasing
compared to decreasing force exertion. To extend on the work of Kimura et al. (2003), we also
manipulated the slope of the sinusoidal curve, resulting in slower or faster scaling of fingertip forces, to
investigate sensorimotor involvement in the rate of force changes. We hypothesized that CSE modulation
would be larger in the fast compared to slow condition. Importantly, if CSE modulation reflects ongoing
task performance, independent of the type of visual information, CSE would be modulated similarly in the
sinusoidal curve and horizontal line conditions. However, Bestmann et al. (2008) showed that CSE
modulation is larger during low uncertainty. Accordingly, we hypothesized that, due to differences in the
type of visual information between the target line conditions, CSE modulation would be larger when
contextual information was more explicitly present, i.e., in the sinusoidal curve condition.

For experiment 2 we used an object lifting task that was similar to Mawase & Karniel (2010).
Participants lifted small objects that were visually identical but had different weights. The total series of
all lifted objects consisted of intermixed series of randomly ordered weights and series of increasing
weights. Briefly, Mawase & Karniel (2010) showed that individuals predictively scale their fingertip forces
to a non-existent heavier weight after lifting a series of increasing weights when this series was hidden
within larger series consisting of random weights. Notably, in their study participants indicated during
debriefing that they were unaware of these hidden series of increasing weights. As such, the authors
argued that participants implicitly embodied the series of increasing weights, leading to extrapolation
during predictive force planning. In this experiment, we argued that sensorimotor control mechanisms
could extrapolate force requirements after the series of increasing weights, which would be reflected in
changes in corticospinal output. That is, motor performance and CSE would not reflect a direct encoding
of force scaling based on the last trials (i.e., the sensorimotor memory) but rather would reflect an
extrapolation process for predicting force scaling based on (the implicit information) related to the series
of increasing weight. As such, we hypothesized that participants would use more force when lifting the
same weight after the implicit series compared to after the random series. Similarly, we hypothesized that
CSE would reflect the extrapolated weight based on the series of increasing weights and not the weight
of the previously lifted object (akin to Loh et al., 2010).

2. **General Methods**

<u>2.1 Participants</u>

Thirty healthy right-handed volunteers were recruited for the experiment, of which 15 took part in
experiment 1 (12 females, mean age 22±2 years) and 15 in experiment 2 (11 females, mean age 22±3
years). Two participants (one for each experiment) were excluded from analyses, causing us to have 14
participants for each experiment. The first one (female) was excluded during data processing as the motor
evoked potentials (MEPs; see below) during the experiment (and thus after defining stimulation intensity)
exceeded the range of our electromyography (EMG) system. The second participant (female) was
excluded as even during baseline assessment of CSE she was not able to relax their hand.

All participants were right-handed (self-reported). TMS eligibility was assessed by the TMS safety
screening questionnaire (Rossi et al., 2011). Participants had no known neurological or sensory
impairments and had normal or corrected-to-normal vision. The study was approved by the ethical
committee of UZ/KU Leuven and all participants signed an informed consent before the experiment.

<u>2.2 Acquisition of force data</u>

Forces were measured using a grip(-lift) manipulandum (see Figure 1). The manipulandum consisted of
two force/torque sensors (ATI Nano17, ATI Industrial Automation) that were mounted on a custom-made
carbon fiber basket. The total weight of the manipulandum was 1.2 N. The graspable surface (17 mm
diameter and 45 mm apart) of the force sensors was covered with fine sandpaper (P600) to increase
friction. In the manipulandum's basket, small objects could be placed. Force sensor data was sampled at
1 kHz frequency through a NI-USB 6343X (National Instruments, USA) which was connected to a personal
computer. Measurement of forces in both experiments was performed using custom written Labview
scripts (National Instruments, USA) and transferred for off-line analysis.

Figure 1. *The (grip-lift) manipulandum consisting of force sensors mounted on a carbon fibre basket as used in the*
experiments. In experiment 1, participants squeezed the force sensors to track a curve on a computer screen. In
experiment 2, small visually identical cubes with different weights were placed in the basket to be lifted by the
participants.
222

<u>2.3 Transcranial magnetic stimulation procedures</u>

For both experiments, two Ag-Al electrodes were placed on the first dorsal interosseus muscle (FDI)

in a typical belly-tendon montage. A third reference electrode was placed on the ulnar styloid processus.

Electrodes were connected to a NL824 AC preamplifier (Digitimer) and a NL820A isolation amplifier

(Digitimer), which in turn was connected to a micro140-3 CED (Cambridge Electronic Design). The CED was

connected to a personal computer. Electromyography (EMG) recordings were amplified with a gain of

1000, high pass filtered with a frequency of 3 Hz, sampled using Signal and Spike2 software (Cambridge

Electronic Design, UK) and were stored for off-line analysis. Spike2 was used for data collection in

experiment 1 and Signal was used for data collection in experiment 2. The sample frequency was 1 kHz

and 5 kHz in Experiment 1 and 2, respectively. A 70 mm figure-of-eight TMS coil [experiment 1: Magstim

rapid 200 (Magstim Co Ltd, United Kingdom); experiment 2: DuoMag XT-100 (Deymed Diagnostic, Czech

Republic)] was used to apply TMS. In both experiments, participants received TMS over M1 only. For M1

stimulation, the coil was tangentially placed over the head to induce a posterior-anterior current flow and

to elicit motor evoked potentials (MEPs) in the FDI muscle. Before the experiment, we defined the optimal

stimulation site (i.e., 'hotspot') and the stimulation intensity to be used in the experimental task for each

participant. The hotspot was defined as the position from which MEPs with maximal amplitude were

produced in the FDI muscle. The hotspot was marked on the scalp. The stimulation intensity (1mV

threshold) was defined as the lowest stimulation intensity that produced MEPs equal to or greater than 1

mV in the FDI muscle in at least five out of ten consecutive trials when stimulating at the predetermined

hotspot.

Specific procedures for experiment 1. The behavioural task of experiment 1 consisted of continuous

force production (i.e., 'squeezing' on the F/T sensors; see below). We assessed the 1 mV threshold while

participants produced a constant total grip force of 2.5 N (sum of horizontal forces applied by thumb and

index finger on the force sensors). As such, the 1 mV threshold was not defined during 'rest' but during

active contraction. The 1 mV threshold for experiment 1 was 42 ± 11 % (group mean ± standard error) of

the maximal stimulator output.

Specific procedures for experiment 2. The behavioural task of experiment 2 consisted of an object-

lifting task and TMS was applied during lift planning (i.e., before the reaching phase, thus while the hand

was still relaxed on the table; see below). Accordingly, the stimulation intensity for experiment 2 was

defined while participants kept their hand relaxed on the table. The 1 mV threshold was 56 ± 8 % (group

mean ± standard error) of the maximum stimulator output.

Figure 2. Drawing of the experimental setup in experiment 1. Participants squeezed the force sensors to track a target curve on the screen with their right hand. During the task, transcranial magnetic stimulation was applied over the left motor cortex and electromyography was measured with electrodes placed on the right first dorsal interosseus muscle.

3. Methods of Experiment 1

<u>3.1. Task and procedures</u>

After initial preparations (i.e., consent, questionnaires and TMS preparations), participants

performed the experimental task. Participants were comfortably seated in a chair. A table was positioned

in front of them on which a computer screen was placed and the grasp-lift manipulandum was placed on

another table on their right side. Participants were instructed to rest their right arm on the table and grasp

the two force sensors of the manipulandum using their right thumb and index finger (precision grip). The

manipulandum itself was fixed on the table to ensure it would not move during the experiment. During

the experiment, the TMS-coil was held above the hotspot by an experimenter (see Figure 2 for a drawing

of the set-up).

Figure 3. *Representative example of a part of a trial in the horizontal line condition.* **A.** *Screen view as shown to the participant during performance of the task in the horizontal line condition. The black line indicates the target line and the red line represents the participant's performance as deviation in exerted force from the target line (in N).* **B.** *Actual performed force (red) and actual target force which was hidden from the participant (blue). The black dashed line represents the force level at which TMS was applied. A TMS event is indicated with a green asterisk. Yellow shadings indicate the area under the curve (AUC) as a measure of performance. Note that in the sinusoidal condition, the screen view looked like the blue (target) and red (performance) lines.* **C.** *Electromyography (EMG) data (black). The time window of 100 ms before TMS that was used to calculate the background EMG (RMS100) is indicated with green vertical lines.*

Sinusoidal curve condition. On the screen a sinusoidal target line was projected with the x- and y-axis representing time and amount of exerted force, respectively (Figure 3B). The target sinusoid had an amplitude of 1.2 N and was centred around 2 N. Given that we wanted to avoid muscle fatigue for the TMS data (Kotan et al., 2015), we decided to have a light muscle contraction (2 N average) and relatively small change in force exertion (1.2 N sinusoidal increase/decrease). The 2 N was arbitrarily defined close to the lightest weights used in our previous studies (e.g., 1.5 N in Rens et al., 2020, and 2.2 N in Van Polanen et al. 2020). The rationale was that we found clear MEPs when participants planned to lift these weights in those studies. As such, we argued that we could investigate a low-intensity muscle contraction

during squeezing (this study) of which we could still anticipate MEP effects based on our previous object-
lifting work. The change in force exertion (1.2 N sinusoidal increase/decrease) was arbitrarily defined as
60 % of 2 N. Given that we applied TMS only above 2 N, task-related MEPs should have been present. A
cursor moved over the target line indicating the participant's location over time. The cursor produced a
red line providing visual feedback of performance over time. Participants controlled the cursor's vertical
displacement by exerting force on the force sensors. Participants could not control the cursor's horizontal
displacement as the cursor moved with a constant speed over the screen. The sinusoid could have a
frequency of either 0.09 ('slow' condition) or 0.25 Hz ('fast' condition) over the positive and negative half
cycles which was indicated by a 'wider' or 'smaller' sinusoidal wave thus causing a more gradual or steep
slope, respectively. The frequency (thus the sinusoid's width) was changed pseudo-randomly when
crossing the midpoint (i.e., when exerting a force equal to 2 N). In this way, four different conditions could
be identified, being fast up (upscaling of forces; 0.25 Hz), fast down (downscaling; 0.25 Hz), slow up
(upscaling; 0.09 Hz) and slow down (downscaling; 0.09 Hz). Our frequencies were based on Kimura et al.
(2003) who used three frequency conditions (slow: 0.08 Hz, medium: 0.17; fast: 0.33Hz). Because we
included different visual conditions (sinusoidal curve vs horizontal line) as well, we decided on two
frequency conditions instead of three. Based on pilot experiments to verify comfortable speeds for our
squeezing task, we chose frequencies of 0.09 and 0.25 Hz. Note that we did not intend to directly compare
our results with Kimura et al. (2003) as we were not particularly interested in the relationship between
MEP amplitude and contraction speed. Instead, we were mainly interested in relative differences (i.e.,
how do MEPs differ between faster and slower contractions) as basis to explore the effects of different
types of visual information. Participants were instructed to stay as close as possible to the target line by
controlling the vertical displacement of the cursor and, thus, by sinusoidally changing the amount of force
they were exerting on the force sensors. As participants could see where they were on the target line,
they could anticipatorily plan to up- or downscale their forces and their speed. Optimal performance in
this task would consist of sinusoidally scaling fingertip forces and would show the red trace (produced by
the cursor's movement) to perfectly overlap with the sinusoidal target line. Last, TMS was applied when
participants exerted 2.5 N (Figure 3B, green asterisk). As such, TMS was only applied during the positive
half-cycle (considering that the sinusoid was centered around 2 N). We argued that we could exclude
stimulating during the negative half cycle. Inclusion of these timings would only allow us to additionally
compare the amounts of exerted force which did not add to our research questions. In addition, excluding
TMS during the negative half cycle allowed us to reduce experimental duration and to minimize potential
confounding effects of muscle fatigue and/or decline in attention. We did decide to stimulate during both

upward (increasing contraction) and downward force scaling (increasing relaxation) during the positive

half cycle to validate our findings with respect to Kimura et al., (2003). In addition, we arbitrarily decided

to stimulate at 2.5 N based on the following rationale: in the fast condition (0.25 Hz), a cycle is completed

in 4 seconds. As such, from the midpoint to the first peak takes approximately 1 second (one fourth of

cycle). If TMS is applied after a 0.5 N increase (± 40 % of maximum increase of 1.2 N), then TMS should be

applied earliest ± 400 ms after reaching the 2 N midpoint in the fast condition. Based on grasping studies

(for a review see: Castiello, 2005), we argued that this would enable participants to integrate feedback if

their performance "faltered" after switching conditions (in particular from slow to fast). To end, the

interval between two TMS pulses was minimally 3.5 seconds. The latency between TMS pulses should not

have resulted in carry-over effects on MEPs (e.g., longer-term inhibition) (Cincotta et al., 2003; Hannah,

2020; Wilson et al., 1993).

Horizontal line condition. The experimental design was nearly identical to the sinusoidal curve

condition. As such, we will only discuss differences. In this task, the sinusoidal target line was not shown

on the screen, but only a horizontal line was shown (Figure 3B). Participants had to perform the same task

of sinusoidally scaling their force. The cursor showed the deviation from the 'hidden' sinusoidal target

curve as a deviation from the horizontal line. Accordingly, participants could not perceive where they were

(e.g., positive, or negative half cycle of sinusoidally scaling their forces), but they could still perceive the

red trace produced by the cursor's movement. As such, participants could not anticipatorily plan their

forces (e.g., by seeing that they were at the peak of the positive half cycle or whether they were

performing a cycle in the slow or fast condition) but they still received on-line feedback about their

performance. Optimal performance in this task would consist of sinusoidally scaling fingertip forces but

would show the red trace (produced by the cursor's movement) overlapping with the straight horizontal

target line.

The top row (Figure 3A) shows a representative example of a participant's on-screen view and their

performance in the horizontal line condition. The sinusoidal curve was hidden (i.e., only a target horizontal

line was shown) although participants still received visual feedback about their ongoing performance. The

second row of Figure 3 (Figure 3B) shows the actual sinusoidal curve, that was hidden from the

participants, and the participant's performance relative to this sinusoidal curve. When a participant would

perform the sinusoidal curve condition, the target curve and the feedback of the participant's

performance would look like the example in Figure 3B. On this figure it can also be seen that in this

example the first two and last cycles are part of the 'fast' condition, whereas the third cycle is part of the

'slow' condition.

The sinusoidal curve and horizontal line conditions were presented in a blocked design which were
counterbalanced across participants to avoid order effects. In each block, 13 trials of 30 seconds were
performed (6 participants) or 5 trials of 120 seconds (8 participants). The length of trials was changed
after the first six participants as it became apparent that participants could easily perform trials of longer
durations. Each trial consisted of a random sequence of the four conditions (frequency: slow-fast;
direction: down-up). In each trial, each condition was equally presented to avoid potential fatigue and
decline-in-attention effects. To ensure that group identity (i.e., either performing short or long trials) did
not impact task performance, we compared performance (AUC parameter as described in Data analysis)
for each condition between our groups using student t-tests. For each comparison (e.g., slow up of the
short trial group vs slow up of the long trial group), we found a p-value of minimally p = 0.144 which are
not corrected for multiple comparisons. Because of this, we argued that there are no meaningful
differences between both groups and, thus, decided to pool both groups for further analyses.

Table 1. Events included in the analysis for each condition on average (after removal of incorrect events).
Mean ± SD over participants for each condition.

	Slow up	Fast up	Slow down	Fast down
Sinusoidal curve	25.3±4.8	28.9±5.6	24.0±7.2	18.7±6.6
Horizontal line	26.1±6.0	26.2±8.3	21.7±8.6	17.5±5.0

At the start of each trial, a preparatory target line at the 2 N level was shown. The cursor took between
5 to 8 seconds (randomized) to complete, allowing participants to move the cursor to the midpoint force
level at the start of the trial. This preparatory target line would then transition into the experimental
sinusoidal curve or horizontal line. The transition between the preparatory and experimental target line
was indicated with a vertical line. Before starting the experimental task, 2 practice trials of 30 seconds
were performed, and a short break was presented between the two blocks.

<u>3.2 Data analysis</u>
Several parameters were extracted when TMS was applied (i.e., when participants generated 2.5 N); the
MEP amplitude (as defined above) was extracted in a window of 5 to 45 ms after the TMS pulse. From the
EMG, we also extracted muscle activity using peak-to-peak amplitude and root mean squared activation
(RMS100) in a 100 ms interval before the TMS pulse (105-5 ms before the TMS pulse) as a measure of
background EMG before TMS.

Figure 4. Example of calculation of Euclidian distance to determine erroneous trials. The blue and red line represent the target and performed force, respectively. From the point at which the TMS was applied (green asterisk) the shortest distance to the target force was determined, which was done in 2D space incorporating both the deviation in time and force. Black lines indicate distances to the target curve if only force or time were considered.

Although participants were instructed to keep the cursor as close as possible to the target line, they often made errors. Particularly in the horizontal line condition, participants could not rely on visual information regarding the upcoming force requirements as they could only rely on their ongoing performance (as indicated by increasing/decreasing deviations from the target horizontal line). Accordingly, participants sometimes scaled their fingertip forces not in line with the current event (e.g., up- instead of downscaling or fast instead of slow scaling). To assess (i) which event should have been performed (up/down – fast/slow) and (ii) performance on this event, we did the following: we calculated the total error in 2D space (force-time space). This allowed us to include errors in force (too much/too little force applied at any given give) and time (where participants should have been on the curve when applying a certain amount of force). The total error for each event was defined as the shortest Euclidean

distance between the target curve and the participant's force generation at the TMS event (see Figure 4).

Specifically, in Figure 4, the TMS event is indicated with a green asterisk on the actual performed force

(red line). The shortest distance from this asterisk to the target curve (i.e., between current performance

and the target sinusoidal curve in blue) was determined. Note that in the horizontal line condition, even

though participants perceived a straight horizontal target curve, we used the 'hidden' sinusoidal curve to

investigate performance (i.e., participants saw on-screen Figure 3A, but their performance was quantified

as in Figure 3B). We decided to include both dimensions (force and time) for assessing general

performance as, for instance, participants could have been generating the appropriate amount of force

although they were performing the incorrect event (e.g., up- instead of downscaling forces). As the TMS

pulse was automatically triggered when participants generated 2.5 N (with a minimum 3.5 s delay

between two consecutive stimulations), we had no direct control over when TMS was applied. We

reasoned that if participants deviated too much in force-time space, the MEP would not reflect the current

event and thus needed to be removed from analysis. Based on visual inspection of all trials, a criterion of

0.55 was used for this total error. This value of 0.55 was chosen as it closely matched the visual judgement

of two observers that independently rated the individual trials. Using this criterion, 2% of all MEPs were

removed. Note that we did not include this parameter (the Euclidian total error) to assess participants'

overall performance but rather to reject incorrectly performed events. In addition, we also removed MEPs

that were smaller than the peak-to-peak EMG in the 100 ms interval before TMS (134 trials, 4%), as these

MEPs lower than background EMG activity likely reflect noise.

Last, MEPs were obtained from TMS applied at the same force level. Since EMG amplitude

depends on the amount of force that is applied, the RMS100 should not differ between the conditions.

However, we found that this was not always the case (see Results). To control for differential force

activation, we also divided the MEP values by the RMS100 before the TMS pulse and analysed this

corrected value (MEP$_{EMG}$) separately. We decided to perform this normalization to be in line with Kimura

et al. (2003) and given that MEP amplitude varies with muscle contraction (Lavoie et al., 1995).

To investigate actual participants' performance, we relied on force-related parameters. The average

force rate (time differentiated force) was determined between 300 ms before the TMS pulse and the TMS

pulse. Furthermore, to examine the deviation from the target line for a longer period, the difference

between the target force and the generated force was calculated from the midpoint of the curve (start of

a condition) until the time point TMS was applied. The absolute area under the curve of this difference

(AUC) was calculated as a measure of deviation from the target curve. To correct for possible different

time durations the AUC was divided by the number of samples included in its calculation. In this way, the

AUC represented the average deviation between the executed force and the target force from the start

of the curve to the time of TMS in a specific condition. Finally, to further evaluate task performance, the

average speed was calculated as the mean force rate in the 300 ms before the TMS pulse.

<u>3.3 Statistical analysis</u>

All variables (MEP, RMS100, average speed, AUC) were calculated for each condition (i.e., fast up, fast

down, slow up or slow down) and each task (sinusoidal curve or horizontal line). Since TMS was applied

3.5 s apart and always at the performed force level, the number of events in each condition differed

among participants (as shown in Table 1 and explained in calculating the total Euclidian error in 'Data

Analysis'). A repeated measures analysis of variance (ANOVA) was performed on the variables with factors

direction (up, down), speed (slow, fast) and task (sinusoidal curve or horizontal line). In line with previous

work of our group (Rens et al., 2020), we z-scored MEP and RMS100 data for each participant separately

before calculating participant averages and performing statistics. Post-hoc tests were performed with

Bonferroni corrected t-tests. A significance level of 0.05 was used and effect sizes for the factors are

reported using partial eta squared (η_p^2). Only if otherwise denoted, Data in text is reported as mean ±

standard error.

4. Methods of Experiment 2

<u>4.1 Task and procedure</u>

After initial preparations (i.e., consent, questionnaires and TMS preparations), participants

performed the experimental task. The behavioural task of experiment 2 comprised of an object-lifting task

based on Mawase & Karniel (2010). Participants were comfortably seated in front of a table with their

right arm resting on the tabletop. The experimenter was seated in front of the participant. Participants

were instructed to grasp and lift the manipulandum on the table and received the following instructions:

(1) Lift the manipulandum to a height of approximately 5 cm at a smooth pace that is natural to you. (2)

Only place thumb and index finger on the graspable surfaces (i.e., use precision grip). (3) After lifting, keep

the manipulandum in the air until the switchable screen (see below) turns opaque. Only then replace the

object on the table. (4) Only start reaching towards the manipulandum after you have received the TMS

pulse. Accordingly, one trial consisted of one reach-grasp-lift-replace action. Before the start of the

experiment, five practice trials were performed with a random sequence of objects included in the

experimental task.

A switchable screen (MagicGlass) was placed between the participant and the table. The screen could

become transparent or opaque to enable/block vision on the manipulandum, respectively. When the

screen was opaque between trials, the experimenter could change the cube in the manipulandum's

basket out of the participant's view. Five possible objects could be placed in the manipulandum with

approximate weights of 1.04, 2.55, 3.11, 3.56 and 5.15 N, which will be referred to as (weight) 1-5. As

mentioned before, the manipulandum weighted approximately 1.2 N, thus giving total weights of

approximately 2.23, 3.75, 4.31, 4.75 and 6.34 N to be lifted. Total weight was assessed by holding the

cube-loaded manipulandum steady in the air for a few seconds which was repeated 5 times for each

weight. These cubes were 3D printed, visually identical and measured 5×5×5 cm. In addition, all objects

were hidden below the same paper cover to ensure that participants could not rely on visual cues

indicating object weight.

Each trial started with the switchable screen becoming transparent. After 400 ms (± 100 ms jitter), a

TMS pulse was applied over M1. Four seconds after the screen becoming transparent, it turned opaque

again indicating the end of the trial and indicating participants to replace the manipulandum on the table.

Force data was only sampled when the screen was transparent, EMG data was continuously sampled but

only an epoch of 5 seconds (from 2 seconds before to 3 seconds after the TMS pulse) was stored.

The TMS pulse served as the 'go' cue for participants to initiate their action of grasping, lifting, and

replacing the manipulandum. We decided to stimulate during planning, not execution, as we did not want

to interfere with the participants' lifting performance which would be caused by the unvoluntary muscle

contraction due to M1 stimulation. To ensure that participants could not anticipate the TMS timing, we

decided to include a jitter of 100 ms. We decided to use 300 ms as the lower threshold (thus 400 ms ± 100

ms jitter) as this would provide minimally 300 ms of visual information to participants. Although TMS

during lift planning was applied substantially later in our study than in the one of Loh et al. (2010), it should

still elicit CSE modulation as shown in previous work of our group (Rens et al., 2021b). In addition, Parikh

et al. (2014) used a similar object lifting paradigm, applying TMS 1000 ms after object presentation. Briefly,

in their study, CSE was also task-specifically modulated. Altogether, these studies indicate that TMS at the

'go cue' probes planning effects on CSE modulation, irrespective of the timing of the cue itself.

Similar to the task of Mawase & Karniel (2010) the weights were presented in an apparently random

order although we used both series of increasing weights and of random weights. A series of increasing

weights consisted of the weights 1, 2, 4 and 5 (i.e., four trials). After this series of increasing weights,

weight 3 was presented in the fifth trial as a 'catch' trial'. The series of random weights varied between 4

and 9 trials of which a randomly chosen trial comprised of lifting weight 3 (weight 3 was presented earliest

trial 2 and latest trial 9). This was to ensure that participants would not anticipate that the catch trial was

the fifth trial after the series of increasing weights.

As weight 3 was presented after both the series of increasing and of random weights, we could

compare force scaling in catch trials between these two conditions. Accordingly, we could investigate

whether participants would predictively scale their forces differently for the same weight based on

implicit information (i.e., increasing vs. random series). Importantly, to ensure that predictive force scaling

differences for weight 3 between series were not driven solely by the previous lift (Johansson & Westling,

1988), we decided that the weight immediately before weight 3 in the series of random weights was the

same as the last weight in the series of increasing weights (i.e., weight 5) (for an example sequence of the

object lifting task see Figure 5). For experiment 2, we decided on 15 catch trials per condition, thus

resulting in 15 series of increasing weights and 15 series of random weights which was close to double the

amount of trials in Mawase & Karniel (2010).

A different experimental sequence was generated for each participant. The 15 series of increasing

weights included 4 weights and a catch trial, giving a total of 15 x 5 = 75 trials. The 15 random series

consisted of 4-9 random weights (including a catch trial), giving on average 6.5 x 15 = 97.5 trials. In

addition, each experimental sequence included 5-10 and 2-4 random trials at the start and end of the

experiment, respectively. In total, participants performed on average 181 trials. Participants received a

break after approximately 60 and 120 trials which varied as the break took place after completing a series

(thus after lifting weight 3). After completing the experiment, participants performed a short

questionnaire (akin to Mawase & Karniel, 2010), which contained questions about their experience of the

objects:

1) Did you experience a change in the weight of the objects?

2) Did you experience this change in object weights as random?

3) Did you experience a pattern or series in the change of object weights?

4) Did you feel like some weights appeared more often or less frequent, or was this equally divided?

Figure 5. *Example of trial sequence in experiment 2. Trials consisted of a series of increasing weights (turquoise) or random order (dark blue). A catch trial was presented in each series: at the end of an increasing series (red) or at a random position in the random series (light blue). Note that a catch trial was always preceded by the heaviest weight in the object set.*

4.2 <u>Data analysis</u>

EMG data. From the EMG recordings, we extracted the MEPs, which were defined as the peak-to-peak

amplitude in the EMG signal, using a custom-written MATLAB script. MEPs were determined between 5

ms to 45 ms after the TMS pulse. Trials were visually inspected for background noise related to muscle

contractions. Trials that showed visual contamination of EMG activity prior to TMS were flagged and

removed from analysis (3.6 %). In addition, MEPs were also considered noise and removed when they

were smaller than 50 µV or smaller than the peak-to-peak EMG signal in the 100 ms before TMS (1.7 %)

as they could not be discerned from background noise.

Force data. Trials in which the incorrect object was lifted (i.e., experimenter placing the wrong

weight), or in which the object was not lifted or lifted twice in one trial (e.g., when the participant's

fingertips slipped from the force sensors) were removed from the force analysis (0.3%). The force data

was filtered with a fourth order low-pass bidirectional Butterworth filter with a cut-off frequency of 15

Hz. This filter was selected to provide an optimal balance between noise removal and signal smoothing to

find the peaks in the force rates (Van Polanen et al., 2020; van Polanen & Davare, 2015). Total load force

(LF) and total grip force (GF) were the sum of the forces parallel and perpendicular to the gravitational

force, respectively. Load and grip force rates (LFR and GFR) were defined as the first time-derivatives of LF and GF. Peak values of the force rates were determined between object contact (defined as GF > 0.1 N) and 50 ms after lift-off (LF > object weight). We decided to use the peak values, and in particular the first peak values (peak1 LFR and peak1 GFR), of the force rates (and not the forces) as this has been considered a more sensitive measure of predictive force scaling (Johansson & Flanagan, 2009; Johansson & Westling, 1988). In line with Van Polanen et al. (2020), we extracted the first peak that was at least 30% of the maximum value to avoid noise extraction. Finally, the loading phase duration (LPD) was determined as the latency between LF onset (LF > 0.1 N) and object lift-off (defined above). To end, we also calculated grip force at lift-off (GFatLO). GFatLO was extracted as this was the behavioural parameter used in (Mawase & Karniel, 2010). This allowed us to directly compare our findings with theirs.

4.3 Statistical analysis

We extracted all parameters (MEP, GFatLO, peak 1 GFR, peak1 LFR, LPD,) for catch trials after the series of increasing and random weights. For each parameter, outliers, defined as being smaller or larger than the mean ± three standard deviations, were removed from the analysis (< 2 % for any parameter). Analyses were performed on the averaged participant data. In line with Rens et al. (2020), MEP data were converted to z-scores to take inter-subject variability in account. Data were then averaged per parameter for each participant and condition and then used for statistical analysis. While data was obtained for longer series (see Figure 7), we only performed contrast comparisons on the last (catch trial) and the second to last lift (weight 5) of the series of increasing and random weights. For this, we used a repeated measures ANOVA with two factors being series (increasing vs random) and weight (weight 5 vs catch trial). Note that we only included the last two lifts due to the series of increasing weights consisting of 5 lifts and the series of random weights of at least 3 and maximally 5 lifts. In addition, while the last two lifts of both series were always the same (weight 5 and catch trial), the second to last lift of the random series varied randomly, providing no meaningful inputs for the statistical analysis. Post-hoc tests were performed with Bonferroni corrected t-tests. A significance level of 0.05 was used and effect sizes for the factors are reported using partial eta squared (η_p^2). Only if otherwise denoted, data is reported as mean ± standard error. To end, we calculated correlations between the weights and the participant averages for each parameter (MEPs and behavioral). This was done to explore a potential linear relationship and was performed once including the catch weight and once excluding it.

5. Results

<u>5.1 Experiment 1</u>

In the first experiment, participants tracked a curve by applying force on two sensors. TMS was
applied when a force of 2.5N was crossed, which could occur when moving up or down on the curve and
at a fast or slow speed.

Corticospinal excitability. The MEPs were compared for all conditions to test whether CSE differed
depending on the task that had to be performed. Results are shown in Figure 6A. The repeated measures
ANOVA showed that MEPs were larger when moving up (up = 0.42 ± 0.04) compared to moving down
(down = -0.42 ± 0.07) on the curve (effect direction, $F(1,13) = 74.9$, $p<0.001$, $\eta_p^2 = 0.85$). Furthermore,
MEPs were also larger when the movement was fast (fast = 0.08 ± 0.03) compared to slow (slow = -0.08 ±
0.04) (effect speed, $F(1,13) = 22.8$, $p<0.001$, $\eta_p^2 = 0.64$). However, an interaction effect for speed x
direction ($F(1,13) = 7.8$, $p = 0.015$, $\eta_p^2 = 0.38$) indicated that the speed effect was only seen in upwards
movements (up fast = 0.57 ± 0.06; up slow = 0.27 ± 0.03; p = 0.002), but not in downwards movements
(down fast = -0.40 ± 0.07; down slow = -0.44 ± 0.09; p = 1.00). The difference in CSE modulation was also
seen between moving up and down within both the fast (up fast compared to down fast) and slow (up
slow compared to down slow) movements (both p<0.001). Finally, an interaction for direction x task was
found ($F(1,13) = 7.2$, $p = 0.019$, $\eta_p^2 = 0.36$), but post-hoc tests only showed differences between up and
downward movements within tasks (sinusoidal curve: up = 0.47 ± 0.06; down = -0.48 ± 0.03; p < 0.001;
horizontal line: up = 0.37 ± 0.07; down = -0.36 ± 0.1; p < 0.001), but no differences within direction
between tasks (e.g., upward movement compared between the sinusoidal curve and horizontal line
conditions) were found (both p = 1.00). To end, the main effect of task ($F(1,13) = 0.35$, $p = 0.855$, $\eta_p^2 =$
0.003) as well as the interaction effects of task x speed ($F(1,1) = 0.43$, $p = 0.524$, $\eta_p^2 = 0.032$), and task x
direction x speed were not significant ($F(1,13) = 0.70$, $p = 0.418$, $\eta_p^2 = 0.05$) .

Background EMG. Since MEPs also depend on background EMG, we compared the RMS of the EMG
obtained 100ms before the TMS was applied (RMS100). Here, an effect of direction ($F(1,13) = 112.90$,
$p<0.001$, $\eta_p^2 = 0.90$) was found. RMS100 was higher when going up (up = 0.28 ± 0.04) compared to going
down (down = -0.44 ± 0.04). The main effects of task ($F(1,13) = 0.53$, $p=0.481$, $\eta_p^2 = 0.039$) and speed
($F(1,13) = 2.37$, $p=0.148$, $\eta_p^2 = 0.15$) were not significant as well as the interaction effects of task x direction
($F(1,13) = 0.11$, $p=0.751$, $\eta_p^2 = 0.008$), task x speed ($F(1,13) = 0.62$, $p=0.45$, $\eta_p^2 = 0.045$), direction x speed
($F(1,13) = 4.42$, $p=0.056$, $\eta_p^2 = 0.254$) and task x direction x speed ($F(1,13) = 0.14$, $p=0.713$, $\eta_p^2 = 0.011$).

Together, these results suggest that MEPs differ depending on movement direction and speed, but

not on visibility of the target curve. However, the direction effect as observed in the MEPs were also seen

in the background EMG, which might have contaminated the MEP results.

Figure 6. *Results for experiment 1, for both the sinusoidal and horizontal line condition.* ***A.*** *Motor evoked potentials*
(MEP). ***B.*** *MEPs divided by background EMG.* ***C.*** *Average speed.* ***D.*** *Area under the curve (AUC). Green and red bars*
represent slow and fast conditions, respectively. Blue dots represent individual values. Main effects are reported and
interactions are indicated with significance bars. Error bars indicate standard errors.

Corticospinal excitability corrected for background EMG. To ensure that the effects of speed and

direction for the MEPs were not due to differences in background EMG, we performed an additional

analysis on the corrected MEPs. To correct for background differences in muscle contraction, we divided

the MEP of each TMS event by the RMS100 in that event (shown in Figure 6B). For this MEP$_{EMG}$ we found

main effects of direction (F(1,13) = 12.5, p = 0.004, η_p^2 = 0.489) and speed (F(1,13) = 11.2, p = 0.005, η_p^2 =

0.464), indicating that MEPs were still higher when going up compared to going down (up = 0.29 ± 0.05;

down = -0.22 ± 0.11) and moving fast compared to slow (fast = 0.10 ± 0.03; down = -0.03 ± 0.05), even

when corrected for background EMG. For this analysis we also found an interaction for direction × task

($F(1,13)$ = 4.8, p = 0.046, η_p^2 = 0.272). Post-hoc tests indicated that the upward movements were only

different from the downward movements in the sinusoidal curve condition (up = 0.38 ± 0.07; down = -

0.28 ± 0.04; p<0.001), but not in the horizontal line condition (up = 0.20 ± 0.09; down = -0.16 ± 0.13; p =

0.292). No differences were seen between the two conditions (between conditions: both p > 0.21). Finally,

no significance for the main effect of task ($F(1,13)$ = 0.16, p = 0.697, η_p^2 = 0.012), nor for the interaction

effects of task x speed ($F(1,13)$ = 2.22, p = 0.159, η_p^2 = 0.146), direction x speed ($F(1,13)$ = 2.42, p = 0.144,

η_p^2 = 0.157) and task x direction x speed ($F(1,13)$ = 1.61, p = 0.226, η_p^2 = 0.110) was found.

Rate of force scaling. To assess whether participants performed the task correctly, we calculated the

average speed prior to the TMS (Figure 6C). The ANOVA on the average speed showed significance for the

main effects of direction ($F(1,13)$ = 518.1, p < 0.004, η_p^2 = 0.98) and task ($F(1,13)$ = 12.3, p = 0.004, η_p^2 =

0.486) but not of speed ($F(1,13)$ = 0.2, p = 0.691, η_p^2 = 0.013). All interaction effects were significant: task

x direction ($F(1,13)$ = 7.0, p = 0.020, η_p^2 = 0.350), task x speed ($F(1,13)$ = 41.9, p < 0.001, η_p^2 = 0.763),

direction x speed ($F(1,13)$ = 8.2, p < 0.001, η_p^2 = 0.877) and task x direction x speed ($F(1,13)$ = 26.5, p <

0.001, η_p^2 = 0.671). Participants moved faster in the fast conditions compared to the slow conditions (all

p<0.001), except in the horizontal line condition when making a downward movement where no

difference was found (p = 1.00). Not surprisingly, the direction also differed based on whether up or down

movements were made (all p<0.001). There were no notable differences between the sinusoidal curve

and horizontal line conditions, except for the fast down condition, which was slower in the horizontal line

compared to the sinusoidal curve condition (p = 0.002). These measures indicated that participants

seemed to perform the task according to the instructions (sinusoidal curve: slow up: 0.83 ± 0.047, fast up:

1.35 ± 0.06, slow down: -0.73 ± 0.048, fast down: -1.60 ± 0.075; horizontal line slow up: 0.89 ± 0.039, fast

up: 1.49 ± 0.039, slow down: -0.76 ± 0.085, fast down: -0.94 ± 0.169).

Motor performance. To determine whether participants' performance differed between conditions,

we measured the AUC of the difference between the target curve and the applied force (Figure 6D). An

ANOVA on the AUC showed that the deviation depended on the direction ($F(1,13)$ = 148.1, p<0.001, η_p^2 =

0.919), speed ($F(1,13)$ = 161.7, p<0.001, η_p^2 = 0.926) and task ($F(1,13)$ = 100.0, p<0.001, η_p^2 = 0.885).

Participants deviated further from the curve when they moved upwards (up = 0.30 ± 0.01) compared to

downwards (down = 0.21 ± 0.02). In addition, a larger AUC was seen for fast movements (fast = 0.32 ±

0.02) compared to slow movements (slow = 0.18 ± 0.01). Finally, larger deviations were seen in the

horizontal line condition (mean = 0.33 ± 0.02) compared to the sinusoidal curve condition (mean = 0.17 ±

0.01). Furthermore, significant interactions for direction x task (F(1,13) = 56.0, p<0.001, η_p^2 = 0.812) and

for speed x task (F(1,13) = 102.3, p<0.001, η_p^2 = 0.887) were observed. However, the post-hoc tests of

these interactions were all significant (all p<0.01), providing no new interpretations to the data. Finally,

no significance was found for the interaction effect task x direction x speed (F(1,13) = 1.4, p = 0.254, η_p^2 =

0.099)

In sum, CSE was primarily modulated by the direction and speed of the movement, even after

correcting for background EMG. This CSE modulation was seen across in the sinusoidal curve and

horizontal line conditions, i.e., when the sinusoidal target curve was visible or when only feedback from a

straight line was present. As such, our findings mainly provide evidence that the motor system encodes

ongoing motor components (i.e., direction and speed) but not that it encodes the differences in available

information between the sinusoidal curve and horizontal line conditions.

5.2 Experiment 2

In experiment 2, participants lifted objects that were presented either in a series of increasing or random

weights. After a series of increasing weights or within the random series, a 'catch trial' (weight 3) was

presented. Immediately prior to this catch trial, in both the series of random and increasing weights, the

heaviest (weight 5) was presented. In addition, we applied TMS prior to object lifting to assess CSE

modulation during lift planning. With this experiment, we aimed to investigate whether lift planning (on

the behavioral and corticospinal level) reflects implicit information, related to a series of random or

increasing object weights, during a discrete object lifting task. For this we investigated differences in

predictive force scaling and CSE modulation for the catch trial only.

Debriefing results. After the experimental task, participants were debriefed and were asked

questions using a standardized questionnaire (see methods) related to the experimental task. All

participants (Question 1: 14/14) reported that the weights of the object changed between trials. All but

one participant (Question 2: 13/14) experienced these changes as random and two participants (Question

3: 2/14) reported to experience a pattern in the object presentation. While most participants (Question

4: 10/14) reported all weights to appear equally frequent, three answered that there were more heavy

objects in the sequence and one participant argued they did not know whether the weights appeared

equally frequent.

Corticospinal excitability. As can be seen in Figure 7A, neither the main effects of series (F(1,13) = 1.5,

p = 0.705, η_p^2 = 0.011) and weight (F(1,13) = 0.65, p = 0.434, η_p^2 = 0.048) nor the interaction effect of series

x weight (F(1,13) = 0.9, p = 0.373, η_p^2 = 0.061) were significant. As such, our results provide no evidence

that objects weight influenced CSE modulation differently during the series of increasing and random

weights. With specific interest to the series of increasing weights, no significant correlation was found

between object weight and MEP amplitude (excluding catch trial: r = -0.134, p = 0.323; including catch

trial: r = -0.115, p = 0.341).

Grip force at lift off. For GFatLO (shown in Figure 7B), the main effect of series (F(1,13) = 0.0, p =

0.851, η_p^2 = 0.003) and the interaction effect of series x weight were not significant (F(1,13) = 1.4, p =

0.258, η_p^2 = 0.097). In contrast, the main effect of weight (F(1,13) = 200.6, p < 0.001, η_p^2 = 0.939) was

significant. Participants exerted more grip force at lift off when lifting the heavy weight (heavy = 16.40 ±

0.85 N) than when lifting the catch weight (catch = 12.16 ± 0.76 N). Notably, the correlation between

object weight and the grip applied at lift off for the series of increasing weight was significant when both

the catch trial was excluded (r = 0.655, p < 0.001) and included (r = 0.614, p < 0.001).

First peak grip force rate. For peak 1 GFR (Figure 7C), all effects were not significant: series F(1,13) =

0.0, p = 0.899, η_p^2 = 0.001), weight F(1,13) = 3.6, p = 0.078, η_p^2 = 0.220) and series x weight F(1,13) = 0.3,

p = 0.586, η_p^2 = 0.023). In addition, correlations between object weight and the first peak of grip force rate

were not significant either (excluding catch weight: r = 0.067, p = 0.624; including catch weight: r = 0.059,

p = 0.792).

*First peak load force rate. F*or peak 1 LFR (Figure 7D), the effects of series F(1,13) = 1.3, p = 0.283, η_p^2

= 0.088) and series x weight F(1,13) = 1.5, p = 0.239, η_p^2 = 0.105) were not significant. However, the main

effect of weight F(1,13) = 8.5, p = 0.012, η_p^2 = 0.395) was significant: participants scaled their load forces

faster (larger peak LFR value) when lifting the catch weight (catch = 31.40 ± 2.43 N/s) than when lifting

weight 5 (weight 5 = 28.64 ± 2.12 N/s). Akin to peak 1 GFR, correlations, correlations between object

weight and the first peak of load force rate were not significant (excluding catch weight: r = 0.036, p =

0.791; including catch weight: r = 0.032, p = 0.792).

Loading phase duration. Similar to peak 1 LFR, the main effect of weight F(1,13) = 157.7, p < 0.001,

η_p^2 = 0.924) was significant for LPD (Figure 7E) whereas the effects of series F(1,13) = 0.3, p = 0.592, η_p^2 =

0.023) and series x weight (F(1,13) = 1.7, p = 0.214, η_p^2 = 0.116) were not. In line with peak 1 LFR,

participants had a shorter loading phase duration when lifting the catch weight (catch = 196.75 ± 9.28 ms)

than when lifting weight 5 (weight 5 = 320.64 ± 14.47 ms). Notably, correlations between object weight

and LPD were significant (excluding catch weight: r = 0.809, p < 0.001; including catch weight: r = 0.776, p

< 0.001).

In sum, these findings show differences (Grip force at lift off and loading phase duration) on the
behavioral level when participants lifted the heaviest weight or the catch one. However, our results
provide no evidence that participants lifted the catch weight differently after the series of increasing or
random weights. In addition, this absence of proof was also present on the corticospinal level (i.e., no
differences between MEPs).

Figure 7. *Results for experiment 2 for the catch trial after a series of increasing (red) or random (blue) weight. X-axis*
represents the catch trial (trial N) and all preceding trials for each series. For the series of increasing weights, the

6. Discussion

In the present study, we investigated how explicit and implicit contextual information affected force

scaling and CSE modulation during hand-object interactions. In experiment 1, participants performed a

continuous force tracking task and in experiment 2, other participants performed a discrete object lifting

task. In both experiments, motor performance was assessed to investigate how participants scale their

fingertip forces during continuous and discrete hand-object interactions. TMS was used to probe the

underlying changes in CSE.

For experiment 1, we found that when a sinusoidal curve was shown (i.e., explicit visual

information of position on curve and performance error), participants performed significantly better

compared to the condition in which the sinusoidal curve was shown as a horizontal line (i.e., explicit visual

information of performance error but implicit information regarding position on curve). Our results

indicate that CSE was modulated primarily by movement-related features (i.e., fast vs slow change in

muscle contraction and up- vs downscaling of fingertip forces) in both the sinusoidal curve and horizontal

line conditions. However, our findings do not robustly show that CSE modulation was altered based on

how visual information was presented. As such, these findings indicate that CSE is primarily modulated by

task requirements that can be directly mapped onto the corticospinal output (i.e., rate and direction of

force scaling) but not by whether this information is explicitly or implicitly provided.

For experiment 2, we found no behavioral and neurophysiological differences when participants

performed the catch trial after the series of increasing weights or after the series of random weights.

These findings generally indicate that participants did not predictively scale their forces by extrapolating

from the implicit weight increasing series.

In experiment 1, participants performed a sinusoidal force tracking task when provided with the

actual sinusoidal target force (sinusoidal condition) or only received feedback about their performance in

the form of a deviation from a horizontal line (horizontal line condition). As can be seen in Figure 6D,

participants performed significantly worse (higher AUC value) in the horizontal line condition compared

to the sinusoidal curve one. These findings indicate performance suffered when participants could not

visually track their location on the sinusoidal curve and thus not predict the required directionality (up- or

downscaling of force scaling) and rate of force scaling (the steepness of the curve's slope). On the neural

level we found that CSE modulation (Figure 6A-B) was primarily driven by ongoing movement-related

features, i.e., direction and the change in force scaling rate (i.e., "speed"). CSE was larger when

participants were upscaling their fingertip forces compared to downscaling (Figure 6A). The latter findings

are in line with Kimura et al. (2003) and Li (2013) who showed that CSE is increased when contracting the

elbow flexor (compared to relaxation) and when dynamically contracting the muscles of four fingers

(without thumb) compared to isometric contraction and relaxation, respectively. These studies and our

results agree with findings that motoneuron inputs having a greater relative contribution during

shortening muscle contractions than relaxations (Semmler et al. 2002) and that CSE decreases prior to

muscle relaxation (Suzuki et al. 2015). Importantly, our results extend on these studies by showing that

(a) CSE is increased when upscaling fingertip forces compared to downscaling during a force tracking task,

and (b) CSE modulation is also driven by movement speed, where it is increased with fast compared to

slow changes in force.

Critically, effects of CSE modulation by the availability of explicit information (i.e., sinusoidal curve

vs horizontal condition) seems to be minimal at best and most likely to be absent. As such, our findings

indicate that CSE modulation reflected ongoing task execution (i.e., rate of force scaling and direction) but

not the availability of explicit/implicit contextual information driving task performance.

Despite information regarding the task being less explicitly available in the horizontal line

condition (i.e., error shown in inverse manner and no explicit information regarding the curve's slope),

CSE modulation was not driven differently than in the sinusoidal curve condition. Interestingly, these

findings are in line with Young et al. (2000): using a reaction time task, they showed that when individuals

respond to stimuli, CSE is automatically driven and not selectively modulated by intention- ('readiness to

respond') and attention-selectivity ('preferential processing of stimuli'). In addition, it has been proposed

by Albert et al. (2022), using reaching tasks, that the amount of implicit learning can be altered by explicit

strategies due to the implicit and explicit learning systems competing for errors as a common source of

information about performance. Finally, Bestmann et al. (2008) showed that CSE modulation during action

preparation becomes smaller when uncertainty increases. Combining the findings of these studies, it is

feasible that, in our experiment 1, participants used the same behavioral strategy across both tracking

conditions: focus on deviations between target line and actual performance and, thus, not necessarily on

the target line's shape. Given that errors were shown differently between conditions (i.e., inverse in the

horizontal line condition leading to higher uncertainty), this could explain lower performance in the

horizontal line condition while the similarity in the behavioral strategy between conditions could then

explain the absence of differences in CSE modulation across conditions. However, this assumption cannot

be verified with the current data but could be investigated in future studies including eye tracking to probe

participants' behavioral strategy and conditions where the visibility of the target line is manipulated. In

sum, our findings for the continuous force tracking experiment suggest that corticospinal outputs are

primarily driven by the ongoing motor commands and do not seem to be influenced by the visual nature

of how feedback about ongoing performance is provided.

790   A potential limitation of our study is that we only performed TMS at the upper part of the curve,

i.e., at 2.5 N. This could create a potential bias regarding the up and down conditions, since TMS was

applied when participants were increasing their forces for a longer time compared to when they were

decreasing the force. Although the forces were dynamic, requiring no adaptation to a static force, and the

rate of change (frequency) could be altered at the 2N midway point, it is possible that participants adapted

differently to specific force directions or force speeds. However, since our differences between upward

and downward forces replicate results of previous studies (Kimura et al. 2003), it seems unlikely that

differences in CSE modulation were only due to adaptation differences in behavior.

798   It should be noted that, since TMS was applied at a fixed force level, participants could have

anticipated the TMS pulse. However, we took care to construct the sinusoidal curves in such a way that

TMS was not always applied at 2.5 N: a minimum duration of 3.5 s was placed between each stimulation

to avoid carry-over effects. As such, TMS was not applied every time participants exerted 2.5 N of force,

thus making the stimulation less predictable. Finally, since we found no differences in CSE modulation

between the horizontal and sinusoidal condition, it seem unlikely that the MEPs were affected by

expectation of the TMS pulse. In support of the continuous motor task in experiment 1, we performed

experiment 2 to investigate how implicit information affects CSE modulation in a discrete object lifting

task. Previous studies have indicated that the CSE modulation represents task requirements (i.e., object

weight) based on the previously lifted object (Loh et al., 2010). Noteworthy, behavioural evidence

suggests that the sensorimotor system can adapt other object lifting strategies based on information

acquired across several lifts (Mawase & Karniel 2010; Cashaback et al. 2017). Here, by relying on a

paradigm similar to Mawase & Karniel (2010), we investigated whether a lifting strategy, acquired by

lifting hidden series of increasing weights, would be reflected in CSE modulation. Participants performed

an object lifting task on differently weighted, but visually identical, cubes. Within the overarching

sequence of presented weights (see Figure 5), two types of series were hidden. One consisted of random

weights whereas the second one consisted of increasing weights. Importantly, in both the random and

increasing weights series, the second to last trial was always performed on the heaviest object and the

last trial on a lighter 'catch weight', as in Mawase & Karniel (2010). The second to last weight was

standardized across increasing and random series to control for sensorimotor memory effects (Baugh et

al., 2012; Castiello, 2005; Johansson & Westling, 1993). With this experiment, we wanted to investigate

whether CSE modulation would reflect the weight series (random or increasing) beyond each lift

separately. Critically, our results provided no evidence that participants lifted the catch weight differently

based on the series of increasing weights. Accordingly, we did not observe CSE modulation either. That is,

we did not find statistical differences for behavioral parameters and CSE between same-weighted lifts

after the series of random and increasing weights. Noteworthy, Mawase & Karniel (2010) reported that

participants extrapolated their fingertip forces based on the hidden series of increasing weights. There

are three major differences between their and our study that could explain why we could not replicate

their behavioural results.

First, they relied on a virtual reality set-up whereas we used real hand-object interactions. In a

recent review, Snow & Culham (2021) argued substantial differences between how virtual and real objects

affect perception, cognition and action. For instance, individuals have a 'real-object preference' when

gazing at real objects and 2D representations of the same object. It is possible that in the study of Mawase

& Karniel (2010) participants relied on a virtual-reality associated lifting strategy that differs from how we

interact with real-world objects. For instance, in a virtual reality setup, some tactile information (e.g.,

friction between object and fingertips) is missing, which contributes to force scaling (Castiello, 2005) thus

potentially altering the grasping strategy. Second, as a proxy for predictive force scaling, Mawase & Karniel

(2010) relied on grip force at lift-off whereas we relied on multiple dependent proxies (grip force, load

force duration and force rates). It has been argued that the force rate provides a higher sensitivity for

probing predictive lift planning (Johansson & Westling, 1993). Critically, neither for grip force at lift-off

(akin Mawase & Karniel, 2010) nor for the other behavioral measures did we find differences between

conditions. Third, Mawase & Karniel (2010) did not include a direct comparison between same-weighted

lifts after the series of increasing and random weights. We argued that, if we wanted to investigate how

predictive force scaling would be altered by the hidden series, a direct comparison between same

weighted objects in different conditions should be made. This is the typical approach we used before (for

example see: Rens & Davare, 2019; Rens et al., 2020; Van Polanen et al., 2020) and which has been used

by other groups in our field as well (for example see Alaerts et al., 2010; Baugh et al., 2012; Buckingham

et al., 2014; Cashaback et al., 2017; Gordon et al., 1993; Johansson & Westling, 1993; Loh et al., 2010;

Reichelt et al., 2013; Senot et al., 2011). As such, it is unclear whether the results of Mawase & Karniel

(2010) reflect (i) an object lifting strategy driven by implicitly learning the hidden series or (ii) solely

sensorimotor effects that are entirely driven by last lifted object. However, it is evident in our experiment

that participants did not develop a lifting strategy based on the hidden series. Potential explanations for

why such strategies were not adopted by participants can be derived from relevant literature. First,

Cashaback et al. (2017) showed that the sensorimotor system, in the case of uncertainty, aims to minimize

prediction errors across many lifts instead of the prediction error for each lift separately. In our study,

participants were instructed to perform the lifts naturally, but they were not specifically instructed to

avoid lifting errors as in previous work from our group (Rens et al., 2021a). As such, participants might

have relied on a general lifting strategy, minimizing prediction errors across all trials (akin to Cashaback et

al., 2017) instead of attempting to "discover" the hidden series of increasing weights and predictively scale

fingertip forces for each upcoming weight specifically (akin to Mawase & Karniel, 2010). Second, another

explanation could be that the sensorimotor system prioritizes implicit information (akin to Albert et al.,

2022) that is not related to the series of increasing weights. For instance, it is feasible that predictive force

scaling, based on sensorimotor memories, relies on implicit motor learning (e.g., Flanagan et al., 2001).

Apparently, single-lift information (i.e., sensorimotor memory) is more readily available than multi-lift

information (i.e., series of increasing weights). As such, it seems feasible that the sensorimotor system

prioritizes the integration of implicit information related to the previous lift but not to the entire series.

Subsequently, if the participant explicitly understands the series to be random (and thus does not discover

the implicit series of increasing weights), they might fall back on a general lifting strategy (again akin to

Cashabak et al., 2017). Future research could tease apart some of the questions raised here. For instance,

by altering the amount of explicit information (e.g., instructing to lift without lifting errors instead of lifting

naturally or providing participants with distinct strategies) or implicit series of weights (e.g., having each

presented multiple times in a row or use outlier weights for catch trials), it might be possible to further

investigate how implicit information is integrated for discrete object lifting as has been done for reaching

(Albert et al., 2022).

With respect to CSE, our results provide no evidence that CSE modulation was altered by the

hidden series of increasing weights. Loh et al., (2010) showed that when individuals plan to lift an object,

CSE was modulated by the previously lifted object. Based on Loh et al. (2010), it seems plausible that CSE

modulation in experiment 2 depended entirely on the sensorimotor memory of the previous lift which

reflected the heaviest weight that was the same in both the increasing and random series. It is unlikely

that this null result is driven by the timing of the TMS pulse during lift planning as we did find CSE

modulation in our previous work using the same TMS timing (Rens et al., 2021b). In addition, other work

reported lift planning effects on CSE modulation when the TMS pulse was applied even longer after object

presentation (Parikh et al., 2014). As such, and in support of our behavioral findings, we argue that the

null results are caused by sensorimotor control mechanisms (i) not being able to detect the hidden series

of increasing weights (ii) prioritizing other implicit information (Albert et al., 2022) or (iii) opting for a more

general lifting strategy that is not trial-specific (Cashaback et al., 2017).

In conclusion, in the present study we investigated how corticospinal outputs could be modulated

by implicit information during the execution of continuous (experiment 1) and discrete (experiment 2)

hand-object interactions. Our results indicate that CSE modulation is primarily concerned with ongoing

force scaling (experiment 1). Noteworthy, our results fail to demonstrate that contextual implicit

information (experiment 1: type of visual feedback; experiment 2: hidden series of increasing weights) is

represented into corticospinal outputs. As such, these findings indicate that sensorimotor control might

not integrate implicit contextual information for hand-object interactions. Accordingly, other implicit

information sources related to ongoing task performance during continuous force exertion and discrete

object lifting might be prioritized.

7. **References**

Alaerts, K., Senot, P., Swinnen, S. P., Craighero, L., Wenderoth, N., & Fadiga, L. (2010). Force

requirements of observed object lifting are encoded by the observer's motor system: a TMS study.

European Journal of Neuroscience, *31*(6), 1144–1153. https://doi.org/10.1111/j.1460-

9568.2010.07124.x

Albert, S. T., Jang, J., Modchalingam, S., Marius, B., Henriques, D., Lerner, G., Maggiore, V. D.-, Haith, A.

M., Krakauer, J. W., & Shadmehr, R. (2022). Competition between parallel sensorimotor learning

systems. *ELife*, 1–81.

Baugh, L. a., Kao, M., Johansson, R. S., & Flanagan, J. R. (2012). Material evidence: interaction of well-

learned priors and sensorimotor memory when lifting objects. *Journal of Neurophysiology*, *108*,

1262–1269. https://doi.org/10.1152/jn.00263.2012

Bestmann, S., Harrison, L. M., Blankenburg, F., Mars, R. B., Haggard, P., Friston, K. J., & Rothwell, J. C.

(2008). Influence of Uncertainty and Surprise on Human Corticospinal Excitability during

Preparation for Action. *Current Biology*, *18*(10), 775–780.

https://doi.org/10.1016/j.cub.2008.04.051

Buckingham, G., Wong, J. D., Tang, M., Gribble, P. L., & Goodale, M. A. (2014). Observing object lifting

errors modulates cortico-spinal excitability and improves object lifting performance. *Cortex*, *50*,

115–124. https://doi.org/10.1016/j.cortex.2013.07.004

Cashaback, J. G. A., McGregor, H. R., Pun, H. C. H., Buckingham, G., & Gribble, P. L. (2017). Does the

sensorimotor system minimize prediction error or select the most likely prediction during object

lifting? *Journal of Neurophysiology*, *117*(1), 260–274. https://doi.org/10.1152/jn.00609.2016

Castiello, U. (2005). The neuroscience of grasping. *Nature Reviews. Neuroscience*, *6*(9), 726–736.

https://doi.org/10.1038/nrn1744

Chouinard, P. A., Leonard, G., & Paus, T. (2005). Role of the primary motor and dorsal premotor cortices

in the anticipation of forces during object lifting. *Journal of Neuroscience*, *25*(9), 2277–2284.

https://doi.org/10.1523/JNEUROSCI.4649-04.2005

Cincotta, M., Borghesi, A., Gambetti, C., Balestrieri, F., Rossi, L., Zaccara, G., Ulivelli, M., Rossi, S.,

Civardi, C., & Cantello, R. (2003). Suprathreshold 0.3 Hz repetitive TMS prolongs the cortical silent

period: Potential implications for therapeutic trials in epilepsy. *Clinical Neurophysiology*, *114*(10),

1827–1833. https://doi.org/10.1016/S1388-2457(03)00181-0

Davare, M., Kraskov, A., Rothwell, J. C., & Lemon, R. N. (2011). Interactions between areas of the cortical

grasping network. *Current Opinion in Neurobiology*, *21*(4), 565–570.

https://doi.org/10.1016/j.conb.2011.05.021

Dupin, L., Carment, L., Guedj, L., Cuenca, M., & Krebs, M. (2019). Predictive Modulation of Corticospinal

Excitability and Implicit Encoding of Movement Probability in Schizophrenia. *Schizophrenia Bulletin*,

45(6), 1358–1366. https://doi.org/10.1093/schbul/sby186

Flanagan, J. R., King, S., Wolpert, D. M., & Johansson, R. S. (2001). Sensorimotor prediction and memory

in object manipulation. *Canadian Journal of Experimental Psychology*, *55*(2), 87–95.

https://doi.org/10.1037/h0087355

Fu, Q., Zhang, W., & Santello, M. (2010). Anticipatory Planning and Control of Grasp Positions and Forces

for Dexterous Two-Digit Manipulation. *Journal of Neuroscience*, *30*(27), 9117–9126.

https://doi.org/10.1523/JNEUROSCI.4159-09.2010

Gerbella, M., Rozzi, S., & Rizzolatti, G. (2017). The extended object-grasping network. *Experimental Brain*

Research, *235*(10), 2903–2916. https://doi.org/10.1007/s00221-017-5007-3

Gordon, A., Westling, G., Cole, K., & Johansson, R. S. (1993). Manipulation of Common and Novel

Objects. *Journal of Neurophysiology*, *69*(6, June).

Gordon, Forssberg, Johansson, & Westling. (1991). Visual size cues in the programming of manipulative

forces during precision grip. *Experimental Brain Research*, *83*(3), 447–482.

Hadjiosif, A. M., & Smith, M. A. (2015). Flexible control of safety margins for action based on

environmental variability. *Journal of Neuroscience*, *35*(24), 9106–9121.

https://doi.org/10.1523/JNEUROSCI.1883-14.2015

Hannah, R. (2020). Transcranial magnetic stimulation: a non-invasive window into the excitatory circuits involved in human motor behavior. *Experimental Brain Research*, *238*(7–8), 1637–1644. https://doi.org/10.1007/s00221-020-05803-0

Heald, J. B., Lengyel, M., & Wolpert, D. M. (2021). Contextual inference underlies the learning of sensorimotor repertoires. *Nature*, *600*(December). https://doi.org/10.1038/s41586-021-04129-3

Johansson, R. S., & Flanagan, J. R. (2009). Coding and use of tactile signals from the fingertips in object manipulation tasks. *Nature Reviews Neuroscience*, *10*(5), 345–359. https://doi.org/10.1038/nrn2621

Johansson, R. S., & Westling, G. (1993). Coordinated isometric muscle commands adequately and erroneously programmed for the weight during lifting task with precision grip. *Experimental Brain Research*, 499–515.

Johansson, & Westling. (1984). Roles of glabrous skin receptors and sensorimotor memory in automatic control of precision grip when lifting rougher or more slippery objects. *Experimental Brain Research*, 550–564.

Johansson, & Westling. (1988). Coordinated isometric muscle commands adequately and erroneously programmed for the weight during lifting task with precision grip. *Experimental Brain Research*, *71*(1), 59–71.

Kimura, T., Yamanaka, K., Nozaki, D., Nakazawa, K., Miyoshi, T., Akai, M., & Ohtsuki, T. (2003). Hysteresis in corticospinal excitability during gradual muscle contraction and relaxation in humans. *Experimental Brain Research*, *152*(1), 123–132. https://doi.org/10.1007/s00221-003-1518-1

Körding, K. P., Ku, S.-P., & Wolpert, D. M. (2004). Bayesian integration in force estimation. *Journal of Neurophysiology*, *92*(5), 3161–3165. https://doi.org/10.1152/jn.00275.2004

Kotan, S., Kojima, S., Miyaguchi, S., Sugawara, K., & Onishi, H. (2015). Depression of corticomotor excitability after muscle fatigue induced by electrical stimulation and voluntary contraction. *Frontiers in Human Neuroscience*, *9*(June), 1–7. https://doi.org/10.3389/fnhum.2015.00363

Landy, M. S., Trommershäuser, J., & Daw, N. D. (2012). Dynamic estimation of task-relevant variance in movement under risk. *Journal of Neuroscience*, *32*(37), 12702–12711. https://doi.org/10.1523/JNEUROSCI.6160-11.2012

Lavoie, B. A., Cody, F. W. J., & Capaday, C. (1995). Cortical control of human soleus muscle during volitional and postural activities studied using focal magnetic stimulation. *Experimental Brain Research*, *103*(1), 97–107. https://doi.org/10.1007/BF00241968

Li, S. (2013). Analysis of increasing and decreasing isometric finger force generation and the possible role

of the corticospinal system in this proces. *Motor Control*, *17*(3), 221–237.

https://doi.org/10.1016/j.biotechadv.2011.08.021.Secreted

Loh, M. N., Kirsch, L., Rothwell, J. C., Lemon, R. N., & Davare, M. (2010). Information about the Weight of

Grasped Objects from Vision and Internal Models Interacts within the Primary Motor Cortex.

Journal of Neuroscience, *30*(20), 6984–6990. https://doi.org/10.1523/JNEUROSCI.6207-09.2010

Mawase, F., & Karniel, A. (2010). Evidence for predictive control in lifting series of virtual objects.

Experimental Brain Research, *203*(2), 447–452. https://doi.org/10.1007/s00221-010-2249-8

Parikh, P., Davare, M., McGurrin, P., & Santello, M. (2014). Corticospinal excitability underlying digit

force planning for grasping in humans. *Journal of Neurophysiology*, *111*(12), 2560–2569.

https://doi.org/10.1152/jn.00815.2013

Parikh, P. J., Fine, J. M., & Santello, M. (2020). Dexterous Object Manipulation Requires Context-

Dependent Sensorimotor Cortical Interactions in Humans. *Cerebral Cortex*, *30*(5), 3087–3101.

https://doi.org/10.1093/cercor/bhz296

Puglisi, G., Leonetti, A., Landau, A., Fornia, L., Cerri, G., & Borroni, P. (2017). The role of attention in

human motor resonance. *PLoS ONE*, 1–21.

Reichelt, A. F., Ash, A. M., Baugh, L. A., Johansson, R. S., & Flanagan, J. R. (2013). Adaptation of lift forces

in object manipulation through action observation. *Experimental Brain Research*, *228*(2), 221–234.

https://doi.org/10.1007/s00221-013-3554-9

Rens, & Davare. (2019). Observation of Both Skilled and Erroneous Object Lifting Can Improve Predictive

Force Scaling in the Observer. *Frontiers in Human Neuroscience*, *13*.

https://doi.org/10.3389/fnhum.2019.00373

Rens, G., Orban de Xivry, J., Davare, M., & van Polanen, V. (2021a). Lift observation conveys object

weight distribution but partly enhances predictive lift planning. *Journal of Neurophysiology*.

Rens, G., Orban de Xivry, J., Davare, M., & van Polanen, V. (2021b). Motor resonance is modulated by an

object's weight distribution. *Neuropsychologia*, *156*(March).

https://doi.org/10.1101/2020.10.11.335000

Rens, G., van Polanen, V., Botta, A., Gann, M., Orban de Xivry, J., & Davare, M. (2020). Sensorimotor

expectations bias motor resonance during observation of object lifting: The causal role of pSTS.

Journal of Neuroscience, *40*(20), 3995–4009. https://doi.org/10.1523/JNEUROSCI.2672-19.2020

Rossi, S., Hallett, M., Rossini, P. M., & Pascual-leone, A. (2011). Screening questionnaire before TMS: An

update. *Clinical Neurophysiology*, *122*(8), 1686. https://doi.org/10.1016/j.clinph.2010.12.037

Schabrun, S. M., Ridding, M. C., & Miles, T. S. (2008). Role of the primary motor and sensory cortex in

precision grasping: A transcranial magnetic stimulation study. *European Journal of Neuroscience*, *27*(3), 750–756. https://doi.org/10.1111/j.1460-9568.2008.06039.x

Semmler, J. G., Kornatz, K. W., Dinenno, D. V., Zhou, S., & Enoka, R. M. (2002). Motor unit synchronisation is enhanced during slow lengthening contractions of a hand muscle. *Journal of Physiology*, *545*(2), 681–695. https://doi.org/10.1113/jphysiol.2002.026948

Senot, P., D'Ausilio, A., Franca, M., Caselli, L., Craighero, L., & Fadiga, L. (2011). Effect of weight-related labels on corticospinal excitability during observation of grasping: A TMS study. *Experimental Brain Research*, *211*(1), 161–167. https://doi.org/10.1007/s00221-011-2635-x

Snow, J. C., & Culham, J. C. (2021). The Treachery of Images: How realism influences brain and behavior. *Trends in Cognitive Sciences*, *xx*(xx), 1–14. https://doi.org/10.1016/j.tics.2021.02.008

Suzuki, T., Sugawara, K., Takagi, M., & Higashi, T. (2015). Excitability changes in primary motor cortex just prior to voluntary muscle relaxation. *Journal of Neurophysiology*, *113*(1), 110–115. https://doi.org/10.1152/jn.00489.2014

Van Elswijk, G., Kleine, B. U., Overeem, S., & Stegeman, D. F. (2007). Expectancy induces dynamic modulation of corticospinal excitability. *Journal of Cognitive Neuroscience*, *19*(1), 121–131. https://doi.org/10.1162/jocn.2007.19.1.121

Van Polanen, V., Rens, G., & Davare, M. (2020). Control of Movement The role of the anterior intraparietal sulcus and the lateral occipital cortex in fingertip force scaling and weight perception during object lifting. *Journal of Neurophysiology*, 557–573. https://doi.org/10.1152/jn.00771.2019

van Polanen, Vonne, & Davare, M. (2015). Sensorimotor Memory Biases Weight Perception During Object Lifting. *Frontiers in Human Neuroscience*, *9*(December), 700. https://doi.org/10.3389/fnhum.2015.00700

Wilson, S. A., Lockwood, R. J., Thickbroom, G. W., & Mastaglia, F. L. (1993). The muscle silent period following transcranial magnetic cortical stimulation. *Journal of the Neurological Sciences*, *114*(2), 216–222. https://doi.org/10.1016/0022-510X(93)90301-E

Wolpert, D. M., & Flanagan, J. R. (2016). Computations underlying sensorimotor learning. *The Journal of Biological Chemistry*, *270*(45), 26723–26726.

Young, M. S., Triggs, W. J., Pendergast, J. F., & Heilman, K. M. (2000). The effects of selective response preparation on corticospinal excitability. *Cortex*, *36*(1), 19–29. https://doi.org/10.1016/S0010-9452(08)70833-5

Zhang, H., Daw, N. D., & Maloney, L. T. (2015). Human representation of visuo-motor uncertainty as mixtures of orthogonal basis distributions. *Nature Neuroscience*, *18*(8), 1152–1158.

1041   https://doi.org/10.1038/nn.4055

1042