Target-agnostic discovery of Rett Syndrome therapeutics by coupling

computational network analysis and CRISPR-enabled *in vivo* disease modeling

Novak, R.[1,†,*], Lin, T.[1,*], Kaushal, S.[1], Sperry, M.[1], Vigneault, F.[1,†], Gardner, E.[1,†], Loomba, S.[1],

Shcherbina, K.[1], Keshari, V.[1], Dinis, A.[1], Vasan, A.[1], Chandrasekhar, V.[1], Takeda, T.[1], Turner,

J.R.[2], Levin, M.[1,3], Ingber, D.E.[1,4,5,#]

[1] Wyss Institute for Biologically Inspired Engineering, Harvard University, Boston, MA, USA

[2] Laboratory of Mucosal Barrier Pathobiology, Department of Pathology, Brigham and Women's
Hospital and Harvard Medical School, Boston, MA, USA

[3] Allen Discovery Center at Tufts University, Medford, MA, USA

[4] Harvard John A. Paulson School of Engineering and Applied Sciences, Harvard University,
Cambridge, MA, USA

[5] Vascular Biology Program and Department of Surgery, Boston Children's Hospital and Harvard
Medical School, Boston, MA, USA

[†] Present address: Unravel Biosciences, Inc., Boston, MA, USA

* Contributed equally to this manuscript

Corresponding author: Donald E. Ingber, MD, PhD

20   Wyss Institute at Harvard University

21   CLSB5, 3 Blackfan Circle

22   Boston, MA, 02115 USA

23   E-mail: don.ingber@wyss.harvard.edu

ABSTRACT

      Many neurodevelopmental genetic disorders, such as Rett syndrome, are caused by a single gene mutation but trigger changes in expression and regulation of numerous other genes. This severely impair functions of multiple organs and organ systems beyond the central nervous system (CNS), adding to the challenge of developing broadly effective treatments based on a single drug target. This challenge is further complicated by the lack of sufficiently broad and biologically relevant drug screens, and the inherent complexity in identifying clinically relevant targets responsible for diverse phenotypes. Here, we combined human gene regulatory network-based computational drug prediction with *in vivo* screening in a population-level diversity, CRISPR-edited, *Xenopus laevis* tadpole model of Rett syndrome to carry out target-agnostic drug discovery, which rapidly led to the identification of the FDA-approved drug vorinostat as a potential repurposing candidate. Vorinostat broadly improved both CNS and non-CNS (e.g., gastrointestinal, respiratory, inflammatory) abnormalities in a pre-clinical mouse model of Rett syndrome. This is the first Rett syndrome treatment to demonstrate pre-clinical efficacy across multiple organ systems when dosed after the onset of symptoms, and network analysis revealed a putative therapeutic mechanism for its cross-organ normalizing effects based on its impact on acetylation metabolism and post-translational modifications of microtubules. Although traditionally considered an inhibitor of histone deacetylases (HDAC), vorinostat unexpectedly restored protein acetylation across both hypo- and hyperacetylated tissues, suggesting non-HDAC-mediated therapeutic mechanisms supported by proteomic analysis.

INTRODUCTION

Rett Syndrome (Rett) is a neurodevelopmental disorder that is also known to clinically

impact multiple organs and systems in the body. It is primarily caused by mutations of the gene

encoding methyl-CpG binding protein (MeCP2), a ubiquitous nuclear protein that acts as a

transcriptional repressor and activator affecting hundreds of genes across the genome

(Chahrour and Zoghbi, 2007). Rett is characterized by a high degree of cellular heterogeneity,

with a mixture of wild type and MeCP2 mutant cells within the tissues of the same individual

(Renthal et al., 2018) that leads to subtle but widespread gene misregulation (Gabel et al.,

2015). Clinical criteria and research on Rett have primarily focused on neuromotor, social, and

cognitive impairments and other CNS symptoms (e.g., seizures, repetitive purposeless

movements, regression, microcephaly, loss of speech, autistic features)(Colvin et al., 2003; Kerr

et al., 2001). However, these patients also commonly exhibit multiple clinical symptoms outside

of the CNS, including respiratory dysfunction, gastrointestinal issues, systemic inflammation,

disrupted metabolism, circadian rhythm and sleep disturbances, abnormal bone density, kidney

function, nociception derangement, which are a major source of morbidity that contribute to a

shortened lifespan (Hagberg, 2002; Vashi and Justice, 2019). It is important to note that

although Rett is considered a neurodevelopmental disorder and therapeutics development has

focused on CNS-related endpoints, these system-wide symptoms are often the most disturbing

to patients and families, and thus there is a need to ameliorate symptoms of this disease across

the entire body.

Monogenic diseases are theoretically amenable to gene therapy treatment, and Rett has

been shown to be reversible in a mouse model of X chromosome reactivation (Przanowski et

al., 2018) and post-natal expression of MeCP2 (Giacometti et al., 2007; Guy et al., 2007).

However, even if current challenges around delivery efficacy and safety could be overcome,

Rett and other X-linked disorders require gene dosage compensation to avoid toxicity due to

overexpression of the corrected gene since even a modest 20% deviation of MeCP2 expression

levels can lead to Rett symptoms (Collins et al., 2004; Luikenhuis et al., 2004). Therefore,

pharmaceutical treatments for Rett and other neurodevelopmental disorders are still needed.

Trofinetide, a synthetic IGF-1 C-terminal tripeptide activating BDNF expression, has shown

initial promise in phase II clinical trials and it progressed to phase III trials but demonstrated only

moderate efficacy (Glaze et al., 2017). Sarizotan, a 5-HT1A and D2 receptor agonist with

promising efficacy in mouse models of Rett (Abdala et al., 2014), was recently terminated after

clinical trial results failed to demonstrate efficacy (NCT02790034). Esketamine, an N-methyl D-

aspartate receptor (NMDAR) agonist, is also being evaluated in the clinic (NCT03633058) due

to its reduction of neuroexcitatory glutamate. But these and other therapeutic paths for Rett

pursued thus far target only a subset of patients' CNS symptoms and have not shown any

effects on extra-CNS morbidity.

Here we present an integrated approach combining a novel gene expression network

computational model to predict existing drugs and screening in an in vivo phenotypically diverse

model of Rett syndrome generated in *Xenopus* tadpoles to assess whole-body efficacy. Our

results led to the identification of vorinostat, which showed significant efficacy in a mouse Rett

syndrome model even when initiating treatment after the onset of symptoms, demonstrating the

utility of the target-agnostic platform for predicting drugs to restore multi-organ system function

in Rett syndrome and potentially other complex diseases.

METHODS

***X. laevis* care and rapid generation of Rett syndrome tadpole models**

Xenopus embryos and tadpoles were housed at 18°C with a 12/12 h light/dark cycle in

0.1X Marc's Modified Ringer's (MMR) medium. Tadpoles were fed 3x/week with sera Micron

Nature fry food. All animal experiments and procedures were reviewed and approved by the

Harvard Medical School (HMS) Institutional Animal Care and Use Committee regulations.

Xenbase (Nenni et al., 2019) (www.xenbase.org) was used to search for MeCP2 homolog in

Xenopus laevis (*Xenopus laevis* J-strain 9.2 version). MeCP2.L (XB-GENE-17346751),

MeCP2.S (XB-GENE-494744), found on Chromosome 8L and 8S, show 75% and 70%

sequence identity with human MeCP2, respectively. An important biological distinction is that

while human MeCP2 is found on the X chromosome, Chromosome 8 is not part of the sex

determination alleles of homomorphic chromosome pairs in *X. laevis*. Cas9 target sites were

selected using CHOPCHOP (Labun et al., 2016) on the *X. laevis* J-strain 9.2 version, which

facilitated the selection of guide RNA with no predicted off-targets. We used the default settings

to generate a list of target sequences. We selected 4 targets from this list for each of the L and

S forms. This selection was based primarily on the ranking provided by CHOPCHOP and as a

function of their location on MeCP2 to target exons coding for the methyl-CpG-binding domain

(MBD, exons 2 and 3) and Transcriptional repression domain (TRD, exon 3). The 8 sgRNA

sequences are presented in **Supplementary Table 9** and were synthesized as modified sgRNA

by Synthego Inc.

The sgRNA were resuspended to 100 µM in 0.1X Tris EDTA (pH 8.0). Then, an

equimolar sgRNA mix was made using 2 µl of each sgRNA, for a total of 16 ul. The Cas9 RNP

complex was formed by mixing 75 pmol of the sgRNA mix with 75 pmol of Cas9 in annealing

buffer (5 mM HEPES, 50 mM KCl, pH 7.5), for a total volume of 100 µL, then incubated at 37°C

for 10 minutes. The RNP was kept frozen at -20°C until the day of the injection. To generate

MeCP2 knockdown models of Rett syndrome, *Xenopus* embryos were fertilized and maintained

at 14°C until the 4-cell stage. Each cell was injected with ~2 nL RNP per injection resulting in a

final amount of 1.5 fmol of RNP per injection per cell, and we did not see any adverse effect

when controlling for Cas9 or sgRNA at that amount. Note that the 1.5 fmol of ribonucleoprotein

(RNP) used per injection is two to five time less than what has been commonly used in

previously published studies for *Xenopus* (Aslan et al., 2017) as we found that higher amounts

of sgRNA had adverse effects (not shown). Following injections, embryos were kept in 0.1X

MMR at 18°C as described above, for the 18 days duration of the experiment.

PCR and fragment analysis to validate MeCP2 gene disruption

MeCP2 editing efficiency was measured using Indel Detection by Amplicon Analysis

(Yang et al., 2015), which leverages PCR using a pair of primers and universal fluorescein

labeled oligo (**Supplementary Table 10**).The primers were synthesized by IDT. The amplicons

were purified (Zymo Research) and submitted for fragment analysis on an ABI 3730XL

(performed by Genewiz) following the IDAA method described previously. ProfileIT IDAA

analysis software (COBO Technologies) was used to analyze the fragments and visualize the

distribution of indels and visualize out-of-frame indels (Lonowski et al., 2017). IDAA is a robust

method with 1-bp resolution suited for the analysis of a tetraploid genome without the need for a

deep sequencing approach.

We analyzed the indels of 3 target sites on MeCP2.S of 14 tadpoles presenting with

seizures and 3 non-edited tadpoles. Using a window of +/- 100 bp from the target site, we

observed varied editing efficiencies depending on the target site. The likely out-of-frame indels

ratio averaged 47% for Xl-mecp2s-g01 on exon 3, with the lowest being 12% for a given tadpole

and highest being 95%, 48% for Xl-mecp2s-g02 on exon 2 (min/max 20-84%), and ~82% for Xl-

mecp2s-g04 on exon 2 (min/max 7-91%). Overall, 98% of indels where between 1 to 30 bp, and

84% were deletions.

Microarray analysis

Tadpoles were sacrificed at stage 50 of development and lysed using a QIAGEN

TissueLyser II bead mill (30 Hz, 2 x 30 sec) in a 2 mL tube with a 2.4 mm steel beads (Omni)

and 1 mL phosphate buffered saline buffer at 4°C. 250 mL of lysate was used for RNA

extraction using a QIAGEN RNeasy mini kit. RNA samples were DNAse treated. *Xenopus laevis*

Genome 2.0 Array (Affymetrix) microarrays were processed by Advanced Biomedical

Laboratories (Cinnaminson, NJ). RNA samples were processed using a Nugen Ovation PICO

WTA System V2 kit. The resulting cDNAs were purified using a Qiagen MinElute PCR

Purification Kit following the modifications outlined in the Nugen protocol. The cDNAs were

fragmented and labeled using a Nugen Encore Biotin Module. Hybridization solutions were

prepared by combining the fragmented, biotin-labeled cDNAs with hybridization cocktail

(Affymetrix Hybridization, Wash, and Stain Kit). The mixtures were incubated in a thermal cycler

at 99 °C for 2 min followed by 45 °C for 5 min. then loaded on *Xenopus laevis* Genome 2.0

arrays and incubated for 16-20 hours at 45 ºC and 60 rpm in an Affymetrix Hybridization Oven

645. Following hybridization, arrays were washed and stained on Affymetrix Fluidics Station

450s using the Affymetrix FS450_0001 protocol with the stains and buffers supplied in the

Affymetrix Hybridization, Wash, and Stain Kit. The stained arrays were scanned at 532 nm

using an Affymetrix GeneChip Scanner 3000.

Transcriptomics Data Processing

Microarray datasets were RMA normalized. RNAseq gene counts were normalized using

size factors similar to the normalization performed by the R package DESeq. Genes missing

values were removed and duplicates were combined by taking the maximum across counts.

Gene expression heatmaps and hierarchical clustering plots were generated for all groups using

the R package limma (version 3.42.2) and Python package seaborn (version 0.9.0). Volcano

plots were generated using the Python package bioinfokit (version 0.7).

nemoCAD Drug Prediction Algorithm

In the present study, we developed the nemoCAD computational tool to predict drugs

that would shift MeCP2-edited tadpoles to a control tadpole state based on transcriptome state

for each condition. nemoCAD utilizes pre-computed interaction probabilities of drug-gene and

gene-gene interactions and differential gene expression signatures of a disease state and

appropriate control to identify compounds capable of changing a transcriptional signature

indicative of one biological state to another state of interest (e.g., reverting a disease state to a

healthy state).

The repurposing algorithm is first used to identify transcriptome-wide differential

expression profiles between two biological states in the input transcriptomic dataset

(experimental or published) and to define the target normalization signature, i.e., the subset of

genes whose expression levels need to be reversed in order to revert one state to the other

(e.g., normalize the diseased state). Pairwise analysis, that implicitly assumes genes are

expressed independently of one another, is carried out based on the comparison between gene

expression profiles of >19,800 compounds found in the LINCS database release v1

(Subramanian et al., 2017) and the target normalization signature. Multiple correlation statistics

(e.g., Pearson correlation, cross-entropy) across all differentially expressed genes are

calculated and a combination score (e.g., Pearson correlation divided by the cross-entropy) is

computed.

Putative predictions that incorporate gene-gene dependencies are then generated

separately based on Bayesian network analysis on a regulatory and drug-gene interaction

network architecture defined using publicly available databases of gene-gene interactions based

on single gene knockout datasets in human cells (KEGG, TRRUST) and reference

transcriptional signatures of drugs (LINCS, CTD), as described below. By combining a directed

unweighted network structure with interaction probabilities for the connecting network edges,

the constructed network is a weighted directed graph consisting of all possible paths that

connect at least 2 genes of interest from the relevant genes within the target transcriptomic

normalization set and the drugs; it therefore encodes the entire region of influence of a given list

of genes and the drugs that can reverse their gene expression profiles in the desired manner.

This network is then used as input for a message-passing algorithm (e.g. loopy belief

propagation algorithm (Forbes, 2021; Pearl, 1982)) in which the marginal probability

distributions of drugs being "on" given the expression state of every gene are computed using

the joint probability distribution encoded in the . This network-based approach leads to a ranking

of all the drugs based on the probability of them inducing the desired transcriptomic signature

across all genes in the network, which is separate from the initial ranking based on correlation

analysis described above. This method has the advantage of incorporating gene-gene

dependencies, but its success relies on the quality of network structures curated in the source

databases.

Ultimately, drugs are prioritized that have both high correlation with the desired

transcriptional signature change, representing desired drug-gene interactions, and high

probabilities of being able to reverse the diseased state predicted by the Bayesian network

analysis, representing gene network-level drug effects that can counter network-level disease

processes. nemoCAD also enables visualization of the architecture of the network and

subnetworks, and thus provides insight into potential molecular targets that can be verified using

additional experiments. nemoCAD also can be used to optimize treatment regiments by

iteratively inputting animal omics data following drug treatment to further tune drug predictions

and develop drug combinations that might synergize in unpredictable ways. The approach

underlying nemoCAD avoids the needs for *a priori* target or pathway inputs and, in fact, enables

their discovery.

Transcriptomics Gene Network Construction

Gene regulatory networks were inferred from gene expression data using the

Bioconductor package GEne Network Inference with Ensemble of trees (GENIE3) release 3.12

in RStudio (R version 3.6.2)(Huynh-Thu et al., 2010). GENIE3 infers a weighted, directed gene

regulatory network from expression data by decomposing the prediction of a network between *k*

genes into *k* different regression problems. For each individual regression analysis, the

expression pattern of one of the genes (target gene) is predicted from the expression patterns of

all the other genes (regulator genes) using ensembles of regression trees. GENIE3 produces an

adjacency matrix representation of the network with *k* nodes in which each node represents a

gene, and an edge directed from one gene *i* to another gene *j* indicates that gene *i* regulates the

expression of gene *j*. Based on known mechanisms involved in Rett Syndrome

(**Supplementary Table 1**), a subset of genes was selected as candidate regulators: BDNF,

FCRL2, FMR1, MeCP2, NTRK2, PER1, PER2, and PUM1. These potential regulators were

calculated over all available target genes in the transcriptomics datasets (10,934 genes) for

MeCP2 knock down and cas9-injected control samples. Digraph objects were constructed from

adjacency matrices and visualized in Matlab R2020a (Mathworks; Natick, MA) for each of the

MeCP2 knock down and control groups. The strongest regulator gene-target gene relationships

were highlighted by filtering each network at a consistent edge weight (e_w) threshold (e_w = 0.3)

prior to plotting.

Cross-Species Transcriptomics Analysis

Gene networks involved in Rett Syndrome in human patients and MeCP2 knock down in

laboratory animal models were compared using previously acquired transcriptomics datasets

(**Supplementary Table 2**). Transcriptomics datasets were processed using the same methods

described for *Xenopus* transcriptomics and normalized using a series of transformations. Since

RNA-seq data typically have a wider dynamic range than microarray datasets, RNA-seq and

microarray data was transformed into a common space (Sekhon et al., 2013). RNA-seq

datasets were transformed using a hyperbolic sine function and the $\log_2$ transformation was

applied to microarray data (Sekhon et al., 2013). Subsequently, a variance stabilizing

transformation was applied to remove potential mean-variance relationships in each dataset

(Russo et al., 2018). Common genes expressed and measured between datasets (926 genes)

were considered for further analysis. Dimensional reduction was performed using Principle

Component Analysis (PCA) and visualized using Uniform Manifold Approximation and

Projection (UMAP) in Python with the help of sklearn and UMAP packages.

Commonalities in gene expression values across species were analyzed by plotting

heatmaps of differential gene expression using the Python packages seaborn and matplotlib. All

datasets were normalized as described above and log fold change for each gene was

calculated by combining samples of a condition using geometric mean. Log fold changes for all

species were combined and the genes (96 genes) with variance greater than the mean variance

(in fold changes) were used to construct the heatmaps.

Transcriptomics gene networks were constructed using GENIE3 and the same methods

previously described. To investigate potential changes across the entire gene network, regulator

genes were *not* down selected. Adjacency matrices were visualized for each network and the

distribution of edge weights in the network was assessed by histogram and boxplots in Matlab.

Digraph objects were constructed from adjacency matrices and visualized in Matlab for the

strongest connections in each network (e_w > 99.99% for each network). Resultant networks

were combined across all datasets for each species and plotted with different colors designating

each dataset. A cross-species network that includes the strongest network connections across

all datasets was formed by combining each species-specific dataset and plotted with different

colors designating each species. To identify the strongest regulator and target gene nodes in

the resultant cross-species network, the in-degree and out-degree were calculated in Matlab.

The in-degree is defined as the number of edges with that node as the target and out-degree of

a node is equal to the number of edges with that node as the source.

Drug screening

Drugs for tadpole screening were purchased from Sigma-Aldrich (St. Louis , MO, USA)

or MedChem Express (Monmouth Junction, NJ, USA) and dissolved in DMSO to a stock

concentration of 100-1,000 μM, depending on the solubility. For screens, compounds were

diluted to their final concentrations with a 0.1% DMSO concentration. Media with dosed drug

were made fresh and exchanged every two days following feeding. Dosing began at stage 45

and lasted for 7 days, after which standard 0.1X MMR was used to evaluate drug washout

effects. Free-swimming tadpoles were imaged in 60 mm dishes using a SONY Alpha a6100

camera with 16mm objective against an illuminated background. Tadpole behavior was scored

manually by noting the most severe seizure-related phenotype described by Hewapathirane et

al. (Hewapathirane et al., 2008) within a 20 min period.

Vorinostat and trofinetide for mouse studies were purchased from MedChem Express

(Monmouth Junction, NJ, USA). For intraperitoneal injection, trofinetide was solubilized in

DMSO, and vorinostat was solubilized in a method previously published(Basu et al., 2019;

Hockly et al., 2003). 2-hydroxypropyl-β-cyclodextrin powder (HPβCD) was purchased from

Acros Organics (ThermoFisher, Waltham, MA, USA). Vorinostat was dissolved in 100 mM

HPβCD by boiling for 5 minutes. Mice were dosed intraperitoneally at 100mg/kg with Trofinetide

and 50 mg/kg with vorinostat. For oral dosing, commercial instructions were followed to add

solubilized vorinostat (100 mg/kg) to Medigel (purchased from ClearH20, Westbrook, ME, USA).

X. laevis **Tissue Processing**

After stage 46 tadpoles were euthanized with 20x Tricaine, tadpoles were washed three

times with PBS (-/-) and fixed in 4% Paraformaldehyde in PBS + Mg + EGTA (MEMFA) (Alfa

Aesar, Tewksbury, MA, USA) for 2 hours at RT. For sections, *Xenopus* tadpoles were

cryoprotected by sequential incubation of 10%, 20% and 30% sucrose in 1x PBS(-/-), only

transferring to the subsequent concentration when the samples sunk to the bottom of the

scintillation vial. Cryoprotected tadpoles were embedded in 5% agarose and were stored in -

80°C. For GI staining, the gastrointestinal tract was microdissected with a stereoscope before

embedding. Embedding tissue were sectioned coronally at 20 μm thickness on a Cryostat

(Leica, CM3050 S, Wetzlar, Germany) and mounted on superfrost plus slides (Thermo Fisher,

Waltham, MA, USA). Slides were dehydrated for 30 minutes at 20°C, before storing at -80°C.

For whole mount preparation, tadpoles were processed in a method previously published

(Willsey, 2021).

***In situ* hybridization**

Following X. *laevis* tissue processing of cryosections, sample pretreatment, RNAscope

Target Retrieval and the RNAscope Assay was conducted by the HMS Neurobiology Imaging

Facility following manufacturer's instructions. Xl.MeCP2 was custom designed by Advanced Cell

Diagnostics. Rpl8 was used as a positive control (Xl-LOC108706872-C3, Cat No. 516501-C3,

Advanced Cell Diagnostics, Hayward, CA) and DapB as a negative control (Cat No. 320758,

Advanced Cell Diagnostics, Hayward, CA).

Immunohistochemistry of *Xenopus laevis* sections and whole mounts

Following tissue processing, cryosections were blocked for 2 hours at room temperature

and subsequently incubated with primary antibodies overnight at 4°C, including mouse anti-

alpha acetylated tubulin (1:1000 dilution, Sigma Aldrich, Burlington, MA USA), mouse anti-beta-

tubulin (1:1000 dilution, E7, Developmental Studies Hybridoma Bank, Iowa City, Iowa, USA),

anti-MeCP2 (1:1000 dilution, Invitrogen, Waltham, MA USA), Hoechst (1:1000 dilution,

ThermoFisher, Waltham, MA, USA), negative controls anti-mouse IgG (1:1000 dilution,

Invitrogen, Waltham, MA, USA), and anti-rabbit (1:1000 dilution, Invitrogen, Waltham, MA USA),

as well as FITC-conjugated isolectin B_4 (1:1000 dilution, Thermo Fisher, Waltham, MA, USA).

Slides were washed twice with TRIS-buffered saline, with 1% Tween-20 (TBT) for 5 minutes

each and incubated in the dark with their corresponding secondary antibody: Goat anti-rabbit

Alexa 647 (1:1000 dilution, Abcam, Waltham, MA, USA) and Goat anti-mouse Alexa 594

(1:1000 dilution, Abcam, Waltham, MA, USA) for 1 hour at room temperature. Slides were

washed three times with TBT for 5 minutes each on an orbital shaker and mounted in ProLong

Gold Antifade Mountant (ThermoFisher, Waltham, MA, USA) with a #1 coverslip and sealed.

Slides were stored in the dark at 4°C before image acquisition. Z-stacks were acquired at $1\mu m$

to visualize brain and gastrointestinal sections.

X. laevis embryos were processed for whole mount immunofluorescence as described

(Willsey, 2021). Fixed tadpoles were quenched for 1 hour in a lightbox in 5% formamide and 4%

hydrogen peroxide in PBS, permeabilized for 1 hour in PBS with 0.1% Triton X-100 (PBT) and

blocked for 1 hour in 10% CAS-Block in PBT (Life Technologies, Carlsbad, CA, USA). Primary

antibodies were incubated overnight at 4°C, including mouse anti-alpha acetylated tubulin

(1:700 dilution Sigma Aldrich, Burlington, MA USA), mouse anti-beta-tubulin (1:100 dilution, E7,

Developmental Studies Hybridoma Bank, Iowa City, Iowa, USA)), Hoechst (1:100 dilution,

ThermoFisher, Waltham, MA, USA) and anti-mouse IgG (1:100 dilution). Slides were washed in

PBT, blocked in 10% CAS-Block for 30 minutes, and incubated with the corresponding

secondary antibody for 2 hours at room temperature including Goat anti-mouse Alexa 546

(1:100 dilution, Thermo Fisher, Waltham, MA, USA) and Goat anti-rabbit 633 (1:100 dilution,

ThermoFisher, Waltham, MA, USA). Tadpoles are washed with PBT for 1 hour and then

washed in PBS for another hour before mounting in Vectashield (Vector Laboratories,

Burlingame, CA, USA). Z-stacks were acquired at 1μm for visualizing epidermal MCC's and

3μm for visualizing entire brain regions.

Drug target discovery in *X. laevis* using thermal proteome profiling

Sample preparation

While we largely followed the protein integral solubility alteration (PISA) thermal

proteome profiling protocol (Gaetani et al., 2019), we adapted the mammalian cell- and tissue-

specific parameters to tadpoles by optimizing the method to tadpoles by reducing thermal

treatment temperatures to account for the lower normal temperature range of *X. laevis*.

Unmodified *X. laevis* tadpoles (stage 47-50) were exposed to 25 μM vorinostat or 0.1 μM

ivermectin for 2 h while freely swimming in 0.1x MMR medium. Following euthanasia using

Tricaine, animals were washed with fresh MMR, cut into thirds (head, abdomen containing all

visceral organs, and tail), and placed into individual Eppendorf tubes covered with MMR

containing the same drug and concentration and incubated for 3 min at one of 9 temperatures

spanning 30-60°C. 30°C was selected as the control temperature, where no appreciable thermal

denaturation should take place, based on the maximum tolerable temperature for *X. laevis*

(Ruthsatz et al., 2018). Following aspiration of remaining liquid, samples were flash frozen in

liquid nitrogen.

Samples were further processed and analyzed by Phenoswitch Bioscence, Inc.

(Sherbrooke, Québec, Canada). Thermally-treated samples were lysed in 100 µL PBS with

0.4% NP-40 with 4 cycles of freeze/thaw. Insoluble material was cleared by centrifugation (10

min, 10,000 G, 4°C). 11µL of supernatant for each of the 9 temperatures were then pooled in

one tube and samples were centrifuged again at 13,000 RPM for 75 minutes, at 4°C to pellet

precipitated proteins. 80 µL of the supernatant were reduced with 10 mM DTT for 15 min at

65°C and alkylated with 15 mM IAA and 30 min at room temperature in the dark. Proteins were

precipitated with 8 volumes of ice-cold acetone and 1 volume of ice cold methanol overnight.

Protein pellets were washed 3 times with 250 µl of ice-cold methanol and resuspended in 100

µL digestion buffer. Digestion was carried for 4 hours in 50 mM Tris ph 8 + 0.75 mM Urea + 1µg

trypsin/LysC at 37°C with agitation. Another 1 µg of trypsin/LysC was added, and digestion was

continued overnight. Peptides were purified by reversed phase SPE and analyzed by LC-

MS/MS.

Mass spectrometry

Acquisition was performed with a TripleTOF 6600 (Sciex, Foster City, CA, USA)

equipped with an electrospray interface with a 25 µm iD capillary and coupled to a Micro LC200

(Eksigent, Redwood City, CA, USA). Analyst TF 1.8 software was used to control the

instrument. Acquisition was performed in data independent acquisition (DIA or SWATH) using

gas phase fractionation (GPF1 from 350 m/z to 800 m/z and GPF2 from 800 m/z to 1250 m/z).

The source voltage was set to 5.5 kV and maintained at 325oC, curtain gas was set at 45 psi,

gas one at 25 psi and gas two at 25 psi. Separation was performed on a reversed phase

Kinetex XB column 0.3 µm i.d., 2.6 µm particles, 150mm long (Phenomenex) which was

maintained at 60°C. Samples were injected by loop overfilling into a 5 µL loop. For the 60

minutes LC gradient, the mobile phase consisted of the following solvent A (0.2% v/v formic acid

and 3% DMSO v/v in water) and solvent B (0.2% v/v formic acid and 3% DMSO in EtOH) at a

flow rate of 3 µL/min. Both GPF files were analyzed on a previously-generated 3D ion library

using the SWATH 2.0 microapp from Peakview (Sciex, Foster City, CA, USA). Each GPF file

was analyzed using 10 peptides per protein, 4 MS/MS transition per peptide, 12.5 min RT

window and 25 ppm XIC width. The reported quantification for a protein is the sum of all the

correctly integrated peptides (FDR<0.05) in both GPF files.

Mouse care

Experiments were performed with MeCP2-null ($MeCP2^{-/Y}$) (strain 003890) and age-

matched WT male littermates ($MeCP2^{+/Y}$), purchased from Jackson Laboratories (Bar Harbor,

ME). All mice were housed in ventilated racks under specific-pathogen-free conditions at a room

temperature in a 12/12 h light/dark cycle with food and water ad libitum. All animal experiments

and procedures were reviewed and approved by the Harvard Medical School (HMS) Institutional

Animal Care and Use Committee regulations. Every effort was made to minimize their suffering.

Mice were assessed for disease severity and endpoint criteria on a daily basis by weighing and

using the phenotypic scoring method developed by the Bird laboratory (Guy et al., 2007;

Szczesna et al., 2014), which includes assessing mobility, gait, breathing, tremor, and general

condition. Endpoint was determined by 20% body mass loss or a score of 2 in criteria D, E, or

F. Dosing and Behavioral procedures began when mice reached 4 weeks of age.

Behavioral Testing

Elevated Plus Maze (EPM)

The elevated plus maze (EPM) consisted of two open and two closed (34 cm long and 5

cm wide) arms extended out from a central platform 50 cm above the floor. The test was carried

out in dim ambient lighting. The room of the experiment and its spatial cues were kept

consistent and minimal. Mice were habituated in the test room for 30 minutes before the start of

the test. Mice were placed near the center compartment of the maze, facing an open arm, and

allowed to explore the apparatus for 5 minutes. After each test, the maze was cleaned

thoroughly with 70% ethanol and left to dry before the start of the next test. A computer-assisted

video-tracking system (EthoVision XT 14, Noldus, Leesburg, VA) was used to record the

number of open and closed arm entries as well as the total time spent in open, closed, and

center compartments. An increase in the percent time spent or entries into the open arms was

used as a surrogate measure of anxiolytic-like behavior (Carobrez and Bertoglio, 2005).

Spatial Novelty Y-Maze (Y-maze)

The spatial novelty y-maze consisted of a Y-shaped maze with three arms at a 120° from

each other as well as a removable blockade used to prevent access to one of the arms during the

Habitutation phase. The test was carried out in dim ambient lighting. The room of the experiment

and its spatial cues were kept consistent. One of the three arms was defined as the start arm,

where the mice would be placed at the start of the experiment and was kept consistent throughout

the entire study. The test consisted of a three-minute Habituation phase where one of the two

non-start arms was blocked off, a two-minute intertrial interval (ITI) outside of the maze and a

three-minute Test phase, where the previously blocked arm was exposed to the mouse. The arm

that was blocked off during the Habituation phase was randomized and noted for each mouse.

Mice were kept in an adjacent room to not disturb the test. The test mouse was placed in

a holding cage and brought to the test room with the maze and immediately placed into the Y-

maze at the designated start arm and allowed to explore the start and familiar arm trial for three

minutes. At the end of the habituation phase, the mouse was placed into the holding cage for 2

minutes ITI. During this time, the blockade was removed, and the maze was cleaned with an

ammonia-based cleaner with a paper towel and left to dry to remove odor cues. The mice were

then placed back in the start arm and left to explore the entire maze for three minutes. After the

test trial, the mouse was transported back to its home cages and the maze was again wiped dry

with an ammonia based cleaner. A computer-assisted video tracking system (EthoVision XT 14,

Noldus, Leesburg, VA) was used to record the number of entries, time spent in each arm and as

distance traveled in each arm during the Habituation and Test Phase. The percentage of time and

distance spent in the novel, or previously blocked arm were used to assess spatial novelty seeking

behavior (Dellu et al., 2000).

Mouse Histology and Immunostaining

Mice were euthanized with CO_2 inhalation, transcardially perfused with PBS (-/-) and 4%

PFA, and tissues of interest were removed. Regions of the gastrointestinal tract and lung

processed as was described previously(Morton and Snider, 2017; Nalle et al., 2019; Tsai et al.,

2017). Paraffin blocks were sectioned into 15 μm thickness. Mouse lung tissues were sectioned

coronally and GI tissues were stained with hematoxylin and eosin (H&E) by the Beth Israel

Deaconess Medical Center's IHC Core Facility. The slides were also stained for rabbit anti-

acetylated α-tubulin (ABclonal, ab179484, Woburn, MA, USA) mouse anti-CD-64 (R&D systems,

AF3628, Minneapolis, MN, USA), chicken anti-βIII-tubulin (Biolegend, 801202, San Diego, CA,

USA) immunofluorescence by Beth Israel Deaconess Medical Center's IHC Core Facility.

Mouse brains were fixed with 4% paraformaldehyde at 4°C overnight, cryoprotected in a

sucrose gradient (10%, 20% and 30%), and subsequently embedded in OCT Tissue Tek. The

embedded tissues were subsequently sectioned coronally at 30μm and the free-floating sections

were stained as previously described(Potts et al., 2020). The sections were blocked with 10%

CAS Block in PBT for 1hr, stained with primary antibody (rabbit anti-iba-1, 1:1000, Wako, #019-

19741; mouse anti-NeuN, 1:1000, Abcam, #ab104224; Hoechst, 1:1000, Thermo Fisher, H1399)

overnight at 4°C. After three 10-minute washes in PBS (-/-) on an orbital shaker, the sections were

stained in secondary antibody (anti-rabbit Alexa 647 for iba1, anti-mouse Alexa 594 for NeuN) for

1 hour at room temperature. After three 10-minute washes on an orbital shaker, the sections were

mounted on superfrost plus slides (ThermoFisher, Waltham, MA, USA) in ProLong Gold Antifade

Mountant (ThermoFisher, Waltham, MA, USA) with a #1 coverslip and sealed. Z-stack images

were acquired at 1μm intervals using the 63x/ glycerol.

Multiplex Chemokine Assay

After mouse euthanization, blood was collected by cardiac puncture using at 23G needle

in ethylendiamine tetra-acetic acid (EDTA) treated tubes (Microvette 200 K3EDTA, CAT#

20.1288.100, Sarstedt, Numbrecht, Germany). To avoid hemolysis, the tubes were kept at room

temperature and quickly processed for plasma. Tubes were centrifuged at 2000 x g for 10

minutes and the plasma was separated and stored in -80C until assayed. Chemokine levels

were measured in mouse plasma using the Bio-Plex Pro Mouse Chemokine Panel 31-Plex

(Cat# 12009159, Bio Rad, Hercules, CA, USA). Measurements were performed using the Bio-

Plex 3D Suspension Array System (BioRad, Hercules, CA, USA), following the manufacturer's

instructions.

Microscopy and Image Processing

H&E slides were imaged with 4x and 20x objectives with the Biotek Cytation 5.

Immunofluorescent *Xenopus* whole mounts and *Xenopus* and mouse stained sections were

imaged with the Leica SP5 X MP Inverted Confocal Microscope using 25x (NA 0.95) water

objective and 63x (NA 1.3) Glycerol objectives. Sections were imaged in white light and diode

laser scanning mode and whole mounts were imaged using multiphoton pulsed IR laser

scanning. Serial scanning was used during acquisition to avoid bleed-through. Image settings

were kept constant between groups, in terms of laser power, bit-depth, z-stack interval,

resolution (pixels), aperture settings, and gain/offset. Images were post-processed and

analyzed using ImageJ (Fiji, NIH).

MCC morphological analysis

2-photon images of X. laevis 25x and 63x MCC's were maximum projected, background

subtracted, and subsequently assessed for cilia orientation and cilia length, as previously

described (Chien et al., 2018).

MCC Functional Assay

We visualized the cilia-driven flow of fluorescent microspheres over the surface of the xenopus

tadpole by taking times series with the Leica SP5 x MP inverted confocal. Our methods were

based on previous reports (Kulkarni et al., 2018) with some modifications. Stage 45 tadpoles

were placed in a 24 well μ-Plate 14mm (ibidi, Gräfelfing, Germany), and underwent anesthesia

by incubation with 1x tricaine before proceeding to image. One tadpole was placed per well for

individual tracking. Fluorescent microspheres beads (Thermo Fisher, Waltham, MA) were added

at a final dilution of 1:100. We chose the bead size of 3μm, because during preliminary testing

and as is reflected in the data, WT *X. laevis* tadpoles consistently was able to clear beads that

were 3μm diameter at this concentration. The Leica SP5 x MP laser scanning mode was used

to visualize the fluorescent microspheres beads at an excitation of 520nm and emission of

560nm with the following settings: 512x512 pixels, bidirectional scanning mode, and at a scan

speed of 800-1600Hz, with the 25x (NA 0.95) water objective, and over the time course of 5

minutes every 5s. Time series were acquired of in the olfactory and lateral region of the

epidermis.

Quantification of bead displacement

The time series stacks were maximum intensity projected to visualize the overall bead flow.

Intensity was measured as a function from the distance from the surface of the epidermis.

Mean Intensity Quantitation

To generate a ROI mask, channels acquired in one z-stack were split and the maximum

intensity projection of a 'control' channel was taken. In the case for assessing αaT expression

mouse GI tract and bronchiole, the βIII-tubulin channel was used, and for assessing MeCP2

expression, the α -acetylated tubulin channel was used in the image stack. Subsequently,

background noise was subtracted, and auto-thresholded, whose settings were held constant

between groups. The ROIs generated from the control channel were used to quantify the

maximum projected channel of interest, to obtain the mean intensity (integrated density per

area). A minimum of 3 serial sections were analyzed per animal.

Counting Ib4+ cells

Confocal Z-stacks of *X. laevis* GI tract images were quantified for IB4+ cells, by taking the

maximum intensity projection of the ib4+ channel. Following noise reduction, and auto-

threshold, ib4+ cells were segmented and counted using the particle analysis function in the

ImageJ software (NIH, Bethesda, MD).

Sholl Analysis

Sholl analysis was performed as previously published[47]. Confocal (at 1um intervals) Z-stacks

were maximum intensity projected in the iba-1 channel, noise de-speckled, subsequently

segmented using the auto-threshold function, and microglia were duplicated into 8-bit TIF files,

in the field of view, ensuring that for each mouse at least 10 microglia were selected in the

ipsilateral and contralateral region of the olfactory bulb. Using the Sholl analysis plugin in

ImageJ (NIH, Bethesda, MD), traces and intersections were generated.

Statistical analysis

Statistical analysis of biological results was performed using Graphpad Prism 9.3. For

comparison of two conditions, Student's t-test was used unless otherwise indicated. ANOVA

with pairwise t-tests and Bonferroni correction or the Holm-Šídák tests was used to evaluate

significance of multiple variables. P values <0.05 were considered significant (*, $p < 0.05$; ** $p <$

0.01; *** $p < 0.001$; **** $p < 0.0001$). Tadpole screens and mouse studies were conducted with

5-8 animals each and repeated, except for the mouse oral dosing study.

RESULTS

Modeling Rett syndrome genetic and phenotypic heterogeneity in _Xenopus laevis_

A major challenge to Rett therapeutics discovery lies in the inherent heterogeneity

observed in patients stemming from gene expression variability due to the gene regulatory

impact of distinct MeCP2 mutations, combined with tissue heterogeneity caused by mosaicism

resulting from X chromosome inactivation. As a result, Rett patients exhibit a spectrum of

disabilities. To establish an animal model of Rett that would incorporate the heterogeneity

observed in patients, we used CRISPR to generate a mosaic knockdown of MeCP2 protein in

Xenopus laevis tadpoles, which have been previously used to model neurodevelopmental

diseases due to their evolutionary proximity to mammals and well-characterized neural

development (Exner and Willsey, 2021; Pratt and Khakhalin, 2013).

Injection of _X. laevis_ embryos at 4- or 8-cell stages using cas9 pre-complexed with

sgRNA targeting 6 sites of the MeCP2 gene, including the regions corresponding to the DNA-

binding domain, resulted in a tunable model of MeCP2 knockdown with a heterogeneous

phenotype when assessed at the population level. Based on our target selection and injection

strategy at the 4-cell stage, each individual animal was edited as a mosaic and presented

varying degree of severity, without apparent toxicity from Cas9 or sgRNA. Indeed, all embryos

developed to the swimming tadpole stages (Nieuwkoop-Faber stages 45-50) with no apparent

loss of viability or morphological defect in earlier stages.

Importantly, swimming tadpoles with MeCP2 knock down ("Rett" tadpoles) exhibited a

broad range of abnormal behavior compared to wild type vehicle-injected control tadpoles

("Control") including darting motions and rapid repetitive swimming in tight circles (**Fig. 1a,b and**

Supplementary Movie 1) reminiscent of the repetitive motions observed in human Rett patients

as well as C-shaped or straight rigor (**Supplementary Movie 1**) that has been shown to

correspond to seizures in prior *Xenopus* studies (Hewapathirane et al., 2008). The cas9 protein

and sgRNA doses were titered to achieve between 5 and 95% gene editing efficiency as

determined by qPCR without resulting in toxicity (**Supplementary Fig. 1a**). All tadpoles

exhibited significant MeCP2 gene insertion/deletion polymorphisms (indels) at each of the

mecp2 gene target sites (**Supplementary Fig. 1b**) and reduced MeCP2 RNA expression (**Fig.**

1c). MeCP2 protein levels were also heterogeneously reduced in the brain (**Fig. 1d,e**) and

olfactory multiciliated cells (MCC) (**Fig. 1d,f**). This this editing approach resulted in an animal

model of Rett with intra-subject and inter-subject diversity at molecular and phenotypic scales,

which is similar to what is observed in humans.

In line with prior studies in humans and other model organisms (Sanfeliu et al., 2019, p.

2), when we analyzed the transcriptome-wide effects of MeCP2 knockdown on the Rett

phenotype in *Xenopus*, only subtle shifts in gene expression were detected with 70 out of

10,935 genes probed undergoing a significant change in gene expression ($p_{adjusted} < 0.05$) (**Fig.**

1g, Supplementary Table 1). Of the 70 differentially expressed genes, 37 are involved in

metabolic processes, 9 operate within developmental processes, and 8 are part of signal

transduction pathways (**Supplementary Fig. 2a**); this aligns with changes in genes that control

metabolism or regulate neuronal processes (e.g., ion transport, nervous system development) in

Rett patients (Renthal et al., 2018).

Interestingly, analysis of gene co-expression networks identified substantial

reorganization from control animals (**Fig. 1h**) following MeCP2 knock down (**Fig. 1i**),

characterized by a loss of strong connectivity among genes involved in the regulation of MeCP2

(FMR1 and TRKB receptor NTRK2 (Abuhatzira et al., 2007; Arsenault et al., 2021)), neuronal

development (BDNF (Chang et al., 2006; Li and Pozzo-Miller, 2014)),, and circadian rhythm

(PER1, PER2 (Martínez de Paz et al., 2015)) all of which are altered in Rett patients (Li and

Pozzo-Miller, 2014; Miller et al., 2019). Notably, after MeCP2 knock down, BDNF strengthened

as a network hub and exhibited greater co-expression connectivity amongst gene nodes (**Fig.**

1i).This was unexpected because although BDNF is involved in neuronal development, synaptic

transmission, and plasticity, it declines with the onset of Rett-like neuropathological and

behavioral phenotypes (Li and Pozzo-Miller, 2014). However, BDNF is a known target of

repression by MeCP2 and our findings are consistent with prior work suggesting that

downregulation of BDNF is a later and indirect outcome of MeCP2 deficiency (Chang et al.,

2006; Li and Pozzo-Miller, 2014).

We then comprehensively compared the effects of MeCP2 knock down in *Xenopus* to

changes observed in published transcriptomic data sets from Rett patients as well as MeCP2

knockout mouse and zebrafish models (**Supplementary Table 2**). After performing

dimensionality reduction in this cross-species analysis, *K*-Means clustering applied to the

principal components revealed 3 clusters, with mouse and human clustering together and

zebrafish and *Xenopus* forming 2 independent clusters (**Supplementary Fig. 2b**). Importantly,

diverse gene-expression abnormalities that occur across cell types in Rett syndrome could

contribute to the differences observed across these datasets (Renthal et al., 2018). In this case,

both the human and mouse transcriptomics were performed on brain tissues, whereas the

zebrafish and *Xenopus* data are from whole organisms, which may account for the clustering

patterns observed.

Despite evidence of species-specific clustering, a common gene signature comprised of

96 genes was identified across all species by searching for genes with a fold change variance

greater than mean variance. However, the common genes within this transcriptomic signature

are expressed at similar low fold change levels in 7 of the 9 datasets analyzed, with slightly

greater variations in expression only observed in 1 out of 5 mouse datasets and 1 out of 3

human datasets (**Fig. 1j**). This finding provides additional evidence that mutation of MeCP2

produces widespread, yet subtle, perturbations in gene expression.

Using gene co-expression network analysis (Huynh-Thu et al., 2010), we also identified

interconnected gene sub-networks that are conserved across species and tissues in the MeCP2

deficient condition (**Fig. 1k, Supplementary Tables 3 and 4**). Comparison across all species

identified several shared major gene targets and regulators despite differences in species-

specific network structure (**Supplementary Fig. 2c**). Interestingly, WNT8B, which is known to

be modulated by the Rett-implicated gene Foxg1 (Aguiar et al., 2014), was found to be the most

interconnected gene target within the MeCP2 cross-species network. The strongest gene

regulators in this network, NR4A3 and SOD2, are both involved in oxidative metabolism, which

is also known to be dysregulated in Rett Syndrome (De Felice et al., 2012)**.** Thus, this CRISPR-

enabled *Xenopus* model of Rett replicates many genetic and phenotypic features of the disease

seen in humans.

Figure 1: MeCP2 knockdown using CRISPR in *Xenopus laevis* tadpoles models Rett

syndrome. a-b, Tadpole models of Rett syndrome exhibit distinct swimming behavior in 60 mm

diameter dishes following MeCP2 knockdown ("Rett") compared to mock-injected controls

("Control"). MeCP2 RNA expression in brain and gastrointestinal (GI) tract using RNAscope, **c**,

and protein in brain and multi-ciliated olfactory cells using immunohistochemistry, **d-f**, are

significantly reduced in Rett syndrome tadpole models while maintaining a large degree of

heterogeneity spatially, **d**, and across tissues, **e-f**; scale bars, 100 μm in brain, 20 μm in

olfactory multi-ciliated cells. **g**, Volcano plot of differentially expressed genes showing that only

70 genes are significantly up- or down-regulated following MeCP2 knockdown (p < 0.05, fold

change >2, t-test with Bonferroni correction, N = 3). Gene regulatory networks for control, **h,** and

Rett, **i,** tadpoles reveal large network rearrangements with an increase in BDNF centrality. **j**,

Comparison against other Rett syndrome models and clinical samples indicates minimal

differential gene expression across the 96-gene signature developed to classify Rett syndrome.

k, Network-level comparison across species and tissues (left) and identified shared target

(middle) and regulator (right) genes.

Computational discovery of drugs that reverse the network-level impact of Rett

In Rett, MeCP2 knockout results in widespread but subtle gene expression patterns, and

most effects are not detectable using standard statistical analysis. Researchers have applied

network-based analytical methods to uncover previously-hidden connections within RNA and

protein interaction networks (Miller et al., 2019; Sanfeliu et al., 2019; Varderidou-Minasian et al.,

2020), but these approaches have not been applied to Rett syndrome therapeutics discovery.

To leverage transcriptomics analysis for drug discovery, we developed a computational

framework we termed nemoCAD (network model for causality-aware discovery)(**Fig. 2a**) that

compares transcriptomic signatures of disease versus healthy control subjects to predict

compounds from the Library of Integrated Network-Based Cellular Signatures (LINCS) database

("NIH LINCS Program," n.d.), including FDA-approved drugs, which have a high likelihood to

reverse the disease signature state back to a healthy state. Though signature-based tools have

been previously developed to enable drug discovery (e.g., CMAP (Lamb et al., 2006), NicheNet

(Browaeys et al., 2020), GPSnet (Cheng et al., 2019)), a major drawback is the impact of the

expression of any single gene or protein on the resulting prediction. To be used effectively,

these tools require well-matched disease datasets to the human cell line-based datasets

present in LINCS, which may not be feasible for many diseases.

In contrast, we pursued an entirely different approach to transcriptomics signature

analysis that is based on a Bayesian network encoding directed gene-gene and drug-gene

interactions. By leveraging data in LINCS containing transcriptomics results following single

gene knockout/knockdown and overexpression as well as addition of drug perturbagens in

human cells, we calculated the probabilities of pairwise interactions among all of the genes and

drugs in a human gene-gene and drug-gene interaction network whose structure was extracted

from Comparative Toxicogenomics Database (CTD) ("The Comparative Toxicogenomics

Database | CTD," n.d.), KEGG (Kanehisa and Goto, 2000), and TRRUST (Han et al., 2018).

Unlike signature based computational tools that take gene expression signatures at face value,

this probabilistic approach offers greater abstraction while prioritizing for the network impact of

perturbations, including those caused by disease or addition of drugs, and thereby reduces the

impact of tissue-specific or species-specific transcriptomic biases as well as the influence of the

expression of any individual gene.

To identify drugs that might reverse disease states, we generated probabilistic network

maps for each drug in the aggregate 19,800+ compound dataset. By inputting and comparing

gene microarray data from MeCP2 knockdown tadpoles versus vehicle-injected controls, we

identified sets of differentially expressed genes contained within the gene-gene interaction

network with a p-value < 0.05 and a log2 fold change between 0.7 and 3.1 in 0.3-fold-change

increments. For each fold-change, we generated a ranked list of compounds predicted to induce

reversal of transcriptome-wide changes of: 1) single gene states using a cross-correlation score,

2) single gene states using a cross-entropy score, and 3) an overall gene network state score

corresponding to the probability of involvement of the drug nodes in the identified subnetwork.

These drug lists were combined to identify drugs that were ranked by being most consistently

predicted to reverse the Rett transcriptome-wide changes and robust across the computational

settings (**Supplementary Table 5**). Drugs were manually down-selected based on available

toxicity data and to prioritize chemical diversity for initial screening. Rett tadpoles or controls

were then dosed with lead compounds for 7 days via the culture medium beginning 1-2 days

after the onset of symptoms and after collecting baseline swimming behavior data. Animal

behaviors (swimming pattern abnormalities and seizure stages) were recorded every other day

during drug exposure and washout phases, and the severity of seizure-like events was used as

a screening metric.

These studies revealed that the FDA-approved drugs, vorinostat and ivermectin,

demonstrated time-dependent reductions in seizure score that reversed after the drugs were

washed out (**Fig. 2b**; **Supplementary Fig. 3**). However, we focused on vorinostat rather than

ivermectin in all subsequent studies due to its higher predicted scoring consistency over time,

more favorable reported pharmacokinetics, and its ability, albeit limited, to cross the blood-brain

barrier. As a positive control with which to gauge relative efficacy, we also evaluated trofinetide,

the synthetic IGF-1 C-terminal tripeptide. Although only moderately effective in recent Phase III

Rett syndrome clinical trial top-line results that aligned with Phase II trial outcomes (Glaze et al.,

2017), trofinetide represents the most relevant clinical-stage benchmark treatment. Importantly,

in our Rett tadpole models, both trofinetide and vorinostat reduced seizure-like phenotypes and

also increased viability (**Fig. 2b**). At day 6 of treatment, there was a statistically significant effect

by both drugs on reducing swimming score relative to baseline before treatment onset (**Fig. 2c**).

As expected, no significant change was seen in WT and Rett vehicle-treated animals. This

head-to-head comparison has supported the possibility that our *Xenopus* population-level model

of Rett can offer predictive value for drug development.

Figure 2: Network-based computational prediction of effective drugs to treat Rett

syndrome in tadpole models. a, Network model for causality-aware discovery (nemoCAD)

combines a directed gene-gene and drug-gene interaction network extracted from CTD,

TRRUST, and KEGG databases with interaction probabilities inferred from single gene and drug

perturbations in LINCS. Transcriptome data from any disease model or patient and

corresponding control are used to identify the relevant subnetwork and disease-specific "node

weights" that account for probabilities of up-/down-regulation of a gene. A drug-gene interaction

probability matrix, inferred from LINCS, is computationally screened against the disease-specific

subnetwork to identify compounds that significantly interact with the subnetwork and are ranked

by their predicted ability to restore the disease transcriptome back to a healthy state based on

single gene and gene network signatures. Downstream analyses can be performed on the

resulting gene-gene interaction subnetwork by interrogating the underlying subnetwork structure

to find control nodes and other network metrics. Additionally, the chemical structures of the

predicted drugs can be clustered by structural similarity based on SMILES notation and

annotated protein targets and pathways from DrugBank data. **b**, Graph showing relative effects

on seizure score over 10 days of treatment in MeCP2 KD tadpoles of vehicle and 25 μM

vorinostat (Rett - Vor) versus 70 μg/mL trofinetide (Rett - Tro), a clinical-stage drug with

demonstrated efficacy (tadpoles per condition and timepoint: N = 12 Rett - Vor and Vehicle, N =

8 Rett - Tro; error bars indicate s.d.; ANOVA P = 0.028 Vehicle-treated MeCP2 KD tadpoles did

not survive past day 3 of the treatment period in one study and past day 6 in a second study. **c**,

Change in swimming score at day 6 of treatment vs. baseline (day 0) is significantly improved

by both drugs. *, P < 0.05.

To understand the protein targets of both drugs, we adapted thermal proteome profiling

(Franken et al., 2015; Huber et al., 2015; Mateus et al., 2017) that identifies molecules that

drugs bind directly to protein targets for use with a whole exothermic animal (*Xenopus* tadpole)

as well as using isolated body segments (head, viscera, tail). While we were able to detect

some of the known histone deacetylase (HDAC) targets of vorinostat, we also found that both

this drug and ivermectin bind to multiple other proteins related more broadly to acetylation

metabolism, beyond histones and HDACs (**Supplementary Table 6**). These findings suggest a

potential mechanism of action for the Rett normalizing responses we observed that involves

restoration of normal acetyl-CoA metabolism and post-translational acetylation.

As these observations initially aligned with results of prior studies of HDAC6 acetylation

of tubulin in cells of Rett patients (Gold et al., 2015; Lebrun et al., 2021) and microtubule

modulation in closely-related CDKL5 deficiency disorder (Barbiero et al., 2019), we analyzed

tadpole tissues for α-tubulin acetylation and observed surprising bidirectional shifts in

acetylation patterns depending on the tissue type. For example, α-tubulin was hypoacetylated in

neurons in sections of the midbrain (**Supplementary Fig. 4a**) and hindbrain (**Fig. 3f**) and

increased in GI tract (**Fig. 3g**) and olfactory multi-ciliated cells (**Supplementary Fig. 4b**) and

following MeCP2 knockdown, resulting in more sparse but dense tangles. Higher resolution

imaging revealed that the cilia were longer and misaligned on multiciliated cells compared to

controls (**Fig. 3a**), and both length (**Fig. 3b**) and orientation (**Fig. 3c**) were restored by

vorinostat treatment.

We then evaluated the impact of ciliary morphology and organization on ciliary function

using fluorescent microparticles (**Supplemental Fig. 5**). The disruption of ciliary beating-

induced flow on the surface of skin (**Fig. 3d**) and olfactory multiciliated cells (**Fig. 3e**) due to

MeCP2 knockdown is reversed by vorinostat treatment, suggesting that ciliary function

restoration is linked to ciliary function.

Due to the prevalence of GI symptoms in Rett patients and our observation of tubulin

acetylation disruption and vorinostat-induced recovery in the GI tract (**Fig. 3g**), we further

explored the role of MeCP2 knockdown in the gut. Staining for isolectin B4-positive (ib4+) cells

to assess GI tract inflammation and nociceptive innervation, which is increased in Rett patients

(Baikie et al., 2014; Downs et al., 2010), revealed that tadpoles exhibited significant ib4+

expression due to MeCP2 knockdown (**Fig. 3h,i)**, which vorinostat again reversed.

Figure 3: Vorinostat treatment of Rett tadpoles restores ciliary function, normalizes hypo-

and hyperacetylated tubulin across tissues, and reduces GI tract inflammation. a,

Olfactory cell ciliary abnormalities due to MeCP2 knockdown were restored by treatment with

vorinostat (magenta-yellow colormap of fluorescence intensity, cilia stained for tubulin),

significantly restoring their **b,** length and **c,** orientation (****, P < 0.0001). Functional rescue of

cilia was also observed in both epithelial, **d**, and olfactory multiciliated cells in a microbead

clearance assay (*, P < 0.05; **, P < 0.01). Despite the canonical HDACi activity of vorinostat,

both hypoacetylated tubulin in the hindbrain, **f**, as well as hyperacetylated tubulin in the GI tract,

g, was normalized. **h**, immunifluorescence images and **i**, plot showing that the % of ib4+ cells in

the GI tract of tadpoles is increased in Rett tadpoles, which is indicative of inflammation and

heightened pain response, and that this can be rescued by vorinostat treatment.

Vorinostat normalizes the Rett phenotype in a mouse model

Because vorinostat was predicted to reverse the Rett state in our computational studies,

we evaluated its efficacy in the MeCP2-deficient male (*MeCP2$^{-/Y}$*) mouse Rett model that is

commonly used in pre-clinical studies for Rett therapeutics, including trofinetide (Guy et al.,

2001). We first treated younger animals starting at day 31 post-partum (p31) as was done in

previous studies and confirmed that daily intraperitoneal (i.p.) administration of trofinetide (100

mg/kg) results in significant amelioration of multiple disease-related parameters, including

diarrhea and motor function, measured by a reduction in the established Bird score (Guy et al.,

2007) that incorporates a broad range of Rett-related CNS and non-CNS symptoms. When

vorinostat (50 mg/kg) and trofinetide were similarly administered i.p. on a daily basis over two

weeks and compared with vehicle control, we found that animals responded favorably to both

drugs (**Fig. 4a**); however, treatment with vorinostat resulted in improved neurological function

compared to trofinetide, as determined by measuring elevated plus maze (EPM) performance

(**Fig. 4b**). Both drugs also resulted in enhanced spatial novelty Y-maze performance (**Fig. 4c**)

and an improved diarrhea score (**Fig. 4d**).

As brain microglia have been implicated in Rett progression and therapeutic effects

(Cronk et al., 2015; Derecki et al., 2012; Schafer et al., 2016) based on their role in

neuroinflammation, waste removal, and synaptic pruning, we immuno-stained for microglia

using ionized calcium binding adaptor molecule (Iba-1), a microglia-specific marker, in sections

of MeCP2$^{-/y}$ mouse brains. Using Sholl analysis to assess microglia morphology (Timmerman et

al., 2018), we found that the MeCP2$^{-/y}$ microglia exhibited significantly stunted morphology with

fewer projection crossings compared to WT littermates, indicative of a heightened inflammatory

state, which vorinostat restored (**Fig. 4e,f**). This result, together with the reduction of multiple

serum inflammatory markers, including TNF-α, IL-4, IL-6, IL-16, CCL2, MIP3b, and CCl1 (**Fig.**

4g), suggests that animals with Rett experience a heightened inflammatory state, and that

treatment with vorinostat can suppress this inflammation.

Figure 4: Vorinostat rescues multiple Rett syndrome-related CNS and somatic

symptoms. a, Bird severity scores measured in MeCP2$^{-/Y}$ mice treated with vehicle (Rett),

trofinetide (100 mg/kg), or vorinostat (50 mg/kg) from day 31 to 51 of age (ANOVA; ****, P <

0.0001). **b,** elevated-plus maze and, **c**, Y-novelty maze cognitive tests of MeCP2$^{-/Y}$ mice

comparing vorinostat treatment efficacy to vehicle and trofinetide (ANOVA and Holm-Šídák test;

*, P < 0.05; **, P < 0.01). **d**, diarrhea scored using a 0-3 scale (ANOVA and Holm-Šídák test;

***, P < 0.001; ****, P < 0.0001). Representative images of Sholl analysis of microglial

arborization in mouse olfactory (**e**) and the Sholl analysis graph showing the branching profile of

microglia (**f**) in control versus Rett mice treated with or without vorinostat (ANOVA and Holm-

Šídák test; Control vs. Rett, P = 0.0004; Control vs. Vor, P < 0.0001). **g**, Levels of indicated

cytokines measured in plasma of control mice versus Rett mice with or without treatment with

trofinetide (Rett – Tro, 100 mg/kg), vorinostat (Rett – Vor, 50 mg/kg), or vehicle (Rett) for 3

weeks (mean values shown, N = 3, scale at right indicates z-score normalized to wild-type

vehicle-treated littermates). The observed baseline inflammatory state in Rett mice agreed with

published effects of MeCP2 knockdown(O'Driscoll et al., 2015).

Oral vorinostat exhibits broad efficacy in Rett mice even after onset of symptoms

Pre-clinical studies for Rett therapies routinely dose MeCP2$^{-/y}$ mice prior to the onset of

symptoms (Castro et al., 2014; Szczesna et al., 2014) due to the rapid onset of symptoms and

short lifespan of animals, in contrast to patients who seek clinical care due to developmental

deficits which may not be diagnosed for several years before treatment could be initiated

(Tarquinio et al., 2015). To better mimic this more relevant clinical scenario, male MeCP2$^{-/y}$

mice were dosed approximately one week after the onset of symptoms. Also, because long term

treatment of Rett patients would be best served by oral administration, and due to the fact that

our results showed a reduction in diarrhea in mice treated with vorinostat via daily i.p., we tested

the efficacy of an oral formulation of vorinostat. Interestingly, when we administered trofinetide

i.p. after symptoms developed in this model, it was found to be ineffective based on overall Rett

score, similar to the moderate efficacy observed in human clinical trials, whereas oral vorinostat

(50 mg/kg) prevented significant worsening of the symptom severity score (**Fig. 5a**), ameliorated

weight gain, and increased performance in EPM (**Fig. 5b**) and Y mazes (**Fig. 5c**). Animals

treated with oral vorinostat showed complete survival after 3 weeks of treatment, which is

comparable to what would be expected for vehicle-treated Rett animals for a study of this

duration, whereas only ~ 60% of trofinetide-treated Rett animals survived (**Fig. 5d**). To our

knowledge, this represents the first effective treatment of MeCP2$^{-/y}$ mice when treatment is

initiated *after* the onset of symptoms.

Rett patients and mouse Rett models exhibit significant declines in fine motor and gait

control as well as GI tract dysfunction(Hagberg, 2002; Motil et al., 2012). Importantly, the

deficiencies in mobility, gait, and hindlimb clasping, as well as increased diarrhea we also

observed in MeCP2$^{-/y}$ animals were again rescued by vorinostat treatment (**Fig. 5e,f**). Breathing

also was significantly improved following vorinostat treatment compared to trofinetide and

vehicle treatments (**Fig. 5g**). Given the putative effects of vorinostat on acetylation metabolism

and post-translational modification of tubulin and its disruption of ciliary function identified in

tadpoles, we also analyzed α-tubulin acetylation in the multi-ciliated cells in mouse lung

bronchioles, which revealed an overall increase of hyperacetylated α-tubulin in MeCP2$^{-/y}$

animals that vorinostat again restored (**Fig. 5h**). In addition, we explored the potential link of

tubulin acetylation with muscle function and found that vorinostat treatment can rescue the

disrupted α-tubulin acetylation observed in femoral muscle sections (**Fig. 5i**). Increased

acetylation due to the loss of MeCP2 may play a role in muscle function as hyperacetylation has

been shown to increase stiffness and resistance in striatal muscles *in vitro* (Coleman et al.,

2021).

We continued to evaluate the possible impact of tubulin acetylation by examining the

colon, a biomechanically-active organ that is often distended in Rett patients and may play a

major role in digestive tract disruption that is a major complaint of these patients clinically (Motil

et al., 2012). Initial histological analysis of the colon indicated a greater degree of vacuolization

and heightened neutrophil infiltrate in MeCP2-null animals, which vorinostat and trofinetide

reversed (**Fig. 5j**). Staining of these sections for βIII- and acetylated α-tubulin also revealed that

acetylated α-tubulin is significantly more colocalized with neuronal staining in the colon in

MeCP2-null mice, and that this can be normalized by treatment with either vorinostat or

trofinetide (**Fig. 5j**). However, while vorinostat also normalized the ratio of acetylated α-tubulin

to βIII-tubulin, it was unexpectedly worsened by trofinetide (**Fig. 5j,k**). While the developing GI

tract of tadpoles prevented detailed analysis of their enteric neurons, the tubulin acetylation-

normalizing effect of vorinostat identified by network analysis in *Xenopus* clearly translated to

this mammalian (mouse) model at the cellular level.

These data show that oral dosing of vorinostat improved classical disease outcome

metrics in MeCP2-null male mice and it ameliorated microglial dysfunction and hyperacetylation

of tubulin in their GI and respiratory tracts as well as in skeletal muscle. Vorinostat also was

recently shown to be effective in treating Fragile X syndrome (Ding et al., 2021), which is closely

related to Rett, sharing both X-linked mutations and symptoms of autism, although it is caused

by a mutation in the FMR1 gene. Interestingly, we identified a loss of gene network connectivity

with FMR1 in the Rett-related *Xenopus* gene network (**Fig. 1h**), and genes with the maximum

variation and fold change differences in humans with Fragile X also occurred in pathway classes

involved in oxidative stress and multiple biosynthesis pathways that align with the major

regulators we identified by the cross-species MeCP2 mutant network analysis (**Supplementary**

Figure 6a and **Supplementary Tables 7 and 8**). Moreover, when we used nemoCAD to predict

drugs that might reverse the Fragile X genotype in humans, both the compound classes and

gene targets were similar to those predicted for *Xenopus* with MeCP2 knockdown, including a

high number of PPAR receptor agonists, HDAC inhibitors, and cyclooxygenase inhibitors

(**Supplementary Figure 6b,c**). Thus, these computational results suggest more broadly that

developmental therapeutics approaches for neurodevelopmental disorders with similar broad

transcriptional effects may benefit from using this form of gene interaction network analysis to

identify network-level signatures characteristic of specific diseases and to predict drugs that

may reverse those changes.

Figure 5: Oral administration of vorinostat in MeCP2$^{-/y}$ mice after the onset of Rett

symptoms rescues CNS and somatic symptoms. a, Bird severity scores measured in

MeCP2$^{-/Y}$ mice treated with vehicle (Rett), trofinetide (100 mg/kg i.p.), or oral vorinostat (50

mg/kg) from day 36, after onset of symptoms, showing significant suppression of symptoms by

vorinostat treatment while trofinetide's effects were indistinguishable from the vehicle control

(ANOVA, ****, P < 0.0001). **b,** elevated-plus maze and, **c**, Y-novelty maze cognitive tests of

MeCP2$^{-/Y}$ mice comparing vorinostat treatment efficacy to vehicle and trofinetide (ANOVA and

Dunnett's multiple comparison test; *, P < 0.05; **, P < 0.01; ****, P < 0.0001). **d**, survival curve

of animals in this study. **e**, mobility scored using a 0-2 scale and **f**, diarrhea scored using a 0-3

scale (ANOVA and Holm-Šídák test; **, P < 0.01 ***, P < 0.001; ****, P < 0.0001). **g**, breathing

difficulty evaluated using a 0-1 score (ANOVA and Dunnett's multiple comparison test; ***, P <

0.001). Graphs showing levels of hyperacetylated α-tubulin in cells of the multi-ciliated cells in

the bronchiolar epithelium of the lung (**h**) and skeletal muscle (**i**) in mice treated as described in

a (data are presented as mean ± s.d.; *, P < 0.05; **, P < 0.01; ***, P < 0.001). **j**, Hematoxylin

and eosin (H&E) stained histological sections of colon from the control and drug-treated mice

(top row) and immunofluorescent staining for acetylated α-tubulin (middle) and βIII-tubulin

(bottom) in these sections. **k**, Graph showing changes in the βIII-tubulin ratio in drug-treated

versus control colon tissues (*, P < 0.05; ****, P < 0.0001).

DISCUSSION

Taken together, our results demonstrate the value of combining computational network

analysis-based drug repurposing algorithms with CRISPR technology to rapidly build novel *in*

vivo disease models using *X. laevis* that can be used to generate data and carry out screening

of drugs with activities of interest predicted computationally. The *Xenopus* tadpole model

developed here, which was generated in our laboratory within 3 weeks, exhibits a large degree

of genetic and phenotypic heterogeneity that reduces risks associated with computational and

screening outcomes by better capturing the disease state. Importantly, the *Xenopus* results

translated well to a mouse Rett model, and hence the tadpole model offers a faster, lower-cost,

and potentially much higher throughput alternative to mammalian models. In addition, our

target-agnostic computational approach enabled rapid discovery of novel mechanisms,

including the counterintuitive ability of an HDAC inhibitor to normalize both hypoacetylated

tubulin in the CNS and hyperacetylated tubulin in other organs, as well as identify drugs that

have broad efficacy across multiple organs. This approach led to the first demonstration of

efficacy in the mouse MeCP2 knockout model of Rett syndrome even when treated after the

onset of symptoms, in contrast with the lack of efficacy of trofinetide. Given the modest efficacy

of trofinetide in the clinic, conducting pre-clinical studies to restore health after the onset of

symptoms as in our *Xenopus* screen may be a more robust benchmark for efficacy in complex

CNS disorders. The drug discovery platform presented here offers Rett and other

neurodevelopmental patients the possibility of single small molecule therapeutics that address

multiple fundamental disease-modifying pathways with a broad clinical impact. Furthermore, by

identifying and rapidly evaluating existing drugs, potentially even in patients, our platform

enables the discovery of hidden therapeutic mechanisms for new drug development.

ACKNOWLEDGEMENTS

We would like to thank Dr. Barbara J. Caldarone for guidance and assistance with the

mouse neurobehavioral assays, Dr. Amanda Graveline, Sarai Bardales, Andyna Vernet and

Melissa Sanchez-Ventura for their assistance with mouse studies, Thomas C. Ferrante for

guidance and assistance with image acquisition and analysis, Dr. Lay-Hong Ang and Suzanne

White for tissue immunostaining support, Dr. Aurelien Begue and Mahmoud El-Rifai for their

assistance with RNAscope experiments, and Erin Switzer and Emma Lederer for help with

Xenopus embryo fertilization and husbandry. All behavioral experiments were conducted at the

HMS Mouse Behavioral Core facility. M.L. gratefully acknowledges support of the Elisabeth

Giauque Trust, London. We are grateful for funding from the Wyss Institute for Biologically

Inspired Engineering at Harvard University through Validation Project support.

COMPETING INTERESTS

RN, FV, EG, ML, and DEI hold equity in Unravel Biosciences, Inc.; RN, FV, and DEI are

members of its board of directors; ML and DEI are members of its scientific advisory board; and

RN, FV, and EG are current employees of the company.

AUTHOR CONTRIBUTIONS

RN, FV, and DEI conceived of and directed the work. RN, TL, MS, FV, EG, KS, KV, and

AV conducted animal studies and in vitro analyses. FV, MS, and VK developed and performed

CRISPR generation of Rett tadpole models with guidance by ML. SK, MS, SL, VC, and TT

performed bioinformatic analysis and computational drug prediction. SL and RN designed and

developed nemoCAD software. JRT guided GI tract and lung immunostaining studies and

evaluated the histology and tissue staining. RN and DEI wrote the manuscript. All authors

reviewed and provided input for the manuscript.

DATA AVAILABILITY

All data used in this study, including the *Xenopus* microarray data, are publicly available

using accession codes listed in **Supplementary Table 2**.

CODE AVAILABILITY

Code is available upon reasonable request to the corresponding author.

REFERENCES

Abdala AP, Lioy DT, Garg SK, Knopp SJ, Paton JFR, Bissonnette JM. 2014. Effect of Sarizotan, a 5-HT1a and D2-Like Receptor Agonist, on Respiration in Three Mouse Models of Rett Syndrome. *Am J Respir Cell Mol Biol* **50**:1031–1039. doi:10.1165/rcmb.2013-0372OC

Abuhatzira L, Makedonski K, Kaufman Y, Razin A, Shemer R. 2007. MeCP2 Deficiency in the Brain Decreases BDNF Levels by REST/CoREST-Mediated Repression and Increases TRKB Production. *Epigenetics* **2**:214–222. doi:10.4161/epi.2.4.5212

Aguiar DP, Sghari S, Creuzet S. 2014. The facial neural crest controls fore- and midbrain patterning by regulating Foxg1 expression through Smad1 activity. *Development* **141**:2494–2505. doi:10.1242/dev.101790

Arsenault J, Hooper AWM, Gholizadeh S, Kong T, Pacey LK, Koxhioni E, Niibori Y, Eubanks JH, Wang L-Y, Hampson DR. 2021. Interregulation between fragile X mental retardation protein and methyl CpG binding protein 2 in the mouse posterior cerebral cortex. *Hum Mol Genet* **29**:3744–3756. doi:10.1093/hmg/ddaa226

Aslan Y, Tadjuidje E, Zorn AM, Cha S-W. 2017. High-efficiency non-mosaic CRISPR-mediated knock-in and indel mutation in F0 Xenopus. *Development* **144**:2852–2858. doi:10.1242/dev.152967

Baikie G, Ravikumara M, Downs J, Naseem N, Wong K, Percy A, Lane J, Weiss B, Ellaway C, Bathgate K, Leonard H. 2014. Gastrointestinal dysmotility in Rett syndrome. *J Pediatr Gastroenterol Nutr* **58**:237–244. doi:10.1097/MPG.0000000000000200

Barbiero I, De Rosa R, Kilstrup-Nielsen C. 2019. Microtubules: A Key to Understand and Correct Neuronal Defects in CDKL5 Deficiency Disorder? *Int J Mol Sci* **20**:4075. doi:10.3390/ijms20174075

Basu T, O'Riordan KJ, Schoenike BA, Khan NN, Wallace EP, Rodriguez G, Maganti RK, Roopra A. 2019. Histone deacetylase inhibitors restore normal hippocampal synaptic plasticity and seizure

threshold in a mouse model of Tuberous Sclerosis Complex. *Sci Rep* **9**:5266.

doi:10.1038/s41598-019-41744-7

Browaeys R, Saelens W, Saeys Y. 2020. NicheNet: modeling intercellular communication by linking

ligands to target genes. *Nat Methods* **17**:159–162. doi:10.1038/s41592-019-0667-5

Carobrez AP, Bertoglio LJ. 2005. Ethological and temporal analyses of anxiety-like behavior: The elevated

plus-maze model 20 years on. *Neurosci Biobehav Rev*, Defensive Behavior **29**:1193–1205.

doi:10.1016/j.neubiorev.2005.04.017

Castro J, Garcia RI, Kwok S, Banerjee A, Petravicz J, Woodson J, Mellios N, Tropea D, Sur M. 2014.

Functional recovery with recombinant human IGF1 treatment in a mouse model of Rett

Syndrome. *Proc Natl Acad Sci U S A* **111**:9941–9946. doi:10.1073/pnas.1311685111

Chahrour M, Zoghbi HY. 2007. The Story of Rett Syndrome: From Clinic to Neurobiology. *Neuron* **56**:422–

437. doi:10.1016/j.neuron.2007.10.001

Chang Q, Khare G, Dani V, Nelson S, Jaenisch R. 2006. The disease progression of Mecp2 mutant mice is

affected by the level of BDNF expression. *Neuron* **49**:341–348.

doi:10.1016/j.neuron.2005.12.027

Cheng F, Lu W, Liu C, Fang J, Hou Y, Handy DE, Wang R, Zhao Y, Yang Y, Huang J, Hill DE, Vidal M, Eng C,

Loscalzo J. 2019. A genome-wide positioning systems network algorithm for in silico drug

repurposing. *Nat Commun* **10**:1–14. doi:10.1038/s41467-019-10744-6

Chien Y-H, Srinivasan S, Keller R, Kintner C. 2018. Mechanical Strain Determines Cilia Length, Motility,

and Planar Position in the Left-Right Organizer. *Dev Cell* **45**:316-330.e4.

doi:10.1016/j.devcel.2018.04.007

Coleman AK, Joca HC, Shi G, Lederer WJ, Ward CW. 2021. Tubulin acetylation increases cytoskeletal

stiffness to regulate mechanotransduction in striated muscle. *J Gen Physiol* **153**:e202012743.

doi:10.1085/jgp.202012743

Collins AL, Levenson JM, Vilaythong AP, Richman R, Armstrong DL, Noebels JL, David Sweatt J, Zoghbi HY.

2004. Mild overexpression of MeCP2 causes a progressive neurological disorder in mice. *Hum*

Mol Genet **13**:2679–2689. doi:10.1093/hmg/ddh282

Colvin L, Fyfe S, Leonard S, Schiavello T, Ellaway C, Klerk N de, Christodoulou J, Msall M, Leonard H.

2003. Describing the phenotype in Rett syndrome using a population database. *Arch Dis Child*

88:38–43. doi:10.1136/adc.88.1.38

Cronk JC, Derecki NC, Ji E, Xu Y, Lampano AE, Smirnov I, Baker W, Norris GT, Marin I, Coddington N, Wolf

Y, Turner SD, Aderem A, Klibanov AL, Harris TH, Jung S, Litvak V, Kipnis J. 2015. Methyl-CpG

Binding Protein 2 Regulates Microglia and Macrophage Gene Expression in Response to

Inflammatory Stimuli. *Immunity* **42**:679–691. doi:10.1016/j.immuni.2015.03.013

De Felice C, Signorini C, Leoncini S, Pecorelli A, Durand T, Valacchi G, Ciccoli L, Hayek J. 2012. The role of

oxidative stress in Rett syndrome: an overview. *Ann N Y Acad Sci* **1259**:121–135.

doi:10.1111/j.1749-6632.2012.06611.x

Dellu F, Contarino A, Simon H, Koob GF, Gold LH. 2000. Genetic Differences in Response to Novelty and

Spatial Memory Using a Two-Trial Recognition Task in Mice. *Neurobiol Learn Mem* **73**:31–48.

doi:10.1006/nlme.1999.3919

Derecki NC, Cronk JC, Lu Z, Xu E, Abbott SBG, Guyenet PG, Kipnis J. 2012. Wild-type microglia arrest

pathology in a mouse model of Rett syndrome. *Nature* **484**:105–109. doi:10.1038/nature10907

Ding Q, Wu X, Li X, Wang H. 2021. Vorinostat Corrects Cognitive and Non-Cognitive Symptoms in a

Mouse Model of Fragile X Syndrome. *Int J Neuropsychopharmacol* pyab081.

doi:10.1093/ijnp/pyab081

Downs J, Géranton SM, Bebbington A, Jacoby P, Bahi-Buisson N, Ravine D, Leonard H. 2010. Linking

MECP2 and pain sensitivity: The example of Rett syndrome. *Am J Med Genet A* **152A**:1197–1205.

doi:10.1002/ajmg.a.33314

Exner CRT, Willsey HR. 2021. Xenopus leads the way: Frogs as a pioneering model to understand the

human brain. *genesis* **59**:e23405. doi:10.1002/dvg.23405

Forbes M. 2021. py-factorgraph.

Franken H, Mathieson T, Childs D, Sweetman GMA, Werner T, Tögel I, Doce C, Gade S, Bantscheff M,

Drewes G, Reinhard FBM, Huber W, Savitski MM. 2015. Thermal proteome profiling for

unbiased identification of direct and indirect drug targets using multiplexed quantitative mass

spectrometry. *Nat Protoc* **10**:1567–1593. doi:10.1038/nprot.2015.101

Gabel HW, Kinde B, Stroud H, Gilbert CS, Harmin DA, Kastan NR, Hemberg M, Ebert DH, Greenberg ME.

2015. Disruption of DNA-methylation-dependent long gene repression in Rett syndrome. *Nature*

522:89–93. doi:10.1038/nature14319

Gaetani M, Sabatier P, Saei AA, Beusch CM, Yang Z, Lundström SL, Zubarev RA. 2019. Proteome Integral

Solubility Alteration: A High-Throughput Proteomics Assay for Target Deconvolution. *J Proteome*

Res **18**:4027–4037. doi:10.1021/acs.jproteome.9b00500

Giacometti E, Luikenhuis S, Beard C, Jaenisch R. 2007. Partial rescue of MeCP2 deficiency by postnatal

activation of MeCP2. *Proc Natl Acad Sci* **104**:1931–1936. doi:10.1073/pnas.0610593104

Glaze DG, Neul JL, Percy A, Feyma T, Beisang A, Yaroshinsky A, Stoms G, Zuchero D, Horrigan J, Glass L,

Jones NE. 2017. A Double-Blind, Randomized, Placebo-Controlled Clinical Study of Trofinetide in

the Treatment of Rett Syndrome. *Pediatr Neurol* **76**:37–46.

doi:10.1016/j.pediatrneurol.2017.07.002

Gold WA, Lacina TA, Cantrill LC, Christodoulou J. 2015. MeCP2 deficiency is associated with reduced

levels of tubulin acetylation and can be restored using HDAC6 inhibitors. *J Mol Med Berl Ger*

93:63–72. doi:10.1007/s00109-014-1202-x

Guy J, Gan J, Selfridge J, Cobb S, Bird A. 2007. Reversal of Neurological Defects in a Mouse Model of Rett

Syndrome. *Science* **315**:1143–1147. doi:10.1126/science.1138389

Guy J, Hendrich B, Holmes M, Martin JE, Bird A. 2001. A mouse *Mecp2*-null mutation causes neurological

symptoms that mimic Rett syndrome. *Nat Genet* **27**:322–326. doi:10.1038/85899

Hagberg B. 2002. Clinical manifestations and stages of rett syndrome. *Ment Retard Dev Disabil Res Rev*

8:61–65. doi:10.1002/mrdd.10020

Han H, Cho J-W, Lee Sangyoung, Yun A, Kim H, Bae D, Yang S, Kim CY, Lee M, Kim E, Lee Sungho, Kang B,

Jeong D, Kim Y, Jeon H-N, Jung H, Nam S, Chung M, Kim J-H, Lee I. 2018. TRRUST v2: an

expanded reference database of human and mouse transcriptional regulatory interactions.

Nucleic Acids Res **46**:D380–D386. doi:10.1093/nar/gkx1013

Hewapathirane DS, Dunfield D, Yen W, Chen S, Haas K. 2008. In vivo imaging of seizure activity in a novel

developmental seizure model. *Exp Neurol* **211**:480–488. doi:10.1016/j.expneurol.2008.02.012

Hockly E, Richon VM, Woodman B, Smith DL, Zhou X, Rosa E, Sathasivam K, Ghazi-Noori S, Mahal A,

Lowden PAS, Steffan JS, Marsh JL, Thompson LM, Lewis CM, Marks PA, Bates GP. 2003.

Suberoylanilide hydroxamic acid, a histone deacetylase inhibitor, ameliorates motor deficits in a

mouse model of Huntington's disease. *Proc Natl Acad Sci U S A* **100**:2041–2046.

doi:10.1073/pnas.0437870100

Huber KVM, Olek KM, Müller AC, Tan CSH, Bennett KL, Colinge J, Superti-Furga G. 2015. Proteome-wide

drug and metabolite interaction mapping by thermal-stability profiling. *Nat Methods* **12**:1055–

1057. doi:10.1038/nmeth.3590

Huynh-Thu VA, Irrthum A, Wehenkel L, Geurts P. 2010. Inferring Regulatory Networks from Expression

Data Using Tree-Based Methods. *PLOS ONE* **5**:e12776. doi:10.1371/journal.pone.0012776

Kanehisa M, Goto S. 2000. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res* **28**:27–

30.

Kerr AM, Nomura Y, Armstrong D, Anvret M, Belichenko PV, Budden S, Cass H, Christodoulou J, Clarke A,

Ellaway C, d'Esposito M, Francke U, Hulten M, Julu P, Leonard H, Naidu S, Schanen C, Webb T,

Engerstrom IW, Yamashita Y, Segawa M. 2001. Guidelines for reporting clinical features in cases

with MECP2 mutations. *Brain Dev* **23**:208–211. doi:10.1016/S0387-7604(01)00193-0

Kulkarni SS, Griffin JN, Date PP, Liem KF, Khokha MK. 2018. WDR5 Stabilizes Actin Architecture to

Promote Multiciliated Cell Formation. *Dev Cell* **46**:595-610.e3. doi:10.1016/j.devcel.2018.08.009

Labun K, Montague TG, Gagnon JA, Thyme SB, Valen E. 2016. CHOPCHOP v2: a web tool for the next

generation of CRISPR genome engineering. *Nucleic Acids Res* **44**:W272-276.

doi:10.1093/nar/gkw398

Lamb J, Crawford ED, Peck D, Modell JW, Blat IC, Wrobel MJ, Lerner J, Brunet J-P, Subramanian A, Ross

KN, Reich M, Hieronymus H, Wei G, Armstrong SA, Haggarty SJ, Clemons PA, Wei R, Carr SA,

Lander ES, Golub TR. 2006. The Connectivity Map: using gene-expression signatures to connect

small molecules, genes, and disease. *Science* **313**:1929–1935. doi:10.1126/science.1132939

Lebrun N, Delépine C, Selloum M, Meziane H, Nectoux J, Herault Y, Bienvenu T. 2021. HDAC inhibitor

ameliorates behavioral deficits in Mecp2308/y mouse model of Rett syndrome. *Brain Res*

147670. doi:10.1016/j.brainres.2021.147670

Li W, Pozzo-Miller L. 2014. BDNF deregulation in Rett syndrome. *Neuropharmacology*, BDNF Regulation

of Synaptic Structure, Function, and Plasticity **76**:737–746.

doi:10.1016/j.neuropharm.2013.03.024

Lonowski LA, Narimatsu Y, Riaz A, Delay CE, Yang Z, Niola F, Duda K, Ober EA, Clausen H, Wandall HH,

Hansen SH, Bennett EP, Frödin M. 2017. Genome editing using FACS enrichment of nuclease-

expressing cells and indel detection by amplicon analysis. *Nat Protoc* **12**:581–603.

doi:10.1038/nprot.2016.165

Luikenhuis S, Giacometti E, Beard CF, Jaenisch R. 2004. Expression of MeCP2 in postmitotic neurons

rescues Rett syndrome in mice. *Proc Natl Acad Sci U S A* **101**:6033–6038.

doi:10.1073/pnas.0401626101

Martínez de Paz A, Sanchez-Mut JV, Samitier-Martí M, Petazzi P, Sáez M, Szczesna K, Huertas D, Esteller

M, Ausió J. 2015. Circadian Cycle-Dependent MeCP2 and Brain Chromatin Changes. *PLoS ONE*

10. doi:10.1371/journal.pone.0123693

Mateus A, Määttä TA, Savitski MM. 2017. Thermal proteome profiling: unbiased assessment of protein

state through heat-induced stability changes. *Proteome Sci* **15**:13. doi:10.1186/s12953-017-

0122-4

Miller RA, Ehrhart F, Eijssen LMT, Slenter DN, Curfs LMG, Evelo CT, Willighagen EL, Kutmon M. 2019.

Beyond Pathway Analysis: Identification of Active Subnetworks in Rett Syndrome. *Front Genet*

10:59. doi:10.3389/fgene.2019.00059

Morton J, Snider TA. 2017. Guidelines for collection and processing of lungs from aged mice for

histological studies. *Pathobiol Aging Age-Relat Dis* **7**:1313676.

doi:10.1080/20010001.2017.1313676

Motil KJ, Caeg E, Barrish JO, Geerts S, Lane JB, Percy AK, Annese F, McNair L, Skinner SA, Lee H-S, Neul JL,

Glaze DG. 2012. Gastrointestinal and nutritional problems occur frequently throughout life in

girls and women with Rett syndrome. *J Pediatr Gastroenterol Nutr* **55**:292–298.

doi:10.1097/MPG.0b013e31824b6159

Nalle SC, Zuo L, Ong MLDM, Singh G, Worthylake AM, Choi W, Manresa MC, Southworth AP, Edelblum

KL, Baker GJ, Joseph NE, Savage PA, Turner JR. 2019. Graft-versus-host disease propagation

depends on increased intestinal epithelial tight junction permeability. *J Clin Invest* **129**:902–914.

doi:10.1172/JCI98554

Nenni MJ, Fisher ME, James-Zorn C, Pells TJ, Ponferrada V, Chu S, Fortriede JD, Burns KA, Wang Y, Lotay

VS, Wang DZ, Segerdell E, Chaturvedi P, Karimi K, Vize PD, Zorn AM. 2019. Xenbase: Facilitating

the Use of Xenopus to Model Human Disease. *Front Physiol* **10**. doi:10.3389/fphys.2019.00154

NIH LINCS Program. n.d. https://lincsproject.org/

O'Driscoll CM, Lima MP, Kaufmann WE, Bressler JP. 2015. Methyl CpG binding protein 2 deficiency

enhances expression of inflammatory cytokines by sustaining NF-κB signaling in myeloid derived

cells. *J Neuroimmunol* **283**:23–29. doi:10.1016/j.jneuroim.2015.04.005

Pearl J. 1982. Reverend bayes on inference engines: a distributed hierarchical approachProceedings of

the Second AAAI Conference on Artificial Intelligence, AAAI'82. Pittsburgh, Pennsylvania: AAAI

Press. pp. 133–136.

Potts EM, Coppotelli G, Ross JM. 2020. Histological-Based Stainings using Free-Floating Tissue Sections. *J*

Vis Exp JoVE. doi:10.3791/61622

Pratt KG, Khakhalin AS. 2013. Modeling human neurodevelopmental disorders in the Xenopus tadpole:

from mechanisms to therapeutic targets. *Dis Model Mech* **6**:1057–1065.

doi:10.1242/dmm.012138

Przanowski P, Wasko U, Zheng Z, Yu J, Sherman R, Zhu LJ, McConnell MJ, Tushir-Singh J, Green MR,

Bhatnagar S. 2018. Pharmacological reactivation of inactive X-linked Mecp2 in cerebral cortical

neurons of living mice. *Proc Natl Acad Sci* **115**:7991–7996. doi:10.1073/pnas.1803792115

Renthal W, Boxer LD, Hrvatin S, Li E, Silberfeld A, Nagy MA, Griffith EC, Vierbuchen T, Greenberg ME.

2018. Characterization of human mosaic Rett syndrome brain tissue by single-nucleus RNA

sequencing. *Nat Neurosci* **21**:1670–1679. doi:10.1038/s41593-018-0270-6

Russo PST, Ferreira GR, Cardozo LE, Bürger MC, Arias-Carrasco R, Maruyama SR, Hirata TDC, Lima DS,

Passos FM, Fukutani KF, Lever M, Silva JS, Maracaja-Coutinho V, Nakaya HI. 2018. CEMiTool: a

Bioconductor package for performing comprehensive modular co-expression analyses. *BMC*

Bioinformatics **19**:56. doi:10.1186/s12859-018-2053-1

Ruthsatz K, Dausmann KH, Peck MA, Drees C, Sabatino NM, Becker LI, Reese J, Hartmann L, Glos J. 2018.

Thyroid hormone levels and temperature during development alter thermal tolerance and

energetics of Xenopus laevis larvae. *Conserv Physiol* **6**. doi:10.1093/conphys/coy059

Sanfeliu A, Hokamp K, Gill M, Tropea D. 2019. Transcriptomic Analysis of Mecp2 Mutant Mice Reveals

Differentially Expressed Genes and Altered Mechanisms in Both Blood and Brain. *Front*

Psychiatry **10**:278. doi:10.3389/fpsyt.2019.00278

Schafer DP, Heller CT, Gunner G, Heller M, Gordon C, Hammond T, Wolf Y, Jung S, Stevens B. 2016.

Microglia contribute to circuit defects in Mecp2 null mice independent of microglia-specific loss

of Mecp2 expression. *eLife* **5**:e15224. doi:10.7554/eLife.15224

Sekhon RS, Briskine R, Hirsch CN, Myers CL, Springer NM, Buell CR, de Leon N, Kaeppler SM. 2013. Maize

gene atlas developed by RNA sequencing and comparative evaluation of transcriptomes based

on RNA sequencing and microarrays. *PloS One* **8**:e61005. doi:10.1371/journal.pone.0061005

Subramanian A, Narayan R, Corsello SM, Peck DD, Natoli TE, Lu X, Gould J, Davis JF, Tubelli AA, Asiedu JK,

Lahr DL, Hirschman JE, Liu Z, Donahue M, Julian B, Khan M, Wadden D, Smith IC, Lam D, Liberzon

A, Toder C, Bagul M, Orzechowski M, Enache OM, Piccioni F, Johnson SA, Lyons NJ, Berger AH,

Shamji AF, Brooks AN, Vrcic A, Flynn C, Rosains J, Takeda DY, Hu R, Davison D, Lamb J, Ardlie K,

Hogstrom L, Greenside P, Gray NS, Clemons PA, Silver S, Wu Xiaoyun, Zhao W-N, Read-Button

W, Wu Xiaohua, Haggarty SJ, Ronco LV, Boehm JS, Schreiber SL, Doench JG, Bittker JA, Root DE,

Wong B, Golub TR. 2017. A Next Generation Connectivity Map: L1000 Platform and the First

1,000,000 Profiles. *Cell* **171**:1437-1452.e17. doi:10.1016/j.cell.2017.10.049

Szczesna K, de la Caridad O, Petazzi P, Soler M, Roa L, Saez MA, Fourcade S, Pujol A, Artuch-Iriberri R,

Molero-Luis M, Vidal A, Huertas D, Esteller M. 2014. Improvement of the Rett Syndrome

Phenotype in a Mecp2 Mouse Model Upon Treatment with Levodopa and a Dopa-Decarboxylase

Inhibitor. *Neuropsychopharmacology* **39**:2846–2856. doi:10.1038/npp.2014.136

Tarquinio DC, Hou W, Neul JL, Lane JB, Barnes KV, O'Leary HM, Bruck NM, Kaufmann WE, Motil KJ, Glaze

DG, Skinner SA, Annese F, Baggett L, Barrish JO, Geerts SP, Percy AK. 2015. Age of diagnosis in

Rett syndrome: patterns of recognition among diagnosticians and risk factors for late diagnosis.

Pediatr Neurol **52**:585-591.e2. doi:10.1016/j.pediatrneurol.2015.02.007

The Comparative Toxicogenomics Database | CTD. n.d. http://ctdbase.org/

Timmerman R, Burm SM, Bajramovic JJ. 2018. An Overview of in vitro Methods to Study Microglia. *Front*

Cell Neurosci **12**. doi:10.3389/fncel.2018.00242

Tsai P-Y, Zhang B, He W-Q, Zha J-M, Odenwald MA, Singh G, Tamura A, Shen L, Sailer A, Yeruva S, Kuo W-

T, Fu Y-X, Tsukita S, Turner JR. 2017. IL-22 Upregulates Epithelial Claudin-2 to Drive Diarrhea and

Enteric Pathogen Clearance. *Cell Host Microbe* **21**:671-681.e4. doi:10.1016/j.chom.2017.05.009

Varderidou-Minasian S, Hinz L, Hagemans D, Posthuma D, Altelaar M, Heine VM. 2020. Quantitative

proteomic analysis of Rett iPSC-derived neuronal progenitors. *Mol Autism* **11**:38.

doi:10.1186/s13229-020-00344-3

Vashi N, Justice MJ. 2019. Treating Rett syndrome: from mouse models to human therapies. *Mamm*

Genome. doi:10.1007/s00335-019-09793-5

Willsey HR. 2021. Whole-Mount RNA In Situ Hybridization and Immunofluorescence of Xenopus

Embryos and Tadpoles. *Cold Spring Harb Protoc* **2021**:pdb.prot105635.

doi:10.1101/pdb.prot105635

Yang Z, Steentoft C, Hauge C, Hansen L, Thomsen AL, Niola F, Vester-Christensen MB, Frödin M, Clausen

H, Wandall HH, Bennett EP. 2015. Fast and sensitive detection of indels induced by precise gene

targeting. *Nucleic Acids Res* **43**:e59. doi:10.1093/nar/gkv126

1191