1   Social information-mediated population dynamics in non-grouping prey

3   Zoltán Tóth[1,2*] & Gabriella Csöppü[1]

5   [1]Department of Zoology, Plant Protection Institute, Centre for Agricultural Research, ELKH,

6   Budapest, Hungary

7   [2]University of Debrecen, Debrecen, Hungary

9   [*]Corresponding author: Zoltán Tóth, toth.zoltan@atk.hu; ORCID: 0000-0002-2634-8393

Abstract

Inadvertent social information (ISI) use, i.e., the exploitation of social cues including the presence and behaviour of others, has been predicted to mediate population-level processes even in the absence of cohesive grouping. However, we know little about how such effects may arise when the prey population lacks social structure beyond the spatiotemporal autocorrelation originating from the random movement of individuals. In this study, we built an individual-based model where predator avoidance behaviour could spread among randomly moving prey through the network of nearby observers. We qualitatively assessed how ISI use may affect prey population size when cue detection was associated with different probabilities and fitness costs, and characterised the structural properties of the emerging detection networks that would provide pathways for information spread in prey. We found that ISI use was among the most influential model parameters affecting prey abundance and increased equilibrium population sizes in most examined scenarios. Moreover, it could substantially contribute to population survival under high predation pressure, but this effect strongly depended on the level of predator detection ability. When prey exploited social cues in the presence of high predation risk, the observed detection networks consisted of a large number of connected components with small sizes and small ego networks; this resulted in efficient information spread among connected individuals in the detection networks. Our study provides hypothetical mechanisms about how temporary local densities may allow information diffusion about predation threats among conspecifics and facilitate population stability and persistence in non-grouping animals.

Keywords: social information use, detection networks, predator-prey relationship, non-grouping animals, equilibrium population size, individual-based model

Significance Statement

The exploitation of inadvertently produced social cues may not only modify individual behaviour but also fundamentally influence population dynamics and species interactions. Using an individual-based model, we investigated how the detection and spread of adaptive antipredator behaviour may cascade to changes in the demographic performance of randomly moving (i.e., non-grouping) prey. We found that social information use contributed to population stability and persistence by reducing predation-related per capita mortality and raising equilibrium population sizes when predator detection ability reached a sufficient level. We also showed that temporary detection networks had structural properties that allowed efficient information spread among prey under high predation pressure. Our work represents a general modelling approach that could be adapted to specific predator-prey systems and scrutinize how temporary local densities allow dynamic information diffusion about predation threats and facilitate population stability in non-grouping animals.

Introduction

Organisms have to gather information about their surroundings to overcome challenges such as finding resources and avoiding danger (Dall and Johnstone 2002). For that, individuals directly interact with the environment to gain up-to-date information about its state ('personal information'), but they can also complement that knowledge by utilizing social information for optimal decision-making (Galef and Giraldeau 2001; Dall et al. 2005; Bonnie and Earley 2007; Hoppitt and Laland 2013). One type of social information is associated with inadvertently produced social cues that include the presence or the behaviour of others, or the product of their behaviour such as scent marks, excretions or food remnants, all of which may provide relevant information about current environmental conditions. Inadvertent social information (ISI) use is known to occur in many ecological contexts, including predator avoidance, foraging and habitat choice (Danchin et al. 2004; Gil et al. 2018). The advantages of living in social groups are thought to include the opportunity to access social information (Krause & Ruxton 2002; Ward & Webster 2016; Goodale et al. 2017), and thus ISI use is usually associated with species where social interactions promote information transmission among group-mates (King and Cowlishaw 2007; Duboscq et al. 2016; Gil et al. 2017).

Under predation risk, dynamic information about threats is transmitted from alarmed group members to naïve ones, a phenomenon that is commonly called collective detection (Lima 1990; Pays et al. 2013). This process often takes place through evolved signals such as alarm calls, but social cues including sudden movements (Coleman 2008; Hingee and Magrath 2009; Boujja-Miljour et al. 2017), fright responses (Chivers and Ferrari 2014; Cruz et al. 2020), or changes in posture (Brown et al. 1999; Pays et al. 2013) have also been found to convey information about the presence of predators in animal collectives. Adjustments to the behaviour of others (also referred to as 'behavioural contagion'; Firth 2020) do not only affect individual fitness by increasing survival probabilities, but can also lead to the

emergence of correlated behaviours and space use in many individuals and thus influence

system-level functions (Goodale et al. 2010; Gil et al. 2018; Tóth 2021). Previous theoretical

models have predicted that ISI use can prevent population collapses under high predation

pressure (Gil et al. 2017, 2018) and facilitate the coexistence of competing species that share

common predators (Parejo and Avilés 2016; Gil et al. 2019). Empirical evidence also

indicates that the utilization of social information can influence the material flux on the

ecosystem level (Gil and Hein 2017). By promoting adaptive behavioural responses to

environmental uncertainties (e.g., due to anthropogenic effects [Greggor et al. 2017], in the

distribution of resources [O'Mara et al. 2014] or predation risk [Crane et al. 2021]), ISI use

has the potential to minimize the impact of morphological, physiological or genetic

adaptations (Laland 1992) or influence genetic change through gene–culture coevolution

(Whitehead et al. 2019).

There are animal species that do not exhibit social attraction toward conspecifics and

therefore do not form permanent or periodical cohesive groups. We refer to these organisms

as non-grouping animals (for more details about to this definition, see Tóth et al. 2020).

Lacking motivation for social cohesion, non-grouping animals do not maintain spatial

proximity with others, and thus direct interactions between conspecifics can be infrequent.

Nevertheless, such individuals may also exploit social cues (e.g., visual, acoustic, chemical or

vibrational cues) when these are within the range of relevant sensory perception. Moreover,

social information may also diffuse among nearby observers via 'detection networks'

(reviewed in Tóth et al. 2020). If so, spatial changes in social cues over time (e.g., relative

differences in activity and associated conspicuousness; Chivers and Ferrari 2014) can provide

dynamic information about predation threats in many terrestrial and aquatic systems (Gil et al.

2017). In accordance with this idea, wood crickets (*Nemobius sylvestris*) adaptively change

their behaviour after having observed the predator avoidance behaviour of knowledgeable

conspecifics, and this information is transmitted to and utilized by other naïve individuals as

well (Coolen et al. 2005). In temporary aggregations, escape responses of Iberian green frogs

(*Rana perezi*) are also influenced by the behaviour of adjacent conspecifics (Martín et al.

2006). In mixed-species aggregations of non-schooling fish, the density (number of fish in the

foraging area) and behaviour (when to feed in and when to flee from the foraging area) of

nearby individuals are being used as inadvertent social information (Gil and Hein 2017). The

resulting behavioural coupling among individuals, in turn, affects both species abundance and

the amount of algae consumed and as a result, determines the total material flow in the coral

reef ecosystem. While such observations prove that threat-related social cues can be exploited

by non-grouping animals in some instances, the general conditions under which ISI use exerts

a positive effect on population stability and persistence in such species have remained largely

unexplored. For example, thresholds associated with the cost of antipredator behaviour and

probabilities of cue detection (i.e., the detection of predators or conspecifics' behaviour) may

set boundaries for social information-modulated population-level effects under different

predation pressure regimes. Similarly, detection networks may have only a limited capacity to

provide efficient information pathways for the emergence of such effects. In a previous work,

Tóth (2021) used an individual-based model to test specifically how ISI use may alter the

relationship between fluctuating predator and non-grouping prey populations. The author

found that ISI use can disrupt population cycles and decrease the strength of the negative

feedback of second-order dependence between predator and prey, thus stabilize their

dynamics and facilitate their long-term coexistence.

In this study, we investigated how the detection and spread of predator avoidance

behaviour among conspecifics affected demographic performance in non-grouping prey. We

constructed an individual-based model of prey and generalist predator populations where

individuals (both prey and predators) moved randomly on the landscape, and social

information could diffuse through the observation of antipredator behaviour in prey. This

model, an extension of our earlier model presented by Tóth (2021), allowed us to assess

qualitatively how ISI use may cascade to changes in the size of noncyclic prey populations

that experience relatively constant predation pressures, and examine the structural properties

of detection networks in prey that lack social structure. Thus, predictions from the presented

model may be applied more generally compared to those from our previous work.

Materials and Methods

Model construction

We simulated a homogeneous, continuous 2D landscape (80×80 spatial units) where both

prey and predators moved randomly by exhibiting correlated random walks (CRW). CRW

considers short-term correlations between successive step orientations and has been

successfully used to model animals' non-orientated search paths for a long time (Benhamou

2006; Codling et al. 2008; Reynolds 2014). In CRW models, habitat must be rather

homogeneous; examples of such natural habitats include beaches and deserts, grasslands,

agricultural crops, or those where resource patch distribution at the large scale is uniform or

random (Byers 2001). At the start of a simulation cycle, 500 prey and 150 predators were

randomly placed on the landscape, and then individuals performed a given set of behaviours

(Fig. 1, Table 1). During movement, each individual's movement distance was randomly

selected between zero and a maximum value given by the parameters d_{prey} and d_P for prey and

predators, respectively. Turning angles were determined by random deviates drawn from

wrapped Cauchy circular distribution with $\mu=0$ and $\rho=0.8$. At the landscape edge, individuals

moved to the opposite side of the landscape when crossing a boundary and continued moving

(i.e., torus landscape with no edge). Both prey and predator could also detect other individuals

through the landscape edge. We assumed that only one individual could survive within the

range of one spatial unit due to competition in both prey and predators (after movement and

dispersion of offspring; see Fig. 1), introducing density-dependent mortality in their

populations. In this system, we assumed non-dynamic predators that can exert high predation

pressure on the prey population, thus predator population size was determined only by their

reproductive rate and density-dependent mortality, but was unaffected by the success of

hunting (as if switching to alternative prey when necessary). Consequently, predator and prey

populations were noncyclic and demographically decoupled (for a similar approach, see Gil et

al. 2019), and prey populations experienced predation pressures that were directly

proportional to the given value of predators' reproduction-related parameter (Table 1).

In the absence of predators, prey moved, competed, fed and reproduced in the

simulated landscape. Prey population size resulted in this scenario was regarded as being in

equilibrium at the carrying capacity of the environment. When present, each predator could

consume a maximum of two prey individuals in a cycle within its hunting range, which was

defined as an r_P distance from the predator's position in any direction. Prey could detect

predators that were r_{prey} distance with a probability given by P_{detect} (determined by individual

Bernoulli trials). Thus, the detection of other individuals (predators, prey or conspecifics)

depended on individual sensory ranges that also crossed through the torus landscape edges,

bringing more reality to the simulation compared to the model presented in Tóth (2021). Upon

successfully detecting a predator, prey became alarmed and hid, and thus were undetectable to

predators. However, these individuals did not feed either and consequently could have a

reduced reproduction rate. Thus, prey animals were capable of behaviourally adjusting their

exposure to predators (with the probability ranging between 0.1 and 0.9; see Table 1), but this

antipredator behaviour potentially incurred a fitness cost. Lima and Dill (1990) summarized

supporting evidence for costly behavioural responses to predation risk in multiple taxa,

including a reduction in feeding, growth or reproduction. Predators hunted on visible, feeding

prey with a 50% success (determined by individual Bernoulli trials). Similarly high success

rates against some prey species has been observed, for instance, in generalist predators such

as red foxes (*Vulpes vulpes*; Červený et al. 2012), *Drassodes lapidosus* spiders (Michálek et

al. 2017), or American kestrels (*Falco sparverius*; Toland 1987). Prey could also detect

predators indirectly by observing alarmed conspecifics within r_{prey} distance with a probability

given by P_{isi} (determined by individual Bernoulli trials). Being alarmed had the same

consequences (i.e., immune to predation, reduced reproduction rate) irrespective of the

detection mode. Wide ranges of possible values for both P_{detect} and P_{isi} (see in Table 1) were

used to cover most scenarios in which ISI use may occur under natural conditions. We did not

manipulate cue reliability in the model, we simply considered that ISI use had a higher cost

when social cues could also be false and individuals responded to those indiscriminately. Prey

feeding occurred once in a cycle in prey that was not hiding. The number of offspring for each

individual was sampled from a Poisson distribution with the shape parameter given by $\lambda_{reduced}$

for alarmed prey, λ_{max} for fed prey, and λ_P for predators in each cycle. Offspring dispersed in

the same cycle 8, 9 or 10 spatial units away (randomly chosen) from the parent in both prey

and predators. These higher step values (10 spatial units is the double of maximum d_{prey} and

d_P) were chosen to reflect that juvenile dispersion distances can far exceed adult movement

ranges.

Detection networks

From the spatial distribution of prey, we defined detection networks based on the range within

which individuals could observe the behaviour of others (i.e., exploit social cues if present) in

each simulation cycle (Fig. 2). In such networks, nodes represent individuals, and edges

denote the possibility of mutual observation. If a prey individual became alarmed because it

successfully detected a predator, information could spread from this individual to other

conspecifics in the network under the following rules. The probability of information

acquisition from one node to another is given by w^k, where w is the edge weight

(corresponding to the probability of information spread from one node to another through the

edge between them and specified by the parameter P_{isi} in the model) and k is the number of

steps on the shortest path between the two nodes. Only shortest paths were used to minimize

the "travel time" of information between nodes in the network. During simulations, the

maximum number of steps between the focal and the observed nodes was set to two and the

total number of observed neighbours to ten (i.e., $k_{max}=2$ and $\sum(n)_{max}=5$ in each k step). Thus,

an individual could receive information from a maximum of ten of its neighbours that were a

maximum of two steps away in the detection network. With such restrictions, ISI use did not

facilitate the emergence of large aggregations in prey and did not occur far outside the hunting

range of predators. If there were more than five nodes at k step to a focal node, we randomly

selected five. For any individual, the total probability of receiving information from its

neighbours was calculated using the inclusion-exclusion principle (Allenby and Slomson

2010).

Analysis of simulation outputs

All simulations and calculations were performed in R 4.0.4 (R Core Team 2021). Instead of

frequentist hypothesis testing, we focused on evaluating the magnitude of differences between

simulation runs with different parameter settings (White et al. 2014). We ran the population

simulations for 200 cycles (this interval was sufficient to reach equilibrium prey population

size in the studied scenarios; see in Fig. 3a) and used the data from the last cycle in all

calculations. R script for model construction and simulated data are available at Figshare

(https://figshare.com/s/34fc714342dab9123193).

We characterised prey population sizes by calculating the mean, standard deviation,

maximum and minimum values in four settings: in the absence of predators, with minimal

P_{detect}, with nominal P_{detect}, and with nominal P_{detect} and P_{isi} parameter values, respectively. All

other parameters were set to their initial values; for each model type, simulations were iterated

50 times. When the predator detection probability was set to its minimal value, the prey

population died out in a single iteration; prey extinction was not observed in other settings.

We examined how predator abundance affected mortality rate due to predation in prey

in the presence of minimal P_{detect}, nominal P_{detect}, and nominal P_{detect} and P_{isi} parameter values,

respectively. All other parameters were set to their initial values. In each setting, simulation

runs were iterated 50 times. If the prey population died out before the 200th simulation cycle,

the given run was omitted from the dataset ($n=209$; only in the 'minimal predator detection'

setting).

Sensitivity analysis

We used Morris's „OAT" elementary effects screening method (Morris 1991) with the

extension introduced by Campolongo et al. (2007) as a global sensitivity analysis (SA) to rank

the model parameters according to their impact on prey population size. We chose this SA

because it produces results comparable to the more complex methods (Confalonieri et al.

2010) and is applicable to uncover the mechanisms and patterns produced by individual-based

models (Imron et al. 2012; Beaudouin et al. 2015; Ten Broeke et al. 2016). The mean of the

absolute value of the elementary effect (μ^*_i) provides a measure for the overall influence of

each input variable on the model output, whereas the standard deviation of the elementary

effect (σ_i) indicates possible non-linear effects or interactions among variables (Campolongo

et al. 2007; Iooss & Lemaître 2015). We also ranked the model parameters using a global

index (GI) (Ciric et al. 2012) calculated as:

$$GI = \sqrt{\left(\mu^*_i\right)^2 + (\sigma_i)^2}$$

For the space-filling sampling strategy proposed by Campolongo et al. (2007), we generated

$r_2 = 1000$ Morris trajectories and then retained $r_1 = 50$ with the highest 'spread' in the input

space to calculate the elementary effect for each model parameter.

Parameter space exploration

We explored a specific part of the parameter space by visualising the combined effect of the

parameters P_{detect}, P_{isi}, λ_P and $\lambda_{reduced}$ on prey population size. Specifically, we investigated the

effect of ISI use at low, intermediate and high levels of predator detection probabilities. In

each scenario, predator avoidance behaviour had either no cost or incurred moderate fitness

cost (i.e., decreased by one third compared to the maximum) and predation pressure was

either low (0.025), intermediate (0.05) or high (0.075). In each setting, we used the complete

range of parameter values for P_{isi} (Table 1). Simulations were iterated 30 times. In the low

predator detection probability scenario coupled with high predation pressure, the prey

population died out in the majority of simulation runs (n=581); these simulation outputs were

omitted from the assembled dataset.

Network characterisation

We generated network data by running the model with λ_P=0.075 (i.e., under a high level of

predation pressure), while all other parameters were set to their nominal values. We compared

emerging detection networks that were generated with the presence of ISI use (P_{isi}=0.5) to

those that were obtained when P_{isi}=0. We calculated the number of components, component

size, average ego network and average global efficiency as structural network properties for

network characterisation. Simulations were repeated 50 times in each parameter setting.

Additionally, we calculated the same characteristics for randomised detection networks as

well (Farine 2017; Hobson et al. 2021). These were constructed from both type of the

observed detection networks by randomly reshuffling the edges between nodes while also

retaining the original degree distributions. Thus, randomized networks represented a

hypothetical scenario where interactions are equally likely between any pair of nodes (Croft et

al. 2011). While the main purpose here was to explore the global structure of the observed

detection networks, the randomized networks helped us to assess whether ISI use could

similarly affect the structural properties of networks that were based on this simplifying

assumption. The number of components represents the number of connected parts in the

detection networks (isolated nodes excluded). We computed component size as the number of

components divided by the number of connected nodes; this measure denotes the average

number of nodes embedded within components. We calculated the average size of ego

networks as the mean number of reachable nodes within two steps in the components. To

estimate transmissibility within components, we used the measure 'global efficiency'

(Pasquaretta et al. 2014; Romano et al. 2018; Latora and Marchiori 2001). Global efficiency

for a graph with N vertices is:

$$E_{\mathrm{global}}(\mathrm{G}) = \frac{1}{N(N-1)}\sum_{i \neq j}\frac{1}{d_{ij}}$$

where d_{ij} is the shortest path length between nodes i and j. The value of this measure ranges

from 0 to 1, and represents how fast information may spread from the source to the most

peripheral network positions with the least number of connections (Romano et al. 2018). We

computed global efficiency for the largest components in the networks. For the calculation of

the above network properties, we used the 'igraph' and 'brainGraph' R packages (Csardi and

Nepusz 2016; Watson 2020).

Results

We found that nominal predation pressure coupled with minimal predator detection

probability (P_{detect}=0.1) led to small prey population size with high variation among runs

compared to the null model when predators were absent and prey population existed at the

carrying capacity of the environment (Fig. 3a, Table 2). Nominal predator detection

probability (P_{detect}=0.5) increased mean prey population size and stabilised the prey

population at higher abundance values, while in the presence of nominal probability of ISI use

in prey (P_{detect}=0.5 & P_{isi}=0.5), prey population size increased further by approx. 53%. The

sensitivity analysis also confirmed that P_{isi} was an influential model input in the constructed

model (Fig. 3b). As expected, the parameters driving antipredator behaviour, i.e. the level of

predation pressure, the probability of predator detection directly or via conspecifics, and the

cost associated with performing antipredator behaviour, were all important and characterised

by non-linear effects on prey abundance and/or strong interactions with other parameters. The

parameters d_{prey} and d_P had considerably less influence on the dispersion of the model output,

and were fixed to their nominal values in the subsequent analyses. The mechanism behind the

effect of P_{isi} was that the presence of ISI use could decrease the per capita mortality due to

predation across the whole range of the examined predation pressure regime and substantially

mitigate the positive relationship between predation-related mortality rate and predator

population size (Fig. 4, Fig. S1).

Consistent with expectations, P_{isi} affected prey number in all examined P_{detect}

scenarios in interaction with the effect of cost and predator pressure (Fig. 5). This relationship

was positive and nonlinear in most cases. When the predation pressure was low, P_{isi} positively

influenced prey abundance to a limited extent, while the effect of the associated cost,

especially at lower P_{isi} values, depended on the value of P_{detect}. When the predation pressure

was intermediate or high, P_{isi} exerted a more substantial influence on prey abundance and had

the capacity to double the number of prey individuals irrespective of the presence or absence

of associated cost. Importantly, ISI use could counteract high predation pressure only when

P_{detect} had a sufficient value (directly dependent on the degree of predation pressure), and did

not compensate for low predator detection ability as indicated by the high prevalence of

population extinctions in prey when high predation pressure was coupled with low predator

detection ability. The presence of associated fitness cost in the high predation pressure

settings greatly reduced the magnitude of the effect of ISI use on prey population size, but P_{isi}

could still increase prey population size even at intermediate values if $P_{detect}>0.25$.

The observed detection networks were characterized by high numbers of components

that consisted of few connected individuals and small ego networks, especially when social

information could spread among individuals (Fig. 6). Global efficiency within the largest

components, therefore, was also high in the observed detection networks, indicating efficient

information transmission among individuals whenever connected prey was able to detect

nearby predators. These attributes of functioning detection networks were not the direct

consequence of higher prey population size in the presence of ISI use, because the

corresponding randomized networks did not show the same degree of structural changes

compared to the $P_{isi}=0$ setting.

Discussion

Social information use has been assumed both to increase individual fitness and to affect

population- and community-level processes (Dall et al. 2005; Gil et al. 2018). We expected

that such effects could emerge in randomly moving non-grouping prey if behavioural

contagion can occur through detection networks, i.e., a dynamic system of temporary

observation-based connections between conspecifics. Correlated random walk has

successfully been used to describe non-oriented movement trajectory in a number of non-

grouping animals, e.g. insects (e.g., Kareiva & Shigesada 1983; McCulloch & Cain 1989;

Byers 2001), zooplankton (e.g., Komin et al. 2004; Uttieri et al. 2004, 2008), echinoderms

(e.g., Lohmann et al. 2016) or mammals (e.g., Johnson et al. 2008). We found that

irrespective of the apparent stochasticity in our model, the sharing of adaptive antipredator

behaviour could contribute to population stability and persistence in prey by mitigating

predation-related per capita mortality and raising equilibrium population sizes. We also

showed that temporary detection networks had structural properties that allowed the efficient

spread of adaptive antipredator behaviour among prey under high predation pressure. In

animal groups, information spreads via social connections among individuals and social

network positions strongly interact with individual spatial behaviour (Firth and Sheldon 2016;

Spiegel et al. 2016; Webber & Vander Wal 2018; Albery et al. 2021), thus movement

characteristics and space use are shaping information transmission by affecting social

connections. Our findings indicate that non-grouping animals, by being embedded in

detection networks based on their perception attributes and spatial locations, can benefit from

similar information transmission processes as well.

Our results corroborate with previous studies indicating that social information may

act as a stabilizing mechanism in systems where predators can exert high pressure on prey

populations (Gil et al. 2017, 2018, 2019). While in those models social information directly

reduced (following a specific function) the per capita mortality (e.g., Gil et al. 2018), the

presented work offers a more mechanistic understanding of how inadvertent social

information could propagate through a population of randomly moving individuals. Our

findings indicate that predator detection ability had to reach a sufficient level, strongly

dependent on the actual level of predation pressure, for ISI use to facilitate prey population

persistence. Notably, when this condition was met, ISI use exerted a detectable positive

influence on prey population size by relaxing predation pressure even at low probabilities and

even if the adaptive antipredator behaviour incurred a fitness cost. Although the depth of our

understanding of the detected non-linear relationships and potential thresholds is limited by

their coarse-grained variation in these parameters examined here, simulations nonetheless

prove that in a substantial part of the parameter space social information use can be expected

to raise non-grouping prey population size and facilitate its persistence. These findings may

have crucial implications in many theoretical and applied ecological contexts, ranging from

the invasive dynamics of predator-prey systems to the efficiency of biological control

practices. For instance, the recognition of novel predators by naïve prey has been associated

with social information use via different perception modalities in fish (Ferrari et al. 2005;

Manassa et al. 2013), and similar utilization of social cues in birds has been shown to

facilitate the spread of novel aposematic prey (Thorogood et al. 2017; Hämäläinen et al.

2021). Such social information-mediated interactions between prey and predators might be

more prevalent in natural ecosystems that include non-grouping species as well, contributing

to deviations from the predictions of theoretical models in the dynamics of trophic

interactions (Polis et al. 2000). When natural enemies are used as biological control agents for

pest management, social diffusion of antipredator responses may substantially reduce

predation rates rendering these practices less effective and profitable. Besides, it may also

mitigate the expected positive impact of the non-consumptive effects of predators (NCEs;

Preisser et al. 2007; Sih et al. 2010) such as decreased crop damage due to reduced feeding

rate in pests (Beleznai et al. 2017; Tholt et al. 2018). This inflation of NCEs due to

information spread can generate discrepancies in the findings of large-scale field studies and

laboratory experiments (see in Weissburg et al. 2014), and should be taken into consideration

in investigations that aim to evaluate how NCEs may trigger trophic cascades in different

ecosystems (Herman and Landis 2017; Haggerty et al. 2018; Pessarrodona et al. 2019).

Detection networks had distinct structural characteristics when prey experienced high

predation pressure and exploited social cues to avoid predators. These networks typically

consisted of many components with few connected individuals and small average ego

networks, and within these small components, social information could spread with relatively

high efficiency. The key to understanding the differences in structural properties of detection

networks in the presence and absence of ISI use lies in identifying the process that generates

more and smaller components. One plausible explanation is that prey distribution in the

simulated landscape could remain more homogeneous due to a decreased susceptibility to

predation in the vicinity of predators as the diffusion of social information greatly enhances

the probability of predator detection even among a few nearby individuals. While high

network efficiency has previously been identified in small groups, cognitive abilities and

strong social affiliations have usually been involved in explaining this emergent property

(Waters and Fewell 2012; Pasquaretta et al. 2014). Our findings indicate that incidental

connections may generate networks that have similar favourable attributes. In addition to

differences in the sizes of connected components, there may be other key differences in how

information spreads through detection or sensory networks among group-living (Strandburg-

Peshkin et al. 2013; Rosenthal et al. 2015; Davidson et al. 2021) and non-grouping

individuals, however. First, behavioural contagion can be complex, and the number of non-

alarmed individuals within the detection range influences the likelihood of adopting a specific

behaviour (Firth 2020). Previous works on social species have provided mounting evidence

for such complex contagion (Hoppitt and Laland 2013; Grüter and Leadbeater 2014; Kendal

et al. 2018). Second, imperfect copying might decrease the intensity of behavioural responses

with each transmission step, and under a given threshold intensity, social cues exert no

response from nearby observers. In this case, individuals' ability to convey information about

predation hazards is related to the extent of behavioural change compared to a baseline level

(Chivers and Ferrari 2014). Third, phenotypic heterogeneity among individuals may influence

information diffusion if individual traits (e.g., related to hunger, age or developmental stage)

or functional traits that transcend species (e.g., similarity in body size that may lead to shared

predators) affects the individual capacity to produce social information (Farine et al. 2015).

To describe how ISI use may affect population dynamics in non-grouping prey, we

constructed a tentative model with naturalistic predator-to-prey ratios (1:1.03 [when predator

detection probabilities was set to minimal]–1:4.23 [with nominal predator detection and ISI

use probabilities]; see in Donald and Anderson 2003). Previous observations indicate that

predator detection probability, which has been found to play a crucial role in the emergence of

social information-mediated effects in our study, can have a value within the upper half of the

range examined here (i.e., >0.5) under relevant conditions (e.g., Tisdale and Fernández-Juricic

2009; Manzur et al. 2018). However, being strongly dependent on the neuronal pathways

underlying detection mode and the processing capacity of the brain (Clark and Dukas 2003,

Pereira and Moita 2016), it can differ significantly between species and even within the same

species as it may also depend on the forager's state of energy reserves (Clark and Mangel

2000). Therefore, to construct a more realistic model, both species-specific and context-

specific information (e.g., movement distances, detection ranges and reproduction rates) for

existing predator-prey relationships need to be incorporated, which can be done only at the

expense of generality. Model precision may be further enhanced by incorporating additional

variables including the functional response of specific predator species (Dunn and Hovel

2020), different non-consumptive effects (other than reduced feeding rate) (Peckarsky et al.

2008), a measure of social cue reliability (Dunlap et al. 2016), social information use in

predators (Hämäläinen et al. 2021), or landscape heterogeneity that could alter the space use

of individuals (Albery et al. 2021). The effects of different transmission modes can also be

tested, for instance, by weighting the probability of information diffusion among conspecifics

by the proportion of alarmed and non-alarmed individuals within the detection zone or

incorporating heterogeneity among individuals in attributes that affect their propensity to act

as social cue producers. Our work, thus, represents a general modelling approach that could

be applied to predator-prey systems in which populations are demographically decoupled, and

prey may mitigate predation hazards through the exploitation of incidentally produced social

information.

Acknowledgements

We are indebted to Béla Keresztfalvi for providing access to the institutional IT resources,

with which we were able to reduce the computational time of the performed simulations

radically.

Statements and Declarations

Funding

ZT was financially supported by the Prémium Postdoctoral Research Programme of the

Hungarian Academy of Sciences (MTA, PREMIUM-2018-198), by the János Bolyai

Research Scholarship of the Hungarian Academy of Sciences (MTA, BO/00634/21/8) and by

the New National Excellence Program of the Ministry for Innovation and Technology (ITM,

ÚNKP-21-5-DE-478) from the source of the National Research, Development and Innovation

Fund. GC was financially supported by the Young Researcher Programme of the Hungarian

Academy of Sciences (MTA, Mv-41/2020).

Conflict of interest

The authors have no conflict of interest to declare.

Ethics approval

Not applicable.

Consent to participate

Not applicable.

Consent for publication

Not applicable.

Data availability

Data files supporting the results are archived and available at Figshare

(https://figshare.com/s/34fc714342dab9123193). Upon reasonable requests, R codes for the

model functions are also available from the corresponding author.

Authors' contributions

ZT and CG conceived and designed the study. ZT constructed the model, performed the

simulations, analysed the model output, wrote the initial manuscript, and revised and edited

the subsequent versions. CG contributed substantially to the text and revisions.

References

Albery GF, Morris A, Morris S, Pemberton JM, Clutton-Brock TH, Nussey DH, Firth JA

(2021) Multiple spatial behaviours govern social network positions in a wild ungulate.

Ecol Lett 24:676–686

Allenby RBJT, Slomson A (2010) How to Count: An Introduction to Combinatorics, Discrete

495   Mathematics and Its Applications. 2nd edn CRC Press, Boca Raton, pp 51–60

496   Beaudouin R, Goussen B, Piccini B, Augustine S, Devillers J, Brion F, Péry AR (2015) An

497   individual-based model of zebrafish population dynamics accounting for energy

498   dynamics. PLoS ONE 10:e0125841

499   Beleznai O, Dreyer J, Tóth Z, Samu F (2017) Natural enemies partially compensate for

500   warming induced excess herbivory in an organic growth system. Sci Rep 7(1):1–9

501   Benhamou S (2006) Detecting an orientation component in animal paths when the preferred

502   direction is individual-dependent. Ecology 87:518–528

503   Bonnie KE, Earley RL (2007) Expanding the scope for social information use. Anim Behav

504   74:171–181

505   Boujja-Miljour H, Leighton PA, Beauchamp G (2017) Spread of false alarms in foraging

506   flocks of house sparrows. Ethology 123:526–531

507   Brown GE, Godin JGJ, Pedersen J (1999) Fin-flicking behaviour: a visual antipredator alarm

508   signal in a characin fish, *Hemigrammus erythrozonus*. Anim Behav 58:469–475

509   Byers JA (2001) Correlated random walk equations of animal dispersal resolved by

510   simulation. Ecology 82:1680–1690

511   Campolongo F, Cariboni J, Saltelli A (2007) An effective screening design for sensitivity

512   analysis of large models. Environ Model Softw 22:1509–1518

513   Červený J, Begall S, Koubek P, Nováková P, Burda H (2011) Directional preference may

514   enhance hunting accuracy in foraging foxes. Biol Lett 7:355–357

515   Chivers DP, Ferrari MC (2014) Social learning of predators by tadpoles: does food restriction

516   alter the efficacy of tutors as information sources? Anim Behav 89:93–97

517   Ciric C, Ciffroy P, Charles S (2012) Use of sensitivity analysis to identify influential and non-

518   influential parameters within an aquatic ecosystem model. Ecol Modell 246:119–130

Clark CW, Dukas R (2003) The behavioral ecology of a cognitive constraint: limited

attention. Behav Ecol 14(2):151–156

Clark CW, Mangel M (2000) Dynamic state variable models in ecology. Oxford University

Press, New York

Codling EA, Plank MJ, Benhamou S (2008) Random walk models in biology. J R Soc

Interface 5:813–834

Coleman SW (2008) Mourning dove (*Zenaida macroura*) wing-whistles may contain threat-

related information for con-and hetero-specifics. Sci Nat 95:981

Confalonieri R, Bellocchi G, Bregaglio S, Donatelli M, Acutis M (2010) Comparison of

sensitivity analysis techniques: a case study with the rice model WARM. Ecol

Modell 221(16):1897–1906

Coolen I, Dangles O, Casas J (2005) Social learning in noncolonial insects? Curr Biol

15:1931–1935

Crane AL, Demers EE, Feyten LE, Ramnarine IW, Brown GE (2021) Exploratory decisions

of Trinidadian guppies when uncertain about predation risk. Anim Cogn DOI:

10.1007/s10071-021-01575-4

Croft DP, Madden JR, FranksDW, James R (2011) Hypothesis testing in animal social

networks. Trend Ecol Evol 26(10):502–507

Cruz A, Heinemans M, Marquez C, Moita MA (2020) Freezing displayed by others is a

learned cue of danger resulting from co-experiencing own freezing and shock. Curr Biol

30:1128–1135

Csardi G, Nepusz T (2006) The igraph software package for complex network research. Int J

Complex Syst 1695:1–9

Dall SR, Johnstone RA (2002) Managing uncertainty: information and insurance under the

risk of starvation. Philos Trans R Soc B 357:1519–1526

Dall SR, Giraldeau L-A, Olsson O, McNamara JM, Stephens DW (2005) Information and its

use by animals in evolutionary ecology. Trends Ecol Evol 20:187–193

Danchin E, Giraldeau L-A, Valone TJ, Wagner RH (2004) Public information: from nosy

neighbors to cultural evolution. Science 305:487–491

Davidson JD, Sosna MMG, Twomey CR, Sridhar VH, Leblanc SP, Couzin ID (2021)

Collective detection based on visual information in animal groups. J R Soc Interface

18:20210142

Donald DB, Anderson RS (2003) Resistance of the prey-to-predator ratio to environmental

gradients and to biomanipulations. Ecology 84(9):2387–2394

Duboscq J, Romano V, MacIntosh A, Sueur C (2016) Social information transmission in

animals: lessons from studies of diffusion. Front Psychol 7:1147

Dunlap AS, Nielsen ME, Dornhaus A, Papaj DR (2016) Foraging bumble bees weigh the

reliability of personal and social information. Curr Biol 26(9):1195–1199

Dunn RP, Hovel KA (2020) Predator type influences the frequency of functional responses to

prey in marine habitats. Biol Lett 16:20190758

Farine DR, Montiglio PO, Spiegel O (2015) From individuals to groups and back: the

evolutionary implications of group phenotypic composition. Trends Ecol Evol 30:609–

621

Farine DR (2017) A guide to null models for animal social network analysis. Methods Ecol

Evol 8:1309–1320

Ferrari MC, Trowell JJ, Brown GE, Chivers DP (2005) The role of learning in the

development of threat-sensitive predator avoidance by fathead minnows. Anim Behav

70(4):777–784

Firth JA, Sheldon BC (2016) Social carry-over effects underpin trans-seasonally linked

structure in a wild bird population. Ecol Lett 19:1324–1332

Firth JA (2020) Considering complexity: animal social networks and behavioural contagions.

Trends Ecol Evol 35:100–104

Galef Jr BG, Giraldeau L-A (2001) Social influences on foraging in vertebrates: causal

mechanisms and adaptive functions. Anim Behav 61:3–15

Gil MA, Hein AM (2017) Social interactions among grazing reef fish drive material flux in a

coral reef ecosystem. Proc Natl Acad Sci USA 114:4703–4708

Gil MA, Emberts Z, Jones H, St. Mary CM (2017) Social information on fear and food drives

animal grouping and fitness. Am Nat 189:227–241

Gil MA, Hein AM, Spiegel O, Baskett ML, Sih A (2018) Social information links individual

behavior to population and community dynamics. Trends Ecol Evol 33:535–548

Gil MA, Baskett ML, Schreiber SJ (2019) Social information drives ecological outcomes

among competing species. Ecology 100:e02835

Goodale E, Beauchamp G, Magrath RD, Nieh JC, Ruxton GD (2010) Interspecific

information transfer influences animal community structure. Trends Ecol Evol 25:354–

361

Goodale E, Beauchamp G, Ruxton GD (2017) Mixed-species groups of animals: behavior,

community structure, and conservation. Academic Press, London

Greggor AL, Thornton A, Clayton NS (2017) Harnessing learning biases is essential for

applying social learning in conservation. Behav Ecol Sociobiol 71(1):1–12

Grüter C, Leadbeater E (2014) Insights from insects about adaptive social information use.

Trends Ecol Evol 29:177–184

Haggerty MB, Anderson TW, Long JD (2018) Fish predators reduce kelp frond loss via a

trait-mediated trophic cascade. Ecology 99(7):1574–1583

Hämäläinen L, Hoppitt W, Rowland HM, Mappes J, Fulford AJ, Sosa S, Thorogood R (2021)

Social transmission in the wild can reduce predation pressure on novel prey signals. Nat

Commun 12:1–11

Hämäläinen L, Rowland HM, Mappes J, Thorogood R (2021) Social information use by

predators: expanding the information ecology of prey defences. Oikos 00:1–15

Hermann SL, Landis DA (2017) Scaling up our understanding of non-consumptive effects in

insect systems. Curr Opin Insect Sci 20:54–60

Hingee M, Magrath RD (2009) Flights of fear: a mechanical wing whistle sounds the alarm in

a flocking bird. Proc R Soc B 276:4173–4179

Hobson EA, Silk MJ, Fefferman NH, Larremore DB, Rombach P, Shai S, Pinter‑Wollman N

(2021) A guide to choosing and implementing reference models for social network

analysis. Biol Rev 96:2716–2734

Hoppitt W, Laland KN (2013) Social Learning: An Introduction to Mechanisms, Methods,

and Models. Princeton University Press, Princeton

Imron MA, Gergs A, Berger U (2012) Structure and sensitivity analysis of individual-based

predator–prey models. Reliab Eng Syst Saf 107:71–81

Iooss B, Lemaître P (2015) A review on global sensitivity analysis methods. In: Dellino G and

Meloni C (eds) Uncertainty management in simulation-optimization of complex

systems. Springer, Boston, pp 101–122

Johnson DS, London JM, Lea MA, Durban JW (2008) Continuous‑time correlated random

walk model for animal telemetry data. Ecology 89:1208–1215

Kareiva PM, Shigesada N (1983) Analyzing insect movement as a correlated random walk.

Oecologia 56:234–238

Kendal RL, Boogert NJ, Rendell L, Laland KN, Webster M, Jones PL (2018) Social learning

strategies: Bridge-building between fields. Trends Cogn Sci 22:651–665

King AJ, Cowlishaw G (2007) When to use social information: the advantage of large group

size in individual decision making. Biol Lett 3(2):137–139

Komin N, Erdmann U, Schimansky-Geier L (2004) Random walk theory applied to daphnia

motion. Fluctuation Noise Lett 4:L151–L159

Krause J, Ruxton GD (2002) Living in groups. Oxford University Press, New York

Laland KN (1992) A theoretical investigation of the role of social transmission in evolution.

Ethol Sociobiol 13:87–113

Latora V, Marchiori M (2001) Efficient behavior of small-world networks. *Phys Rev*

Lett 87(19):198701

Lima SL (1990) The influence of models on the interpretation of vigilance. In: Bekoff M and

Jamieson D (eds) Interpretation and explanation in the study of animal behavior:

Explanation, evolution and adaptation. Vol 2. Westview Press, Boulder, pp 246–267

Lima SL, Dill LM (1990) Behavioral decisions made under the risk of predation: a review and

prospectus. Can J Zool 68(4):619–640

Lohmann AC, Evangelista D, Waldrop LD, Mah CL, Hedrick TL (2016) Covering ground:

Movement patterns and random walk behavior in *Aquilonastra anomala* sea stars. Biol

Bull 231:130–141

Manassa RP, McCormick MI, Chivers DP, Ferrari MCO (2013) Social learning of predators

in the dark: understanding the role of visual, chemical and mechanical information. Proc

R Soc B 280(1765):20130720

Manzur T, Gonzalez-Mendez A, Broitman BR (2018) Scales of predator detection behavior

and escape in Fissurella limbata: A field and laboratory assessment. Mar Ecol

39(2):e12492

Martín J, Luque-Larena JJ, López P (2006) Collective detection in escape responses of

temporary groups of Iberian green frogs. Behav Ecol 17:222–226

McCulloch CE, Cain ML (1989) Analyzing discrete movement data as a correlated random

walk. Ecology 70(2):383–388

Michálek O, Petráková L, Pekár S (2017) Capture efficiency and trophic adaptations of a

specialist and generalist predator: a comparison. Ecol Evol 7:2756–2766

Morris MD (1991) Factorial sampling plans for preliminary computational experiments.

Technometrics 33:161–174

O'Mara MT, Dechmann DK, Page RA (2014) Frugivorous bats evaluate the quality of social

information when choosing novel foods. Behav Ecol 25(5):1233–1239

Parejo D, Avilés JM (2016) Social information use by competitors: resolving the enigma of

species coexistence in animals? Ecosphere 7:e01295

Pasquaretta C, Levé M, Claidiere N, Van de Waal E, Whiten A, MacIntosh AJ, et al (2014).

Social networks in primates: smart and tolerant species have more efficient networks.

Sci Rep 4:1–8

Pays O, Beauchamp G, Carter AJ, Goldizen AW (2013) Foraging in groups allows collective

predator detection in a mammal species without alarm calls. Behav Ecol 24:1229–1236

Peckarsky BL, Abrams PA, Bolnick DI, Dill LM, Grabowski JH, Luttbeg B, et al (2008)

Revisiting the classics: considering nonconsumptive effects in textbook examples of

predator–prey interactions. Ecology 89:2416–2425

Pereira AG, Moita MA (2016) Is there anybody out there? Neural circuits of threat detection

in vertebrates. Curr Opin Neurobiol 41:179–187

Pessarrodona A, Boada J, Pagès JF, Arthur R, Alcoverro T (2019) Consumptive and non-

consumptive effects of predators vary with the ontogeny of their prey. Ecology

100(5):e02649

Polis GA, Sears AL, Huxel GR, Strong DR, Maron J (2000) When is a trophic cascade a

trophic cascade? Trends Ecol Evol 15(11):473–475

Preisser EL, Orrock JL, Schmitz OJ (2007) Predator hunting mode and habitat domain alter

nonconsumptive effects in predator–prey interactions. Ecology 88(11):2744–2751

R Core Team (2021) R: A language and environment for statistical computing. R foundation

for statistical computing, Vienna, Austria.

Reynolds AM (2014) Towards a mechanistic framework that explains correlated random walk

behaviour: correlated random walkers can optimize their fitness when foraging under

the risk of predation. Ecol Complex 19:18–22

Romano V, Shen M, Pansanel J, MacIntosh AJ, Sueur C (2018) Social transmission in

networks: global efficiency peaks with intermediate levels of modularity. Behav Ecol

Sociobiol 72:1–10

Rosenthal SB, Twomey CR, Hartnett AT, Wu HS, Couzin ID (2015) Revealing the hidden

networks of interaction in mobile animal groups allows prediction of complex

behavioral contagion. Proc Natl Acad Sci USA 112:4690–4695

Sih A, Hanser SF, McHugh KA (2009) Social network theory: new insights and issues for

behavioral ecologists. Behav Ecol Sociobiol 63:975–988

Spiegel O, Leu ST, Sih A, Bull CM (2016) Socially interacting or indifferent neighbours?

Randomization of movement paths to tease apart social preference and spatial

constraints. Methods Ecol Evol 7:971–979

Strandburg-Peshkin A, Twomey CR, Bode NW, Kao AB, Katz Y, Ioannou CC, et al (2013)

Visual sensory networks and effective information transfer in animal groups. Curr Biol

23:R709–R711

Ten Broeke G, Van Voorn G, Ligtenberg A (2016) Which sensitivity analysis method should

I use for my agent-based model? J Artif Soc Soc Simul 19:5

Tholt G, Kis A, Medzihradszky A, Szita É, Tóth Z, Havelda Z, Samu F (2018) Could vectors'

fear of predators reduce the spread of plant diseases? Sci Rep 8(1):1–10

Thorogood R, Kokko H, Mappes J (2018) Social transmission of avoidance among predators

facilitates the spread of novel prey. Nat Ecol Evol 2(2):254–261

Tisdale V, Fernández-Juricic E (2009) Vigilance and predator detection vary between avian

species with different visual acuity and coverage. Behav Ecol 20(5):936–945

Toland BR (1987) The effect of vegetative cover on foraging strategies, hunting success, and

nesting distribution of American kestrels in central Missouri. J Raptor Res 21:14–20

Tóth Z (2021) The hidden effect of inadvertent social information use on fluctuating

predator–prey dynamics. Evol Ecol 35:101–114

Tóth Z, Jaloveczki B, Tarján G (2020) Diffusion of social information in non-grouping

animals. Front Ecol Evol 8:586058

Uttieri M, Mazzocchi MG, Nihongi A, D'Alcalà MR, Strickler JR, Zambianchi E (2004)

Lagrangian description of zooplankton swimming trajectories. J Plankton Res 26:99–

105

Uttieri M, Paffenhöfer GA, Mazzocchi MG (2008) Prey capture in *Clausocalanus furcatus*

(Copepoda: Calanoida). The role of swimming behaviour. Mar Biol 153:925–935

Ward A, Webster M (2016) Sociality: The Behaviour of Group-Living Animals. Springer

International Publishing, New York

Waters JS, Fewell JH (2012) Information processing in social insect networks. PLoS ONE

7:e40337

Watson CG (2020) brainGraph: Graph Theory Analysis of Brain MRI Data. R package

version 3.0.0. https://CRAN.R-project.org/package=brainGraph

Webber QM, Vander Wal E (2018) An evolutionary framework outlining the integration of

individual social and spatial ecology. J Anim Ecol 87:113–127

Weissburg M, Smee DL, Ferner MC (2014) The sensory ecology of nonconsumptive predator

effects. Am Nat 184(2):141–157

White JW, Rassweiler A, Samhouri JF, Stier AC, White C (2014) Ecologists should not use

statistical significance tests to interpret simulation model results. Oikos 123:385–388

Whitehead H, Laland KN, Rendell L, Thorogood R, Whiten A (2019) The reach of gene–

culture coevolution in animals. Nat Comm 10:1–10

Table 1. Model parameters and their range for sensitivity analysis (SA). Maximum movement

distances indicate the maximum number of spatial units that an individual could travel on the

landscape in a simulation cycle; the actual integer value was randomly selected between zero

and this maximum value.

Symbol	Description	Nominal value	Unit/Scale	Range for SA
d_{prey}	Prey maximum movement distance	5	1	3-8
d_P	Predator maximum movement distance	5	1	3-8
r_{prey}	Prey detection range	2.5 (constant)	-	-
r_P	Predator hunting range	3.5 (constant)	-	-
r_c	Competition range within which only one individual could survive	1 (constant)	-	-
P_{detect}	Probability of prey detecting a nearby predator; determined by individual Bernoulli trials	0.5	0.1	0.1-0.9
P_{isi}	Probability of prey ISI use (i.e., copying the defensive behaviour of others); determined by individual Bernoulli trials	0.5	0.1	0-0.9
λ_{max}	Prey reproduction-related shape parameter when prey feed; used to draw a random number of offspring for each individual from a Poisson distribution	0.75 (constant)	-	-
$\lambda_{reduced}$	Prey reproduction-related shape parameter	0.75	5×10^{-2}	0.5-0.75

	when prey is alarmed (i.e., does not feed); used to draw a random number of offspring for each individual from a Poisson distribution			
λ_P	Predator reproduction-related shape parameter; used to draw a random number of offspring for each individual from a Poisson distribution	5×10^{-2}	5×10^{-3}	2.5×10^{-2}- 7.5×10^{-2}

726   Table 2. Descriptive statistics of the simulated prey populations calculated from 50 replicates

727   at the 200^{th} simulation cycle (49 in the case of the second model type as prey population died

728   out in a single iteration).

Models	Parameters		Prey population size	
	P_{detect}	P_{isi}	Mean ± SD	Range
No predators (n=50)	-	-	1198.16 ± 24.37	1115-1251
Minimal predator detection (n=49)	0.1	0	154.45 ± 56.49	24-257
Nominal predator detection (n=50)	0.5	0	516.18 ± 50.36	408-637
Nominal predator detection & ISI use (n=50)	0.5	0.5	789.52 ± 48.05	632-891

 Figure legends

Figure 1. Model flowchart for a single simulation cycle. Sequential prey and predator

behaviours are listed together with the model parameter(s) associated with the given steps.

Behavioural steps resulting in a decrease in population size, i.e., mortality due to intraspecific

competition (in rounded rectangles) or predation (in diamond) are shown in light and dark

grey, respectively.

Figure 2. Schematic figure of a detection network (a) and segment of an individual ego

network embedded within that network (b). Nodes represent individuals and edges denote the

possibility of mutual observation. The probability of information acquisition from one node to

another is given by w^k, where w is the edge weight and k is the number of steps on the shortest

path between the two nodes. For any individual, the total probability of receiving information

from neighbours is calculated using the inclusion-exclusion principle. In our model, we used

the settings $k_{max}=2$ and $\sum n_{max}=5$ in each k step, so the focal individual (black circle) could

receive social information from a maximum of ten neighbours that were a maximum of two

steps away in the detection network (orange circles).

Figure 3. Effects of the P_{detect} and P_{isi} model parameters on the prey population. (a) Temporal

fluctuations in prey abundance (means with range) without predators (grey) and under

nominal parameter settings (with nominal P_{detect} & P_{isi} – orange, with nominal P_{detect} – dark

blue, with minimal P_{detect} – light blue). When the predator detection probability was set to its

minimal value, the prey population died out in a single iteration; in all other cases, the number

of iterations was set to 50. (b) Results of the global sensitivity analysis (SA) depicting the

impact of each model parameter on the mean (x-axis) and standard deviation (y-axis) of prey

abundance; mean $\pm$ SD values for each parameter were calculated from five independent SA

runs. Inset shows the model parameters ordered according to their overall influence on the

model output.

Figure 4. The relationship between per capita mortality due to predation and the number of

predators using the same parameter settings as in Fig. 3a (but without the 'No predators'

group). Trend lines were fitted using second-order polynomial approximation. Simulation

results from incomplete runs (i.e., simulation cycles were less than 200) were omitted from

the dataset ($n=204$; only in the 'minimal predator detection' model type).

Figure 5. Interactive effects of the probability of ISI use (P_{isi}), predation pressure (λ_P) and the

presence of fitness cost (associated with the defensive behaviour; $\lambda_{reduced}$) on prey population

size in three P_{detect} scenarios. The colour of the boxplots indicates the level of predation

pressure (purple: high, blue: intermediate, green: low), while the colour tone is associated

with the presence of cost (dark: costly defensive behaviour, light: no cost). Trend lines were

fitted using the 'LOESS' regression method for smoothing with the default value of span

(0.75); presented only for illustration purposes. Simulation results from incomplete runs were

omitted from the dataset ($n=581$; only in the '$P_{detect}=0.25$' setting).

Figure 6. Four structural network properties (a: number of components, b: component size, c:

average size of ego networks, d: average global network efficiency) calculated for the

observed detection networks (circles) and corresponding randomized networks (diamonds).

Predation pressure was set to 'high' (i.e., $\lambda_P=0.075$). The colour of the boxplots indicates the

absence (gray; $P_{isi}=0$) or presence of ISI use (orange; $P_{isi}=0.5$).

Cycle starts

Cycle ends

(a)

(b)

$k=1$ $k=2$ $k=3$

w w w

w w

w w

w w

w w

$n_{k=1}=2$ $n_{k=2}=3$ $n_{k=3}=3$

Prey abundance (y-axis), P_{isi} (x-axis)

$P_{detect} = 0.75$ $P_{detect} = 0.5$ $P_{detect} = 0.25$

Low predation pressure & costly defensive behaviour ($\lambda_P = 0.025$; $\lambda_{reduced} = 0.5$)

Low predation pressure & no cost ($\lambda_P = 0.025$; $\lambda_{reduced} = \lambda_{max} = 0.75$)

Intermediate predation pressure & costly defensive behaviour ($\lambda_P = 0.05$; $\lambda_{reduced} = 0.5$)

Intermediate predation pressure & no cost ($\lambda_P = 0.05$; $\lambda_{reduced} = \lambda_{max} = 0.75$)

High predation pressure & costly defensive behaviour ($\lambda_P = 0.075$; $\lambda_{reduced} = 0.5$)

High predation pressure & no cost ($\lambda_P = 0.075$; $\lambda_{reduced} = \lambda_{max} = 0.75$)