Complete chloroplast genomes and phylogeny in three *Euterpe* palms (*E. edulis*, *E. oleraceae* and *E. precatoria*) from different Brazilian biomes

Ana Flávia Francisconi[1], Luiz Augusto Cauz dos Santos[2], Jonathan Andre Morales Marroquín[1], Cássio van den Berg[3,4], Alessandro Alves-Pereira[5], Luciano Delmondes de Alencar[1], Doriane Picanço-Rodrigues[6], Cesar Augusto Zanello[1], Marcones Ferreira Costa [1,7], Maria Teresa Gomes Lopes[8], Elizabeth Ann Veasey[4], Maria Imaculada Zucchi[9]

[1] Programa de pós-gradução em Genética e Biologia Molecular Universidade Estadual de Campinas, Campinas, São Paulo, Brasil

[2] Department of Botany and Biodiversity Research, University of Vienna, Wien, Austria

[3] Departamento de Ciências Biológicas, Universidade Estadual de Feira de Santana, Feira de Santana, Bahia, Brasil

[4] Departamento de Genética, Universidade de São Paulo, Piracicaba, São Paulo, Brasil

[5] Departament de Biologia Vegetal, Universidade Estadual de Campinas, Campinas, São Paulo, Brasil

[6] Departamento de Biologia, Universidade Federal do Amazonas (ICB-UFAM), Manaus, Amazonas, Brasil

[7] Universidade Federal do Piauí, Floriano, Piauí, Brasil

[8] Departamento de Produção Animal e Vegetal, Universidade Federal do Amazonas, Manaus, Amazonas, Brasil

[9] Agência Paulista de Tecnologia dos Agronegócios, Piracicaba, São Paulo, Brasil

*Corresponding author
E-mail: mizucchi@sp.gov.br (MIZ), anaf_f@hotmail.com (AFF)

Abstract

The Brazilian palm fruits and hearts-of-palm of *Euterpe edulis, E. oleracea and E. precatoria* are an important source for agro-industrial production, but overexploitation requires conservation strategies to maintain genetic diversity. Chloroplast genomes have conserved sequences, which are useful to explore evolutionary questions. Besides the plastid DNA, genome skimming allows the identification of other genomic resources, such as single nucleotide polymorphisms (SNPs), providing information about the genetic diversity of species. We sequenced the chloroplast genome and identified gene content in the three *Euterpe* species. We performed comparative analyses, described the polymorphisms among the chloroplast genome sequences (repeats, indels and SNPs) and performed a phylogenomic inference based on 55 palm species chloroplast genomes. Finally, using the remaining data from genome skimming, the nuclear and mitochondrial reads, we identified SNPs and estimated the genetic diversity among these *Euterpe* species. The *Euterpe* chloroplast genomes varied from 159,232 to 159,275 bp and presented a conserved quadripartite structure with high synteny with other palms. In a pairwise comparison, we found a greater number of insertions/deletions (indels = 93 and 103) and SNPs (284 and 254) between *E. edulis*/*E. oleracea* and *E. edulis*/*E. precatoria* when compared to *E. oleracea*/*E. precatoria* (58 indels and 114 SNPs). Also, the phylogeny indicated a closer relationship between *E. oleracea*/*E. precatoria*. The nuclear and mitochondrial genome analyses identified 1,077 SNPs and high divergence among species ($F_{ST} = 0.77$), especially between *E. edulis* and *E. precatoria* ($F_{ST} = 0.86$). These results showed that, despite the few structural differences among the chloroplast genomes of these *Euterpe* palms, a differentiation between *E. edulis* and the other *Euterpe* species can be identified by point mutations. This study not only brings new knowledge about the

69 evolution of *Euterpe* chloroplast genomes, but also these new resources open the way for

70 future phylogenomic inferences and comparative analyses within Arecaceae.

71
72 **Key-Words**
73 Arecaceae, SNPs, genome-skimming, genome comparison, polymorphism

Introduction

The palm family (Arecaceae) comprises 188 genera and ca. 2,400 species, distributed

throughout tropical and subtropical regions [1,2]. Palms are often keystone species,

providing ecosystem services, shaping their environment and offering many products

used for fabrics, fuel, food, medicine and as ornamentals [2–4]. Within Arecaceae, the

genus *Euterpe* originated in South America and includes seven species distributed

throughout Central America and tropical South America [2,5,6]. In Brazil, among the

native species of *Euterpe*, *E. edulis* Mart., *E. oleracea* Mart. and *E. precatoria* Mart. are

the most important from an agro-industrial point of view [7].

Euterpe edulis is endemic to the Atlantic Forest and has substantial importance

for the functioning of the ecosystem [8,9]. However, the overexploitation of heart-of-

palm [10,11] has placed this palm species in the Brazilian Flora Red Book [12] as

"vulnerable (VU)", which makes conservation strategies for it extremely necessary.

Recently, as an alternative to palm heart-of-palm extraction, local communities have

started to cultivate this species for fruit pulp extraction, due to the similarity with the

Amazonian açaí (*E. oleracea* and *E. precatoria*) [9,10,13]. This practice has been

fundamental to capture and conserve the species genetic diversity present in older forests

[10].

Euterpe oleracea and *E. precatoria* are the main sources for the production of açaí

fruit pulp, estimated to be 760,000 tons in 2017, mostly in the northern region of Brazil

(98,36% of the national production) [14,15]. *E. oleracea* is a multiple-stemmed palm,

commonly found in the Amazon estuary, especially prevalent in the Brazilian states of

Pará, Maranhão and Amapá [15–17]. However, in the last 20 years, the intense demand

for açaí fruit in Pará resulted in a significant loss of local tree species richness, which

presumably is occurring after decades of thinning to reduce interspecific competition with

açai palm trees [18]. This resulted in a reduction of pollinators, which in turn decreased

the fruit production [19]. *E. precatoria* is a single stemmed palm, predominantly found

in the Brazilian states of Amazonas, Acre and Rondônia [20], in non-flooded areas [21].

This species is highly promising for açaí fruit extraction, as the state of Amazonas is

already the second largest producer of Brazil [15]. Therefore, the establishment of

strategies for *E. precatoria* domestication, conservation and management of populations

is extremely necessary, in addition to the support of riverside communities and farmers

to conserve this species based on their genetic diversity [15].

The fast growth of açaí (*E. oleracea* and *E. precatoria*) pulp production in the last

decades and the threats of extinction for *E. edulis* indicate an urgent demand for new

genomic resources in this genus, contributing to the understanding of the evolution and

population dynamics of the species, which can result in the best planning of sustainable

development and conservation [9,15]. Fortunately, high-throughput sequencing

technologies have revolutionized how genomic data can be obtained for plant species and

have allowed the complete assembly of the chloroplast genome for an increasing number

of non-model species [22–24], while providing useful information for the study of genetic

variation of native species.

The common structure of the chloroplast genomes typically has four parts: two

single copy regions, one large (LSC) and one small (SSC), and a pair of inverted regions

(IRs) [25] which basically consist of long circular or linear molecules (120 to 180 kb),

usually conserved across palms species [26]. Apart from the conservative nature and low

nucleotid substitution rates of the chloroplast sequences [27], the description of this

organellar genome is important to identify repetitive sequences and polymorphic regions,

useful to further obtain molecular markers, which can be applied to assess genetic

structure and diversity of natural populations [28,29]. Whole chloroplast genomes and

chloroplast sequences have been used to infer phylogenetic relationships and investigate

diversification patterns [24,30–32]. The tribe Euterpeae was already analyzed with

combined chloroplast and nuclear sequence data [33], and whole chloroplast genomes

have been reported in reference for two species in the *Euterpe* genus (*E. oleracea* and *E.*

edulis) [34]. These priors research emphasize the differentiation of *E. edulis*, due to its

distribution in a different Brazilian biome (Atlantic Forest biome). But, the addition of

new chloroplast genomes can give more consistency to the evolutionary processes of the

genus.

Moreover, the amount of plastid sequences present in a data from total genomic

DNA extraction depends on different factors and can vary substantially, from 0.4% to

29.5% [23], and the recovery of these sequences can occur through the use of genome

skimming. Therefore, the remaining nuclear and mitochondrial sequences can provide

genomic resources [27,35], representing useful information to study the genetic variation

of native species.

Considering the effectiveness of genome skimming to produce genomic data, we

employed this approach to obtain three newly sequenced complete chloroplast genomes,

thereby facilitating the study of genetic diversity from *Euterpe* species. We were able to

investigate: i) the plastid organization in *Euterpe* and the synteny level with other

available Arecaceae species; ii) the repetitive sequences and polymorphic regions in the

chloroplast genomes; iii) the Arecaceae family evolution based on a phylogenomic study

with complete chloroplast genomes; and iv) the species divergence using the remaining

nuclear and mitochondrial data from genome-skimming.

Material and Methods

Leaf material

The leaf material from *Euterpe edulis* and *E. oleracea* were collected from the *ex*

situ collection at the Escola Superior de Agricultura "Luiz de Queiroz" (ESALQ),

Universidade de São Paulo (USP) in Piracicaba, São Paulo, Brazil

(http://www.esalq.usp.br/trilhas/palm/) and the sample of *E. precatoria* was collected at

the Instituto Nacional de Pesquisas da Amazônia (INPA), in Manaus, AM, Brazil. The

collections were registered according to the Brazilian laws (SISGEN number A411583,

Brazil).

Intact chloroplast isolation in sucrose gradient and chloroplast

DNA extraction

The chloroplast organelles were isolated using the sucrose gradient method [36].

About 20 g of fresh leaves from each species were frozen with liquid nitrogen and

macerated. The material was resuspended in 200 mL of isolation buffer (50 mM Tris-HCl

pH 8.0, 0.35 M sucrose, 7 mM EDTA, 5 mM 2-mercaptoethanol and 0.1% BSA) and

incubated for 10 min in the dark. The suspension was filtered through two layers of

Miracloth (Merck), and the filtrate was centrifuged at $1,000 \times g$ for 10 min.

The pellet was resuspended in 5 mL of isolation buffer and the suspension slowly

laid out in the density gradients of 20/45% sucrose in 50 mM Tris-HCl (pH 8.0), 0.3 M

sorbitol and 7 mM EDTA. The gradients were centrifuged at $2000 \times g$ for 30 min, and

the green band formed at the interface containing intact chloroplasts were collected. The

solution containing the chloroplasts were diluted in three volumes of buffer and

centrifuged at 3,000 × g for 10 minutes to obtain the purified chloroplasts in the pellet.

The pellet was resuspended in 2% CTAB buffer to promote lysis. The suspension

was incubated and stirred at 65°C for 1 h. The supernatant was extracted twice with an

equal volume of chloroform: isoamyl alcohol (24: 1) and centrifuged at 10,000 × g for 20

min. An equal volume of isopropanol was added and incubated at 20°C for 1h. Finally,

the aqueous phase was centrifuged for 10,000 × g for 20 min, and the chloroplast DNA

(cpDNA) pellet was washed with ethanol (70%), dried and resuspended with 40 µL TE

(1 M Tris-HCl, 0.5 M EDTA, pH 8).

Chloroplast genome sequencing, assembly and annotation

The genomic libraries were constructed using 100 ng of cpDNA and the Nextera

DNA Flex kit (Illumina), following the manufacturer's instructions. Paired-end

sequencing (2x 150 bp) was performed on the Illumina NextSeq550 platform (Fundação

Hemocentro de Ribeirão Preto, Brazil).

The assembly was conducted in three steps: First, the filtered reads from *Euterpe*

oleracea   were   assembled   in   NOVOPlasty   v   4.2   [37]

(https://github.com/ndierckx/NOVOPlasty) using the *rbcL* gene sequence as a seed

(NCBI accession number: MN621452.1) and the chloroplast genome of *Acrocomia*

aculeata (NCBI accession number: NC_037084.1), a native Brazilian palm, as a reference

to ordinate the contigs. Subsequently, all the three chloroplast genomes (*E. precatoria, E.*

edulis and *E. oleracea*) were assembled in GetOrganelle v 1.7.3.1 [38]

(https://github.com/Kinggerm/GetOrganelle/) using the *E. oleracea* chloroplast sequence

obtained in NOVOPlasty as seed. Briefly, in GetOrganelle, the chloroplast genomes were

assembled using default settings, starting with the recruitment of target-associated reads

using Bowtie2 [39]. In this step, the seed chloroplast genome was used as target for the

extension in five iterations. Then, the total target-associated reads were *de novo*

assembled into a fasta assembly graph using SPAdes [40]. The *de novo* assemblies were

trimmed and used to calculate all possible paths of a complete organelle genome [38].

Finally, the correctness and coverage of the assembly was assessed and confirmed in

Geneious v2020 2.4. (https://www.geneious.com/, last assessed January, 2021). We used

the "Map to reference" function to map the paired-end raw data onto the final assembled

chloroplast genomes.

The chloroplast genome annotation was performed in GeSeq (Organellar Genome

Annotation) [41] from the Chlorobox platform, with settings for the identification of

protein coding sequence (CDS), rRNAs and tRNAs based on reference chloroplast

sequences and homologies through BLAST search. Following the automatic annotation,

a manual correction of start and stop codons and a verification of pseudogenes and intron

positions were performed using GenomeView [42]. We then obtained the chloroplast

circular genome maps using OGDRAW [43].

Chloroplast genome structure comparison

To perform a comparative study and access the synteny between the obtained

chloroplast sequences, we used a Perl script of MUMmer4 [44]

(https://github.com/mummer4/mummer) to align the chloroplast genome from *E. edulis,*

E. oleracea and *E. precatoria* with the function NUCmer. This analysis enables the

identification of the conserved regions among sequences between species. The results

were visualized in dot plots created by the function MUMmerplot. Additionally, to

compare and align the *Euterpe* chloroplast sequences obtained with the GenBank

sequences [34], MAFFT v.7 [45] was used as a way to identify differences between them.

The location of the additional sequences found between them was also annotated in

GeSeq [41].

Taking into account the species relationships from Arecoideae subfamily, and the

previous evidence from palms chloroplast genome structures [28,46], we decided to

conduct two multiple progressive sequence alignment in Mauve v.2.4.0 [47]. The first

one only included chloroplast genomes from Brazilian native species of subfamily

Arecoideae: *Euterpe edulis*, *Euterpe oleracea*, *Syagrus coronata*, *Astrocaryum*

aculeatum, *Astrocaryum murumuru* and *Acrocomia aculeata*. The second analysis was

carried out using 17 chloroplast genomes from different palm species (S1 Table) available

in GenBank. Taking into the account the evolution of the group, we selected the species

that represented the five palm subfamilies: *Phytelepas aequatoriallis* and *Pseudophoenix*

vinifera (Subfamily: Ceroxyloideae); *Trachycarpus fortune* and *Caryota mitis*

(Subfamily: Coryphoideae); *Nypa fruticans* (Subfamily: Nypoideae); *Calamus*

caryotoides and *Eresmopatha macrocarpa* (Subfamily: Calamoideae); *Veitchia arecina*

(Subfamily: Arecoideae) and the Brazilian native species also from the Arecoideae

subfamily.

Expansions and contractions in the inverted repeats (IR) regions from the

chloroplast genome structure were also explored. The chloroplast genome junctions

(IRB/LSC; IRB/SSC, SSC/IRA; IRA/LSC) from the *Euterpe* species, and other Brazilian

palm species from subfamily Arecoideae were examined to identify differences between

individuals within the same genus and among subfamilies.

Identification of SSRs, dispersed repeats, indels, SNPs and nucleotide divergence hotspots

Simple sequence repeats (SSR) consisting of 1-6 nucleotide units were carefully

determined using the web package MISA (available at https://webblast.ipk-

gatersleben.de/misa/) [48]. The criteria to search SSR motifs were: SSR of one to six

nucleotides long, with a minimum repeat number of 10, 5, and 4 units for mono-, di-, and

trinucleotide SSRs, respectively, and three units for tetra-, penta- and hexanucleotide

SSRs. The SSRs sequences and location were compared among the species.   The

dispersed repeats (forward, reverse, palindrome and complement sequences) identified in

REPuter [49] were based on the following criteria: minimum repetition size $\geq$ 30 bp and

sequence identity $\geq$ 90% (Hamming distance = 3). Posteriorly, the position of the SSRs

and repeats were manually compared with the gene annotation of each chloroplast

genome.

MAFFT v.7 [45] was used to obtain pairwise alignments between the chloroplast

genomes to pinpoint small insertions/deletions (indels) in the sequences. The alignment

between the three species was also used to identify single nucleotide polymorphisms

(SNPs) and nucleotide divergence hotspots with DnaSP v.5 [50]. Specific coding genes

with a high number of SNPs were aligned with the software Muscle [51] in the Selecton

Server [52], which was also used to identify the synonymous (Ka) or non-synonymous

(Ks) mutations. With this analysis, according with the Ka/Ks ratios, it was possible to

determine positive (Ka/Ks > 1) or purifying (Ka/Ks < 1) selection, estimated under the

evolutionary model M8.

A sliding window analysis (window length of 200 bp and step size of 50 bp) was

conducted to find nucleotide divergence hotspots. All the positions of indels, SNPs and

divergence hotspots were manually identified using the annotation of the aligned

chloroplast genomes, performed previously in GeSeq (Organellar Genome Annotation)

[41].

Phylogenomic studies

Plastid sequences of 54 palm species plus one outgroup (*Daypogon bromeliifolius*,

Dasypogonaceae) (S1 Table) curated annotated features in Genbank format were

separated into genes and entered into a SQLITE database using a custom Python script

(available on demand from the corresponding author). Sequences of all putative coding

proteins were then grouped by species and aligned with MAFFT, and later assembled into

an interleaved NEXUS file for phylogenetic analyses. Species with missing sequences

were filled with missing data in each gene matrix. Each block of gene sequences was then

manually checked for start and stop codons and evidence of non-coding behaviour. For

the coding genes, we annotated first, second and third positions for each codon, but the

regions *cemA* and *rpl16* presented strange start codons and non-triplet insertions and were

not annotated for codon positions. We tested three *ad hoc* partitioning schemes for models

of molecular evolution: i) single model for the whole matrix, ii) four partitions: 1^{st}, 2^{nd},

3^{rd} codon positions and a partition for *cemA +rpl16* (not split into codon positions), iii)

five partitions: 1^{st}, 2^{nd}, 3^{rd} codon positions, *cemA* (not split into codon positions)*, rpl16*

(not split into codon positions). The assessment of molecular evolution model for each

partition was calculated in the different partition schemes with the Akaike Information

Criterion (AIC) in MrModelTest 2.0 [53], but they were all GTR+I+G, probably due to

the complexity of mixing different genes in each partition. The assessment of the best

partition scheme was then made using stepping stone (SS) [54] which is more accurate

than taking the harmonic means of the likelihoods for model comparison. Bayesian

inference were carried out with MrBAYES 3.2.7 [55] in the CIPRES platform [56], with

one run of two chains, and 10×10^{6} generations, sampling one tree every 1,000. Marginal

likelihoods of the SS runs were then compared using the standard Bayes Factors scale

[57]. The same partition schemes were run separately to estimate the phylogeny with two

runs of four chains (three hot and one cold), and 20×10^6 generations, sampling one tree

every 1,000 and discarding 25% initial generations for burn-in. The remaining trees were

checked for estimated sample sized (ESS) > 200 in all parameters, and the final majority-

rule was computed with MrBayes, rooted with the single outgroup *D. bromeliifolius* and

the tree was exported to FigTree 1.4 [58] for drawing, and later improved with InkScape

[59] for designing the figures.

Genetic variation and comparative analyses with nuclear and

mitochondrial sequences

After the assembly of the chloroplast genomes, the reads were also used to obtain

SNPs for preliminary comparative analysis between species. Initially, the raw reads of

each species were aligned in their respective assembled chloroplast genome using

Bowtie2 [39]. Subsequently, the non-aligned sequences (non-chloroplast) were obtained

with SAMTools [60], and converted to fastq file using Picard Tools program [61].

The reads corresponding to non-chloroplast genome (nuclear and mitochondrial

data) were cleaned using *process_shortreads* from the Stacks package v.1.42 [62].

Sequences from each species were used to build loci with a minimum depth of three and

up to two mismatches, applying the *ustacks* program (-m 3, -M 2) [62]. Subsequently, a

catalog was built from each species, allowing for up to two mismatches (-n 2, cstacks)

[62], and using *sstacks* [62] loci were matched with the catalog of each species. *Rxstacks*

[62] was used as a correction step, assuming a mean log likelihood of -10 to discard loci

with lower probabilities. The *populations* [62] program was administered for a final

filtering to retain loci with a maximum missing data of 5%, and also used to verify the

number of private alleles (*Ap*), observed heterozygosity (H_O) and expected heterozygosity

(H_E; S1 Fig).

Finally, the mean sequence depth and the mutation counts of the SNPs were

determined in vcftools [63]. The number of alleles (*A*) in each species and the genetic

differentiation (F_{ST}) between them were estimated with *Adegenet* v 1.3 -1 [64] and

Genpop v 1.1.7 [65] for the platform R v 4.0.3 [66] .

Results

Organization of the three *Euterpe* species chloroplast genomes

The chloroplast genomes of *E. edulis, E. oleracea* and *E. precatoria* had the

typical quadripartite structure (Fig 1), with the presence of two copies of inverted repeat

regions (Inverted Repeats A and B = IRA and IRB) and two single copy regions (Large

Single Copy = LSC and Small Single Copy = SSC). *E. precatoria* had the largest

chloroplast genome size (159,275 bp) and the largest LSC and SSC (87,282 bp and 17,756

bp, respectively), while *E. edulis* had the largest IR regions and *E. oleracea* a slightly

larger amount of GC content (37,3%) (Table1). All three chloroplast genome annotations

resulted in the identification of 113 unique genes, with 30 tRNAs, 4 rRNAs and 79 protein

coding genes (Fig 1, Table 2).

Table 1. General features of chloroplast genomes of three *Euterpe* species.

Species	Total cpDNA size (bp)	Length of LSC region (bp)	Length of IR region (bp)	Length of SSC region (bp)	GC content (%)	Protein coding genes (bp)	tRNA coding genes (bp)	rRNA coding genes (bp)	Introns (bp)	Intergenic Regions (bp)
Euterpe edulis	159,232	87,237	54,280	17,715	37.20	80,462	2,880	9,052	18,068	48,770
Euterpe	159,260	87,273	54,232	17,755	37.30	80,415	2,881	9,052	18,062	48,850

oleracea									
Euterpe precatoria	159,275	87,282	54,234	17,759	37.20	80,411	2,881	9,052	18,064	48,867

338 **Table 2. Gene content in *Euterpe edulis*, *E. oleracea* and *E. precatoria* chloroplast**

339 **genomes according to each respective category.**

Category	Gene
Subunits of photosystem I	*psaA, psaB, psaC, psaI, psaJ*
Subunits of photosystem II	*psbA, psbB, psbC, psbD, psbE, psbF, psbH, psbI, psbJ, psbK, psbL, psbM, psbN, psbT, psbZ*
Subunits of cytochrome b/f complex	*petA, petB[a], petD[a], petG, petL, petN*
Subunits of ATP synthase	*atpA, atpB, atpE, atpF[a], atpH, atpI*
Large subunit of rubisco	*rbcL*
Subunits of NADH-dehydrogenase	*ndhA[a], ndhB[a,b], ndhC, ndhD, ndhE, ndhF, ndhG, ndhH, ndhI, ndhJ, ndhK*
Proteins of large ribosomal subunit	*rpl2[a,b], rpl14, rpl16[a], rpl20, rpl22, rpl23[b], rpl32, rpl33, rpl36*
Proteins of small ribosomal subunit	*rps2, rps3, rps4, rps7[b], rps8, rps11, rps12[a,b,c], rps14, rps15, rps16[a], rps18, rps19[b]*
Subunits of RNA polymerase	*rpoA, rpoB, rpoC1[a], rpoC2*
Maturase	*matK*
Translational initiation factor	*infA*
Protease	*clpP[a]*
Envelope membrane protein	*cemA*
Subunit of acetyl-CoA carboxylase	*accD*
Cytochrome c biogenesis	*ccsA*
Conserved hypothetical genes	*ycf3[a], ycf4*
Component of TIC complex	*ycf1*
Component of 2-MD heteromeric AAAATPase complex	*ycf2[b]*
Ribosomal RNAs	*rrn4.5[b], rrn5[b], rrn16[b], rrn23[b]*
Transfer RNAs	*trnA –UGC[a,b]; trnC –GCA; trnD –GUC; trnE –UUC; trnF –GAA; trnfM –CAU; trnG –UCC[a] trnG –GCC; trnH –GUG[b]; trnI –CAU[b]; trnI –GAU[a,b]; trnK –UUU[a]; trnL –CAA[b]; trnL –UAA[a]; trnL –UAG; trnM –CAU; trnN –GUU[b]; trnP –UGG; trnQ –UUG; trnR –ACG[b]; trnR–UCU; trnS –GCU; trnS –UGA; trnS –GGA; trnT –UGU; trnT –GGU; trnV –GAC[b]; trnV–UAC[a]; trnW –CCA; trnY –GUA*

[a]Intron-containing gene; [b]Two gene copies in the Irs; [c]Gene divided into two independent transcription units.

341 **Fig 1. Gene map of the *Euterpe edulis*, *E. oleracea* and *E. precatoria* chloroplast**

342 **genomes**. Genes represented inside the large circle are oriented clockwise and the ones

outside are oriented counter clockwise. The different colors represent functional groups, and the darker gray in the inner circle indicates the GC content. The quadripartite structure is also reported as: LSC = Large Single Copy, SSC = Small Single Copy, IRA/IRB = Inverted Repeats A and B.

Considering the duplicated genes in the IRs, more than 50% of the three chloroplast genome sequences are from protein coding regions (86 genes, Table 1 and Table 2). Among genes, *ycf2* presented a slight difference between species, 6,903 bp in *Euterpe edulis* and 6,879 bp in *E. oleracea* and *E. precatoria*. Furthermore, chloroplast tRNA and rRNA were conserved among species, constituting 1.8% and 5.7% of their sequences, respectively. Introns constituted ca. 12% and intergenic regions represented almost 31% of the chloroplast genomes (Table 1).

All chloroplast genomes had 17 unique genes (11 protein- and six tRNA-coding genes) with introns, five duplicated genes in the inverted repeats regions with introns, and two introns in the *ycf3* and *clpP* genes. Accounting for the lengths, the largest intron was identified in *trnK – UUU* (2,621 bp – *E. edulis*; 2,615 bp – *E. oleracea*; 2,618 bp – *E. precatoria*) and the smallest in *trnL-UAA* (519 bp – *E. edulis*, 519 bp – *E. oleracea*, 521 bp – *E. precatoria*).

Chloroplast genome structures and comparative analyses

The comparative analysis enabled us to identify a high level of synteny between the three *Euterpe* chloroplast genomes, with large conserved blocks. The only structural difference among *Euterpe* chloroplast genomes were three small inversions and single nucleotide polymorphisms (SNPs) between *E. edulis* and *E. oleracea* chloroplast sequences (S2A, B and C Figs). In a local alignment we detected these divergences in the region between 66,081 to 69,440 bp; the first corresponding to an intergenic region

between *petA* and *psbJ* genes from *E. oleracea*; the second found in the *psbJ* gene and the third in the intergenic region of *trnW-CCA* and *trnP-UGG* from *E. oleracea* (S2D Fig).

The parallel between the chloroplast genomes of *Euterpe edulis* and *Euterpe oleracea* with the ones available in Genbank [34] showed differences in the total number of base pairs. *E. edulis* and *E. oleracea* were 835 bp and 458 bp, respectively, larger than the others. The alignment revealed that these increases were due to insertions that occurred mainly in intergenic spacers (83.3% of insertions in *E. edulis* and 100% in *E. oleracea*; S2 Table).

Although exhibiting high synteny and only a few structural rearrangements in their chloroplast genome, the most notable divergence among the three Brazilian native species from subfamily Arecoideae was in the length of the LSC, between 40,000 and 50,000 bp (Fig 2). A high level of synteny was also observed in a comparison of palm chloroplast genomes from 12 genera, with five subfamilies (S3 Fig). *Astrocaryum aculeatum*, *A. murumuru* and *Syagrus coronata* are native palms from subfamily Arecoideae with smaller LSCs compared with *Euterpe* palms (Fig 2 and S3 Fig). *Acrocomia aculeata,* also a Brazilian palm, presented a reduction in the size of the region comprising the genes *ndhJ*, *ndhK*, *ndhC*, *trnF-GAA* and *trnL-UAA* (in lime green Fig 2). Additionally, *Astrocaryum aculeatum* and *A. murumuru* showed a flip-flop recombination in this same region (in lime green, Fig 2) [67].

Fig 2. Synteny and divergence in the SSC size detected in Arecaceae chloroplast genomes using the Mauve multiple-genome alignment program. A sample of nine different chloroplast genomes is shown. Color bars indicate syntenic blocks and the lines indicate the correspondence between them. Blocks on the top row are in the same orientation, while blocks on the bottom row are in inverse orientation.

In addition, we explored in more details the chloroplast IR expansions and

contractions of the Brazilian palm species from the subfamily Arecoideae (Fig 3).

Comparing the IR borders from Genbank available for *Euterpe edulis* chloroplast genome,

it was possible to observe a small difference in the *rpl22-rps19* location and an increase

in the length of *rps19-psbA* intergenic spacers. Among the chloroplast genomes from

Euterpe genus, we identified an expansion in the *ndhF* gene at the IRB region from

Euterpe precatoria. Among the Brazilian palm species, a contraction in the IR from

Syagrus coronata influenced the position of *rps19*. Among all species analyzed, the most

common variations were identified around the positions of *rpl22-rps19* (IRB-SSC), the

ycf1 length (IRB and IRA), and the *rps19-ycf1* (IRA-LSC) (Fig 3).

Fig 3. Comparison of the IRA and IRB borders among Brazilian palms from the

Arecoideae species. The numbers indicate the lengths of IGSs, genes, and spacers

between IR-LSC and IR-SSC junctions. The *ycf1** and *rps19** genes have incomplete

CDSs.

Sequence repeats and polymorphisms between *Euterpe* chloroplast genomes

We identified a total of 323 SSRs (*E. edulis*: 111, *E. oleracea*: 105, *E. precatoria*:

107 SSRs) in the chloroplast genomes from the three species. Most of them were located

in intergenic spacer regions (IGS: 72.07% *E. edulis*, 72.38% *E. oleracea*, 71.96 % *E.*

precatoria; Fig 4A), especially in the IGS between the tRNAs trnS-GCU/trnG-UCC (S3

Table). The SSRs were more abundant in the LSC region of the chloroplast genomes

(78.38% *E. edulis*, 78.10% *E. oleracea*, 77.57% *E. precatoria*; S3 Table) and less frequent

in the IR regions. They were mostly mononucleotides (62.16% *E. edulis*, 62.86% *E.*

oleracea, 60.75% *E. precatoria*; Fig 4B) and composed with A/T motifs (60.36% *E.*

edulis, 60.00% *E. oleracea*, 59.81% *E. precatoria*; S4A Fig). From the 323 SSRs, 47 had

the same sequence and gene location among *E. edulis*, *E. oleracea* and *E. precatoria*. We

also identified 62 SSRs between *E. edulis/E. oleracea* and *E. edulis/E. precatoria,* and 68

between *E. oleracea and E. precatoria* (S4 Table).

Considering the dispersed repeats, we observed 66 in *E. edulis*, 55 in *E. oleracea*

and 49 in *E. precatoria*, ranging from 30 to 77 bp (S4B Fig). These repeats were

distributed mostly in the IGS regions (65.15% *E. edulis*, 70.91% *E. oleracea*, 73.47% *E.*

precatoria; Fig 4C). Comparing the species, in *E. precatoria*, most of the repeats were

found in the intergenic region of *psaC/ndhE*. However, in *E. edulis* and *E. oleracea* the

repeats were concentrated in the *ycf2* gene (S5 Table). Regarding repeat types, most of

the repeats are palindrome or forward sequences, 40% and 30%, respectively (Fig 4D).

They were also mainly found in the LSC region (48.48% *E. edulis*, 52.73% *E. oleracea*,

44.90% *E. precatoria*; S5 Table).

Fig 4. Distribution and classification of SSR and dispersed repeats in the chloroplast

genomes of *Euterpe edulis, E. oleracea* and *E. precatoria*. (A) Proportion of coding and

non-coding regions containing SSRs; (B) Proportion of different types of SSR present in

the chloroplast genomes; (C) Proportion of regions containing repeats; (D) Frequency

distribution of different types of repeats: F = Forward, P = Palindrome, R = Reverse and

C = Complement. CDS = Coding sequence, IGS = Intergenic spacer.

Based on the pairwise alignment between the species chloroplast genomes, we

identified 93 indels between *E. edulis/E. oleracea,* 103 between *E. edulis/E. precatoria*

and 58 between *E. oleracea/E. precatoria*. Most indels were distributed in the IGS (Fig

5A), especially between the *E. oleracea/E. precatoria* chloroplast genome, located

mainly in the LSC region (80.65% ~ 72.41%; S6 Table). The highest occurrence of indels

was identified in the *trnS-GCU/trnG-UCC* of the three alignments (S6 Table).

A total of 325 SNPs were detected between the three species (Fig 5B). In a

pairwise comparison, 283 SNPs were detected between *E. edulis/E. oleracea* and 254

SNPs between *E. edulis/E. precatoria*, which were most frequent in the CDS regions

(50.35% and 54.72%, respectively; Fig 5C). Between *E. oleracea/E. precatoria*, a lower

number of SNPs was found (113 SNPs), having the greatest number concentrated in the

IGS (50.44%; Fig 5C). Only one SNP was found shared among the three species (Fig

5B).

Fig 5. Indels and single nucleotide polymorphisms (SNPs) detected in comparisons

between *Euterpe* chloroplast genomes. (A) Proportion of indels in different coding and

non-coding regions; (B) Comparison of the number of SNPs found in the alignment (C)

Proportion of SNPs in different coding and non-coding regions of the chloroplast

genomes.

The large number of SNPs in the CDS region occurs mainly in the *atpE*, *ycf1* and

psbJ genes (S7 Table). These *atpE*, *psbJ* and *ycf1* genes presented respectively 19, 18 and

18 SNPs for *E. edulis/E. oleracea* and 20, 19 and 21 SNPs for *E. edulis/E. precatoria*.

For *E. oleracea/E. precatoria* there were only 1 (*atpE*) and 7 (*ycf1*) SNPs (S7 Table).

The LSC region had the highest number of SNPs in the chloroplast genomes

(69.01% *E. edulis/E. oleracea*, 66.54% *E. edulis/E. oleracea*, 64.91% *E. oleracea/E.*

precatoria; S7 Table). For *E. edulis/E. oleracea* the greatest number of SNPs

corresponded to C/T - T/C (74) substitutions, the same was saw for *E. edulis/E. precatoria*

T/C - C/T (66), and 29 substitutions of G/A - A/G and C/T - T/C was observed for *E.*

oleracea/E. precatoria (S5 Fig).

Considering pi > 0.02, the sliding window analysis revealed six hotspots of high

nucleotide polymorphism among the *E. edulis*, *E. oleracea* and *E. precatoria* chloroplast

genome sequence. The six hotspots were in IGS regions (Fig 6). Among them, the two

hotspots with the highest polymorphism were located between the *trnM*/*atpE* (Pi > 0.06;

Fig 6) and *psbJ*/*psbL* (Pi > 0.06; Fig 6). Comparing with the SNP identification, the *atpE*

gene had the highest number of SNPs among the species *E. edulis/E. oleracea* and *E.*

edulis/E. precatoria. The *psbJ* gene had 18 SNPs in *E. edulis/E. oleracea* and 19 SNPs

in *E. edulis/E. precatoria*.

Fig 6. Sliding window analysis of the alignment from the chloroplast genome of

***Euterpe edulis*, *E. oleracea* and *E. precatoria* chloroplast genomes**. The regions with

high nucleotide variability (Pi > 0.02) are indicated. Pi is the nucleotide diversity of each

window, and the window length was 200 bp with 50 bp step sizes.

Both genes, *atpE* and *psbJ*, were analyzed for positive or purifying selection

across the palm phylogeny using the estimation of Ka/Ks ratio. In both cases, the Ka/Ks

< 1 revealed that no positively selected sites were found in the corresponding protein

sequences, indicating purifying selection. The analysis of the amino acid sequence

resulted in Ka/Ks rates varying from 0.17 to 0.87 for *atpE* (S8 Table) and ranging from

0.50 to 0.96 for *psbJ*. Also, from the 134 amino acids of the *atpE* protein, 68 had Ka/Ks

< 0.19 and from the 40 amino acids, 37 had Ka/Ks < 0.55, in the case of *psbJ* (S9 Table).

Phylogenomic of *Euterpe* based on chloroplast sequences

In the phylogenomic analysis, the marginal mean likelihoods of partition schemes

for the two runs were i) -186887.39, ii) -178507.00, and iii) -178497.59. Therefore, the

best partition schemes were by far the ones considering codon positions, although 'iii'

was strongly better than 'ii' with the 2*log difference = 18,82 (>10 is strong in Kass and

Raftery 1995). We also compared the majority-rule consensus trees of the three partition

scheme analyses, which did not differ in topology, but in general the scheme 'iii'

presented better support in some nodes. For that reason, the figure presented and used in

our discussion will be based on partition scheme 'iii'.

The phylogeny using coding sequences from 54 whole chloroplast genomes from

palm species rooted in the outgroup species *Dasypogon bromeliifolius* revealed the same

relationships in subfamilies previously reported [34], with highly supported nodes (Fig

7). According to our results, the species from *Euterpe* were placed in subfamily

Arecoideae, tribe Euterpeae, and were sister to tribe Areceae. The new samples of *E.*

oleracea and *E. edulis* were grouped with the respective previously published chloroplast

genomes and the *E. precatoria* was placed in a branch sister to *E. oleracea*.

Fig 7. Majority-rule consensus tree of 30,000 trees obtained from a Bayesian

inference analysis of chloroplast protein coding genes of 55 taxa. Posterior

probabilities (PP) for each are indicated above branches. Co = Coryphoideae, Ar =

Arecoideae, Ny = Nypoideae, Ce = Ceroxyloideae, Ca = Calamoideae.

Species divergence with nuclear and mitochondrial genome

From the 11,738,792 reads that were generated for *E. edulis*, 553,857 (4.71%)

aligned to the assembled chloroplast genome. The remaining (ca. 95%) reads correspond

to nuclear and mitochondrial regions. Similar proportions were also found for *E. oleracea*

and *E. precatoria* for which 12,987,827 and 12,612,001 reads were generated,

respectively. Moreover, 249,387 (1.92%) and 132,227 (1.04%) were aligned to the

assembled chloroplast sequence for both species, respectively. Using Stacks, we found

1,077 SNPs in the nuclear and mitochondrial genome sequences, with 10X mean

sequencing depth per locus (S6A Fig). Among *Euterpe* species, more transitions than

transversions were detected, and the most frequent mutations were A-G and C-T

transitions (S6C Fig).

516   The highest number of alleles (A = 1,309) and highest observed heterozygosity

517 (H_O = 0.162) was identified in *E. oleracea*, while the highest number of private alleles

518 was observed in *E. precatoria* (Ap = 421). All the three species had a higher value of H_O

519 compared with the expected heterozygosity (H_E, Table 3).

520 **Table 3. Parameters of genetic diversity using 1,077 SNPs found the nuclear and**

521 **mitochondrial genome sequences from three *Euterpe* species.**

	E. edulis	*E. oleracea*	*E. precatoria*
A	1,251	1,309	1,263
Ap	292	260	421
H_O	0.162 (±0.011)	0.215 (± 0.013)	0.173 (±0.012)
H_E	0.081 (±0.006)	0.108 (±0.006)	0.086 (±0.006)

522 A = number of alleles, Ap = number of private alleles, H_O = observed heterozygosity, H_E

523 = expected heterozygosity.

524   The pairwise F_{ST} showed high and significant divergence between species, with

525 values higher than 0.770. Although all the F_{ST} values were high, the species *E. edulis* and

526 *E. precatoria* presented the greatest genetic divergence (0.860, S7 Fig) among the three

527 pairs of species, and *E. edulis* and *E. oleracea* had the lowest divergence (0.779, S7 Fig).

Discussion

High level of conservation between Brazilian palms

chloroplast genomes

532   Among the chloroplast genomes from the *Euterpe* genus, we found a conserved

533 organization and gene content, with small differences between species, as in the length of

534 the duplicated *ycf2* gene. The *ycf2* is the largest plastid gene reported in angiosperms [68],

535 and is part of the 2-MD heteromeric AAA-ATPase complex directly interacting with

536 various translocating preproteins [69]. Particularly, *ycf2* has an insertion in the *E. edulis*

537 chloroplast genome that is absent in *E. oleracea* and *E. precatoria*, a feature previously

observed in *Euterpe* chloroplast genomes [34]. Across palm species, the chloroplast

genome structure is highly conserved as reported in other studies [28,34,70]. The multiple

alignments with 15 different palms from five subfamilies did not exhibit significant

rearrangements in the chloroplast structure. The only rearrangement observed was an

inversion of 4.6 kb in the *Astrocaryum* chloroplast genome. It was possibly to have a

lineage-specific structural variation, creating a isoform from a flip-flop recombination

between inverted repeats [67]. However, differences in the total size of chloroplast

genomes were observed in the same species, such as in *E. edulis* and *E. oleracea*, which

may be due to the assembly method used. Different from the approach of all possible

paths' calculation to complete target organelle genome, as applied by GetOrganelle, other

widely used pipelines utilize reference genomes to select and filter scaffolds/contigs for

further concatenation or post-assembly gap filling and closing [38].

Besides these conserved general features, similar occurrences of SSRs were

identified among the three *Euterpe* species. Currently, SSR markers are very useful in

studies of population structure, genetic mapping, and evolutionary processes [71]. The

SSR identified here can be a valuable resource for studies aiming at the conservation of

the species or their sustainable exploitation. These SSR are mostly composed of A/T

motifs, and are more frequently located in the IGS of the single-copy regions from the

chloroplast genomes [1,72,73], which is expected since these regions evolve faster than

the CDS [74]. Also, the limited occurrence of SSRs in the IRs is related to a lower

mutation rate observed in these regions, caused by an efficient mechanism of gene copy-

correction, as observed in other palm chloroplast genomes [34,75].

The dispersed repeats (forward, palindrome, reverse and complement) showed the

same pattern of the SSRs and were frequently distributed in the IGS of the single copy

regions, differing from a predominant occurrence of reverse repeats (18,37%) in *E.*

precatoria. Actually, the number of repeats found in the *Euterpe* chloroplast genomes

were not considered high [76]. According to Milligan et al. [77], repetitive sequences are

substrates for recombination and chloroplast genome rearrangements. Since non-

structural rearrangements were detected in the *Euterpe* chloroplast genomes, low

frequency of repetitive elements was also expected.

568   Regarding the structure patterns of the IRs, we could identify most of the variation

related to expansion/contractions when comparing only Brazilian palm species from

subfamily Arecoideae. Events of expansion and contraction at the IR boundaries was

previously observed in the *ycf1* gene at IR-SSC/LSC junction and *rps19* intergenic spacer

with *rpl22*/*psbA* in IR-SSC/LSC junctions of other species [28,75].

Indels and single nucleotide polymorphisms clearly

demonstrate differentiation among the three *Euterpe* species,

which was reflected in the phylogeny

577   We identified six hotspots of high nucleotide polymorphism with the largest

polymorphism (Pi > 0.06) in intergenic regions composed mainly by genes with a great

number of SNPs. The genes *atpE* and *psbJ* had the greatest number of SNPs, especially

when considering the nucleotide substitutions between *E. edulis* and the *Euterpe*

Amazonian species, *E. oleracea* and *E. precatoria*. This divergence between *E. edulis* in

relation to the two Amazonian species was also observed in the number of indels and

plastid SNPs and was almost 50% higher than the number of polymorphisms detected in

a pairwise analysis with *E. oleracea* and *E. precatoria.*

585   Indels display no ambiguity in complex mutation patterns and SNPs are the most

abundant type of markers [78], giving us a clear pattern of divergence between *E. edulis*

in relation to *E. oleracea* and *E. precatoria*. In our study, these mutations were frequently

found in the CDS than in the intergenic spacers. This high number of indels and SNPs

between *E. edulis/E. oleracea* had already been identified in a previous study [34], but

they were mainly found in non-coding regions, which presumably may be related to a

more recent event of divergence.

Regarding the phylogenetic relationships among the palm species, we identified

a clear distinction among subfamilies using the informative chloroplast coding sequences,

corroborating previous studies [28,34,67,79]. Our phylogeny suggests that *E. oleracea*

and *E. precatoria* are sister species with an early split of *E. edulis* from these two,

differently from what was observed previously, having four nuclear low copy markers

and one plastid region [33]. Our result reveals the possibility of incongruences between

nuclear and plastid phylogenies in *Euterpe*, since the previous study using nuclear

markers reported paraphyly in *E. precatoria* and showed one of the *E. precatoria* varieties

as sister to *E. edulis* [33].

The closest relationship between *E. oleracea* and *E. precatoria* could reflect their

environmental occupation since both species occur in the Brazilian Amazon. *E. edulis*,

nonetheless, evolved via vicariant split from its common ancestor (around 1.4 Mya), and

the speciation was intensified by the savanna barrier composed from the Cerrado and

Caatinga biomes [33,34,80]. Furthermore, the phenotypic plasticity and local adaptation

may have favored the successful expansion of *E. edulis* throughout the Atlantic Forest

[11,80].

Regarding the coding sequences, the *rpl22, rpl32, rpl14, rps14, matK, accD, rbcl,*

ccsA, ycf2 and ycf1 genes have been reported to harbor positive selection in an analysis

using 41 species of the Arecaceae family, including the *E. oleracea* and *E. edulis*

chloroplast genomes [34]. This was highlighted as indicative of convergent evolution,

associated with environmental adaptation in *E. edulis*. However, using 52 palm species,

we could only identify a tendency of purifying selection for *psbJ* and *atpE*, the genes with

higher SNPs and polymorphisms hotspots. This may reflect the typically conservative

nature of the chloroplast genome sequences across most angiosperms [81,82]. These

genes are essential for plant development, since *atpE* encodes a subunit of the ATP

synthase complex that participates in the photosynthetic phosphorylation [83] and *psbJ* is

associated with photosystem II (PSII) [84].

620 A comparison using nuclear and mitochondrial genome
621 sequences detected high divergence between *Euterpe* species

The reads from nuclear and mitochondrial DNA obtained from the sequencing

were used in this first-time comparison of the genetic diversity and genomic divergence

among the three species from the *Euterpe* genus. The advances in next-generation

sequencing (NGS) is allowing the discovery of a large number of single nucleotide

polymorphism (SNPs), even for non-model species [85–88]. This generates a valuable

source of bi-allelic genetic markers that are appropriate to evaluate the genetic diversity

even when sample sizes are small [89]. Additionally, SNPs occurs at higher density across

the genome, with lower genotyping error rates than other markers [90], enabling robust

estimates of genetic diversity and structuring [9].

The three *Euterpe* species had similar number of alleles and low observed

heterozygosity. Yet, they are higher than the expected heterozygosity, with *E. oleracea*

presenting slightly higher results. However, despite this enlightening information

obtained with the remaining nuclear and mitochondrial sequences, we highlight that these

findings could be biased because they were based on only one individual per species.

Considering assessments of genome-wide diversity, previous studies identifying SNPs in

palms were only performed for *E. edulis* [9,80]. Furthermore, the genetic diversity

estimates from our *E. edulis* sample, that was also from the Atlantic rainforest of São

Paulo state, were in agreement with previously published studies. This observed genetic

variability could be influenced by the type of vegetation and the altitude where the

samples were collected [9,80].

In summary, our results open the way for future genome-skimming studies in

palms. Also, the addition of larger sample sizes from different sampling locations will

provide a better understanding on the evolution and diversification of this important group

of plants. However, the high differentiation found between the three *Euterpe* species may

indicate that they are evolving independently and that this differentiation is caused by a

private pool of alleles from each species, which is in accordance with the phylogenetic

analyses.

Acknowledgements

The authors would like to thank Dr. Charles R. Clement for sending samples of *Euterpe*

precatoria from Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Brazil

for the genomic extraction and for reading the manuscript.

Supporting information

S1 Fig. Flowchart of the SNPs identification methodology using nuclear and

mitochondrial genomes sequences.

(TIFF)

S2 Fig. Dot-plot analyses comparing three *Euterpe* chloroplast genomes. (A) *Euterpe*

edulis (X – axis) x *Euterpe Oleracea* (Y-axis), (B) *E. edulis* (X-axis) x *Euterpe precatoria*

(Y-axis), (C) *E. oleracea* (X – axis) x *E. precatoria* (Y-axis). The positive slope, in purple,

represents the pair of sequences aligned and in the same orientation. The negative slope,

in blue, represents the pair of sequence aligned, but in opposite orientation. The blue

arrow in A highlights the region with inversions and SNPs between *E. edulis* and *E.*

oleracea. (D) Local alignment with the chloroplast genomes of *E. edulis*, *E. oleracea* and

E. precatoria in the region where inversions and SNPs were detected in A.

(TIFF)

S3 Fig. Synteny and divergence in the size of the SSC detected in Arecaceae

chloroplast genome sequences. A sample of 20 different chloroplast genomes is shown.

Color bars indicate syntenic blocks and the lines indicate the correspondence between

them. Blocks on the top row are in the same orientation, while blocks on the bottom row

are in inverse orientation.

(TIFF)

Fig S4. SSR and dispersed repeats in the chloroplast genomes of *Euterpe edulis, E.*

oleracea* and *E. precatoria. (A) The number of SSR motifs found in the three

chloroplast genomes, considering sequence complementarities; (B) Number of

dispersed repeats present in different size classes.

(TIFF)

S5 Fig. Number and type of SNPs found in the alignment of three *Euterpe*

chloroplast genomes

(TIFF)

S6 Fig. Description of 1,077 SNPs found among the nuclear and mitochondrial

genome sequences from three *Euterpe* species. A) Mean depth per loci; B) Mean

depth per species sequenced; C) Mutation count in the present in the 1,077 SNPS.

(TIFF)

S7 Fig. Pairwise F_{ST} based on 1,077 SNPs of nuclear and mitochondrial genomes

sequences from three *Euterpe* specie

(TIFF)

S1 Table. List of complete chloroplast genomes used in comparative analysis.

Synteny and rearrangement, border comparison and phylogenomics.

(XSLX)

S2 Table. Insertions identified in the alignment of *Euterpe edulis* and *E. oleracea*

chloroplast genomes with sequences from GenBank (de Santana Lopes et al. 2021).

CDS = Coding sequence; IGS = intergenic spacer; LSC = Large single copy; SSC =

Small single copy.

(XSLX)

S3 Table. Simple sequence repeats (SSR) identified in the complete chloroplast

genomes of three *Euterpe* species. CDS = Coding sequence; IGS = intergenic spacer;

LSC = Large single copy; SSC = Small single copy; IRs = Inverted Repeats.

(XSLX)

S4 Table. Simple sequence repeats (SSR) among the chloroplast genomes of

Euterpe edulis, *E. oleracea* and *E. precatoria* **according to their location.**

(XSLX)

S5 Table. Repeats identified in the chloroplast genomes of *Euterpe edulis, E.*

oleracea **and** *E. precatoria*. F = Forward; P = Palindrome; R = Reverse; C =

Complement; CDS= Coding sequence; IGS = intergenic spacer; LSC = Large single

copy; SSC = Small single copy; IRs = Inverted Repeats.

(XSLX)

S6 Table. Indels identified in the pairwise comparison of three *Euterpe* **species**

complete chloroplast genomes. CDS = Coding sequence; IGS = intergenic spacer;

LSC = Large single copy; SSC = Small single copy; IRs = Inverted Repeats.

(XSLX)

S7 Table. Single nucleotide polymorphisms (SNPs) identified in the pairwise

comparison of three *Euterpe* **species complete chloroplast genomes.** CDS = Coding

sequence; IGS = intergenic spacer; LSC = Large single copy; SSC = Small single copy;

IRs = Inverted Repeats.

(XSLX)

S8 Table. Bayesian selection results based on synonymous (Ka)/non-synonymous

(Ks) ratio of amino acid substitutions from the *atpE* **gene in 54 palm species**.

(XSLX)

S9 Table. Bayesian selection results based on synonymous (Ka)/non-synonymous

(Ks) ratio of amino acid substitutions from the *psbJ* gene in 54 palm species.

(XSLX)

739
740

Author Contributions

Conceptualization: Maria Imaculada Zucchi

Data curation: Ana Flávia Francisconi, Jonathan Andre Morales Marroquín, Cássio

van den Berg, Luciano Delmondes de Alencar, Alessandro Alves-Pereira

Formal Analysis: Ana Flávia Francisconi, Luiz Augusto Cauz dos Santos, Cássio van

den Berg, Alessandro Alves-Pereira, Luciano Delmondes de Alencar

Funding Acquisition: Maria Imaculada Zucchi

Investigation: Ana Flávia Francisconi, Luiz Augusto Cauz dos Santos

Methodology: Luiz Augusto Cauz dos Santos, Jonathan Andre Morales Marroquín,

Doriane Picanço-Rodrigues, Luciano Delmondes de Alencar, Cesar Augusto

Zanello, Marcones Ferreira Costa

Resources: Maria Imaculada Zucchi, Doriane Picanço-Rodrigues

Software: Cássio van den Berg

Supervision: Maria Imaculada Zucchi

Writing – Original Draft Preparation: Ana Flávia Francisconi, Luiz Augusto Cauz

dos Santos

Writing – Review & Editing: Ana Flávia Francisconi, Luiz Augusto Cauz dos Santos,

Jonathan Andre Morales Marroquín, Cássio van den Berg, Alessandro Alves-

Pereira, Doriane Picanço-Rodrigues, Luciano Delmondes de Alencar, Cesar

Augusto Zanello, Marcones Ferreira Costa, Maria Teresa Gomes Lopes, Elizabeth

Ann Veasey, Maria Imaculada Zucchi

References

1. Uthaipaisanwong P, Chanprasert J, Shearman JR, Sangsrakru D, Yoocha T, Jomchai N, et al. Characterization of the chloroplast genome sequence of oil palm (*Elaeis guineensi*s Jacq.). Gene. 2012;500: 172–180. doi:10.1016/j.gene.2012.03.061

2. Baker WJ, Couvreur TLP. Global biogeography and diversification of palms sheds light on the evolution of tropical lineages. I. Historical biogeography. J Biogeogr. 2013;40: 274–285. doi:10.1111/j.1365-2699.2012.02795.x

3. Brokamp G, Valderrama N, Mittelbach M, Grandez CA, Barfod AS, Weigend M. Trade in palm products in North-Western South America. Bot Rev. 2011;77: 571–606. doi:10.1007/s

4. Eiserhardt WL, Pintaud J, Asmussen-lange C, Hahn J, Bernal R, Balslev H, et al. Phytogeny and divergence times of Bactridinae (Arecaceae, Palmae) based on plastid and nuclear DNA sequences. 2011;60: 485–498.

5. Baker WJ, Couvreur TLP. Global biogeography and diversification of palms sheds light on the evolution of tropical lineages. II. Diversification history and origin of regional assemblages. J Biogeogr. 2013;40: 286–298. doi:10.1111/j.1365-2699.2012.02794.x

6. Henderson A, Galeano G. Euterpe, Prestoea, and Neonicholsonia (Palmae). Flora Neotrop. 1996;72: 1–89.

7. Mart E, Bussmann RW, Zambrana NYP. Acta Societatis Botanicorum Poloniae Facing global markets – usage changes in Western Amazonian plants: the example of *Euterpe precatoria* Mart. and *E. oleracea* Mart. 2012;81: 257–261. doi:10.5586/asbp.2012.032

8. Galetti M, Guevara R, Côrtes MC, Fadini R, Matter S Von, Leite AB, et al. Functional extinction of birds drives rapid evolutionary changes in seed size. Science (80-). 2013;340: 1086–1091. doi:DOI: 10.1126/science.1233774

9. Alves-Pereira A, Novello M, Dequigiovanni G, Pinheiro JB, Brancalion PHS, Veasey EA, et al. Genomic diversity of three Brazilian native food crops based on double-digest restriction site-associated DNA sequencing. Trop Plant Biol. 2019;12: 268–281. doi:10.1007/s12042-019-09229-z

10. Novello M, Viana JPG, Alves-Pereira A, de Aguiar Silvestre E, Nunes HF, Pinheiro JB, et al. Genetic conservation of a threatened Neotropical palm through community-management of fruits in agroforests and second-growth forests. For Ecol Manage. 2017;407: 200–209. doi:10.1016/j.foreco.2017.06.059

11. Coelho GM, Santos AS, de Menezes IPP, Tarazi R, Souza FMO, Silva M das GCPC, et al. Genetic structure among morphotypes of the endangered Brazilian palm *Euterpe edulis* Mart (Arecaceae). Ecol Evol. 2020;10: 6039–6048. doi:10.1002/ece3.6348

12. Martinelli G, Moraes MA. Livro vermelho da flora do Brasil. 1st ed. Rio de Janeiro: CNCFlora, Centro Nacional de Conservação da Flora Rio de Janeiro; 2013.

13. Ball AA, Brancalion PHS. Governance challenges for commercial exploitation of a non-timber forest product by marginalized rural communities. Environ Conserv. 2016;43: 208–220. doi:10.1017/S0376892916000072

14. IBGE. Censo agropecuário 2017: resultados definitivos. Censo agropecuário. 2019;8: 1–105. Available from: https://biblioteca.ibge.gov.br/visualizacao/periodicos/3096/agro_2017_resultados _definitivos.pdf

812   15.   Ramos SLF, Dequigiovanni G, Lopes MTG, Aguiar AV de, Lopes R, Veasey
813   EA, et al. Genetic structure in populations of *Euterpe precatoria* Mart. in the
814   Brazilian Amazon. Front Ecol Evol. 2021;8: 1–11.
815   doi:10.3389/fevo.2020.603448
816   16.   Shanley P, Medina G. Frutíferas e plantas úteis na vida amazônica. 1st ed.
817   Belém-PA: Cifor, Imazon; 2005.
818   17.   Vedel-sørensen M, Tovaranonte J, Bøcher PK, Balslev H, Barfod AS. Spatial
819   distribution and environmental preferences of 10 economically important forest
820   palms in western South America. For Ecol Manage. 2013;307: 284–292.
821   doi:10.1016/j.foreco.2013.07.005
822   18.   Freitas MAB, Vieira ICG, Albernaz ALKM, Magalhães JLL, Lees AC. Floristic
823   impoverishment of Amazonian floodplain forests managed for açaí fruit
824   production. For Ecol Manage. 2015;351: 20–27.
825   doi:10.1016/j.foreco.2015.05.008
826   19.   Campbell AJ, Carvalheiro LG, Maués MM, Jaffé R, Giannini TC, Freitas MAB,
827   et al. Anthropogenic disturbance of tropical forests threatens pollination services
828   to açaí palm in the Amazon river delta. J Appl Ecol. 2018; 1725–1736.
829   doi:10.1111/1365-2664.13086
830   20.   Kahn F. Palms as key swamp forest resources in Amazonia. For Ecol Manage.
831   1991;38: 133–142. doi:https://doi.org/10.1016/0378-1127(91)90139-M
832   21.   Ramos SLF, Dequigiovanni G, Magno A, Pimentel P, Teresa M, Lopes G, et al.
833   Paternity analysis, pollen flow, and spatial genetic structure of a natural
834   population of *Euterpe precatoria* in the Brazilian Amazon. Ecol Evol. 2018;8:
835   11143–11157. doi:10.1002/ece3.4582
836   22.   Du FK, Lang T, Lu S, Wang Y, Li J, Yin K. An improved method for chloroplast
837   genome sequencing in non-model forest tree species. Tree Genet Genomes.
838   2015;11. doi:10.1007/s11295-015-0942-2
839   23.   Dodsworth S. Genome skimming for next-generation biodiversity analysis.
840   Trends Plant Sci. 2015;20: 525–527. doi:10.1016/j.tplants.2015.06.012
841   24.   Sobreiro MB, Vieira LD, Nunes R, Novaes E, Coissac E, Silva-Junior OB, et al.
842   Chloroplast genome assembly of *Handroanthus impetiginosus*: Comparative
843   analysis and molecular evolution in Bignoniaceae. Planta. 2020;252: 1–16.
844   doi:10.1007/s00425-020-03498-9
845   25.   Sugiura M. The chloroplast genome. Plant Mol Biol. 1992;19: 149–168.
846   doi:10.1007/978-94-011-2656-4_10
847   26.   Barrett CF, Bacon CD, Antonelli A, Cano Á, Hofmann T. An introduction to
848   plant phylogenomics with a focus on palms. Bot J Linn Soc. 2016;182: 234–255.
849   doi:10.1111/boj.12399
850   27.   Dabral A, Shamoon A, Meena RK, Kant R, Pandey S, Ginwal HS, et al. Genome
851   skimming-based simple sequence repeat (SSR) marker discovery and
852   characterization in *Grevillea robusta*. Physiol Mol Biol Plants. 2021;27: 1623–
853   1638. doi:10.1007/s12298-021-01035-w
854   28.   de Santana Lopes A, Gomes Pacheco T, Nimz T, do Nascimento Vieira L,
855   Guerra MP, Nodari RO, et al. The complete plastome of macaw palm
856   [*Acrocomia aculeata* (Jacq.) Lodd. ex Mart.] and extensive molecular analyses of
857   the evolution of plastid genes in Arecaceae. Planta. 2018;247: 1011–1030.
858   doi:10.1007/s00425-018-2841-x
859   29.   Aljohi HA, Liu W, Lin Q, Zhao Y, Zeng J, Alamer A, et al. Complete sequence
860   and analysis of coconut palm (*Cocos nucifera*) mitochondrial genome. PLoS
861   One. 2016;11: 1–18. doi:10.1371/journal.pone.0163990

30. Yang Y, Zhou T, Duan D, Yang J, Feng L, Zhao G. Comparative analysis of the
complete chloroplast genomes of five quercus species. Front Plant Sci. 2016;7:
1–13. doi:10.3389/fpls.2016.00959
31. Cauz-santos LA, Munhoz CF, Rodde N, Cauet S, Santos AA, Penha HA, et al.
The chloroplast genome of *Passiflora edulis* (Passifloraceae) assembled from
long sequence reads: structural organization and phylogenomic studies in
malpighiales. 2017;8: 1–17. doi:10.3389/fpls.2017.00334
32. Machado L de O, Vieira L do N, Stefenon VM, Oliveira Pedrosa F de, Souza EM
de, Guerra MP, et al. Phylogenomic relationship of feijoa (*Acca sellowiana*
(O.Berg) Burret) with other Myrtaceae based on complete chloroplast genome
sequences. Genetica. 2017;145: 163–174. doi:10.1007/s10709-017-9954-1
33. Pichardo-Marcano FJ, Nieto-Blázquez ME, MacDonald AN, Galeano G, Roncal
J. Phylogeny, historical biogeography and diversification rates in an
economically important group of Neotropical palms: Tribe Euterpeae. Mol
Phylogenet Evol. 2019;133: 67–81. doi:10.1016/j.ympev.2018.12.030
34. de Santana Lopes A, Gomes Pacheco T, Nascimento da Silva O, do Nascimento
Vieira L, Guerra MP, Pacca Luna Mattar E, et al. Plastid genome evolution in
Amazonian açaí palm (*Euterpe oleracea* Mart.) and Atlantic forest açaí palm
(*Euterpe edulis* Mart.). Plant Mol Biol. 2021;105: 559–574. doi:10.1007/s11103-
020-01109-5
35. Liu L, Zhang Y, Li P. Development of genomic resources for the genus Celtis
(Cannabaceae) based on genome skimming data. Plant Divers. 2021;43: 43–53.
doi:10.1016/j.pld.2020.09.005
36. Takamatsu T, Baslam M, Inomata T, Oikawa K. Optimized method of extracting
rice chloroplast DNA for high-quality plastome resequencing and de novo
assembly. 2018;9: 1–13. doi:10.3389/fpls.2018.00266
37. Dierckxsens N, Mardulyn P, Smits G. NOVOPlasty: De novo assembly of
organelle genomes from whole genome data. Nucleic Acids Res. 2017;45.
doi:10.1093/nar/gkw955
38. Jin JJ, Yu W Bin, Yang JB, Song Y, DePamphilis CW, Yi TS, et al.
GetOrganelle: A fast and versatile toolkit for accurate de novo assembly of
organelle genomes. Genome. 2020;21: 1–31. doi:10.1101/256479
39. Langdon WB. Performance of genetic programming optimised Bowtie2 on
genome comparison and analytic testing (GCAT) benchmarks. BioData Min.
2015;8: 1–7. doi:10.1186/s13040-014-0034-0
40. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al.
SPAdes: A new genome assembly algorithm and its applications to single-cell
sequencing. J Comput Biol. 2012;19: 455–477. doi:10.1089/cmb.2012.0021
41. Tillich M, Lehwark P, Pellizzer T, Ulbricht-Jones ES, Fischer A, Bock R, et al.
GeSeq - Versatile and accurate annotation of organelle genomes. Nucleic Acids
Res. 2017;45: W6–W11. doi:10.1093/nar/gkx391
42. Abeel T, Van Parys T, Saeys Y, Galagan J, Van De Peer Y. GenomeView: A
next-generation genome browser. Nucleic Acids Res. 2012;40: 1–10.
doi:10.1093/nar/gkr995
43. Greiner S, Lehwark P, Bock R. OrganellarGenomeDRAW (OGDRAW) version
1.3.1: Expanded toolkit for the graphical visualization of organellar genomes.
Nucleic Acids Res. 2019;47: W59–W64. doi:10.1093/nar/gkz238
44. Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A.
MUMmer4: A fast and versatile genome alignment system. PLoS Comput Biol.
2018;14: 1–14. doi:10.1371/journal.pcbi.1005944

45.   Katoh K, Standley DM. MAFFT: Iterative refinement and additional methods.
913   Methods Mol Biol. 2014;1079: 131–146. doi:10.1007/978-1-62703-646-7_8

914   46.   Da Silva RS, Clementi CR, Balsanelli E, de Baura VA, de Souza EM, de Freitas
915   Fraga HP, et al. The plastome sequence of *Bactris gasipaes* and evolutionary
916   analysis in tribe Cocoseae (Arecaceae). PLoS One. 2021;16: 1–15.
917   doi:10.1371/journal.pone.0256373

918   47.   Darling ACE, Mau B, Blattner FR, Perna NT. Mauve: Multiple alignment of
919   conserved genomic sequence with rearrangements. Genome Res. 2004;14: 1394–
920   1403. doi:10.1101/gr.2289704

921   48.   Beier S, Thiel T, Münch T, Scholz U, Mascher M. MISA-web: A web server for
922   microsatellite prediction. Bioinformatics. 2017;33: 2583–2585.
923   doi:10.1093/bioinformatics/btx198

924   49.   Kurtz S, Schleiermacher C. REPuter: fast computation of maximal repeats in
925   complete genomes. Bioinforma Appl Note. 1999;15: 426–427.
926   doi:10.1093/bioinformatics/15.5.426

927   50.   Librado P, Rozas J. DnaSP v5: A software for comprehensive analysis of DNA
928   polymorphism data. Bioinformatics. 2009;25: 1451–1452.
929   doi:10.1093/bioinformatics/btp187

930   51.   Edgar RC. MUSCLE: Multiple sequence alignment with high accuracy and high
931   throughput. Nucleic Acids Res. 2004;32: 1792–1797. doi:10.1093/nar/gkh340

932   52.   Stern A, Doron-Faigenboim A, Erez E, Martz E, Bacharach E, Pupko T. Selecton
933   2007: Advanced models for detecting positive and purifying selection using a
934   Bayesian inference approach. Nucleic Acids Res. 2007;35: 506–511.
935   doi:10.1093/nar/gkm382

936   53.   Nylander JAA. MrModeltest v2. Program distributed by the author. Evolutionary
937   Biology Center, Uppsala University; 2004.

938   54.   Xie W, Lewis PO, Fan Y, Kuo L, Chen MH. Improving marginal likelihood
939   estimation for bayesian phylogenetic model selection. Syst Biol. 2011;60: 150–
940   160. doi:10.1093/sysbio/syq085

941   55.   Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, et al.
942   Mrbayes 3.2: Efficient bayesian phylogenetic inference and model choice across
943   a large model space. Syst Biol. 2012;61: 539–542. doi:10.1093/sysbio/sys029

944   56.   Miller MA, Pfeiffer W, Schwartz T. Creating the CIPRES Science Gateway for
945   inference of large phylogenetic trees. 2010 Gatew Comput Environ Work GCE
946   2010. 2010. doi:10.1109/GCE.2010.5676129

947   57.   Kass RE, Raftery AE. Bayes Factors. J Am Stat Assoc. 1995;90: 773–795.

948   58.   Rambaut A. FigTree v1. 3.1. 2012. Available from:
949   http://tree.bio.ed.ac.uk/software/figtree

950   59.   Inkscape. Inkscape Project. 2021. Available from: https://inkscape.org/pt-br/

951   60.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The
952   Sequence Alignment/Map format and SAMtools. Bioinformatics. 2009;25: 2078–
953   2079. doi:10.1093/bioinformatics/btp352

954   61.   Broad Institute. Picard Tools. Broad Institute GitHub Repository. 2018.
955   Available from: http://broadinstitute.github.io/picard/

956   62.   Catchen J, Hohenlohe PA, Bassham S, Amores A. Stacks: an analysis tool set for
957   population genomics. 2013; 3124–3140. doi:10.1111/mec.12354

958   63.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The
959   variant call format and VCFtools. Bioinformatics. 2011;27: 2156–2158.
960   doi:10.1093/bioinformatics/btr330

64.   Jombart T, Ahmed I. adegenet 1.3-1: New tools for the analysis of genome-wide

SNP data. Bioinformatics. 2011;27: 3070–3071.
doi:10.1093/bioinformatics/btr521

65. Rousset F, Lopez L, Belkhir K. R package: genepop. 2020. p. 16. Available
from: http://kimura.univ-montp2.fr/~rousset/Genepop.htm

66. R Core Team. R: A language and environment for statistical computing. Vienna,
Austria: R Foundation for Statistical Computing; 2021. Available from:
https://www.r-project.org/

67. de Santana Lopes A, Gomes Pacheco T, Nascimento da Silva O, Magalhães Cruz
L, Balsanelli E, Maltempi de Souza E, et al. The plastomes of *Astrocaryum*
aculeatum G. Mey. and *A. murumuru* Mart. show a flip-flop recombination
between two short inverted repeats. Planta. 2019;250: 1229–1246.
doi:10.1007/s00425-019-03217-z

68. Drescher A, Stephanie R, Calsa T, Carrer H, Bock R. The two largest chloroplast
genome-encoded open reading frames of higher plants are essential genes. Plant
J. 2000;22: 97–104. doi:10.1046/j.1365-313X.2000.00722.x

69. Kikuchi S, Asakura Y, Imai M, Nakahira Y, Kotani Y, Hashiguchi Y, et al. A
Ycf2-FtsHi heteromeric AAA-ATPase complex is required for chloroplast
protein import. Plant Cell. 2018;30: 2677–2703. doi:10.1105/tpc.18.00357

70. Barrett CF, Davis JI, Leebens-Mack J, Conran JG, Stevenson DW. Plastid
genomes and deep relationships among the commelinid monocot angiosperms.
Cladistics. 2013;29: 65–87. doi:10.1111/j.1096-0031.2012.00418.x

71. Vieira MLC, Santini L, Diniz AL, Munhoz C de F. Microsatellite markers: What
they mean and why they are so useful. Genet Mol Biol. 2016;39: 312–328.
doi:10.1590/1678-4685-GMB-2016-0027

72. Eguiluz M, Rodrigues NF, Guzman F, Yuyama P, Margis R. The chloroplast
genome sequence from *Eugenia uniflora*, a Myrtaceae from Neotropics. Plant
Syst Evol. 2017;303: 1199–1212. doi:10.1007/s00606-017-1431-x

73. Ping J, Feng P, Li J, Zhang R, Su Y, Wang T. Molecular evolution and SSRs
analysis based on the chloroplast genome of *Callitropsis funebri*s. Ecol Evol.
2021;11: 4786–4802. doi:10.1002/ece3.7381

74. Rogalski M, Vieira LDN, Fraga HP, Guerra MP. Plastid genomics in
horticultural species: Importance and applications for plant population genetics,
evolution, and biotechnology. Front Plant Sci. 2015;6: 1–17.
doi:10.3389/fpls.2015.00586

75. Zhu A, Guo W, Gupta S, Fan W, Mower JP. Evolutionary dynamics of the
plastid inverted repeat: The effects of expansion, contraction, and loss on
substitution rates. New Phytol. 2016;209: 1747–1756. doi:10.1111/nph.13743

76. Zeng S, Zhou T, Han K, Yang Y, Zhao J, Liu ZL. The complete chloroplast
genome sequences of six rehmannia species. Genes (Basel). 2017;8.
doi:10.3390/genes8030103

77. Milligan BG, Hampton JN, Palmer JD. Dispersed repeats and structural
reorganization in subclover chloroplast DNA. Mol Biol Evol. 1989;6: 355–368.
doi:10.1093/oxfordjournals.molbev.a040558

78. Song Y, Dong W, Liu B, Xu C, Yao X, Gao J, et al. Comparative analysis of
complete chloroplast genome sequences of two tropical trees *Machilus*
yunnanensis and *Machilus balansae* in the family Lauraceae. Front Plant Sci.
2015;6: 1–8. doi:10.3389/fpls.2015.00662

79. Barrett CF, Baker WJ, Comer JR, Conran JG, Lahmeyer SC, Leebens-Mack JH,
et al. Plastid genomes reveal support for deep phylogenetic relationships and
extensive rate variation among palms and other commelinid monocots. New

Phytol. 2016;209: 855–870. doi:10.1111/nph.13617

80. Brancalion PHS, Oliveira GCX, Zucchi MI, Novello M, van Melis J, Zocchi SS,
et al. Phenotypic plasticity and local adaptation favor range expansion of a
Neotropical palm. Ecol Evol. 2018;8: 7462–7475. doi:10.1002/ece3.4248

81. Du X, Zeng T, Feng Q, Hu L, Luo X, Weng Q, et al. The complete chloroplast
genome sequence of yellow mustard (*Sinapis alba* L.) and its phylogenetic
relationship to other Brassicaceae species. Gene. 2020;731.
doi:10.1016/j.gene.2020.144340

82. Liu Q, Li X, Li M, Xu W, Schwarzacher T, Heslop-Harrison JS. Comparative
chloroplast genome analyses of Avena: Insights into evolutionary dynamics and
phylogeny. BMC Plant Biol. 2020;20: 1–20. doi:10.1186/s12870-020-02621-y

83. Huang Y, Yang Z, Huang S, An W, Li J, Zheng X. Comprehensive analysis of
Rhodomyrtus tomentosa chloroplast genome. Plants. 2019;8.
doi:10.3390/plants8040089

84. Williams-Carrier R, Brewster C, Belcher SE, Rojas M, Chotewutmontri P,
Ljungdahl S, et al. The *Arabidopsis pentatricopeptide* repeat protein LPE1 and its
maize ortholog are required for translation of the chloroplast psbJ RNA. Plant J.
2019;99: 56–66. doi:10.1111/tpj.14308

85. Díaz BG, Zucchi MI, Alves-Pereira A, de Almeida CP, Moraes ACL, Vianna
SA, et al. Genome-wide SNP analysis to assess the genetic population structure
and diversity of Acrocomia species. PLoS One. 2021;16: 1–24.
doi:10.1371/journal.pone.0241025

86. Viana JPG, Siqueira MVBM, Araujo FL, Grando C, Sujii PS, De Aguiar
Silvestre E, et al. Genomic diversity is similar between Atlantic Forest
restorations and natural remnants for the native tree *Casearia sylvestris* Sw.
PLoS One. 2019;13: 1–14. doi:10.1371/journal.pone.0192165

87. Cordeiro EMG, Macrini CM, Sujii PS, Schwarcz KD, Pinheiro JB, Rodrigues
RR, et al. Diversity, genetic structure, and population genomics of the tropical
tree *Centrolobium tomentosum* in remnant and restored Atlantic forests. Conserv
Genet. 2019;20: 1073–1085. doi:10.1007/s10592-019-01195-z

88. Allendorf FW, Hohenlohe PA, Luikart G. Genomics and the future of
conservation genetics. Nat Rev Genet. 2010;11: 697–709. doi:10.1038/nrg2844

89. Willing EM, Dreyer C, van Oosterhout C. Estimates of genetic differentiation
measured by fst do not necessarily require large sample sizes when using many
snp markers. PLoS One. 2012;7: 1–7. doi:10.1371/journal.pone.0042649

90. Hamblin MT, Warburton ML, Buckler ES. Empirical comparison of simple
sequence repeats and single nucleotide polymorphisms in assesment of maize
diversity and relatedness. PLoS One. 2007;2. doi:10.1371/journal.pone.0001367

1050

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5

Fig 6

Fig 7