Deep topographic proteomics of a human brain tumour

Simon Davis[1,2], Connor Scott[3], Janina Oetjen[4], Philip D Charles[1,5], Benedikt M Kessler[1,2], Olaf Ansorge[3], Roman Fischer[1,2] *

[1] Target Discovery Institute, Centre for Medicines Discovery, Nuffield Department of Medicine, University of Oxford, Roosevelt Drive, Oxford, OX3 7FZ, UK

[2] Chinese Academy for Medical Sciences Oxford Institute, Nuffield Department of Medicine, University of Oxford, Roosevelt Drive, Oxford, OX3 7FZ, UK

[3] Academic Unit of Neuropathology, Nuffield Department of Clinical Neurosciences, University of Oxford, John Radcliffe Hospital, Oxford, OX3 9DU, UK

[4] Bruker Daltonics GmbH & Co. KG, Fahrenheitstraße 4, 28359 Bremen, Germany

[5] Big Data Institute, Nuffield Department of Medicine, University of Oxford, Roosevelt Drive, Oxford, OX3 7FZ, UK

* Corresponding author. E-mail: roman.fischer@ndm.ox.ac.uk. Phone: +44 1865 743639

Running title: Spatial brain tumour proteomics

Key words: brain tumour, spatial proteomics, mass spectrometry, imaging

1 Abstract

The spatial organisation of cellular protein expression profiles within tissue determines

cellular function and is key to understanding disease pathology. To define molecular

phenotypes in the spatial context of tissue, there is a need for unbiased, quantitative

technology capable of mapping proteomes within tissue structures. Here, we present a

workflow for spatially-resolved, quantitative proteomics of tissue that generates maps of

protein abundance across tissue slices derived from a human atypical teratoid-rhabdoid

tumour (AT/RT). We employ spatially-aware algorithms that do not require prior knowledge

of the fine tissue structure to detect proteins and pathways with spatial abundance

patterns. We identified PYGL, ASPH and CD45 as spatial markers for tumour boundary and

reveal immune response-driven, spatially-organised protein networks of the extracellular

tumour matrix. Overall, this work informs on methods for spatially resolved deep proteo-

phenotyping of tissue heterogeneity, to push the boundaries of understanding tissue

biology and pathology at the molecular level.

1   Main

Tissues contain various microscopic features, cell types, and phenotypically diverse

subpopulations while the location of cells within a tissue and their spatial neighbourhood is

crucial for determining their identity and function[1–6]. The cellular composition of tissue has

substantial influence on measured co-expression signals within the molecular profiles of

bulk tissue and their microenvironment, contributing to the influence of cellular function,

signalling and different disease outcomes[7–11]. Recent technology developments in spatially-

resolved sequencing technologies have enabled the characterisation of spatially

heterogeneous gene expression profiles within tissues[12,13]. However, while genomic and

transcriptomic alterations act as drivers of disease, the proteins they encode regulate

essentially all cellular processes and therefore also need consideration when investigating

tissue spatial heterogeneity[14].

A range of mass spectrometry (MS)-based techniques are available to map the distribution

of proteins throughout tissues and cells. Mass spectrometry imaging (MSI) enables the

determination of proteins or other molecules within a sample by rastering an ion source

over a sample in a grid pattern. MSI techniques to visualise bio-molecules *in situ* cannot

generate in-depth proteome data and may require prior knowledge of measurement

targets[15–18].

Laser capture microdissection (LCM) is well-placed to address the limitations of the

spatially-resolved mass spectrometry methods described above[19]. LCM can extract regions

from a tissue slice ranging from single cells to square millimetres of tissue[20,21]. We and

others have previously described several methods coupling LCM to MS-based proteomics[22–

26]. The approach has been used to investigate a wide range of tissue biology, generally in a

'feature-driven' approach, extracting tissue regions based on visible features[27–30]. Mund et

al. recently developed the concept of Deep Visual Proteomics by combining high-resolution

imaging and machine learning-based image analysis to classify, isolate and analyse cells with

a sensitive proteomics workflow[31].

However, sampling in a systematic manner, like MSI, could reveal novel tissue fine structure

and give insights into the spatial protein expression patterns. For example, Piehowski et al.

used LCM-proteomics to sample mouse uterine tissue in a rastered grid with a resolution of

100 µm and their custom, robotic, nanolitre-scale nanoPOTS sample preparation platform to

quantify over 2,000 proteins within the tissue[32].

Here, we systematically performed spatially-resolved measurements of a human brain

tumour proteome using laser capture microdissection to a depth of over 5,000 proteins. We

use spatially-aware statistical methods to identify proteins and pathways displaying

differential spatial and clustered expression within tissue sections. This did not require prior

knowledge of tissue structures, features, or pathology. Furthermore, clustering of protein

expression reveals new, spatially defined proteo-phenotypes within the otherwise

homogeneous macrostructure of the analysed tumour revealing spatially resolved ECM

biology correlating with immune response.

Results

In order to establish required tissue areas to achieve a target proteome depth of 4000

quantified proteins in low (Orbitrap Fusion Lumos, 60-gradient) and medium throughput

proteomics platforms (TimsTOF Pro, 17-minute gradient) we collected tissue areas ranging

from 316 µm^2 to 1,000,000 µm^2 from 10 µm-thick sections of human brain (Figure S1). We

observed that areas above 316,000 µm^2 result in diminishing returns with protein

identifications scaling between 282 and 3480 on the Orbitrap and 127 and 3318 on the

timsTOF Pro platforms, respectively.

Proteomic topography of a human brain tumour

After characterising the upper and lower limits of the workflow, we sampled a 10 µm thick

section of an atypical teratoid-rhabdoid tumour (AT/RT) block (~20 x 15 mm). The tissue was

subdivided into 384 (24 x 16) square 'pixels' with an area of ~694,000 µm^2 (side length of

833 µm); each pixel was isolated by LCM and processed with our LCM-SP3 protocol[24] (Figure

1). Each sample was analysed on the 17-minute timsTOF Pro setup. In total, 5,321 proteins

were identified, with 32 – 4,741 proteins identified per sample (Figure S2). This range

includes empty pixels where no tissue was visible, demonstrating a low level of

contamination throughout the workflow. Quantified proteins can be mapped back to their

original positions within the tissue grid. Figure 2A shows proteomic maps for four example

proteins, liver glycogen phosphorylase (PYGL), peripherin (PRPH), haemoglobin (HBB) and

histone H4 (HIST1H4A). Glycogen phosphorylase releases glucose from glycogen for entry

into glycolysis, and its expression in cancer is associated with malignant phenotypes,

hypoxia resistance and cancer cell survival[33]. Peripherin is an intermediate filament protein

without a clear function and is highly expressed during development and after nerve injury;

its expression pattern is consistent with the tumour growth into surrounding normal brain

tissue[34–36].

We tested the 4,306 proteins quantified in at least 9 pixels for spatial autocorrelation[37]

using the unsupervised Moran's *I* test. Moran's I returns values between -1 for complete

dispersion and +1 for complete correlation, with 0 indicating random distribution. Removing

empty pixels and areas of haemorrhage from the analysis, we found that 3,212 proteins

demonstrated correlated spatial expression profiles with ($q \leq 0.05$), while excluding any

impact of sampling and processing on spatial variability (Figure S3).

In order to increase spatial resolution particularly in the 'brain/tumour' interface as

identified by H&E staining, we sampled with 96 pixels with 350 μm x 350 μm side-length.

This region of tissue contains predominantly normal and neoplastic cells along with a large,

prominent blood vessel. In total, 3,994 proteins were quantified in at least one pixel. The

increased resolution proteomic maps for PYGL, PRPH, haemoglobin and histone H4 are

shown in Figure 2B. Their expression is consistent with the large field-of-view data, with

PYGL and PRPH showing opposite expression patterns across the margin between solid

tumour (high PYGL, low PRPH) and brain/tumour interface (low PYGL, high PRPH) and

haemoglobin co-localising with the visible blood vessels. Histone H4 shows even expression

across the two annotated areas, with a region of lower expression corresponding with a

visibly diffuse patch of tissue. Of the 3,050 proteins quantified in at least 9 pixels, 1,375

show evidence for significant spatial autocorrelation (Moran's *I* test, $q \leq 0.05$).

Three candidate proteins showing significant spatial variation were selected for follow-up

immunohistochemistry (IHC) staining: glycogen phosphorylase, aspartate beta-hydroxylase

(ASPH) and CD45 (PTPRC) to validate the spatially resolved protein expression data

generated above. The IHC staining images closely resemble the protein intensity

distributions (Figure 3A, C, E) within the proteomic maps for these three proteins. Both PYGL

and ASPH show intense IHC staining in the region of solid tumour (Figure 3B,D), and CD45

shows intense staining in the region of tissue corresponding to the upper-left pixels in the proteomic map (Figure 3F).

Spatial proteomic mapping highlights molecular pathways underlying tissue heterogeneity

As Moran's *I* measures global autocorrelation, it does not indicate where the locations that drive the autocorrelation occur. To investigate which regions of the sampled tissue show similar expression, the data were clustered and the cluster labelling of pixels were mapped back to their spatial location (Figure 4A). The clusters generally form contiguous regions in space, with some long-range co-clustering in smaller clusters. The margin between solid tumour and brain/tumour interface is well-represented by the border between cluster 1 (solid tumour) and cluster 3 (brain/tumour interface). In addition to the cluster map, the assigned clusters were plotted onto a uniform manifold approximation and projection (UMAP) visualisation (Figure 4B). The clusters visible in the UMAP plot correspond well to the spatial clusters in Figure 4A.

This clustering approach generates spatially well-defined clusters, allowing for a feature-driven approach without prior knowledge of the histopathological details as the clustering is performed only on the protein quantitation data, without information on the spatial relationship between samples. Proteins quantified were tested for significantly differing intensities across the clusters using a one-way ANOVA test. There are 3,512 proteins with significant evidence ($q \leq 0.05$) for differential intensity between two or more clusters. Pairwise Tukey *post-hoc* tests were used to generate fold-change estimates between pairs of clusters, and these fold-changes were used for gene set enrichment analysis (GSEA). Functional analysis of clusters 1 & 3 indicates that proteins and processes involved in protein translation and modification, extracellular matrix organisation, energy pathways, mRNA processing, steroid synthesis, neuronal cell adhesion, and neuronal differentiation are differentially abundant between the region of solid tumour and brain/tumour interface (Figure 4C).

In order to discover new spatial relationships between phenotypically similar and distinct areas, we performed unbiased spatial clustering by integrating a proteomic similarity matrix (rank correlation of top 25 % most variable proteins) with a complementary spatial similarity

network (based on Euclidean distances) using Affinity Network Fusion (ANF)[38]. This spatially-

aware clustering method resulted in six clusters (Cluster A-F) covering the regions of solid

tumour, brain/tumour interface, immune infiltration, and haemorrhage (Figure 4D).

The functional differences between these clusters were investigated by first determining

'marker proteins' for each cluster by testing for differential abundance of a protein in one

cluster versus all other clusters, iteratively for each protein-cluster pair. Protein markers for

each cluster were then tested for functional enrichment against the MSigDB Hallmark gene

sets (Figure 4E)[39]. Cluster A shows enrichment for blood-related hallmarks, consistent with

this cluster comprising predominantly haemorrhage. Clusters B, D & F show enrichment for

cell cycle and growth-related hallmarks, consistent with the 'solid tumour' histology

evaluation. Cluster C shows enrichment of immune-related hallmarks and the Epithelial-

Mesenchymal Transition hallmark, consistent with the presence of immune cell-marker

proteins above. Cluster E shows enrichment for the "KRAS Signalling Dn" and "Epithelial-

Mesenchymal Transition" hallmarks.

Clustering in high- and low-resolution maps was broadly consistent (Figure 5A) with

generally contiguous clusters that represent the solid tumour (cluster 1), the brain/tumour

interface (cluster 3), the margin between (clusters 2 & 5), and blood vessels (cluster 6). A

volcano plot between cluster 3 and cluster 1 reflects the previously observed differential

abundance of PYGL, ASPH and PRPH and other marker candidates (Figure 5B). Functional

analysis of clusters 1 & 3 indicates that proteins involved in extracellular matrix, cell

adhesion & motility, angiogenesis, immune processes, epidermis function, and neuronal

development are differentially abundant between solid tumour and brain/tumour interface

(Figure 5C).

We investigated the expression of other immune cell-marker proteins because of the above

differences above in immune processes, and highly localised CD45 abundance and staining

(Figure 3A). Two neutrophil markers, neutrophil cytosolic factor 2 & 4 (NCF2 & NCF4), show

co-localisation with CD45 in the upper left of the sampled region along with two marker

proteins for pro-tumour M2 macrophages, CD163 & mannose receptor C-type 1 (CD163 &

MRC1)[40,41]. These proteins' peak expression locations correspond with clusters 13 and 6 for

the neutrophil and macrophage markers, respectively (Figure S4) and shows increased

abundance for many proteins involved in neutrophil function and other immune-related

processes such as B-cell differentiation and the JNK cascade within cluster 13. Within cluster

6, proteins involved in cell death, the cell cycle, morphogenesis, hedgehog signalling,

collagen, and cytoskeletal organisation show increased abundance (Figure S5).

MALDI imaging visualises lipid distributions consistent with proteomic maps

Due to the relatively coarse resolution of out proteomics approach, we also performed mass

spectrometry imaging to investigate the spatial distribution of lipids within the tissue on an

adjacent section using MALDI (Figure S6). Bisecting K-means clustering of the MALDI

imaging pixels derived from 498 molecular ions broadly reflects the proteomic and cluster

maps' patterns, but with higher spatial resolution (20 μm). The upper-left region is a distinct

cluster, and the boundary between the solid tumour and brain/tumour interface is visible.

The ion image of haeme can be used to visualise the vascularisation of the tissue, and shows

variability across the tissue, potentially indicating areas of nutrient gradients within the

tissue.

Plotting the ion image of the 689.56 and 673.58 m/z ions, which correspond to the m/z of

cholesterol ester[42] (18:1) $[M+K]^+$ and $[M+Na]^+$ respectively, shows the highest intensity in

the region of the tissue that corresponds to the area of high macrophage marker expression

(cluster 6) in the proteomic maps (Figure S6), consistent with M1/M2 macrophage

accumulation at the tumour interface visible in Figure S6 and Figures 2&3[43].

Spatial network clustering reveals extracellular matrix and integrin receptor

heterogeneity

Throughout the spatial analysis and manual inspection of the data we noticed the common

presence of many extracellular matrix-related proteins in the resulting enriched pathways

and highly spatially-correlated proteins. Because of the relevance of the ECM within many

cellular pathways and tumour development/progression, we performed ANF clustering

based on the proteins annotated as 'Core Matrisome' proteins within MatrisomeDB[44]. This

resulted in nine clusters (A-I) which are generally spatially contiguous across the tissue

(Figure 6A). ECM-defined clusters were used for functional enrichment of proteins within

each cluster, (Figure 6B) and are generally consistent with the functional enrichments in

Figure 4E without ECM focus, linking overall spatial proteome distribution with ECM

architecture in the tumour structure. To determine whether differential ECM abundance or

composition was driving this clustering, we plotted the summed (Figure S7A) and mean (Figure S7B) abundance of the core matrisome proteins. These aggregate ECM spatial abundances show that both total ECM abundance (in the haemorrhage and immune infiltration area; clusters A & C in Figure 6A) and ECM composition (in the remaining clusters) are contributing to these cluster definitions.

Plotting the spatial distributions of detected collagens reveals broad heterogeneity in abundance of collagen proteins across the different collagen subfamilies (Figure 6C). The fibrillar collagens COL2A1, COL11A1, and COL11A2 show higher expression in the solid tumour region and other fibrillar collagens, COL1A2 and COL3A1, show higher abundance in the brain/tumour interface region. Other collagen subfamily members also show differential spatial abundance: COL6A1, COL6A2, COL26A1 (filament forming collagens); COL12A1, COL14A1 (fibril associated collagens). Collagens 4A1, 4A2 (network forming collagens), 15A1 and 18A1 (multiplexins) show spatially homogeneous expression. These spatial distributions broadly map to the ANF clusters. In addition, several integrin proteins also show spatial heterogeneity within the tissue (Figure 6C). Integrin α5, α7, α10, αV, αM and β2 show spatial heterogeneity whereas integrin β1 shows relatively homogeneous expression. Other proteins associated with the ECM also show spatial heterogeneity such as CD44, Lysyl oxidase (LOX), Lumican (LUM) (lower in solid tumour region), Versican (VCAN), and Cathepsin D & G (higher in immune cell-enriched region). In general ECM and ECM-associated proteins as defined in MatrisomeDB show a higher mean Moran's I value than non-ECM proteins (ECM: I = 0.334, ECM-associated: I = 0.320, non-ECM: I = 0.219) (Figure S8)[44].

Discussion

The presented spatial proteomics workflow addresses the need to generate highly multiplexed, quantitative, spatially resolved measurements of proteins within tissue to understand the spatial organisation of molecular pathways in health and pathology. We demonstrate methods for the systematic sampling of tissue sub-sections using laser capture microdissection (LCM), sample preparation, liquid chromatography-mass spectrometry and advanced statistical analysis of topographic data. We have shown that these optimised methods are capable of detecting proteins and pathways with spatially variable abundance

within tumour and tumour-infiltrated normal tissue. The application of spatially aware data analysis pinpoints processes with deep molecular resolution without prior knowledge of the tissue composition, thus creating an objective, unbiased way of deep phenotyping pathological tissue in its biological context.

A growing number of studies are currently focused on feature-driven LCM-coupled proteomics. In contrast, here, we propose a systematic approach, which allows the unbiased creation of comprehensive proteomic maps at the individual protein or pathway level. These maps can fulfil the requirements for feature-driven analysis by reconstituting features from systematic sampling and allow the discovery of new proteo-phenotypes without the requirement of visually identifiable parameters.

Through our generation of proteomic maps, we have demonstrated the presence of molecular heterogeneity at multiple scales within ATRT tumour tissue sections, revealing proteomic differences between areas of tissue that appear visually homogeneous with good agreement with immunohistochemistry staining. Additionally, we demonstrate that this approach can also generate information on immune cell infiltration and state within the tissue, by the detection of neutrophil and pro-tumour M2 macrophage markers at different distances from the solid tumour.

We also have demonstrated that this approach provides spatial information on the cellular microenvironment across the tissue. The increased level of fibrillar collagens within the solid tumour is consistent with observations of increased fibrillar collagen deposition in glioblastoma tumours and the measurement of increased tissue stiffness of tumour types including brain tumours[45,46]. The spatial abundance of integrin subunits αM and β2 correlates with neutrophil marker abundance, strongly suggesting localised neutrophil accumulation[47]. Integrin α10 abundance correlates well with the abundance of COL2A1 and COL11A1, consistent with the observation that the α10β1 integrin receptor binds to type 2 and type 11 collagens[48]. Although functional interaction between these proteins cannot be determined from the methods used here, further investigation into integrin receptor signalling in ATRT could be a worthwhile avenue of investigation as integrin α10/β1 has shown potential as a therapeutic target in a glioblastoma xenograft model[49].

The work we present here has several avenues for future improvement to generate more comprehensive information about biological structures. Integrating multiple data modalities and patient data is critical to a more comprehensive understanding of disease processes relying on advanced bioinformatics tools[50,51]. The same is expected to be true of spatially-resolved data. Further development of analytical and statistical methods that are aware of spatial relationships between samples is also required to maximise the utility of spatially resolved proteomics. Missing value imputation and machine learning approaches could benefit from taking samples' spatial relationships into account[52,53].

Spatial resolution should be balanced with the expectation for a meaningful depth to cover pathways of interest and technical limitations for sensitivity and throughput. Increasing the spatial resolution demonstrated here towards the single-cell level and covering comparable areas would be a formidable analytical challenge, further escalating when analysing tissue in three dimensions, despite the development of novel high-throughput LC-MS platforms that can now robustly analyse 1000s of samples relatively quickly[54–56]. Therefore, future approaches are likely to use systematic spatial proteomic analysis, possibly compromising spatial resolution but incorporating elements of machine learning, using orthogonal higher resolution 'omics and imaging data to infer protein abundance towards individual cell resolution, as can be done on spatial transcriptomics data[57–60]. With the detection of LCM-based and cell-type resolved deep proteomes, these data will be highly complementary to current imaging technologies and increase the understanding of spatially resolved biological and pathological processes at the molecular level[24,31,61].

Acknowledgements

SD acknowledges support from the Nuffield Department of Medicine. SD, PDC, BMK & RF acknowledge support from the Chinese Academy of Medical Sciences Medical Sciences 2018-I2M-2-002. PDC was supported by Pfizer funding awarded to BMK.

We acknowledge the Oxford Brain Bank, supported by the Medical Research Council (MRC, MR/L022656/1) and Brains for Dementia Research (BDR) (Alzheimer Society and Alzheimer Research UK). This research project was funded by the NIHR Oxford Biomedical Research Centre (to OA, BRC-1215-20008). The views expressed are those of the authors and not

necessarily those of the NHS, the NIHR, or the Department of Health. This work uses data

provided by patients and collected by the NHS as part of their care and support and would

not have been possible without access to this data. The NIHR recognises and values the role

of patient data, securely accessed and stored, both in underpinning and leading to

improvements in research and care.

Competing interests

J.O is an employee of Bruker Daltonics GmbH & Co. KG. All other authors have no competing

interests.

References

1. Asp, M. *et al.* A Spatiotemporal Organ-Wide Gene Expression and Cell Atlas of the Developing Human Heart. *Cell* **179,** 1647–1660.e19 (2019).

2. Baccin, C. *et al.* Combined single-cell and spatial transcriptomics reveal the molecular, cellular and spatial bone marrow niche organization. *Nat. Cell Biol.* **22,** 38–48 (2020).

3. Moncada, R. *et al.* Integrating microarray-based spatial transcriptomics and single-cell RNA-seq reveals tissue architecture in pancreatic ductal adenocarcinomas. *Nat. Biotechnol.* **38,** 333–342 (2020).

4. Shah, S., Lubeck, E., Zhou, W. & Cai, L. In Situ Transcription Profiling of Single Cells Reveals Spatial Organization of Cells in the Mouse Hippocampus. *Neuron* **92,** 342–357 (2016).

5. Thrane, K., Eriksson, H., Maaskola, J., Hansson, J. & Lundeberg, J. Spatially resolved transcriptomics enables dissection of genetic heterogeneity in stage III cutaneous malignant melanoma. *Cancer Res.* **78,** 5970–5979 (2018).

6. Berglund, E. *et al.* Spatial maps of prostate cancer transcriptomes reveal an unexplored landscape of heterogeneity. *Nat. Commun.* **9,** 1–13 (2018).

7. Keren, L. *et al.* A Structured Tumor-Immune Microenvironment in Triple Negative Breast Cancer Revealed by Multiplexed Ion Beam Imaging. *Cell* **174,** 1373–1387 (2018).

8. Farahbod, M. & Pavlidis, P. Untangling the effects of cellular composition on coexpression analysis. *Genome Res.* **30,** 849–859 (2020).

9. Bloom, A. B. & Zaman, M. H. Influence of the microenvironment on cell fate determination and migration. *Physiological Genomics* **46,** 309–314 (2014).

10. De Palma, M., Biziato, D. & Petrova, T. V. Microenvironmental regulation of tumour angiogenesis. *Nature Reviews Cancer* **17,** 457–474 (2017).

11. Balkwill, F. R., Capasso, M. & Hagemann, T. The tumor microenvironment at a glance. *J. Cell Sci.* **125,** 5591–5596 (2012).

12. Mamlouk, S. *et al.* DNA copy number changes define spatial patterns of heterogeneity in colorectal cancer. *Nat. Commun.* **8,** 14093 (2017).

13. Marek, S. *et al.* Spatial and Temporal Organization of the Individual Human Cerebellum. *Neuron* **100,** 977–993 (2018).

14. Aebersold, R. & Mann, M. Mass-spectrometric exploration of proteome structure and function. *Nature* **537,** 347–355 (2016).

15. Caprioli, R. M., Farmer, T. B. & Gile, J. Molecular Imaging of Biological Samples: Localization of Peptides and Proteins Using MALDI-TOF MS. *Anal. Chem.* **69,** 4751–4760 (1997).

16. Todd, P. J., Gregory Schaaff, T., Chaurand, P. & Caprioli, R. M. Organic ion imaging of biological tissue with secondary ion mass spectrometry and matrix-assisted laser desorption/ionization. *J. Mass Spectrom.* **36,** 355–369 (2001).

17. Angelo, M. *et al.* Multiplexed ion beam imaging of human breast tumors. *Nat. Med.* **20,** 436–442 (2014).

18. Giesen, C. *et al.* Highly multiplexed imaging of tumor tissues with subcellular resolution by mass cytometry. *Nat. Methods* **11,** 417–422 (2014).

19. De Marchi, T. *et al.* The advantage of laser-capture microdissection over whole tissue analysis in proteomic profiling studies. *Proteomics* **16,** 1474–1485 (2016).

20. Zhu, Y. *et al.* Spatially Resolved Proteome Mapping of Laser Capture Microdissected Tissue with Automated Sample Transfer to Nanodroplets. *Mol. Cell. Proteomics* **17,** 1864–1874 (2018).

21. Buczak, K. *et al.* Spatial tissue proteomics quantifies inter- and intratumor heterogeneity in hepatocellular carcinoma (HCC). *Mol. Cell. Proteomics* **17,** 810–825 (2018).

22. Longuespée, R. *et al.* A laser microdissection-based workflow for FFPE tissue microproteomics: Important considerations for small sample processing. *Methods* **104,** 154–162 (2016).

23. Zhu, Y. *et al.* Development of a laser capture microscope-based single-cell-type

proteomics tool for studying proteomes of individual cell layers of plant roots. *Hortic. Res.* **3,** (2016).

24. Davis, S., Scott, C., Ansorge, O. & Fischer, R. Development of a Sensitive, Scalable Method for Spatial, Cell-Type-Resolved Proteomics of the Human Brain. *J. Proteome Res.* **18,** 1787–1795 (2019).

25. Coscia, F. *et al.* A streamlined mass spectrometry–based proteomics workflow for large-scale FFPE tissue analysis. *J. Pathol.* **251,** 100–112 (2020).

26. Griesser, E. *et al.* Quantitative Profiling of the Human Substantia Nigra Proteome from Laser-capture Microdissected FFPE Tissue. *Mol. Cell. Proteomics* **19,** 839–851 (2020).

27. Waanders, L. F. *et al.* Quantitative proteomic analysis of single pancreatic islets. *Proc. Natl. Acad. Sci. U. S. A.* **106,** 18902–7 (2009).

28. Clair, G. *et al.* Spatially-Resolved Proteomics: Rapid Quantitative Analysis of Laser Capture Microdissected Alveolar Tissue Samples. *Sci. Rep.* **6,** 39223 (2016).

29. Eckert, M. A. *et al.* Proteomics reveals NNMT as a master metabolic regulator of cancer-associated fibroblasts. *Nature* **569,** 723–728 (2019).

30. Herrera, J. A. *et al.* Laser capture microdissection coupled mass spectrometry (LCM-MS) for spatially resolved analysis of formalin-fixed and stained human lung tissues. *Clin. Proteomics* **17,** 24 (2020).

31. Mund, A. *et al.* Deep Visual Proteomics defines single-cell identity and heterogeneity. *Nat. Biotechnol.* **40,** 1231–1240 (2022).

32. Piehowski, P. D. *et al.* Automated mass spectrometry imaging of over 2000 proteins from tissue sections at 100-µm spatial resolution. *Nat. Commun.* **11,** 1–12 (2020).

33. Favaro, E. *et al.* Glucose Utilization via Glycogen Phosphorylase Sustains Proliferation and Prevents Premature Senescence in Cancer Cells. *Cell Metab.* **16,** 751–764 (2012).

34. Porlier, M. M., de Néchaud, B. & Gros, F. Peripherin, a new member of the intermediate filament protein family. *Dev. Neurosci.* **6,** 335–344 (1983).

35. Yuan, A. *et al.* Peripherin is a subunit of peripheral nerve neurofilaments: Implications for differential vulnerability of cns and peripheral nervous system axons. *J. Neurosci.* **32,** 8501–8508 (2012).

36. Troy, C. M., Brown, K., Greene, L. A. & Shelanski, M. L. Ontogeny of the neuronal intermediate filament protein, peripherin, in the mouse embryo. *Neuroscience* **36,** 217–237 (1990).

37. Moran, P. A. P. Notes on Continuous Stochastic Phenomena. *Biometrika* **37,** 17–23 (1950).

38. Ma, T. & Zhang, A. Affinity network fusion and semi-supervised learning for cancer patient clustering. *Methods* **145,** 16–24 (2018).

39. Liberzon, A. *et al.* The Molecular Signatures Database Hallmark Gene Set Collection. *Cell Syst.* **1,** 417–425 (2015).

40. Orecchioni, M., Ghosheh, Y., Pramod, A. B. & Ley, K. Macrophage Polarization: Different Gene Signatures in M1(LPS+) vs. Classically and M2(LPS−) vs. Alternatively Activated Macrophages. *Front. Immunol.* **10,** (2019).

41. Hu, J. M. *et al.* CD163 as a marker of M2 macrophage, contribute to predict aggressiveness and prognosis of Kazakh esophageal squamous cell carcinoma. *Oncotarget* **8,** 21526–21538 (2017).

42. Guerrini, V. & Gennaro, M. L. Foam Cells: One Size Doesn't Fit All. *Trends Immunol.* **40,** 1163–1179 (2019).

43. Murray, P. J. & Wynn, T. A. Protective and pathogenic functions of macrophage subsets. *Nat. Rev. Immunol.* **11,** 723–737 (2011).

44. Shao, X., Taha, I. N., Clauser, K. R., Gao, Y. (Tom) & Naba, A. MatrisomeDB: the ECM-protein knowledge database. *Nucleic Acids Res.* **48,** D1136–D1144 (2020).

45. Huijbers, I. J. *et al.* A Role for Fibrillar Collagen Deposition and the Collagen Internalization Receptor Endo180 in Glioma Invasion. *PLoS One* **5,** (2010).

46. Nia, H. T., Munn, L. L. & Jain, R. K. Physical traits of cancer. *Science (80-.).* **370,** (2020).

47. Zhang, L. The αMβ2 integrin and its role in neutrophil function. *Cell Res. 1999 93* **9,** 171–178 (1999).

48. Lundgren-Åkerlund, E. & Aszòdi, A. Integrin α10β1: A Collagen Receptor Critical in Skeletal Development. 61–71 (2014). doi:10.1007/978-94-017-9153-3_4

49. Thorén, M. M. *et al.* Integrin α10, a Novel Therapeutic Target in Glioblastoma, Regulates Cell Migration, Proliferation, and Survival. *Cancers 2019, Vol. 11, Page 587* **11,** 587 (2019).

50. Santos, A. *et al.* A knowledge graph to interpret clinical proteomics data. *Nat. Biotechnol.* (2022). doi:10.1038/s41587-021-01145-6

51. Subramanian, I., Verma, S., Kumar, S., Jere, A. & Anamika, K. Multi-omics Data Integration, Interpretation, and Its Application. *Bioinform. Biol. Insights* **14,** (2020).

52. Li, Z., Song, T., Yong, J. & Kuang, R. Imputation of spatially-resolved transcriptomes by graph-regularized tensor completion. *PLoS Comput. Biol.* **17,** e1008218 (2021).

53. Boluki, S., Zamani Dadaneh, S., Qian, X. & Dougherty, E. R. Optimal clustering with missing values. *BMC Bioinformatics* **20,** 1–10 (2019).

54. Bache, N. *et al.* A novel LC system embeds analytes in pre-formed gradients for rapid, ultra-robust proteomics. *Mol. Cell. Proteomics* **17,** 2284–2296 (2018).

55. Bian, Y. *et al.* Robust, reproducible and quantitative analysis of thousands of proteomes by micro-flow LC–MS/MS. *Nat. Commun.* **11,** 1–12 (2020).

56. Messner, C. B. *et al.* Ultra-fast proteomics with Scanning SWATH. *Nat. Biotechnol.* **39,** 846–854 (2021).

57. Pang, M., Su, K. & Li, M. Leveraging information in spatial transcriptomics to predict super-resolution gene expression from histology images in tumors. *bioRxiv* 2021.11.28.470212 (2021). doi:10.1101/2021.11.28.470212

58. Monjo, T., Koido, M., Nagasawa, S., Suzuki, Y. & Kamatani, Y. Efficient prediction of a spatial transcriptomics profile better characterizes breast cancer tissue sections without costly experimentation. *Sci. Rep.* **12,** 4133 (2022).

59. Bergenstråhle, L. *et al.* Super-resolved spatial transcriptomics by deep data fusion. *Nat. Biotechnol.* (2021). doi:10.1038/s41587-021-01075-3

60. He, B. *et al.* Integrating spatial gene expression and breast tumour morphology via deep learning. *Nat. Biomed. Eng.* **4,** 827–834 (2020).

61. Bhatia, H. S. *et al.* Proteomics of spatially identified tissues in whole organs. *bioRxiv* (2021). doi:10.1101/2021.11.02.466753

62. Hughes, C. S. *et al.* Ultrasensitive proteome analysis using paramagnetic bead technology. *Mol. Syst. Biol.* **10,** 757 (2014).

63. Sielaff, M. *et al.* Evaluation of FASP, SP3 and iST Protocols for Proteomic Sample Preparation in the Low Microgram Range. *J. Proteome Res.* **16,** 4060–4072 (2017).

64. Sandow, J., Infusini, G., Dagley, L., Larsen, R. & Webb, A. Simplified high-throughput methods for deep proteome analysis on the timsTOF Pro. *bioRxiv* 657908 (2019). doi:10.1101/657908

65. Cox, J. *et al.* Accurate Proteome-wide Label-free Quantification by Delayed Normalization and Maximal Peptide Ratio Extraction, Termed MaxLFQ. *Mol. Cell. Proteomics* **13,** 2513–2526 (2014).

66. Tyanova, S., Temu, T. & Cox, J. The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. *Nat. Protoc.* **11,** 2301–2319 (2016).

67. Bivand, R. S. & Wong, D. W. S. Comparing implementations of global and local indicators of spatial association. *TEST* **27,** 716–748 (2018).

68. Bivand, R. S., Pebesma, E. J., Gomez-Rubio, V. & Pebesma, E. J. *Applied spatial data analysis with R*. **2,** (Springer, 2013).

69. Hijmans, R. J. raster: Geographic Data Analysis and Modeling. (2020).

70. Langfelder, P., Zhang, B. & Horvath, S. Defining clusters from a hierarchical cluster tree: The Dynamic Tree Cut package for R. *Bioinformatics* **24,** 719–720 (2008).

71. McInnes, L., Healy, J., Saul, N. & Großberger, L. UMAP: Uniform Manifold Approximation and Projection. *J. Open Source Softw.* **3,** 861 (2018).

72.   Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R Package for Comparing

Biological Themes Among Gene Clusters. *Omi. A J. Integr. Biol.* **16,** 284–287 (2012).

73.   Trede, D. *et al.* Exploring Three-Dimensional Matrix-Assisted Laser

Desorption/Ionization Imaging Mass Spectrometry Data: Three-Dimensional Spatial

Segmentation of Mouse Kidney. *Anal. Chem.* **84,** 6079–6087 (2012).

74.   Liebisch, G. *et al.* Update on LIPID MAPS classification, nomenclature, and shorthand

notation for MS-derived lipid structures. *J. Lipid Res.* **61,** 1539–1555 (2020).

Methods

Tissue retrieval and processing

Post-mortem brain tissue was retrieved by the Oxford Brain Bank; a research ethics committee (REC) approved and HTA regulated research tissue bank (REC reference 15/SC/0639). The retrieved brain was sectioned into 1 cm thick coronal sections starting at the level of the mammillary bodies. Due to its large size, tumour tissue was present within multiple of these coronal slices. The tumour tissue was dissected from the first coronal and second posterior coronal slices (P1 and P2). The tumour from the P2 slice was split into quadrants. Cryosections were taken from all pieces to determine the tissue block with the best morphological and cellular preservation. Cryosections were stained with H&E (see below) and examined by a Neuropathologist (OA). The upper-right quadrant from the P2 coronal slice was selected for use in further experiments.

Relevant tissue blocks of the AT/RT tumour were acclimatised to -20 °C and mounted onto a cryostat block using OCT Compound (Cell Path, ARG1180). Careful consideration was taken to ensure cut sections were not contaminated with OCT. Sections were cut at 10 µm and mounted onto UV irradiated (254 nm, 30 minutes) 1.0 PEN membrane slides (Zeiss) at -18 °C for LCM or Superfrost glass slides for histology. Sections were then air-dried for several minutes and placed onto a Shandon Linistain for automated H&E staining. Sections were fixed in 70 % denatured alcohol, hydrated, stained with Harris' Haematoxylin, incubated in 0.4 % acid alcohol, placed in Scot's tap water, and stained with Eosin containing 0.25 % acetic acid with regular washing steps in between. Stained sections were then dehydrated in increasing concentrations of denatured alcohol and air-dried without coverslips and stored at -80 °C until processing by laser-capture microdissection.

Laser capture microdissection

Areas of tissue analysed were annotated and isolated from the prepared slides using a laser-capture microscope equipped with laser pressure catapulting (PALM Microbeam, Zeiss). Cutting and capturing the annotated tissue areas were performed automatically and used the 10x objective lens. The settings in the control software for cutting were Energy: 43, Focus: 55; and for capturing were Energy 20, Focus -15. Samples were collected into 20 µL

1   RIPA buffer (Pierce #89900) in the cap of 200 µL PCR tubes or PCR-cap strips of 8. Collected

2   samples were immediately placed in dry ice. Samples were stored at -80 °C until further use.

Proteomic sample processing

Samples were thawed, incubated at room temperature for 30 minutes and briefly

centrifuged. Caps were rinsed with 20 µL of RIPA buffer (#89900, Pierce) containing 25 units

of Benzonase (E1014, Merck) to collect any remaining tissue and briefly centrifuged,

followed by incubation at room temperature for 30 minutes to degrade DNA and RNA.

Proteins were reduced by adding DTT to 5 mM and incubated at room temperature for 30

minutes, followed by the addition of iodoacetamide to 20 mM and incubation at room

temperature for 30 minutes.

Paramagnetic SP3 beads (GE45152105050250 & GE65152105050250, Cytiva) were prepared

as described by Hughes *et al.* and processed by a modified SP3 protocol [24,62,63]. Three µL of

SP3 beads were mixed with the samples, and acetonitrile added to a final concentration of

70 % (v/v). Samples were mixed with 1000 rpm orbital shaking for 18 minutes, followed by

bead immobilisation on a magnet for 2 minutes. The supernatant was discarded, and beads

were washed twice with 70 % (v/v) ethanol in water and once with 100 % acetonitrile

without removal from the magnet. Beads were resuspended in 50 mM ammonium

bicarbonate containing 25 ng of Trypsin (V5111, Promega) and digested overnight at 37 °C.

After digestion, the beads were resuspended by bath sonication. Acetonitrile was added to

the samples to 95 % (v/v) and shaken at 1000 rpm for 18 minutes. Beads were immobilised

on a magnet for 2 minutes, and the supernatant discarded. Beads were resuspended in 2 %

acetonitrile and immobilised on a magnet for 5 minutes. Peptides were transferred to glass

LC-MS vials or 96-well PCR plates containing formic acid in water, resulting in a final formic

acid concentration of 0.1 %.

LC-MS/MS

26   Peptides from 833 µm resolution samples were analysed by LC-MS/MS using a Dionex

27   Ultimate 3000 (Thermo Scientific) coupled to a timsTOF Pro (Bruker) using a 75 µm x 150

28   mm C18 column with 1.6 µm particles (IonOpticks) at a flow rate of 400 nL/min. A 17-

29   minute linear gradient from 2 % buffer B to 30 % buffer B (A: 0.1 % formic acid in water. B:

30   0.1 % formic acid in acetonitrile) was used[64]. The TimsTOF Pro was operated in PASEF mode.

The TIMS accumulation and ramp times were set to 100 ms, and mass spectra were recorded from 100 – 1700 m/z, with a 0.85 – 1.30 Vs/cm^2 ion mobility range. Precursors were selected for fragmentation from an area of the full TIMS-MS scan that excludes most ions with a charge state of 1+. Those selected precursors were isolated with an ion mobility dependent collision energy, increasing linearly from 27 – 45 eV over the ion mobility range. Three PASEF MS/MS scans were collected per full TIMS-MS scan, giving a duty cycle of 0.53 s. Ions were included in the PASEF MS/MS scan if they met the target intensity threshold of 2000 and were sampled multiple times until a summed target intensity of 10000 was reached. A dynamic exclusion window of 0.015 m/z by 0.015 Vs/cm^2 was used, and sampled ions were excluded from reanalysis for 24 seconds.

Peptides from 350 μm resolution samples were analysed by nano-UPLC-MS/MS using a Dionex Ultimate 3000 coupled to an Orbitrap Fusion Lumos (Thermo Scientific) using a 75 μm x 500 mm C18 EASY-Spray Columns with 2 μm particles (Thermo Scientific) at a flow rate of 250 nL/min. A 60-minute linear gradient from 2 % buffer B to 35 % buffer B (A: 5 % DMSO, 0.1 % formic acid in water. B: 5 % DMSO, 0.1 % formic acid in acetonitrile). MS1 scans were acquired in the Orbitrap between 400 and 1500 m/z with a resolution of 120,000 and an AGC target of 4 x 10^5. Precursor ions between charge state 2+ and 7+ and above the intensity threshold of 5 x 10^3 were selected for HCD fragmentation at a normalised collision energy of 28 %, an AGC target of 4 x 10^3, a maximum injection time of 80 ms and a dynamic exclusion window of 30 s. MS/MS spectra were acquired in the ion trap using the rapid scan mode.

Proteomic data analysis

Raw data files were searched against the UniProtKB human database (Retrieved 17/01/2017, 92527 sequences) using MaxQuant version 1.6.14.0, allowing for tryptic specificity with up to 2 missed cleavages. Cysteine carbamidomethylation was set as a fixed modification. Methionine oxidation and protein N-terminal acetylation were set as variable modifications and the "match between runs (MBR)" option was used (MBR was not used for tissue titration data). All other settings were left as default. Label-free quantification was performed using the MaxLFQ algorithm within MaxQuant[65,66]. Protein and peptide false discovery rate (FDR) levels were set to 1 %.

Spatial data analysis

The mapping of the mass spectrometry raw files to their relative spatial locations is shown in Supplementary Table 1. The spatial analysis uses functions within the spdep and raster R packages[67–69]. MaxQuant's protein level output files ('proteingroups.txt') were filtered to remove reverse hits, 'Only identified by site' hits and potential contaminants. The 'LFQ intensity' columns were $\log_2$ transformed and then normalised by median subtraction. Protein groups that did not meet a cut-off of having at least 9 pixels with normalised LFQ values are not taken forward for further analysis. The following steps occur independently for each protein group. Normalised LFQ intensities were then coerced into a matrix reflecting the rastered pattern of sample acquisition. The quantification matrix was converted into a raster object and then to a polygon object using the raster R package. From this polygon object, a neighbour list was built for each pixel of the raster using the 'Queen's Case' where cells are considered neighbours if they share an edge or a vertex. The neighbour list was then supplemented with a spatial weights matrix using a binary coding scheme where neighbours are given a weighting of '1' and non-neighbours a weighting of '0' in the spatial weights matrix. The raster object and the weighted neighbour list were then used as inputs to a permutation test for the Moran's *I* statistic, calculated using 999 random spatial permutations of the raster object to calculate pseudo-*p*-values. Moran's *I* statistics and the associated *p*-values are collected for every protein group. The *p*-values were then corrected for multiple testing using the Benjamini-Hochberg FDR method.

Immunohistochemistry

Sections were cut as above and were mounted to Superfrost glass slides for IHC and air-dried. Slides were fixed in ice-cold acetone for 10 minutes, washed twice with TBS/T (20 mM Tris, 150 mM NaCl, 0.05 % Tween 20) and blocked with 10 % goat serum in TBS/T for 60 minutes at room temperature. Primary antibodies were diluted in 5 % goat serum in TBS/T and incubated at RT for 60 minutes or 4 °C overnight. Sections were washed three times with TBS/T. Staining visualisation was performed by incubating with a cocktail of anti-mouse and anti-rabbit secondary antibodies conjugated to horseradish peroxidase (Envision Kit, Agilent) for 60 minutes at room temperature. Sections were then washed with TBS/T three times and incubated with 2 % 3,3'-diaminobenzidine for 5 minutes, immersed in water and then counterstained with Harris' Haematoxylin for 1 minute. Primary antibodies used and

1 dilutions: rabbit anti-PYGL, 1:100, 4 °C overnight, HPA000962 (Atlas Antibodies); rabbit anti-

2 ASPH, 1:1000, RT 60 minutes, NBP2-34125 (Novus Biologicals); mouse anti-CD45 (PD7/26 +

3 2B11), 1:200, RT 60 minutes, ab781 (Abcam).

Clustering

Empty pixels and pixels covering the large region of haemorrhage were not included. A

distance matrix was built containing the Euclidean distance between each pixel's set of

protein LFQ values. Hierarchical clustering of the distance matrix was performed in R using

the "average" agglomeration method. Dendrograms were cut using the Dynamic Tree Cut

method at a height setting of 100[70].

A complete input matrix is required for UMAP visualisation[71], so proteins with fewer than 70

% valid values across the experiment were removed. The remaining missing values were

imputed in on a per-sample basis by random draws from a normal distribution using a width

of 0.3 and a downshift of 1.8. UMAP dimensionality reduction was performed on this

imputed data with default settings, and the first two embedding components plotted, and

samples coloured according to their cluster assigned by Dynamic Tree Cut at a height of 100.

Pathway Analysis

A one-way ANOVA test was performed in R to test for differences in means between the

clusters generated by Dynamic Tree Cut at a height of 100. A pairwise post-hoc correction

was applied using Tukey's Honestly Significant Difference method. The resulting pairwise

comparisons were used as inputs to ClusterProfiler's gene set enrichment analysis to test for

enrichment of Gene Ontology Biological Process terms using a Fisher's exact test with a 5 %

FDR threshold[72].

Lipid MALDI imaging

Vacuum dried sections were scanned with a TissueScout scanner at 3200 dpi (Bruker) to

generate a reference image for later position teaching. Dihydroxybenzoic acid (DHB) was

dissolved at a concentration of 15 mg/ml in 90 % ACN, 0.1 % TFA and sprayed on top of the

AT/RT sections using a TM-sprayer (HTX Technologies). The matrix was applied in a criss-

cross pattern with 3 mm track spacing at a 1200 mm/min nozzle velocity. Fourteen layers

were sprayed at a 0.125 ml/ml flow rate using 50 % ACN as the liquid phase at 10 psi

pressure. The nozzle temperature was set to 60 °C and the distance of the nozzle to the

section was 40 mm. All imaging data were acquired on a timsTOF fleX instrument (Bruker)

which is equipped with a dual ESI and MALDI source in positive Q-TOF mode. External

calibration was performed using red phosphorous which was spotted next to the section.

The laser was operated in beam scan mode, ablating an area of 15x15 μm resulting in a pixel

size of 20 μm. The repetition rate of the laser was set to 10 kHz and 400 laser shots were

acquired per pixel. Data were acquired in the mass range 300-1400 m/z.

The software SCiLS Lab (version 2020a; Bruker) was used for MALDI Imaging data analysis.

All data were root-mean-square (RMS) normalised. After importing the data, an

unsupervised segmentation was calculated using the bisecting k-means algorithm and a

peak list containing 302 m/z-intervals with correlation used as a distance metric[73]. The

resulting segmentation map was split into several clusters that resemble the histopathology

of the tumour section. The identification of cholesterol ester, observed as 689.5607 and

673.5843 m/z ions, which correspond to the m/z adducts of cholesterol ester (18:1) $[M+K]^+$

and $[M+Na]^+$, was performed using LIPID-MAPS database searches using exact mass[74].

Expected m/z were 689.5633 ($M+K^+$), 673.5894 ($M+Na^+$) and delta masses were 0.0026

(3.7ppm) and 0.0051 m/z (7.5ppm), respectively. The next closest match corresponding to a

species with both Na^+ and K^+ adducts is DG 38:1 with delta masses of 0.0102 ($M+Na^+$) and

0.0126 ($M+K^+$) m/z.

Affinity Network Fusion

For the network fusion approach, we used a minimally processed expression matrix, log2

transformed and median centred, removing all empty pixels. We selected the top 25% of

proteins by variance across all pixels and used this to calculate a proteomic similarity matrix

of Spearman's rank correlation coefficients. For the focussed extracellular matrix protein

analysis, all detected proteins annotated as "Core Matrisome" in MatrisomeDB were used

to calculate the protein similarity matrix. We converted this to a proteomic distance matrix

(by taking 1-similarity). Separately, we created a complementary spatial distance matrix

representing the Euclidean distance from each pixel location to each other pixel location

(where horizontally and vertically adjacent neighbours are distance 1, diagonal neighbours

are distance √2 and so on). We converted both matrices into affinity matrices and fused

them using Affinity Network Fusion[38]. We then performed spectral clustering on the fused

affinity matrix, where the number of clusters (6) was selected using the maximal eigengap

heuristic.

Data availability

4   The mass spectrometry proteomics data have been deposited to ProteomeXchange

5   Consortium via the PRIDE partner repository and will be made available to reviewers upon

6   submission.

Figures

Figure 1 – Overview of the spatially resolved proteomics workflow
Tissue is segmented into a regular grid shape, and each element of the grid is isolated by
laser capture microdissection (LCM). Proteins from each individually lysed sample are
digested before analysis by LC-MS/MS. The quantitative information for each protein can be
mapped back to its location within the gridded tissue and visualised in a topographic protein
map, with one map per protein quantified.

Figure 2 – Spatial proteomic maps of AT/RT tumour tissue reveal regional boundaries
Normalised protein intensity of four example proteins mapped back to the original spatial
positions within the atypical teratoid-rhabdoid tumour (AT/RT) tumour tissue at a resolution
of 833 μm (A) and 350 μm (B) with their corresponding Moran's Index of spatial
autocorrelation (*I*). Box in (A) represents the area analysed in an adjacent tissue section (B).
Scale bar = 1 mm. Normalised protein intensities are scaled separately for each protein.
Grey = not detected.

Figure 3 - Immunohistochemistry validation of AT/RT proteomic maps
Proteomic maps of proteins targeted for follow-up IHC staining and IHC images. (A,C,E)
Normalised protein intensity maps with their corresponding Moran's Index of spatial
autocorrelation (*I*). Normalised protein intensities are scaled separately for each protein.
Grey = not detected. Rectangles depict the approximate location displayed in IHC images.
(B,D,F) AT/RT tissue stained and visualised by IHC. All scale bars = 1 mm.

Figure 4 - Clustering and functional analysis displays regional AT/RT proteomic maps in the
tumour microenvironment
(A) Map of cluster assignment based on hierarchical clustering and the dynamic tree cut
algorithm. This clustering is spatially-unaware and is based only on protein abundance
values. White pixels represent excluded empty regions and regions of haemorrhage (B)
UMAP embedding of data coloured by cluster assignment in (A). (C) GSEA of clusters 1 & 3
from (A). Gene set membership is indicated by colouring the cell with that protein's $\log_2$
fold-change. (D) Cluster-map of ATRT tissue generated by affinity network fusion (ANF). This
clustering is spatially-aware, meaning the relative spatial locations of samples and protein
abundance values are used to generate these clusters. (E) Enriched MSigDB Hallmark gene
sets within marker proteins of the clusters shown in the ANF cluster-map. Significantly
enriched hallmarks for each cluster are indicated by the presence of circles. The size and

1   colour of the circles represent the number of proteins contributing to that term and the

2   adjusted *p* value of the enrichment, respectively.

Figure 5 - High-resolution proteomic map clustering defines functional layers at the
tumour periphery
(A) Map of cluster assignment based on hierarchical clustering and the dynamic tree cut
algorithm. Cluster 3 corresponds to brain/tumour interface. Cluster 1 corresponds to solid
tumour. Clusters 2 & 5 correspond to the transition between solid tumour and
brain/tumour interface (B) Volcano plot of clusters 3 against cluster 1 from (A). The
horizontal dashed line represents an FDR of 5 %. The vertical dashed line represents -/+ 2-

1   fold-change. (C) GSEA of clusters 1 & 3 from (A). Gene set membership is indicated by
2   colouring the cell with that protein's $\log_2$ fold-change.

2
3
Figure 6 – Spatial heterogeneity of extracellular matrix proteins across the ATRT tumour
(A) Cluster-map of ATRT tissue generated by affinity network fusion (ANF) of core
matrisome proteins as defined in MatrisomeDB. This clustering is spatially-aware, meaning
the relative spatial locations of samples and protein abundance values are used to generate
these clusters. The grey pixel represents one sample where no core matrisome proteins
were detected. (B) Enriched MSigDB Hallmark gene sets within marker proteins of the
clusters shown in the ANF cluster-map. Significantly enriched hallmarks for each cluster are
indicated by the presence of circles. The size and colour of the circles represent the number
of proteins contributing to that term and the adjusted *p* value of the enrichment,
respectively. (C) Proteomic maps of selected extracellular matrix proteins and integrin
receptors at 833 μm resolution. Normalised protein intensities are scaled separately for
each protein. Grey = not detected

1 Supplementary Figures

AT/RT H&E

	Side Length (μm)	Area (μm²)	cells
$10^{2.5}$	18	316	2.7
10^{3}	32	1,000	8.6
$10^{3.5}$	56	3,162	27
10^{4}	100	10,000	86
$10^{4.5}$	178	31,623	270
10^{5}	316	100,000	860
$10^{5.5}$	562	316,228	2700
10^{6}	1000	1,000,000	8600

3 **Figure S1 – Tissue area titration across two LC-MS/MS systems**
4 The number of proteins identified from a titration of AT/RT tissue area on two LC-MS/MS
5 systems.
6

2 **Figure S2 – Distribution of identified and quantified proteins at 833 μm resolution**
3 (A) Violin plots showing distributions of identified and quantified proteins per pixel. Solid
4 vertical lines represent the median value. Dashed vertical lines represent upper and lower
5 quartiles. (B) Histogram of quantified pixels per protein map.
6

A

B

Figure S3 – Aggregate intensity distributions
Maps of the $\log_2$ transformed (A) summed and (B) mean intensities of each pixel.

Figure S4 – Proteomic maps of immune cell markers

Proteomic maps of immune cell-marker proteins at 833 µm resolution. Normalised protein intensities are scaled separately for each protein. Grey = not detected.

Figure S5 – Pathway enrichment in immune-cell clusters

GSEA of clusters 6 & 13 from Figure 4A. Gene set membership is indicated by colouring the cell with that protein's log2 fold-change.

Figure S6 – AT/RT brain tissue MALDI imaging shows regional enrichment of haeme and lipids at 20 μm resolution.
(Left) Image of haematoxylin and eosin-stained tissue after MALDI acquisition. (Middle-left) MALDI Imaging pixels were clusters by Bisecting K-Means clustering and the generated clusters are indicated by colour. m/z images for ions corresponding to the masses of haeme (middle), cholesterol ester (18:1) [M+K]+ (middle-right) and [M+Na]+ (right) ions. Ion intensities are scaled separately for each ion.

A

B

Figure S7 – Aggregate intensity distribution of core matrisome proteins
Maps of the log_2 transformed (A) summed and (B) mean intensities of core matrisome
proteins as defined by MatrisomeDB for each pixel. Grey pixels indicate no core matrisome
proteins were detected.

2 **Figure S8 – Distribution of Moran's I in matrisome and non-matrisome proteins**
3 Moran's *I* distribution of proteins annotated in Matrisome DB as core matrisome (blue),
4 matrisome associated (yellow), and all other proteins (grey).