1

Lipolysis of bone marrow adipocytes is required to fuel bone and the marrow niche during energy deficits

Ziru Li[1], Emily Bowers[2], Junxiong Zhu[3,4], Hui Yu[1], Julie Hardij[1], Devika P. Bagchi[1], Hiroyuki Mori[1], Kenneth T. Lewis[1], Katrina Granger[1], Rebecca L. Schill[1], Steven M. Romanelli[1], Simin Abrishami[2], Kurt D. Hankenson[3], Kanakadurga Singer[1,2], Clifford J. Rosen[5], and Ormond A. MacDougald[1,6]*

Affiliations:

1. University of Michigan Medical School, Department of Molecular & Integrative Physiology, Ann Arbor, MI

2. University of Michigan Medical School, Department of Pediatrics, Ann Arbor, MI

3. Department of Orthopedic Surgery, University of Michigan Medical School, Ann Arbor, MI

4. Department of Orthopedic Surgery, The Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, China.

5. Maine Medical Center Research Institute, Scarborough, ME

6. University of Michigan Medical School, Department of Internal Medicine, Ann Arbor, MI

*Correspondence to: Ormond A. MacDougald

NCRC Building 25, Rm. 3686

2800 Plymouth Rd., Ann Arbor, MI 48109

Tel:(734) 647-4880

E-mail: macdouga@umich.edu

Summary

To investigate roles for bone marrow adipocyte (BMAd) lipolysis in bone homeostasis, we

created a BMAd-specific Cre mouse model in which we knocked out adipose triglyceride lipase

(ATGL, *Pnpla2*). BMAd-*Pnpla2*$^{-/-}$ mice have impaired BMAd lipolysis, and increased size and

number of BMAds at baseline. Although energy from BMAd lipid stores is largely dispensable

when mice are fed *ad libitum*, BMAd lipolysis is necessary to maintain myelopoiesis and bone

mass under caloric restriction. BMAd-specific *Pnpla2* deficiency compounds the effects of

caloric restriction on loss of trabecular bone, likely due to impaired osteoblast expression of

collagen genes and reduced osteoid synthesis. RNA sequencing analysis of bone marrow

adipose tissue reveals that caloric restriction induces dramatic elevations in extracellular matrix

organization and skeletal development genes, and energy from BMAd is required for these

adaptations. BMAd-derived energy supply is also required for bone regeneration upon injury,

and maintenance of bone mass with cold exposure.

Keywords: bone marrow adipocyte (BMAd), bone marrow adipose tissue (BMAT), lipolysis,

adipose triglyceride lipase (ATGL), caloric restriction (CR), energy supply, hematopoiesis, bone

Introduction

Bone marrow adipocytes (BMAds) are a heterogeneous cell population that form depots of bone marrow adipose tissue (BMAT) distinct from white, brown, and other adipose tissues (Z. Li, Hardij, Bagchi, Scheller, & MacDougald, 2018). Clinical associations generally demonstrate inverse relationships between BMAT and bone mass (Shen et al., 2007), or BMAT and circulating immune cells (Polineni et al., 2020), which may be due to the interactions between cells within the bone marrow niche. In addition to BMAds, the bone marrow niche contains osteoblasts, osteoclasts, hematopoietic cells, stromal/mesenchymal cells, blood vessels, and nerves (Vogler & Murphy, 1988). The relationships between BMAds, bone cells, and hematopoietic cells are influenced by their shared location within bone, an anatomically restricted system, such that expansion of one cell type is by necessity at the expense of others. For example, elevated BMAT is negatively correlated with low bone mass of aging and diabetes, whereas expansion of BMAT is associated with multiple hematopoietic disorders (Z. Li & MacDougald, 2021). Mechanistic links underlying these associations have proven challenging to investigate because they often involve complex intercellular, endocrine, and/or central mechanisms.

In addition to serving as an energy source, BMAds potentially influence the marrow niche through cell-to-cell contact, release of extracellular vesicles, and secretion of adipokines (e.g adiponectin) and cytokines (e.g. stem cell factor). Removal of these stimuli in mouse models of lipodystrophy, which lack BMAT, results in increased bone mass (Corsa et al., 2021; Zou et al., 2020; Zou et al., 2019). However, use of these models to investigate the direct effects of BMAT depletion is confounded by concurrent loss of white and brown adipose depots, which also regulate bone mass through myriad mechanisms, including secretion of adipokines (Riddle & Clemens, 2017; Zou et al., 2019). Although loss of BMAT in lipodystrophic mice was integral to the original finding that BMAT is a negative regulator of

hematopoiesis, positive effects of BMAT on hematopoiesis have also been observed; these

differences are attributed to use of distinct animal models and analysis of different skeletal

sites (Ambrosi et al., 2017; Naveiras et al., 2009; Zhou et al., 2017). In rodents, there are two

readily identifiable BMAd populations: constitutive BMAT (cBMAT) and regulated BMAT

(rBMAT) (Scheller et al., 2015). cBMAT typically exists in distal tibia and caudal vertebrae,

appears early in life, and has the histological appearance of white adipose tissue. rBMAT is

found in proximal tibia and distal femur, and is comprised of single or clustered BMAds

interspersed with hematopoietic cells. Whereas cBMAT generally resists change in response

to altered physiological states, rBMAT expands with aging, obesity, diabetes, caloric

restriction (CR), irradiation, and estrogen deficiency, and is reduced by cold exposure,

fasting, β3-agonist, exercise, and vertical sleeve gastrectomy (Z. Li et al., 2018; Z. Li et al.,

2019). These treatments also cause alterations to the skeleton and/or formation of blood

cells, some of which may be secondary to effects on BMAds.

BMAds have long been believed to fuel maintenance of bone and hematopoietic

cellularity because of their shared physical location in the marrow niche. However, this

hypothesis has not been formally tested because current methods to target BMAds lack

penetrance or cause recombination in other cell types, such as white adipocytes, osteoblasts,

or bone marrow stromal cells, complicating the interpretation of interactions between BMAds

and cells of the marrow niche. To circumvent this problem, we created a BMAd-specific Cre

mouse model based on expression patterns of endogenous *Osterix* and *Adipoq*, and

investigated roles for BMAds as a local energy source by deleting ATGL/*Pnpla2*, the rate-

limiting enzyme of lipolysis. Consistent with *Pnpla2* deficiency in white and brown adipocytes

(Ahmadian et al., 2011), BMAd-*Pnpla2*$^{-/-}$ mice have impaired BMAd lipolysis, resulting in

hypertrophy and hyperplasia of BMAT. Despite significant increases in bone marrow

adiposity, hematopoietic abnormalities of BMAd-*Pnpla2*$^{-/-}$ mice are negligible under basal

conditions. However, the recovery of bone marrow myeloid lineages following sublethal

irradiation is impaired with CR, and further reduced by BMAd-*Pnpla2* deficiency. Similarly,

proliferative capacity of myeloid progenitors is also decreased with CR, and further inhibited

in mice lacking BMAd-*Pnpla2*. Whereas alterations in bone parameters were not observed in

ad libitum fed BMAd-*Pnpla2*$^{-/-}$ mice, bone loss occurred under conditions of elevated energy

needs such as bone regeneration or cold exposure, or reduced energy availability such as

CR. Reduction of bone mass in CR BMAd-*Pnpla2*$^{-/-}$ mice is likely due to impaired osteoblast

functions such as expression of extracellular matrix genes and creation of osteoid. Gene

profiling reveals that *Pnpla2* deletion largely blocks CR-induced genes within pathways of

extracellular matrix organization and skeletal development, indicating that BMAd-derived

energy contributes to skeletal homeostasis under conditions of negative energy balance.

Results

Generation of a BMAd-specific Cre mouse model (BMAd-Cre). Based on previous

studies showing that *Osterix* traces to osteoblasts and BMAds, but not to white adipocytes

(Chen & Long, 2013; Mizoguchi et al., 2014), we used CRISPR/Cas9 to create *Osterix-FLPo*

mice with an in-frame fusion of *Osterix* and optimized *FLPo*, separated by a *P2A* self-

cleaving sequence to allow independent functioning of the two proteins (Figure 1A). To

validate tissue-specific expression and FLPo efficiency, we bred *Osterix-FLPo* mice to FLP-

dependent EGFP reporter mice, and observed EGFP-positive osteocytes, osteoblasts,

BMAds, and a subset of marrow stromal cells within the bone (Figure 1B and 1C).

We next created F<u>LPo</u>-dependent <u>A</u>dipoq-<u>C</u>re (FAC) mice, which contain an internal

ribosome entry sequence (IRES) followed by FLPo-dependent *Cre* in reverse orientation

within the 3'-untranslated region (UTR) of endogenous *Adipoq* gene (Figure 1D). FLPo

expressed from the *Osterix* locus recombines *Cre* to the correct orientation in progenitors of

osteoblasts and BMAds. However, because *Adipoq* is selectively expressed in adipocytes

(Eguchi et al., 2011), Cre is expressed in BMAds, but not in osteoblasts or other adipose

depots. Consistent with this schema, *Cre* in the correct orientation (flipped band, Figure 1 -

figure supplement 1A-1C) is only observed in caudal vertebral DNA of mice positive for at

least one copy each of *Osterix* (Mut band) and FAC (Ori band). The correct insertion of the

sequence in *Adipoq* 3'-UTR was validated by Sanger sequencing following genomic PCRs

that spanned endogenous *Adipoq* sequences, homology arms, IRES and Cre (Figure 1 -

figure supplement 1D). To investigate the cell type-specificity of Cre activity, we next bred

BMAd-Cre mice to mT/mG reporter mice (Muzumdar, Tasic, Miyamichi, Li, & Luo, 2007)

(Figure 1E), in which all tissues and cells express red fluorescence (membrane-targeted

tdTomato; mT) at baseline, and will express membrane-targeted EGFP in the presence of

cell-specific Cre. We observed loss of tdTomato and gain of EGFP in BMAT depots of caudal

vertebrae (tail) and tibiae, but not in brown or white adipose depots, or other tissues/organs

such as liver, pancreas, muscle and spleen (Figure 1E and Figure 1 - figure supplement 1E).

The correct orientation of *Cre* (flipped band) was also observed in mRNA isolated from distal

tibiae and caudal vertebrae, but not WAT depots, of *Osterix-FLPo* mice positive for FAC

(Figure 1 - figure supplement 1F). To evaluate conditions optimal for Cre-induced

recombination, we visualized conversion of tdTomato to EGFP in *BMAd-Cre* mice at various

ages and with different FAC copy numbers (Figure 1 - figure supplement 1G-1H) and found

that the proportion of EGFP-positive BMAds increases with age and number of FAC alleles.

Indeed, Cre efficiency is ~80% in both male and female mice over 16 weeks of age with one

Cre allele, and over 90% at 12 weeks of age in mice with two Cre alleles.

Previous studies found that a randomly inserted *Adipoq-Cre* bacterial artificial

chromosome (BAC) causes recombination in osteoblasts (Bozec et al., 2013; Eguchi et al.,

2011; Mukohira et al., 2019); thus, we tested whether this observation is true in *BMAd-Cre*

mice, which rely on endogenous *Adipoq* expression that is restricted by *Osterix*. When

BMAd-Cre mice are bred to mT/mG reporter mice, only tdTomato-positive cells are

detectable on trabecular bone surfaces (Figure 1F), suggesting that osteoblasts are not

targeted. However, labeling of a small subset of stromal/dendritic cells is consistently

observed, likely due to expression of *Adipoq* within this cell population (Mukohira et al.,

2019).

Although insertion of the FLPo-activated Cre cassette into the 3'-UTR of endogenous

Adipoq overcomes potential problems that arise from random genomic integration, we

considered the possibility that placement of the IRES-Cre cassette within the 3'-UTR might

alter the expression and/or secretion of adiponectin. Although insertion of the IRES-Cre

cassette may cause a slight reduction in mRNA expression in whole bone, the adiponectin

mRNA in white adipose tissue is elevated by more than 3 fold (Figure 1 - figure supplement

2A). Thus, the cassette itself does not limit expression of mRNA. Instead, it appears that the

2 kb IRES-Cre cassette within the 3'-UTR impairs translatability of the mRNA since

expression of adiponectin protein in BMAT and WAT is decreased by ~50% (Figure 1 - figure

supplement 2B and 2C). Circulating adiponectin concentrations appear to be decreased even

further (Figure 1 - figure supplement 2D-2F), perhaps suggesting that flux of translated

adiponectin protein is impaired through the secretion pathway. Of note, hypoadiponectinemia

is associated with insulin resistance (Cook & Semple, 2010; N. Li et al., 2021), with

discrepant reports on osteogenesis (Lewis, Edwards, Naylor, & McGettrick, 2021). Thus, we

evaluated systemic metabolism in BMAd-Cre mice and found that body weight, glucose

tolerance, WAT depot weights, and tibial trabecular and cortical bone variables are

unaffected by this degree of hypoadiponectinemia (Figure 1 - figure supplement 2G-2P). To

minimize variability between treatments, all mice used in the following studies were positive

for both *Osterix-FLPo* and FAC. Gene knockout mice and their controls were determined by

the presence or absence of floxed alleles, respectively.

Ablation of adipose triglyceride lipase (ATGL, *Pnpla2*) causes BMAT expansion. A

fundamental function of adipocytes is to store excess energy as triacylglycerols and to

release non-esterified fatty acids and glycerol during times of negative energy balance. ATGL

is the first and rate-limiting enzyme in the lipolytic process; thus, to determine the

physiological functions of BMAd lipolysis in bone metabolism and hematopoiesis, we

generated BMAd-*Pnpla2*$^{-/-}$ mice, in which the gene encoding ATGL, *Pnpla2*, is specifically

knocked out in BMAds. We validated the deletion of ATGL in caudal vertebrae, because

cBMAT is abundant in this location. Mutant *Pnpla2* mRNA is observed in caudal vertebrae of

BMAd-*Pnpla2* mice, but not in WAT (Figure 2A). Possible sources for the remaining wildtype

(WT) *Pnpla2* signal include periosteal WAT, non-adipocyte cells, or perhaps from incomplete

deletion of *Pnpla2* in BMAds. Immunofluorescent staining confirms loss of ATGL in proximal

tibial rBMAds of BMAd-*Pnpla2*$^{-/-}$ mice (Figure 2B). ATGL protein is reduced substantially in

caudal vertebrae, but not in subcutaneous WAT (Figure 2C), which further confirms

specificity of BMAd-Cre recombinase for BMAT.

To quantify effects of *Pnpla2*-deficiency on bone marrow adiposity and cellularity, we

used osmium tetroxide, and hematoxylin and eosin staining to evaluate BMAT quantity and

cellular details, respectively. We found that BMAT volume is significantly increased in

proximal tibiae and throughout the endocortical compartment of BMAd-*Pnpla2*$^{-/-}$ mice,

whereas differences in BMAT volume are not observed in distal tibiae, which is almost

completely occupied by BMAT in WT mice (Figure 2D-2F). Histological analyses reveal

increased BMAd number in proximal tibiae of BMAd-*Pnpla2*$^{-/-}$ mice (Figure 2G and 2H).

Depletion of *Pnpla2* also causes BMAd hypertrophy in both proximal and distal tibia, with an

increased proportion of BMAds larger than 500 μm^2 observed in proximal tibia, and larger

than 1000 μm^2 observed in distal tibia (Figure 2I and 2J). Secretion of basal glycerol and

non-esterified fatty acids (NEFA) from *ex vivo* cultured explants of distal tibial BMAT was not

different between genotypes. Whereas forskolin treatment greatly increased lipolysis in

BMAT explants from control mice, secretion of glycerol and NEFA from BMAd-*Pnpla2*$^{-/-}$

explants remained unchanged from basal rates (Figure 2K and 2L). These results confirm

that BMAds of BMAd-*Pnpla2*$^{-/-}$ mice have impaired lipolysis.

BMAd lipolysis is required to maintain bone homeostasis in male mice under

conditions of CR, but not when mice are fed *ad libitum*. We next evaluated whether loss

of BMAd lipolysis in male BMAd-*Pnpla2*$^{-/-}$ mice is sufficient to influence systemic physiology,

or function of bone cells within the marrow niche. In mice fed normal chow *ad libitum*, we did

not observe differences in body weight, glucose tolerance, or weights of soft tissue, including

subcutaneous WAT (sWAT), epididymal WAT (eWAT) and liver (Figure 3 - figure supplement

1A-1E), which further confirms the tissue-specificity of our BMAd-knockout model. As

observed above, *Pnpla2* deficiency causes expansion of proximal tibial rBMAT (Figure 3 -

figure supplement 1F and 1G). Interestingly, when dietary energy is readily available,

trabecular bone volume fraction, bone mineral density and trabecular number tend to be

lower in BMAd-*Pnpla2*$^{-/-}$ mice, but no significant differences were observed (Figure 3A-3C).

These data also indicate that expansion of BMAT is not sufficient to cause bone loss, and

that the correlation between elevated fracture risk and expansion of BMAT under a variety of

clinical situations is not necessarily a causal relationship.

Our experiments in *ad libitum* fed mice suggest that energy released from BMAds is

dispensable for bone cell functioning when dietary energy is plentiful, which is supported by

the comparable concentrations of glycerol and NEFA in circulation and in bone marrow

supernatant in BMAd-*Pnpla2*[+/+] and BMAd-*Pnpla2*[-/-] mice (Figure 3 - figure supplement 1H

and 1I). To determine whether male BMAd-*Pnpla2*[-/-] mice have impaired bone homeostasis

when dietary energy is limited, we next challenged mice with a 30% CR for six weeks. BMAd-

Pnpla2[-/-] mice did not have altered body weight, glucose tolerance, or tissue weights

comparing with their controls (Figure 3 - figure supplement 1J-1N), suggesting that BMAds

are not a critical source of circulating energy under these conditions. Although *ad libitum* fed

BMAd-*Pnpla2*[-/-] mice have increased rBMAT compared to controls (Figure 3 - figure

supplement 1F and 1G), CR stimulates rBMAT expansion in control mice such that levels are

comparable between genotypes (Figure 3 - figure supplement 1O and 1P). Interestingly,

despite this similarity in BMAT volumes, BMAd-*Pnpla2*[-/-] mice fed CR diet have reduced

trabecular bone volume fraction, connective density, and bone mineral density when

compared to controls (Figure 3E). This reduction in bone mass is due to decreased

trabecular bone number and increased trabecular spacing, without effects on trabecular

thickness (Figure 3D-3F). BMAds are a source of NEFA for vicinal osteoblasts (Maridas et

al., 2019); thus, the reduction in trabecular bone in CR BMAd-*Pnpla2*[-/-] mice likely results

from impaired osteoblast function due to insufficient energy available in the bone marrow

niche. Consistent with this notion, we observed a trend ($P = 0.07$) towards reduced NEFA

concentrations in bone marrow supernatant of CR BMAd-*Pnpla2*[-/-] mice, whilst circulating

glycerol and NEFA were not different between genotypes (Figure 3 - figure supplement 1Q

and 1R). Of note, changes in cortical bone area and thickness were not observed in BMAd-

Pnpla2[-/-] mice with either *ad libitum* or CR diet (Figure 3 - figure supplement 2A-2D), perhaps

because cortical bone surfaces are less active.

To investigate mechanisms underlying trabecular bone loss in CR BMAd-*Pnpla2*[-/-] mice,

we next measured circulating markers of bone cell activity and performed histomorphometry.

Osteoblast and osteoclast numbers are not changed by BMAd-*Pnpla2* deficiency under *ad*

libitum conditions (Figure 3 - figure supplement 2E and 2F). Although circulating markers of

bone formation (P1NP) and osteoclast activation (RANK Ligand) are not altered in BMAd-

Pnpla2$^{-/-}$ mice, markers for bone resorption (CTX-1) and osteoclast activation (TRACP5b) are

decreased (Figure 3 - figure supplement 2G-2J). Although with CR, osteoblast numbers are

not altered by a deficiency of BMAT lipolysis, the bone formation marker P1NP is decreased

(Figure 3G and 3I), suggesting that osteoblast functions are impaired. Osteoclast numbers

are increased in CR BMAd-*Pnpla2*$^{-/-}$ mice, but the osteoclast surface per bone surface

(Oc.S/BS) is not changed and osteoclast activation marker, TRACP5b, is decreased (Figure

3H and 3J), suggesting the osteoclast functions are inhibited. No differences were observed

in RANK Ligand and CTX-1 with CR (Figure 3 - figure supplement 2K and 2L). We then

visualized and quantified osteoid thickness with Goldner's trichrome staining and observed a

trend towards thinner osteoid layers without affecting osteoid surface (Figure 3K and Figure 3

- figure supplement 2M), which could be secondary to impaired secretion of collagen matrix

by osteoblasts or to enhanced bone mineralization. Further, we injected mice with calcein to

label those bone surfaces undergoing active mineralization. Dynamic histomorphometry data

suggests that BMAd-*Pnpla2*$^{-/-}$ mice have less bone-forming surface, as indicated by reduced

single-labelled surface and mineral surface (Figure 3L). There were no differences in double-

labelled bone surface, inter-label width, mineral apposition rate or osteoid maturation time

(Figure 3M and Figure 3 - figure supplement 2N), suggesting that bone mineralization is not

affected by loss of BMAd lipolysis. Taken together, these data support a model in which

reduced trabecular bone in BMAd-*Pnpla2*$^{-/-}$ mice is likely due to impaired ability of osteoblasts

to secrete osteoid.

BMAd lipolytic deficiency impairs myelopoiesis during regeneration. A single rBMAd

connects to almost 100 hematopoietic cells (Robles et al., 2019), suggesting that BMAds

potentially serve as important local energy sources for hematopoiesis. However, when fed *ad*

libitum, male BMAd-*Pnpla2*$^{-/-}$ mice do not exhibit differences in mature blood cells

populations, as assessed by complete blood cell counts (Table 1) and bone marrow flow

cytometry for hematopoietic populations (Table 2 and Figure 4 - figure supplement 1A and

1B). Although most hematopoietic stem and progenitor cells (HSPCs) do not depend on

BMAd lipolysis, reduced numbers of granulocyte-monocyte progenitors (GMP) are observed

in marrow of BMAd-*Pnpla2*$^{-/-}$ mice fed *ad libitum* (Table 2). Of note, with CR, reduced

numbers of circulating and bone marrow mature neutrophils are observed in BMAd-*Pnpla2*$^{-/-}$

mice, although GMP numbers, which are the progenitor population, are not altered in CR

BMAd-*Pnpla2*$^{-/-}$ mice (Tables 1 and 2), perhaps due to impaired maturation of granulocytes.

These data suggest that although myeloid cell defects are observed in BMAd-*Pnpla2*$^{-/-}$ mice,

other hematopoietic cell populations can metabolically compensate for the lack of BMAd

lipolytic products.

To investigate further whether hematopoiesis is dependent upon BMAd lipolysis, we

administered a sublethal dose of whole-body irradiation, and evaluated hematopoietic cell

recovery. Following irradiation, white and red blood cell depletion and recovery were

monitored by complete blood cell counts every 2-3 days (Figure 4 - figure supplement 1C-

1E). In *ad libitum* fed mice, HSPCs (Figure 4 - figure supplement 1F and 1G) and

mature/immature hematopoietic cells (Figure 4A-4H) are not influenced by deficiency of

BMAd-lipolysis. However, CR alone decreases bone marrow cellularity (Figure 4A), and

HSPC (Figure 4 - figure supplement 1F and 1G) and neutrophil numbers, without affecting

monocytes and lymphocytes (Figure 4B-4E). In CR BMAd-*Pnpla2*$^{-/-}$ mice, HSPCs are not

altered compared to CR controls, but bone marrow cellularity, monocytes and neutrophils are

decreased further. Preneutrophils and immature neutrophils are also reduced by CR, and

show additional decline in CR mice lacking BMAd-*Pnpla2* (Figure 4F-4G), but mature

neutrophils are only affected by CR (Figure 4H). These reductions in monocytes and

neutrophils suggest that BMAd-lipolysis is required for myeloid cell lineage regeneration

when energy supply from circulation is limited. CFU assays, which are optimized for the

growth of myeloid progenitor cells (CFU-Granulocytes, CFU-Macrophages and CFU-GM),

demonstrate fewer granulocyte progenitor colonies (CFU-G) from CR BMAd-*Pnpla2*[-/-] mice

(Figure 4I). The proliferative capacity of macrophage progenitors (CFU-M) is impaired when

derived from BMAd-*Pnpla2*[-/-] mice fed *ad libitum*, or when derived from CR mice of either

genotype (Figure 4J). CR marrow produced fewer granulocyte-macrophage progenitors

(CFU-GM), which was further reduced when isolated from mice with impaired BMAd-lipolysis

(Figure 4K). These data suggest that proliferative and differentiation capacities of cultured

myeloid cell progenitors may have been reprogrammed in the marrow niche when their

energy supply is restricted either by diet or impaired BMAd lipolysis (Figure 4L).

BMAd lipolysis is not required to maintain bone homeostasis under calorie-restricted

conditions in female mice. To determine whether sex influences responses of BMAd-

Pnpla2[-/-] mice to CR, we performed similar experiments in female mice at 20 weeks of age.

As expected, after six weeks of CR, both control and knockout mice exhibited comparable

reduced body weights, random blood glucose concentrations, and tissue weights (Figure 3 -

figure supplement 3A and 3B). Although CR and *Pnpla2* deficiency cause expansion of

proximal tibial rBMAT (Figure 3 - figure supplement 3C), neither CR nor BMAd *Pnpla2*

deletion cause alterations in trabecular or cortical bone variables in female mice (Figure 3 -

figure supplement 3D and 3E). Consistent with these findings, previous studies showed that

CR adult or aged female mice have milder bone loss than in males (Z. Li & MacDougald,

2021). Although male BMAd-*Pnpla2*[-/-] mice exhibited deficiencies in circulating neutrophils,

experiments with female mice did not reveal differences in white or red blood cell populations

(Figure 3 - figure supplement 3F). We next considered whether estrogen protects female

mice from CR-induced osteoporosis. Thus, we ovariectomized control and BMAd-*Pnpla2*[-/-]

330 mice two weeks prior to initiation of CR. Following initiation of CR, both control and knockout

331 mice demonstrated rapid reduction in body weight for two weeks, then gradually stabilized

332 during the following ten weeks (Figure 3 - figure supplement 4A). We did not observe

333 significant differences in glucose tolerance or bone length with CR or *Pnpla2* deletion (Figure

334 3 - figure supplement 4B and 4C). Additionally, although CR caused reduction in tissue

335 weights, these variables were not different between genotypes (Figure 3 - figure supplement

336 4D). Interestingly, despite the dramatic increases of BAMT responding to CR or BMAd-

337 *Pnpla2* deficiency, bone volume fraction and trabecular number were increased by CR in

338 ovariectomized mice (Figure 3 - figure supplement 4E and 4F), suggesting that the metabolic

339 benefits of CR diet combat detrimental effects of estrogen deficiency and/or aging. Although

340 we observed a trend toward reduced trabecular thickness in *Pnpla2*-deficient mice, other

341 skeletal parameters were not affected (Figure 3 - figure supplement 4F).

343 **Coupling of BMAd *Pnpla2* deletion and CR results in extensive alterations to the bone**

344 **marrow transcriptome.** To determine mechanisms by which BMAd *Pnpla2* deficiency

345 causes bone loss in CR male mice, we profiled overall gene expression using bulk RNAseq

346 in bone marrow plugs from distal tibiae, a skeletal location highly enriched with BMAT. PCA

347 plots show that RNA profiles from CR groups are distinct from *ad libitum* controls (Figure 5 -

348 figure supplement 1A). Whereas *Pnpla2* deficiency does not cause gene expression to

349 diverge substantially in mice fed *ad libitum*, loss of *Pnpla2* interacts with CR to cause a well-

350 segregated pattern of gene expression (Figure 5 - figure supplement 1A). By our criteria (padj

351 < 0.05, |log2 fold change|>1), CR changes expression of 1,027 genes compared to *ad libitum*

352 controls. Although *Pnpla2* deficiency alone only alters 10 genes in BMAd-*Pnpla2*$^{-/-}$ mice fed

353 *ad libitum*, loss of *Pnpla2* in CR mice causes alterations in 1,060 genes (Figure 5 - figure

354 supplement 1B). Analyses of genes regulated in BMAd-*Pnpla2*$^{+/+}$ mice with CR reveals four

355 distinct clusters (Figure 5A). Approximately 80% of genes fall in cluster 1, which are

upregulated by CR in control mice, with induction largely blocked by *Pnpla2* deficiency in CR

mice. Pathway analyses of cluster 1 with Metascape (https://metascape.org) reveals that

regulated genes are associated with vasculature development and cellular response to

growth factor stimulus, which likely reflects adaptation mechanisms to compensate for energy

insufficiency (Figure 5 - figure supplement 1C). In addition, genes associated with skeletal

system development, extracellular matrix organization, and adipogenesis pathways are

upregulated by CR in control mice, but these effects are blunted by *Pnpla2* deletion (Figure 5

- figure supplement 1C). Cluster 2 includes genes that are mildly upregulated by CR in

control mice and are further increased with *Pnpla2*-deficiency. However, no specific

pathways are enriched in this gene set (Figure 5 - figure supplement 1D). Cluster 3 highlights

genes that are downregulated by CR treatment regardless of genotype (Figure 5 - figure

supplement 1E), and which are associated with B cell proliferation and interleukin-8

production, which may partially explain the changes observed in hematopoietic cellularity.

Cluster 4 contains 80 genes that are down-regulated independently by CR and *Pnpla2*-

deficiency, and are associated with regulation of ossification and calcium-mediated signaling

(Figure 5 - figure supplement 1F).

To evaluate further whether BMAd-*Pnpla2* is required for the adaptation of BMAT to CR,

we graphically ordered genes from those maximally induced to those most repressed by CR

in control mice (Figure 5B; red dots). Of these genes, 67.7% do not meet our criteria for

regulated expression by CR in BMAd-*Pnpla2*$^{-/-}$ mice (blue dots). Pathway analyses on the

32.3% of genes regulated by CR regardless of *Pnpla2* deficiency reveals association with

ribonuclease activity, response to hormones and other transmembrane signaling pathways

(Figure 5 - figure supplement 2A). Pathway analyses of genes for which *Pnpla2* is required

for response to CR identifies vasculature development and cellular response to growth factor

stimulus, similar to cluster 1, followed closely by skeletal system development, adipogenesis

genes, and extracellular matrix organization pathways (Figure 5C). A heatmap shows that the

adipogenic genes, including *Adipoq, Fabp4, Cebpa, Lipe, Lpl, Pparα, Pparγ, Scd1, Plin1,*

Cd36, Fasn and *Acaca*, are upregulated by CR in control mice (Figure 5D), a subset of which

were confirmed by qPCR (Figure 5 - figure supplement 2B). These changes may help explain

molecular mechanisms underlying BMAT expansion following CR. It is important to note that

whereas CR BMAd-*Pnpla2*$^{+/+}$ and BMAd-*Pnpla2*$^{-/-}$ mice had comparable amounts of BMAT

(Figure 3 - figure supplement 1O and 1P), adipocyte gene expression is greatly suppressed

in mice with BMAT lacking *Pnpla2 (Figure 5D).* This observation is consistent with prior work

showing that adipocyte-specific deletion of *Pnpla2* results in decreased expression of genes

associated with lipid uptake, synthesis, and adipogenesis (Schoiswohl et al., 2015), perhaps

because NEFA and associated metabolites act as PPAR ligands are decreased (Mottillo,

Bloch, Leff, & Granneman, 2012). In addition, genes related to endogenous fatty acid

biosynthesis are also increased by CR in BMAd-*Pnpla2*$^{+/+}$ mice, but not in mice lacking

Pnpla2 (Figure 5 - figure supplement 2C). Interestingly, myeloid leukocyte differentiation

genes follow a similar pattern (Figure 5 - figure supplement 2D), which may contribute to

reduced neutrophil production and impaired myeloid cell proliferation of CR BMAd-*Pnpla2*$^{-/-}$

mice (Tables 1 and 2, and Figure 4).

We previously observed increased bone loss in BMAd-*Pnpla2*$^{-/-}$ mice challenged with CR

(Figure 3). To investigate potential mechanisms underlying this bone loss, we analyzed

pathways from Cluster 1 related to bone metabolism, skeletal system development and

extracellular matrix organization. Interestingly, osteoblast-derived alkaline phosphatase

(*Alpl), Col1a1* and *Col1a2*, and bone marrow Fgf/Fgfr and Wnt signaling-related molecules

are highly induced by CR in control mice but not in *Pnpla2*-deficient mice (Figure 5E); many

of these genes are also found in the ossification pathway (data not shown). Multiple collagen

genes, *Lox*, and *Adamts* (A Disintegrin and Metalloproteinase with Thrombospondin motifs)

family members, which are multidomain extracellular protease enzymes and play key roles in

extracellular matrix remodeling, are also upregulated by CR in control mice, with effects

largely eliminated by *Pnpla2* deficiency (Figure 5F). These findings may partially explain why

osteoid thickness tends to be thinner in BMAd-*Pnpla2*$^{-/-}$ mice (Figure 3K). In this regard, 20

collagen genes are significantly up-regulated in control mice following CR (Figure 5 - figure

supplement 2E), whereas only four collagen genes are elevated in CR BMAd-*Pnpla2*$^{-/-}$ mice.

Taken together, these data suggest that under conditions of limited dietary energy, BMAds

provide energy to maintain osteoblast functions, including the secretion of collagen matrix for

osteoid synthesis.

BMAd lipolysis is required for trabecular and cortical bone regeneration. To test

whether BMAds are a critical source of local energy under conditions where energy needs

are elevated, we investigated the impact of CR and BMAd *Pnpla2* deficiency on bone

regeneration. To do this, we created a 0.7 mm hole in the proximal tibia, approximately 1 to 2

mm distal from the growth plate, and evaluated trabecular and cortical parameters nine days

later (Figure 6A). As expected, CR impairs formation of new trabecular bone in the region of

interest (ROI) by decreasing trabecular bone volume fraction, bone mineral density, and bone

mineral content (Figure 6B). Importantly, these effects of CR are mimicked by BMAd-specific

Pnpla2 deletion; however, gene deletion does not compound effects of CR on impaired bone

regeneration, perhaps because bone volume is already low with either CR or *Pnpla2*

deficiency alone. We then visualized newly formed cortical bone with Safranin O/ Fast Green

(SO/FG) staining of paraffin-embedded proximal tibia sections (Figure 6C). Of note, both

bone marrow stromal and periosteal cells contribute to the cortical bone regeneration and are

derived from common mesenchymal progenitors (Duchamp de Lageneste et al., 2018).

Although deficiency of BMAd-lipolysis is unlikely to affect the periosteal cell functions when

the cortical bone is intact, it may interact with this cell population during bone regeneration.

CR also impaired formation of new cortical bone by decreasing cortical bone volume fraction,

bone mineral density, and bone mineral content. As with trabecular bone formation, effects of

CR on cortical bone are mimicked by BMAd-specific depletion of *Pnpla2*, but effects of gene

deletion do not exacerbate CR-induced impaired bone regeneration (Figure 6D). Taken

together, these data provide compelling evidence that under conditions where energy

requirements are high, BMAd provide a critical local energy source for bone regeneration.

Energy from BMAd protects against bone loss caused by chronic cold exposure. To

explore further under what conditions BMAd lipolysis may be critical for bone cell functions,

control and BMAd-*Pnpla2$^{-/-}$* female mice at 20 weeks of age were housed at room (22°C) or

cold (5°C) temperatures for three weeks. Cold exposure is well-documented to increase

energy expenditure and adaptive thermogenesis, largely fueled by energy stored in WAT

depots. As expected (Scheller et al., 2015; Scheller et al., 2019), cold exposure results in

smaller regulated BMAds within the proximal tibia of BMAd-*Pnpla2$^{+/+}$* mice (Figure 6E and

6F). In contrast, BMAd size is increased at baseline in BMAd-*Pnpla2$^{-/-}$* mice, with no

reduction observed with cold exposure (Figure 6E and 6G). These data indicate that intact

BMAd lipolysis is required for reduction of BMAd size with cold exposure. We then evaluated

bone mass by μCT and found that whereas trabecular bone of the proximal tibia is

maintained in BMAd-*Pnpla2$^{+/+}$* mice with cold stress, trabecular bone volume fraction and

bone mineral density decline with cold exposure in BMAd-*Pnpla2$^{-/-}$* mice (Figure 6I). These

results indicate that lipolysis from vicinal BMAds is required for maintenance of bone when

energy needs are high or when energy supply is limited.

Discussion

To date, investigation of the physiological functions of BMAT have been hampered by the

lack of a BMAd-specific mouse model. Although a number of BMAT depletion models show

high bone mass, including A-ZIP (Naveiras et al., 2009), *Adipoq*-driven DTA (Zou et al.,

2019), and *Adipoq*-driven loss of *Pparγ* (Wang, Mullican, DiSpirito, Peed, & Lazar, 2013),

Lmna (Corsa et al., 2021), or *Bscl2* (McIlroy et al., 2018), these lipodystrophic mice also lack

white and brown adipose depots and thus exhibit global metabolic dysfunction, including fatty

liver, hyperlipidemia and insulin resistance. Of note, *Adipoq*-driven Cre used in those studies

also causes recombination in bone marrow stromal cells (Mukohira et al., 2019). Similarly,

mouse models or treatments that result in BMAT expansion, such as *Prx1*-driven *Pth1r*

knockout mice (Fan et al., 2017), CR, thiazolidinedione administration, estrogen-deficiency,

and irradiation (Z. Li & MacDougald, 2021), also cause bone loss, but again, effects on bone

may be secondary to lack of promoter specificity or systemic effects. Lineage tracing studies

have previously been performed to determine cell-specific markers for BMAds. For example,

both *Prx1* and *Osterix* are restricted to bone and trace to 100% of BMAds, but are also

expressed in mesenchymal cells (Logan et al., 2002; Mizoguchi et al., 2014). *Nestin* and

leptin receptor (*LepR*) label over 90% of BMAds, but also trace to stromal cells (Zhou et al.,

2017). Whereas Pdgf-receptor α (*PdgfRα)*-driven Cre expression causes recombination in all

white adipocytes, only 50-70% of BMAds are traced (Horowitz et al., 2017). Thus, our

strategy for targeting BMAds using dual expression of *Osterix* and *Adipoq* will be critical for

improving interpretability of experiments on roles for BMAds in the marrow niche.

After generating this novel BMAd-specific Cre mouse model, we successfully ablated

Pnpla2 in BMAds, and demonstrated the necessity of ATGL in BMAd lipolysis. Our studies

directly demonstrate for the first time the importance of BMAd lipolysis in myelopoiesis and

bone homeostasis under conditions of energetic stress, including CR, irradiation, bone

regeneration and cold exposure. With the loss of peripheral WAT in CR mice, there is a

dramatic increase in BMAT throughout the tibia, and this is accompanied by increased

expression of adipocyte genes, including those involved in lipid uptake, de novo lipogenesis,

and lipolysis. BMAT expansion with CR has been observed in both rodents and humans

(Cawthorn et al., 2014; Devlin et al., 2010; Fazeli et al., 2021), and mechanisms underlying

this observation are still not fully understood. We speculate that CR promotes lipolysis in

peripheral adipose tissues, and the released non-esterified fatty acids have increased flux to

bone marrow, where they are used either directly to maintain hematopoiesis and bone

homeostasis, or used indirectly after having been stored and released from BMAT. Thus, we

speculate that BMAds have elevated rates of lipid uptake, lipogenesis and lipolysis with CR.

However, when BMAd lipolysis is impaired, this dynamic cycle is halted, which is reflected by

the blunted induction of adipocyte genes in response to CR. This BMAd quiescence causes a

shortfall in local energy supply and contributes to hematopoietic cell and osteoblast

dysfunction. In this regard, two major collagens secreted by osteoblasts, *Col1a1* and *Col1a2*,

and *Alpl* were increased by CR in WT mice, but not in calorie-restricted BMAd-*Pnpla2*$^{-/-}$ mice.

These data are consistent with the trend for thinner osteoid observed in CR BMAd-*Pnpla2*$^{-/-}$

mice and reduced circulating concentrations of the bone formation marker P1NP. Although

bone mineralization is not impaired, new bone forming surface is reduced in CR mice that

lack BMAd lipolysis. In addition, the number of osteoclasts on trabecular bone surface is

higher in CR BMAd-*Pnpla2*$^{-/-}$ mice; however, osteoclast surface per bone surface and a

circulating marker of bone turnover (CTX-1) are not increased. A serum marker for osteoclast

activity, TRACP5b, is decreased in BMAd-*Pnpla2*$^{-/-}$ mice either fed *ad libitum* or a CR diet.

Taken together, our studies suggest that bone mass reduction in CR BMAd-*Pnpla2*$^{-/-}$ mice is

less likely due to degradation of bone by osteoclasts, and more likely due to impaired

osteoblast function, including osteoid production.

Interestingly, the contributions of BMAd lipolysis to bone homeostasis appear to be more

important in male mice compared to females. Although we considered that a stronger

phenotype might be revealed in female mice following estrogen depletion, the low bone mass

observed with ovariectomy or CR may represent a critical threshold that is strongly defended

through mechanisms independent of BMAd lipolysis. Alternatively, it is possible that

androgens increase energy requirements of bone such that male mice are more dependent

on BMAd lipolysis under stressful conditions. Importantly, we observed that bone volume

fraction and trabecular number were increased by 12 weeks of CR in OVX mice of both

genotypes, suggesting that the metabolic and anti-aging benefits of CR somehow block the

bone loss associated with estrogen deficiency as mice age to 40 weeks.

Surprisingly, despite expansion of BMAT in *ad libitum*-fed BMAd-*Pnpla2*[-/-] male and

female mice, we did not observe differences in skeletal parameters. Indeed, the critical role of

BMAd lipolysis in fueling osteoblast was observed in BMAd-*Pnpla2*[-/-] male mice only when

energy needs were high, or the availability of dietary energy was low. Of note, cellular protein

synthesis typically uses 25 to 30% of the oxygen consumption coupled to ATP synthesis

(Rolfe & Brown, 1997), and this percentage may be higher in osteoblasts since protein

synthesis and translation are integral to cell function. Additionally, fatty acids contribute

substantially to the energy demands of bone tissue and cells (Adamek, Felix, Guenther, &

Fleisch, 1987), and in the absence of BMAd lipolysis, these energy needs must be met from

circulating lipids, glucose, lactate and amino acids. Uptake of fatty acids by osteoblasts is

likely mediated by the CD36 fatty acid translocase, which is required in mice to maintain

osteoblast numbers and activity (Kevorkova et al., 2013). Further, impairment of fatty acid

oxidation in osteoblasts and osteocytes led to reduced postnatal bone acquisition in female,

but not male mice. Interestingly, significant increases in the osteoid thickness, osteoid

volume per bone volume, and the osteoid maturation time suggest that a mineralization

defect occurs in mice unable to oxidize fatty acids obtained not only from BMAds, but also

from circulation (Kim et al., 2017).

Given extensive literature describing interactions between BMAds and hematopoietic

cells (Lee, Al-Sharea, Dragoljevic, & Murphy, 2018; Valet et al., 2020), we were surprised at

the lack of substantial changes in hematopoietic progenitors, white blood cells, and red blood

cells in BMAd-*Pnpla2*$^{-/-}$ mice under basal and CR conditions. One possibility is that skeletal

sites containing low BMAd numbers, including vertebrae, sternum, ribs and pelvis, may allow

compensatory formation of circulating mature blood cells (Kricun, 1985). In addition,

hematopoietic progenitors and mature blood cell populations were not substantially altered in

bone marrow of long bones, with the exception of myeloid differentiation into neutrophils,

suggesting that hematopoietic cellularity is generally maintained despite expansion of BMAT

volume. Although hematopoietic progenitors and mature cells can use lipids for energy,

glucose serve as the major source of energy for glycolysis and oxidative metabolism in these

cell populations (Jeon, Hong, Kim, & Lee, 2020; Roy, Biswas, Verfaillie, & Khurana, 2018). It

is perhaps unsurprising, given the importance of blood cell production in maintaining life, that

there is a great deal of metabolic flexibility when it comes to fuel sources for hematopoiesis.

While HSPC numbers are unaltered in our BMAd-*Pnpla2*$^{-/-}$ mice, myeloid lineage cell

recovery is significantly blunted by CR and deficiency of BMAd-lipolysis following sublethal

irradiation and in *in vitro* myeloid CFU assays. Myeloid progenitors from CR BMAd-*Pnpla2*$^{-/-}$

mice had an impaired capacity to expand, suggesting that progenitors were reprogrammed in

response to energy deficiency in the bone marrow niche. We did not profile the metabolic

changes of HSPCs in CR BMAd-*Pnpla2*$^{-/-}$ mice, but it has been reported that fatty acid

oxidation is required for HSC asymmetric division to retain the stem cell properties (Ito et al.,

2012).

In summary, we have developed a novel mouse model to specifically evaluate the

importance of BMAds as a local energy source. We report that BMAds are a local energy

source that support myeloid cell lineage regeneration following irradiation, and maintain

progenitor differentiation capacity when systemic energy is limiting. In addition, we find that

BMAT serves a highly specialized function to maintain bone mass and osteoblast function in

times of elevated local energy needs such as with bone regeneration, increased whole body

energy needs from cold exposure, or when dietary energy is limited due to CR.

Limitations of Study

There are some limitations to the BMAd-Cre mice that should be noted. For instance,

expression of Cre from the IRES within the 3'-UTR of *Adipoq* is relatively low, and thus rates

of recombination are less frequent in young mice than optimal. Mice that are older than 16

weeks are suggested for future usage of this Cre mouse model. However, bone formation

and turnover rates decrease with age both in mice and humans (Fatayerji & Eastell, 1999;

Ferguson, Ayers, Bateman, & Simske, 2003), which may provide challenges for mechanistic

studies. The IRES-Cre cassette also causes hypoadiponectinemia so experimental design

must include appropriate controls with BMAd-Cre positive mice with or without floxed genes.

Although a total absence of adiponectin reduces bone mass (Naot et al., 2016; Yang et al.,

2019), mice with hypoadiponectinemia described here did not exhibit metabolic or bone

phenotypes. In retrospect, a better approach might have been to accept loss of one *Adipoq*

allele and insert the FAC cassette at the start site of the adiponectin coding region to promote

high levels of Cre expression. This strategy might also have allowed inducible knockout of

BMAd genes with tamoxifen through expression of CreERT2, which we found to be too

inefficient to be functional when inserted into the 3'-UTR of *Adipoq*, with only 5 to 8% BMAds

labeled in tamoxifen-treated adult mice. Of note, all mice used in these studies were on a

mixed SJL and C57BL/6J background, and mouse strain influences bone mass, and

responsiveness of bone to stressors.

Materials and Methods

 KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit monoclonal anti-adiponectin	Sigma-Aldrich	A6354
Mouse/Rat FABP4/A-FABP Antibody	R&D Systems	AF1443
alpha Tubulin Monoclonal Antibody	Thermo Fisher Scientific	MA180017
Anti-ERK 2 Antibody (D-2)	Santa Cruz Biotechnology	sc-1647
Anti-Albumin antibody	Abcam	Ab207327
ATGL Antibody	Cell Signaling Technology	2138S
ATGL Antibody	Abcam	ab207799
Anti-GFP antibody	Abcam	ab13970
RFP Antibody Pre-adsorbed	Rockland	600-401-379
Goat anti Rabbit IgG (H+L) Secondary Antibody, Alexa Fluor 594	Invitrogen	A11012
Alexa Fluor 488 goat anti-chicken IgG (H+L)	Invitrogen	A11039
Ly6G FITC	BD Biosciences	551460
CD11b APC	Invitrogen	17-0112-82
CD115 APC-Cy7	Biolegend	135531
CD3e PE-Cy7	Biolegend	100319
CD19 Pacific Blue	Invitrogen	48-0193-82
CD45 AlexaFluor700	BD Biosciences	560510
Gr-1 Biotin	Biolegend	79750
CD11b Biotin	Biolegend	79749
B220 Biotin	Biolegend	79752
CD3e Biotin	Biolegend	79751
TER119 Biotin	Biolegend	79748

Sca1 PE-Cy7	Invitrogen	25-5981-82
cKit (CD117) APC-Cy7	Biolegend	105826
CD150 BrilliantViolet 421	Biolegend	115925
CD48 FITC	Invitrogen	11-0481-85
CD16/32 PerCP-Cy5.5	Biolegend	101324
CD105 APC	Biolegend	120413
CXCR2 PE	Biolegend	149304
CXCR4 PE-Dazzle	Biolegend	146514
CD62L BrilliantViolet 421	Biolegend	104435
Streptavidin BrilliantViolet 510	Biolegend	405233
Chemicals, peptides, and recombinant proteins		
Calcein	Sigma-Aldrich	C0875
EDTA	DOT Scientific Inc	dse57020
Tetroxide Osmium	Electron Microscopy Sciences	19190
Forskolin	Cayman Chemical Company	11018
Bovine Serum Albumin (BSA), Fraction V	Gold Biotechnology	A-421-250
qPCRBio SyGreen Mix Hi-ROX Blue	Innovative Solutions	4SPB20.16
PCRBio HS Taq Mix Red	Innovative Solutions	4SPB10.23
Agarose	Thermo Fisher Scientific	BP160-500
RNA STAT-60	AMSBIO	CS-502
M-MLV Reverse Transcriptase	Thermo Fisher Scientific	28025013
100 bp DNA Ladder	NEB	N3231S
Critical commercial assays		
Acid Phosphatase Leukocyte (TRAP) Kit	Sigma-Aldrich	387A-1KT
Free Glycerol Determination Kit	Sigma-Aldrich	FG0100
NEFA Reagent (NEFA-HR(2))	FUJIFILM Wako Diagnostics	NC9517309
BCA Protein Assay Kit	Thermo Fisher Scientific	23225

RAT/MOUSE P1NP ELISA KIT	Immunodiagnostic Systems Inc	NC9666468
Mouse TRANCE/RANK L/TNFSF11 Quantikine ELISA Kit	R&D Systems	MTR00
MOUSE TRAP ASSAY	Immunodiagnostic Systems Inc	NC9360739
MethoCult™ GF M3534	ATEMCELL	03534
DNA-free™ Kit	Life Technologies	AM1906
Deposited data		
RNA-seq	This study	GSE183784
Experimental models: Organisms/strains		
Mouse: *Osterix-FLPo*	Generated in University of Michigan Transgenic Animal Model Core	N/A
Mouse: FLPo-dependent *Adipoq-Cre* (FAC)	Generated in University of Michigan Transgenic Animal Model Core	N/A
Mouse: Frt-floxed EGFP	Generated in University of Michigan Transgenic Animal Model Core	N/A
Mouse: mT/mG	Jackson Laboratory	Stock No. 007676
Mouse: *Pnpla2*flox/flox	Jackson Laboratory	Stock No. 024278
Oligonucleotides		
Primers for genotyping PCR and qPCR	See Table 3	N/A
Software and algorithms		
Microsoft Office	Microsoft	https://its.umich.edu /communication/coll aboration/microsoft- office-365/getting- started

Adobe photoshop	Adobe	https://www.adobe.com/creativecloud/desktop-app.html
Prism 9	GraphPad software	https://www.graphpad.com/
Image J	Image J	https://imagej.nih.gov/ij/
MetaMorph	BioVision Technologies	https://www.biovis.com/metamorph.html
Scano μCT 100	SCANCO Medical AG	https://www.scanco.ch/
BIOQUANT OSTEO	BIOQUANT Image analysis corporation	https://bioquant.com/
STAR	PMID 23104886	https://github.com/alexdobin/STAR
DESeq2	PMID 25516281	https://bioconductor.org/packages/release/bioc/html/DESeq2.html
QualiMap-2	PMID 26428292	https://qualimap.conesalab.org
R	The R Foundation	https://www.r-project.org/
Metascape	PMID 30944313	https://metascape.org/gp/index.html

Dragonfly	ORS - OBJECT RESEARCH SYSTEMS	https://www.theobjects.com/dragonfly/index.html
FlowJo	BD Biosciences	https://www.flowjo.com/solutions/flowjo
Other		
Element HT5 Veterinary Hematology Analyzer	Heska	https://www.heska.com/product/element-ht5/
FACSAria III cell sorter	BD Biosciences	N/A
LSRFortessa Cell Analyzer	BD Biosciences	N/A
Nikon A1 Confocal Microscope	Nikon	N/A
Bayer Contour Next Test Glucose Strips	Diabetic Corner	ByrContournext
Scanco µCT 100 micro-computed tomography system	SCANCO Medical AG	https://www.scanco.ch/
Olympus BX51	Olympus	N/A
StepOnePlus™ Real-Time PCR System	Thermo Fisher Scientific	4376600
Microtome	Leica	RM2235
Cryostat	Leica	N/A
Slide scanner	Nikon	N/A

Resource Availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled

by the Lead Contact, Ormond A. MacDougald (macdouga@umich.edu).

Materials availability

Our *Osterix-FLPo* and <u>FLPo</u>-dependent *<u>A</u>dipoq-<u>C</u>re* (FAC) mouse models will be available to

investigators upon request. All the other data and materials that support the findings of this study are

available within the article and supplemental information, or available from the authors upon request.

Data availability

The accession number for the BMAT bulk RNA seq data reported in this paper is GEO: GSE183784.

This paper does not report original code.

Animal

Generation of BMAd-specific Cre mouse model. To create a BMAd-specific Cre mouse

model, we expressed mouse codon-optimized FLP (FLPo) (Raymond & Soriano, 2007) from

the *Osterix* (*Sp7*) locus to recombine and activate a Cre expressed from the *Adipoq* locus.

After designing sgRNA target sequences against 3'-UTR of endogenous *Osterix* (sgRNA:

gatctgagctgggtagaggaagg) and *Adipoq* (sgRNA1: tgaacaagtgagtacacgtgtgg; sgRNA2:

cagtgagcagaaaaatagcatgg) genes with the prediction algorithm available at

http://crispor.tefor.net, we cloned sgRNA sequences into an expression plasmid bearing both

sgRNA scaffold backbone (BB) and Cas9, pSpCas9(BB) (Ran et al., 2013), which is also

known as pX330 (Addgene plasmid ID: 42230; Watertown, MA). Modified pX330 plasmids

were injected into fertilized ova. After culture to the blastocyst stage, Cas9 activity was

confirmed by sequencing the predicted cut site. We then inserted a DNA fragment containing

an in-frame fusion protein between endogenous *Osterix* and FLPo, separated by coding

sequence for P2A (porcine teschoviral-1) self-cleavage site to allow the full-length proteins to

function independently. We also inserted into the 3'-UTR of *Adipoq* an IRES-F3-Frt-reversed

Cre-F3-Frt cassette with ~1kb of 5' and 3' flanking homology-arm sequence. The targeting

vectors (Cas9 expression plasmid: 5 ng/ul; donor DNA: 10 ng/ul) were injected into fertilized

mouse eggs, which were transferred into pseudopregnant recipients to obtain pups. Tail DNA

was obtained for genomic PCR and Sanger sequencing to screen F0 founders for desired

genetic modifications at the expected locations. These F0 generation chimeric mice were

mated with normal mice to obtain mice that are derived exclusively from the modified

fertilized eggs. Germline transmitted mice (F1 generation) carrying the designated genetic

modifications were bred with reporter mice to validate the activity and efficiency of FLPo and

Cre recombinase. These mice were generated by University of Michigan Transgenic Animal

Model Core and MDRC Molecular Genetics Core.

Validation of Osterix-FLPo specificity and efficiency. *Osterix-FLPo* mice were bred with FLP-

dependent EGFP reporter mice (derived from https://www.jax.org/strain/012429; Bar Harbor,

ME), provided by Dr. David Olson from the University of Michigan MDRC Molecular Genetics

Core. Fresh tissue confocal was performed to validate EGFP expression.

Validation of the specificity and efficiency of FLPo-activated Adipoq-Cre (FAC) mice. Among

80 F0 generation pups, 13 mice carried the FAC cassette. These founder mice were bred

with *Osterix-FLPo* mice and mT/mG reporter mice (Stock No. 007676, Jackson Laboratory;

Bar Harbor, ME) to further validate Cre recombinase activity and specificity via fresh tissue

confocal and immunofluorescent histology. Every single mouse in F1 generation was

confirmed and separated for future breeding. We finally selected F0-#693→ F1-#4376 for

BMAd-specific knockout lines. Of note, *Osterix-FLPo* and *FAC* mice were generated on a

mixed SJL and C57BL/6J background. *Pnpla2*$^{flox/flox}$ mice purchased from Jackson laboratory

(Stock No. 024278; Bar Harbor, ME) were on a C57BL/6J background.

Animal housing

Mice were housed in a 12 h light/dark cycle in the Unit for Laboratory Animal Medicine at the

University of Michigan, with free access to water. Mice were fed *ad libitum* or underwent

caloric restriction, as indicated. All procedures were approved by the University of Michigan

Committee on the Use and Care of Animals with the protocol number as PRO00009687.

Animal Procedures

1) 30% CR. After acclimation to single-housing for 2 weeks, and control diet

(D17110202; Research Diets; New Brunswick, NJ) for a week, daily food intake was

measured for another week. Mice were then fed a 30% CR diet (D19051601; Research

Diets; New Brunswick, NJ) daily at ~6 pm, prior to onset of the dark cycle.

2) Ovariectomy. Female mice at 16 weeks of age underwent ovariectomy (OVX). After 2

weeks recovery, they were either fed an *ad libitum* or 30% CR diet for 12 weeks.

3) Proximal tibial defect. Surgeries were performed under isoflurane anesthesia, and

subcutaneous 0.1 mg/kg buprenorphine was given in 12-h intervals for peri-/post-operative

pain management. The proximal tibial defects were obtained by drilling a hole through

anterior cortical and trabecular bone, 1 to 2 mm below the epiphyseal growth plate, with a 0.7

mm low-speed drill.

4) Cold exposure. Mice were single-housed without nesting materials in thermal

chambers for 3 weeks at 5ºC.

5) Glucose Tolerance Test (GTT). Mice were fasted overnight (16-18 hours). Body

weight and fasting glucose levels were measured, followed by an intraperitoneal (i.p.)

injection of glucose (1g glucose/kg body weight). Blood glucose was measured with Bayer

Contour test strips at 15, 30, 60, 90 and 120 min time points by cutting the tip of tails.

6) Whole body irradiation. Irradiations were carried out using a Kimtron IC320 (Kimtron

Medical; Oxford, CT) at a dose rate of ~4 Gy/minute with total dosage of 6 Gy in the

University of Michigan Rogel Cancer Center Experimental Irradiation Shared Resource.

Dosimetry was carried out using an ionization chamber connected to an electrometer system

that is directly traceable to a National Institute of Standards and Technology calibration.

Fresh tissue confocal microscopy. *Osterix*-driven EGFP expression and BMAd-Cre-driven

mT/mG reporter mice were sacrificed. Fresh tissues were collected immediately and put in

ice-cold PBS, which was protected from light. Soft tissues including adipose tissues and liver

were cut into small pieces and placed in a chamber for confocal imaging (Nikon Ti-E Inverted

Microscope; Minato City, Tokyo, Japan). Bones were bisected, and butterflied on a coverslip

for imaging. Both white field and fluorescent images were taken under 200x magnification.

Histology and histomorphometry. Tissue histology was performed essentially as described

previously (Z. Li et al., 2019). Briefly, soft tissues were fixed in formalin, and embedded in

paraffin for sectioning. Tibiae were fixed in paraformaldehyde, decalcified in EDTA for at least

2 weeks, and followed by post-decalcification fixation with 4% paraformaldehyde. Bone

tissues were then embedded in paraffin, sectioned. After staining with hematoxylin and eosin

(H&E), soft tissues were imaged with an Olympus BX51 microscope. Bones were stained

with H&E or Tartrate-Resistant Acid Phosphatase (TRAP; Sigma-Aldrich, MO) as indicated,

and slides were scanned at 200X magnification. Static measurements include bone volume

fraction, trabecular bone microstructure parameters, osteoblast number and surface, and

osteoclast number and eroded surface. Undecalcified tibia was used for plastic sectioning.

Mineralized trabecular bone and osteoid were evaluated with Goldner's Trichrome Staining.

For dynamic studies, calcein (C0857; Sigma-Aldrich, MO) dissolved in 0.02 g/ml sodium

bicarbonate with 0.9% saline at 20 mg/kg was injected intraperitoneally nine- and two-days

before sacrifice for quantification of mineral surface (MS), inter-label width (Ir. L. Wi), and

mineral apposition rate (MAR) in tibia. Calculations were made with Bioquant Osteo 2014

(Nashville, TN) software in a blinded manner (Merceron et al., 2014; Morse et al., 2014).

µCT analysis. Tibiae were placed in a 19 mm diameter specimen holder and scanned over

the entire length of the tibiae using a µCT system (µCT100 Scanco Medical, Bassersdorf,

Switzerland). Scan settings were: voxel size 12 µm, 70 kVp, 114 µA, 0.5 mm AL filter, and

integration time 500 ms. Density measurements were calibrated to the manufacturer's

hydroxyapatite phantom. Analysis was performed using the manufacturer's evaluation

software with a threshold of 180 for trabecular bone and 280 for cortical bone.

Marrow fat quantification by osmium tetroxide staining and μCT. After analyses of bone

variables, mouse tibiae were decalcified for osmium tetroxide staining, using our previously

published method (Scheller et al., 2015). In addition, a lower threshold (300 grey-scale units)

was used for proximal tibial rBMAT quantification because density of osmium staining is low

due to smaller adipocyte size, and with threshold as 400 grey-scale units for cBMAT in distal

tibia.

Immunofluorescent staining. Decalcified tibiae were embedded in OCT compound and

used for frozen sectioning at 15 μm. Excess OCT was removed, and tibial tissues were

blocked with 10% goat serum for one hour at room temperature. Primary antibodies for GFP

(1:500) and RFP (1:200; Figure 1F) or ATGL (1:100; Figure 2B) were then added to slides

and incubated with bone tissues overnight at 4°C. Secondary antibodies (goat anti Rabbit

IgG, Alexa Fluor 594 & goat anti-chicken IgG Alexa Fluor 488 or donkey anti rabbit IgG

(H+L), Alexa Fluor 488) were added to slides following three washes. Two hours later, DAPI

staining was performed. Slides were mounted with prolong gold antifade reagent and

imaged.

RNA extraction and quantitative real-time PCR (qPCR). RNA was extracted from tissues

after powdering in liquid nitrogen and lysis in RNAStat60 reagent in a pre-cooled dounce

homogenizer. Quantitative PCR was performed using an Applied Biosystems QuantStudio 3

qPCR machine (Waltham, MA). Gene expression was calculated based on a cDNA standard

curve within each plate and normalized to expression of the geometric mean of

housekeeping genes *Hprt, Rpl32A* and *Tbp*.

Immunoblot. Detection of proteins by immunoblot was as described previously (Mori et al.,

2021).

***Ex vivo* lipolysis.** Distal tibial BMAT plugs were flushed out from the bone, and four plugs

were pooled into one well of a 96-well plate for each n. Pre-warmed 2% BSA HBSS was

added to BMAT explants with vehicle (DMSO) or forskolin (FSK, 5 μM), to activate lipolysis

(Litosch, Hudson, Mills, Li, & Fain, 1982). Cultured media was collected hourly for 4 hours,

and glycerol and NEFA concentrations were measured with commercially available kits as

listed above in reagents.

CFU assay. Bone marrow cells of BMAd-*Pnpla2*$^{+/+}$ and BMAd-*Pnpla2*$^{-/}$ mice were obtained

from femur and tibia. To isolate bone marrow, the bones were flushed with IMDM (Gibco

12440; Waltham, MA) containing penicillin-streptomycin antibiotic (Gibco 15270-063;

Waltham, MA). Pelleted cells were counted with a hemocytometer, $1X10^4$ cells were plated in

Methocult medium (Stem cell M3534; Vancouver, Canada) in 35 mm culture dishes, and cells

were then incubated at 37°C in 5% CO_2 with ≥ 95% humidity for seven days. CFU-G, CFU-M

and CFU-GM colonies on each plate were counted using a microscope with a 4X objective

lense.

Bulk RNA sequencing. Distal tibial plugs (cBMAT) from two animals were pooled together

for each bulk RNAseq sample. Total RNA was isolated from BMAT for strand specific mRNA

sequencing (Beijing Genomics Institute, China). Over twenty million reads were obtained

using a paired-end 100 bp module on DNBSEQ platform. The quality of the raw reads data

was checked using FastQC (v.0.11.9) and the filtered reads were aligned to reference

genome (UCSC mm10) using STAR with default parameters. All samples passed the post-

alignment quality check (QualiMap (v.2.2.1). The DEseq2 method was used for differential

expression analysis with genotype (*Pnpla2*$^{+/+}$ vs. *Pnpla2*$^{-/-}$) and treatment (CR vs. *ad libitum*)

as the main effects. Gene ontology analysis was done using MetaScape. The Principal

Component Analysis (PCA) plot was generated using the PlotPCA function building in

DESeq2 package. To compare the number of differentially expressed genes between

genotype/treatment groups, volcano plots were constructed using the Enhanced Volcano

package under R environment. Heatmap plots were generated using pheatmap package

under R environment and the complete-linkage clustering method was used for the

hierarchical cluster of genes. These data are available through NCBI GEO with the following

accession number, GSE183784.

Flow cytometry. Femurs were isolated from mice. Bone marrow was harvested by flushing

the femurs with 1 mL of ice-cold PEB (1X PBS with 2 mM EDTA and 0.5% bovine serum

albumin). Red blood cells were lysed once by adding 1 mL of RBC Lysis Buffer (155 mM

NH_4Cl, 10 mM $KHCO_3$, 0.1 mM EDTA) and gently pipetting to mix. Cells were immediately

pelleted by centrifugation and resuspended in 1 mL of ice-cold PEB. Cells were stained for

30 minutes in PEB buffer with the indicated antibodies below and analyzed on the BD

LSRFortessa or BD FACSAria III. Data was analyzed using FlowJo software (BD

Biosciences, version 10.8). Dead cells and doublets were excluded based on FSC and SSC

distribution. To stain for mature leukocytes antibodies used were against CD45, Ly6G,

CD11b, CD115, CD19, and CD3e. All CD45$^+$ cells were gated first for further identification.

Neutrophils were defined as Ly6G$^+$CD11b$^+$, monocytes were defined as Ly6G$^-$

CD11b$^+$CD115$^+$, B cells were defined as Ly6G$^-$CD11b$^-$CD19$^+$, and T cells were defined as

Ly6G$^-$CD11b$^-$CD3e$^+$. To stain for bone marrow neutrophil populations, antibodies used were

against Ly6G, CD11b, cKit, CXCR2, CXCR4, and CD62L. Pre-neutrophils were defined as

Ly6G$^+$CD11b$^+$cKit$^+$, immature neutrophils were defined as Ly6G$^+$CD11b$^+$cKit$^-$CXCR2^{lo}, and

mature neutrophils were defined as Ly6G$^+$CD11b$^+$cKit$^-$CXCR2^{hi}. To stain for hematopoietic

stem and progenitor cells (HSPCs) antibodies used were against a lineage panel (Gr-1,

CD11b, B220, CD3e, TER119), cKit, Sca-1, CD150, CD48, CD105, and CD16/32. After

gating on the lineage⁻ population, HSPCs were defined as follows; HSCs as

LSKCD150$^+$CD48$^-$, MPPs as LSKCD150$^-$CD48$^-$, HPC1 as LSKCD150$^-$CD48$^+$, HPC2 as

LSKCD150$^+$CD48$^+$, GMPs as LKCD150$^-$CD16/32$^+$, PreGMs as LKCD150$^-$CD105$^-$, PreMegEs

as LKCD150$^+$CD105$^-$, and PreCFUe as LKCD150$^+$CD105$^+$. LSK = lineage$^-$Sca1$^+$cKit$^+$, LK =

lineage$^-$cKit$^+$.

Bone regeneration analysis. Nine days after the proximal tibial defect surgery, proximal

tibiae were collected for μCT scanning. After construction, new generated trabecular and

cortical bone were quantified by Dragonfly software (Montréal, Canada). Under the full view

of 3D and 2D images, a cylinder-shaped region of interest (ROI) was defined in trabecular or

cortical bone defect area (as shown in Figure 5A) with 3D dimension as 0.3 mm (diameter) x

0.7 mm (height) for trabecular bone and 0.3 mm (diameter) x 0.2 mm (height) for cortical

bone. An automatic split at otsu threshold for each bone was collect first, and then mean

threshold for a whole cohort was calculated. This average threshold was applied to each

bone to normalize bone volume fraction and mineral content.

Statistics. We calculated the minimal animal number required for studies based on the mean

and SD values to make sure we had adequate animals per group to address our hypothesis.

All the mice were randomly assigned to the indicated groups. Although the investigators

responsible for group allocation were not blinded to the allocation scheme, they were blinded

to group allocation during data collection, and the investigators responsible for analyses were

blinded to the allocation scheme.

Significant differences between groups were assessed using a two-sample *t*-test or

ANOVA with post-tests as appropriate: one-way ANOVA with Tukey's multiple comparisons

test, two-way ANOVA with Sidak's multiple comparisons test and three-way ANOVA analysis

as appropriate. All analyses were conducted using the GraphPad Prism version 9. All

graphical presentations are mean +/- SD. For statistical comparisons, a *P*-value of < 0.05

was considered significant. All experiments were repeated at least twice.

Acknowledgements

This work was supported by grants or fellowships from the NIH to OAM (R01 DK62876; R24

DK092759; R01 DK126230; R01 AG069795), SMR (T32 GM835326; F31 DK12272301),

DPB (T32 HD007505; T32 GM007863), KS (R01DK115583), KTL (T32 DK071212; F32

DK122654), RLS (T32 DK101357; F32 DK123887), KDH (R01AR066028), and CJR

(R24DK092759). ZL was supported by a fellowship from the American Diabetes Association

(1-18-PDF-087) and EB from the American Heart Association (20-PAF00361). This research

was also supported by a Pilot & Feasibility grant and core facilities of the Michigan Integrative

Musculoskeletal Health Core Center (P30 AR069620), Michigan Diabetes Research Center

(P30 DK020572), Michigan Nutrition and Obesity Center (P30 DK089503), and University of

Michigan Comprehensive Cancer Center (P30 CA046592).

Author contributions

ZL, KS, KDH, CJR, and OAM conceived the studies and planned the experimental design.

ZL, JH, JZ, EB, DPB, HM, KG, KTL, RLS, SMR, and SA performed the experiments. ZL, HY,

and OAM analyzed the data. ZL and OAM wrote the manuscript, while all other authors

edited and approved the final manuscript.

Declaration of Interests

The authors declare no conflicting interests.

References

Adamek, G., Felix, R., Guenther, H. L., & Fleisch, H. (1987). Fatty acid oxidation in bone

tissue and bone cells in culture. Characterization and hormonal influences. *Biochem*

J, 248(1), 129-137. doi:10.1042/bj2480129

Ahmadian, M., Abbott, M. J., Tang, T., Hudak, C. S., Kim, Y., Bruss, M., . . . Sul, H. S. (2011).

Desnutrin/ATGL is regulated by AMPK and is required for a brown adipose

phenotype. *Cell Metab, 13*(6), 739-748. doi:10.1016/j.cmet.2011.05.002

Ambrosi, T. H., Scialdone, A., Graja, A., Gohlke, S., Jank, A. M., Bocian, C., . . . Schulz, T. J.

(2017). Adipocyte Accumulation in the Bone Marrow during Obesity and Aging

Impairs Stem Cell-Based Hematopoietic and Bone Regeneration. *Cell Stem Cell,*

20(6), 771-784 e776. doi:10.1016/j.stem.2017.02.009

Bozec, A., Bakiri, L., Jimenez, M., Rosen, E. D., Catala-Lehnen, P., Schinke, T., . . . Wagner,

E. F. (2013). Osteoblast-specific expression of Fra-2/AP-1 controls adiponectin and

osteocalcin expression and affects metabolism. *J Cell Sci, 126*(Pt 23), 5432-5440.

doi:10.1242/jcs.134510

Cawthorn, W. P., Scheller, E. L., Learman, B. S., Parlee, S. D., Simon, B. R., Mori, H., . . .

MacDougald, O. A. (2014). Bone marrow adipose tissue is an endocrine organ that

contributes to increased circulating adiponectin during caloric restriction. *Cell Metab,*

20(2), 368-375. doi:10.1016/j.cmet.2014.06.003

Chen, J., & Long, F. (2013). beta-catenin promotes bone formation and suppresses bone

resorption in postnatal growing mice. *J Bone Miner Res, 28*(5), 1160-1169.

doi:10.1002/jbmr.1834

Cook, J. R., & Semple, R. K. (2010). Hypoadiponectinemia--cause or consequence of human

"insulin resistance"? *J Clin Endocrinol Metab, 95*(4), 1544-1554. doi:10.1210/jc.2009-

2286

Corsa, C. A. S., Walsh, C. M., Bagchi, D. P., Foss Freitas, M. C., Li, Z., Hardij, J., . . .

MacDougald, O. A. (2021). Adipocyte-specific deletion of lamin A/C largely models

human familial partial lipodystrophy type 2. *Diabetes*. doi:10.2337/db20-1001

Devlin, M. J., Cloutier, A. M., Thomas, N. A., Panus, D. A., Lotinun, S., Pinz, I., . . . Bouxsein,

M. L. (2010). Caloric restriction leads to high marrow adiposity and low bone mass in

growing mice. *J Bone Miner Res, 25*(9), 2078-2088. doi:10.1002/jbmr.82

Duchamp de Lageneste, O., Julien, A., Abou-Khalil, R., Frangi, G., Carvalho, C., Cagnard,

N., . . . Colnot, C. (2018). Periosteum contains skeletal stem cells with high bone

regenerative potential controlled by Periostin. *Nat Commun, 9*(1), 773.

doi:10.1038/s41467-018-03124-z

Eguchi, J., Wang, X., Yu, S., Kershaw, E. E., Chiu, P. C., Dushay, J., . . . Rosen, E. D.

(2011). Transcriptional control of adipose lipid handling by IRF4. *Cell Metab, 13*(3),

249-259. doi:10.1016/j.cmet.2011.02.005

Fan, Y., Hanai, J. I., Le, P. T., Bi, R., Maridas, D., DeMambro, V., . . . Lanske, B. (2017).

Parathyroid Hormone Directs Bone Marrow Mesenchymal Cell Fate. *Cell Metab,

25*(3), 661-672. doi:10.1016/j.cmet.2017.01.001

Fatayerji, D., & Eastell, R. (1999). Age-related changes in bone turnover in men. *J Bone

Miner Res, 14*(7), 1203-1210. doi:10.1359/jbmr.1999.14.7.1203

Fazeli, P. K., Bredella, M. A., Pachon-Pena, G., Zhao, W., Zhang, X., Faje, A. T., . . .

Klibanski, A. (2021). The dynamics of human bone marrow adipose tissue in

response to feeding and fasting. *JCI Insight, 6*(12). doi:10.1172/jci.insight.138636

Ferguson, V. L., Ayers, R. A., Bateman, T. A., & Simske, S. J. (2003). Bone development and

age-related bone loss in male C57BL/6J mice. *Bone, 33*(3), 387-398.

doi:10.1016/s8756-3282(03)00199-6

Horowitz, M. C., Berry, R., Holtrup, B., Sebo, Z., Nelson, T., Fretz, J. A., . . . Rosen, C. J.

(2017). Bone marrow adipocytes. *Adipocyte, 6*(3), 193-204.

doi:10.1080/21623945.2017.1367881

Ito, K., Carracedo, A., Weiss, D., Arai, F., Ala, U., Avigan, D. E., . . . Pandolfi, P. P. (2012). A

PML-PPAR-delta pathway for fatty acid oxidation regulates hematopoietic stem cell

maintenance. *Nat Med, 18*(9), 1350-1358. doi:10.1038/nm.2882

Jeon, J. H., Hong, C. W., Kim, E. Y., & Lee, J. M. (2020). Current Understanding on the

Metabolism of Neutrophils. *Immune Netw, 20*(6), e46. doi:10.4110/in.2020.20.e46

Kevorkova, O., Martineau, C., Martin-Falstrault, L., Sanchez-Dardon, J., Brissette, L., &

Moreau, R. (2013). Low-bone-mass phenotype of deficient mice for the cluster of

differentiation 36 (CD36). *PLoS One, 8*(10), e77701.

doi:10.1371/journal.pone.0077701

Kim, S. P., Li, Z., Zoch, M. L., Frey, J. L., Bowman, C. E., Kushwaha, P., . . . Riddle, R. C.

(2017). Fatty acid oxidation by the osteoblast is required for normal bone acquisition

in a sex- and diet-dependent manner. *JCI Insight, 2*(16). doi:10.1172/jci.insight.92704

Kricun, M. E. (1985). Red-yellow marrow conversion: its effect on the location of some

solitary bone lesions. *Skeletal Radiol, 14*(1), 10-19. doi:10.1007/BF00361188

Lee, M. K. S., Al-Sharea, A., Dragoljevic, D., & Murphy, A. J. (2018). Hand of FATe: lipid

metabolism in hematopoietic stem cells. *Curr Opin Lipidol, 29*(3), 240-245.

doi:10.1097/MOL.0000000000000500

Lewis, J. W., Edwards, J. R., Naylor, A. J., & McGettrick, H. M. (2021). Adiponectin signalling

in bone homeostasis, with age and in disease. *Bone Res, 9*(1), 1.

doi:10.1038/s41413-020-00122-0

Li, N., Zhao, S., Zhang, Z., Zhu, Y., Gliniak, C. M., Vishvanath, L., . . . Scherer, P. E. (2021).

Adiponectin preserves metabolic fitness during aging. *Elife, 10*.

doi:10.7554/eLife.65108

Li, Z., Hardij, J., Bagchi, D. P., Scheller, E. L., & MacDougald, O. A. (2018). Development,

regulation, metabolism and function of bone marrow adipose tissues. *Bone, 110*, 134-

140. doi:10.1016/j.bone.2018.01.008

Li, Z., Hardij, J., Evers, S. S., Hutch, C. R., Choi, S. M., Shao, Y., . . . MacDougald, O. A.

(2019). G-CSF partially mediates effects of sleeve gastrectomy on the bone marrow

niche. *J Clin Invest, 129*(6), 2404-2416. doi:10.1172/JCI126173

Li, Z., & MacDougald, O. A. (2021). Preclinical models for investigating how bone marrow

adipocytes influence bone and hematopoietic cellularity. *Best Pract Res Clin*

Endocrinol Metab, 101547. doi:10.1016/j.beem.2021.101547

Litosch, I., Hudson, T. H., Mills, I., Li, S. Y., & Fain, J. N. (1982). Forskolin as an activator of

cyclic AMP accumulation and lipolysis in rat adipocytes. *Mol Pharmacol, 22*(1), 109-

115. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/6289066

Logan, M., Martin, J. F., Nagy, A., Lobe, C., Olson, E. N., & Tabin, C. J. (2002). Expression

of Cre Recombinase in the developing mouse limb bud driven by a Prxl enhancer.

Genesis, 33(2), 77-80. doi:10.1002/gene.10092

Maridas, D. E., Rendina-Ruedy, E., Helderman, R. C., DeMambro, V. E., Brooks, D., Guntur,

A. R., . . . Rosen, C. J. (2019). Progenitor recruitment and adipogenic lipolysis

contribute to the anabolic actions of parathyroid hormone on the skeleton. *FASEB J,*

33(2), 2885-2898. doi:10.1096/fj.201800948RR

McIlroy, G. D., Suchacki, K., Roelofs, A. J., Yang, W., Fu, Y., Bai, B., . . . Rochford, J. J.

(2018). Adipose specific disruption of seipin causes early-onset generalised

lipodystrophy and altered fuel utilisation without severe metabolic disease. *Mol Metab,*

10, 55-65. doi:10.1016/j.molmet.2018.01.019

Merceron, C., Mangiavini, L., Robling, A., Wilson, T. L., Giaccia, A. J., Shapiro, I. M., . . .

Risbud, M. V. (2014). Loss of HIF-1alpha in the notochord results in cell death and

complete disappearance of the nucleus pulposus. *PLoS One, 9*(10), e110768.

doi:10.1371/journal.pone.0110768

Mizoguchi, T., Pinho, S., Ahmed, J., Kunisaki, Y., Hanoun, M., Mendelson, A., . . . Frenette,

P. S. (2014). Osterix marks distinct waves of primitive and definitive stromal

progenitors during bone marrow development. *Dev Cell, 29*(3), 340-349.

doi:10.1016/j.devcel.2014.03.013

Mori, H., Dugan, C. E., Nishii, A., Benchamana, A., Li, Z., Cadenhead, T. S. t., . . .

MacDougald, O. A. (2021). The molecular and metabolic program by which white

adipocytes adapt to cool physiologic temperatures. *PLoS Biol, 19*(5), e3000988.

doi:10.1371/journal.pbio.3000988

Morse, A., McDonald, M. M., Kelly, N. H., Melville, K. M., Schindeler, A., Kramer, I., . . . Little,

D. G. (2014). Mechanical load increases in bone formation via a sclerostin-

independent pathway. *J Bone Miner Res, 29*(11), 2456-2467. doi:10.1002/jbmr.2278

Mottillo, E. P., Bloch, A. E., Leff, T., & Granneman, J. G. (2012). Lipolytic products activate

peroxisome proliferator-activated receptor (PPAR) alpha and delta in brown

adipocytes to match fatty acid oxidation with supply. *J Biol Chem, 287*(30), 25038-

25048. doi:10.1074/jbc.M112.374041

Mukohira, H., Hara, T., Abe, S., Tani-Ichi, S., Sehara-Fujisawa, A., Nagasawa, T., . . . Ikuta,

K. (2019). Mesenchymal stromal cells in bone marrow express adiponectin and are

efficiently targeted by an adiponectin promoter-driven Cre transgene. *Int Immunol,*

31(11), 729-742. doi:10.1093/intimm/dxz042

Muzumdar, M. D., Tasic, B., Miyamichi, K., Li, L., & Luo, L. (2007). A global double-

fluorescent Cre reporter mouse. *Genesis, 45*(9), 593-605. doi:10.1002/dvg.20335

Naot, D., Watson, M., Callon, K. E., Tuari, D., Musson, D. S., Choi, A. J., . . . Cornish, J.

(2016). Reduced Bone Density and Cortical Bone Indices in Female Adiponectin-

Knockout Mice. *Endocrinology, 157*(9), 3550-3561. doi:10.1210/en.2016-1059

Naveiras, O., Nardi, V., Wenzel, P. L., Hauschka, P. V., Fahey, F., & Daley, G. Q. (2009).

Bone-marrow adipocytes as negative regulators of the haematopoietic

microenvironment. *Nature, 460*(7252), 259-263. doi:10.1038/nature08099

Polineni, S., Resulaj, M., Faje, A. T., Meenaghan, E., Bredella, M. A., Bouxsein, M., . . .

Fazeli, P. K. (2020). Red and White Blood Cell Counts Are Associated With Bone

Marrow Adipose Tissue, Bone Mineral Density, and Bone Microarchitecture in

Premenopausal Women. *J Bone Miner Res, 35*(6), 1031-1039. doi:10.1002/jbmr.3986

Ran, F. A., Hsu, P. D., Wright, J., Agarwala, V., Scott, D. A., & Zhang, F. (2013). Genome

engineering using the CRISPR-Cas9 system. *Nat Protoc, 8*(11), 2281-2308.

doi:10.1038/nprot.2013.143

Raymond, C. S., & Soriano, P. (2007). High-efficiency FLP and PhiC31 site-specific

recombination in mammalian cells. *PLoS One, 2*(1), e162.

doi:10.1371/journal.pone.0000162

Riddle, R. C., & Clemens, T. L. (2017). Bone Cell Bioenergetics and Skeletal Energy

Homeostasis. *Physiol Rev, 97*(2), 667-698. doi:10.1152/physrev.00022.2016

Robles, H., Park, S., Joens, M. S., Fitzpatrick, J. A. J., Craft, C. S., & Scheller, E. L. (2019).

Characterization of the bone marrow adipocyte niche with three-dimensional electron

microscopy. *Bone, 118*, 89-98. doi:10.1016/j.bone.2018.01.020

Rolfe, D. F., & Brown, G. C. (1997). Cellular energy utilization and molecular origin of

standard metabolic rate in mammals. *Physiol Rev, 77*(3), 731-758.

doi:10.1152/physrev.1997.77.3.731

Roy, I. M., Biswas, A., Verfaillie, C., & Khurana, S. (2018). Energy Producing Metabolic

Pathways in Functional Regulation of the Hematopoietic Stem Cells. *IUBMB Life,*

70(7), 612-624. doi:10.1002/iub.1870

Scheller, E. L., Doucette, C. R., Learman, B. S., Cawthorn, W. P., Khandaker, S., Schell,

B., . . . MacDougald, O. A. (2015). Region-specific variation in the properties of

skeletal adipocytes reveals regulated and constitutive marrow adipose tissues. *Nat*

Commun, 6, 7808. doi:10.1038/ncomms8808

Scheller, E. L., Khandaker, S., Learman, B. S., Cawthorn, W. P., Anderson, L. M., Pham, H.

A., . . . MacDougald, O. A. (2019). Bone marrow adipocytes resist lipolysis and

remodeling in response to beta-adrenergic stimulation. *Bone, 118*, 32-41.

doi:10.1016/j.bone.2018.01.016

Schoiswohl, G., Stefanovic-Racic, M., Menke, M. N., Wills, R. C., Surlow, B. A., Basantani,

M. K., . . . Kershaw, E. E. (2015). Impact of Reduced ATGL-Mediated Adipocyte

Lipolysis on Obesity-Associated Insulin Resistance and Inflammation in Male Mice.

Endocrinology, 156(10), 3610-3624. doi:10.1210/en.2015-1322

Shen, W., Chen, J., Punyanitya, M., Shapses, S., Heshka, S., & Heymsfield, S. B. (2007).

MRI-measured bone marrow adipose tissue is inversely related to DXA-measured

bone mineral in Caucasian women. *Osteoporos Int, 18*(5), 641-647.

doi:10.1007/s00198-006-0285-9

Valet, C., Batut, A., Vauclard, A., Dortignac, A., Bellio, M., Payrastre, B., . . . Severin, S.

(2020). Adipocyte Fatty Acid Transfer Supports Megakaryocyte Maturation. *Cell Rep,*

32(1), 107875. doi:10.1016/j.celrep.2020.107875

Vogler, J. B., 3rd, & Murphy, W. A. (1988). Bone marrow imaging. *Radiology, 168*(3), 679-

693. doi:10.1148/radiology.168.3.3043546

Wang, F., Mullican, S. E., DiSpirito, J. R., Peed, L. C., & Lazar, M. A. (2013). Lipoatrophy and

severe metabolic disturbance in mice with fat-specific deletion of PPARgamma. *Proc*

Natl Acad Sci U S A, 110(46), 18656-18661. doi:10.1073/pnas.1314863110

Yang, J., Park, O. J., Kim, J., Han, S., Yang, Y., Yun, C. H., & Han, S. H. (2019). Adiponectin

Deficiency Triggers Bone Loss by Up-Regulation of Osteoclastogenesis and Down-

Regulation of Osteoblastogenesis. *Front Endocrinol (Lausanne), 10*, 815.

doi:10.3389/fendo.2019.00815

Zhou, B. O., Yu, H., Yue, R., Zhao, Z., Rios, J. J., Naveiras, O., & Morrison, S. J. (2017).

Bone marrow adipocytes promote the regeneration of stem cells and haematopoiesis

by secreting SCF. *Nat Cell Biol, 19*(8), 891-903. doi:10.1038/ncb3570

Zou, W., Rohatgi, N., Brestoff, J. R., Li, Y., Barve, R. A., Tycksen, E., . . . Teitelbaum, S. L.

(2020). Ablation of Fat Cells in Adult Mice Induces Massive Bone Gain. *Cell Metab,*

32(5), 801-813 e806. doi:10.1016/j.cmet.2020.09.011

Zou, W., Rohatgi, N., Brestoff, J. R., Zhang, Y., Scheller, E. L., Craft, C. S., . . . Teitelbaum,

S. L. (2019). Congenital lipodystrophy induces severe osteosclerosis. *PLoS Genet,*

15(6), e1008244. doi:10.1371/journal.pgen.1008244

1015

Figures and figure supplement list:

Figure 1 is associated with 2 supplements:

Figure 1 - figure supplement 1

Figure 1 - figure supplement 2

Figure 3 is associated with 4 supplements:

Figure 3 - figure supplement 1

Figure 3 - figure supplement 2

Figure 3 - figure supplement 3

Figure 3 - figure supplement 4

Figure 4 is associated with 1 supplement:

Figure 4 - figure supplement 1

Figure 5 is associated with 2 supplements:

Figure 5 - figure supplement 1

Figure 5 - figure supplement 2

Figure 1. Generation of a BMAd-specific Cre mouse model (BMAd-Cre).
A. Efficacy of *Osterix-FLPo* evaluated by crossing with FLP-dependent EGFP reporter to yield *Osterix-*EGFP.

B. *Osterix-*EGFP male mice at 16 weeks were sacrificed. Fresh tissue confocal microscopy was performed on bisected caudal vertebrae, distal tibia and proximal tibia. Red arrows indicate singly dispersed BMAds. Scale bar; 100 μm.

C. Frozen-sections of proximal tibiae slides from *Osterix-*EGFP were stained with Anti-GFP (green) and DAPI (blue). Scale bar; 50 μm.

D. Schematic of how *Osterix-FLPo* recombines FLPo-activated *Adipoq-Cre* (FAC) in BMAd-Cre mice to restrict expression of Cre to BMAds.

E. BMAd-Cre mice were bred with mT/mG reporter mice and resulting BMAd-mT/mG mice were sacrificed at 16 weeks of age and cellular fluorescence evaluated by fresh tissue confocal microscopy. Scale bar; 100 μm.

F. Proximal tibia sections from BMAd-mT/mG mice were stained with antibodies to tdTomato, (red) and EGFP (green), and nuclei were counterstained with DAPI (blue). * indicates BMAds. Scale bar; 50 μm.
Figure 1 is associated with 2 supplements: Figure 1 - figure supplement 1; Figure 1 - figure supplement 2.

Figure 2. Ablation of adipose triglyceride lipase (ATGL; gene name *Pnpla2*) increases size and number of BMAd.

A-J. Male mice of the indicated genotypes at 24 weeks of age were euthanized for investigation of white adipose tissue (WAT) and bone.

A. RNA was extracted from WAT and caudal vertebrae and converted to cDNA. PCR products for wildtype (WT; 1257bp) *Pnpla2* and exon 2-7 knockout (Mut; 553 bp) bands were visualized.

B. Decalcified proximal tibiae were sectioned and used for immunofluorescent staining for ATGL (green) expression. Slides were counterstained with DAPI (blue) for nuclei. Scale bar; 50 μm.

C. Immunoblot analyses of ATGL, α-tubulin, and ERK1/2 in lysates from subcutaneous WAT and caudal vertebral.

D-F. Decalcified tibiae were stained with osmium tetroxide and visualized by μCT (D). BMAT of proximal (E) and distal (F) tibia was quantified. Data are expressed as mean ± SD. * indicates *P* < 0.05 with a two-sample *t*-test.

G-H. Decalcified tibiae were paraffin-sectioned and stained with Hematoxylin & Eosin. Representative pictures were taken from proximal (G) and distal (H) tibia. Scale bar; 200 μm.

I-J. BMAd sizes from proximal (I) and distal (J) tibiae were quantified with MetaMorph software. Data are expressed as mean ± SD. * indicates *P* < 0.05 with two-way ANOVA with Sidak's multiple comparisons test.

K-L. Distal tibial BMAT was flushed from female BMAd-*Pnpla2*-/- and their wildtype littermates at 24 weeks of age. For each n, distal tibial explants from 2 mice were combined per well and cultured in 2% BSA-HBSS solution (K). Subgroups from each genotype were treated with forskolin (FSK, 5 μM) or vehicle (Veh, DMSO). Released glycerol and non-esterified fatty acid (NEFA) in culture media at indicated time points were measured by colorimetric assay (n = 3-4 per treatment) (L). * indicates *Pnpla2*+/+ (FSK) different from *Pnpla2*+/+ (Veh) and from *Pnpla2*-/- (FSK) with *P* < 0.05 with two-way ANOVA with Sidak's multiple comparisons test.

Figure 3. BMAd lipolysis is required to maintain bone homeostasis in male mice under CR conditions, but not when mice are fed *ad libitum*.

A-C. Male BMAd-*Pnpla2*⁻/⁻ and their BMAd-*Pnpla2*⁺/⁺ littermates with *ad libitum* feeding were euthanized at 24 weeks of age. Tibiae from *ad libitum* mice were analyzed by μCT for indicated trabecular bone variables. Scale bars indicate 500 μm.

D-M. Male mice at 18 weeks of age underwent 30% CR for 6 weeks. Two independent age- and sex-matched cohorts were plotted together for μCT parameters (D-F), one of those two cohorts was used for ELISA, and static or dynamic histomorphometry (G-M).

D-F. Tibiae from CR mice were analyzed by μCT for indicated trabecular bone variables. Scale bar; 500 μm.

G-H. Static histomorphometry analyses were performed to calculate osteoblast number (Ob. N), osteoclast number (OC. N) and osteoclast surface (Oc. S) per bone surface (BS).

I-J. Concentrations of circulating P1NP and TRACP5b in CR mice.

K. Osteoid quantification was performed on undecalcified plastic sections with Goldner's Trichrome staining.

L-M. Dynamic histomorphometry was performed on calcein-labelled trabecular bone from proximal tibia. sLS: single-labelled surface; MS: mineralized surface; Ir.L.Wi: inter-label width; MAR: mineral apposition rate. Data are expressed as mean ± SD. * indicates $P < 0.05$ with a two-sample *t*-test.

Figure 3 is associated with 4 supplements: Figure 3 - figure supplement 1; Figure 3 - figure supplement 2; Figure 3 - figure supplement 3; Figure 3 - figure supplement 4.

Figure 4. BMAd-*Pnpla2* deficiency impairs myelopoiesis.

A-H. BMAd-*Pnpla2*[-/-] mice and littermate controls (*Pnpla2*[+/+]) were caloric restricted (CR) for 20 weeks or remained on an *ad libitum* (AL) diet, and then received whole-body irradiation (6 Gy). Mice were euthanized 9 days post-irradiation. Femurs were collected for flow cytometry to measure the regeneration of hematopoietic cells. Bone marrow mononuclear cells (BMNCs), monocytes, B and T lymphocytes and neutrophils were quantified.

I-L. CFU assays. Femora and tibial bone marrow cells were isolated from BMAd-*Pnpla2*[-/-] mice and littermate controls (*Pnpla2*[+/+]), which had been fed ad libitum (AL) or a caloric restricted (CR) diet for 20 weeks. After counting, 1×10^4 cells were plated for CFU assays. Colonies were counted by an independent expert in a blinded manner 7 days after plating.

Data are expressed as mean ± SD. * indicates $P < 0.05$ with one-way ANOVA with Tukey's multiple comparisons test.

Figure 4 is associated with 1 supplement: Figure 4 - figure supplement 1.

Figure 5. BMAd-*Pnpla2* deficiency causes extensive alterations to the bone marrow transcriptome only when coupled with CR.

Male control and BMAd-*Pnpla2*-/- mice at 24 weeks of age were either fed AL or underwent 30% CR for 6 weeks. Distal tibial cBMAT was flushed and cBMAT from two mice was pooled as one sample for RNAseq analyses (n of 3 or 4 per treatment).

A. Differential genes with our criteria (P adj<0.05 and |Log2 fold change|>1) between BMAd-*Pnpla2*+/+ CR and BMAd-*Pnpla2*+/+ *ad libitum* (AL) were grouped into 4 clusters.

B. Genes different between BMAd-*Pnpla2*+/+ CR and BMAd-*Pnpla2*+/+ AL were ordered from maximum to minimum log2 fold change (red dots), and compared to corresponding data from BMAd-*Pnpla2*-/- CR versus BMAd-*Pnpla2*+/+ AL (blue dots). Venn diagram shows the differential genes between BMAd-*Pnpla2*+/+ CR versus BMAd-*Pnpla2*+/+ AL BMAT; and BMAd-*Pnpla2*-/- CR versus BMAd-*Pnpla2*+/+ AL BMAT.

C. Pathway analyses of genes significantly changed by CR in BMAd-*Pnpla2*+/+ mice, but not in CR mice lacking *Pnpla2* (indicated by * area in panel B). Pathways further analyzed by heatmap indicated with blue arrows.

D-F. Expression Z-scores of genes related to adipogenesis (D), skeletal system development (E) and extracellular matrix organization (F) were shown as heatmap. Effects of genotype and diet, and their interactions were analyzed by three-way ANOVA.

Figure 5 is associated with 2 supplements: Figure 5 - figure supplement 1; Figure 5 - figure supplement 2.

Figure 6. Energy from BMAd is required for trabecular bone regeneration and protects against bone loss caused by chronic cold exposure.

A-D. BMAd-*Pnpla2*^+/+ and BMAd-*Pnpla2*^-/- male mice at 24 weeks of age fed with chow diet (-) or underwent 30% CR for 6 weeks (+). A 0.7 mm proximal tibial defect was created 1 to 2 mm distal to the growth plate. Tibiae were collected 9 days after surgery. MicroCT was performed to analyze trabecular and cortical bone regeneration.

A. Representative analyzing images of bone defect. Example of μCT showing trabecular bone region of interest (ROI; green) and newly generated trabecular bone (red); cortical bone ROI (yellow) and new-formed cortical bone (purple).

B. Quantification of regenerated trabecular bone volume fraction (BV/TV), mineral density (BMD) and mineral content (BMC). Data are expressed as mean ± SD. * indicates *P* < 0.05 with one-way ANOVA with Tukey's multiple comparisons test.

C. Safranin O/ Fast Green (SO/FG) staining of new cortical bone formation in defect sites. Scale bar; 200μm.

D. Quantification of regenerated cortical bone volume fraction (BV/TV), mineral density (BMD) and mineral content (BMC). Data are expressed as mean ± SD. * indicates *P* < 0.05 with one-way ANOVA with Tukey's multiple comparisons test.

E-I. Female BMAd-*Pnpla2*^+/+ and BMAd-*Pnpla2*^-/- mice at 20 weeks of age were singly housed at 5°C for 3 weeks without enrichments. Tibiae were collected for sectioning and μCT analyses.

E. Proximal tibiae were decalcified and paraffin-sectioned for H&E staining. Representative images for proximal tibia are shown. Scale bar, 100 μm.

F-G. Quantification of BMAds from H&E-stained slides using MetaMorph software. Comparison of BMAd size at room temperature (RT) versus cold exposure (CE) in mice of indicated genotypes. Data are expressed as mean ± SD. * indicates *P* < 0.05 with two-way ANOVA with Sidak's multiple comparisons test.

H. Trabecular bone volume fraction (BV/TV) and mineral density (BMD) were quantified by μCT. Data are expressed as mean ± SD. * indicates *P* < 0.05 with one-way ANOVA with Tukey's multiple comparisons test.

Figure 1 - figure supplement 1. Genotyping strategy and validation of BMAd-Cre.

A-C. Schematic of genotyping primer designing for *Osterix-FLPo* (A) and FLPo activated *Adipoq-Cre* (FAC; B), and representative genotyping results (C). Mut= Mutation; WT=wildtype; Ori=original band without conversion of reversed Cre cassette; Flipped=FLPo activated Cre (BMAd-Cre).

D. Representative image of genomic PCR sequencing aligned with the endogenous allele.

E. BMAd-Cre mice were bred with mT/mG reporter mice and the resulting BMAd-mT/mG mice were sacrificed at 16 weeks of age. Cellular fluorescence was evaluated by fresh tissue confocal microscopy. Scale bar; 100 μm.

F. The flipped recombination band is found in mRNA of bone, but not WAT depots. Mice expressing *Osterix-FLPo* with or without FAC were sacrificed and mRNA was isolated. cDNAs from distal tibia, caudal vertebrae, epididymal and subcutaneous WATs were used for PCR and agarose gel electrophoresis.

G, H. Efficiency of BMAd-Cre recombinase is age- and allele-dependent. Fresh caudal vertebrae collected from BMAd-mT/mG mice at indicated ages were bisected and used for confocal imaging. Scale bar; 100 μm.

Figure 1 - figure supplement 2. Insertion of IRES-Cre cassette in the 3'UTR of endogenous *Adipoq* decreases expression of adiponectin but does not cause a metabolic or bone phenotype.
A-F. Endogenous adiponectin expression is decreased by IRES-Cre insertion. Mice expressing *Osterix-FLPo* with or without FAC were sacrificed. Epididymal WAT (eWAT), caudal vertebrae, distal tibiae and serum were collected.
A. mRNA expression of adiponectin in eWAT, caudal vertebrae and distal tibiae from BMAd-Cre (OSX+FAC+) WT, Heterozygous, and Homozygous mice.
B-C. Lysates from eWAT (B) and caudal vertebrae (C) were used for immunoblot analyses of adiponectin and FABP4, with α-tubulin and ERK1/2 as loading controls. Quantification was performed using Image J.
D-F. Non-denatured (D) and denatured (E) serum from BMAd-Cre WT, heterozygous, or homozygous mice was used for immunoblot analyses of adiponectin, with albumin as a reference protein. High, medium and low molecular weight forms of adiponectin were quantified by Image J.
G-P. Hypoadiponectinemia does not cause detectable phenotypes. Male BMAd-Cre WT, heterozygote, or homozygote mice were sacrificed at 18 weeks of age. Soft tissues and bones were collected.
G-K. Body weight, glucose tolerance, and WAT depot weights are not changed in BMAd-Cre mice.
L-P. Trabecular and cortical bone parameters were determined by μCT.
Data are expressed as mean ± SD. * indicates $P < 0.05$ with one-way ANOVA with Tukey's multiple comparisons test.

Figure 3 - figure supplement 1. Blocking BMAd-lipolysis does not influence global metabolism when mice are fed *ad libitum* or calorically restricted.

A-I. BMAd-*Pnpla2*[-/-] male mice and littermate controls (*Pnpla2*[+/+]) were fed *ad libitum* until 24 weeks of age. Body weight (A), glucose tolerance test (B), and weights of subcutaneous WAT (sWAT), epididymal WAT (eWAT) and liver (C-E) were recorded. Decalcified tibiae were used for osmium tetroxide-staining and quantified by μCT analyses to measure the BMAT volume at proximal and distal ends, as indicated by boxed regions (F-G). Concentrations of glycerol and NEFA in serum (H) and bone marrow supernatant (I) were measured with colorimetric assay kits. Glycerol and NEFA contents in bone marrow supernatant were normalized to protein concentrations.

J-R. BMAd-*Pnpla2*[-/-] male mice and littermate controls (*Pnpla2*[+/+]) at 18 weeks of age and underwent a 30% CR for 6 weeks. Body weight changes (J) and glucose tolerance (K) were recorded. sWAT (L), eWAT (M), and liver (N) weights were measured during dissection. BMAT volume at proximal and distal tibia were quantified in osmium tetroxide-stained bones following μCT scanning (O-P). Concentrations of glycerol and NEFA in serum (Q) and bone marrow supernatant (R) were measured using colorimetric assays. Glycerol and NEFA contents in bone marrow supernatant were normalized to protein concentrations.

Data are expressed as mean ± SD. * indicates $P < 0.05$ with a two-sample *t*-test.

Figure 3 - figure supplement 2. Cortical bone variables in BMAd-*Pnpla2*-/- mice and other possible mechanisms for bone loss in BMAd-*Pnpla2*-/- CR mice.

A-D. Mouse tibiae from 24 weeks old *ad libitum* (A-B) or CR (C-D) mice were collected. Cortical bone area (CT. BA/TA) and thickness (Ct. Th) were measured by μCT. Scale bar; 500 μm.

E-J. BMAd-*Pnpla2*-/- male mice and littermate controls (*Pnpla2*+/+) were fed *ad libitum* until 24 weeks of age. E-F. Proximal tibial static histomorphometry was performed to calculate osteoblast number (Ob. N), osteoclast number (OC. N) and osteoclast surface (Oc. S) per bone surface (BS). G-J. Circulating P1NP, RANKL, CTX-1 and TRACP5b were measured with commercially available ELISA kits.

K-N. BMAd-*Pnpla2*-/- male mice and littermate controls (*Pnpla2*+/+) at 18 weeks of age underwent a 30% CR for 6 weeks. K-L. Circulating RANKL and CTX-1 were measured with commercially available ELISA kits. M-N. Histomorphometry analysis for osteoid surface and osteoid maturation time (Omt).

Data are expressed as mean ± SD. * indicates *P* < 0.05 with a two-sample *t*-test.

Figure 4 - figure supplement 1. Flow cytometry strategies for hematopoietic cells and sublethal irradiation-induced hematopoietic regeneration in BMAd-*Pnpla2⁻ᐟ⁻* mice.

A-B. Hematopoietic cell gating strategies in hematopoietic stem/progenitor cells (HSPC, A) and mature leukocytes (B).

C-G. BMAd-*Pnpla2⁻ᐟ⁻* mice and littermate controls (*Pnpla2⁺ᐟ⁺*) underwent 30% caloric restriction (CR) for 20 weeks or remained on *ad libitum* (AL) diet, and then received a whole-body irradiation (6 Gy). Tail vein blood (~50 ul) was collected every 2-3 days to monitor hematopoietic cell recovery. Mice were euthanized at day 9 post irradiation.

C-E. Data from complete blood cell counts shows dynamic changes of white- and red blood cells before and after irradiation.

F-G. Hematopoietic cells from two femurs were stained with cell markers to identify hematopoietic stem and progenitor cells.

Data are expressed as mean ± SD. * indicates $P < 0.05$ with one-way ANOVA with Tukey's multiple comparisons test.

Figure 3 - figure supplement 3. BMAd lipolysis is not required in female mice to maintain bone homeostasis under CR conditions.

BMAd-*Pnpla2*-/- female mice and their wildtype controls (*Pnpla2*+/+) at 18 weeks of age were fed a*d libitum* (AL) or underwent a 30% CR for another 6 weeks.

A-B. Final body weight and random glucose levels were measured (A). sWAT, parametrial WAT (pmWAT), liver and spleen weights were recorded during dissection (B).

C. Representative images from proximal tibiae were collected from decalcified and paraffin-sectioned bones. Scale bar; 200 μm.

D-E. Trabecular and cortical bone variables were determined by μCT analysis.

F. Complete blood counts (CBC) were performed to measure white- and red- blood cells in circulation.

Data are expressed as mean ± SD. * indicates *P* < 0.05 with one-way ANOVA with Tukey's multiple comparisons test.

Figure 3 - figure supplement 4. BMAd-lipolysis impairment in estrogen-deficient female mice does not affect CR-induced bone changes.

16 weeks old female mice underwent ovariectomy and recovered for 2 weeks, which were followed by 30% CR for 12 weeks. Changes in body weight (A) and glucose tolerance test (B) were recorded. Femoral and tibial lengths were measured (C). Weights of sWAT, pmWAT, liver, spleen, and uterus (green dots indicate sham mice) were measured during dissection (D). Trabecular bone parameters were determined by μCT (E). Tb.: trabecular bone; BV/TV: bone volume fraction; BMD: bone mineral density; N: number; Th: thickness; Sp: separation.

Data are expressed as mean ± SD. * indicates $P < 0.05$ with one-way ANOVA with Tukey's multiple comparisons test.

Figure 5 - figure supplement 1. CR causes profound changes in BMAT transcriptome.

24 weeks old male mice underwent 30% CR for 6 weeks. Distal tibial cBMAT was flushed and cBMAT from two mice were pooled as one sample for RNA sample preparation. High-quality RNA samples were submitted for RNAseq analyses.

A. Principal component analysis (PCA) plot shows the distinct transcriptional characters in CR groups with (purple dots) or without (aqua dots) *Pnpla2* in BMAds.

B. Volcano plots show the differential genes with P adj<0.05 & |Log2 fold change|>1 in comparisons between BMAd-*Pnpla2*+/+ CR versus AL (left), BMAd-*Pnpla2*-/- versus BMAd-*Pnpla2*+/+ at AL (middle) and BMAd-*Pnpla2*-/- versus BMAd-*Pnpla2*+/+ at CR (right).

C-F. Differential genes from comparison between BMAd-*Pnpla2*+/+ CR versus AL were grouped into 4 clusters according to the alteration patterns. Pathway analyses were performed on each cluster except cluster 2, which gene set was not enriched in any pathways.

Figure 5 - figure supplement 2. BMAd-*Pnpla2* deficiency causes extensive alterations to the bone marrow transcriptome only when coupled with CR.

24 weeks old male mice underwent 30% CR for 6 weeks. Distal tibial cBMAT was flushed and cBMAT from two mice were pooled as one sample for RNA sample preparation. High-quality RNA samples were submitted for RNAseq analyses.

A. Pathway analysis of gene set that respond to CR independent of *Pnpla2* deficiency in BMAds, indicated by * area.

B. qPCRs were performed to confirm the changes of adipogenesis genes in BMAT. Data are expressed as mean ± SD. * indicates *P* < 0.05 with one-way ANOVA with Tukey's multiple comparisons test.

C-D. Heat maps for genes related to fatty acid biosynthesis and myeloid leukocyte differentiation.

E. Collagen genes that upregulated by CR in BMAd-*Pnpla2*+/+ mice.