Molecular drivers of insecticide resistance in the Sahelo-Sudanian populations of a major malaria vector

Sulaiman S. Ibrahim[1, 2, 3,] *, Abdullahi Muhammad[1, 4], Jack Hearn [1], Gareth D. Weed all [5], Sanjay C. Nagi [1], Muhammad M. Mukhtar [2], Amen N. Fadel[3], Leon M. J. Mug enzi[3], Edward I. Patterson[6], Helen Irving[1] and Charles S. Wondji [1,3]

[1] Vector Biology Department, Liverpool School of Tropical Medicine (LSTM), L3 5QA, UK

[2] Department of Biochemistry, Bayero University, PMB 3011, Kano, Nigeria

[3] LSTM Research Unit, Centre for Research in Infectious Diseases (CRID), P.O. Box 13591, Yaoundé, Cameroon

[4] Centre for Biotechnology Research, Bayero University, PMB 3011, Kano, Nigeria

[5] School of Biological and Environmental Sciences, Liverpool John Moores University, Liverpool L3 3AF, UK

[6] Department of Biological Sciences, Brock University, St. Catharines, Ontario, L2S 3A1, Canada.

* Correspondence: suleiman.ibrahim@lstmed.ac.uk ; Tel.: +44-74-4044-3871; ORCID: 0000-0003-0744-802X

Authors email addresses

Sulaiman S. Ibrahim: suleiman.ibrahim@lstmed.ac.uk

Abdullahi Muhammad: abdullahi.muhammad@lstmed.ac.uk

Jack Hearn: jack.hearn@lstmed.ac.uk

Gareth D. Weedall: g.d.weedall@ljmu.ac.uk

Sanjay C. Nagi: sanjay.nagi@lstmed.ac.uk

Muhammad M. Mukhtar: muhammadmahemukhtar@gmail.com

Amen N. Fadel: amen.fadel@crid-cam.net

Leon M. J. Mugenzi: leon.mugenzi@crid-cam.net

Edward I. Patterson: ipatterson@brocku.ca

Helen Irving: helen.irving@lstmed.ac.uk

Charles S. Wondji: charles.wondji@lstmed.ac.uk

Abstract

Information on common markers of metabolic resistance in malaria vectors from countries/regions sharing similar eco-climatic characteristics can facilitate coordination of malaria control. Here, we characterized populations of the major malaria vector *Anopheles coluzzii* from Sahelo-Sudanian region, spanning four sub-Saharan African countries, Nigeria, Niger, Chad and Cameroon. Genome-wide transcriptional analysis identified major genes previously implicated in pyrethroid and/or cross resistance to other insecticides, overexpressed across the Sahel. These include CYP450s, glutathione S-transferases, carboxylesterases, and cuticular proteins. Several variants identified recently as markers of resistance were found in high frequencies, including in the voltage-gated sodium channel (V402L, I940T, L995F, I1527T and N1570Y), the *acetylcholinesterase*-1 gene (G280S) and the *CYP4J5*-L43F. High frequencies of the phenotypically important chromosomal inversions, 2La, 2Rb and 2Rc were observed, with 2La alternative arrangement fixed across the Sahel, and some of the most commonly overexpressed metabolic resistance genes sitting in these inversions. Two most commonly overexpressed genes, *GSTe2* and *CYP6Z2* were functionally validated. Transgenic *Drosophila melanogaster* expressing *GSTe2* exhibited extremely high DDT and permethrin resistance (mortalities < 10% in 24 h). Serial deletion of the 5′ intergenic region, to identify putative nucleotide(s) associated with *GSTe2* overexpression, revealed that simultaneous insertion of adenine nucleotide and a transition (T->C), between Fork-head box L1 and c-EST putative binding sites are responsible for the high overexpression of *GSTe2* in the resistant mosquitoes. Transgenic flies expressing *CYP6Z2* exhibited marginal resistance towards 3-phenoxybenzyl alcohol and α-cypermethrin. However, significantly higher mortalities were obtained in *CYP6Z2* transgenic flies compared with controls, on exposure to the neonicotinoid, clothianidin. This suggests a possible bioactivation of clothianidin into a toxic intermediate, making it an ideal insecticide against populations of *An. coluzzii* overexpressing *CYP6Z2*. These findings will facilitate regional collaborations within the Sahel region, and refine implementation strategies through refocussing interventions, improving evidence-based, cross-border policy towards local and regional malaria pre-elimination.

Keywords: *Anopheles coluzzii*, Sahel, insecticides, pyrethroid, DDT, metabolic, resistance, genes, inversions.

Background

Since the year 2000 the massive scale-up in vector control interventions and treatment with antimalarial drugs had cut malaria incidence by ~40 % across Africa (Bhatt et al., 2015). Bolstered by this progress the World Health Organization (WHO) has been pushing to eliminate malaria, as proposed in the Global Technical Strategy (GTS) 2016-2030 (WHO, 2015). Unfortunately, the GTS, an ambitious framework with targets to reduce global malaria burden by 90 % in 15 years was dealt an immediate blow by a rebound in malaria transmission, with increased cases between 2016 and 2019 (WHO, 2018b, WHO, 2021). This stark warning of the risk posed to control and elimination efforts was a reflection of the lack of significant progress in the primary regions of interest in sub-Saharan Africa, which constitute 96 % of the 627,000 malaria-related deaths in 2020 alone (WHO, 2019, WHO, 2021). Indeed, as the WHO widens the net of malaria elimination (E-2025) its important to acknowledge that no meaningful progress will be made without progress in sub-Saharan Africa, where none of the six countries having the highest global burden of malaria (e.g. Nigeria alone contributing ~27 % of all cases) is on the path to elimination (WHO, 2018a).

The recent escalation of insecticide resistance in the major malaria vectors (Ibrahim et al., 2019b, Riveron et al., 2019, Toe et al., 2015) makes the development of molecular tools to anticipate emergence and predict spread of resistance across the African continent imperative, in order to achieve malaria control and elimination. For decades no simple molecular assays were available to track metabolic resistance and assess its impact on malaria control and transmission, until the recent discovery of two DNA markers in the *cis*-regulatory region of two cytochrome P450s, *CYP6P9a* (Weedall et al., 2019) and *CYP6P9b* (Mugenzi et al., 2019) in the major malaria vector *Anopheles funestus*; findings which allowed the design of simple PCR assays to detect and track metabolic resistance in the field. Unfortunately, these markers explain resistance only in southern Africa for *An. funestus*, since genetic basis of pyrethroid resistance, and cross-resistance with other insecticides is complex in other African regions with different fronts, driven by distinct genes in the presence of barriers to gene flow (Barnes et al., 2017, Weedall et al., 2019, Weedall et al., 2020). No major metabolic resistance markers for P450s and/or GSTs, functionally validated exist for the major malaria vectors of the *Anopheles gambiae* Complex, though it is the omnipresent vector species with widespread

presence across Africa (Bruce-Chwatt, 1951). This is hindering early detection and tracking of

molecular drivers of resistance in this species, slowing down optimal evidence-based control

measures and resistance management.

Major genomic regions associated with metabolic resistance to pyrethroids in *An. gambiae*

sensu lato include the *CYP6* P450 clusters on the 2R and 3R chromosomes, and a *GST* epsilon

(*GSTe*) cluster on chromosome 3R (Anopheles gambiae Genomes et al., 2017), with some key

resistance-associated genes functionally validated. These genes include *CYP6P3* shown to

confer cross resistance to pyrethroids and organophosphates (Muller et al., 2008, Yunta et al.,

2019), *CYP6M2* conferring cross resistance to pyrethroids (Stevenson et al., 2011, Mitchell et

al., 2012b) and DDT (Mitchell et al., 2012b), as well as *GSTe2* shown to confer resistance to

DDT (Mitchell et al., 2014). For *GSTe2*, in addition to the I114T marker (Mitchell et al., 2014),

a recent study has found a novel mutation (*Gste2*-119V) associated with resistance (Lucas et

al., 2019) using a high-throughput genotypic panel for markers. Functional validation is

required establish the empirical evidence of the role of this mutation in resistance. What has

been missing in the case of *Anopheles gambiae/coluzzii* is reliable molecular markers of

resistance for major metabolic gene families, e.g. P450s (with field and laboratory validated

data) to aid creation of DNA-based diagnostic assays which will allow (i) easy tracking of

resistance in the field (e.g. the case of 119F-*GSTe2* mutation in *An. funestus* (Riveron et al.,

2014b), and (ii) determination of the operational impact of the resistance markers in the field,

as recently done for 119F-*GSTe2* (Menze et al., 2020) and *CYP6P9a*_R/*CYP6P9b*_R markers

(Mugenzi et al., 2019, Weedall et al., 2019) in *An. funestus*.

The *An. gambiae s.l.*, especially populations in the semi-arid steppe, exhibit high frequency of

paracentric chromosomal inversions [one of the most effective instruments for speciation and

local adaptations (Ayala et al., 2014, Dobzhansky, 1971, Kirkpatrick, 2010)], maintained in

spatially and temporally heterogenous environment, and which segregate along climatic

gradients of increasing aridity (Coluzzi et al., 1979)]. The 2L inversions are associated with

resistance to desiccation in adults (Fouet et al., 2012, Gray et al., 2009) and thermal stress in

larvae (Cassone et al., 2011). It was also shown that inversion 2La assorts with insecticide

resistance, e.g., dieldrin plus fipronil (Brooke et al., 2000), and is associated with

thermotolerance and permethrin resistance in the Sahelain *An. coluzzii* (Ibrahim et al., 2021).

To support malaria pre-elimination effort in sub-Saharan Africa we targeted the Sahelo-
Sudanian region, which represent northern-most limit of malaria endemicity in sub-Saharan
Africa, and where malaria is highly seasonal [offering excellent target for pre-elimination
effort through sustained seasonal vector control and seasonal malaria chemoprevention
(Cairns et al., 2012)]. Focusing on the predominant malaria vector, *An. coluzzii* from
Sudan/Sahel transects of four countries, Nigeria, Niger, Chad and Cameroon (Ibrahim et al.,
2019b, Ibrahim et al., 2019c, Fadel et al., 2019, Ibrahim et al., 2019a) we identified the major
metabolic resistance genes mediating pyrethroid resistance and cross-resistance in this region,
established the genetic variants which explained the resistance. We also functionally validated
the roles of two major candidate genes in the resistance (*CYP6Z2* and *GSTe2*), as well as single
nucleotide polymorphism in the 5′ regulatory elements of *GSTe2*, responsible for its
overexpression in the resistant population.

Materials and Methods

Study Site and Mosquito Sampling

Blood fed female *An. coluzzii* mosquitoes, resting indoor were collected at one locality each in
Sudan/Sahel (Figure S1): Hadiyau (HAD: 12°21′38″N, 9°59′15″E), a Sudan/sub-Sahel village in
northern Nigeria; Takatsaba (TAK: 13°44'01.8"N 7°59'05.2"E), a Sahel village in southern
Niger; Simatou (SIMAT: 10°50'40.7"N 14°56'40.9"E), a Sudan/sub-Sahel village in Maga
Department, far north of Cameroon; and Massakory (CHAD: 12° 6′ N, 15° 02′ E), a Sahel town
in Chad Republic. Details of sampling approaches and resistance profiles of mosquitoes
collected from Nigeria, Niger, and Chad are available in previously published articles
(Ibrahim et al., 2019b, Ibrahim et al., 2019c, Ibrahim et al., 2019a). As in the above countries
the Simatou F_1 females were also highly pyrethroid resistant, with a mortality of only 3.7%
from WHO tube bioassays using 0.05% deltamethrin, and no mortality at all with 0.75%
permethrin (data not published).

Genome-wide transcriptional analysis for common insecticide resistance genes in the

Sahel/sub-Sahel regions

RNA extraction, library preparation and sequencing

The RNA was extracted using the Arcturus PicoPure RNA isolation Kit (Applied Biosystems,

CA, USA) from three pools of 8 F_1 *An. coluzzii* females (2-4 day old) alive after exposure to

deltamethrin (resistant, R), unexposed (control, C), and also from unexposed females of the

fully susceptible laboratory colony of *An. coluzzii*, Ngoussou (Mitchell et al., 2012a)

(susceptible, S). RNA isolation was carried out following the manufacturer's protocol with

Dnase I-treatment to remove contaminating DNA. The quantity and quality of RNA was

measured using a NanoDrop spectrophotometer (ThermoFisher, MA, USA) and Bioanalyzer

(Agilent, CA, USA).

Library preparation, sequencing and data quality control were carried out by the Centre for

Genomic Research (CGR), University of Liverpool, UK. RNA samples were subjected to

poly(A) mRNA enrichment and libraries prepared from the poly(A) mRNA-enriched

materials (dual-indexed, strand-specific RNAseq libraries were prepared using the NEBNext

polyA selection and Ultra Directional RNA library preparation kits). Libraries were

sequenced on a single lane of an Illumina HiSeq 4000 (paired-end, 2x150 bp sequencing,

generating data from >280 M clusters per lane). Basecalling and de-multiplexing of indexed

reads were performed by CASAVA version 1.8.2 (Illumina). De-multiplexed fastq files were

trimmed to remove Illumina adapter sequences using Cutadapt version 1.2.1 (Martin, 2011).

Option -O 3 was used, so that the 3' end of any reads which matched the adapter sequence for

3 bp or more were trimmed. Reads were further trimmed to remove low quality bases

using Sickle version 1.200 (Joshi and Fass, 2011) with a minimum window quality score of 20.

Reads shorter than 20 bp after trimming were removed. If both reads from a pair passed this

filter, each was included in either the R1 (forward reads) or R2 (reverse reads) file. If only one

of a read pair passed this filter, it is included in the R0 (unpaired) file. Statistics were generated

using fastq-stats from EAUtils (Aronesty, 2011). Summary of total number of reads for each

sample and distribution of trimmed read length for forward (R1) and reverse (R2) reads and

reads unpaired after trimming (R0) are provided in Figure S2.

Data analysis and estimation of transcript abundance by tag counting and differential gene expression

Paired data for each replicate per country was aligned to the *An. gambiae* reference transcriptome AgamP4.10 downloaded from VectorBase (https://vectorbase.org/) in salmon (0.11.4), using 'validate mappings', 'seqBias', 'gcBias' and 'rangeFactorizationBins 4' flags. Read mapping results (pre-alignment and post-alignment descriptive statistics (flagstat output files) showing sequencing depth and coverage are given in Tables S1 and -S2, respectively. Salmon results were converted into a gene expression matrix using the Bioconductor package 'tximport' for input to DESeq2 1.26.0 (Love, 2014). Differential gene expression was tested for the three possible combinations of Exposed (R, deltamethrin resistant), Unexposed (C, control) and Susceptible replicates (S). For results interpretation $\log_2$-fold change thresholds of 1 was imposed, with false discovery rate adjusted p-values of 0.05 applied to accept significance. Principal components analysis implemented in DESeq2 was used to examine relationships between respective replicates and treatments. This was carried out based on the 500 most variable genes, with data transformation (normalisation/scaling) as implemented in VST (DESeq2). For visualisation of expression results, volcano plots were created using the Enhanced Volcano package (Blighe, 2018) using the top most overexpressed genes from lists which were prepared with $\log_2$FC cut off of 1 and p value of 0.01. Heatmaps were generated (Gu et al., 2016) using the list of the top 50 most overexpressed metabolic resistance genes with $\log_2$FC cut off of 1 and p value of 0.01.

Gene Ontology Enrichment and Mapping/Functional annotations

The enrichment analysis for GO terms was carried out using a topGO package (Alexa A., 2021) against *An. gambiae* (AgamP4.10), with data reannotated using EggNOG v5.0 (http://eggnog5.embl.de/#/app/home). Gene lists used in topGO are from the $\log_2$FC = 0, p <0.05 analysis (the default in DESeq2). For the data from each site (country) six set of results were generated (R C and S contrasts) each for genes either up- or down-regulated in each contrast, for molecular function (MF) and biological process (BP) ontologies. For analysis the GO terms lists for MF were used as input into the Revigo (Supek et al., 2011) for interpretation

and visualisation, querying the Whole Uniprot database, and SimRel setting for semantic
similarity measurements.

Quantitative PCR measurement of expression profiles of metabolic resistance genes

The level of expression of 12 resistance-associated genes was validated by qRT-PCR using the primers provided in Table S3. These include the *GSTe2* (AGAP009194) and *GSTZ1* (AGAP002898), *CYP6Z2* (AGAP008218), *CYP6Z3* (AGAP008217), *CYP4C27* (AGAP009246), *CYP4G16* (AGAP001076), *CYP4G17* (AGAP000877), *CYP6P3* (AGAP002865), *CYP6M2* (AGAP008212), *CYP9K1* (AGAP000818), a *UGT-B19* (AGAP007920) and *COEBE3C* (AGAP005372). The qRT-PCR was carried out using cDNA extracted from 1 µg of total RNA from each of three biological replicates (and three technical replicates each) from the Resistant (R), Control (C) and Ngoussou (S), for each of the four countries. Protocol followed was as established in previous studies (Riveron et al., 2014a), with relative expression level and fold change (FC) of each target gene in R and C relative to S calculated according to the $2^{-\Delta\Delta CT}$ method incorporating the PCR efficiency (Schmittgen and Livak, 2008) after normalization with the housekeeping genes ribosomal protein S7, *RPS7* (AGAP010592) and glycerol-3-phosphate dehydrogenase (NAD+), *GPDH* (AGAP007593). Significant differences were calculated using ANOVA with Dunnett's post hoc test.

Detection of Signatures of Selective Sweep

To detect signature of select sweeps in the major metabolic resistance genes of interest, a RNAseq population genetics pipeline, the Snakemake workflow (https://zenodo.org/record/6078337) was utilised. F_{st} (Hudson et al., 1992) per gene between population pairs and Tajima's D per gene within population were estimated after calling genomic variants with *Freebayes*. This was performed against all SNPs passing quality and missingness filters. Population branch statistic (PBS) scans may also be performed, conditional on the presence of three suitable populations (Yi et al., 2010). It is also possible to run Hudson's F_{st} and PBS scans, taking the average for each protein-coding gene, as opposed to in windows. The population genetics statistical analyses are calculated in scikit-allel v1.2.1 (Alistair Miles, 2020).

Establishment of allele frequencies of variants in genes of interest

After genome alignment, RNA-Seq-Pop utilises samtools (https://zenodo.org/record/6078337)
to cross check specific positions of the genome, calculating raw allele frequencies at those sites
with a custom R script.

Detection of chromosomal inversion polymorphisms and metabolic genes within its
breakpoint

A modified version of the Python 3 program, compkaryo (Love et al., 2019) was used to
karyoptype the major, phenotypically important inversion polymorphisms in chromosome 2,
and calculate its frequencies, *in silico*, using the previously identified tag SNPs significantly
associated with inversions. This allows to predict with high confidence genotypes of the six
common polymorphic inversions on chromosome 2 in individually sequenced genomes of the
field *An. coluzzii*, as well as in the Ngoussou. Compkaryo uses the Ag1000 database
(The *Anopheles gambiae* 1000 Genomes Consortium 2017) (Anopheles gambiae Genomes et al.,
2017) by leveraging a subset of cytologically karyotyped specimens to develop a
computational approach for karyotyping applicable to whole genome sequence.
Modifications in the Snakemake pipeline allows for variable ploidy (useful in the case of
replicates from our pooled RNAsequencing samples) here.

Functional validation of the role of commonly overexpressed resistance-associated genes

Comparative analysis of coding sequences of major resistance genes

To establish presence of allelic variants which could impact catalytic activities, full length
coding sequences (cDNAs) of the two resistance-associated genes, *GSTe2* and *CYP6Z2* were
amplified and sequenced from alive mosquitoes in the four Sahel countries, as well as from
Ngoussou. This was done using total RNA extracted from 5 individual pools of 8 F$_1$ *An.*
coluzzii females (2-4 day old) alive after exposure to deltamethrin (resistant, R for *CYP6Z2*) or
DDT (R for *GSTe2*). Protocol for RNA extraction was as described in previous section, above.
Amplification was done using Phusion HotStart II Taq Polymerase (ThermoFisher
SCIENTIFIC, MA, USA) and the Full primers listed in Table S4. The PCR mix comprised 5X

Phusion HF Buffer (containing 1.5 mM MgCl$_2$), 85.7 µM deoxynucleotides (dNTPs), 0.34 µM each of forward and reverse primers, 0.015 U of Phusion HotStart II DNA Polymerase (Fermentas, MA, USA), 10.71 µL of ddH$_2$0 and 1 µL cDNA. Thermocycling conditions were 1 cycle at 95 °C for 5 min; followed by 35 cycles each of 94 °C for 20 s, 60 °C for 30 s, 72 °C for 2 min (1 min for *GSTe2*); and finally, one cycle at 72 °C for 5 min. PCR products were cleaned with a QIAquick® PCR Purification Kit (QIAGEN, Hilden, Germany) and ligated into the pJET1.2/blunt cloning vector using the CloneJET PCR Cloning Kit (ThermoFisher SCIENTIFIC, MA, USA). These were then cloned into *E. coli DH5α*, plasmids miniprepped with the QIAprep® Spin Miniprep Kit (QIAGEN) and sequenced on both strands using pJET1.2 primers.

Polymorphisms were detected through examination and manual editing of sequence traces using BioEdit version 7.2.3.0 (Hall, 1999) and nucleotide differences in sequences aligned using CLC Sequence Viewer 7.0 (http://www.clcbio.com/). Different haplotypes were compared by constructing a phylogenetic maximum likelihood tree using MEGA X (Kumar et al., 2018). Genetic parameters of polymorphism including number of haplotypes (h) and its diversity (H$_d$), number of polymorphic sites (S) and nucleotide diversity (π) were computed using DnaSP v6.12.03 (Rozas et al., 2017). The DNA sequences were deposited in GenBank, with accession numbers: XXXXX-YYYYY.

Characterization of the 5' regulatory regions of GSTe2 and CYP6Z2

Amplification, cloning and sequence characterisation of 5' regulatory element

To investigate presence of genetic variants in the regulatory elements, which could be responsible for overexpression of *GSTe2*, 351 bp intergenic regions (spanning the 43 bp 3' UTR of *GSTe1,* 248 bp flanking sequence and 60 bp 5'UTR of *GSTe2*) preceding the start codon were amplified from 10 each of DDT-alive and -dead females from the 4 Sahel countries, as well as from the Ngoussou females (primers provided in Table S4). For *CYP6Z2*, a 1078 bp intergenic region was retrieved from the VectorBase and used for amplification of the putative 5'-regulatory elements. Primers spanning 38 bp 3' UTR of *CYP6Z1,* a 937 bp flanking sequence and 103 bp 5'UTR of *CYP6Z2*, preceding the start codon of *CYP6Z2* were used to amplify

fragments from 10 each of deltamethrin-alive and -dead females from Nigeria and Niger, as

well as from the Ngoussou females. Amplification was carried out using HotStart II

Polymerase (ThermoFisher SCIENTIFIC, MA, USA) with similar thermocycling conditions as

outlined above for coding region of *GSTe2* and *CYP6Z2,* respectively. Purification of PCR

amplicons, cloning into pJET1.2 vector, sequencing and polymorphism analysis were done as

outlined above. The DNA sequences were deposited in GenBank, with accession numbers:

XXXXX-YYYYY.

The 351 bp 5'-UTR fragments of *GSTe2* and 1078 bp fragment of the *CYP6Z2* were analysed

with the Gene Promoter Miner (http://gpminer.mbc.nctu.edu.tw/) and MatInspector

(Cartharius et al., 2005) to identify putative promoter elements, predict transcription start

(TSS) and potential transcription factor binding sites.

Cloning of GSTe2 and CYP6Z2 5' regulatory elements in PGL3-Basic vector and dual luciferase

reporter assay

Following analysis of the above sequences the 351 bp intergenic fragments of *GSTe2* were

amplified from the most predominant sequences of DDT-alive, DDT-dead and Ngoussou.

Same was done for for *CYP6Z2* amplifying 1087 bp fragment from deltamethrin-alive and -

dead, and Ngoussou. Primers bearing *kpn*I and *BgL*II sites (Table S4) allowed incorporation

into pGL3-Basic reporter vector containing luciferase gene from the firefly *Photinus pyralis*

(Promega, Wisconsin, USA). Amplification was carried out using Phusion HotStart II

Polymerase, with conditions as above, followed by purification of PCR amplicons, and

cloning into pJET1.2 vector. Positive colonies (sequencing primers for pGL3-Basic provided

in Table S4) were miniprepped; the minipreps digested with the above restriction enzymes,

gel-purified and ligated upstream of luciferase gene in pGL3-Basic vector already linearized

with the same restriction enzymes. Positive colonies were cloned and miniprepped and

concentrations of the recombinant plasmids adjusted to 200 ng/μL before co-transfection.

Protocols for maintenance of the *An. gambiae* cell line 4a-3B (MRA-919,

https://www.beiresources.org/), co-transfection of 200 ng of either *GSTe2* or *CYP6Z2* promoter

constructs, LRIM promoter in pGL3-Basic vector (Lynd and Lycett, 2011), or promoter-less

pGL3-Basic control, with 1 ng/μL of internal control, sea pansy *Renilla reniformis* luciferase

containing the *Drosophila* Actin *5C* promoter in pRL-null (Lynd and Lycett, 2011), are provided

in File S1 (additional text for Methods and Results). Protocol for cell lysis and measurement

of the activities of the firefly luciferase (Dual-Luciferase Reporter Assay kit, Promega), with

normalisation using the Renilla luciferase activity, were as outlined in the Promega Quick

Protocol, and provided in detail in File S1.

Assay background was also measured using lysate from non-transfected control cells. Results

were compared using a two-tailed Chi-Square test of independence as implemented in

GraphPad Prism 7.02 (GraphPad Inc., La Jolla, CA, USA).

Generation of GSTe2 promoter deletion constructs and assays

The intergenic region separating *GSTe1* from *GSTe2* was progressively delineated. Partial

fragments of the 5' regions (encompassing the 3'-UTR of *GSTe1*, the *GSTe1*/*GSTe2* flanking

region and the 5'-UTR of the *GSTe2*) were created by sequential deletion using forward

primers (provided in Table S4, with numbers in primer names denoting distance from the

AUG start codon of *GSTe2*). Reverse primers were those initially used for amplification of the

full 351 bp nucleotide fragments. These fragments include (i) 308 nucleotides fragment

generated from deletion of the 43 bp 3'-UTR of the *GSTe1* (-308 primers) obliterating cellular-

Myb (c-Myb) transcriptional factor binding site; (ii) 291 nucleotide fragments generated from

deletion of *GSTe1* 3'-UTR plus 38 nucleotides of the flanking region (-270 primers), which

removed in addition δ elongation factor 1 (δEF1) binding site; (iii) 262 nucleotide fragments

produced from deletion of *GSTe1* 3'-UTR plus 46 nucleotides of the flanking region (-262

primers), which obliterated in addition Forkhead box L1 (FOX-L1) putative binding site; as

well as (iv) 232 nucleotide fragments produced from deletion of *GSTe1* 3'-UTR plus 75

nucleotides of the flanking region (-232 primers), located 11 nucleotides from the c-EST

binding site/grainy head transcription factor binding site. Also, to assess the impact of

removing the 5'UTR of *GSTe2*, 43 nucleotides upstream the AUG codon were deleted (from

the 60 nucleotide 5'-UTR of both the Sahel-alive and Ngoussou), leaving the putative

transcription start site position motif untouched. This was done using the original forward

primer for the 351 nucleotides intergenic region, with a newly designed primer,

GSTe2_minus_5'-UTR-R (Table S4).

Characterisation of major resistance genes using transgenic analysis

Cloning and microinjection of GSTe2 and CYP6Z2 in Drosophila melanogaster flies

Transgenic flies expressing recombinant GSTe2 and CYP6Z2 were generated using
GAL4/UAS system and used in contact bioassays to confirm if over-expression of these genes
alone can confer resistance to insecticides. Transgenic flies were produced out as outlined in
a previous publication (Ibrahim et al., 2015). Details of amplification of full-length *GSTe2* and
CYP6Z2 from the predominant alleles of sequences of cDNA, cloning into pUASattB vector,
microinjection into germ-line *D. melanogaster* lines, crossing of the UAS-lines with the GAL4-
Actin driver strain are outlined in File S1.

To confirm overexpression of *GSTe2* and *CYP6Z2* in the transgenic flies three replicates each
of 6 females (both experimental and control group) were used for qRT-PCR using a previously
established protocol (Riveron et al., 2014a). Total RNA and cDNA were extracted as described
above, and the relative expression levels of the transgene were assessed by qRT-PCR, with
normalization using the *RPL11* housekeeping gene. The qtrg primers used for the two genes
and the *RPL11* primers are provided in Table S4.

Insecticide Susceptibility Contact Bioassay

For insecticide bioassays, 3- to 4-day old experimental and control F_1 females were exposed to
0.15 % deltamethrin, 2 % permethrin, 0.05 % α-cypermethrin, 4 % DDT and 2 % clothianidin-
impregnated papers prepared in acetone and Dow Corning 556 Silicone Fluid (BHD/Merck,
Hesse, Germany). Flies overexpressing *CYP6Z2* were also exposed to the primary product of
hydrolysis of pyrethroids: 4 % and 20 % (5x) 3-phenoxybenzaldehyde (PBAld) and 3-
phenoxybenzyl alcohol (PBAlc). Transgenic flies expressing *GSTe2* were exposed to 2 %
permethrin, 0.15 % deltamethrin, 0.05 % α-cypermethrin and 4 % DDT only. Impregnated
papers were rolled and introduced into 45 cc plastic vials to cover the entire wall and the vials
plugged with cotton soaked in 10 % sucrose (Riveron et al., 2014a). 20–25 flies were placed in
each vial, and the mortality plus knockdown scored at 1 h, 3 h, 6 h, 12 h and 24 h of exposure
to the insecticides. For each insecticide, assays were performed in 6 replicates and Student's

t-test used to compare the mortality plus knockdown between the experimental groups and

the control.

Results

Genome-wide transcriptional profile of the Sahelian *An. coluzzii* populations

A three-way pairwise comparison was conducted for the data from each country: resistant *vs*

susceptible (R-S), resistant *vs* unexposed control (R-C) and unexposed control *vs* susceptible

(C-S). This captures background variations due to geographical differences in the resistant vs

susceptible (R-S) comparison, accounts for genes overexpressed due to induction (R-C

comparison), as well as genes that are constitutively overexpressed (C-S comparison). A total

of 1384 genes were significantly differentially expressed (FDR-adjusted p <0.05 and $\log_2$ fold

change threshold of 1/FC ≥ 2) in R-S comparison in Nigeria (1077 upregulated and 307 down-

regulated); 1185 genes were differentially expressed in C-S (1002 upregulated and 183

downregulated); and 295 genes in R-C (129 upregulated and 166 downregulated). Of these,

52 genes were commonly differentially expressed in all 3 comparisons (Figure S3, panel a),

including the upregulated genes, *COEAE80* (AGAP006700), *CYP4H18* (AGAP028019),

CYP4H17 (AGAP008358) and cuticular proteins, *CPLCX3* (AGAP006149), *CPR59*

(AGAP006829), *CPR76* (AGAP009874) and *CPR75* (AGAP009871). The Figure S3, panels a-d

depicts the differentially expressed genes for the four countries. For Niger, 881 genes were

differentially expressed in R-S comparison (619 upregulated and 262 down-regulated) (Figure

S3b); 1256 genes were differentially expressed in C-S (986 upregulated and 270

downregulated); and 196 genes in R-C (81 upregulated and 115 downregulated). Of these, 22

genes were commonly differentially expressed in all 3 comparisons [including the

upregulated *aminopeptidase N1* (AGAP012757), *CYP6Z2* (AGAP008218), *chymotrypsin-3*

(AGAP006711) and *chymotrypsin-2* (AGAP006710), and an acid trehalase (AGAP008547)]. For

Chad, 1392 genes were differentially expressed in R-S comparison (975 upregulated and 417

down-regulated) (Figure S3c); 1284 in C-S (105 upregulated and 269 downregulated); and 526

genes in R-C (270 upregulated and 256 downregulated). Of these, 97 genes were commonly

differentially expressed in all 3 comparisons [including *CYP4C27* (AGAP009246), *SULTD1*

(AGAP012672), *aminopeptidase N1*, and diverse cuticular proteins, e.g. chitinase (*Cht24*,

AGAP006191, *CPFL1* (AGAP010902), *CPCFC1* (AGAP007980), *CPLCX3*, *CPR24*

(AGAP005999), *CPR106* (AGAP006095) and *CPR130* (AGAP000047). Finally, for Cameroon,

376 genes were differentially expressed in R-S comparison (204 upregulated and 172 down-

regulated) (Figure S3d); 932 in C-S (778 upregulated and 154 downregulated); and 116 genes

in R-C [only 9 upregulated and 107 downregulated (probably due to a single, low quality

replicate in the raw data from Cameroon (Figure S2). Not surprising, only 7 genes were

commonly differentially expressed in all 3 comparisons. These include the highly upregulated

gene, *GSTe2* (AGAP009194) and *chymotrypsin-1* (AGAP006709).

All data analysed together (Figure S3e) revealed no single gene differentially expressed in

common; possibly due to the low quality with the Chad unexposed (C) data. Analysis of data

from Nigeria, Niger and Chad, revealed a single gene (AGAP000046, transporter major

facilitator superfamily) differentially expressed across all countries (Figure S3f). Niger and

Chad shared *hexamerin* (AGAP010658), *aminopeptidase N1*, and unknown protein,

AGAP0290967; Nigeria and Niger share only a single gene, AGAP003248; while seven genes

were common to Nigeria and Chad, including *CPLCX3*.

Principal component analysis for the top 500 most variable genes in all experimental arms

revealed data from field samples (R and C) from all four countries clustering closer in PC1

and PC2 axes, away from the data from the susceptible females, Ngoussou (Figure S4).

Analysis of the common differentially expressed genes across the Sahel

The most differentially expressed genes were presented in Figure 1, a volcano plot of fold

change *vs* significance levels for Nigeria and Niger, and Figure 2, for Chad and Cameroon.

Exhaustive lits of these genes of interest is provided in File S2. Comparisons of genes

commonly, upregulated and/or downregulated in R-S/R-C/C-S, from the four countries

revealed similar transcriptomic profiles between R and C compared with the S. The most

commonly and consistently overexpressed genes across the Sahel (taking account mean

expressions) are the chymotrypsins, 3, -2 and -1 (*CHYM3*/AGAP006711 and

CHYM2/AGAP006710 and *CHYM1*/AGAP006709) (Figures 1 and 2, File S2), the glutathione

S-transferase, *GSTe2* (AGAP009194), an aquaporin, *AQP3* (AGAP010326), *CYP6Z2*

(AGAP008218), *CYP6Z3* (AGAP008217), *CYP4C27* (AGAP009246), a chitinase, *Cht24*

(AGAP006191), a thioester-containing protein-1, *TEP-1* (AGAP010815),), a trehalose 6-

phosphate synthase/phosphatase, *TPS*1/2 (AGAP008227), a lipase (AGAP002353), and

AGAP012818 (V-type protein ATPase subunit A). More on these genes is provided in the

sections, below.

Other genes commonly overexpressed in data from two or three countries include

AGAP005501 (dehydrogenase/reductase SDR family 11) upregulated in Nigeria and Niger (R-

S and C-S comparisons), and in R-S for Chad and Cameroon; AGAP008091 (CLIP-domain

serine protease, *CLIPE1*), upregulated in all countries in R-S and C-S comparisons; a chitinase,

Cht5-5, AGAP013260) upregulated in Nigeria, Nigeria and Chad R-S and C-S, with both

induced upregulated in R-C in Chad, as heme peroxidase (*HPX15*), upregulated in R-S and C-

S comparisons in Nigeria, Chad and Cameroon, as well as malate dehydrogenase

(AGAP000184), upregulated in Nigeria, Chad and Cameroon, R-S and C-S.

However, the most overexpressed genes in Nigeria are a cubilin, a histone (H2B), carbonic

anhydrase I, and a cuticular protein, *CPR131* (File S2), while for Cameroon its H2B, Serpin 9

inhibitory serine protease inhibitor, galectin 4 and a Protease m1 zinc metalloprotease.

Several genes were significantly downregulated across the four countries, particularly the

cytochrome P450s and the ATP-binding cassette transporters. The most consistently down-

regulated P450s were *CYP6AK1* (AGAP010961), downregulated in R-S and C-S comparisons

in Nigeria, Niger and Cameroon, while down-regulated in all comparisons in Chad; *CYP4H24*

(AGAP013490), downregulated in R-S and C-S comparisons for Nigeria and Cameroon, and

downregulated in all comparisons in Niger and Chad; *CYP6M4* (AGAP008214),

downregulated in R-S and C-S comparisons in Nigeria, Niger and Cameroon, while

downregulated in all comparisons in Chad; and finally *CYP6P5* (AGAP002866),

downregulated in R-S and C-S comparisons in Niger and Chad, but only in C-S in Nigeria.

Analysis of commonly overexpressed metabolic resistance genes resistance across the Sahel

Special attention was given to the known metabolic resistance genes, implicated in insecticide

resistance in *Anopheles* and/or other insects, including, CYP450s, GSTs, carboxylesterases,

cuticular proteins, chemosensory proteins/SAPs, uridine diphosphoglucuronosyltransferases,

etc, in addition to other important genes, e.g., the immune proteins. Analysis of the data from

the list of genes significantly, differentially expressed (FDR-adjusted $p < 0.05$ and $\log_2 FC = 1/$

$FC \geq 2$) revealed the major, commonly upregulated metabolic resistance genes in the four

Sahel countries. Top 50 genes from these lists (File S3) from each country are displayed as

heatmaps of fold changes in Figure S5a, -b, -c and -d). The most common genes linked with

resistance and/or other physiologically important phenotypes are tabulated in Table 1,

showing fold changes (R-S and C-S comparisons) of 55 genes, common to all countries, or

three or two countries at least. The most commonly and consistently overexpressed genes

across the Sahel (taking account mean expressions) are the chymotrypsins, -1, -2 and -3

(*CHYM1*/AGAP006709, *CHYM3*/AGAP006711 and *CHYM2*/AGAP006710) (Table 1, Figure S5,

File S2 Results), with fold change for *CHYM1* in R-S and C-S comparisons of 31.62 and 54.41

for Nigeria, 12.31 and 41.59 for Niger, and 18.00 and 18.08 for Chad, and 5.69 and 11.77 for

Cameroon; for *CHYM3*, 32.45 and 58.43 for Nigeria R-S and C-S, respectively, 7.30 and 43.27

for Niger, 13.40 and 7.24 for Chad, and 2.59 and 8.15 for Cameroon. Of *Anopheles* metabolic

resistance genes, *GSTe2* (AGAP009194) was the most consistently upregulated gene across the

Sahel, FC = 9.61 and 7.16, respectively for Nigeria R-S and C-S comparisons; 6.55 and 11.32 for

Niger; 9.09 and 10.02 for Chad, 9.68 and 20.67 for Cameroon. Other common GST was *GSTZ1*,

upregulated in three countries (Nigeria, Niger and Chad) are *GSTe4*, *GSTU1*, *GSTD1-4* and

GSTD3.

Other most consistently and highly upregulated genes include a lipase (AGAP002353, FC of

26.73 and 20.80 for Nigeria R-S and C-S comparisons; 16.53 and 29.76 for Niger, 14.95 and 9.95

for Chad, 14.58 and 14.53 for Cameroon); a chitinase, *Cht24* (AGAP006191, FC = 21.12 and

38.55 for Nigeria R-S and C-S, 40.08 and 67.18 for Niger, 91.42 and 5.71 for Chad, 44.26 and

40.53 for Cameroon); a V-type protein ATPase subunit a (AGAP012818, FC = 22.12 and 14.61

for Nigeria, 31.05 and 25.43 for Nigeri, 6.00 and 4.31 for Chad, 44.92 and 43.4 for Cameroon),

and an aquaporin, *AQP3* (AGAP010326, FC = 14.81 and 11.03 for Nigeria, 12.09 and 21.94 for

Niger, 14.86 and 1.21 for Chad, 14.17 and 17.3 for Cameroon).

The most overrepresented gene family, and most consistently upregulated across the Sahel

were the cytochrome P450s, with *CYP6Z2*, *CYP6Z3*, and *CYP4C27* taking the front seat, in

addition to the two P450s linked to cuticular resistance, *CYP4G16* (upregulated in Nigeria and

Chad) and *CYP4G17* (upregulated in Nigeria, Niger and Chad). The three P450s, *CYP6Z2,*

CYP6Z3 and CYP4C27 are consistently overexpressed: FC for *CYP6Z2* is 6.55 and 2.10 for

Nigeria R-S and C-S comparisons, 6.73 and 2.32 for Niger, 13.96 and 1.48 for Chad, 11.84 and

8.39 for Cameroon; FC for *CYP6Z3* of 4.73 and 1.82 for Nigeria,   6.09 and 1.42 for Niger,

3.51 and 1.11 for Chad, 7.11 and 3.79 for Cameroon; and FC for *CYP4C27* is 7.36 and 4.6 for

Nigeria, 4.03 and 5.64 for Niger, 6.04 and 2.62 for Chad, 2.01 and 2.63 for Cameroon. The two

well-known insecticide resistance genes *CYP6P3* and *CYP9K1* did not future prominently

across the Sahel of these countries, neither did *CYP6M2*.

The second most commonly overrepresented gene families belong to cuticular proteins,

including cuticular proteins Rebers and Riddiford (RR), several cuticular proteins of low

complexity (CPLC) and chitin-binding cuticular proteins (CPAP), with the commonly

upregulated ones across all four countries being *CPR75* (AGAP009871) and *CPR76*

(AGAP009874).

Two carboxylesterases picture prominently across the Sahel – the beta esterase, *COEBE3C*

(AGAP005372) upregulated in all four countries, while *COEJHE5E* was common to Nigeria,

Niger and Chad. For Phase II metabolism enzymes, several uridine-diphospho-

glucuronosyltransferases, UGTs were upregulated, including the *UGT-B19*, which is

upregulated in Nigeria and Niger (Table 1), and unknown UGT (AGAP006222) upregulated

across all the countries (File S2).

Other genes in the top 55 metabolic genes include the malate dehydrogenase, *TPS* (*TPS1/2*),

AQP2, thioester-containing proteins, *TEP1, -12* and *-14,* a cyanogenic-beta-glucosidase

(AGAP006422) and three chemosensory proteins, *CSP5,* sensory appendage proteins, -1 and -

2.

547 **Table 1: Common detoxification and metabolic genes differentially and constitutively**
548 **upregulated in Sahel *An. coluzzii* at FDR-adjusted p < 0.05.**

	NIGERIA		NIGER		CHAD		CAMEROON		
Gene	**R-S**	**C-S**	**R-S**	**C-S**	**R-S**	**C-S**	**R-S**	**C-S**	**Gene Description**
AGAP006400	2.19	4.18	1.72	4.47	-	-	2.11	2.82	alkaline phosphatase 2
AGAP028491	4.34	2.83	1.42	2.66	2.98	2.57	-	-	aquaporin, AQP2
AGAP010326	14.81	11.03	12.09	21.94	14.86	1.21	14.17	17.3	Aquaporin, AQP3
AGAP005372	9.45	12.19	3.34	4.25	6.60	3.62	1.39	2.63	carboxylesterase, COEBE3C
AGAP005837	4.70	5.02	1.41	3.29	1.89	2.64			carboxylesterase, COEJHE5E
AGAP004904	3.88	2.19	-	-	2.34	1.74	-	-	catalase, CAT1
AGAP029127	4.14	3.17	-	-	1.73	2.22	-	-	chemosensory protein 5, CSP5
AGAP006191	21.12	38.55	40.08	67.18	91.42	5.71	44.26	40.53	chitinase, Cht24
AGAP013260	4.40	11.47	1.26	3.89	3.13	0.75	-	-	chitinase, Cht5-5
AGAP006898	8.97	5.52	-	-	2.23	3.93	-	-	chitinase, Cht6
AGAP006709	31.62	54.41	12.31	41.59	18.00	18.08	5.69	11.77	chymotrypsin-1, CHYM1
AGAP006710	9.85	19.44	2.07	8.75	4.21	3.87	-	-	chymotrypsin-2, CHYM2
AGAP006711	32.54	58.43	7.30	43.27	13.40	7.24	2.59	8.15	chymotrypsin-3, CHYM3
AGAP000987	4.28	2.70	-	-	1.34	2.61	-	-	cuticular protein, CPAP3-A1b
AGAP000988	5.33	1.56	-	-	0.74	5.44	-	-	cuticular protein, CPAP3-A1c
AGAP000986	7.31	4.60	-	-	1.73	3.46	-	-	cuticular protein, CPAP3-D
AGAP006149	3.67	0.78	-	-	0.74	5.78	-	-	cuticular protein, CPLCX3
AGAP010123	33.33	54.97	2.35	13.2	-	-	-	-	cuticular protein, CPR131
AGAP010123	33.33	54.97	2.35	13.20					cuticular protein, CPR131
AGAP005456	5.24	1.78	-	-	1.24	2.21	1.07	2.07	cuticular protein, CPR15
AGAP006009	-	-	2.88	1.30	-	-	3.89	1.62	cuticular protein, CPR30
AGAP009871	8.67	2.72	1.07	2.63	2.23	4.18	1.67	2.56	cuticular protein, CPR75
AGAP009874	10.64	2.22	1.05	2.55	3.69	5.56	1.09	2.03	cuticular protein, CPR76
AGAP009879	7.43	2.57	-	-	1.87	4.6	-	-	cuticular protein, CPR81
AGAP006422	7.50	9.41	-	-	8.44	7.33	2.11	1.96	cyanogenic beta-glucosidase
AGAP002417	5.53	4.17	0.95	1.22	1.90	1.87	-	-	cytochrome P450, CYP4AR1
AGAP009246	7.36	4.6	4.03	5.64	6.04	2.62	2.01	2.63	cytochrome P450, CYP4C27
AGAP012957	5.01	2.83	-	-	2.88	2.45	2.04	2.61	cytochrome P450, CYP4D17
AGAP001076	5.79	4.69	1.45	1.80	3.81	2.41	0.88	1.48	cytochrome P450, CYP4G16
AGAP000877	3.81	1.96	2.34	1.85	3.76	2.05	1.20	2.05	cytochrome P450, CYP4G17
AGAP008358	-	-	2.96	4.08	3.43	3.28	3.61	1.99	cytochrome P450, CYP4H17
AGAP007480	5.36	3.99	1.31	2.73	2.49	2.01	-	-	cytochrome P450, CYP6AH1
AGAP002865	1.11	0.50	2.29	1.87	0.66	12.83	2.06	2.26	cytochrome P450, CYP6P3
AGAP008207	3.56	1.96	-	-	2.08	1.64	1.74	2.35	cytochrome P450, CYP6Y2
AGAP008218	6.55	2.10	6.73	2.32	13.96	1.48	11.84	8.39	cytochrome P450, CYP6Z2
AGAP008217	4.73	1.82	6.09	1.42	3.51	1.11	7.11	3.79	cytochrome P450, CYP6Z3
AGAP000818	2.97	1.15	0.75	0.54	0.90	1.59	3.38	2.21	cytochrome P450, CYP9K1
AGAP010400	3.58	2.56	1.62	2.83	3.29	1.44	-	-	flavin-containing monooxygenase
AGAP007920	5.72	2.40	2.84	1.84	-	-	-	-	glucuronosyltransferases, UGT-B19
AGAP004164	2.19	1.85	1.68	2.34	1.39	1.30	-	-	glutathione S-transferase, GSTD1-4
AGAP009194	9.61	7.16	6.55	11.32	9.09	10.02	9.68	20.67	glutathione S-transferase, GSTe2
AGAP009193	3.26	2.29	1.14	2.09	1.46	1.6	-	-	glutathione S-transferase, GSTe4
AGAP000947	4.36	2.31	1.23	2.18	1.91	3.15	-	-	glutathione S-transferase, GSTU1
AGAP002898	5.52	3.51	1.66	2.61	2.76	2.38	-	-	glutathione S-transferase, GSTZ1
AGAP002198	-	-	2.00	5.77	4.75	6.58	-	-	glycine-N-methyltransferase
AGAP013327	-	-	1.16	7.28	7.89	20.11	5.11	14.82	heme peroxidase, HXP15
AGAP002353	26.73	20.80	16.53	29.76	14.95	9.95	14.58	14.53	lipase
AGAP000184	5.16	6.26	-	-	4.87	4.31	1.21	1.90	malate dehydrogenase
AGAP008051	3.65	3.96	1.14	1.30	1.73	1.68	-	-	sensory appendage protein 1, SAP1
AGAP008052	5.47	10.84	1.02	1.45	3.45	1.06	-	-	sensory appendage protein 2, SAP2
AGAP010815	5.68	14.41	4.07	8.55	7.75	3.79	1.65	3.06	thioester-containing protein 1, TEP1
AGAP008368	3.34	6.12	1.39	3.85	3.42	1.97	-	-	thioester-containing protein, TEP14
AGAP008654	4.59	17.36	2.12	10.03	-	-	-	-	thioester-containing protein, TEP12
AGAP008227	4.07	4.00	3.99	2.00	4.59	1.47	2.58	4.12	trehalose 6-phosphate synth/phosph
AGAP012818	22.12	14.61	31.05	25.43	6.00	4.31	44.92	43.4	V-type proton ATPase, subunit a

Gene Ontology Enrichment Analysis

Gene ontology enrichment analysis, for genes significant in comparisons ($\log_2$FC = 1, p < 0.05)
revealed crucial differences in overrepresented GO terms between the up- and -
downregulated genes, in data from the four countries. For example, the most over-
represented, semantically similar GO terms associated with xenobiotics metabolism is
oxidoreductase activity ((highest frequency of 12.9 %, Figure S6a, REVIGO Table View), which
cluster together in Nigerian R vs S comparison (for genes upregulated in R). Other over-
represented GO terms in this comparison include glutathione S-transferase activity (the most
enriched/specific term), peroxidase activity, odorant binding and chitin binding. In contrast,
GO terms over-represented in R vs S (down-regulated in R) were mostly involved in
neurotransmission, metal ions binding and receptor channelling activities (Figure S6b).

For R vs C comparison (upregulated in C) the over-represented GO terms include
oxidoreductase (frequency = 12.88 %), glutathione S-transferase activity (the most
enriched/specific term), peroxidase activity, glucuronosyltransferase activity, aldehyde
oxidase activity, chitin binding, carbohydrate binding activities, etc (Figure S6c). In contrast
the most enriched GO terms, downregulated in C were endopeptidase activities, Toll binding,
oxidoreductase (frequency = 1.21 %) and monooxygenase activities (Figure S6d). For the rest
of the three countries contrasts were also observed, between phenotypes, and detailed in File
S1, Results section.

Quantitative PCR validation of expression profiles of metabolic resistance genes

The relative expression levels of 12 metabolic resistance genes were validated using the qRT-
PCR. The qRT-PCR results support the transcriptomic patterns obtained from the RNAseq
analysis, with *GSTe2* as the most overexpressed gene, followed by *CYP6Z2* (Figure 3). For
example, for R-S and C-S comparisons *GSTe2* had fold changes of 49.25 ± 5.05 and 28.45 ± 4.45,
for Nigeria ($p \leq 0.0001$); 37.01 ± 2.22 and 33.34 ± 6.61 for Niger ($p \leq 0.0001$); 29.97 ± 5.39 and
30.16 ± 6.16 for Chad ($p \leq 0.0001$); and 28.30 ± 3.30 ($p \leq 0.0001$) and 17.03 ± 7.03 for Cameroon
($p \leq 0.001$). *CYP6Z2* has R-S and C-S fold changes of 15.93 ± 2.54 ($p \leq 0.001$) and 7.35 ± 1.52
($p \leq 0.05$), for Nigeria; 24.21 ± 2.17 ($p \leq 0.001$) and 20.83 ± 3.02 for Niger ($p \leq 0.001$); 15.02

($p \le 0.001$) ± 2.01 and 6.09 ± 2.08 for Chad ($p \le 0.05$); and 10.58 ± 3.00 ($p \le 0.01$) and 3.43 ± 1.43

for Cameroon. Correlation analyses for R-S comparisons for all genes revealed positive and

significant association (Figure S7) between the RNA-seq and qRT-PCR data for these genes in

data from Niger ($R^2 = 0.58$, p = 0.03) and Chad ($R^2 = 0.408$, p = 0.025), with positive, but non-

significance seen in Cameron ($R^2 = 0.58$ p = 0.08) and Nigerian ($R^2 = 0.503$, p = 0.06) data.

Detection of Signatures of Selective Sweeps

Signatures of selection were investigated in the major metabolic resistance genes, by

estimating Tajima's D per gene within populations, and Fst per gene between population

pairs. These tests of neutrality revealed several genes exhibiting high genetic differentiation,

or possibly undergoing expansion. Among the top 100 most overexpressed metabolic genes

across the four countries (File S2), 13 genes were possibly undergoing genetic differentiation.

These include *TEP1*, and *TEP3*, with average Tajima's D of -1.4 and -1.6 respectively, in

populations from Chad, and average Fst values of 0.58 and 0.49 for all countries (File S3,

provide the number of SNPs and chromosomal location for each gene), *CYP9K1* (average

Tajima's D = -1.00 for Cameroon, and combined Fst = 0.55), *CPR15* (average Tajima's D = -

0.94, and Fst = 0.17); *CPAP3-A1b* (Tajima's D = -0.70 in Chad, -0.95 in Nigeria, and -1.32 in

Cameroon, with combined Fst of 0.55) and *CPLCX3*. Several genes from the GST family

exhibited strong genetic differentiation with reduced variations prominently in *GSTe2* (< 3

SNPs in Chad and Cameroon samples), with Tajima's D of -1.45 and -1.15 for Nigeria and

Niger, and combined Fst of 0.02. Detailed analysis is provided in File S1, and File S3 contain

the respective frequencies of these variables.

Identificaton of genetics variants associated with insecticide resistance

The RNA-Seq-Pop workflow calculated allele frequencies of variants of interest found in the

raw RNA-sequencing read data. Figure 4 displays allele frequencies of each variant in

respective treatment replicates used for the RNAseq, as well as Ngoussou. Several mutaions

were found within the voltage-gated sodium channel. In addition to the L995F *kdr* mutation,

the recently identified V402L and I1527T mutations exist in all the field populations across the

Sahel, at frequencies ranging from 29 % to 67 %. We also detected the N1570Y mutation, which

shares the same haplotype as L995F. The G280S *acetylcholinesterase*-1 mutation was also

observed in low to moderate frequencies in all populations except for Ngoussou and samples

from Niger. The L43F pyrethroid resistance marker of *CYP4J5* was also seen, fixed in most

populations.

Detection of chromosomal inversion polymorphisms and metabolic genes within the

inversion breakpoints

The frequency of the major inversion polymorphisms in chromosome 2 were calculated,

considering ploidy (8 individual female mosquitoes were pooled for RNA extraction, for each

replicate). Table S5 provided the frequencies of the respective inversions for the populations

from each country, as well as for Ngoussou. High frequency of three Anopheles

phenotypically important inversions, the 2La, 2Rb and 2Rc, in contrast to the 2Rd, 2Rj and

2Ru which were in lower frequencies [except for the 2Rj in Nigeria (30.38 %) and 2Ru in

Cameroon (24.29 %)]. The 2La inversion was found fixed in the field populations (100 % in

Nigeria and Cameroon, 99.92 % in Niger and 99.21 % in Chad), in contrast with Ngoussou,

with frequency of only 6.26 %. Similar pattern was observed with the 2Rb inversion, with high

frequencies in three field populations (79.01 %, 84.85 %, 80.22 % in Nigeria, Niger and Chad),

but lower in Cameroon (29.08 %) and Ngoussou (5.35 %). Frequencies of the 2Rc inversion

were 88.54 %, 85.52 %, 88.82 %, 46.99 % and 9.89 % for Nigeria, Niger, Chad, Cameroon and

Ngoussou respectively. Some genes among the top 100 most overexpressed metabolic

resistance genes (above) and which were likely undergoing recent population expansion were

located within the 2La, 2Rb and 2Rc inversions (File S4). These include the *hsp83*

(AGAP006958) and molecular chaperone hptG (AGAP006961), *CPLCX3* (AGAP006149),

CPLCA1 (AGAP006145,) within the inverted 2La chromosomal arm. Other genes within 2La

inversion include a group of chitinase enzymes (*Cht6*, *-8*, *-11* and *-24*, most notably the *Cht24*

and *Cht6*, which are among the top 50 most overexpressed genes in all countries. Other genes

include *COEJHE5E*, one of the 14 carboxylesterases in the 2La region, and which was among

the top 100 most overexpressed genes across the Sahel; the most overexpressed chymotrypsin

genes, *CHYM1* and *CHYM2*; three ionotropic receptors, *IR136* (AGAP006440), *IR139*

(AGAP006691) and *IR142* (AGAP006407); as well as five P450s, *CYP301A1*, *CYP302A1*,

CYP4J5, CYP4J9 and *CYP4J10*. The most overrepresented genes within 2La inversions are the

CPR cuticular proteins (73 in total), with *CPR21, -26, -30, -59* and *-140* from the list of the top

100 resistance genes from the Sahel.

Carbonic anhydrase I (the most overexpressed metabolic resistance in Nigeria, AGAP013402

is sit within 2Rb inversion region). Four chitinases (*Cht4*, *Cht5-1*, *Cht5-3*, and *Cht5-5*) are

located within 2Rb inversion; *Cht5-1* and *Cht5-5* were among the observed top 50 most

overexpressed metabolic genes in the Sahel. Lipase (*AGAP002353*) which is the among the top

6 overexpressed metabolic gene in the four countries sit within the 2Rb inversion. Eight

CYP450s reside in the 2Rb inversion, including *CYP4D15*, *CYP4D17* and *CYP4AR1*, which

were among the top 100 metabolic resistance genes. Three cuticular proteins, *CPR7, -8* and *-9*

were also within the 2Rb inversion.

Interesting genes sitting within 2Rc inversion include *GSTZ1*; 5 ionotropic receptor genes, *IR7i*

(AGAP013363), *IR7u* (AGAP013285), *IR7t* (AGAP002763), *IR7w* (AGAP013416) and *IR41a*

(AGAP002904); a carboxylesterase, *COEAE6O* (AGAP002863); and 11 CYP450s, including

CYP6AA1 (AGAP002862), *CYP6AA2* (AGAP013128), *CYP6P15P* (AGAP002864), *CYP6P3*

(AGAP002865), *CYP6P5* (AGAP002866), *CYP6P4* (AGAP002867), *CYP6P1* (AGAP002868),

CYP6P2 (AGAP002869), *CYP6AD1* (AGAP002870) and *CYP6Z4* (AGAP002894)

Investigation of the polymorphism in the coding sequences of Sahelian GSTe2 and CYP6Z2

Analysis of the polymorphism patterns of full-length cDNA sequences of *GSTe2* (666 bp) and

CYP6Z2 (1479 bp) from the Sahel region of Africa revealed complete homogeneity for *GSTe2*,

with no single polymorphism detected in the field populations as well as Ngoussou (all

sequences identical to AGAP009194 reference). This suggests a fixed allele, consistent with the

observation from the analyses from the fixation index (File S3). For *CYP6Z2* (1479 bp),

homogeneity was observed within each country and Ngoussou, except for Niger,

characterised by an unusually high polymorphism (Figure S8a, -b). *CYP6Z2* is polymorphic

with 10 haplotypes across the Sahel, with 75 polymorphic sites of which 65 were synonymous,

and 12 led to amino acids substitutions. The bulk of the polymorphisms were contributed

from larger variations in the Niger and Cameroon (S = 42 and 23 respectively), while highest

homogeneity was observed in Chad, with a single haplotype. Haplotype diversity is high

(H_d = 0.921), from 10 haplotypes out of only 20 sequences, with the lowest H_d in the Chad

sequences, suggesting a directional selection/fixed allele. The haplotypes cluster according to

origin on the maximum likelihood phylogenetic tree, except for Niger (Figure S8c).

Investigation of the role of intergenic region regulatory elements in overexpression of GSTe2

and CYP6Z2

Investigation of polymorphism in the intergenic region/regulatory elements

To investigate polymorphisms in the regulatory elements of the above genes, the 351

intergenic regions of *GSTe2* (spanning the 43 bp 3'-UTR of *GSTe1*, 248 bp flanking sequence

and 60 bp 5'-UTR of *GSTe2*) preceding the start codon were amplified from 10 each of DDT-

alive and -dead females from the 4 Sahel countries, as well as from Ngoussou females,

successfully. For *CYP6Z2*, 1078 bp (spanning 38 bp 3' UTR of *CYP6Z1*, 937 bp flanking

sequence and 103 bp 5'UTR of *CYP6Z2*, preceding the start codon of *CYP6Z2*) were used to

amplify fragments from 10 each of deltamethrin-alive and -dead females from Nigeria, and

from Ngoussou females.

Out of the 90 *GSTe2* 5'-UTR sequences analysed differences were observed between the alive

and dead mosquitoes, with a total of 65 sequences from alive and dead females (regardless of

country of origin) identical to the 10 sequences of Ngoussou. From the 40 sequences of the

dead females, 39 were identical to Ngoussou (regardless of the country). Twelve sequences,

all from the alive females (from across the four countries) were similar, with several mutations

in putative transcriptional factors binding sites, which may impact overexpression of the

GSTe2. In short, 8 mutations were shared in common between these 12 sequences of DDT-

alive females from across the countries (3 each from Nigeria, Chad and Cameroon) and

additional mutations in the Niger samples (4 sequences). These mutations include (i) T->A

transition withing the cellular myb-DNA (c-myb) binding domain (Figure S9), (ii) a T->C

transversion in the zinc-finger homeodomain, δEF1 (δ elongation factor 1) binding site, (iii) T-

>A transition in the nuclear matric protein 4 (NMP4), (iv) simultaneous insertion of adenine

and a transition T -> C, in positions 113 and 114 respectively, between the Fork-head box L1

and c-EST binding sites, (v) an A -> C transversion in a second NMP4, (vi) a T -> C transition,

six nucleotides downstream the nuclear factor κB (NF-κB), (vii) followed by an A -> G

transversion 3 nucleotides downstream the NF-κB, (viii) a G -> A transversion, 7 nucleotides

upstream the GC box, (ix) a C -> G transition, 2 nucleotides downstream the arthropod

initiator (Inr consensus) sequence, and finally (x) a C -> G transition, within the 5'-UTR of the

GSTe2, 28 nucleotides downstream the transcriptional start site/49 nucleotides upstream the

GSTe2, start codon.

Analysis of the 90 sequences revealed a very low polymorphism in the dead mosquitoes (S =

0, for Niger-dead, Chad-dead and Ngoussou), but high polymorphism in the alive (S = 16 for

Nigeria-alive, and 13 each for Chad-alive and Cameroon-alive) (Table S6). All sequences

produced 17 polymorphic sites, with 6 haplotypes (Figure S10a and -b). High haplotype

diversities were obtained from the alive mosquitoes (for example, Nigeria alive, Hd = 0.71,

Niger-alive 0.53). Regardless of country of origin, the haplotypes cluster according to

phenotype on the maximum likelihood phylogenetic tree, with the alive haplotypes forming

a distinct/separate clade (Figure S10c).

With regard to *CYP6Z2* no major differences were observed when comparing the

deltamethrin-alive and -dead sequences with the Ngoussou.

Comparative measurement of activities of the 5' regulatory elements of the GSTe2 and CYP6Z2

Initial promoter analyses were conducted with the 5'- regulatory element sequences of *GSTe2*,

for the DDT-alive females (representative sequence with the 8 common polymorphic

positions, designated, Sahel-alive), comparing it with the sequence from the DDT-

dead/Ngoussou (designated Ngoussou/Sahel-dead). For *CYP6Z2*, a predominant sequence

from the Nigeria field sample (with 8 nucleotide insertion) was compared with the Ngoussou

sequence.

The ability to drive heterologous expression of the firefly luciferase was determined, with the

Sahel-alive construct and Ngoussou/Sahel-dead producing increased luciferase activity

[~3090× (normalized luciferase activity = 2.58) and 479× (normalized activity = 0.400),

respectively] compared with the promoter pGL3-Basic vector. But the Sahel-alive construct

was significantly more active than the Ngoussou/Sahel dead counterpart [promoter activity

~6-fold higher (Tukey HSD Q = 8.11, p = 0.004)]. In contrast, for *CYP6Z2* no significant

differences were observed when comparing the field construct (normalized luciferase activity

= 5.95 for alive/dead field construct), compared with 5.84 for the Ngoussou construct.

GSTe2 promoter delineation and measurement of activity

Sequential deletion of the intergenic region of the *GSTe2* resulted in progressive reduction in

luciferase activity. Deletion of the 43 bp 3'-UTR of the *GSTe1* (-308 from the start codon of

GSTe2) reduced activity of the Sahel-alive construct by only 18.6 % suggesting that the c-Myb

transcriptional factor binding site may not be critical for overexpression (Figure 5). Deletion

of an additional 38 nucleotides from the flanking region (-270 fragment, obliterating the δEF1

binding site) had comparable impact as above, with activities reducing by 15.3 % only.

Removing the *GSTe-1* 3'-UTR and an additional 46 nucleotides from the flanking region (-262

fragment, which obliterated in addition the FOX-L1 putative binding site) reduced activity by

64.1 % (Q = 6.18, p = 0.01) indicating the importance of this binding site. Deletion of *GSTe1* 3'-

UTR plus 75 nucleotides of the flanking region (-232 fragment, located 10 nucleotides from

the c-EST binding site) significantly reduced the activity by ~70% (Q = 6.14, p = 0.01),

suggesting the importance of the simultaneous insertion of adenine and transition of T to C,

in position 113 and 114 respectively, between the FOX-L1 and c-EST binding sites. Removing

the fragment of the 5'UTR of *GSTe2* (43 nucleotides preceding the AUG codon obliterated the

promoter activity, reducing the luciferase expression by 98 % (Q = 9.41, p = 0.002). This is

despite presence of all the above binding sites and transcriptional start site in this fragment.

Deletion of the 5'-UTR from Ngoussou significantly reduced activities as well (reduction by

88 % compared with the full Ngoussou intergenic region construct, Q = 13.75, p = 0.001).

Overall, these findings suggest that the essential binding sites for overexpression of *GSTe2*

span the FOX-L1 and c-EST binding sites, with the 5'-UTR essential for activity.

Investigating the role of GSTe2 and CYP6Z2 in insecticides resistance using transgenic

analysis

A qRT-PCR was conducted using the transgenic flies to first establish overexpression of the above genes. Relative fold changes (FC) of 32.7 ± 4.6 and 25.30 ± 2.8 were obtained in flies overexpressing the *GSTe2* and *CYP6Z2* respectively, compared with control flies (progenies of crosses between the parental line flies with no gene insertion, crossed with GAL4/UAS driver line) (Figure S11).

Contact bioassays carried out using 0.05 % α-cypermethrin revealed a high susceptibility in the transgenic flies expressing *GSTe2* (Act5C-GAL4-UAS-GSTe2) and controls (Figure 6a), with mortalities increasing from 62 % and 70 % respectively in 1 h, to 94 % and 99 % in 24 h. However, significantly reduced mortalities were observed in transgenic flies expressing *CYP6Z2* (Act5C-GAL4-UAS-CYP6Z2) compared to control flies, at 1 h (mortality = 35 % *vs* 70 % in control, $p < 0.001$) and 3 h (mortality = 52 % vs 89 % in the control, $p < 0.001$). High susceptibilities were also seen in all the experimental flies exposed to 0.05 % deltamethrin, except for 1 h with flies expressing *CYP6Z2* (mortality = 47 % compared with 58 % in the control flies, $p < 0.010$). The *GSTe2* transgenic flies were highly resistant to 2 % permethrin, with no mortality at all in 1 h (Figure 6b), and average mortalities of only 8.5 % at 24 h ($p < 0.0001$), compared with 85 % at 1 h for control flies, which increased to 100 % from 3 h.

Initial exposure to 4 % of 3-phenoxybenzaldehyde (PBAld) and 3-phenoxybenzyl alcohol (PBAlc) had no toxic effect on flies (Figure not shown). However, 5x concentration of these primary products of pyrethroid hydrolysis induced mortalities, albeit low in all flies (Figure 6c). For PBAld, mortalities ranged from 15.5 % and 17.5 % at 1 h, for *CYP6Z2* transgenic flies and control flies respectively, to 30 % and 27.5 % at 24 h. Although low mortalities were observed as well with PBAlc, but at 1 h and 3 h exposure times the mortalities in the transgenic flies expressing *CYP6Z2* were significantly lower compared with mortalities from the control flies (1 h mortality = 2.5 % *vs* 12.5 %, $p < 0.001$; 3 h mortality = 9.5 % *vs* 17.5 %, $p < 0.001$).

Marginal tolerance towards DDT was observed in the *CYP6Z2* transgenic flies when compared with the control flies, but only at 3 h ($p < 0.001$) and 6 h ($p < 0.01$) (Figure 6d). This is in contrast with the *GSTe2* transgenic flies, with very low mortalities, in ranges of 0 % to

9.52 % for 1 h to 24 h, when compared with control flies (mortality = 34 % in 1 h and 100 % in

6 – 24 h, p < 0.0001).

Susceptibility to clothianidin was very high (Figure 6d), but surprisingly the *CYP6Z2*

transgenic flies exhibited a contrasting phenotype, with significantly higher mortalities

compared with the control flies, at 1 h (mortality = 67 % vs 42 %, p < 0.001), 3 h (mortality = 71

% vs 53 %, p < 0.001) and 6 h (mortality = 78 % vs 62 %, p < 0.01).

Taken together, these results confirmed that over-expression of *GSTe2* alone is sufficient to

confer resistance to type I pyrethroid (permethrin) and DDT, while overexpression of *CYP6Z2*

alone, may confer marginal resistance to the α-cypermethrin and PBAlc.

Discussion

Escalating insecticide resistance across Africa (Ibrahim et al., 2019b, Riveron et al., 2019, Toe

et al., 2015, Riveron, 2018), if not tackled, will probably compromise the malaria control and

elimination efforts. Molecular markers of metabolic resistance e.g., (Lucas et al., 2019,

Mugenzi et al., 2019, Weedall et al., 2019), will support evidence-based control and resistance

management. Identification and validation of resistance markers in malaria vectors across

regions of sub-Saharan Africa can promote communication, cooperation and coordination

among malaria control/elimination programs, and allow control efforts to be tailored to the

vector species involved in transmission across borders (Shiff, 2002), and tracking of the

evolution and spread of resistance markers across regions (Riveron et al., 2014b). To support

malaria pre-elimination efforts in Africa, in this study we targeted the Sudan savannah and

Sahel (regions of Africa sharing similar eco-climatic conditions, and characterised by high

seasonal transmission), which are ideal for control and elimination of malaria using seasonal

vector control (Cairns et al., 2012) and chemoprevention.

***Anopheles coluzzii* is a major malaria vector across the Sudano-Sahelian transects**

Contrary to the previous observations that *An. arabiensis* tends to predominate in arid

savannas, while *An. gambiae* is the dominant species in humid forest zones (Lindsay, 1998,

Coetzee, 2000, Ranson et al., 2009), *An. coluzzii* has repeatedly been identified as the major

malaria vector in the Sudan savannah and Sahel of several, neighbouring countries (including

in northern Nigeria, southern Niger, central Chad, and northern Cameroon) (Ibrahim et al.,

2019a, Ibrahim et al., 2019b, Ibrahim et al., 2019c, Fadel et al., 2019, Ibrahim et al., 2021) in

recent years, suggesting that this vector has adapted well in drier regions of the Sahel

transects, and is probably predominating over *An. arabiensis* and *An. gambiae s.s*. This is not

surprising as this species exhibits higher exploitation of breeding sites associated with

anthropogenic activities, and behavioural plasticity to avoid predators (Gimonneau et al.,

2010), and is known to survive a long dry season *in situ*/aestivation, which allows it to

predominate, becoming the primary force of malaria transmission (Dao et al., 2014, Lehmann

et al., 2017). Also, photoperiod and lower nightly temperature have been shown to

significantly increase the longevity of the *An. coluzzii*; mechanisms which allow it to diapause

in the dry season and re-establish first in the early rainy season (Huestis et al., 2017).

Common metabolic genes mediate multiple resistance in the Sahelian *An. coluzzii*

Several genes shown to confer metabolic resistance in *Anopheles* mosquitoes and other insects

were found constitutively overexpressed and/or induced in this study. *GSTe2* (AGAP009194)

is one of the most regularly encountered metabolic genes in resistant populations of the major

malaria vectors *An. gambiae*, *An. coluzzii* and *An. funestus* (Lucas et al., 2019, Mitchell et al.,

2014, Riveron et al., 2014b, Menze et al., 2020). This gene has been validated in conferring

resistance to DDT (Mitchell et al., 2014) and fenitrothion (Adolfi et al., 2019) in *An. gambiae*. It

was extensively studied in *An. funestus*, in which it was shown to confer cross-resistance to

DDT and permethrin (Riveron et al., 2014b), reduce efficacy of the LLINs, PermaNet 2.0 and

PermaNet 3.0 (side panels) (Menze et al., 2020), and even increase the longevity of the resistant

populations carrying the 119F mutation (Tchouakui et al., 2018). These and our findings

suggest that the overexpression of this GST alone confer resistance to three insecticides from

three different classes (DDT, permethrin and fenitrothion).

Several cytochrome P450s previously linked with insecticide resistance were found

overexpressed across the Sahel. For example, *An. gambiae CYP6Z2* (AGAP008218) known to

metabolize carbaryl (Chiu et al., 2008), the insect juvenile hormone analogue insecticide,

pyriproxyfen (Yunta et al., 2016) and mitochondrial complex I inhibitors, fenazaquin,

pyridaben and tolfenpyrad (Lees et al., 2020). This P450 also plays a pivotal role in the

clearance of pyrethroid insecticides via further catabolism of pyrethroid derivatives (PBAld

and PBAlc) obtained by the action of carboxylesterases (Chandor-Proust et al., 2013), in line

with our findings of this gene conferring marginal resistance to PBAlc and α-cypermethrin.

However, our findings suggest that overexpression of this P450 may enhance the efficacy of

clothianidin, which will be epidemiologically advantageous in terms of control. Indeed,

bioactivation by P450s is known to be a requirement for insecticidal toxicity of several classes

of insecticides, e.g., the organophosphates and chlorpenafyr. Other important CYP450s found

to be commonly overexpressed across the Sahel include the *CYP4C27* (AGAP009246), *CYP6Z3*

*(*AGAP008217) and *CYP9K1* (AGAP000818), all three shown to be consistently overexpressed

in field populations of *An. gambiae/coluzzii* and *An. funestus* across Africa (Antonio-Nkondjio

et al., 2016, Toe et al., 2015, Riveron et al., 2017). *CYP9K1* has been shown to be

epidemiologically important pyrethroid-metabolising P450 linked with metabolism of

deltamethrin and pyriproxyfen (Vontas et al., 2018).

Several carboxylesterases were also upregulated/induced across the Sahel, with the *COEBE3C*

(AGAP005372) upregulated in all four countries. This beta esterase is enriched in the legs

(where xenobiotic detoxification probably occur) of pyrethroid-resistant *An. coluzzii* (Kefi et

al., 2021). Other genes consistently overexpressed across the Sahel, include the chymotrypsin

genes (*CHYM3*/AGAP006711 and *CHYM1*/AGAP006709) and a lipase (AGAP002353).

Chymotrypsins are known to defend insects against plants' proteinase inhibitors (Dunse,

2010); and previous transcriptional studies have shown *CHYM1* and *CHYM3* overexpressed

in insecticide resistant populations of *An. gambiae* and *An. coluzzii,* respectively (Toe et al.,

2015, Antonio-Nkondjio et al., 2016). Using *in vitro* and *in vivo* tools, lipases have been linked

with deltamethrin resistance in *Culex pipiens* pallens (Hu et al., 2020).

Common metabolic resistance markers probably exacerbate resistance across the Sahel

Several well-known genetic variants implicated in resistance, as well as the recently

discovered ones exist in high frequencies in the Sahelian *An. coluzzii*. For example, the

pyrethroid resistance L43F marker of *CYP4J5* (Weetman et al., 2018) was found fixed across

the Sahel. The G280S/G119S *ace*-1 mutation, found in high frequencies across Sahel confers

organophosphate an carbamates resistance (Weill et al., 2004), and is shown recently to confer

resistance to pirimiphos-methyl, in *An. coluzzii/gambiae* (Grau-Bove et al., 2021). Several

mutations found within the VGSC have recently been described/validated. For example, the

resistance mutation, L995F (Williams et al., 2022) and the V402L/I15227T haplotypes have

been observed across Africa (Clarkson et al., 2021). Recently, the two mutations

(V402L/I15227T) are described to be in tight linkage and mutually exclusive to the classical

L995F/S mutations (Williams et al., 2022), but our results suggest haplotypes carrying the

V402L/I1527T combination plus the L995 replacement do exist in the Sahel *An. coluzzii*. Not

only that, it is also in addition to the N1570Y replacement.

Evidence of cuticular resistance mechanism in the Sahelian *An. coluzzii*

Our results suggest cuticular mechanism playing role in pyrethroid resistance in these

populations. For example, the findings of *CYP4G16* (AGAP001076) and *CYP4G17*

(AGAP000877) overexpressed across the Sahel. The former P450 was previously shown to be

involved in epicuticular hydrocarbon biosynthesis associated with resistance (Balabanidou et

al., 2016).

The three major classes of the insect cuticular proteins - the CPR, CPLC and CPAP, were found

overrepresented in the top overexpressed genes across the Sahel, with the commonly

upregulated ones being *CPR76*, *CPR15* and *CPR30*, and with the *CPR15*, a chitin-binding

cuticular protein, *CPAP3-A1b*, and cuticular proteins of low complexity, *CPLCX3* and

CPLCA1. Indeed, *CPAP3-A1b* (AGAP000987) have been shown to be highly overexpressed in

deltamethrin-resistant Sahelian population of *An. coluzzii* from Burkina Faso (Toe et al., 2015)

and induced by blood feeding in *An. gambiae* (Vannini et al., 2014). The CPR and CPLC

cuticular proteins have been described to potentially play a crucial role in insecticide

resistance through leg cuticle remodelling/thickening, regulating penetration rate of

insecticides in *An. coluzzii* (Kefi et al., 2021). Furthermore, a recent study has found most of

the cuticular proteins we have described here, as highly overexpressed in

permethrin/malathion resistant populations of Ethiopian *An. arabiensis* (Messenger et al.,

2021), e.g., *CPR30*, *CPR75*, *CPR81*, and *CPLCP11*. Out of the several chitinases overexpressed

across the Sahel, four were amongst the top 50 most overexpressed metabolic genes. These include the *Cht24* (AGAP006191) and *Cht6* (AGAP006898) that have been shown to be overexpressed in the *An. arabiensis* from the above study (Messenger et al., 2021), with the ortholog of *Cht24*, AARA007329 among the top 10 most overexpressed genes in *An. arabiensis*, in line with our observations in *An. coluzzii* across the Sahel. There is an overwhelming need to functionally investigate the role/contribution towards insecticide resistance of these cuticular proteins, chitinases and a chitin synthase (AGAP001748) significantly overexpressed in the field *An. coluzzii* from Nigeria and Chad.

Evidence of insecticide resistance- and thermotolerance-associated genes sitting within chromosomal inversions

In this study, the findings of high frequency of 2La, 2Rb and 2Rc inversion polymorphisms in the resistant field populations of *An. coluzzii* across the Sahel, compared with the Ngoussou, suggested strong phenotypic adaptations in this species, across the Sahel. Most importantly, in addition to several of the cuticular protein genes associated with resistance (chitinases, chitin synthase, CPR, CPLC and CPAP proteins), several other genes previously implicated in thermotolerance and/or desiccation resistance in *An. gambiae/coluzzii*, and which were highly overexpressed in this study sit within these inversions. For example, the heat shock proteins, *hsp83* (AGAP006958) and *hsp90* hptG (AGAP006961), both of which are known to be heat and insecticides stress inducible (Zhao, 2012) and were among the core set of hsp genes involved in a common and immediate response to thermal stress in *An. gambiae* populations (Cassone et al., 2011) sit within the 2La inversion. These two genes were among the overexpressed genes in both heat-hardened and permethrin-resistant *An. coluzzii* populations from northern Nigeria (Ibrahim et al., 2021). Several ionotropic glutamate receptors were found within the 2La inversion breakpoints: *IR136* (AGAP006440), *IR139* (AGAP006691) and *IR142* (AGAP006407). This is not surprising as ionotropic receptors are commonly associated with chemosensation, thermosensation, and hygrosensation (VanHook, 2016, Enjin et al., 2016), characteristics which can confer advantages in xeric environments. The *IR25a* (AGAP010272) and *IR21a* (AGAP008511) are known to mediate both humidity and temperature preference in the fruit fly, *D. melanogaster* (Enjin et al., 2016, Greppi et al., 2020),

930 in addition to *IR21a* driving heat seeking and heat-stimulated blood feeding in *An.*
931 *gambiae* (Greppi et al., 2020), with both shown to be overexpressed/induced in
932 thermotolerant/permethrin resistant populations of *An. coluzzii* (Ibrahim et al., 2021).

934 Conclusions

Information on molecular basis of resistance and/or resistance genes and its markers facilitates
evidence-based control measures. In this study we characterised a major malaria vector, *An.*
coluzzii from the Sahel region of four countries, with findings which could promote evidence-
based, cross-border policy towards local and regional malaria control. The study found that
across Sahel (where malaria is highly seasonal, reaching its peak in the rainy season), the major
malaria vector is *An. coluzzii*. And that a handful of common cross-resistance genes are
responsible for multiple insecticide resistance in this species. Findings from this study suggest
pleotropic role of some key genes – able to confer insecticide resistance and/or stabilize the
insecticide resistance gene, at the same time conferring environmental adaptations, such as
the ability to survive thermal stress (thermotolerance). From operational vector control
perspective this study provided evidence of the role of key insecticide metabolism gene,
CYP6Z2 in bioactivation of a major insecticide, clothianidin, into more insecticidal
intermediates, which could increase the efficacy of the ingredients in malaria control tools
utilizing it to target field populations overexpressing this key P450.

950 Data Availablity

951 The raw sequence reads from this study have been deposited in the European Nucleotide
952 Archive, with accession PRJEB51644, and secondary accession of ERP136291.

954 Conflict of Interest

The authors declare no conflict of interest.

Acknowledgments

A Wellcome Trust International Training Fellowship in Public Health and Tropical Medicine

(WT201918/Z/16/Z) to SSI supported this work.

Reference

ADOLFI, A., POULTON, B., ANTHOUSI, A., MACILWEE, S., RANSON, H. & LYCETT, G. J. 2019. Functional
genetic validation of key genes conferring insecticide resistance in the major African malaria
vector, Anopheles gambiae. *Proc Natl Acad Sci U S A,* 116**,** 25764-25772.
ALEXA A., R. J. 2021. *topGO: Enrichment Analysis for Gene Ontology. R package version 2.46.0.*
[Online]. [Accessed].
ALISTAIR MILES, M. F. R., PETER RALPH, NICK HARDING, RAHUL PISUPATI, AND SUMMER RAE. 2020.
cggh/scikit-allel: A Python package for exploring and analysing genetic variation data
[Online]. [Accessed].
ANOPHELES GAMBIAE GENOMES, C., DATA ANALYSIS, G., PARTNER WORKING, G., SAMPLE, C.-A.,
BURKINA, F., CAMEROON, GABON, GUINEA, GUINEA, B., KENYA, UGANDA, CROSSES,
SEQUENCING, DATA, P., WEB APPLICATION, D. & PROJECT, C. 2017. Genetic diversity of the
African malaria vector Anopheles gambiae. *Nature,* 552**,** 96-100.
ANTONIO-NKONDJIO, C., POUPARDIN, R., TENE, B. F., KOPYA, E., COSTANTINI, C., AWONO-AMBENE,
P. & WONDJI, C. S. 2016. Investigation of mechanisms of bendiocarb resistance in Anopheles
gambiae populations from the city of Yaounde, Cameroon. *Malar J,* 15**,** 424.
ARONESTY, E. 2011. ea-utils: Command-line tools for processing biological sequencing data. Durham,
NC.
BALABANIDOU, V., KAMPOURAKI, A., MACLEAN, M., BLOMQUIST, G. J., TITTIGER, C., JUAREZ, M. P.,
MIJAILOVSKY, S. J., CHALEPAKIS, G., ANTHOUSI, A., LYND, A., ANTOINE, S., HEMINGWAY, J.,
RANSON, H., LYCETT, G. J. & VONTAS, J. 2016. Cytochrome P450 associated with insecticide
resistance catalyzes cuticular hydrocarbon production in Anopheles gambiae. *Proc Natl Acad*
Sci U S A, 113**,** 9268-73.
BARNES, K. G., WEEDALL, G. D., NDULA, M., IRVING, H., MZIHALOWA, T., HEMINGWAY, J. & WONDJI,
C. S. 2017. Genomic Footprints of Selective Sweeps from Metabolic Resistance to
Pyrethroids in African Malaria Vectors Are Driven by Scale up of Insecticide-Based Vector
Control. *PLoS Genet,* 13**,** e1006539.
BHATT, S., WEISS, D., CAMERON, E., BISANZIO, D., MAPPIN, B., DALRYMPLE, U., BATTLE, K., MOYES,
C., HENRY, A. & ECKHOFF, P. 2015. The effect of malaria control on Plasmodium falciparum
in Africa between 2000 and 2015. *Nature,* 526**,** 207.
BLIGHE, K., S. RANA, AND M. LEWIS. 2018. "EnhancedVolcano: Publication-ready volcano plots with
enhanced colouring and labeling.".
BRUCE-CHWATT, L. J. 1951. Malaria in Nigeria. *Bulletin of the World Health Organization,* 4**,** 301-327.
CAIRNS, M., ROCA-FELTRER, A., GARSKE, T., WILSON, A. L., DIALLO, D., MILLIGAN, P. J., GHANI, A. C.
& GREENWOOD, B. M. 2012. Estimating the potential public health impact of seasonal
malaria chemoprevention in African children. *Nat Commun,* 3**,** 881.
CARTHARIUS, K., FRECH, K., GROTE, K., KLOCKE, B., HALTMEIER, M., KLINGENHOFF, A., FRISCH, M.,
BAYERLEIN, M. & WERNER, T. 2005. MatInspector and beyond: promoter analysis based on
transcription factor binding sites. *Bioinformatics,* 21**,** 2933-42.

CASSONE, B. J., MOLLOY, M. J., CHENG, C., TAN, J. C., HAHN, M. W. & BESANSKY, N. J. 2011.
Divergent transcriptional response to thermal stress by Anopheles gambiae larvae carrying
alternative arrangements of inversion 2La. *Mol Ecol,* 20**,** 2567-80.
CHANDOR-PROUST, A., BIBBY, J., REGENT-KLOECKNER, M., ROUX, J., GUITTARD-CRILAT, E.,
POUPARDIN, R., RIAZ, M. A., PAINE, M., DAUPHIN-VILLEMANT, C., REYNAUD, S. & DAVID, J.
P. 2013. The central role of mosquito cytochrome P450 CYP6Zs in insecticide detoxification
revealed by functional expression and structural modelling. *Biochem J,* 455**,** 75-85.
CHIU, T. L., WEN, Z., RUPASINGHE, S. G. & SCHULER, M. A. 2008. Comparative molecular modeling of
Anopheles gambiae CYP6Z1, a mosquito P450 capable of metabolizing DDT. *Proc Natl Acad*
Sci U S A, 105**,** 8855-60.
CLARKSON, C. S., MILES, A., HARDING, N. J., O'REILLY, A. O., WEETMAN, D., KWIATKOWSKI, D.,
DONNELLY, M. J. & ANOPHELES GAMBIAE GENOMES, C. 2021. The genetic architecture of
target-site resistance to pyrethroid insecticides in the African malaria vectors Anopheles
gambiae and Anopheles coluzzii. *Mol Ecol,* 30**,** 5303-5317.
COETZEE, M., CRAIG, M. AND LE SUEUR, D. 2000. Distribution of African malaria mosquitoes
belonging to the Anopheles gambiae complex. *Parasitology today,* 16**,** 74-77.
COLUZZI, M., SABATINI, A., PETRARCA, V. & DI DECO, M. A. 1979. Chromosomal differentiation and
adaptation to human environments in the Anopheles gambiae complex. *Trans R Soc Trop*
Med Hyg, 73**,** 483-97.
DAO, A., YARO, A. S., DIALLO, M., TIMBINE, S., HUESTIS, D. L., KASSOGUE, Y., TRAORE, A. I., SANOGO,
Z. L., SAMAKE, D. & LEHMANN, T. 2014. Signatures of aestivation and migration in Sahelian
malaria mosquito populations. *Nature,* 516**,** 387-90.
DUNSE, K. M., KAAS, Q., GUARINO, R.F., BARTON, P.A., CRAIK, D.J. AND ANDERSON, M.A. 2010.
Molecular basis for the resistance of an insect chymotrypsin to a potato type II proteinase
inhibitor. *Proceedings of the National Academy of Sciences,* 107**,** 15016-15021.
ENJIN, A., ZAHARIEVA, E. E., FRANK, D. D., MANSOURIAN, S., SUH, G. S., GALLIO, M. & STENSMYR, M.
C. 2016. Humidity Sensing in Drosophila. *Curr Biol,* 26**,** 1352-8.
FADEL, A. N., IBRAHIM, S. S., TCHOUAKUI, M., TERENCE, E., WONDJI, M. J., TCHOUPO, M., WANJI, S.
& WONDJI, C. S. 2019. A combination of metabolic resistance and high frequency of the
1014F kdr mutation is driving pyrethroid resistance in Anopheles coluzzii population from
Guinea savanna of Cameroon. *Parasit Vectors,* 12**,** 263.
GIMONNEAU, G., BOUYER, J., MORAND, S., BESANSKY, N. J., DIABATE, A. & SIMARD, F. 2010. A
behavioral mechanism underlying ecological divergence in the malaria mosquito Anopheles
gambiae. *Behav Ecol,* 21**,** 1087-1092.
GRAU-BOVE, X., LUCAS, E., PIPINI, D., RIPPON, E., VAN 'T HOF, A. E., CONSTANT, E., DADZIE, S.,
EGYIR-YAWSON, A., ESSANDOH, J., CHABI, J., DJOGBENOU, L., HARDING, N. J., MILES, A.,
KWIATKOWSKI, D., DONNELLY, M. J., WEETMAN, D. & ANOPHELES GAMBIAE GENOMES, C.
2021. Resistance to pirimiphos-methyl in West African Anopheles is spreading via
duplication and introgression of the Ace1 locus. *PLoS Genet,* 17**,** e1009253.
GREPPI, C., LAURSEN, W. J., BUDELLI, G., CHANG, E. C., DANIELS, A. M., VAN GIESEN, L., SMIDLER, A.
L., CATTERUCCIA, F. & GARRITY, P. A. 2020. Mosquito heat seeking is driven by an ancestral
cooling receptor. *Science,* 367**,** 681-684.
GU, Z., EILS, R. & SCHLESNER, M. 2016. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics,* 32**,** 2847-9.
HALL, T. A. BioEdit: a user-friendly biological sequence alignment editor and analysis program for
Windows 95/98/NT. Nucleic acids symposium series, 1999. 95-98.
HU, H. X., ZHOU, D., MA, L., SHEN, B., SUN, Y. & ZHU, C. L. 2020. Lipase is associated with
deltamethrin resistance in Culex pipiens pallens. *Parasitol Res,* 119**,** 23-30.
HUDSON, R. R., SLATKIN, M. & MADDISON, W. P. 1992. Estimation of levels of gene flow from DNA
sequence data. *Genetics,* 132**,** 583-9.

HUESTIS, D. L., ARTIS, M. L., ARMBRUSTER, P. A. & LEHMANN, T. 2017. Photoperiodic responses of
Sahelian malaria mosquitoes Anopheles coluzzii and An. arabiensis. *Parasit Vectors,* 10**,** 621.
IBRAHIM, S. S., FADEL, A. N., TCHOUAKUI, M., TERENCE, E., WONDJI, M. J., TCHOUPO, M., KERAH-
HINZOUMBE, C., WANJI, S. & WONDJI, C. S. 2019a. High insecticide resistance in the major
malaria vector Anopheles coluzzii in Chad Republic. *Infect Dis Poverty,* 8**,** 100.
IBRAHIM, S. S., MUKHTAR, M. M., DATTI, J. A., IRVING, H., KUSIMO, M. O., TCHAPGA, W., LAWAL, N.,
SAMBO, F. I. & WONDJI, C. S. 2019b. Temporal escalation of Pyrethroid Resistance in the
major malaria vector Anopheles coluzzii from Sahelo-Sudanian Region of northern Nigeria.
Scientific reports, 9**,** 7395.
IBRAHIM, S. S., MUKHTAR, M. M., IRVING, H., LABBO, R., KUSIMO, M. O., MAHAMADOU, I. &
WONDJI, C. S. 2019c. High Plasmodium infection and multiple insecticide resistance in a
major malaria vector Anopheles coluzzii from Sahel of Niger Republic. *Malaria journal,* 18**,**
181.
IBRAHIM, S. S., MUKHTAR, M. M., MUHAMMAD, A. & WONDJI, C. S. 2021. 2La Paracentric
Chromosomal Inversion and Overexpressed Metabolic Genes Enhance Thermotolerance and
Pyrethroid Resistance in the Major Malaria Vector Anopheles gambiae. *Biology (Basel),* 10.
IBRAHIM, S. S., RIVERON, J. M., BIBBY, J., IRVING, H., YUNTA, C., PAINE, M. J. I. & WONDJI, C. S. 2015.
Allelic Variation of Cytochrome P450s Drives Resistance to Bednet Insecticides in a Major
Malaria Vector. *PLoS Genet,* 11**,** e1005618.
JOSHI, N. & FASS, J. 2011. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ
files (Version 1.33)[Software].
KEFI, M., CHARAMIS, J., BALABANIDOU, V., IOANNIDIS, P., RANSON, H., INGHAM, V. A. & VONTAS, J.
2021. Transcriptomic analysis of resistance and short-term induction response to
pyrethroids, in Anopheles coluzzii legs. *BMC Genomics,* 22**,** 891.
KUMAR, S., STECHER, G., LI, M., KNYAZ, C. & TAMURA, K. 2018. MEGA X: Molecular Evolutionary
Genetics Analysis across Computing Platforms. *Mol Biol Evol,* 35**,** 1547-1549.
LEES, R. S., ISMAIL, H. M., LOGAN, R. A. E., MALONE, D., DAVIES, R., ANTHOUSI, A., ADOLFI, A.,
LYCETT, G. J. & PAINE, M. J. I. 2020. New insecticide screening platforms indicate that
Mitochondrial Complex I inhibitors are susceptible to cross-resistance by mosquito P450s
that metabolise pyrethroids. *Sci Rep,* 10**,** 16232.
LEHMANN, T., WEETMAN, D., HUESTIS, D. L., YARO, A. S., KASSOGUE, Y., DIALLO, M., DONNELLY, M.
J. & DAO, A. 2017. Tracing the origin of the early wet-season Anopheles coluzzii in the Sahel.
Evol Appl, 10**,** 704-717.
LINDSAY, S., PARSON, L. AND THOMAS, C. 1998. Mapping the range and relative abundance of the
two principal African malaria vectors, Anopheles gambiae sensu stricto and An. arabiensis,
using climate data. *Proceedings of the Royal Society of London B: Biological Science,* 265**,**
847–854.
LOVE, M. I., HUBER, W., & ANDERS, S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology,* 15.
LOVE, R. R., REDMOND, S. N., POMBI, M., CAPUTO, B., PETRARCA, V., DELLA TORRE, A., ANOPHELES
GAMBIAE GENOMES, C. & BESANSKY, N. J. 2019. In Silico Karyotyping of Chromosomally
Polymorphic Malaria Mosquitoes in the Anopheles gambiae Complex. *G3 (Bethesda),* 9**,**
3249-3262.
LUCAS, E. R., ROCKETT, K. A., LYND, A., ESSANDOH, J., GRISALES, N., KEMEI, B., NJOROGE, H.,
HUBBART, C., RIPPON, E. J., MORGAN, J., VAN'T HOF, A. E., OCHOMO, E. O., KWIATKOWSKI,
D. P., WEETMAN, D. & DONNELLY, M. J. 2019. A high throughput multi-locus insecticide
resistance marker panel for tracking resistance emergence and spread in Anopheles
gambiae. *Sci Rep,* 9**,** 13335.
LYND, A. & LYCETT, G. J. 2011. Optimization of the Gal4-UAS system in an Anopheles gambiae cell
line. *Insect Mol Biol,* 20**,** 599-608.

MARTIN, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet. journal, 17**,** 10-12.
MENZE, B. D., KOUAMO, M. F., WONDJI, M. J., TCHAPGA, W., TCHOUPO, M., KUSIMO, M. O.,
MOUHAMADOU, C. S., RIVERON, J. M. & WONDJI, C. S. 2020. An Experimental Hut
Evaluation of PBO-Based and Pyrethroid-Only Nets against the Malaria Vector Anopheles
funestus Reveals a Loss of Bed Nets Efficacy Associated with GSTe2 Metabolic Resistance.
Genes (Basel), 11.
MESSENGER, L. A., IMPOINVIL, L. M., DERILUS, D., YEWHALAW, D., IRISH, S. & LENHART, A. 2021. A
whole transcriptomic approach provides novel insights into the molecular basis of
organophosphate and pyrethroid resistance in Anopheles arabiensis from Ethiopia. *Insect*
Biochem Mol Biol, 139**,** 103655.
MITCHELL, S. N., RIGDEN, D. J., DOWD, A. J., LU, F., WILDING, C. S., WEETMAN, D., DADZIE, S.,
JENKINS, A. M., REGNA, K., BOKO, P., DJOGBENOU, L., MUSKAVITCH, M. A., RANSON, H.,
PAINE, M. J., MAYANS, O. & DONNELLY, M. J. 2014. Metabolic and target-site mechanisms
combine to confer strong DDT resistance in Anopheles gambiae. *PLoS One,* 9**,** e92662.
MITCHELL, S. N., STEVENSON, B. J., MULLER, P., WILDING, C. S., EGYIR-YAWSON, A., FIELD, S. G.,
HEMINGWAY, J., PAINE, M. J., RANSON, H. & DONNELLY, M. J. 2012a. Identification and
validation of a gene causing cross-resistance between insecticide classes in Anopheles
gambiae from Ghana. *Proc Natl Acad Sci U S A,* 109**,** 6147-52.
MITCHELL, S. N., STEVENSON, B. J., MÜLLER, P., WILDING, C. S., EGYIR-YAWSON, A., FIELD, S. G.,
HEMINGWAY, J., PAINE, M. J., RANSON, H. & DONNELLY, M. J. 2012b. Identification and
validation of a gene causing cross-resistance between insecticide classes in Anopheles
gambiae from Ghana. *Proceedings of the National Academy of Sciences,* 109**,** 6147-6152.
MUGENZI, L. M. J., MENZE, B. D., TCHOUAKUI, M., WONDJI, M. J., IRVING, H., TCHOUPO, M., HEARN,
J., WEEDALL, G. D., RIVERON, J. M. & WONDJI, C. S. 2019. Cis-regulatory CYP6P9b
P450 variants associated with loss of insecticide-treated bed net efficacy against Anopheles
funestus. *Nat Commun,* 10**,** 4652.
MULLER, P., WARR, E., STEVENSON, B. J., PIGNATELLI, P. M., MORGAN, J. C., STEVEN, A., YAWSON, A.
E., MITCHELL, S. N., RANSON, H., HEMINGWAY, J., PAINE, M. J. & DONNELLY, M. J. 2008.
Field-caught permethrin-resistant Anopheles gambiae overexpress CYP6P3, a P450 that
metabolises pyrethroids. *PLoS Genet,* 4**,** e1000286.
RANSON, H., ABDALLAH, H., BADOLO, A., GUELBEOGO, W. M., KERAH-HINZOUMBE, C., YANGALBE-
KALNONE, E., SAGNON, N., SIMARD, F. & COETZEE, M. 2009. Insecticide resistance in
Anopheles gambiae: data from the first year of a multi-country study highlight the extent of
the problem. *Malar J,* 8**,** 299.
RIVERON, J. M., HUIJBEN, S., TCHAPGA, W., TCHOUAKUI, M., WONDJI, M. J., TCHOUPO, M., IRVING,
H., CUAMBA, N., MAQUINA, M., PAAIJMANS, K. & WONDJI, C. S. 2019. Escalation of
Pyrethroid Resistance in the Malaria Vector Anopheles funestus Induces a Loss of Efficacy of
Piperonyl Butoxide-Based Insecticide-Treated Nets in Mozambique. *J Infect Dis,* 220**,** 467-
475.
RIVERON, J. M., IBRAHIM, S. S., CHANDA, E., MZILAHOWA, T., CUAMBA, N., IRVING, H., BARNES, K.
G., NDULA, M. & WONDJI, C. S. 2014a. The highly polymorphic CYP6M7 cytochrome P450
gene partners with the directionally selected CYP6P9a and CYP6P9b genes to expand the
pyrethroid resistance front in the malaria vector Anopheles funestus in Africa. *BMC*
Genomics, 15**,** 817.
RIVERON, J. M., IBRAHIM, S. S., MULAMBA, C., DJOUAKA, R., IRVING, H., WONDJI, M. J., ISHAK, I. H.
& WONDJI, C. S. 2017. Genome-Wide Transcription and Functional Analyses Reveal
Heterogeneous Molecular Mechanisms Driving Pyrethroids Resistance in the Major Malaria
Vector Anopheles funestus Across Africa. *G3 (Bethesda),* 7**,** 1819-1832.

RIVERON, J. M., TCHOUAKUI M., MUGENZI L., MENZE B. D., MU-CHUN C. AND WONDJI C. S. 2018.
Insecticide Resistance in Malaria Vectors: An Update at a Global Scale. *In:* S. MANGUIN, V. D.
(ed.) *Towards Malaria Elimination - A Leap Forward.* London: IntechOpen.
RIVERON, J. M., YUNTA, C., IBRAHIM, S. S., DJOUAKA, R., IRVING, H., MENZE, B. D., ISMAIL, H. M.,
HEMINGWAY, J., RANSON, H., ALBERT, A. & WONDJI, C. S. 2014b. A single mutation in the
GSTe2 gene allows tracking of metabolically based insecticide resistance in a major malaria
vector. *Genome Biol,* 15**,** R27.
ROZAS, J., FERRER-MATA, A., SANCHEZ-DELBARRIO, J. C., GUIRAO-RICO, S., LIBRADO, P., RAMOS-
ONSINS, S. E. & SANCHEZ-GRACIA, A. 2017. DnaSP 6: DNA Sequence Polymorphism Analysis
of Large Data Sets. *Mol Biol Evol,* 34**,** 3299-3302.
SCHMITTGEN, T. D. & LIVAK, K. J. 2008. Analyzing real-time PCR data by the comparative C(T)
method. *Nat Protoc,* 3**,** 1101-8.
SHIFF, C. 2002. Integrated approach to malaria control. *Clin Microbiol Rev,* 15**,** 278-93.
STEVENSON, B. J., BIBBY, J., PIGNATELLI, P., MUANGNOICHAROEN, S., O'NEILL, P. M., LIAN, L. Y.,
MULLER, P., NIKOU, D., STEVEN, A., HEMINGWAY, J., SUTCLIFFE, M. J. & PAINE, M. J. 2011.
Cytochrome P450 6M2 from the malaria vector Anopheles gambiae metabolizes
pyrethroids: Sequential metabolism of deltamethrin revealed. *Insect Biochem Mol Biol,* 41**,**
492-502.
SUPEK, F., BOSNJAK, M., SKUNCA, N. & SMUC, T. 2011. REVIGO summarizes and visualizes long lists
of gene ontology terms. *PLoS One,* 6**,** e21800.
TCHOUAKUI, M., RIVERON, J. M., DJONABAYE, D., TCHAPGA, W., IRVING, H., SOH TAKAM, P.,
NJIOKOU, F. & WONDJI, C. S. 2018. Fitness Costs of the Glutathione S-Transferase Epsilon 2
(L119F-GSTe2) Mediated Metabolic Resistance to Insecticides in the Major African Malaria
Vector Anopheles Funestus. *Genes (Basel),* 9.
TOE, K. H., N'FALE, S., DABIRE, R. K., RANSON, H. & JONES, C. M. 2015. The recent escalation in
strength of pyrethroid resistance in Anopheles coluzzi in West Africa is linked to increased
expression of multiple gene families. *BMC Genomics,* 16**,** 146.
VANHOOK, A. M. 2016. Understanding hygrosensation: How flies sense changes in humidity. *Science
Signaling,* 9**,** ec127-ec127.
VANNINI, L., AUGUSTINE DUNN, W., REED, T. W. & WILLIS, J. H. 2014. Changes in transcript
abundance for cuticular proteins and other genes three hours after a blood meal in
Anopheles gambiae. *Insect Biochem Mol Biol,* 44**,** 33-43.
VONTAS, J., GRIGORAKI, L., MORGAN, J., TSAKIRELI, D., FUSEINI, G., SEGURA, L., NIEMCZURA DE
CARVALHO, J., NGUEMA, R., WEETMAN, D., SLOTMAN, M. A. & HEMINGWAY, J. 2018. Rapid
selection of a pyrethroid metabolic enzyme CYP9K1 by operational malaria control activities.
Proc Natl Acad Sci U S A, 115**,** 4619-4624.
WEEDALL, G. D., MUGENZI, L. M. J., MENZE, B. D., TCHOUAKUI, M., IBRAHIM, S. S., AMVONGO-
ADJIA, N., IRVING, H., WONDJI, M. J., TCHOUPO, M., DJOUAKA, R., RIVERON, J. M. &
WONDJI, C. S. 2019. A cytochrome P450 allele confers pyrethroid resistance on a major
African malaria vector, reducing insecticide-treated bednet efficacy. *Sci Transl Med,* 11.
WEEDALL, G. D., RIVERON, J. M., HEARN, J., IRVING, H., KAMDEM, C., FOUET, C., WHITE, B. J. &
WONDJI, C. S. 2020. An Africa-wide genomic evolution of insecticide resistance in the
malaria vector Anopheles funestus involves selective sweeps, copy number variations, gene
conversion and transposons. *PLoS Genet,* 16**,** e1008822.
WEETMAN, D., WILDING, C. S., NEAFSEY, D. E., MULLER, P., OCHOMO, E., ISAACS, A. T., STEEN, K.,
RIPPON, E. J., MORGAN, J. C., MAWEJJE, H. D., RIGDEN, D. J., OKEDI, L. M. & DONNELLY, M. J.
2018. Candidate-gene based GWAS identifies reproducible DNA markers for metabolic
pyrethroid resistance from standing genetic variation in East African Anopheles gambiae. *Sci
Rep,* 8**,** 2920.

WEILL, M., MALCOLM, C., CHANDRE, F., MOGENSEN, K., BERTHOMIEU, A., MARQUINE, M. &
RAYMOND, M. 2004. The unique mutation in ace-1 giving high insecticide resistance is easily
detectable in mosquito vectors. *Insect Mol Biol,* 13**,** 1-7.
WHO 2015. Global Technical Strategy for Malaria 2016-2030. World Health Organisation. . Geneva,
Switzerland.
WHO 2018a. Update on the E-2020 initiative of 21 malaria-eliminating countries: report and country
briefs. Geneva. Switzerland: World Health Organization.
WHO 2018b. World Malaria Report. World Health Organization. . Geneva Switzerland.
WHO 2019. World Malaria Report 2019. Geneva, Switzerland.
WHO 2021. World Malaria Report. *20 years of global progress and challenges.* Geneva: Switzerland.
WILLIAMS, J., COWLISHAW, R., SANOU, A., RANSON, H. & GRIGORAKI, L. 2022. In vivo functional
validation of the V402L voltage gated sodium channel mutation in the malaria vector An.
gambiae. *Pest Manag Sci,* 78**,** 1155-1163.
YI, X., LIANG, Y., HUERTA-SANCHEZ, E., JIN, X., CUO, Z. X., POOL, J. E., XU, X., JIANG, H.,
VINCKENBOSCH, N., KORNELIUSSEN, T. S., ZHENG, H., LIU, T., HE, W., LI, K., LUO, R., NIE, X.,
WU, H., ZHAO, M., CAO, H., ZOU, J., SHAN, Y., LI, S., YANG, Q., ASAN, NI, P., TIAN, G., XU, J.,
LIU, X., JIANG, T., WU, R., ZHOU, G., TANG, M., QIN, J., WANG, T., FENG, S., LI, G., HUASANG,
LUOSANG, J., WANG, W., CHEN, F., WANG, Y., ZHENG, X., LI, Z., BIANBA, Z., YANG, G., WANG,
X., TANG, S., GAO, G., CHEN, Y., LUO, Z., GUSANG, L., CAO, Z., ZHANG, Q., OUYANG, W., REN,
X., LIANG, H., ZHENG, H., HUANG, Y., LI, J., BOLUND, L., KRISTIANSEN, K., LI, Y., ZHANG, Y.,
ZHANG, X., LI, R., LI, S., YANG, H., NIELSEN, R., WANG, J. & WANG, J. 2010. Sequencing of 50
human exomes reveals adaptation to high altitude. *Science,* 329**,** 75-8.
YUNTA, C., GRISALES, N., NASZ, S., HEMMINGS, K., PIGNATELLI, P., VOICE, M., RANSON, H. & PAINE,
M. J. 2016. Pyriproxyfen is metabolized by P450s associated with pyrethroid resistance in An.
gambiae. *Insect Biochem Mol Biol,* 78**,** 50-57.
YUNTA, C., HEMMINGS, K., STEVENSON, B., KOEKEMOER, L. L., MATAMBO, T., PIGNATELLI, P., VOICE,
M., NASZ, S. & PAINE, M. J. I. 2019. Cross-resistance profiles of malaria mosquito P450s
associated with pyrethroid resistance against WHO insecticides. *Pestic Biochem Physiol,* 161**,**
61-67.
ZHAO, L. A. J., W.A. 2012. Expression of heat shock protein genes in insect stress responses. .
Invertebrate Survival Journal, 9**,** 93-101.

1230

1231 Figure Legends

1232 **Figure 1: A volcano plot of differentially expressed genes showing fold changes and**
1233 **significance in R-S, C-S and R-C comparisons, for Nigeria and Niger populations of** *An.*
1234 *coluzzii.*

1235

1236 **Figure 2: A volcano plot of differentially expressed genes showing fold changes and**
1237 **significance in R-S, C-S and R-C comparisons, for Chad and Cameroon populations of** *An.*
1238 *coluzzii.*

1239

1240 **Figure 3: Validation of candidate resistance genes.** qRT-PCR of twelve metabolic resistance
1241 genes (R-S and C-S comparisons) from Sahel *An. coluzzii.*

1242

1243 **Figure 4: Identification of resistance variants.** A heatmap showing frequencies of the
1244 resistance variants (haplotypes) in key genes of interest.

1245

1246 **Figure 5: Characterization of 5′ regulatory element of** *GSTe2.* Results of dual-luciferase
1247 reporter assays of the promoter (intergenic region) constructs, showing progressive loss of
1248 activity following sequential deletion of the constructs.

1249

1250 **Figure 6: Validation of the role of metabolic resistance genes in insecticide resistance.**
1251 Results of insecticide susceptibility bioassays with transgenic flies expressing *GSTe2* and
1252 *CYP6Z2.* **a.** α-cypermethrin and deltamethrin, b. permethrin, **c.** 3-phenoxybenzaldehyde and
1253 3-phenoxybenzyl alcohol, **d.** DDT and clothianidin.

1254