Molecular drivers of insecticide resistance in the Sahelo-Sudanian populations of a major malaria vector *Anopheles coluzzii*

Sulaiman S. Ibrahim[1,2,3,*], Abdullahi Muhammad[1,4], Jack Hearn[1], Gareth D. Weedall[5], Sanjay C. Nagi[1], Muhammad M. Mukhtar[2], Amen N. Fadel[3], Leon J. Mugenzi[3], Edward I. Patterson[6], Helen Irving[1] and Charles S. Wondji[1,3]

[1] Vector Biology Department, Liverpool School of Tropical Medicine (LSTM), L3 5QA, UK

[2] Department of Biochemistry, Bayero University, PMB 3011, Kano, Nigeria

[3] LSTM Research Unit, Centre for Research in Infectious Diseases (CRID), P.O. Box 13591, Yaoundé, Cameroon

[4] Centre for Biotechnology Research, Bayero University, PMB 3011, Kano, Nigeria

[5] School of Biological and Environmental Sciences, Liverpool John Moores University, Liverpool L3 3AF, UK

[6] Department of Biological Sciences, Brock University, St. Catharines, Ontario, L2S 3A1, Canada.

* Correspondence: suleiman.ibrahim@lstmed.ac.uk ; Tel.: +44-74-4044-3871; ORCID: 0000-0003-0744-802X

Authors email addresses

Sulaiman S. Ibrahim: suleiman.ibrahim@lstmed.ac.uk

Abdullahi Muhammad: abdullahi.muhammad@lstmed.ac.uk

Jack Hearn: jack.hearn@lstmed.ac.uk

Gareth D. Weedall: g.d.weedall@ljmu.ac.uk

Sanjay C. Nagi: sanjay.nagi@lstmed.ac.uk

Muhammad M. Mukhtar: muhammadmahemukhtar@gmail.com

Amen N. Fadel: amen.fadel@crid-cam.net

Leon M. J. Mugenzi: leon.mugenzi@crid-cam.net

Edward I. Patterson: ipatterson@brocku.ca

Helen Irving: helen.irving@lstmed.ac.uk

Charles S. Wondji: charles.wondji@lstmed.ac.uk

Abstract

Information on common markers of metabolic resistance in malaria vectors from countries

sharing similar eco-climatic characteristics can facilitate coordination of malaria control.

Here, we characterized populations of the major malaria vector *Anopheles coluzzii* from Sahel

region, spanning four sub-Saharan African countries: Nigeria, Niger, Chad and Cameroon.

Genome-wide transcriptional analysis identified major genes previously implicated in

pyrethroid and/or cross resistance to other insecticides, overexpressed across the Sahel,

including CYP450s, glutathione S-transferases, carboxylesterases, and cuticular proteins.

Several, well-known markers of insecticide resistance were found in high frequencies -

including in the voltage-gated sodium channel (V402L, I940T, L995F, I1527T and N1570Y),

the *acetylcholinesterase*-1 gene (G280S) and the *CYP4J5*-L43F (fixed). High frequencies of the

epidemiologically important chromosomal inversions, 2La, 2Rb and 2Rc were observed

(~80% for 2Rb and 2Rc). The 2La alternative arrangement is fixed across the Sahel. Low

frequencies of these inversions (<10%) were observed in the fully insecticide susceptible

laboratory colony of *An. coluzzii* (Ngoussou). Several of the most commonly overexpressed

metabolic resistance genes sit in these three inversions. Two commonly overexpressed

genes, *GSTe2* and *CYP6Z2* were functionally validated. Transgenic *Drosophila melanogaster*

expressing *GSTe2* exhibited extremely high DDT and permethrin resistance (mortalities <

10% in 24 h). Serial deletion of the 5' intergenic region, to identify putative nucleotide(s)

associated with *GSTe2* overexpression, revealed that simultaneous insertion of adenine

nucleotide and a transition (T->C), between Fork-head box L1 and c-EST putative binding

sites were responsible for the high overexpression of *GSTe2* in the resistant mosquitoes.

Transgenic flies expressing *CYP6Z2* exhibited marginal resistance towards 3-

phenoxybenzylalcohol (a primary product of pyrethroid hydrolysis by carboxylesterases) and a type II pyrethroid, α-cypermethrin. However, significantly higher mortalities were observed in *CYP6Z2* transgenic flies compared with controls, on exposure to the neonicotinoid, clothianidin. This suggests a possible bioactivation of clothianidin into a toxic intermediate, which if true make it an ideal insecticide against populations of *An. coluzzii* overexpressing this P450. These findings will facilitate regional collaborations within the Sahel region, and refine implementation strategies through re-focusing interventions, improving evidence-based, cross-border policy towards local and regional malaria pre-elimination.

Keywords: *Anopheles coluzzii*, Sahel, insecticides, pyrethroid, DDT, metabolic, resistance, genes, inversions.

Background

Since the year 2000 the massive scale-up in vector control interventions and treatment with antimalarial drugs had cut malaria incidence by ~40 % across Africa (1). Bolstered by this progress the World Health Organization (WHO) has been pushing to eliminate malaria, as proposed in the Global Technical Strategy (GTS) 2016-2030 (2). Unfortunately, the GTS, an ambitious framework with targets to reduce global malaria burden by 90 % in 15 years was dealt an immediate blow by a rebound in malaria transmission, with increased cases between 2016 and 2019 (3, 4). This stark warning of the risk posed to control and elimination efforts was a reflection of the lack of progress in the primary regions of interest in sub-Saharan Africa, which constitute 96 % of the 627,000 malaria-related deaths in 2020 alone (4, 5). Indeed, as the WHO widens the net of malaria elimination (E-2025) its important to acknowledge that no meaningful progress will be made without progress in sub-Saharan Africa, where none of the six countries having the highest global burden of malaria (e.g. Nigeria alone contributing ~27 % of all cases) is on the path to elimination (6).

The recent escalation of insecticide resistance in the major malaria vectors (7-9) makes the development of molecular tools to anticipate emergence and predict spread of resistance across the African continent imperative, in order to achieve malaria control and elimination. For decades no simple molecular assays were available to track metabolic resistance and assess its impact on malaria control and transmission, until the recent discovery of two DNA markers in the *cis*-regulatory region of two cytochrome P450s, *CYP6P9a* (10) and *CYP6P9b* (11) in the major malaria vector *Anopheles funestus*; findings which allowed the design of simple PCR assays to detect and track metabolic resistance in the field. Unfortunately, these markers explain resistance only in southern Africa for *An. funestus*, since genetic basis of

pyrethroid resistance, and cross-resistance with other insecticides is complex in other

African regions with different fronts, driven by distinct genes in the presence of barriers to

gene flow (10, 12, 13). No major metabolic resistance markers for P450s or GSTs (functionally

validated and characterised for epidemiological impact in the natural populations) exist for

the major malaria vectors of the *Anopheles gambiae* Complex, though it is the omnipresent

vector species with widespread presence across Africa (14). This is hindering early detection

and tracking of molecular drivers of resistance in this species, slowing down evidence-based

control measures and resistance management.

Major genomic regions associated with metabolic resistance to pyrethroids in *An. gambiae*

sensu lato include the *CYP6* P450 clusters on the 2R and 3R chromosomes, and a *GST*

epsilon (*GSTe*) cluster on chromosome 3R (15), with some key resistance-associated genes

functionally validated. These genes include *CYP6P3* shown to confer cross resistance to

pyrethroids and organophosphates (16, 17), *CYP6M2* conferring cross resistance to

pyrethroids (18, 19) and DDT (19), as well as *GSTe2* shown to confer resistance to DDT (20).

For *GSTe2*, in addition to the I114T marker (20), a recent study has found a novel mutation

(*Gste2*-119V) associated with resistance (21) using a high-throughput genotypic panel for

markers. Functional validation and fild data is required establish the empirical evidence of

the role of this mutation in resistance. What has been missing in the case of *Anopheles*

gambiae and *An. coluzzii* is reliable molecular markers of resistance for major metabolic gene

families, e.g. P450s (with field and laboratory validated data) to aid creation of DNA-based

diagnostic assays which will allow (i) easy tracking of resistance in the field (e.g. the case of

119F-*GSTe2* mutation in *An. funestus* (22), and (ii) determination of the operational impact of

the resistance markers in the field, as recently done for 119F-*GSTe2* (23), *CYP6P9a*_R and

*CYP6P9b*_R markers (10, 11) in *An. funestus*.

The *An. gambiae s.l.*, especially populations in the semi-arid steppe, exhibit high frequency of

paracentric chromosomal inversions [one of the most effective instruments for speciation

and local adaptations (Ayala et al., 2014, Dobzhansky, 1971, Kirkpatrick, 2010)], maintained

in spatially and temporally heterogenous environment, and which segregate along climatic

gradients of increasing aridity (24)]. The 2La inversions are associated with resistance to

desiccation in adults (Fouet et al., 2012, Gray et al., 2009) and thermal stress in larvae

(Cassone et al., 2011). It was also shown that inversion 2La assorts with insecticide

resistance, e.g., dieldrin plus fipronil (Brooke et al., 2000), and is associated with

thermotolerance and permethrin resistance in the Sahelain *An. coluzzii* (Ibrahim et al., 2021).

To support malaria pre-elimination effort in sub-Saharan Africa we targeted the Sahelo-

Sudanian region, which represent northern-most limit of malaria endemicity in sub-Saharan

Africa, and where malaria is highly seasonal [offering excellent target for pre-elimination

effort through sustained seasonal vector control and seasonal malaria chemoprevention

(25)]. Focusing on the predominant malaria vector, *An. coluzzii* from Sudano-Sahelian

transects of four countries, Nigeria, Niger, Chad and Cameroon (7, 26-28) we identified the

major metabolic resistance genes mediating pyrethroid resistance and cross-resistance in this

region, established the genetic variants which explained the resistance. We also functionally

validated the roles of two major candidate genes in the resistance (*CYP6Z2* and *GSTe2*) using

transgenic *Drosophila melanogaster* flies (GAL4/UAS system), as well as identifying single

nucleotide polymorphisms in the 5' regulatory elements of *GSTe2*, responsible for its

overexpression in the resistant population.

Materials and Methods

Study Site and Mosquito Sampling

Blood fed female *An. coluzzii* mosquitoes, resting indoor were collected at one locality each

in sub-Sahel (Additional Figure S1): Hadiyau (HAD: 12°21′38″N, 9°59′15″E), a sub-Sahel

village in northern Nigeria; Takatsaba (TAK: 13°44'01.8"N 7°59'05.2"E), a Sahel village in

southern Niger; Simatou (SIMAT: 10°50'40.7"N 14°56'40.9"E), a sub-Sahel village in Maga

Department, far north of Cameroon; and Massakory (CHAD: 12° 6′ N, 15° 02′ E), a Sahel

town in Chad Republic. Details of sampling approaches and resistance profiles of

mosquitoes collected from Nigeria, Niger, and Chad are available in previously published

articles (7, 26, 28). As in the above countries the Simatou F_1 females were also highly

pyrethroid resistant, with a mortality of only 3.7% from WHO tube bioassays using 0.05%

deltamethrin, and no mortality at all with 0.75% permethrin (data not published).

Genome-wide transcriptional analysis for common insecticide resistance genes in

the Sahel regions

RNA extraction, library preparation and sequencing

The RNA was extracted using the Arcturus PicoPure RNA isolation Kit (Applied

Biosystems, CA, USA) from three pools of 8 F_1 *An. coluzzii* females (2-4 day old from the

same population) alive after exposure to deltamethrin (resistant, R), unexposed (control, C),

and also from unexposed females of the fully susceptible laboratory colony of *An. coluzzii*,

Ngoussou (susceptible, S) (29). The RNA isolation was carried out following the

manufacturer's protocol with *Dnase* I-treatment to remove contaminating DNA. The

quantity and quality of RNA was measured using a NanoDrop spectrophotometer

(ThermoFisher, MA, USA) and Bioanalyzer (Agilent, CA, USA).

Library preparation, sequencing and data quality control were carried out by the Centre for

Genomic Research (CGR), University of Liverpool, UK. RNA samples were subjected to

poly(A) mRNA enrichment and libraries prepared from the poly(A) mRNA-enriched

materials (dual-indexed, strand-specific RNAseq libraries were prepared using the NEBNext

polyA selection and Ultra Directional RNA library preparation kits). Libraries were

sequenced on a single lane of an Illumina HiSeq 4000 (paired-end, 2x150 bp sequencing,

generating data from >280 M clusters per lane). Basecalling and de-multiplexing of indexed

reads were performed by CASAVA version 1.8.2 (Illumina). De-multiplexed fastq files were

trimmed to remove Illumina adapter sequences using Cutadapt version 1.2.1 (30). Option -O

3 was used, so that the 3' end of any reads which matched the adapter sequence for 3 bp or

more were trimmed. Reads were further trimmed to remove low quality bases

using Sickle version 1.200 (31) with a minimum window quality score of 20. Reads shorter

than 20 bp after trimming were removed. If both reads from a pair passed this filter, each

was included in either the R1 (forward reads) or R2 (reverse reads) file. If only one of a read

pair passed this filter, it is included in the R0 (unpaired) file. Statistics were generated using

fastq-stats from EAUtils (32). Summary of total number of reads for each sample and

distribution of trimmed read length for forward (R1) and reverse (R2) reads and reads

unpaired after trimming (R0) are provided in Additional Figure S2.

Data analysis and estimation of transcript abundance by tag counting and differential gene

expression

Paired data for each replicate per country was aligned to the *An. gambiae* reference transcript

ome AgamP4.10 downloaded from VectorBase (https://vectorbase.org/) in salmon (0.11.4),

using 'validate mappings', 'seqBias', 'gcBias' and 'rangeFactorizationBins 4' flags. Read

mapping results (pre-alignment and post-alignment descriptive statistics (flagstat output

files) showing sequencing depth and coverage are given in Additional Tables 1 and -2,

respectively. Salmon results were converted into a gene expression matrix using the

Bioconductor package 'tximport' for input to DESeq2 1.26.0 (33). Differential gene

expression was tested for the three possible combinations of Exposed (R, deltamethrin

resistant), Unexposed (C, control) and Susceptible replicates (S). For results interpretation

$\log_2$-fold change thresholds of 1 was imposed, with false discovery rate adjusted p-values of

0.05 applied to accept significance. Principal components analysis implemented in DESeq2

was used to examine relationships between respective replicates and treatments. This was

carried out based on the 500 most variable genes, with data transformation

(normalisation/scaling) as implemented in VST (DESeq2). For visualisation of expression

results, volcano plots were created using the Enhanced Volcano package (34) using the top

most overexpressed genes from lists which were prepared with $\log_2$FC cut off of 1 and p

value of 0.01. Heatmaps were generated (35) using the list of the top 50 most overexpressed

metabolic resistance genes with $\log_2$FC cut off of 1 and p value of 0.01.

Gene Ontology Enrichment and Mapping/Functional annotations

The enrichment analysis for GO terms was carried out using a topGO package (36) against

An. gambiae (AgamP4.10), with data reannotated using EggNOG v5.0

(http://eggnog5.embl.de/#/app/home). Gene lists used in topGO are from the $\log_2 FC = 0$, p

<0.05 analysis (the default in DESeq2). For the data from each site (country) six set of results

were generated (R, C and S contrasts) for genes either up- or down-regulated in each

contrast, for molecular function (MF) and biological process (BP) ontologies. For analysis the

GO terms lists for MF were used as input into the Revigo (37) for interpretation and

visualisation, querying the Whole Uniprot database, and SimRel setting for semantic

similarity measurements.

Quantitative PCR measurement of expression profiles of metabolic resistance

genes

The level of expression of 12 resistance-associated genes was validated by qRT-PCR, using

the primers provided in Additional Table 3. These include the *GSTe2* (AGAP009194), *GSTZ1*

(AGAP002898), *CYP6Z2* (AGAP008218), *CYP6Z3* (AGAP008217), *CYP4C27* (AGAP009246),

CYP4G16 (AGAP001076), *CYP4G17* (AGAP000877), *CYP6P3* (AGAP002865), *CYP6M2*

(AGAP008212), *CYP9K1* (AGAP000818), a *UGT-B19* (AGAP007920) and *COEBE3C*

(AGAP005372). The qRT-PCR was carried out using three technical replicates each of cDNA

extracted from 1 µg of total RNA of three biological replicates each from the Resistant (R),

Control (C) and Ngoussou (S). Protocol followed was as established in previous studies (38),

with relative expression level and fold change (FC) of each target gene in R and C relative to

S calculated according to the $2^{-\Delta\Delta CT}$ method incorporating the PCR efficiency (39), after

normalization with the housekeeping genes ribosomal protein S7, *RPS7* (AGAP010592) and

glycerol-3-phosphate dehydrogenase, *GPDH* (AGAP007593). Significant differences were

calculated using ANOVA with Dunnett's post hoc test.

Detection of Signatures of Selective Sweep

To detect signature of selective sweeps in the major metabolic resistance genes of interest, a

Snakemake RNAseq population genetics pipeline, the RNAseq-Pop was utilised (40). The

workflow aligns RNA-Seq reads to the reference genome, and calls genomic variants with

Freebayes, at a user-provided level of ploidy, in our case 16 (8 diploid pooled mosquitoes). F_{st}

(41) per gene between population pairs and Tajima's D per gene within population were

estimated. This was performed against all SNPs passing quality and missingness filters.

Population branch statistic (PBS) scans was performed with the Snakemake, conditional on

the presence of three suitable populations (42). Hudson's F_{st} and PBS scans were ran, taking

the average for each protein-coding gene, as opposed to in windows. The population

genetics statistical analyses were calculated in scikit-allel v1.2.1 (43).

Establishment of allele frequencies of variants in genes of interest

After genome alignment, RNA-Seq-Pop utilises samtools (44) to query specific positions of

the genome, calculating raw allele frequencies at those sites with a custom R script.

Detection of chromosomal inversion polymorphisms and metabolic genes within its

breakpoint

A modified version of the Python 3 program, compkaryo (45) was used to karyoptype the

major, *An. coluzzii/gambiae* phenotypically important inversion polymorphisms in

chromosome 2, and calculate its frequencies, *in silico*, using the previously identified tag

SNPs significantly associated with inversions. This allows to predict with high confidence

genotypes of the six common polymorphic inversions on chromosome 2, in the sequenced

field *An. coluzzii,* as well as in the Ngoussou. Compkaryo uses the Ag1000 database

(The *Anopheles gambiae* 1000 Genomes Consortium 2017) (15) by leveraging a subset of

cytologically karyotyped specimens to develop a computational approach for karyotyping

applicable to whole genome sequence. Modifications in the Snakemake pipeline allows for

variable ploidy (useful in the case of replicates from our pooled RNAsequencing samples)

here.

Functional validation of the commonly overexpressed resistance-associated genes

Comparative analysis of coding sequences of major resistance genes

Two resistance-associated genes, *GSTe2* and *CYP6Z2* which feature most premoninently in

the resistant *An. coluzzii* populations across the Sahel were chosen for functional validation.

To establish presence of allelic variants which could impact catalytic activities, full length

coding sequences (cDNAs) of *GSTe2* and *CYP6Z2* were amplified and sequenced from alive

mosquitoes in the four Sahel countries, as well as from the Ngoussou. This was done using

total RNA extracted from 5 individual pools of 8 F_1 *An. coluzzii* females (2-4 day old) alive

after exposure to deltamethrin (resistant, R for *CYP6Z2*) or DDT (R for *GSTe2*). Protocol for

RNA extraction was as described in previous section, above. Amplification was done using

Phusion HotStart II Taq Polymerase (ThermoFisher SCIENTIFIC, MA, USA) and the Full

primers listed in Additional Table 4. The PCR mix comprised 5x Phusion HF Buffer

(containing 1.5 mM $MgCl_2$), 85.7 μM deoxynucleotides (dNTPs), 0.34 μM each of forward

and reverse primers, 0.015 U of Phusion HotStart II DNA Polymerase (Fermentas, MA,

USA), 10.71 μL of ddH_2O and 1 μL cDNA. Thermocycling conditions were 1 cycle at 95 °C

for 5 min; followed by 35 cycles each of 94 °C for 20 s, 60 °C for 30 s, 72 °C for 2 min (1 min

for *GSTe2*); and finally, one cycle at 72 °C for 5 min. PCR products were cleaned with a

QIAquick® PCR Purification Kit (QIAGEN, Hilden, Germany) and ligated into the

pJET1.2/blunt cloning vector using the CloneJET PCR Cloning Kit (ThermoFisher

SCIENTIFIC, MA, USA). These were then cloned into *E. coli DH5α*, plasmids miniprepped

with the QIAprep® Spin Miniprep Kit (QIAGEN) and sequenced on both strands using

pJET1.2 primers.

Polymorphisms were detected through examination and manual editing of sequence traces

using BioEdit version 7.2.3.0 (46) and nucleotide differences in sequences aligned using CLC

Sequence Viewer 7.0 (http://www.clcbio.com/). Different haplotypes were compared by

constructing a maximum likelihood phylogenetic tree using MEGA X (47). Genetic

parameters of polymorphism including number of haplotypes (h) and its diversity (H_d),

number of polymorphic sites (S) and nucleotide diversity (π) were computed using DnaSP

v6.12.03 (48).

Characterization of the 5′ regulatory regions of GSTe2 and CYP6Z2

Amplification, cloning and sequence characterisation of 5′ regulatory element

To investigate presence of genetic variants in the regulatory elements, which could be

responsible for overexpression of *GSTe2*, 351 bp intergenic regions (spanning the 43 bp 3′

UTR of *GSTe1*, 248 bp flanking sequence and 60 bp 5′UTR of *GSTe2*) preceding the start

codon were amplified from 10 each of DDT-alive and -dead females from the 4 Sahel

countries, as well as from the Ngoussou females (primers provided in Additional Table 4).

For *CYP6Z2*, a 1078 bp intergenic region was retrieved from the VectorBase and used for

amplification of the putative 5′-regulatory elements. Primers spanning 38 bp 3′ UTR of

CYP6Z1, a 937 bp flanking sequence and 103 bp 5′UTR of *CYP6Z2*, preceding the start codon

of *CYP6Z2* were used to amplify fragments from 10 each of deltamethrin-alive and -dead

females from Nigeria and Niger, as well as from the Ngoussou females. Amplification was

carried out using HotStart II Polymerase (ThermoFisher SCIENTIFIC, MA, USA) with

similar thermocycling conditions as outlined above for coding region of *GSTe2* and *CYP6Z2*,

respectively. Purification of PCR amplicons, cloning into pJET1.2 vector, sequencing and

polymorphism analysis were done as outlined above.

The 351 bp 5′-UTR fragments of *GSTe2* and 1078 bp fragment of the *CYP6Z2* were analysed

with the Gene Promoter Miner (http://gpminer.mbc.nctu.edu.tw/) and MatInspector (49) to

identify putative promoter elements, predict transcription start (TSS) and potential

transcription factor binding sites.

Cloning of GSTe2 and CYP6Z2 5' regulatory elements in PGL3-Basic vector and dual luciferase

reporter assay

Following analysis of the above sequences the 351 bp intergenic fragments of *GSTe2* were

amplified from the most predominant sequences of DDT-alive, DDT-dead and Ngoussou.

Same was done for for *CYP6Z2* amplifying 1087 bp fragment from deltamethrin-alive and -

dead, and Ngoussou. Primers bearing *kpn*I and *BgL*II sites (Additional Table 4) allowed

incorporation into pGL3-Basic reporter vector containing luciferase gene from the firefly

Photinus pyralis (Promega, Wisconsin, USA). Amplification was carried out using Phusion

HotStart II Polymerase, with conditions as above, followed by purification of PCR

amplicons, and cloning into pJET1.2 vector. Positive colonies (sequencing primers for pGL3-

Basic provided in Additional Table 4) were miniprepped; the minipreps digested with the

above restriction enzymes, gel-purified and ligated upstream of luciferase gene in pGL3-

Basic vector already linearized with the same restriction enzymes. Positive colonies were

cloned and miniprepped and concentrations of the recombinant plasmids adjusted to 200

ng/μL before co-transfection. Protocols for maintenance of the *An. gambiae* cell line 4a-3B

(MRA-919, https://www.beiresources.org/), co-transfection of 200 ng of either *GSTe2* or

CYP6Z2 promoter constructs, LRIM promoter in pGL3-Basic vector (50), or promoter-less

pGL3-Basic control, with 1 ng/μL of internal control, sea pansy *Renilla reniformis* luciferase

containing the *Drosophila* Actin *5C* promoter in pRL-null (50), are provided in Additional

File 1 (additional text for Methods and Results). Protocol for cell lysis and measurement of

the activities of the firefly luciferase (Dual-Luciferase Reporter Assay kit, Promega), with

normalisation using the Renilla luciferase activity, were as outlined in the Promega Quick

Protocol, and provided in detail in Additional File 1.

Assay background was also measured using lysate from non-transfected control cells.

Results were compared using a two-tailed Chi-Square test of independence using GraphPad

Prism 7.02 (GraphPad Inc., La Jolla, CA, USA).

Generation of GSTe2 promoter deletion constructs and assays

The intergenic region separating *GSTe1* from *GSTe2* was progressively delineated. Partial

fragments of the 5′ regions (encompassing the 3′-UTR of *GSTe1*, the *GSTe1*/*GSTe2* flanking

region and the 5′-UTR of the *GSTe2*) were created by sequential deletion using forward

primers (provided in Additional Table 4, with numbers in primer names referring to

distances from the AUG start codon of *GSTe2*). Reverse primers were those initially used for

amplification of the full 351 bp nucleotide fragments. These fragments include (i) 308

nucleotides fragment generated from deletion of the 43 bp 3′-UTR of the *GSTe1* (-308

primers) obliterating cellular-Myb (c-Myb) transcriptional factor binding site; (ii) 291

nucleotide fragments generated from deletion of *GSTe1* 3′-UTR plus 38 nucleotides of the

flanking region (-270 primers), which removed in addition δ elongation factor 1 (δEF1)

binding site; (iii) 262 nucleotide fragments produced from deletion of *GSTe1* 3′-UTR plus 46

nucleotides of the flanking region (-262 primers), which obliterated in addition Forkhead

box L1 (FOX-L1) putative binding site; as well as (iv) 232 nucleotide fragments produced

from deletion of *GSTe1* 3′-UTR plus 75 nucleotides of the flanking region (-232 primers),

located 11 nucleotides from the c-EST/grainy head transcription factor binding site. Also, to

assess the importance of the *GSTe2* 5′UTR, 43 nucleotides upstream the AUG codon were

deleted (from the 60 nucleotide 5′-UTR of both the Sahel-alive and Ngoussou), leaving the

putative transcription start site motif untouched. This was done using the original forward

primer for the 351 nucleotides intergenic region, with a newly designed primer,

GSTe2_minus_5′-UTR-R (Additional Table 4).

Characterisation of major resistance genes using transgenic analysis

Cloning and microinjection of GSTe2 and CYP6Z2 into Drosophila melanogaster

Transgenic flies expressing recombinant GSTe2 and CYP6Z2 were created using GAL4/UAS

system, as outlined in a previous publication (51) and used in contact bioassays, to confirm if

over-expression of these genes alone can confer resistance to insecticides.   Details of

amplification of full-length *GSTe2* and *CYP6Z2* from the predominant alleles of sequences of

cDNA, cloning into pUASattB vector, microinjection into germ-line *D. melanogaster* lines,

crossing of the UAS-lines with the GAL4-Actin driver strain are outlined in Additional File

1.

To confirm overexpression of *GSTe2* and *CYP6Z2* in the transgenic flies three replicates each

of 6 females (both experimental and control group) were used for qRT-PCR using a

previously established protocol (38). Total RNA and cDNA were extracted as described

above, and the relative expression levels of transgenes were assessed, with normalization

using the *RPL11* housekeeping gene. The qtrg primers used for the two genes and the *RPL11*

primers are provided in Additional Table 4.

Insecticide Susceptibility Contact Bioassay

For insecticide bioassays, 3- to 4-day old experimental and control F_1 females were exposed

to 0.15 % deltamethrin, 2 % permethrin, 0.05 % α-cypermethrin, 4 % DDT and 2 %

clothianidin-impregnated papers prepared in acetone and Dow Corning 556 Silicone Fluid

(BHD/Merck, Hesse, Germany). Flies overexpressing *CYP6Z2* were also exposed to the

primary product of hydrolysis of pyrethroids: 4 % and 20 % (5x) 3-phenoxybenzaldehyde

(PBAld) and 3-phenoxybenzylalcohol (PBAlc). Transgenic flies expressing *GSTe2* were

exposed to 2 % permethrin, 0.15 % deltamethrin, 0.05 % α-cypermethrin and 4 % DDT only.

Impregnated papers were rolled and introduced into 45 cc plastic vials to cover the entire

wall and the vials plugged with cotton soaked in 10 % sucrose (38). 20–25 flies were placed

in each vial, and the mortality plus knockdown scored at 1 h, 3 h, 6 h, 12 h and 24 h of

exposure to the insecticides. For each insecticide, assays were performed in 6 replicates and

Student's t-test used to compare the mortality plus knockdown between the experimental

groups and the control. Controls were adult F_1 progeny with the same genetic background

as the experimental group but without the *GSTe2* or *CYP6Z2* inserts. They were obtained by

crossing virgin females from the driver strain Act5C-GAL4 and the UAS recipient male lines

with white eyes (not carrying the pUASattB-GSTe2 or pUASattB-CYP6Z2 insertions).

Results

Genome-wide transcriptional profile of the Sahelian *An. coluzzii* populations

A three-way pairwise comparison was conducted for the data from each country: resistant

vs susceptible (R-S), resistant *vs* unexposed control (R-C) and unexposed control *vs*

susceptible (C-S). This captures background variations due to geographical differences in the

resistant vs susceptible (R-S) comparison, accounts for genes overexpressed due to induction

(R-C comparison), as well as genes that are constitutively overexpressed (C-S comparison).

A total of 1384 genes were significantly differentially expressed (FDR-adjusted p <0.05 and

log₂ fold change threshold of 1/FC ≥ 2) in R-S comparison in Nigeria (1077 upregulated and

307 down-regulated); 1185 genes were differentially expressed in C-S (1002 upregulated and

183 downregulated); and 295 genes in R-C (129 upregulated and 166 downregulated). Of

these, 52 genes were commonly differentially expressed in all 3 comparisons (Figure S3,

panel a), including the upregulated genes, *COEAE80* (AGAP006700), *CYP4H18*

(AGAP028019), *CYP4H17* (AGAP008358) and cuticular proteins, *CPLCX3* (AGAP006149),

CPR59 (AGAP006829), *CPR76* (AGAP009874) and *CPR75* (AGAP009871). The Additional

Figure S3, panels a-d depicts the differentially expressed genes for the four countries. For

Niger, 881 genes were differentially expressed in R-S comparison (619 upregulated and 262

down-regulated) (Additional Figure S3b); 1256 genes were differentially expressed in C-S

(986 upregulated and 270 downregulated); and 196 genes in R-C (81 upregulated and 115

downregulated). Of these, 22 genes were commonly differentially expressed in all 3

comparisons [including the upregulated *aminopeptidase N1* (AGAP012757), *CYP6Z2*

(AGAP008218), *chymotrypsin-3* (AGAP006711) and *chymotrypsin-2* (AGAP006710), and an

acid trehalase (AGAP008547)]. For Chad, 1392 genes were differentially expressed in R-S

comparison (975 upregulated and 417 down-regulated) (Additonal Figure S3c); 1284 in C-S

(105 upregulated and 269 downregulated); and 526 genes in R-C (270 upregulated and 256

downregulated). Of these, 97 genes were commonly differentially expressed in all 3

comparisons [including *CYP4C27* (AGAP009246), *SULTD1* (AGAP012672), *aminopeptidase*

N1, and diverse cuticular proteins, e.g. chitinase (*Cht24*, AGAP006191, *CPFL1*

(AGAP010902), *CPCFC1* (AGAP007980), *CPLCX3*, *CPR24* (AGAP005999), *CPR106*

(AGAP006095) and *CPR130* (AGAP000047). Finally, for Cameroon, 376 genes were

differentially expressed in R-S comparison (204 upregulated and 172 down-regulated)

(Additional Figure S3d); 932 in C-S (778 upregulated and 154 downregulated); and 116

genes in R-C [only 9 upregulated and 107 downregulated (probably due to a single, low

quality replicate in the raw data from Cameroon (Additional Figure S2). Not surprising,

only 7 genes were commonly differentially expressed in all 3 comparisons. These include the

highly upregulated gene, *GSTe2* (AGAP009194) and *chymotrypsin-1* (AGAP006709).

All data analysed together (Additional Figure S3e) revealed no single gene differentially

expressed in common; possibly due to the low quality with the Chad unexposed (C) data.

Analysis of data from Nigeria, Niger and Chad, revealed a single gene (AGAP000046,

transporter major facilitator superfamily) differentially expressed across all countries

(Additional Figure S3f). Niger and Chad shared *hexamerin* (AGAP010658), *aminopeptidase N1*,

and unknown protein, AGAP0290967; Nigeria and Niger share only a single gene,

AGAP003248; while seven genes were common to Nigeria and Chad, including *CPLCX3*.

Principal component analysis for the top 500 most variable genes in all experimental arms

revealed data from field samples (R and C) from all four countries clustering closer in PC1

and PC2 axes, away from the data from the susceptible females, Ngoussou (Additional

Figure S4).

Analysis of the common differentially expressed genes across the Sahel

The most differentially expressed genes were presented in Figure 1, a volcano plot of fold

change *vs* significance levels for Nigeria and Niger, and Figure 2, for Chad and Cameroon.

Detailed lists of these genes of interest is provided in Additional File 2. Comparisons of

genes commonly, upregulated and/or downregulated in R-S/R-C/C-S, from the four

countries revealed similar transcriptomic profiles between R and C compared with the S.

The most commonly and consistently overexpressed genes across the Sahel (taking account

mean expressions) are the chymotrypsins, 3, -2 and -1 (*CHYM3*/AGAP006711 and

CHYM2/AGAP006710 and *CHYM1*/AGAP006709) (Figures 1 and 2, Additional File 2), the

glutathione S-transferase, *GSTe2* (AGAP009194), an aquaporin, *AQP3* (AGAP010326),

CYP6Z2 (AGAP008218), *CYP6Z3* (AGAP008217), *CYP4C27* (AGAP009246), a chitinase, *Cht24*

(AGAP006191), a thioester-containing protein-1, *TEP-1* (AGAP010815),), a trehalose 6-

phosphate synthase/phosphatase, *TPS*1/2 (AGAP008227), a lipase (AGAP002353), and

AGAP012818 (V-type protein ATPase subunit A). More on these genes is provided in the

sections, below.

Other genes commonly overexpressed in data from two or three countries include

AGAP005501 (dehydrogenase/reductase SDR family 11) upregulated in Nigeria and Niger

(R-S and C-S comparisons), and in R-S for Chad and Cameroon; AGAP008091 (CLIP-domain

serine protease, *CLIPE1*), upregulated in all countries in R-S and C-S comparisons; a

chitinase, *Cht5-5*, AGAP013260) upregulated in Nigeria, Nigeria and Chad R-S and C-S, with

both induced upregulated in R-C in Chad, as heme peroxidase (*HPX15*), upregulated in R-S

and C-S comparisons in Nigeria, Chad and Cameroon, as well as malate dehydrogenase

(AGAP000184), upregulated in Nigeria, Chad and Cameroon, R-S and C-S.

However, the most overexpressed genes in Nigeria are a cubilin, a histone (H2B), carbonic

anhydrase I, and a cuticular protein, *CPR131* (Additional File 2), while for Cameroon its

H2B, Serpin 9 inhibitory serine protease inhibitor, galectin 4 and a Protease m1 zinc

metalloprotease.

Several genes were significantly downregulated across the four countries, particularly the

cytochrome P450s and the ATP-binding cassette transporters. The most consistently down-

regulated P450s were *CYP6AK1* (AGAP010961), downregulated in R-S and C-S comparisons

in Nigeria, Niger and Cameroon, while down-regulated in all comparisons in Chad;

CYP4H24 (AGAP013490), downregulated in R-S and C-S comparisons for Nigeria and

Cameroon, and downregulated in all comparisons in Niger and Chad; *CYP6M4*

(AGAP008214), downregulated in R-S and C-S comparisons in Nigeria, Niger and

Cameroon, while downregulated in all comparisons in Chad; and finally *CYP6P5*

(AGAP002866), downregulated in R-S and C-S comparisons in Niger and Chad, but only in

C-S in Nigeria.

Analysis of commonly overexpressed metabolic resistance genes across the Sahel

Special attention was given to the known metabolic resistance genes, implicated in

insecticide resistance in *Anopheles* and/or other insects, including, CYP450s, GSTs,

carboxylesterases, cuticular proteins, chemosensory proteins/SAPs, uridine

diphosphoglucuronosyltransferases, etc, in addition to other important genes, e.g., the

immune proteins. Analysis of the data from the list of genes significantly, differentially

expressed (FDR-adjusted $p < 0.05$ and $\log_2 FC = 1/ FC \geq 2$) revealed the major, commonly

upregulated metabolic resistance genes across the Sahel. Top 50 genes from these lists

(Additional File 3) from each country are displayed as heatmaps of fold changes in

Additional Figure S5a, -b, -c and -d). The most common genes linked with resistance and/or

other physiologically important phenotypes are tabulated in Table 1, showing fold changes

of 55 genes (R-S and C-S comparisons), common to all countries, or three or two countries at

least. The most commonly and consistently overexpressed genes across the Sahel (taking

account mean expressions) are the chymotrypsins, -1, -2 and -3 (*CHYM1*/AGAP006709,

CHYM3/AGAP006711 and *CHYM2*/AGAP006710) (Table 1, Additional Figure S5, Additional

File 2, Results section), with fold change for *CHYM1* in R-S and C-S comparisons of 31.62

and 54.41 for Nigeria, 12.31 and 41.59 for Niger, and 18.00 and 18.08 for Chad, and 5.69 and

11.77 for Cameroon; for *CHYM3*, 32.45 and 58.43 for Nigeria R-S and C-S, respectively, 7.30

and 43.27 for Niger, 13.40 and 7.24 for Chad, and 2.59 and 8.15 for Cameroon. Of *Anopheles*

metabolic resistance genes, *GSTe2* (AGAP009194) was the most consistently upregulated

gene across the Sahel, FC = 9.61 and 7.16, respectively for Nigeria R-S and C-S comparisons;

6.55 and 11.32 for Niger; 9.09 and 10.02 for Chad, 9.68 and 20.67 for Cameroon. Other

common GSTs were *GSTZ1*, upregulated in three countries (Nigeria, Niger and Chad),

GSTe4, *GSTU1*, *GSTD1-4* and *GSTD3*.

Other most consistently and highly upregulated genes include a lipase, AGAP002353: FC of

26.73 and 20.80 for Nigeria R-S and C-S comparisons, 16.53 and 29.76 for Niger, 14.95 and

9.95 for Chad, 14.58 and 14.53 for Cameroon; a chitinase, *Cht24* (AGAP006191), FC = 21.12

and 38.55 for Nigeria R-S and C-S, 40.08 and 67.18 for Niger, 91.42 and 5.71 for Chad, 44.26

and 40.53 for Cameroon); a V-type protein ATPase subunit a (AGAP012818, FC = 22.12 and

14.61 for Nigeria, 31.05 and 25.43 for Nigeri, 6.00 and 4.31 for Chad, 44.92 and 43.4 for

Cameroon), and an aquaporin, *AQP3* (AGAP010326, FC = 14.81 and 11.03 for Nigeria, 12.09

and 21.94 for Niger, 14.86 and 1.21 for Chad, 14.17 and 17.3 for Cameroon).

The most overrepresented gene family, and most consistently upregulated across the Sahel

were the cytochrome P450s, with *CYP6Z2*, *CYP6Z3*, and *CYP4C27*, in addition to the two

P450s linked to cuticular resistance, *CYP4G16* (upregulated in Nigeria and Chad) and

CYP4G17 (upregulated in Nigeria, Niger and Chad). The three P450s, *CYP6Z2, CYP6Z3 and*

CYP4C27 are consistently overexpressed: FC for *CYP6Z2* is 6.55 and 2.10 for Nigeria R-S

and C-S comparisons, 6.73 and 2.32 for Niger, 13.96 and 1.48 for Chad, 11.84 and 8.39

for Cameroon; FC for *CYP6Z3* is 4.73 and 1.82 for Nigeria, 6.09 and 1.42 for Niger, 3.51 and

1.11 for Chad, 7.11 and 3.79 for Cameroon; and FC for *CYP4C27* is 7.36 and 4.6 for Nigeria,

4.03 and 5.64 for Niger, 6.04 and 2.62 for Chad, 2.01 and 2.63 for Cameroon. The two well-

known insecticide resistance genes *CYP6P3* and *CYP6M2* were not

overexpressedprominently across the Sahel of these countries.

The second most commonly overrepresented gene families belong to cuticular proteins,

including cuticular proteins Rebers and Riddiford (RR), several cuticular proteins of low

complexity (CPLC) and chitin-binding cuticular proteins (CPAP), with the commonly

upregulated ones across all four countries being *CPR75* (AGAP009871) and *CPR76*

(AGAP009874).

Two carboxylesterases picture prominently across the Sahel – the beta esterase, *COEBE3C*

(AGAP005372) upregulated in all four countries, while *COEJHE5E* was common to Nigeria,

Niger and Chad. For Phase II metabolism enzymes, several uridine-diphospho-

glucuronosyltransferases, UGTs were upregulated, including the *UGT-B19*, which is

upregulated in Nigeria and Niger (Table 1), and unknown UGT (AGAP006222) upregulated

across all the countries (Additional File 2).

Other genes in the top 55 metabolic genes include the malate dehydrogenase, *TPS* (*TPS1/2*),

AQP2, thioester-containing proteins, *TEP1*, *-12* and *-14*, a cyanogenic-beta-glucosidase

(AGAP006422) and three chemosensory proteins, *CSP5*, sensory appendage proteins, -1 and

-2.

552 **Table 1: The common detoxification and metabolic genes differentially upregulated in**
553 **Sahel *An. coluzzii* (log$_2$FC values and FDR-adjusted p < 0.05).**

Gene	NIGERIA		NIGER		CHAD		CAMEROON		Gene Description
	R-S	C-S	R-S	C-S	R-S	C-S	R-S	C-S	
AGAP006400	2.2	4.2	1.7	4.5	-	-	2.1	2.8	alkaline phosphatase 2
AGAP028491	4.3	2.8	1.4	2.7	2.9	2.6	-	-	aquaporin, AQP2
AGAP010326	14.8	11.0	12.1	21.9	14.9	1.2	14.2	17.3	Aquaporin, AQP3
AGAP005372	9.5	12.2	3.3	4.3	6.6	3.6	1.4	2.6	carboxylesterase, COEBE3C
AGAP005837	4.7	5.0	1.4	3.3	1.9	2.6			carboxylesterase, COEJHE5E
AGAP004904	3.9	2.2	-	-	2.3	1.7	-	-	catalase, CAT1
AGAP029127	4.1	3.2	-	-	1.7	2.2	-	-	chemosensory protein 5, CSP5
AGAP006191	21.1	38.6	40.1	67.2	91.4	5.7	44.3	40.5	chitinase, Cht24
AGAP013260	4.4	11.5	1.3	3.9	3.1	0.8	-	-	chitinase, Cht5-5
AGAP006898	8.9	5.5	-	-	2.2	3.9	-	-	chitinase, Cht6
AGAP006709	31.6	54.4	12.3	41.6	18.0	18.1	5.7	11.8	chymotrypsin-1, CHYM1
AGAP006710	9.9	19.4	2.1	8.8	4.2	3.9	-	-	chymotrypsin-2, CHYM2
AGAP006711	32.5	58.4	7.3	43.3	13.4	7.2	2.6	8.2	chymotrypsin-3, CHYM3
AGAP000987	4.3	2.7	-	-	1.3	2.6	-	-	cuticular protein, CPAP3-A1b
AGAP000988	5.3	1.6	-	-	0.7	5.4	-	-	cuticular protein, CPAP3-A1c
AGAP000986	7.3	4.6	-	-	1.7	3.5	-	-	cuticular protein, CPAP3-D
AGAP006149	3.8	0.8	-	-	0.7	5.8	-	-	cuticular protein, CPLCX3
AGAP010123	33.3	54.9	2.4	13.2	-	-	-	-	cuticular protein, CPR131
AGAP005456	5.2	1.8	-	-	1.2	2.2	1.1	2.1	cuticular protein, CPR15
AGAP006009	-	-	2.9	1.3	-	-	3.9	1.6	cuticular protein, CPR30
AGAP009871	8.7	2.7	1.1	2.6	2.2	4.2	1.7	2.6	cuticular protein, CPR75
AGAP009874	10.6	2.2	1.1	2.6	3.7	5.6	1.1	2.0	cuticular protein, CPR76
AGAP009879	7.4	2.6	-	-	1.9	4.6	-	-	cuticular protein, CPR81
AGAP006422	7.5	9.4	-	-	8.4	7.3	2.1	2.0	cyanogenic beta-glucosidase
AGAP002417	5.5	4.2	0.9	1.2	1.9	1.9	-	-	cytochrome P450, CYP4AR1
AGAP009246	7.4	4.6	4.0	5.6	6.0	2.6	2.0	2.6	cytochrome P450, CYP4C27
AGAP012957	5.0	2.8	-	-	2.9	2.5	2.0	2.6	cytochrome P450, CYP4D17
AGAP001076	5.8	4.7	1.5	1.8	3.8	2.4	0.9	1.5	cytochrome P450, CYP4G16
AGAP000877	3.8	2.0	2.3	1.9	3.8	2.1	1.2	2.1	cytochrome P450, CYP4G17
AGAP008358	-	-	2.9	4.1	3.4	3.3	3.6	2.0	cytochrome P450, CYP4H17
AGAP007480	5.4	3.9	1.3	2.7	2.5	2.0	-	-	cytochrome P450, CYP6AH1
AGAP002865	1.11	0.5	2.3	1.9	0.7	12.8	2.1	2.3	cytochrome P450, CYP6P3
AGAP008207	3.6	1.9	-	-	2.1	1.6	1.7	2.4	cytochrome P450, CYP6Y2
AGAP008218	6.6	2.1	6.7	2.3	14.0	1.5	11.8	8.4	cytochrome P450, CYP6Z2
AGAP008217	4.7	1.8	6.1	1.4	3.5	1.1	7.1	3.8	cytochrome P450, CYP6Z3
AGAP000818	3.0	1.2	0.8	0.5	0.9	1.6	3.4	2.2	cytochrome P450, CYP9K1
AGAP010400	3.6	2.6	1.6	2.8	3.3	1.4	-	-	flavin-containing monooxygenase
AGAP007920	5.7	2.4	2.8	1.8	-	-	-	-	glucuronosyltransferases, UGT-B19
AGAP004164	2.2	1.9	1.7	2.3	1.4	1.3	-	-	glutathione S-transferase, GSTD1-4
AGAP009194	9.6	7.2	6.6	11.3	9.1	10.0	9.7	20.7	glutathione S-transferase, GSTe2
AGAP009193	3.3	2.3	1.1	2.1	1.5	1.6	-	-	glutathione S-transferase, GSTe4
AGAP000947	4.4	2.3	1.2	2.2	1.9	3.2	-	-	glutathione S-transferase, GSTU1
AGAP002898	5.5	3.5	1.7	2.6	2.8	2.4	-	-	glutathione S-transferase, GSTZ1
AGAP002198	-	-	2.0	5.8	4.8	6.6	-	-	glycine-N-methyltransferase
AGAP013327	-	-	1.2	7.3	7.9	20.1	5.1	14.8	heme peroxidase, HXP15
AGAP002353	26.7	20.8	16.5	29.8	15.0	10	14.6	14.5	lipase
AGAP000184	5.2	6.3	-	-	4.9	4.3	1.2	1.9	malate dehydrogenase
AGAP008051	3.7	4.0	1.1	1.3	1.7	1.7	-	-	sensory appendage protein 1, SAP1
AGAP008052	5.5	10.8	1.0	1.5	3.6	1.1	-	-	sensory appendage protein 2, SAP2
AGAP010815	5.7	14.4	4.1	8.6	7.8	3.8	1.7	3.1	thioester-containing protein, TEP1
AGAP008368	3.3	6.1	1.4	3.9	3.4	2.0	-	-	thioester-containing protein, TEP14
AGAP008654	4.6	17.4	2.1	10.0	-	-	-	-	thioester-containing protein, TEP12
AGAP008227	4.1	4.0	4.0	2.0	4.6	1.5	2.6	4.1	trehalose 6-phosphate synthase/phosphatase
AGAP012818	22.1	14.6	31.1	25.4	6.0	4.3	44.9	43.4	V-type proton ATPase, subunit a

Gene Ontology Enrichment Analysis

Gene ontology enrichment analysis, for genes significant in comparisons ($\log_2$FC = 1, p <

0.05) revealed crucial differences in overrepresented GO terms between the up- and -

downregulated genes, in data from the four countries. For example, the most over-

represented, semantically similar GO terms associated with xenobiotics metabolism is

oxidoreductase activity ((highest frequency of 12.9 %, Additional Figure S6a, REVIGO Table

View), which cluster together in Nigerian R vs S comparison (for genes upregulated in R).

Other over-represented GO terms in this comparison include glutathione S-transferase

activity (the most enriched/specific term), peroxidase activity, odorant binding and chitin

binding. In contrast, GO terms over-represented in R vs S (down-regulated in R) were

mostly involved in neurotransmission, metal ions binding and receptor channelling

activities (Additional Figure S6b).

For R vs C comparison (upregulated in C) the over-represented GO terms include

oxidoreductase (frequency = 12.88 %), glutathione S-transferase activity (the most

enriched/specific term), peroxidase activity, glucuronosyltransferase activity, aldehyde

oxidase activity, chitin binding, carbohydrate binding activities, etc, (Additional Figure S6c).

In contrast the most enriched GO terms, downregulated in C were endopeptidase activities,

Toll binding, oxidoreductase (frequency = 1.21 %) and monooxygenase activities (Additional

Figure S6d). For the rest of the three countries similar contrasts were also observed, between

phenotypes, and detailed in Additional File 1, Results section.

Quantitative PCR validation of expression profiles of metabolic resistance genes

The relative expression levels of 12 metabolic resistance genes were validated. The qRT-PCR results support the transcriptomic patterns obtained from the RNAseq analysis, with *GSTe2* as the most overexpressed gene, followed by *CYP6Z2* (Figure 3). For example, for R-S and C-S comparisons *GSTe2* had fold changes of 49.25 ± 5.05 and 28.45 ± 4.45, for Nigeria ($p \leq 0.0001$); 37.01 ± 2.22 and 33.34 ± 6.61 for Niger ($p \leq 0.0001$); 29.97 ± 5.39 and 30.16 ± 6.16 for Chad ($p \leq 0.0001$); and 28.30 ± 3.30 ($p \leq 0.0001$) and 17.03 ± 7.03 for Cameroon ($p \leq 0.001$). *CYP6Z2* has R-S and C-S fold changes of 15.93 ± 2.54 ($p \leq 0.001$) and 7.35 ± 1.52 ($p \leq 0.05$), for Nigeria; 24.21 ± 2.17 ($p \leq 0.001$) and 20.83 ± 3.02 for Niger ($p \leq 0.001$); 15.02 ($p \leq 0.001$) ± 2.01 and 6.09 ± 2.08 for Chad ($p \leq 0.05$); and 10.58 ± 3.00 ($p \leq 0.01$) and 3.43 ± 1.43 for Cameroon. Correlation analyses for R-S comparisons for all genes revealed positive and significant association (Additional Figure S7) between the RNA-seq and qRT-PCR data in data from Niger ($R^2 = 0.58$, p = 0.03) and Chad ($R^2 = 0.408$, p = 0.025), with positive, but non-significance seen in Cameron ($R^2 = 0.58$ p = 0.08) and Nigerian ($R^2 = 0.503$, p = 0.06).

Detection of Signatures of Selective Sweeps

Signatures of selection were investigated in the major metabolic resistance genes, by estimating Tajima's D per gene within populations, and *F*st per gene between population pairs. These tests of neutrality revealed several genes exhibiting genetic differentiation, or possibly undergoing expansion. Among the top 100 most overexpressed metabolic genes across the four countries (Additional File 2), 6 genes were possibly undergoing genetic differentiation. These include *TEP1,* and *TEP3*, with average Tajima's D of -1.4 and -1.6

respectively, in populations from Chad, and Fst values of ≥ 0.85 for *TEP1*, for Nigeria, Chad

and Cameroon samples, compared to the average Fst of 3L chromosomal, which was

calculated as 0.24 (Additional File 3), *CYP9K1* (average Tajima's D = -1.00 for Cameroon,

and Fst within ranges of 0.86 - 1 for Nigeria, Cameroon and Niger, compared with 0.21

calculated for the chromosome X), *CPR15* (average Tajima's D = -0.94); *CPAP3-A1b* (Tajima's

D = -0.70 in Chad, -0.95 in Nigeria, and -1.32 in Cameroon, with Fst values in range of 0.7 -

0.9 for the four countries, compared with the average Fst of 0.21 for chromosome X) and

GSTU1. Several genes from the GST family exhibited strong genetic differentiation, with

reduced variations prominently in *GSTe2* (< 3 SNPs in Chad and Cameroon samples), with

Tajima's D of -1.45 and -1.15 for Nigeria and Niger, and combined. Information on more

genes is provided in Additional File 1, and detailed information on all genes can be found

in the Additional File 3.

Identificaton of genetic variants associated with insecticide resistance

The RNA-Seq-Pop workflow calculated allele frequencies of variants of interest found in the

raw RNA-sequencing read data. Figure 4 displays allele frequencies of each variant in

respective treatment replicates used for the RNAseq, as well as Ngoussou. Several mutaions

were found within the voltage-gated sodium channel. In addition to the L995F *kdr*

mutation, the recently identified V402L and I1527T mutations exist in all the field

populations across the Sahel, at frequencies ranging from 29 % to 67 %. We also detected the

N1570Y mutation, which shares the same haplotype as L995F. The G280S *acetylcholinesterase-*

1 mutation was also observed in low to moderate frequencies in all populations except for

Ngoussou and samples from Niger. The L43F pyrethroid resistance marker of *CYP4J5* was

also seen, fixed in most populations.

623 Detection of chromosomal inversion polymorphisms and metabolic genes within

624 its breakpoints

The frequency of the major inversion polymorphisms in chromosome 2 were calculated,

considering ploidy (8 individual female mosquitoes were pooled for RNA extraction, for

each replicate). Additional Table 5 provided the frequencies of the respective inversions for

the populations from each country, as well as for Ngoussou. High frequency of the 2La, 2Rb

and 2Rc were observed, in contrast to the 2Rd, 2Rj and 2Ru which were in lower frequencies

[except for the 2Rj in Nigeria (30.38 %) and 2Ru in Cameroon (24.29 %)]. The 2La inversion

was found fixed in the field populations (100 % in Nigeria and Cameroon, 99.92 % in Niger

and 99.21 % in Chad), in contrast with Ngoussou, with frequency of only 6.26 %. Similar

pattern was observed with the 2Rb inversion, with high frequencies in three field

populations (79.01 %, 84.85 %, 80.22 % in Nigeria, Niger and Chad), but lower in Cameroon

(29.08 %) and Ngoussou (5.35 %). Frequencies of the 2Rc inversion were 88.54 %, 85.52 %,

88.82 %, 46.99 % and 9.89 % for Nigeria, Niger, Chad, Cameroon and Ngoussou respectively.

Some genes among the top 100 most overexpressed metabolic resistance genes (above) and

which were likely undergoing recent population expansion were located within the 2La, 2Rb

and 2Rc inversions (Additional File 4). These include the *hsp83* (AGAP006958) and

molecular chaperone hptG (AGAP006961), *CPLCX3* (AGAP006149), *CPLCA1* (AGAP006145,)

within the inverted 2La chromosomal arm. Other genes within 2La inversion include a

group of chitinase enzymes (*Cht6*, *-8*, *-11* and *-24*), most notably the *Cht24* and *Cht6*, which

are among the top 50 most overexpressed genes in all countries. Other genes include:

COEJHE5E, one of the 14 carboxylesterases in the 2La region, and which was among the top

100 most overexpressed genes across the Sahel; the most overexpressed chymotrypsin genes,

CHYM1 and *CHYM2*; three ionotropic receptors, *IR136* (AGAP006440), *IR139* (AGAP006691)

and *IR142* (AGAP006407); as well as five P450s, *CYP301A1*, *CYP302A1*, *CYP4J5*, *CYP4J9* and

CYP4J10. The most overrepresented genes within 2La inversions are the CPR cuticular

proteins (73 in total), with *CPR21*, *-26*, *-30*, *-59* and *-140* from the list of the top 100 resistance

genes from the Sahel.

Carbonic anhydrase I (the most overexpressed metabolic resistance in Nigeria,

AGAP013402) sits within 2Rb inversion. Four chitinases (*Cht4*, *Cht5-1*, *Cht5-3*, and *Cht5-5*)

are located within 2Rb inversion; *Cht5-1* and *Cht5-5* were among the observed top 50 most

overexpressed metabolic genes in the Sahel. Lipase (*AGAP002353*) which is the among the

top 6 overexpressed metabolic gene in the four countries sit within the 2Rb inversion. Eight

CYP450s reside in the 2Rb inversion, including *CYP4D15*, *CYP4D17* and *CYP4AR1*, which

were among the top 100 metabolic resistance genes. Three cuticular proteins, *CPR7*, *-8* and *-9*

were also within the 2Rb inversion.

Interesting genes sitting within 2Rc inversion include *GSTZ1*; 5 ionotropic receptor genes,

IR7i (AGAP013363), *IR7u* (AGAP013285), *IR7t* (AGAP002763), *IR7w* (AGAP013416) and

IR41a (AGAP002904); a carboxylesterase, *COEAE6O* (AGAP002863); and 11 CYP450s,

including *CYP6AA1* (AGAP002862), *CYP6AA2* (AGAP013128), *CYP6P15P* (AGAP002864),

CYP6P3 (AGAP002865), *CYP6P5* (AGAP002866), *CYP6P4* (AGAP002867), *CYP6P1*

(AGAP002868), *CYP6P2* (AGAP002869), *CYP6AD1* (AGAP002870) and *CYP6Z4*

(AGAP002894)

Investigation of the polymorphism in the coding sequences of *GSTe2* and *CYP6Z2*

Analysis of the polymorphism patterns of full-length cDNA sequences of *GSTe2* (666 bp)

and *CYP6Z2* (1479 bp) from the Sahel region of Africa revealed complete homogeneity for

GSTe2, with no polymorphism detected in the field populations (all sequences were identical

to those from Ngoussou and the AGAP009194 reference). This suggests a fixed allele,

consistent with the observation from the analyses from the fixation index (Additional File 3).

For *CYP6Z2* (1479 bp), homogeneity was observed within each country and Ngoussou,

except for Niger, characterised by an unusually high polymorphism (Additional Figure S8a,

-b). *CYP6Z2* is polymorphic with 10 haplotypes across the Sahel, with 75 polymorphic sites

of which 65 were synonymous, and 12 led to amino acids substitutions. The bulk of the

polymorphisms were contributed from larger variations in the Niger and Cameroon (S = 42

and 23 respectively), while highest homogeneity was observed in Chad, with a single

haplotype. Haplotype diversity is high (H_d = 0.921), from 10 haplotypes out of only 20

sequences, with the lowest H_d in the Chad sequences, suggesting a directional

selection/fixed allele. The haplotypes cluster according to origin on the maximum likelihood

phylogenetic tree, except for Niger (Additional Figure S8c).

Investigation of the role of intergenic region elements in overexpression of *GSTe2*

and *CYP6Z2*

Investigation of polymorphism in the intergenic region/regulatory elements

To investigate polymorphisms in the regulatory elements of the above genes, the 351

intergenic regions of *GSTe2* (spanning the 43 bp 3'-UTR of *GSTe1*, 248 bp flanking sequence

and 60 bp 5′-UTR of *GSTe2*) preceding the start codon were amplified from 10 each of DDT-

alive and -dead females from the 4 Sahel countries, as well as from Ngoussou females,

successfully. For *CYP6Z2*, 1078 bp (spanning 38 bp 3′ UTR of *CYP6Z1*, 937 bp flanking

sequence and 103 bp 5′UTR of *CYP6Z2*, preceding the start codon of *CYP6Z2*) were used to

amplify fragments from 10 each of deltamethrin-alive and -dead females from Nigeria, and

from Ngoussou females.

Out of the 90 *GSTe2* 5′-UTR sequences analysed differences were observed between the alive

and dead mosquitoes, with a total of 65 sequences from alive and dead females (regardless

of country of origin) identical to the 10 sequences of Ngoussou. From the 40 sequences of the

dead females, 39 were identical to Ngoussou (regardless of the country). Twelve sequences,

all from the alive females (from across the four countries) were similar, with several

mutations in putative transcriptional factors binding sites, which may impact

overexpression of the *GSTe2.* In short, 8 mutations were shared in common between these 12

sequences of DDT-alive females from across the countries (3 each from Nigeria, Chad and

Cameroon) and additional mutations in the Niger samples (4 sequences). These mutations

include (i) T->A transition withing the cellular myb-DNA (c-myb) binding domain

(Additional Figure S9), (ii) a T->C transversion in the zinc-finger homeodomain, δEF1 (δ

elongation factor 1) binding site, (iii) T->A transition in the nuclear matric protein 4 (NMP4),

(iv) simultaneous insertion of adenine and a transition T -> C, in positions 113 and 114

respectively, between the Fork-head box L1 and c-EST binding sites, (v) an A -> C

transversion in a second NMP4, (vi) a T -> C transition, six nucleotides downstream the

nuclear factor κB (NF-κB), (vii) followed by an A -> G transversion 3 nucleotides

downstream the NF-κB, (viii) a G -> A transversion, 7 nucleotides upstream the GC box, (ix)

a C -> G transition, 2 nucleotides downstream the arthropod initiator (Inr consensus)

sequence, and finally (x) a C -> G transition, within the 5'-UTR of the *GSTe2*, 28 nucleotides

downstream the transcriptional start site (49 nucleotides upstream the *GSTe2* start codon).

Analysis of the 90 sequences revealed a very low polymorphism in the dead mosquitoes (S =

0, for Niger-dead, Chad-dead and Ngoussou), but high polymorphism in the alive (S = 16 for

Nigeria-alive, and 13 each for Chad-alive and Cameroon-alive) (Additional Table 6). All

sequences produced 17 polymorphic sites, with 6 haplotypes (Additional Figure S10a and -

b). High haplotype diversities were obtained from the alive mosquitoes (for example,

Nigeria alive, Hd = 0.71, Niger-, Hd = 0.53). Regardless of country of origin, the haplotypes

cluster according to phenotype on the maximum likelihood phylogenetic tree, with the alive

haplotypes forming a distinct (separate) clade (Additional Figure S10c).

With regard to *CYP6Z2* no major differences were observed when comparing the

deltamethrin-alive and -dead sequences with the Ngoussou.

Measurement of activities of the 5' regulatory elements of the GSTe2 and CYP6Z2

Initial promoter analyses were conducted with the 5'- regulatory element sequences of

GSTe2, for the DDT-alive females (representative sequence with the 8 common polymorphic

positions, designated, Sahel-alive), comparing it with the sequence from the DDT-

dead/Ngoussou (designated Ngoussou/Sahel-dead). For *CYP6Z2*, a predominant sequence

from the Nigeria field sample (with 8 nucleotide insertion) was compared with the

Ngoussou sequence.

The ability to drive heterologous expression of the firefly luciferase was determined, with

the Sahel-alive construct and Ngoussou/Sahel-dead producing increased luciferase activity

[~3090$\times$ (normalized luciferase activity = 2.58) and 479$\times$ (normalized activity = 0.400),

respectively] compared with the promoterless pGL3-Basic vector. But the Sahel-alive

construct was significantly more active than the Ngoussou/Sahel dead counterpart

[promoter activity ~6-fold higher (Tukey HSD Q = 8.11, p = 0.004)]. In contrast, for *CYP6Z2*

no significant differences were observed when comparing the field construct (normalized

luciferase activity = 5.95 for alive/dead field construct), compared with 5.84 for the

Ngoussou construct.

GSTe2 promoter delineation and measurement of activity

Sequential deletion of the intergenic region of the *GSTe2* resulted in progressive reduction in

luciferase activity. Deletion of the 43 bp 3'-UTR of the *GSTe1* (-308 from the start codon of

GSTe2) reduced activity of the Sahel-alive construct by only 18.6 % suggesting that the c-

Myb transcriptional factor binding site may not be critical for overexpression (Figure 5).

Deletion of an additional 38 nucleotides from the flanking region (-270 fragment,

obliterating the δEF1 binding site) had comparable impact as above, with activities reducing

by 15.3 % only. But removing the *GSTe-1* 3'-UTR and an additional 46 nucleotides from the

flanking region (-262 fragment, which obliterated in addition the FOX-L1 putative binding

site) reduced activity by 64.1 % (Q = 6.18, p = 0.01) indicating the importance of this binding

site. Deletion of *GSTe1* 3'-UTR plus 75 nucleotides of the flanking region (-232 fragment,

located 10 nucleotides from the c-EST binding site) significantly reduced the activity by

~70% (Q = 6.14, p = 0.01), suggesting the importance of the simultaneous insertion of adenine

and transition T -> C, in positions 113 and 114 respectively, between the FOX-L1 and c-EST

binding sites. Removing the fragment of the 5′UTR of *GSTe2* (43 nucleotides preceding the

AUG codon obliterated the promoter activity, reducing the luciferase expression by 98 % (Q

= 9.41, p = 0.002). This is despite presence of all the above binding sites and transcriptional

start site in this fragment. Deletion of the 5′-UTR from Ngoussou significantly reduced

activities as well (reduction by 88 % compared with the full Ngoussou intergenic region

construct, Q = 13.75, p = 0.001). Overall, these findings suggest that the essential binding sites

for overexpression of *GSTe2* span the FOX-L1 and c-EST binding sites, with the 5′-UTR

essential for activity.

Investigating the role of *GSTe2* and *CYP6Z2* in insecticides resistance using

transgenic analysis

A qRT-PCR was conducted using the transgenic flies to first establish overexpression of the

above genes. Relative fold changes (FC) of 32.7 ± 4.6 and 25.30 ± 2.8 were obtained in flies

overexpressing the *GSTe2* and *CYP6Z2* respectively, compared with control flies (progenies

of crosses between the parental line flies with no gene insertion, crossed with GAL4/UAS

driver line) (Additional Figure S11).

Contact bioassays carried out using 0.05 % α-cypermethrin revealed a high susceptibility in

the transgenic flies expressing *GSTe2* (Act5C-GAL4-UAS-GSTe2) and controls (Figure 6a),

with mortalities increasing from 62 % and 70 % respectively in 1 h, to 94 % and 99 % in 24 h.

However, significantly reduced mortalities were observed in transgenic flies expressing

CYP6Z2 (Act5C-GAL4-UAS-CYP6Z2) compared to control flies, at 1 h (mortality = 35 % *vs* 70

% in control, p < 0.001) and 3 h (mortality = 52 % vs 89 % in the control, p < 0.001). High

susceptibilities were also seen in all the experimental flies exposed to 0.05 % deltamethrin

(both for *GSTe2* and *CYP6Z2* flies), except for 1 h with flies expressing *CYP6Z2* (mortality =

47 % compared with 58 % in the control flies, p < 0.010).

The *GSTe2* transgenic flies were highly resistant to 2 % permethrin, with no mortality at all

in 1 h (Figure 6b), and average mortalities of only 8.5 % at 24 h (p < 0.0001), compared with

85 % at 1 h for control flies, which increased to 100 % from 3 h.

Initial exposure to 4 % of 3-phenoxybenzaldehyde (PBAld) and 3-phenoxybenzyl alcohol

(PBAlc) had no toxic effect on *CYP6Z2* transgenic flies (Figure not shown). However, 5x

concentration of these primary products of pyrethroid hydrolysis induced mortalities, albeit

low in all flies (Figure 6c). For PBAld, mortalities ranged from 15.5 % and 17.5 % at 1 h, for

CYP6Z2 transgenic flies and control flies respectively, to 30 % and 27.5 % at 24 h. Although

low mortalities were observed with PBAlc, but at 1 h and 3 h exposure times the mortalities

in the transgenic flies expressing *CYP6Z2* were significantly lower compared with

mortalities from the control flies (1 h mortality = 2.5 % *vs* 12.5 %, p < 0.001; 3 h mortality = 9.5

% *vs* 17.5 %, p < 0.001).

Marginal tolerance towards DDT was observed in the *CYP6Z2* transgenic flies when

compared with the control flies, but only at 3 h (p < 0.001) and 6 h (p < 0.01) (Figure 6d). This

is in contrast with the *GSTe2* transgenic flies, with very low mortalities, in ranges of 0 % to

9.52 % for 1 h to 24 h, when compared with control flies (mortality = 34 % in 1 h and 100 % in

6 – 24 h, p < 0.0001).

Susceptibility to clothianidin was very high (Figure 6d), but surprisingly the *CYP6Z2*

transgenic flies exhibited a contrasting phenotype, with significantly higher mortalities

compared with the control flies, at 1 h (mortality = 67 % vs 42 %, $p < 0.001$), 3 h (mortality =

71 % vs 53 %, $p < 0.001$) and 6 h (mortality = 78 % vs 62 %, $p < 0.01$).

Taken together, these results confirmed that over-expression of *GSTe2* alone is sufficient to

confer resistance to type I pyrethroid (permethrin) and DDT, while overexpression of

CYP6Z2 alone, may confer marginal resistance to the α-cypermethrin and PBAlc.

Discussion

Escalating insecticide resistance across Africa (7-9, 52), if not tackled will compromise the

malaria control and elimination efforts. Molecular markers of metabolic resistance e.g., (10,

11, 21), will support evidence-based control and resistance management. Identification and

validation of resistance markers in malaria vectors across regions of sub-Saharan Africa can

promote communication, cooperation and coordination among malaria control/elimination

programs, and allow control efforts to be tailored to the vector species involved in

transmission across borders (53), and tracking of the evolution and spread of resistance

markers across regions (22). To support malaria pre-elimination efforts in Africa, in this

study we targeted the Sudan savannah and Sahel (regions of Africa sharing similar eco-

climatic conditions, and characterised by high seasonal transmission), which are ideal for

control and elimination of malaria using seasonal vector control (25) and chemoprevention.

Anopheles coluzzii **is a major malaria vector across the Sudano-Sahelian transects**

Contrary to the previous observations that *An. arabiensis* tends to predominate in arid

savannas, while *An. gambiae* is the dominant species in humid forest zones (54-56), *An.*

coluzzii has repeatedly been identified as the major malaria vector in the Sudan savannah

and Sahel of several, neighbouring countries (including in northern Nigeria, southern Niger,

central Chad, and northern Cameroon) (7, 26-28, 57) in recent years, suggesting that this

vector has adapted well in drier regions of the Sahel transects, and is probably

predominating over *An. arabiensis* and *An. gambiae s.s*. This is not surprising as this species

exhibits higher exploitation of breeding sites associated with anthropogenic activities, and

behavioural plasticity to avoid predators (58), and is known to survive a long dry season *in*

situ/aestivation, which allows it to predominate, becoming the primary force of malaria

transmission (59, 60).

Common metabolic genes mediate multiple resistance in the Sahelian *An. coluzzii*

Several genes shown to confer metabolic resistance in *Anopheles* mosquitoes and other

insects were found constitutively overexpressed and/or induced in this study. *GSTe2*

(AGAP009194) is one of the most regularly encountered metabolic genes in resistant

populations of the major malaria vectors *An. gambiae*, *An. coluzzii* and *An. funestus* (20-23).

Anopheles gambiae GSTe2 has been validated, using transgenic flies to confer DDT (20) and

fenitrothion (61) resistance. It was extensively studied in *An. funestus*, in which it was shown

to confer cross-resistance to DDT and permethrin (22), reduce efficacy of the LLINs,

PermaNet 2.0 and PermaNet 3.0 (side panels) (23), and even increase the longevity of the

resistant populations carrying the 119F mutation (62). These and our findings suggest that

the overexpression of this GST alone can confer resistance to three insecticides from three

different classes (DDT, permethrin and fenitrothion). The absence of mutations in the cDNA

coding sequences of *GSTe2* in *An. coluzzii* from these four countries, suggest that

overexpression of this GST is enough to confer resistance.

Several cytochrome P450s previously linked with insecticide resistance were found

overexpressed across the Sahel. For example, *An. gambiae CYP6Z2* (AGAP008218) known to

metabolize carbaryl (63), the insect juvenile hormone analogue insecticide, pyriproxyfen (64)

and mitochondrial complex I inhibitors, fenazaquin, pyridaben and tolfenpyrad (65). This

P450 also plays a pivotal role in the clearance of pyrethroid insecticides via further

catabolism of pyrethroid derivatives (PBAld and PBAlc) obtained by the action of

carboxylesterases (66), in line with our findings of this gene conferring marginal tolerance to

high concentration of PBAlc, and α-cypermethrin. However, our findings suggest that

overexpression of this P450 may enhance the efficacy of clothianidin, which will be

epidemiologically advantageous in terms of control. Indeed, bioactivation by P450s is

known to be a requirement for insecticidal toxicity of several classes of insecticides, e.g., the

organophosphates and chlorpenafyr. In contrast to *GSTe2*, the *CYP6Z2* from across Sahel

contain three cDNA mutations, which makes it different from Ngoussou coding sequences.

These are $K^{211}N$ and $T^{218}S$ mutations, both within the substrate recognitions site 2, and an

$A^{282}E$. Other important CYP450s found to be commonly overexpressed across the Sahel

include the *CYP4C27* (AGAP009246), *CYP6Z3* (AGAP008217) and *CYP9K1* (AGAP000818),

all three shown to be consistently overexpressed in field populations of *An. gambiae/coluzzii*

and *An. funestus* across Africa (9, 67, 68). *CYP9K1* has been shown to be epidemiologically

important pyrethroid-metabolising P450 linked with metabolism of deltamethrin and

pyriproxyfen in *An. gambiae* (69). We also have recently shown that his P450 is involved in

pyrethroids in *An. funestus* (70).

Several carboxylesterases were also upregulated/induced across the Sahel, with the

COEBE3C (AGAP005372) upregulated in all four countries. This beta esterase is enriched in

the legs (where xenobiotic detoxification probably occur) of pyrethroid-resistant *An. coluzzii*

(71). Other genes consistently overexpressed across the Sahel, include the chymotrypsins

(*CHYM3*/AGAP006711 and *CHYM1*/AGAP006709) and a lipase (AGAP002353).

Chymotrypsins are known to defend insects against plants' proteinase inhibitors (72); and

previous transcriptional studies have shown *CHYM1* and *CHYM3* overexpressed in

insecticide resistant populations of *An. gambiae* and *An. coluzzii*, respectively (9, 67). Using *in*

vitro and *in vivo* tools, lipases have been linked with deltamethrin resistance in *Culex pipiens*

pallens (73).

Common metabolic resistance markers probably exacerbate resistance across the

Sahel

Several well-known genetic variants implicated in resistance, as well as the recently

discovered ones exist in high frequencies in the Sahelian *An. coluzzii*, compared with the

Ngoussou. For example, the pyrethroid resistance *CYP4J5*-L43F marker (74) was found fixed

across the Sahel. The G280S/G119S *ace*-1 mutation, found in high frequencies across Sahel

confers organophosphate an carbamates resistance (75), and is shown recently to confer

resistance to pirimiphos-methyl, in *An. coluzzii/gambiae* (76). Several mutations found within

the VGSC have recently been described/validated. For example, the resistance mutation,

L995F (77) and the V402L/I15227T haplotypes have been observed across Africa (78).

Recently, the two mutations (V402L/I15227T) are described to be in tight linkage and

mutually exclusive to the classical L995F/S mutations (77). Our results suggest haplotypes

carrying the V402L/I1527T combination plus the L995 replacement do exist in the Sahel *An.*

coluzzii. Not only that, it is also in addition to the N1570Y replacement. However, isolation

of Ngoussou colony over generations in the artificial insectary conditions could have led to

genetic isolation and drift, from inbreeding, leading to strong population differentiation in

comparison to the natural populations studied here.

In contrast to *An. funestus GSTe2*, where overexpression and 119F mutation combined to

confer extreme DDT resistance (22), the absence of amino acid replacements in the *An.*

coluzzii GSTe2 from the Sahel suggests that overexpression alone is the key mediator of DDT

and permethrin resistance. This is supported by the higher activity in the regulatory regions,

harbouring an insertion and nucleotide substitutions in the alive mosquitoes. Indeed, some

of the mutations we have found within the intergenic region of *GSTe2* are similar to those

observed in a previous study (79).

Cuticular resistance mechanism probably playing a key role in Sahelian *An.*

coluzzii

Our results suggest cuticular mechanism play a role in pyrethroid resistance in these

populations. For example, the findings of *CYP4G16* (AGAP001076) and *CYP4G17*

(AGAP000877) overexpressed across the Sahel. The former P450 was previously shown to be

involved in epicuticular hydrocarbon biosynthesis associated with resistance (80).

The three major classes of the insect cuticular proteins - the CPR, CPLC and CPAP, were

found overrepresented in the top overexpressed genes across the Sahel, with the commonly

upregulated ones being *CPR76*, *CPR15* and *CPR30*, a chitin-binding cuticular protein,

CPAP3-A1b, and cuticular proteins of low complexity, *CPLCX3* and *CPLCA1*. Indeed,

CPAP3-A1b (AGAP000987) have been shown to be highly overexpressed in deltamethrin-

resistant Sahelian population of *An. coluzzii* from Burkina Faso (9) and induced by blood

feeding in *An. gambiae* (81). The CPR and CPLC cuticular proteins have been described to

potentially play a crucial role in insecticide resistance through leg cuticle

remodelling/thickening, regulating penetration rate of insecticides in *An. coluzzii* (71).

Furthermore, a recent study has found most of the cuticular proteins we have described

here, as highly overexpressed in permethrin/malathion resistant populations of Ethiopian

An. arabiensis (82), e.g., *CPR30*, *CPR75*, *CPR81*, and *CPLCP11*. Out of the several chitinases

overexpressed across the Sahel, four were amongst the top 50 most overexpressed metabolic

genes. These include the *Cht24* and *Cht6* that have been shown to be overexpressed in the

An. arabiensis from the above study (82), with the ortholog of *Cht24*, AARA007329 among the

top 10 most overexpressed genes in *An. arabiensis*, in line with our observations in *An.*

coluzzii across the Sahel. There is an overwhelming need to functionally investigate the

role/contribution towards insecticide resistance of these cuticular proteins, chitinases and a

chitin synthase (AGAP001748) significantly overexpressed in the field *An. coluzzii* from

Nigeria and Chad.

Insecticide resistance- and thermotolerance-associated genes sit within

chromosomal inversions

In this study, the findings of high frequency of 2La, 2Rb and 2Rc inversion polymorphisms

in the populations of *An. coluzzii*, compared with the Ngoussou, suggested strong

phenotypic adaptations in this species, across the Sahel. Most importantly, in addition to

several of the cuticular protein genes associated with resistance (chitinases, chitin synthase,

CPR, CPLC and CPAP proteins), several other genes previously implicated in

thermotolerance and/or desiccation resistance in *An. gambiae/coluzzii*, and which were highly

overexpressed in this study sit within these inversions. For example, the heat shock proteins,

hsp83 (AGAP006958) and *hsp90* hptG (AGAP006961), both of which are known to be heat-

and insecticide-stress inducible (83) and were among the core set of hsp genes involved in a

common and immediate response to thermal stress in *An. gambiae* populations (84), sit

within the 2La inversion. These two genes were among the overexpressed genes in both

heat-hardened and permethrin-resistant *An. coluzzii* populations from northern Nigeria (57).

Several ionotropic glutamate receptors were found within the 2La inversion breakpoints:

IR136 (AGAP006440), *IR139* (AGAP006691) and *IR142* (AGAP006407). This is not surprising

as ionotropic receptors are commonly associated with chemosensation, thermosensation,

and hygrosensation (85, 86), characteristics

which can confer adaptive advantages in xeric environs. The *IR25a* (AGAP010272) and *IR21*

a (AGAP008511) are known to mediate both humidity and temperature preference in the

fruit fly, *D. melanogaster* (86, 87), in addition to *IR21a* driving heat seeking and heat-

stimulated blood feeding in *An. gambiae* (87). These two genes have been shown to be

overexpressed/induced in thermotolerant/permethrin-resistant populations of *An. coluzzii*

(57).

Conclusions

Information on molecular basis of resistance and/or resistance genes and its markers

facilitates evidence-based control measures. In this study we characterised a major malaria

vector, *An. coluzzii* from the Sahel region of four countries, with findings which could

promote evidence-based, cross-border policy towards local and regional malaria control.

The study found that across Sahel (where malaria is highly seasonal, reaching its peak in the

rainy season), *An. coluzzii* is a dominant vector. And that a handful of common cross-

resistance genes are responsible for multiple insecticide resistance in this species. Findings

from this study suggest pleotropic role of some key genes – able to confer insecticide

resistance and/or stabilize the insecticide resistance gene, at the same time conferring

environmental adaptations, such as the ability to survive thermal stress (thermotolerance),

as expected in this Sahelian region. From operational vector control perspective this study

provided evidence of the role of key insecticide metabolism gene, *CYP6Z2* in increasing

insecticidal potency of clothianidin, which could increase the efficacy of the ingredients in

malaria control tools, when targeting field populations overexpressing this key P450.

Declarations

Ethics approval and consent to participate

This study did not use human participants, human data or human tissue. Ethical approval

for collection of indoor resting female mosquitoes in Nigeria, Niger, Chad and Cameroon

have been provided in the references cited in the Study Site and Mosquito Sampling

(Materials and Methods).

Consent for publication

Not Applicable.

Availability of data and materials

The dataset(s) supporting the conclusions of this article are available in the European

Nucleotide Archive, with accession PRJEB51644, and secondary accession of ERP136291, for

the RNA-seqe raw sequence reads. cDNA sequences of *GSTe2* and *CYP6Z2* were deposited

in GenBank (accession numbers: ON169006 - ON169030 for *GSTe2* and ON169031 -

ON169050 for *CYP6Z2*) and 5'-UTR DNA fragment sequences were deposited in GenBank

(accession numbers: ON169051 - ON169140).

Competing interests

The authors declare that they have no competing interests.

Funding

995 This research was funded in whole, by the Wellcome Trust [Grant numbers:

996 WT201918/Z/16/Z to SSI and WT217188/Z/19/Z to CSW]. For the purpose of open access, the

997 author has applied a CC BY public copyright licence to any Author Accepted Manuscript

998 version arising from this submission. The Wellcome Trust had no role in the design of this

999 study and collection, analysis, and interpretation of data and in writing of this manuscript.

1000

Authors' contributions

1001

Conceived and designed by SSI and CSW. SSI carried out the molecular analyses, with

support from AM, LMJM, EIP and HI. ANF and MMM, participated in field collection of

mosquitoes in Nigeria, Chad and Cameroon. SSI carried out data analysis with the support

of JH, GDW, SCN for the RNAseq component. SSI wrote the manuscript with inputs from

CSW, JH and SCN. All authors contributed to corrections of the final draft and approved

final version of the manuscript.

1008

Acknowledgements

1009

1010 Not Applicable

1011

1012

1013

Reference

1.   Bhatt S, Weiss D, Cameron E, Bisanzio D, Mappin B, Dalrymple U, et al. The effect of malaria control on Plasmodium falciparum in Africa between 2000 and 2015. Nature. 2015;526(7572):207.

2.   WHO. Global Technical Strategy for Malaria 2016-2030. World Health Organisation. . Geneva, Switzerland; 2015 May 2015. Report No.: ISBN 978 92 4 156499 1.

3.   WHO. World Malaria Report. World Health Organization. . Geneva Switzerland; 2018. Report No.: Licence: CC BY-NC-SA 3.0 IGO Contract No.: ISBN 978-92-4-156565-3.

4.   WHO. World Malaria Report. Geneva: Switzerland; 2021. Report No.: 978-92-4-004049-6 Contract No.: 978-92-4-004049-6

5.   WHO. World Malaria Report 2019. Geneva, Switzerland; 2019. Report No.: 978-92-4-156572-1.

6.   WHO. Update on the E-2020 initiative of 21 malaria-eliminating countries: report and country briefs. Geneva. Switzerland: World Health Organization; 2018. Report No.: WHO/CDS/GMP/2018.10.

7.   Ibrahim SS, Mukhtar MM, Datti JA, Irving H, Kusimo MO, Tchapga W, et al. Temporal escalation of Pyrethroid Resistance in the major malaria vector Anopheles coluzzii from Sahelo-Sudanian Region of northern Nigeria. Scientific reports. 2019;9(1):7395.

8.   Riveron JM, Huijben S, Tchapga W, Tchouakui M, Wondji MJ, Tchoupo M, et al. Escalation of Pyrethroid Resistance in the Malaria Vector Anopheles funestus Induces a Loss of Efficacy of Piperonyl Butoxide-Based Insecticide-Treated Nets in Mozambique. J Infect Dis. 2019;220(3):467-75.

9.   Toe KH, N'Fale S, Dabire RK, Ranson H, Jones CM. The recent escalation in strength of pyrethroid resistance in Anopheles coluzzi in West Africa is linked to increased expression of multiple gene families. BMC Genomics. 2015;16:146.

10.   Weedall GD, Mugenzi LMJ, Menze BD, Tchouakui M, Ibrahim SS, Amvongo-Adjia N, et al. A cytochrome P450 allele confers pyrethroid resistance on a major African malaria vector, reducing insecticide-treated bednet efficacy. Sci Transl Med. 2019;11(484).

11.   Mugenzi LMJ, Menze BD, Tchouakui M, Wondji MJ, Irving H, Tchoupo M, et al. Cis-regulatory CYP6P9b P450 variants associated with loss of insecticide-treated bed net efficacy against Anopheles funestus. Nat Commun. 2019;10(1):4652.

12.   Barnes KG, Weedall GD, Ndula M, Irving H, Mzihalowa T, Hemingway J, et al. Genomic Footprints of Selective Sweeps from Metabolic Resistance to Pyrethroids in African Malaria Vectors Are Driven by Scale up of Insecticide-Based Vector Control. PLoS Genet. 2017;13(2):e1006539.

13.   Weedall GD, Riveron JM, Hearn J, Irving H, Kamdem C, Fouet C, et al. An Africa-wide genomic evolution of insecticide resistance in the malaria vector Anopheles funestus involves selective sweeps, copy number variations, gene conversion and transposons. PLoS Genet. 2020;16(6):e1008822.

14.   Bruce-Chwatt LJ. Malaria in Nigeria. Bulletin of the World Health Organization. 1951;4(3):301-27.

15.   Anopheles gambiae Genomes C, Data analysis g, Partner working g, Sample c-A, Burkina F, Cameroon, et al. Genetic diversity of the African malaria vector Anopheles gambiae. Nature. 2017;552(7683):96-100.

16.   Muller P, Warr E, Stevenson BJ, Pignatelli PM, Morgan JC, Steven A, et al. Field-caught permethrin-resistant Anopheles gambiae overexpress CYP6P3, a P450 that metabolises pyrethroids. PLoS Genet. 2008;4(11):e1000286.

17.   Yunta C, Hemmings K, Stevenson B, Koekemoer LL, Matambo T, Pignatelli P, et al. Cross-resistance profiles of malaria mosquito P450s associated with pyrethroid resistance against WHO insecticides. Pestic Biochem Physiol. 2019;161:61-7.

18.   Stevenson BJ, Bibby J, Pignatelli P, Muangnoicharoen S, O'Neill PM, Lian LY, et al. Cytochrome P450 6M2 from the malaria vector Anopheles gambiae metabolizes pyrethroids:

Sequential metabolism of deltamethrin revealed. Insect biochemistry and molecular biology.
2011;41(7):492-502.
19. Mitchell SN, Stevenson BJ, Müller P, Wilding CS, Egyir-Yawson A, Field SG, et al. Identification
and validation of a gene causing cross-resistance between insecticide classes in Anopheles gambiae
from Ghana. Proceedings of the National Academy of Sciences. 2012;109(16):6147-52.
20. Mitchell SN, Rigden DJ, Dowd AJ, Lu F, Wilding CS, Weetman D, et al. Metabolic and target-
site mechanisms combine to confer strong DDT resistance in Anopheles gambiae. PLoS One.
2014;9(3):e92662.
21. Lucas ER, Rockett KA, Lynd A, Essandoh J, Grisales N, Kemei B, et al. A high throughput multi-
locus insecticide resistance marker panel for tracking resistance emergence and spread in Anopheles
gambiae. Sci Rep. 2019;9(1):13335.
22. Riveron JM, Yunta C, Ibrahim SS, Djouaka R, Irving H, Menze BD, et al. A single mutation in
the GSTe2 gene allows tracking of metabolically based insecticide resistance in a major malaria
vector. Genome Biol. 2014;15(2):R27.
23. Menze BD, Kouamo MF, Wondji MJ, Tchapga W, Tchoupo M, Kusimo MO, et al. An
Experimental Hut Evaluation of PBO-Based and Pyrethroid-Only Nets against the Malaria Vector
Anopheles funestus Reveals a Loss of Bed Nets Efficacy Associated with GSTe2 Metabolic Resistance.
Genes (Basel). 2020;11(2).
24. Coluzzi M, Sabatini A, Petrarca V, Di Deco MA. Chromosomal differentiation and adaptation
to human environments in the Anopheles gambiae complex. Trans R Soc Trop Med Hyg.
1979;73(5):483-97.
25. Cairns M, Roca-Feltrer A, Garske T, Wilson AL, Diallo D, Milligan PJ, et al. Estimating the
potential public health impact of seasonal malaria chemoprevention in African children. Nat
Commun. 2012;3:881.
26. Ibrahim SS, Mukhtar MM, Irving H, Labbo R, Kusimo MO, Mahamadou I, et al. High
Plasmodium infection and multiple insecticide resistance in a major malaria vector Anopheles
coluzzii from Sahel of Niger Republic. Malaria journal. 2019;18(1):181.
27. Fadel AN, Ibrahim SS, Tchouakui M, Terence E, Wondji MJ, Tchoupo M, et al. A combination
of metabolic resistance and high frequency of the 1014F kdr mutation is driving pyrethroid
resistance in Anopheles coluzzii population from Guinea savanna of Cameroon. Parasit Vectors.
2019;12(1):263.
28. Ibrahim SS, Fadel AN, Tchouakui M, Terence E, Wondji MJ, Tchoupo M, et al. High insecticide
resistance in the major malaria vector Anopheles coluzzii in Chad Republic. Infect Dis Poverty.
2019;8(1):100.
29. Mitchell SN, Stevenson BJ, Muller P, Wilding CS, Egyir-Yawson A, Field SG, et al. Identification
and validation of a gene causing cross-resistance between insecticide classes in Anopheles gambiae
from Ghana. Proceedings of the National Academy of Sciences of the United States of America.
2012;109(16):6147-52.
30. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet journal. 2011;17(1):10-2.
31. Joshi N, Fass J. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files
(Version 1.33)[Software]. 2011.
32. Aronesty E. ea-utils: Command-line tools for processing biological sequencing data. Durham,
NC; 2011.
33. Love MI, Huber, W., & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology. 2014;15(550).
34. Blighe K, S. Rana, and M. Lewis. . "EnhancedVolcano: Publication-ready volcano plots with
enhanced colouring and labeling.". 2018.
35. Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. Bioinformatics. 2016;32(18):2847-9.
36. Alexa A. RJ. topGO: Enrichment Analysis for Gene Ontology. R package version 2.46.0. 2021 [

37.   Supek F, Bosnjak M, Skunca N, Smuc T. REVIGO summarizes and visualizes long lists of gene
ontology terms. PLoS One. 2011;6(7):e21800.
38.   Riveron JM, Ibrahim SS, Chanda E, Mzilahowa T, Cuamba N, Irving H, et al. The highly
polymorphic CYP6M7 cytochrome P450 gene partners with the directionally selected CYP6P9a and
CYP6P9b genes to expand the pyrethroid resistance front in the malaria vector Anopheles funestus
in Africa. BMC Genomics. 2014;15(1):817.
39.   Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method.
Nature protocols. 2008;3(6):1101-8.
40.   Nagi CS, Oruni A, Weetman D, Donnelly J. M. RNA-Seq-Pop: Exploiting the sequence in RNA-
Seq - a Snakemake workflow reveals patterns of insecticide resistance in the malaria vector
Anopheles gambiae. 2022;2022.06.17.493894.
41.   Hudson RR, Slatkin M, Maddison WP. Estimation of levels of gene flow from DNA sequence
data. Genetics. 1992;132(2):583-9.
42.   Yi X, Liang Y, Huerta-Sanchez E, Jin X, Cuo ZX, Pool JE, et al. Sequencing of 50 human exomes
reveals adaptation to high altitude. Science. 2010;329(5987):75-8.
43.   Alistair Miles MFR, Peter Ralph, Nick Harding, Rahul Pisupati, and Summer Rae. cggh/scikit-
allel: A Python package for exploring and analysing genetic variation data 2020 [cggh/scikit-allel:
v1.3.1 (v.3.1). Zenodo.].
44.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.
45.   Love RR, Redmond SN, Pombi M, Caputo B, Petrarca V, Della Torre A, et al. In Silico
Karyotyping of Chromosomally Polymorphic Malaria Mosquitoes in the Anopheles gambiae Complex.
G3 (Bethesda). 2019;9(10):3249-62.
46.   Hall TA, editor BioEdit: a user-friendly biological sequence alignment editor and analysis
program for Windows 95/98/NT. Nucleic acids symposium series; 1999.
47.   Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular Evolutionary Genetics
Analysis across Computing Platforms. Molecular biology and evolution. 2018;35(6):1547-9.
48.   Rozas J, Ferrer-Mata A, Sanchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE, et
al. DnaSP 6: DNA Sequence Polymorphism Analysis of Large Data Sets. Molecular biology and
evolution. 2017;34(12):3299-302.
49.   Cartharius K, Frech K, Grote K, Klocke B, Haltmeier M, Klingenhoff A, et al. MatInspector and
beyond: promoter analysis based on transcription factor binding sites. Bioinformatics.
2005;21(13):2933-42.
50.   Lynd A, Lycett GJ. Optimization of the Gal4-UAS system in an Anopheles gambiae cell line.
Insect Mol Biol. 2011;20(5):599-608.
51.   Ibrahim SS, Riveron JM, Bibby J, Irving H, Yunta C, Paine MJI, et al. Allelic Variation of
Cytochrome P450s Drives Resistance to Bednet Insecticides in a Major Malaria Vector. PLoS Genet.
2015;11(10):e1005618.
52.   Riveron JM, Tchouakui M., Mugenzi L., Menze B. D., Mu-Chun C. and Wondji C. S. Insecticide
Resistance in Malaria Vectors: An Update at a Global Scale. In: S. Manguin VD, editor. Towards
Malaria Elimination - A Leap Forward. London: IntechOpen; 2018. p. 452.
53.   Shiff C. Integrated approach to malaria control. Clinical microbiology reviews.
2002;15(2):278-93.
54.   Lindsay S, Parson, L. and Thomas, C. . Mapping the range and relative abundance of the two
principal African malaria vectors, Anopheles gambiae sensu stricto and An. arabiensis, using climate
data. Proceedings of the Royal Society of London B: Biological Science. 1998; 265((1399)): 847–54.
55.   Coetzee M, Craig, M. and Le Sueur, D. . Distribution of African malaria mosquitoes belonging
to the Anopheles gambiae complex. Parasitology today. 2000;16(2):74-7.
56.   Ranson H, Abdallah H, Badolo A, Guelbeogo WM, Kerah-Hinzoumbe C, Yangalbe-Kalnone E,
et al. Insecticide resistance in Anopheles gambiae: data from the first year of a multi-country study
highlight the extent of the problem. Malar J. 2009;8:299.

57.   Ibrahim SS, Mukhtar MM, Muhammad A, Wondji CS. 2La Paracentric Chromosomal Inversion and Overexpressed Metabolic Genes Enhance Thermotolerance and Pyrethroid Resistance in the Major Malaria Vector Anopheles gambiae. Biology (Basel). 2021;10(6).

58.   Gimonneau G, Bouyer J, Morand S, Besansky NJ, Diabate A, Simard F. A behavioral mechanism underlying ecological divergence in the malaria mosquito Anopheles gambiae. Behav Ecol. 2010;21(5):1087-92.

59.   Dao A, Yaro AS, Diallo M, Timbine S, Huestis DL, Kassogue Y, et al. Signatures of aestivation and migration in Sahelian malaria mosquito populations. Nature. 2014;516(7531):387-90.

60.   Lehmann T, Weetman D, Huestis DL, Yaro AS, Kassogue Y, Diallo M, et al. Tracing the origin of the early wet-season Anopheles coluzzii in the Sahel. Evol Appl. 2017;10(7):704-17.

61.   Adolfi A, Poulton B, Anthousi A, Macilwee S, Ranson H, Lycett GJ. Functional genetic validation of key genes conferring insecticide resistance in the major African malaria vector, Anopheles gambiae. Proceedings of the National Academy of Sciences of the United States of America. 2019;116(51):25764-72.

62.   Tchouakui M, Riveron JM, Djonabaye D, Tchapga W, Irving H, Soh Takam P, et al. Fitness Costs of the Glutathione S-Transferase Epsilon 2 (L119F-GSTe2) Mediated Metabolic Resistance to Insecticides in the Major African Malaria Vector Anopheles Funestus. Genes (Basel). 2018;9(12).

63.   Chiu TL, Wen Z, Rupasinghe SG, Schuler MA. Comparative molecular modeling of Anopheles gambiae CYP6Z1, a mosquito P450 capable of metabolizing DDT. Proceedings of the National Academy of Sciences of the United States of America. 2008;105(26):8855-60.

64.   Yunta C, Grisales N, Nasz S, Hemmings K, Pignatelli P, Voice M, et al. Pyriproxyfen is metabolized by P450s associated with pyrethroid resistance in An. gambiae. Insect biochemistry and molecular biology. 2016;78:50-7.

65.   Lees RS, Ismail HM, Logan RAE, Malone D, Davies R, Anthousi A, et al. New insecticide screening platforms indicate that Mitochondrial Complex I inhibitors are susceptible to cross-resistance by mosquito P450s that metabolise pyrethroids. Sci Rep. 2020;10(1):16232.

66.   Chandor-Proust A, Bibby J, Regent-Kloeckner M, Roux J, Guittard-Crilat E, Poupardin R, et al. The central role of mosquito cytochrome P450 CYP6Zs in insecticide detoxification revealed by functional expression and structural modelling. Biochem J. 2013;455(1):75-85.

67.   Antonio-Nkondjio C, Poupardin R, Tene BF, Kopya E, Costantini C, Awono-Ambene P, et al. Investigation of mechanisms of bendiocarb resistance in Anopheles gambiae populations from the city of Yaounde, Cameroon. Malar J. 2016;15(1):424.

68.   Riveron JM, Ibrahim SS, Mulamba C, Djouaka R, Irving H, Wondji MJ, et al. Genome-Wide Transcription and Functional Analyses Reveal Heterogeneous Molecular Mechanisms Driving Pyrethroids Resistance in the Major Malaria Vector Anopheles funestus Across Africa. G3 (Bethesda). 2017;7(6):1819-32.

69.   Vontas J, Grigoraki L, Morgan J, Tsakireli D, Fuseini G, Segura L, et al. Rapid selection of a pyrethroid metabolic enzyme CYP9K1 by operational malaria control activities. Proceedings of the National Academy of Sciences of the United States of America. 2018;115(18):4619-24.

70.   Hearn J, Tagne CD, Ibrahim SS, Tene-Fossog B, Mugenzi LJ, Irving H, et al. Multi-omics analysis identifies a CYP9K1 haplotype conferring pyrethroid resistance in the malaria vector Anopheles funestus in East Africa. bioRxiv 2022;2021.10.21.465247.

71.   Kefi M, Charamis J, Balabanidou V, Ioannidis P, Ranson H, Ingham VA, et al. Transcriptomic analysis of resistance and short-term induction response to pyrethroids, in Anopheles coluzzii legs. BMC Genomics. 2021;22(1):891.

72.   Dunse KM, Kaas, Q., Guarino, R.F., Barton, P.A., Craik, D.J. and Anderson, M.A. Molecular basis for the resistance of an insect chymotrypsin to a potato type II proteinase inhibitor. Proceedings of the National Academy of Sciences. 2010;107(34):15016-21.

73.   Hu HX, Zhou D, Ma L, Shen B, Sun Y, Zhu CL. Lipase is associated with deltamethrin resistance in Culex pipiens pallens. Parasitol Res. 2020;119(1):23-30.

74. Weetman D, Wilding CS, Neafsey DE, Muller P, Ochomo E, Isaacs AT, et al. Candidate-gene
based GWAS identifies reproducible DNA markers for metabolic pyrethroid resistance from standing
genetic variation in East African Anopheles gambiae. Sci Rep. 2018;8(1):2920.
75. Weill M, Malcolm C, Chandre F, Mogensen K, Berthomieu A, Marquine M, et al. The unique
mutation in ace-1 giving high insecticide resistance is easily detectable in mosquito vectors. Insect
Mol Biol. 2004;13(1):1-7.
76. Grau-Bove X, Lucas E, Pipini D, Rippon E, van 't Hof AE, Constant E, et al. Resistance to
pirimiphos-methyl in West African Anopheles is spreading via duplication and introgression of the
Ace1 locus. PLoS Genet. 2021;17(1):e1009253.
77. Williams J, Cowlishaw R, Sanou A, Ranson H, Grigoraki L. In vivo functional validation of the
V402L voltage gated sodium channel mutation in the malaria vector An. gambiae. Pest Manag Sci.
2022;78(3):1155-63.
78. Clarkson CS, Miles A, Harding NJ, O'Reilly AO, Weetman D, Kwiatkowski D, et al. The genetic
architecture of target-site resistance to pyrethroid insecticides in the African malaria vectors
Anopheles gambiae and Anopheles coluzzii. Mol Ecol. 2021;30(21):5303-17.
79. Ding Y, Hawkes N, Meredith J, Eggleston P, Hemingway J, Ranson H. Characterization of the
promoters of Epsilon glutathione transferases in the mosquito Anopheles gambiae and their
response to oxidative stress. Biochem J. 2005;387(Pt 3):879-88.
80. Balabanidou V, Kampouraki A, MacLean M, Blomquist GJ, Tittiger C, Juarez MP, et al.
Cytochrome P450 associated with insecticide resistance catalyzes cuticular hydrocarbon production
in Anopheles gambiae. Proceedings of the National Academy of Sciences of the United States of
America. 2016;113(33):9268-73.
81. Vannini L, Augustine Dunn W, Reed TW, Willis JH. Changes in transcript abundance for
cuticular proteins and other genes three hours after a blood meal in Anopheles gambiae. Insect
biochemistry and molecular biology. 2014;44:33-43.
82. Messenger LA, Impoinvil LM, Derilus D, Yewhalaw D, Irish S, Lenhart A. A whole
transcriptomic approach provides novel insights into the molecular basis of organophosphate and
pyrethroid resistance in Anopheles arabiensis from Ethiopia. Insect biochemistry and molecular
biology. 2021;139:103655.
83. Zhao LaJ, W.A. . Expression of heat shock protein genes in insect stress responses. .
Invertebrate Survival Journal. 2012;9(1):93-101.
84. Cassone BJ, Molloy MJ, Cheng C, Tan JC, Hahn MW, Besansky NJ. Divergent transcriptional
response to thermal stress by Anopheles gambiae larvae carrying alternative arrangements of
inversion 2La. Mol Ecol. 2011;20(12):2567-80.
85. VanHook AM. Understanding hygrosensation: How flies sense changes in humidity. Science
Signaling. 2016;9(430):ec127-ec.
86. Enjin A, Zaharieva EE, Frank DD, Mansourian S, Suh GS, Gallio M, et al. Humidity Sensing in
Drosophila. Curr Biol. 2016;26(10):1352-8.
87. Greppi C, Laursen WJ, Budelli G, Chang EC, Daniels AM, van Giesen L, et al. Mosquito heat
seeking is driven by an ancestral cooling receptor. Science. 2020;367(6478):681-4.

1255

Figure Titles

Figure 1: A volcano plot of differentially expressed genes showing fold changes and levels of significance in R-S, C-S and R-C comparisons, for Nigeria and Niger populations of *An. coluzzii*. The plot depicts the top upregulated and top downregulated genes for each

comparison, with several genes including chymotrypsins, a malate dehudrogenrase, CYP450s, a glutathione S-transferase (*GSTe2*), and a lipase, commonly upregulated in R vs S and C vs S comparisons. R stands for the female, resistant mosquitoes (survivors of 0.05% deltamethrin, 24 h after exposure), C stands for control (mosquitoes from the same population and of the same age not exposed to deltamethrin, and S stands for susceptible (females fromfully insecticide susceptible colony, Ngoussou).

Figure 2: A volcano plot of differentially expressed genes showing fold changes and significance in R-S, C-S and R-C comparisons, for Chad and Cameroon populations of *An. coluzzii.* The plot depicts the top upregulated and the top downregulated genes for each comparison with several genes including chymotrypsins, a malate dehudrogenrase, an aquaporn, CYP450s (particulary *CYP6Z2*), a glutathione S-transferase (*GSTe2*) and a lipase, commonly upregulated in R vs S and C vs S comparisons. R stands for the female, resistant mosquitoes (survivors of 0.05% deltamethrin, 24 h after exposure), C stands for control (mosquitoes from the same population and of the same age not exposed to deltamethrin, and S stands for susceptible (females from fully insecticide susceptible colony, Ngoussou).

Figure 3: Validation of candidate resistance genes. qRT-PCR profile of twelve metabolic resistance genes (in R-S and C-S comparisons) from the Sahelian *An. coluzzii*. Each bar (data point) is analysis from three biological replicates and three technical replicates for each gene, with error bars representing standard deviations. Significant differences are also indicated with the symbols †, †, † and ‡, for $p < 0.0001$, $p < 0.001$, $p < 0.01$ and $p < 0.05$, respectively.

Figure 4: Identification of resistance variants of interest. A heatmap showing frequencies

of the resistance variants (haplotypes) in some key genes of interest.

Figure 5: Characterization of intergenic region (5' regulatory element) of *GSTe2*. Results of

dual-luciferase reporter assays of the promoter (intergenic region) constructs, showing

progressive loss of activity following sequential deletion of the constructs.

Figure 6: Validation of the role of metabolic resistance genes in insecticide resistance.

Results of insecticide susceptibility bioassays with transgenic flies expressing *GSTe2* and

CYP6Z2. **a.** α-cypermethrin and deltamethrin, **b.** permethrin, **c.** 3-phenoxybenzaldehyde and

3-phenoxybenzyl alcohol, **d.** DDT and clothianidin.

1291

Additional Files

Additional File 1: Supplementary text for methods and results.

Additional File 2: Common differentially expressed genes across the Sahel.

Additional File 3: Population genetics analyses.

Additional File 4: Genes within 2La, 2Rb and 2Rc inversion polymorphisms.

Additional Tables: Additional tables, S1-S6.

Additional Figures: Additional figures, S1-S11.

Supplementary Figures

Figure S1: Sampling sites in the Sahel region of the four sampled countries.

Red line indicates median length
Box indicates interquartile range
Whiskers indicate minimum and maximum read lengths

Figure S2: Summary of total number of reads for each sample and distribution of trimmed read length for forward (R1) and reverse (R2) reads and reads unpaired after trimming (R0).

Figure S3: Venn diagrams summarizing the differentially expressed (DE) genes between resistant (R), unexposed (C) and susceptible (S) samples with a transcription ratio log₂FC ≥ 1 in either direction, and a corrected p value < 0.05.

Figure S4: Principal component analysis of the 500 most variable genes from all experimental arms in the data from the four countries.

Figure S5: A heatmap showing the top 50 overexpressed genes in *An. coluzzii* populations from the Sahel region of each country. a. for Nigeria **b.**, **c.** and **d.** for Niger, Chad and Cameroon. Several genes including chymotrypsins, CYP450s, aquaporins, glutathione S-transferases and carboxylesterases are commonly upregulated across the Sahel of the four countries.

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0004867	serine-type endopeptidase inhibitor activity	0.121%		0.0039	0.91	0.00
GO:0015184	L-cystine transmembrane transporter activity	0.001%		0.0030	0.98	0.00
GO:0016616	oxidoreductase activity, acting on the CH-OH group of donors, NAD or NADP as acceptor	1.855%		0.0095	0.80	0.00
GO:0030246	carbohydrate binding	0.797%		0.0023	0.98	0.00
GO:0003796	lysozyme activity	0.043%		0.0078	0.93	0.03
GO:0016408	C-acyltransferase activity	0.080%		0.0040	0.95	0.03
GO:0016846	carbon-sulfur lyase activity	0.134%		0.0030	0.94	0.03
GO:0016854	racemase and epimerase activity	0.375%		0.0042	0.93	0.03
GO:0005549	odorant binding	0.051%		0.0023	0.99	0.04
GO:0008061	chitin binding	0.116%		0.0000	0.98	0.05
GO:0016491	oxidoreductase activity	12.887%		0.0001	0.95	0.05
GO:0050660	flavin adenine dinucleotide binding	1.631%		0.0014	0.97	0.06
GO:0005507	copper ion binding	0.292%		0.0002	0.98	0.14
GO:0008483	transaminase activity	0.728%		0.0039	0.94	0.20
GO:0004364	glutathione transferase activity	0.086%		0.0040	0.95	0.20
GO:0018685	alkane 1-monooxygenase activity	0.001%		0.0040	0.88	0.23
GO:0005319	lipid transporter activity	0.146%		0.0017	0.98	0.30
GO:0015036	disulfide oxidoreductase activity	0.326%		0.0005	0.83	0.37
GO:0016645	oxidoreductase activity, acting on the CH-NH group of donors	0.350%		0.0030	0.83	0.37
GO:0004601	peroxidase activity	0.427%		0.0000	0.77	0.38

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0005085	guanyl-nucleotide exchange factor activity	0.309%		0.0070	0.90	0.00
GO:0016917	GABA receptor activity	0.027%		0.0011	0.95	0.00
GO:0022843	voltage-gated cation channel activity	0.155%		0.0060	0.43	0.00
GO:0031402	sodium ion binding	0.009%		0.0048	0.91	0.00
GO:0070006	metalloaminopeptidase activity	0.153%		0.0000	1.00	0.00
GO:0061980	regulatory RNA binding	0.013%		0.0031	0.99	0.03
GO:0005516	calmodulin binding	0.135%		0.0009	0.99	0.03
GO:0042277	peptide binding	0.100%		0.0001	0.99	0.04
GO:0008273	calcium, potassium:sodium antiporter activity	0.006%		0.0048	0.49	0.47
GO:0046961	proton-transporting ATPase activity, rotational mechanism	0.115%		0.0023	0.53	0.57
GO:0099095	ligand-gated anion channel activity	0.006%		0.0008	0.49	0.64
GO:0022848	acetylcholine-gated cation-selective channel activity	0.024%		0.0000	0.43	0.70
GO:0005283	amino acid:sodium symporter activity	0.045%		0.0031	0.45	0.72
GO:0005328	neurotransmitter:sodium symporter activity	0.001%		0.0000	0.58	0.73
GO:0008028	monocarboxylic acid transmembrane transporter activity	0.052%		0.0013	0.54	0.74
GO:0022839	ion gated channel activity	0.054%		0.0000	0.53	0.74
GO:0005254	chloride channel activity	0.123%		0.0000	0.45	0.80
GO:0005245	voltage-gated calcium channel activity	0.031%		0.0011	0.44	0.81
GO:0015171	amino acid transmembrane transporter activity	0.268%		0.0000	0.59	0.83
GO:0030955	potassium ion binding	0.068%		0.0048	0.91	0.86
GO:0005096	GTPase activator activity	0.247%		0.0003	0.90	0.87
GO:0005249	voltage-gated potassium channel activity	0.119%		0.0022	0.41	0.90

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0008010	structural constituent of chitin-based larval cuticle	0.002%		0.0000	1.00	0.00
GO:0016811	hydrolase activity, acting on carbon-nitrogen (but not peptide) bonds, in linear amides	0.737%		0.0084	0.93	0.00
GO:0042277	peptide binding	0.100%		0.0051	0.98	0.00
GO:0042626	ATPase-coupled transmembrane transporter activity	1.389%		0.0000	0.83	0.00
GO:0004732	pyridoxal oxidase activity	0.000%		0.0055	0.87	0.02
GO:0015020	glucuronosyltransferase activity	0.043%		0.0051	0.96	0.03
GO:0016491	oxidoreductase activity	12.887%		0.0032	0.97	0.05
GO:0050660	flavin adenine dinucleotide binding	1.631%		0.0044	0.96	0.05
GO:0008061	chitin binding	0.116%		0.0000	0.98	0.05
GO:0004364	glutathione transferase activity	0.086%		0.0000	0.96	0.16
GO:0003995	acyl-CoA dehydrogenase activity	0.152%		0.0038	0.92	0.16
GO:0018685	alkane 1-monooxygenase activity	0.001%		0.0000	0.93	0.20
GO:0070006	metalloaminopeptidase activity	0.153%		0.0006	0.89	0.23
GO:0030170	pyridoxal phosphate binding	1.138%		0.0038	0.96	0.31
GO:0005355	glucose transmembrane transporter activity	0.012%		0.0000	0.86	0.33
GO:0008239	dipeptidyl-peptidase activity	0.046%		0.0004	0.89	0.65
GO:0005351	carbohydrate:proton symporter activity	0.026%		0.0000	0.80	0.71
GO:0046961	proton-transporting ATPase activity, rotational mechanism	0.115%		0.0000	0.83	0.71
GO:0004031	aldehyde oxidase activity	0.002%		0.0055	0.86	0.80

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0004497	monooxygenase activity	1.211%		0.0081	0.88	0.00
GO:0005125	cytokine activity	0.080%		0.0079	0.38	0.00
GO:0004252	serine-type endopeptidase activity	0.893%		0.0000	0.98	0.04
GO:0016705	oxidoreductase activity, acting on paired donors, with incorporation or reduction of molecular oxygen	1.285%		0.0040	0.88	0.41
GO:0005121	Toll binding	0.000%		0.0000	0.59	0.53
GO:0004867	serine-type endopeptidase inhibitor activity	0.121%		0.0002	0.69	0.58
GO:0005126	cytokine receptor binding	0.094%		0.0025	0.50	0.74
GO:0008083	growth factor activity	0.106%		0.0005	0.37	0.85

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0004568	chitinase activity	0.057%		0.0084	0.93	0.00
GO:0004867	serine-type endopeptidase inhibitor activity	0.121%		0.0000	1.00	0.00
GO:0005319	lipid transporter activity	0.146%		0.0076	0.94	0.00
GO:0005507	copper ion binding	0.292%		0.0093	0.99	0.00
GO:0016209	antioxidant activity	0.725%		0.0036	0.91	0.00
GO:0004601	peroxidase activity	0.427%		0.0008	0.73	0.91
GO:0080019	fatty-acyl-CoA reductase (alcohol-forming) activity	0.018%		0.0033	0.81	0.02
GO:0009922	fatty acid elongase activity	0.028%		0.0078	0.96	0.02
GO:0004300	enoyl-CoA hydratase activity	0.051%		0.0002	0.97	0.02
GO:0008061	chitin binding	0.116%		0.0004	0.99	0.04
GO:0051082	unfolded protein binding	0.424%		0.0045	0.98	0.05
GO:0030246	carbohydrate binding	0.797%		0.0027	0.98	0.05
GO:0004364	glutathione transferase activity	0.086%		0.0000	0.96	0.16
GO:0052689	carboxylic ester hydrolase activity	0.694%		0.0055	0.92	0.21
GO:0004252	serine-type endopeptidase activity	0.893%		0.0001	0.92	0.27
GO:0004497	monooxygenase activity	1.211%		0.0026	0.80	0.27
GO:0015036	disulfide oxidoreductase activity	0.326%		0.0014	0.82	0.35
GO:0016614	oxidoreductase activity, acting on CH-OH group of donors	2.116%		0.0023	0.79	0.43
GO:0034041	ABC-type sterol transporter activity	0.000%		0.0058	0.94	0.57
GO:0016620	oxidoreductase activity, acting on the aldehyde or oxo group of donors, NAD or NADP as acceptor	0.593%		0.0001	0.77	0.70

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0004181	metallocarboxypeptidase activity	0.086%		0.0081	0.90	0.00
GO:0008270	zinc ion binding	3.638%		0.0092	0.96	0.00
GO:0008324	cation transmembrane transporter activity	2.405%		0.0090	0.53	0.00
GO:0008020	G protein-coupled photoreceptor activity	0.003%		0.0078	0.91	0.01
GO:0042277	peptide binding	0.100%		0.0000	0.98	0.05
GO:0004383	guanylate cyclase activity	0.024%		0.0059	0.97	0.10
GO:0005184	neuropeptide hormone activity	0.013%		0.0003	0.87	0.25
GO:0005328	neurotransmitter:sodium symporter activity	0.001%		0.0000	0.63	0.43
GO:0015347	sodium-independent organic anion transmembrane transporter activity	0.001%		0.0048	0.60	0.44
GO:0005096	GTPase activator activity	0.247%		0.0002	0.91	0.54
GO:0022848	acetylcholine-gated cation-selective channel activity	0.024%		0.0078	0.49	0.56
GO:0008028	monocarboxylic acid transmembrane transporter activity	0.052%		0.0042	0.54	0.57
GO:0005254	chloride channel activity	0.123%		0.0073	0.51	0.63
GO:0046961	proton-transporting ATPase activity, rotational mechanism	0.115%		0.0005	0.56	0.65
GO:0022839	ion gated channel activity	0.054%		0.0022	0.63	0.65
GO:0005283	amino acid:sodium symporter activity	0.045%		0.0009	0.50	0.74
GO:0099095	ligand-gated anion channel activity	0.006%		0.0022	0.55	0.80
GO:0070006	metalloaminopeptidase activity	0.153%		0.0000	0.90	0.83
GO:0015171	amino acid transmembrane transporter activity	0.268%		0.0001	0.60	0.83

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0003697	single-stranded DNA binding	0.255%		0.0079	0.86	0.00
GO:0016538	cyclin-dependent protein serine/threonine kinase regulator activity	0.069%		0.0012	1.00	0.00
GO:0016830	carbon-carbon lyase activity	1.062%		0.0074	0.98	0.00
GO:0018685	alkane 1-monooxygenase activity	0.001%		0.0045	0.98	0.02
GO:0004035	alkaline phosphatase activity	0.021%		0.0008	0.98	0.03
GO:0004364	glutathione transferase activity	0.086%		0.0003	0.98	0.03
GO:0004467	long-chain fatty acid-CoA ligase activity	0.034%		0.0004	0.98	0.04
GO:0042393	histone binding	0.086%		0.0012	0.97	0.04
GO:0000976	transcription cis-regulatory region binding	0.373%		0.0053	0.75	0.33
GO:0003688	DNA replication origin binding	0.043%		0.0024	0.77	0.81

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0008121	ubiquinol-cytochrome-c reductase activity	0.036%		0.0094	0.66	0.00
GO:0008569	minus-end-directed microtubule motor activity	0.040%		0.0011	0.96	0.00
GO:0051959	dynein light intermediate chain binding	0.026%		0.0039	0.90	0.00
GO:0005200	structural constituent of cytoskeleton	0.076%		0.0041	0.99	0.01
GO:0016831	carboxy-lyase activity	0.619%		0.0055	0.99	0.03
GO:0004181	metallocarboxypeptidase activity	0.086%		0.0014	0.91	0.03
GO:0042277	peptide binding	0.100%		0.0000	0.99	0.04
GO:0045503	dynein light chain binding	0.003%		0.0011	0.91	0.31
GO:0005184	neuropeptide hormone activity	0.013%		0.0002	0.84	0.34
GO:0045505	dynein intermediate chain binding	0.030%		0.0039	0.90	0.36
GO:0022824	transmitter-gated ion channel activity	0.062%		0.0092	0.55	0.45
GO:0022848	acetylcholine-gated cation-selective channel activity	0.024%		0.0002	0.54	0.94

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0004601	peroxidase activity	0.427%		0.0048	0.96	0.00
GO:0008061	chitin binding	0.116%		0.0002	1.00	0.00
GO:0005355	glucose transmembrane transporter activity	0.012%		0.0021	0.80	0.01
GO:0032450	maltose alpha-glucosidase activity	0.006%		0.0004	0.93	0.02
GO:0016863	intramolecular oxidoreductase activity, transposing C=C bonds	0.050%		0.0004	0.97	0.03
GO:0016491	oxidoreductase activity	12.887%		0.0012	0.96	0.04
GO:0004252	serine-type endopeptidase activity	0.893%		0.0000	0.92	0.18
GO:0005351	carbohydrate:proton symporter activity	0.026%		0.0021	0.80	0.71

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0001227	DNA-binding transcription repressor activity, RNA polymerase II-specific	0.041%		0.0013	1.00	0.00
GO:0008157	protein phosphatase 1 binding	0.009%		0.0096	0.71	0.00
GO:0016791	phosphatase activity	1.292%		0.0076	1.00	0.00
GO:0000978	RNA polymerase II cis-regulatory region sequence-specific DNA binding	0.193%		0.0053	0.95	0.03
GO:0008134	transcription factor binding	0.111%		0.0096	0.68	0.36
GO:0046982	protein heterodimerization activity	0.275%		0.0012	0.67	0.46

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0003690	double-stranded DNA binding	0.682%		0.0065	0.87	0.00
GO:0004252	serine-type endopeptidase activity	0.893%		0.0074	0.90	0.00
GO:0005355	glucose transmembrane transporter activity	0.012%		0.0066	0.91	0.00
GO:0016538	cyclin-dependent protein serine/threonine kinase regulator activity	0.069%		0.0000	1.00	0.00
GO:0004300	enoyl-CoA hydratase activity	0.051%		0.0051	0.98	0.03
GO:0004364	glutathione transferase activity	0.086%		0.0000	0.98	0.03
GO:0015036	disulfide oxidoreductase activity	0.326%		0.0022	0.94	0.03
GO:0042393	histone binding	0.086%		0.0056	0.98	0.05
GO:0017110	nucleoside-diphosphatase activity	0.016%		0.0047	0.94	0.19
GO:0004035	alkaline phosphatase activity	0.021%		0.0005	0.94	0.20
GO:0003756	protein disulfide isomerase activity	0.028%		0.0038	0.86	0.31
GO:0004601	peroxidase activity	0.427%		0.0015	0.94	0.32
GO:0003697	single-stranded DNA binding	0.255%		0.0028	0.88	0.35
GO:0000976	transcription cis-regulatory region binding	0.373%		0.0071	0.82	0.36
GO:0016863	intramolecular oxidoreductase activity, transposing C=C bonds	0.050%		0.0019	0.90	0.69
GO:0005351	carbohydrate:proton symporter activity	0.026%		0.0066	0.91	0.71
GO:0003688	DNA replication origin binding	0.043%		0.0010	0.83	0.81

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0003779	actin binding	0.467%		0.0087	0.80	0.00
GO:0004016	adenylate cyclase activity	0.077%		0.0054	0.96	0.00
GO:0005085	guanyl-nucleotide exchange factor activity	0.309%		0.0057	0.88	0.00
GO:0008569	minus-end-directed microtubule motor activity	0.040%		0.0012	0.98	0.00
GO:0015347	sodium-independent organic anion transmembrane transporter activity	0.001%		0.0057	0.72	0.00
GO:0035591	signaling adaptor activity	0.023%		0.0006	1.00	0.00
GO:0042165	neurotransmitter binding	0.001%		0.0057	0.98	0.03
GO:0031402	sodium ion binding	0.009%		0.0021	0.93	0.04
GO:0009528	G protein-coupled neurotransmitter receptor activity	0.007%		0.0006	0.84	0.14
GO:0005509	calcium ion binding	1.117%		0.0010	0.95	0.21
GO:0099094	ligand-gated cation channel activity	0.080%		0.0050	0.58	0.36
GO:0045503	dynein light chain binding	0.003%		0.0000	0.85	0.37
GO:0005328	neurotransmitter:sodium symporter activity	0.001%		0.0025	0.70	0.40
GO:0045296	cadherin binding	0.018%		0.0052	0.83	0.42
GO:0051959	dynein light intermediate chain binding	0.026%		0.0052	0.83	0.43
GO:0045505	dynein intermediate chain binding	0.030%		0.0052	0.83	0.44
GO:0001664	G protein-coupled receptor binding	0.082%		0.0018	0.79	0.48
GO:0008020	G protein-coupled photoreceptor activity	0.003%		0.0001	0.88	0.48
GO:0016917	GABA receptor activity	0.027%		0.0004	0.90	0.48
GO:0005516	calmodulin binding	0.135%		0.0027	0.81	0.49
GO:0046982	protein heterodimerization activity	0.275%		0.0062	0.81	0.52
GO:0046961	proton-transporting ATPase activity, rotational mechanism	0.115%		0.0006	0.64	0.54

n.

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0004467	long-chain fatty acid-CoA ligase activity	0.034%		0.0039	0.93	0.00
GO:0042393	histone binding	0.086%		0.0066	0.96	0.00
GO:0004364	glutathione transferase activity	0.086%		0.0037	0.90	0.02
GO:0003730	mRNA 3'-UTR binding	0.026%		0.0001	0.91	0.04
GO:0005515	protein binding	4.557%		0.0047	0.94	0.05
GO:0008146	sulfotransferase activity	0.121%		0.0024	0.90	0.17
GO:0004017	adenylate kinase activity	0.059%		0.0039	0.85	0.20
GO:0003677	DNA binding	11.354%		0.0038	0.88	0.26

p.

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0008307	structural constituent of muscle	0.004%		0.0025	0.96	0.00
GO:0015662	P-type ion transporter activity	0.107%		0.0065	0.48	0.00
GO:0016776	phosphotransferase activity, phosphate group as acceptor	0.361%		0.0076	0.99	0.00
GO:0016917	GABA receptor activity	0.027%		0.0011	0.96	0.00
GO:0005184	neuropeptide hormone activity	0.013%		0.0005	0.97	0.01
GO:0004553	hydrolase activity, hydrolyzing O-glycosyl compounds	1.613%		0.0033	0.97	0.03
GO:0042277	peptide binding	0.100%		0.0001	1.00	0.03
GO:0070006	metalloaminopeptidase activity	0.153%		0.0000	0.97	0.24
GO:0015217	ADP transmembrane transporter activity	0.020%		0.0051	0.57	0.43
GO:0008020	G protein-coupled photoreceptor activity	0.003%		0.0000	0.93	0.46
GO:0003735	structural constituent of ribosome	2.348%		0.0001	0.96	0.50
GO:0015252	proton channel activity	0.315%		0.0051	0.49	0.52
GO:0099095	ligand-gated anion channel activity	0.006%		0.0008	0.60	0.58
GO:0005283	amino acid:sodium symporter activity	0.045%		0.0034	0.44	0.61
GO:0008137	NADH dehydrogenase (ubiquinone) activity	0.260%		0.0021	0.45	0.61
GO:0044769	ATPase activity, coupled to transmembrane movement of ions, rotational mechanism	0.115%		0.0012	0.52	0.62
GO:0046961	proton-transporting ATPase activity, rotational mechanism	0.115%		0.0003	0.44	1.00

q.

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0004252	serine-type endopeptidase activity	0.893%		0.0096	0.90	0.00
GO:0008307	structural constituent of muscle	0.004%		0.0039	0.98	0.00
GO:0015662	P-type ion transporter activity	0.107%		0.0040	0.55	0.00
GO:0016917	GABA receptor activity	0.027%		0.0073	1.00	0.00
GO:0042277	peptide binding	0.100%		0.0018	1.00	0.00
GO:0016776	phosphotransferase activity, phosphate group as acceptor	0.361%		0.0037	0.96	0.03
GO:0016616	oxidoreductase activity, acting on the CH-OH group of donors, NAD or NADP as acceptor	1.855%		0.0061	0.85	0.04
GO:0008061	chitin binding	0.116%		0.0001	1.00	0.04
GO:0004568	chitinase activity	0.057%		0.0058	0.92	0.21
GO:0008483	transaminase activity	0.728%		0.0036	0.96	0.23
GO:0008010	structural constituent of chitin-based larval cuticle	0.002%		0.0000	0.98	0.33
GO:0016645	oxidoreductase activity, acting on the CH-NH group of donors	0.350%		0.0028	0.86	0.37
GO:0005313	L-glutamate transmembrane transporter activity	0.004%		0.0028	0.65	0.39
GO:0009055	electron transfer activity	1.194%		0.0011	0.77	0.42
GO:0032450	maltose alpha-glucosidase activity	0.006%		0.0028	0.91	0.47
GO:0005328	neurotransmitter:sodium symporter activity	0.001%		0.0000	0.65	0.47
GO:0015252	proton channel activity	0.315%		0.0028	0.57	0.52
GO:0022857	transmembrane transporter activity	9.001%		0.0033	0.61	0.53
GO:0070006	metalloaminopeptidase activity	0.153%		0.0000	0.91	0.54

r.

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0003729	mRNA binding	0.276%		0.0056	0.78	0.00
GO:0008186	ATP-dependent activity, acting on RNA	0.334%		0.0000	0.92	0.00
GO:0140097	catalytic activity, acting on DNA	2.626%		0.0028	0.79	0.00
GO:0005139	nuclear localization sequence binding	0.010%		0.0000	0.96	0.04
GO:0051082	unfolded protein binding	0.424%		0.0028	0.87	0.05
GO:0003697	single-stranded DNA binding	0.255%		0.0035	0.81	0.22
GO:0003677	DNA binding	11.354%		0.0048	0.75	0.33
GO:0008296	3'-5'-exodeoxyribonuclease activity	0.010%		0.0031	0.78	0.39
GO:0030515	snoRNA binding	0.048%		0.0000	0.80	0.40
GO:0042393	histone binding	0.086%		0.0005	0.88	0.47
GO:0003723	RNA binding	5.630%		0.0023	0.76	0.53
GO:0003724	RNA helicase activity	0.000%		0.0000	0.73	0.54
GO:0008297	single-stranded DNA exodeoxyribonuclease activity	0.006%		0.0011	0.78	0.70

Panel s. table

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0008061	chitin binding	0.116%		0.0000	1.00	0.00
GO:0008307	structural constituent of muscle	0.004%		0.0013	0.96	0.00
GO:0008331	high voltage-gated calcium channel activity	0.001%		0.0081	0.93	0.00
GO:0008483	transaminase activity	0.728%		0.0100	0.92	0.00
GO:0004300	enoyl-CoA hydratase activity	0.051%		0.0039	0.97	0.03
GO:0004568	chitinase activity	0.057%		0.0020	0.92	0.03
GO:0016861	intramolecular oxidoreductase activity, interconverting aldoses and ketoses	0.309%		0.0081	0.96	0.03
GO:0004601	peroxidase activity	0.427%		0.0042	0.86	0.03
GO:0016879	ligase activity, forming carbon-nitrogen bonds	1.271%		0.0015	0.90	0.04
GO:0004364	glutathione transferase activity	0.086%		0.0000	0.93	0.20
GO:0005355	glucose transmembrane transporter activity	0.012%		0.0044	0.90	0.22
GO:0016746	acyltransferase activity	3.062%		0.0079	0.91	0.29
GO:0003995	acyl-CoA dehydrogenase activity	0.152%		0.0001	0.87	0.30
GO:0016646	oxidoreductase activity, acting on the CH-NH group of donors, NAD or NADP as acceptor	0.269%		0.0039	0.87	0.31
GO:0008010	structural constituent of chitin-based larval cuticle	0.002%		0.0006	0.96	0.33
GO:0016616	oxidoreductase activity, acting on the CH-OH group of donors, NAD or NADP as acceptor	1.855%		0.0004	0.85	0.38

Panel t. table

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0008028	monocarboxylic acid transmembrane transporter activity	0.052%		0.0014	1.00	0.00
GO:0008569	minus-end-directed microtubule motor activity	0.040%		0.0023	1.00	0.00
GO:0016811	hydrolase activity, acting on carbon-nitrogen (but not peptide) bonds, in linear amides	0.737%		0.0014	1.00	0.00
GO:0051959	dynein light intermediate chain binding	0.026%		0.0080	0.80	0.00
GO:0044183	protein folding chaperone	0.019%		0.0000	1.00	0.01
GO:0005200	structural constituent of cytoskeleton	0.076%		0.0008	0.91	0.01
GO:0004812	aminoacyl-tRNA ligase activity	0.895%		0.0006	1.00	0.03
GO:0003743	translation initiation factor activity	0.394%		0.0001	0.84	0.04
GO:0008020	G protein-coupled photoreceptor activity	0.003%		0.0000	0.94	0.16
GO:0045503	dynein light chain binding	0.003%		0.0037	0.82	0.31
GO:0003723	RNA binding	5.630%		0.0001	0.89	0.32
GO:0005184	neuropeptide hormone activity	0.013%		0.0006	0.76	0.34
GO:0045505	dynein intermediate chain binding	0.030%		0.0080	0.80	0.36
GO:0051082	unfolded protein binding	0.424%		0.0000	0.77	0.43
GO:0003735	structural constituent of ribosome	2.348%		0.0001	0.92	0.63

Panel u. table

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0003954	NADH dehydrogenase activity	0.374%		0.0094	0.76	0.00
GO:0015171	amino acid transmembrane transporter activity	0.268%		0.0079	0.68	0.00
GO:0042277	peptide binding	0.100%		0.0001	1.00	0.00
GO:0044183	protein folding chaperone	0.019%		0.0043	0.99	0.01
GO:0005200	structural constituent of cytoskeleton	0.076%		0.0076	0.99	0.01
GO:0052745	inositol phosphate phosphatase activity	0.046%		0.0067	0.89	0.02
GO:0008236	serine-type peptidase activity	1.328%		0.0051	0.75	0.21
GO:0004252	serine-type endopeptidase activity	0.892%		0.0001	0.76	0.94
GO:0005328	neurotransmitter:sodium symporter activity	0.001%		0.0064	0.69	0.25
GO:0016624	oxidoreductase activity, acting on the aldehyde or oxo group of donors, disulfide as acceptor	0.123%		0.0004	0.82	0.29
GO:0016646	oxidoreductase activity, acting on the CH-NH group of donors, NAD or NADP as acceptor	0.269%		0.0067	0.81	0.31
GO:0009055	electron transfer activity	1.194%		0.0071	0.69	0.36
GO:0044769	ATPase activity, coupled to transmembrane movement of ions, rotational mechanism	0.115%		0.0015	0.64	0.37
GO:0046961	proton-transporting ATPase activity, rotational mechanism	0.115%		0.0001	0.57	1.00
GO:0016616	oxidoreductase activity, acting on the CH-OH group of donors, NAD or NADP as acceptor	1.855%		0.0053	0.78	0.42

Panel v. table

Term ID	Name	Frequency	Pin?	Value	Uniqueness	Dispensability
GO:0005126	cytokine receptor binding	0.094%		0.0096	1.00	0.00
GO:0005201	extracellular matrix structural constituent	0.037%		0.0070	1.00	0.00
GO:0015036	disulfide oxidoreductase activity	0.326%		0.0096	0.95	0.00
GO:0015318	inorganic molecular entity transmembrane transporter activity	2.924%		0.0062	0.84	0.00
GO:0031386	protein tag	0.011%		0.0070	1.00	0.00
GO:0016209	antioxidant activity	0.725%		0.0019	0.99	0.01
GO:0004467	long-chain fatty acid-CoA ligase activity	0.034%		0.0010	0.98	0.02
GO:0018685	alkane 1-monooxygenase activity	0.001%		0.0048	0.95	0.21
GO:0005355	glucose transmembrane transporter activity	0.012%		0.0009	0.78	0.35
GO:0005351	carbohydrate:proton symporter activity	0.026%		0.0009	0.77	0.71

Figure S6: Revigo Scatter and Table Views of the GO terms over-represented in all comparisons (R vs S, R vs C and C vs S) for data from Nigeria, Niger, Chad and Cameroon.

Figure S7: Validation of RNA-seq results using qRT-PCR analysis: correlation between RNA-seq fold changes and those obtained from qRT-PCR for 12 metabolic resistance genes.

a.

Population	n	S	Syn	Nonsyn	h	H_d	π (k)	D (Tajima)	D* (Fu and Li)
Nigeria	4	2	2	0	2	0.667	0.00090	1.893^{ns}	1.893^{ns}
Niger	4	42	36	6	3	0.833	0.01848	2.006^{ns}	2.006^{ns}
Chad	4	0	0	0	1	0.00	0.0000	0.000	0.000
Cameroon	4	23	20	3	2	0.500	0.0078	-0.857^{ns}	0.857^{ns}
Ngoussou	4	1	1	0	2	0.500	0.00034	-0.612^{ns}	-0.612^{ns}
All	20	75	65	12	10	0.921	0.01648	0.501^{ns}	1.298^{ns}

b.

Hap #	Freq.	Sequences
Hap_1:	2	>Nigeria2 >Nigeria3
Hap_2:	2	>Nigeria1 >Nigeria7
Hap_3:	1	>Niger1
Hap_4:	1	>Niger2
Hap_5:	2	>Niger3 >Niger4
Hap_6:	4	>Chad1 >Chad2 >Chad3 >Chad6
Hap_7:	3	>Cameroon1 >Cameroon2 >Cameroon3
Hap_8:	1	>Cameroon4
Hap_9:	3	>Ngoussou2 >Ngoussou4 >Ngoussou6
Hap_10:	1	>Ngoussou3

Figure S8: Polymorphism analysis of full length cDNA of *CYP6Z2*. a. Summary statistics for polymorphisms of *CYP6Z2* cDNA across the Sahel countries. n = number of sequences (n); S, number of polymorphic sites; h, haplotype; H_d, haplotype diversity Syn, Synonymous mutations; Nonsyn, Non-synonymous mutations; π, nucleotide diversity (k= mean number of nucleotide differences); Tajima's D and Fu and Li's D statistics, ns, not significant. **b.** *CYP6Z2* cDNA haplotypes and its frequencies, **c.** a maximum likelihood phylogenetic tree of *CYP6Z2* cDNA sequences.

Figure S9: Comparative alignment of the *GSTe2* 5'-UTR fragments from the various haplotypes, showing putative transcription factors binding sites (blue boxes), and polymorphic positions (in red and highlighted in pink). The arthropod Inr consensus sequence is in green box, the transcription start site indicated with red arrow, and the 3'-UTR of *GSTe1* and 5'-UTR of *GSTe2* indicated with dashed lines.

a.

Hap# Freq. Sequences

Hap_1: 75   NigeriaDE1 NigeriaDE2 NigeriaDE3 NigeriaDE4 NigeriaDE11
NigeriaDE15 NigeriaDE16 NigeriaDE20 NigeriaDE26 Ngoussou1
Ngoussou2 Ngoussou4 Ngoussou5 Ngoussou7 Ngoussou9 Ngoussou11 Ngoussou13
Ngoussou15 Ngoussou16 NigerAL3 NigerAL4 NigerAL5 NigerAL6 NigerAL7 NigerAL8
NigerDE1 NigerDE2 NigerDE3 NigerDE4 NigerDE5 NigerDE6 NigerDE7 NigerDE9
NigerDE11 NigerDE12 ChadAL1 ChadAL3 ChadAL4 ChadAL5 ChadAL6 ChadAL7
ChadAL9 ChadAL10 ChadDE1 ChadDE2 ChadDE3 ChadDE4 ChadDE6 ChadDE7 ChadDE9
ChadDE11 ChadDE13 ChadDE24 CameroonAL1 CameroonAL2 CameroonAL3 CameroonAL4
CameroonAL8 CameroonAL9 CameroonAL11 CameroonDE1 CameroonDE2 CameroonDE3
CameroonDE4 CameroonDE14 CameroonDE12 CameroonDE11 CameroonDE8
CameroonDE10 CameroonDE6 NigeriaAL7 NigeriaAL4 NigeriaAL8 NigeriaAL17 NigeriaAL2

Hap_2: 1   NigeriaDE29

Hap_3: 1   NigeriaAL3

Hap_4: 1   NigeriaAL5

Hap_5: 4   NigerAL1 NigerAL9 NigerAL10 NigerAL11

Hap_6: 8   NigeriaAL1 NigeriaAL6 NigeriaAL9 ChadAL2 ChadAL13
CameroonAL5 CameroonAL25 CameroonAL7

b.

```
        0000001111 2222223
        0246680116 11 23993
        927095246848941 33
Hap_1  TTGTTATTCATAGATCG
Hap_2  ..........................G.....
Hap_3  ........... C.....................
Hap_4  ..ACAG. CACCGA .AG.
Hap_5  AAA...........................
Hap_6  AAACA. .CACCGA...GC
```

c.

Figure S10: Genetic variability of the *GSTe2* 5'-UTR fragment from Nigeria, Niger, Chad, Cameroon and Ngoussou populations. a. the six haplotypes of the 5'-UTR with respective frequencies, **b.** polymorphic positions with numbers on top indicating the nucleotide position in the 351 bp fragment, **c.** maximum likelihood phylogenetic tree of the 90 sequences, showing the DDT-alive sequences with mutations forming a clade specific to its phenotype.

Figure S11: qRT-PCR validation of the overexpression of *GSTe2* and *CYP6Z2*.
The relative expression of the *GSTe2* and *CYP6Z2* in the experimental transgenic *D. melanogaster* in relation to control flies. Data shown are the mean ± standard deviation (*n* = 6).