Regionally distinct trophoblast regulate barrier function and invasion in the human placenta

Bryan Marsh[1, 2, 3, 4]**, Yan Zhou**[1, 3, 5]**, Mirhan Kapzidic**[1, 3, 5]**, Susan Fisher**[1, 3, 5*]**, and Robert Blelloch**[1,2, 3*]

[1]The Eli and Edythe Broad Center of Regeneration Medicine and Stem Cell Research, University of California, San Francisco, San Francisco, CA 94143, USA
[2]Department of Urology, University of California, San Francisco, San Francisco, CA 94143, USA
[3]Center for Reproductive Sciences, University of California, San Francisco, San Francisco, CA 94143, USA
[4]Developmental and Stem Cell Biology Graduate Program, University of California, San Francisco, San Francisco, CA 94143, USA
[5]Department of Obstetrics, Gynecology, and Reproductive Sciences, University of California, San Francisco, San Francisco, CA 94143, USA
*Correspondence: robert.blelloch@ucsf.edu; susan.fisher@ucsf.edu

Abstract

The human placenta contains two specialized regions: the villous chorion where gases and nutrients are exchanged between maternal and fetal blood, and the smooth chorion which surrounds more than 70% of the developing fetus but whose cellular composition and function is poorly understood. Here, we use single cell RNA sequencing to compare the cell types and molecular programs between these two regions in the second trimester human placenta. Each region consists of progenitor cytotrophoblasts (CTBs) and extra-villous trophoblasts (EVTs) with similar gene expression programs. While CTBs in the villous chorion differentiate into syncytiotrophoblasts, they take an alternative trajectory in the smooth chorion producing a previously unknown CTB population which we term smooth-chorion-specific CTBs (SC-CTBs). Marked by expression of region-specific cytokeratins, the SC-CTBs form a stratified epithelium above a basal layer of progenitor CTBs. They express epidermal and metabolic transcriptional programs consistent with a primary role in defense against physical stress and pathogens. Additionally, we show that SC-CTBs closely associate with EVTs and secrete factors that inhibit the migration of the EVTs. This restriction of EVT migration is in striking contrast to the villous region where EVTs migrate away from the chorion and invade deeply into the decidua. Together, these findings add an important new dimension to our understanding of CTB differentiation in these distinct regions of the human placenta. This knowledge has broad implications for studies of the development, functions, and diseases of the human placenta.

Impact Statement

Single cell RNA-sequencing of distinct regions of the human placenta identifies a smooth chorion-specific cytotrophoblast population responsible for unique functions of the smooth chorion, including acting as a barrier and restricting invasion.

Introduction

The human placenta is the first organ to develop and forms the essential bridge between maternal and fetal tissues beginning at implantation (Knöfler et al., 2019; Turco and Moffett, 2019). The placenta must develop rapidly upon conception as it performs myriad functions that remain rudimentary in the developing embryo/fetus such as nutrient and oxygen transport and protection from mechanical and pathogenic insults. The placental also performs unique functions such as modulation of maternal tolerance and hormone production (Maltepe and Fisher, 2015;

Knöfler et al., 2019; Turco and Moffett, 2019). Beginning at implantation, placental development
proceeds asymmetrically to produce two distinct regions. At the human implantation site, the
embryonic pole, trophectoderm and associated mesoderm cells give rise to the villous chorion
including the placental villi and basal plate, which are essential for the exchange of gases and
nutrients. The chorionic villi that arise from the opposite side, the abembyronic pole, undergo a
cycle of generation and degeneration eventually forming the smooth chorion, which fuses with
the amnion and constitutes the fetal membranes. (Hamilton and Boyd, 1960; Boyd and Hamilton,
1967; Benirshcke et al., 2006). Compared to the villous chorion, little effort has been made to
analyze types and functions of the CTBs that comprise the smooth chorion (Benirshcke et al.,
2006, Garrido-Gomez et al., 2017).
58
The anatomical processes associated with the formation of the fetal membranes have been well
described (Benirschke et al., 2006; Hamilton and Boyd, 1960; Boyd and Hamilton, 1967). The
first month of placental development proceeds in a uniform fashion, with the expansion and
establishment of villi surrounding the entire developing embryo. These villi have the same
complement of trophoblasts as more mature villi including diploid cytotrophoblast (CTB) and
multinucleate syncytiotrophoblast (STB) [Knöfler et al., 2019]. The villi surrounding the embryo
expand, branch, and are colonized by the fetal vasculature until 6-7 weeks of gestation, when
degeneration begins in the future smooth chorion. In this region, villi continue to recede towards
the basal plate. By the time the process is complete in the second trimester, villi are restricted
to the discoid placenta that gives rise to the CTBs that invade the decidua (Hamilton and Boyd,
1960; Knöfler et al., 2019). As the developing fetus expands into the uterine cavity, the fetal
membranes become juxtaposed to the decidual wall away from the discoid placenta, eventually
fusing with these maternal cells at approximately 18 weeks of gestation. (Boyd and Hamilton,
1967).
73
The fetal membranes have a complex cellular makeup, which includes amniocytes,
mesenchymal stromal cells, immune cells, and CTBs (Benirschke et al., 2006). The CTBs of the
smooth chorion persist until term as an epithelial like structure. Non-functional "ghost villi" can
occasionally be found in the smooth chorion, but lack STBs and fetal vasculature, and thus
cannot function in a manner comparable to the chorionic villi (Benirschke et al., 2006).
Furthermore, in contrast to the villous chorion where CTBs invade and remodel the maternal
arteries, the CTBs of the smooth chorion do not invade the adjacent decidua and the maternal
blood vessels it contains (Genbacev et al., 2015). Thus, the function of these CTBs remains
unclear.
83
Several pieces of evidence suggest that the smooth chorion is not simply a vestigial structure.
First, an intact CTB layer, which contains proliferating cells persists through the final 20 weeks
of gestation (Yeh et al., 1989; Benirschke et al., 2006, 1979; Garrido-Gomez et al., 2017).
Second, the histological heterogeneity among smooth chorion CTBs suggests functional
distinctions. Yeh et al. (1989) characterize two distinct populations of vacuolated and
eosinophilic CTBs. Vacuolated CTBs were positive for placental lactogen and placental alkaline
phosphatase, while the eosinophilic subpopulations was not. Both populations were rich in
keratin and neither had the known characteristics of villous CTBs. Bou-Resli et al. (1981) also

note high levels of variation among CTBs in the smooth chorion and the existence of a
vacuolated population. A more molecular characterization was carried out by Garrido-Gomez et
al. (2017), which demonstrated heterogeneity of ITGA4 and HLA-G expression, markers
previously associated with stemness and invasion, respectively (Genbacev et al., 2016,
McMaster et al., 1995). This study also uncovered an expansion of the smooth chorion in cases
of severe pre-eclampsia, along with a disease-specific gene expression pattern. In sum, these
results suggest the smooth chorion CTBs are a heterogeneous and dynamic collection of cells
with important functions in development and disease.
100
Single cell RNA-sequencing (scRNA-seq) has emerged as the standard for transcriptional
characterization of complex organs. This methodology was previously applied to the placenta,
but with a focus on the maternal-fetal interface, specifically the chorionic villi and basal plate (Liu
et al., 2018; Suryawanshi et al. 2018; Vento-Tormo et al., 2018). Recently, scRNA-seq was used
to profile the fetal membranes at term. However, in this study CTBs were not identified in the
smooth chorion, possibly due to apoptosis and/or changes in the membrane at term (Pique-Regi
et al., 2019; Yuan et al., 2006; Yuan et al., 2008; Yuan et al., 2009). To fill this information gap,
we applied scRNA-seq to matching samples of cells isolated from the villous and smooth chorion
regions of human samples from late in the second trimester. We used scRNA-seq to compare
the composition and developmental trajectories of CTBs in the villous and smooth chorion. The
data were validated and extended with functional studies. These results identified a novel
smooth chorion-specific CTB population important for establishment of a protective barrier and
the suppression of trophoblast invasion. In addition, these data represent a resource of CTB
types, proportions, and gene expression at mid-gestation against which age related and
pathogenic alterations can be measured.
116

Results
The transcriptional landscape of the villous and smooth chorion at mid-gestation.
To understand the cellular composition of the smooth chorion, we isolated and profiled cells from
both the villous (VC) and the smooth chorion (SC) regions of four second trimester human
placentas spanning gestational weeks 18 to 24 (GW18-24) using single cell RNA-sequencing
(Figure 1a). We chose to analyze second trimester samples because the maturation of the
smooth chorion is complete but the inflammation and apoptosis associated with membrane
rupture and parturition is absent (Benirshcke et al., 2006; Yuan et al., 2006; Yuan et al., 2008;
Yuan et al., 2009; Figure 1a). For the purpose of establishing internal and patient controls
purposes we isolated SC and VC cells from each human placental sample. CTBs were enriched
over stromal and immune cells by using our previously published method (Garrido-Gomez et al.
2017). The transcriptomes of the resulting cells were captured using the 10x Genomics scRNA-
seq platform.
130
Each of the eight datasets (GW17.6, 18.2, 23.0, 24.0; VC and SC) were captured independently,
then integrated computationally (Figure 1b; Stuart et al., 2019). We classified the cells of the
integrated dataset into broad cell type clusters according to functional identities and annotated
each by expression of canonical markers. CTBs were annotated as $KRT7+$, $HLA-G^-$; extravillous
trophoblasts (EVTs) as $KRT7+$, $HLA-G+$; immune cells as $CD45+$, $VIM-$; stromal cells as $VIM+$,

CD45-; and the uterine epithelium by expression of *EPCAM+*, *MUC1+* and *MUC16+* (Figure 1c;
Lee et al., 2016; McMaster et al., 1995; Vento-Tormo et al., 2018). Cells expressing exclusive
markers of disparate cell types (co-expression of *KRT7, HLA-G, HLA-A, VIM, ACTA2*) were
labelled as doublets and excluded from further analysis. The complete dataset used for further
analysis contained 50,496 cells that passed quality control (between 500 and 6500 unique
genes, fewer than 15% mitochondrial reads, doublets removed) [Figure 1-S1a; McGinnis et al.,
2019]. Cells originating from each placental region (VC - 25,367 and SC – 25,129) and each
placental sample (7,181-17,705 cells per sample) were well represented (Figure 1b; Figure S1b
and c). The number of cells in each broad cell type cluster demonstrated enrichment for CTB,
which represent more than 60% of the cells in the integrated dataset (Figure 1-S1d).
146

Figure 1 – The transcriptional landscape of the villous and smooth chorion at mid-gestation. (a) Left - Schematic of the placenta at mid-gestation, highlighting the regions sampled, together with the methods used for cell isolation and characterization. Right – schematic of the cell types and their organization in each region. **(b)** UMAP projections of integrated samples, shown by region of origin (Left – VC, Right – SC), and colored according to broad cell type clusters. **(c)** Heatmap of the transcript expression of select cell identity markers across broad cell type clusters and regions. Values are scaled expression across the clusters of each region independently.

While the enrichment protocol was designed to enrich for CTBs cells, several immune and
stromal subtypes were recovered, providing insight into the composition of these cells from each
region. The immune and stromal cells were subset and re-clustered, allowing the annotation of
subtypes within each group (Figure 1-S2a and b; Figure 1-S3a and b). The majority, if not all,
immune cells were of maternal origin (Figure 1-S2c). Almost two-fold more immune cells were
recovered from the VC than the SC, although it cannot be ruled out that this change in proportion
is not an artifact of dissection or enrichment protocol (Figure 1-S2d). Comparing the immune cell
types identified in each region revealed a higher proportion of macrophages in the VC as
compared to the SC, which contained a greater proportion of NK/T cells (Figure 1-S2e). These
differences may reflect the fact that SC samples contain a higher proportion of decidua and its

resident immune cells than their VC counterparts. Additionally, this result may suggest
differences in the maternal immune cells that interact with the two compartments.
159
While few stromal cells were isolated in the preparations, sub-clustering still revealed a
differential composition of fetal stromal cells between the VC and SC (Figure 1-S3a and b). The
majority of stromal cells recovered originated from the SC (2,941 compared to 942 from VC).
These cells included lymphatic endothelium (Pique-Regi et al., 2019) and two largely SC-specific
mesenchymal cell populations of fetal origin, Mesenchyme 1 and Mesenchyme 3 (Figure 1-S3c-
e). These two clusters are marked by elevated expression of *EGFL6*, *DLK1*, and uniquely by
expression of *COL11A1*, which is observed only in the SC (Figure 1-S3 b and f). Interestingly,
several canonical CTB support factors including HGF, WNT2, and RSPO3, were expressed in
fetal stromal populations in both placental regions, suggesting shared requirements for WNT
and MET signaling (Figure 1-S3g). Taken together these data demonstrate the identification of
broad classes of CTBs, immune, and support cells from both the VC and SC regions of the
human placenta.
172
Identification of a smooth chorion-specific cytotrophoblast population.
CTBs are the fetal cells that perform the specialized functions of the VC, and are required for
normal fetal growth and development (Maltepe and Fisher, 2015; Turco and Moffett, 2019;
Knöfler et al., 2019). To better understand the composition of CTBs in the SC versus the VC, we
sub-clustered this population (*KRT7+*, *VIM-*, *CD45-*, *MUC1-*). The CTB subset is comprised of
29,668 cells with similar representation and cell quality control metrics across all eight samples
(Figure 2-S1a, b, and c). This analysis identified 13 clusters including several CTB, EVT, and
STB subtypes (Figure 2a).
181
Broad classes of trophoblast were annotated by established markers (STBs - *CGA*, *CYP19A1*,
CSH1, *CSH2*; EVTs - *HLA-G, DIO2*; CTBs - *PAGE4*, *PEG10,* and no expression of EVT and
STB markers) [Figure 2b and Figure 2-S2a; McMaster et al., 1995; Lee et al. 2016, Suryawanshi
et al. 2018, Liu et al. 2018]. Two cell clusters showed high expression of canonical phasic
transcripts, including *MKI67*, with the S-phase cluster denoted by expression of *PCNA* and the
G2/M-phase cluster by expression of *TOP2A* (Figure 2-S2b; Tirosh et al. 2015). Both populations
share gene expression with all clusters of CTBs, and therefore, were identified as actively cycling
(Figure 2-S2a). No STB or EVT markers were identified in the cycling clusters as was expected
due to the requirement for cell cycle exit upon terminal differentiation to these lineages (Lu et al.
2017, Genbacev et al., 1997).
192
CTBs separated into four clusters, CTB 1-4. CTB 1 cells highly expressed *PAGE4*, *PEG10*, and
CDH1 (Figure 2b, Figure 2-S2a, Figure 2-S3a), which have been shown to be canonical markers
of villous CTB (Lee et al. 2016, Suryawanshi et al. 2018). Overall, CTB 2-4 were more
transcriptionally similar to each other than to CTB 1, indicating a transcriptional program that
was distinct from canonical villous CTBs (Figure 2-S2a and c). CTB 2-4 existed along a gradient
of gene expression changes suggestive of various stages of a common differentiation pathway.
However, each population expressed distinct transcripts corresponding to important proposed
functions of the smooth chorion. CTB 2 cells highly expressed *CLU*, *CFD*, and *IFIT3*, suggesting

roles in responding to bacterial or viral infection through the innate arm of the immune system (Thurman and Hollers., 2006; Liu et al., 2011). CTB 3 cells upregulated *EGLN3* and *SLC2A3*, indicating a HIF-mediated response to hypoxia and a switch toward glucose metabolism, likely as an adaptation to the decreased oxygen levels in the largely avascular smooth chorion region (del Peso et al. 2003, Maxwell et al., 1997). Finally, CTB 4 specifically expressed several cytokeratins — *KRT6A*, *KRT17*, and *KRT14*, found in many epithelial barrier tissues and important for maintenance of integrity in response to mechanical stressors (Karantza, 2011; Figure 2b, Figure 2-S2a). These results establish a previously underappreciated transcriptional diversity of CTB subpopulations.

Quantification of cells showed a strong regional bias in the number and proportion contributing to each CTB cluster (Figure 2c and d). In the VC samples, 53.8% of CTBs clustered in CTB 1 compared to 11.1% in the SC (Supplementary Table 1). In contrast, the SC had a much larger proportion of the CTB 2-4 clusters. In this region, 71.3% of CTBs were nearly equally distributed among CTB 2 (25.6%), CTB 3 (24.9%), and CTB 4 (20.8%). In the VC, only 24.1% of CTBs were found in the same clusters, with the majority in CTB 2 (16.1%). The contribution to cycling clusters was consistent across regions: 22.1% and 17.5% of CTBs in the VC and SC, respectively. The relative proportions of CTB 1-4 in the VC and SC were consistent across individual samples indicating that this difference was biological rather than driven by sample variability or an artifact of sample integration (Figure 2-S1b-c).

Next, we immunolocalized the protein products of genes that distinguished the subpopulations. At the mRNA level, *PAGE4* expression was highest in CTB 1 and decreased across CTB 2-4 (Figure 2b and e). In the VC, the CTB monolayer between the fetal mesenchymal villous core and the overlying STB layer showed strong PAGE4 immunoreactivity, which diminished upon differentiation to EVT (Figure 2f - left). PAGE4 mRNA and protein expression matched that of known villous CTB marker CDH1 (Figure 2-S3a and b; Zhou et al., 1997). In the SC, the PAGE4 signal was strong in the epithelial layer directly adjacent to the fetal mesenchyme and then decreased in cells distant from the basal layer, again matching CDH1 (Figure 2f - right, Figure 2-S3b - right). Both RNA expression and protein localization were consistent with CTB 1 cells existing on both sides of the placenta and occupying a similar niche.

Staining for KRT6, a marker highly enriched in CTB 4 cells (Figure 2g) showed a strikingly different result. Cells occupying the upper layers of SC epithelium showed a strong KRT6 signal, a pattern opposite to CDH1 (Figure 2h and i - right). KRT6 was absent from either the floating or anchoring villi of the VC (Figure 2h and i - left). Rare decidual resident KRT6 positive CTBs were also identified in the VC region. (Figure 2-S3c, Figure 6-S1 - top). KRT6 isoforms, KRT6B and KRT6C, were not expressed, confirming KRT6A transcript and protein as highly specific markers of a CTB population found only in the smooth chorion (Figure 2-S3d). These data describe a novel subpopulation of CTBs unique to the SC, which going forward we term CTB 4 or SC-CTBs for smooth chorion-specific CTBs.

Figure 2 – Identification of a smooth chorion-specific cytotrophoblast. (a) UMAP projection of subclustered trophoblasts (n=29,668). Colors correspond to the clusters at the right. **(b)** Dot plot showing average expression and percent of cells in each cluster as identified by the marker genes listed on the x-axis. The clusters are listed on the y-axis. **(c)** UMAP projection of subclustered trophoblasts from the VC (Left) or the SC (Right). Clusters and colors are the same as in panel a. **(d)** Quantification of the number of cells in each trophoblast cluster from each region. Cells from the VC are shown in black. Cells from the SC are shown in blue. **(e)** Violin plot of *PAGE4* transcript expression across all trophoblast clusters. **(f)** Immunofluorescence co-localization of PAGE4 with pan-cytokeratin (marker of all trophoblast) in the VC (Left) or SC (Right). **(g)** Violin plot of *KRT6A* transcript expression across all trophoblast clusters. **(h)** Immunofluorescence co-localization of KRT6 with pan-cytokeratin (marker of all trophoblast) in the VC (Left) or SC (Right). **(i)** Immunofluorescence co-localization of CDH1 and KRT6 in the VC (Left) or SC (Middle). High magnification inset is denoted by the white box (Right). For all images nuclei were visualized by DAPI stain; Scale bar = 100μm. Abbreviations: AV = Anchoring Villi; FV = Floating Villi; SC = Smooth Chorion epithelium; Amn. = Amnion; Dec. = Decidua.

A common CTB progenitor gives rise to STBs in the VC and SC-CTBs in the SC.

Next, we investigated the developmental origin of the SC-CTBs. We performed RNA velocity analysis to predict the relationships between cells based on the proportion of exonic and intronic reads. These predictions are shown as vectors representing both the magnitude (predicted rate) and the direction of differentiation (Bergen et al. 2020). We first asked whether RNA velocity could recapitulate the well-established differentiation trajectories of trophoblasts in the VC (Knöfler et al., 2019; Turco and Moffett, 2019; Vento-Tormo et al. 2018). In accordance with previous results, RNA velocity projections identified CTB 1 as the root for three differentiation trajectories: self-renewal, differentiation to STBs, and differentiation to EVTs (Figure 3a). Cells at the boundary of the CTB 1 cluster showed differentiation vectors of high magnitude toward STB Precursors and upregulated canonical drivers of STB differentiation and fusion (*ERVW-1* and *ERVFRD-1*). These cells also expressed transcription factors (*GCM1* and *HOPX*) and hormones (*CSH1*) necessary for STB function (Figure 3-S1a; Baczyk et al., 2009; Mi et al., 2000; Blaise et al., 2003; Yabe et al., 2016).

Figure 3 – A common CTB progenitor gives rise to STBs in the VC and SC-CTBs in the SC. RNA velocity vector projections overlaid on to UMAP projections for trophoblast cells isolated from the **(a)** VC and **(b)** SC. Arrows denote direction and magnitude is represented by line thickness. **(c)** Pseudotime reconstruction of SC derived CTB 1-4 clusters from the scVelo dynamical model of latent time. Each column represents one cell. Cells at the left are clustered in CTB 1 and progress through CTB 2, 3, and 4 along the x-axis. Select genes that were the major drivers of the pseudotime alignment are shown on the y-axis. Expression ranged from dark blue (lowest) to yellow (highest). **(d)** Violin plots of select factors from (c) demonstrated shared or region-specific expression for genes associated with the CTB 4 differentiation trajectory.

In the SC, CTB 1 cells once again were identified as the root for differentiation. However, CTB 1 cells showed strong directionality and magnitude toward CTB 2-4 (Figure 3b). All cells in CTB 1-4 clusters displayed uniform directionality indicating a robust differentiation trajectory ending at CTB 4. High levels of transcriptional similarity between CTB 2-4 compared to CTB 1 suggested CTB 2 and 3 are intermediate states between CTB 1 and CTB 4 (Figure 2-S2c). Mitotic KRT6+ cells were identified, suggesting that differentiation to SC-CTB does not require cell cycle exit (Figure 3-S1b). In contrast to the VC, we observed no velocity vectors with

directionality toward the STB lineage from the CTB clusters in the SC samples. Further, the
smaller number of STB precursors (460 cells) and STBs (14 cells) exhibited reduced expression
of STB canonical markers such as ERVFRD-1 and GCM1, and notably, a near absence of
ERVW-1 (Figure 3-S1a). These cells may be associated with ghost villi (Benirschke et al., 2006).
In sum, these data show differential developmental trajectories for the CTB 1 cells in the SC and
VC, with the former largely giving rise to SC-CTBs and the latter to STBs.
272
To identify the genes that were correlated with progression from CTB 1-4 in the SC, we used the
velocity vector predictions to construct a pseudotemporal model of differentiation. All the cells in
these clusters were plotted in one dimension from the least to the most differentiated according
the pseudotime model (Figure 3c). Genes that were highly expressed at the start of the
pseudotemporal differentiation included pan-trophoblast factors such as *EGFR*, which was
expressed throughout all four CTB populations in both the VC and SC. Progression along the
pseudotime trajectory identified regulators of cell fate and function, including the transcription
factor *KLF4* and extracellular matrix (ECM) components *COL5A1* and *LAMA3* (Figure 3c-d); all
demonstrated SC-specific expression. Elevated expression of ECM transcripts (*COL4A2*, *FN1*)
and transcription factors responsive to cell contact and mechanical stress *(HES1, YAP1)* were
coordinately upregulated, potentially highlighting the effects of the extracellular environment on
fate specification (Figure 3-S1c). These data demonstrated that SC-CTBs originate from CTB 1
progenitors common to both the VC and SC. In the SC, instead of upregulating syncytialization
factors such as *GCM1* and *ERVFRD-1*, CTB 1 progenitors upregulate transcription factors such
as *KLF4*, *YAP1*, and *HES1*, which drive an epithelial cell fate in other contexts (Segre et al.,
1999, Harvey et al., 2013; Rock et al., 2011).
289

SC-CTBs express a distinct epidermal transcriptional program.
Next, we sought a better understanding of the physiological functions of the SC trophoblast
clusters. We performed gene ontology analysis as a summary of functional processes (Figure
4a, Figure 4-S1; Yu et al., 2012). We focused on the progenitor CTB 1 and terminally
differentiated SC-CTBs as they showed enrichment for strikingly different functional categories.
In CTB 1, we identified enrichment for WNT signaling, epithelial morphogenesis, and membrane
transport, categories commonly associated with progenitors (Figure 4a, Figure 4-S1). We
validated the activity of WNT signaling and the location of these cells by immunolocalization of
non-phosphorylated CTNNB1 (intact form of the protein). β-Catenin was localized to the most
basal epithelial layer nearest to the stroma in both the VC and SC regions (Figure 4b), matching
expression of the CTB 1 marker CDH1 (Figure 4-S2a). WNT signaling has an important role in
the maintenance of villous CTBs *in vivo* and in the derivation and culture of self-renewing human
trophoblast stem cells (Knöfler et al., 2019; Haider et al., 2018; Okae et al., 2018). We
investigated proliferation of CTNNB1 expressing cells in both regions using KI67 as a mitotic
marker. This revealed a similar percentage of KI67+ CTB 1 cells, suggesting similar proliferative
capacity across regions (Figure 4b and c). We next analyzed regional differences within CTB 1.
Gene ontology identified an enrichment for oxidative phosphorylation and epithelial signaling
cues in VC CTB 1 cells (Figure 4-S2b). This is in direct contrast to SC CTB 1 cells that displayed
elevated levels of hypoxia response genes (Figure 4-S2b-c). KLF4 was identified in the RNA
velocity analysis as gaining expression from CTB1-4, but also showed greater expression in

CTB1 in the SC compared to the VC (Figure 3d). In accordance with the mRNA expression data,
KLF4 protein often co-localized to CTB1 in the SC, but was only observed in rare cells in the VC
(Figure 4d). These data further support a similar location and function for CTB 1 in the SC and
VC, with transcriptional and metabolic differences that presage distinct developmental
trajectories.
315
CTB 4 ontological analysis strongly supported important roles for these cells in the formation of
a protective barrier. The greatest enrichment was for cell junction and cell substrate adhesion
genes that included numerous integrin and laminin subunits as well as junctional proteins *PATJ*,
DSP, and *JUP*. An enrichment for both skin and epidermal development correlated with
upregulation of the ECM and junctional transcripts. These categories included many cytokeratins
(*KRT6A, 7, 8, 14, 17, 18,* and *19*) and transcription factors (*KLF4, YAP1)* required for epidermal
identity and maintenance (Segre et al., 1999; Schlegelmilch et al., 2011). The organization of
the SC is reminiscent of stratified epidermal cells of the skin, with progenitors adherent to the
basal lamina and more differentiated cells progeny forming the upper layers. In many tissues,
the specific domains of cytokeratin expression correspond to stratified cell layers with different
functions. We asked if this was also the case in the SC epithelium. Cytokeratins 7, 8, and 18

Figure 4 – SC-CTBs express a distinct epidermal transcriptional program. (a) Heatmap of gene ontology analysis adjusted p-values. Dark red corresponds to the lowest p-values and white represents p-values greater than 0.0005. Ontology categories are organized by hierarchical clustering along the y-axis. Marker genes for each cluster were used as inputs for the analysis. A subset of genes in selected categories are listed at the right. Categories and genes enriched in CTB 1 or CTB 4 are in black or blue, respectively. **(b)** Representative immunofluorescence co-localization of non-phosphorylated CTNNB1 and KI67 in the VC (Left) and SC (Right). **(c)** Quantification of the percent of np-CTNNB1 cells with KI67 expression in each region in three placental samples. Each dot represents the percentage in one field of view (at least 7 per region per sample) as an estimate of mitotic cells per population. Percentages for the VC region are shown in black and the SC region in blue. **(d)** Immunofluorescence co-localization of CDH1 and KLF4 protein in the VC (Left) or SC (Right). Arrowheads denote CDH1+/KLF4+ cells. **(e)** Immunofluorescence co-localization of KRT14 and KRT6 protein in the VC (Left) or SC (Right). The outline of the Anchoring Villi (AV) is denoted by the white dashed line. For all images nuclei were visualized by DAPI stain; Scale bar = 100µm. Abbreviations: AV = Anchoring Villi; FV = Floating Villi; SC = Smooth Chorion epithelium; Amn. = Amnion; Dec. = Decidua; GV = Ghost Villi.

were expressed in all trophoblast regardless of region (Figure 4-S3a), but cytokeratin 6A, 14, and 17 displayed smooth chorion-specific expression that increased with differentiation towards CTB 4 (Figure 4-S3b). Immunofluorescence localization confirmed expression of KRT14 as specific to the CTBs in the SC and inclusive of all KRT6 expressing cells (Figure 4e). These data support a model of accumulated cytokeratin expression that begins with CTB 1 (*KRT7, KRT8, KRT18*), increases in CTB 2-3 (*KRT14, KRT17*), and culminates in CTB 4 (*KRT6A*). Together these data are consistent with a central role for the CTBs of the SC in establishing a protective epithelial barrier for the rapidly growing fetus.

Beyond forming a physical barrier, important placental functions include protection from bacterial and viral infections. We identified an enrichment for genes involved in antiviral response in SC-CTBs, and in SC cells more broadly. For example, IFITM3, a restriction factor preventing entry of viruses into cells, is highly expressed in CTBs from the SC compared to the VC (Figure 4-S4; Bailey et al., 2014; Spence et al., 2019). IFITM proteins have also been reported to inhibit syncytialization (Buchrieser et al. 2019), suggesting IFITM3 may also function to block differentiation of CTBs into STBs in the SC. Taken together, these data establish SC-CTBs as the building blocks and critical regulators of the SC barrier, responsible for both protection against physical forces and pathogen infection.

EVTs of the VC and SC regions are behaviorally distinct but transcriptionally similar.
EVTs are the invasive trophoblasts of the placenta (Knöfler et al., 2019; Turco and Moffett, 2019; Red-Horse et al., 2004). While the EVTs of the VC migrate away from the villi, invade the decidua, and replace the endothelial lining of the uterine arteries, the EVTs of the SC adhere to the decidua and do not home to the maternal vasculature (Genbacev et al., 2015). To understand the basis for these differences, we compared EVT subpopulations isolated from the VC and SC. Expression of the canonical marker of EVTs, *HLA-G*, was similarly abundant among the cells isolated from both placental regions (Figure 5a and b). A greater number (VC - 6572; SC - 5021) and larger proportion (VC – 46.83%; SC – 32.12%) of CTBs from the VC expressed HLA-G as compared to the analogous population from the SC (Figure 5c). Consistent with the transcript expression data, immunolocalization of HLA-G showed strong staining of cells in both the VC and SC. However, VC derived HLA-G positive cells were found deep in the maternal decidua. In contrast, HLA-G positive cells in the SC remained in the epithelial layer (Figure 5b).

EVTs were further divided into clusters 1-4. No clusters were specific to the VC or SC, except for the small number of putative EVT Precursors (VC – 99 cells; SC – 8 cells) [Figure 5d, Figure 3a]. Previous work has classified EVTs on the villous side of the placenta as columnar, interstitial, or endovascular based on behavior and location in the decidua (Tilburgs et al., 2015; Knöfler et al., 2019). The columnar subpopulation is believed to represent newly differentiated EVTs, which lie at the base of the columns that connect the anchoring villi to the uterine wall. Interstitial EVTs migrate through the decidua homing to maternal arteries, which they invade. Within the arterial wall, this subpopulation further differentiates to endovascular EVTs and replaces the maternal arterial endothelium (Harris et al. 2009, Red-Horse et al. 2004). We expected to capture both columnar and interstitial EVT populations, but few endovascular EVTs as the cellular preparations were largely devoid of arteries. We investigated the four clusters of EVT identified

in our analysis in the context of the VC and the SC. Both placental regions contained cells from all clusters, although distinct regional biases were evident (Supplementary Table 1). The SC region contained almost twice as many EVT 1 cells as the VC (VC – 684; SC – 1256), whereas the VC contained almost twice as many EVT 2-4 as the SC (VC – 5789; SC – 3757). We next asked which of the EVT clusters corresponded to known EVT classifications, indicating maturation state or invasive capacity. Gene ontology analysis showed an enrichment for placental development and antigen presentation categories in EVT 1 (Figure 5-S1a). Representative marker genes for these categories in EVT 1 included several lineage-specific transcription factors including *GCM1*, *PPARG*, and *CEBPB* (Figure 5-S1b; Knöfler et al., 2019, Ferreira et al., 2016). In contrast, EVT 2-4 showed enrichment for extracellular structure and matrix organization, glycosylation, and peptidase activity. EVT clusters 3-4 specifically showed increased expression for transcripts of proteases such as *HTRA1*, *MMP2*, and *MMP11* (Figure 5-S1c). Based on the GO and specific gene enrichments, EVT 1 appeared most similar to columnar EVTs, while EVT 2-4 were consistent with interstitial EVTs that gain invasive capacity. The relative enrichment for EVT 1 in the SC (VC – 10.57% of EVT; SC – 25.05% of EVT) suggested an expansion in columnar-like EVTs at the expense of interstitial EVT. Conversely, the relative enrichment for EVT 2-4 in the VC region suggested an expansion of the invasive subpopulation (Figure 5d).

As almost 75% of EVTs from the SC region were in EVT clusters 2-4, we next asked if there were differences in gene expression between interstitial EVTs isolated from each region. This analysis identified relatively few differentially expressed genes, with the vast majority identified having a log fold change less than 1 (Figure 5-S2a). The small number of differentially expressed genes with a fold change greater than 1 was typically identified across multiple EVT clusters, suggesting an overriding transcriptional program that might supersede subcluster designations. To identify genes associated with this program, we combined all EVT subclusters into one cluster for analysis (Figure 5e). Differential expression analysis revealed the hormone CSH1 and several proteases including HTRA1, HTRA4, and PAPPA to be expressed at significantly higher levels in the more invasive VC EVTs. However, of the most differentially expressed transcripts only CSH1, ITM2C, and DNASE1L3 were exclusive to the VC (Figure 5-S2b). Immunolocalization of CSH1 confirmed this result (Figure 5f). CSH1, also known as human prolactin, is a secreted hormone that signals through the receptor PRLR found on maternal cells. As such, the primary effect of CSH1 is likely non-cell autonomous, raising the possibility that the largest differences in EVTs between regions may not be in inherent invasive capacity, but rather their ability to communicate with the neighboring cells.

We next asked whether the gene expression differences between the VC and SC were smaller for EVT clusters than for all other trophoblast clusters. Calculating the Pearson Correlation coefficient across each cluster, EVT 3, EVT 4, and CTB 1 (which we have established as common to both regions) were most similar and the only clusters with a coefficient greater than 0.75 (Figure 5-S2c). Since this analysis does not account for cluster heterogeneity or size, we quantified the number of differentially expressed genes between 100 randomly selected cells from each placental region within each cluster. Across 100 permutations, the number of differentially expressed genes between regions was the lowest for all four EVT clusters (Figure

415 5g). Together, these results showed that the EVTs of the VC and SC are surprisingly similar even
416 though they have distinct behaviors *in vivo*, suggesting a cell non-autonomous, rather than cell
417 autonomous, regulation of invasion.
418

Figure 5 – EVTs of the VC and SC regions are behaviorally distinct but transcriptionally similar. (a) Expression of *HLA-G* transcript per cell projected in UMAP space for the VC (Left) and SC (Right). Expression ranged from low in light gray to high in dark red. **(b)** Immunofluorescence localization of HLA-G in the VC (Left) and SC (Right). The Anchoring Villi (AV) are outlined in white (Left). The boundaries of the smooth chorion epithelium are denoted by the white lines (Right) **(c)** Quantification of the number of *HLA-G* expressing EVT (Left) and the percent of total trophoblast that express *HLA-G* (Right) for each placental region. **(d)** UMAP projection of all trophoblast cells including the EVT clusters (Left). The percent of EVT cells in each cluster from each region (Right). **(e)** Volcano plot of the differentially expressed genes between regions for all EVTs. All genes with a log fold change greater than an absolute value of 0.25 and a p-value of less than 0.05 were plotted. Those with greater expression in VC EVT are shown in black. Those with greater expression in SC EVT are shown in blue. **(f)** Immunofluorescence localization of CSH1 in the VC Top) and SC (Bottom). Arrowheads denote CSH1 expressing cells **(g)** Violin plots of the number of differentially expressed genes between 100 cells from each placental region within each cluster (100 permutations). Clusters with less than 100 cells per region were omitted due to the small sample size. For all images nuclei were visualized by DAPI stain; Scale bar = 100µm. Abbreviations: AV = Anchoring Villi; FV = Floating Villi; SC = Smooth Chorion epithelium; Amn. = Amnion; Dec. = Decidua; STB = Syncytiotrophoblast; CC = Cell Column.

420 **CTBs of the SC inhibit EVT invasion.**
421 Given the transcriptional similarity between EVT clusters originating from the VC and SC, we
422 wondered if a non-cell autonomous program could explain their distinct migratory properties.
423 Immunofluorescence co-localization of CTB 1, SC-CTBs, and EVTs showed striking differences
424 in the relative positioning of the EVTs and CTBs in the two placental regions (Figure 6a and b).

On the VC side, which lacks SC-CTBs cells, HLA-G+ EVTs were distant from the np-CTNNB1+
CTB 1 cell population as expected (Genbacev et al., 2015). By the second trimester they had
migrated away from the cell column of the anchoring villi into the uterine stroma (Figure 6a). In
contrast, HLA-G+ EVTs in the SC were adjacent to and in physical contact with KRT6+ SC-CTBs
(Figure 6b). The interactions between SC-CTBs and EVTs were numerous and widespread
throughout the SC epithelium. Contacts between CTB 1 and EVTs in the SC were not observed
and the VC contain only rare and irregular KRT6+ cells (Figure 6-S1). This close association
suggested possible paracrine signaling events between SC-CTBs and EVTs, which might impact
each cell type.

Figure 6 – CTBs of the SC inhibit EVT invasion. (a) Immunofluorescence co-localization of np-CTNNB1 and HLA-G in the VC. **(b)** Immunofluorescence co-localization of KRT6 and HLA-G in the SC. Arrowheads denote CTB and EVT interactions. **(c)** Schematic of the transwell invasion assay (top). Cells from either placental region were plated in the upper chamber of the transwell. After 39 hours of culture, the transwell membrane was fixed and stained with a Pan-cytokeratin antibody. The projections through the membrane are denoted by black arrowheads and quantified below. Results from the VC derived cells are shown in black, and the results from the SC derived cells are shown in blue. **(d)** Results from three biological replicates from each placental region cultured with conditioned medium from either VC or SC cells. The gestational ages of the plated cells is shown adjacent to the y-axis. The gestational ages of the cells from which conditioned medium was collected are noted in the legends at the right. The results for cells cultured with VC cell conditioned medium are denoted by black dots, and the results for those cultured in SC cell conditioned medium are denoted by open squares. P-values were determined by t-test and are listed above each comparison. For all images nuclei were visualized by DAPI staining ; Scale bar = 100μm. Abbreviations: AV = Anchoring Villi; FV = Floating Villi; SC = Smooth Chorion epithelium; Dec. = Decidua.

To test this theory, we attempted to recapitulate the differential invasive properties of VC and SC
EVTs in a transwell migration assay. Briefly, these cells were enriched using the same protocol
as described for the scRNA-seq experiments. The cells were plated on Matrigel coated transwell
filters and cultured for 39 hours. Trophoblast projections that reached the underside of the filter,
a proxy for invasion, were visualized by immunostaining with a pan-cytokeratin antibody (Figure
6c, Figure 6-S2). Consistent with the differences observed *in vivo*, there was greater invasion of
VC as compared to SC trophoblast (Figure 6c). Next, we asked whether EVTs from the SC
region secreted soluble factors that inhibited invasion. The invasion assays were repeated with
VC cells cultured with conditioned medium from SC cells and vice versa. While VC conditioned
medium had no impact on either VC or SC cells, SC conditioned medium significantly reduced
migration of VC cells (Figure 6d). Neither conditioned medium impacted the density of VC or SC
cells (Figure 6-S3). Therefore, a secreted factor from SC cells inhibited the migration of VC
EVTs. Given that SC-CTBs were unique to the SC side, it is highly likely that they produced
secreted factors that inhibit the invasion of EVTs to which they are juxtaposed.

Discussion

This study profiled the developing 2^nd^ trimester placenta, providing a high-resolution molecular description of the trophoblast in the smooth chorion and placing them in context of the trophoblast from the VC of the same placenta. In doing so, we addressed long standing observations concerning apparent similarities and differences between VC and SC trophoblast, deconstruct the trophoblast populations in the SC, and ascribe functions to the cells of this understudied region. Our study characterized a novel trophoblast population distinct from both extant CTB in the VC and from EVT found in either placental region. These trophoblasts reside only in stratified epithelium of the smooth chorion, express a specific cytokeratin (KRT6), and comprise at least 25% of all trophoblast in the region. Lineage reconstruction by RNA velocity suggested the progenitors for these smooth chorion-specific CTBs are a resident proliferative progenitor trophoblast with a transcriptional profile that matches villous CTB. We identified several potential regulators of the transition from a common villous CTB-like cell to SC-CTB, including epidermal transcription factors such as KLF4 and YAP1. These molecules along with region-specific signaling events drive an epidermal transcriptional program creating the building blocks for a barrier against both physical and pathogen stressors. Finally, we identified CTB-EVT interactions occurring only in the smooth chorion and provide evidence that paracrine signaling between these populations restricts trophoblast invasion.

Since the smooth chorion is created by a degenerative process, consensus has been that these cells are remnants of this process, making the smooth chorion a vestigial structure (Benirschke et al., 2006). However, several previous observations suggested this might be an oversimplification. First, these cells were noted to be proliferative, suggesting either cellular expansion or turnover (Benirschke et al., 2006). We confirmed the presence of proliferative trophoblasts in the smooth chorion. Each cycling CTB cluster expressed markers of CTB 1-4 in both regions (Figure 2-S2a) and both CTB 1 (Figure 4b-c) and SC-CTBs (Figure 3-S1b) also expressed KI67 protein. Proliferation of both the progenitor and differentiated cells suggested a need for expansion coordinated with the growth of the developing fetus. Second, the importance of the smooth chorion was suggested by the noted heterogeneity of the trophoblasts within this region (Yeh et al., 1989; Bou-Resli et al., 1981; Garrido-Gomez et al., 2017; Benirschke et al., 2006). In agreement, our scRNA-seq results showed the smooth chorion to be a complex tissue with several trophoblast types. We identified, by differential transcription and function, at least two distinct CTB cell types, columnar and interstitial EVT, and a small number of STB Precursors. As discussed, all CTB and EVT populations either matched those found in the VC or had new region-specific functions. The only evidence of degeneration was the STB Precursor population within the smooth chorion, which lacked expression of both *ERVW-1* and *ERVFRD-1*, the fusogens necessary to form STB (Mi et al., 2000; Blaise et al., 2003; Liu et al., 2018). In sum, these studies suggested that the SC is a complex and functional tissue, not simply the remnant of villous degeneration.

We focused on CTB 4, a novel population found exclusively in the SC, in the context of the unique functions of this placental region. CTB 4 were enriched for epidermal, skin development, and antiviral gene categories suggesting concerted roles in the creation of a barrier against

external forces and pathogens. Previous studies have noted that CTBs residing in the SC have high levels of keratin expression and are associated with extensive ECM deposits rich in laminin, collagen, and fibronectin (Yeh et al., 1989; Bou-Resli et al., 1981; Benirschke et al., 2006). Among all smooth chorion CTB populations, but most notably in CTB 4, we identified the upregulation of ECM components, including *LAMA3*, *COL5A1* and *FN1*. Expression of ECM molecules in the SC is essential to protect against premature rupture of the fetal membranes, but its deposition has previously been ascribed to stromal cells in the chorion (Parry and Strauss, 1998). This result also suggested that these trophoblasts create a specific ECM environment distinct from other placental regions. This altered composition likely has wide-ranging effects on trophoblast fate, gene expression, and behavior. It will be interesting to explore the impact of advancing gestational age on the SC epithelium, especially with respect to the ECM. Presumably, cellular and structural changes precede the normal process of membrane rupture prior to delivery at term. Whether premature rupture of the membranes phenocopies these events or is a unique process is an important open question.

Pathogen defense is another important function of the placenta. Infections of the chorion and amnion (chorioamnionitis) often result in adverse outcomes for mother and fetus, primarily preterm birth (Romero et al., 2014). We identified specific expression of the antiviral gene IFITM3 in CTB 4. IFITM3 expression is particularly interesting due to the recent finding that it blocks STB fusion mediated by the endogenous retroviral elements, ERVW-1 and ERVFRD-1. Therefore, IFITM3 expression in SC-CTBs may have a dual role – restricting viral entry into the cell and blocking the formation of STBs in the SC. Its expression also provides a mechanism by which the lack of STBs in the smooth chorion is maintained after degradation of the villi is complete.

A distinct feature of the CTBs of the SC was remodeling of the cytokeratin network coordinated with differentiation from CTB 1 to SC-CTBs. The progressive expression of KRT14, KRT17, and KRT6A is reminiscent of the cytokeratin code found in the epithelial cell layers of many tissues (Karantza et al., 2011). KRT14 is recognized as a marker of all stratified epithelium including subsets of basal progenitor and stem cells in several tissues, usually alongside KRT5 (Moll et al., 1982; Nelson et al., 1983; Rock et al., 2009). While we did not identify expression of KRT5 in any trophoblast, the KRT14 expressing proliferative cells of the SC fit the profile of stratified epithelia in other locations. The roles of KRT17 and KRT6 are less clear. Most knowledge about the function of these molecules comes from the epidermis in the context of injury and disease. KRT6 and KRT17 are upregulated rapidly upon epidermal injury and are expressed through the repair phase, each contributing specific functions. (Takahashi et al., 1998; McGowan and Coulombe, 1998). KRT17 promotes proliferation and increases in cell size through Akt/mTOR and STAT3, respectively (Kim et al., 2006; Yang et al., 2018). KRT6 is a negative regulator of cell migration through the inhibition of Src kinase and associations with myosinIIA (Wong and Coulombe, 2003; Rotty and Coulombe, 2012; Wang et al., 2018). Finally, this molecule promotes expression of Desmoplakin and the maintenance of desmosomes, the latter, a long-established feature of SC trophoblasts (Bou-Resli et al., 1981; Bartels and Wang, 1983; Benirschke et al., 2006). Taken together, these observations are consistent with data on SC-CTBs. We found no evidence of their invasion, but did identify frequent interactions with the ECM, CTBs, and EVTs.

537 The coordinated expression of these cytokeratins and the transcriptional profile writ large,
538 suggested the SC has the properties of a highly specialized epidermis with a robust proliferative
539 capacity and strong cohesive properties, but lacking migratory or invasive behavior.
540
As previous work has demonstrated or assumed, the majority of trophoblast identified are
represented in both the VC and SC (Benirschke et al., 2006; Garrido-Gomez et al., 2017; Pique-
Regi et al., 2019). We identified two cell types with striking transcriptional similarity between
regions, CTB 1 and EVT. CTB 1 cells in the VC matched canonical villous CTB in transcriptional
profile, behavior, and niche. Surprisingly, we identified a similar population in the SC. They are
localized to an epithelial sheet juxtaposed to the fetal mesenchyme, separated by a thin basal
lamina. These cells are supported by similar signaling pathways in both regions, including HGF
and WNT (Figure 1-S3g; Dokras et al., 2001; Okae et al., 2018; Zhou et al., 2002). The
similarities in location and growth factor requirements demonstrate that a niche similar to the
villous trophoblast membrane, exists in the SC. Additionally, we identified CTB 1 as the
progenitor population for SC-CTB, and possibly for EVT in the SC. Thus, CTB 1 in both regions
function as multipotent progenitors. Despite the many similarities, the developmental trajectories
emerging from CTB 1 differ in a region-specific manner. At present, it is unclear whether CTB 1
cells in both regions originally have the same potential or if fate restriction occurs as the fetal
membranes form. Future experiments will need to functionally address whether CTB 1 from the
VC can be coerced to generate SC-CTBs and if CTB 1 from the SC can efficiently generate
STBs and invasive EVTs. Multipotent trophoblasts have been generated from first trimester
smooth chorion cells, however, these cells have a distinct transcriptional profile from the second
trimester cells we characterized (Genbacev et al., 2016).
560
Similarities between the EVT of each region, as defined by contact with the decidua, has been
documented in observational, molecular, and most recently, transcriptomic studies (Benirschke
et al., 2006, Pique-Regi et al., 2019). Our data confirmed a strong correlation between gene
expression for all EVT populations, regardless of regional identity. Based on previously
established EVT subtypes, we identified a reduction in the proportion of interstitial EVTs and a
concomitant increase in columnar-like EVTs of the SC. However, we did not find evidence of
specific EVT subpopulations or intrinsic gene expression programs that would explain the
differences in the depth of invasion of EVT in each placental region. However, we did uncover
CTB-EVT interactions specific to the SC epithelium. Rather than physically separating as they
do in the VC, CTBs and EVTs co-occupy the stratified epithelium of the SC, providing a
contained environment for paracrine signaling, cell contact mediated signaling, and cell-ECM
interactions that may impact trophoblast invasion. In this study we verified that soluble signaling
factors from SC cells restrict the invasion of their counterparts from the VC. We can speculate
on potential candidates that may function across multiple biological systems to restrict invasion.
CTB 4 cells express high levels of TIMP1, TIMP3, and SERPINE1 (PAI-1), which inhibit the
function and/or activation of MMPs necessary for trophoblast invasion (Fisher et al., 1989; Zhu
et al., 2012). Addition of recombinant TIMPs or a function blocking antibody against plasminogen
(whose processing is inhibited by SERPINE1) reduced trophoblast invasion *in vitro* (Lala and
Graham, 1990). While these molecules are expressed by decidual cells, we identified their
expression within the SC epithelium directly adjacent to EVTs. Another candidate regulator of

invasion expressed by CTB 4 is TNFα. Treatment of trophoblasts with TNFα increases EVT apoptosis, decreases protease expression, increases SERPINE1 expression, and decreases invasion *in vitro* (Otun et al., 2011; Xu et al., 2011). These data suggest that SC-CTBs secrete multiple molecules that could decrease the survival and invasion of EVTs in the SC. In this study we explored the role of secreted factors, but SC-CTBs may also influence EVTs through cell contact mediated signaling and the deposition of an ECM distinct from the basal plate. Future experiments will be necessary to explore the contributions of these distinct signaling pathways and matrices.

In summary, this study provides a high-resolution molecular accounting of the trophoblast that occupy smooth chorion. By comparison with trophoblasts isolated from the VC of the same placentas, we identified key similarities and differences to better understand the molecular determinants of trophoblast function and behavior in the SC. We characterized a novel CTB population, marked by KRT6A expression, that is central to three key functions of the SC: formation of an epidermal-like barrier, blockage of aberrant STB differentiation, and restriction of EVT invasion. These data provide a better understanding of molecular and cellular pathways that control human placental development and against which disease related changes can be identified and therapeutic targets discovered.

599 **Key Resources Table**

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Antibody	Rat anti E-cadherin Monoclonal Antibody (ECCD-2)	Thermofisher Scientific	13-1900	IF (1:250)
Antibody	Rabbit Anti-PAGE4 antibody (Polyclonal)	Sigma Aldrich	HPA023880	IF (1:100)
Antibody	Rabbit Recombinant Anti-Cytokeratin 6 Monoclonal antibody [EPR1603Y]	Abcam	ab52620	IF (1:100)
Antibody	Rabbit anti Non-phospho (Active) β-Catenin (Ser33/37/Thr41) Monoclonal Antibody	Cell Signaling	8814	IF (1:100)
Antibody	Mouse anti Cytokeratin 14 Monoclonal Antibody (LL002)	Invitrogen	MA5-11599	IF (1:100) with Antigen Retrieval
Antibody	Rabbit Anti-KLF4 antibody (polyclonal)	Sigma Aldrich	HPA00292	IF (1:100)
Antibody	Cytokeratin (7D3) Monoclonal Antibody	Susan Fisher/University of California, San Francisco Cat# Fisher_001-clone7D3,RRID:AB_2631235 https://antibodyregistry.org/search?q=AB2631235	AB_2631235	IF (1:100)
Antibody	HLA-G (4H84) Monoclonal Antibody	Susan Fisher/University of California, San Francisco Cat# Fisher_002-clone4H84, RRID:AB_2631236 https://antibodyregistry.org/search?q=AB2631236	AB_2631236	IF (1:20) with Antigen Retrieval
Software, algorithm	R	https://www.r-project.org/		
Software, algorithm	ImageJ	ImageJ (http://imagej.nih.gov/ij/)		
Software, algorithm	Seurat (3.1.3)	https://satijalab.org/seurat/		
Software, algorithm	cellranger (3.0.2)	https://support.10xgenomics.com/single-cell-gene-expression/software/pipelines/latest/feature-bc		
Software, algorithm	ClusterProfiler	https://guangchuangyu.github.io/software/clusterProfiler/		
Software, algorithm	scVelo	https://github.com/theislab/scvelo		
Software, algorithm	Prism 6.0	https://www.graphpad.com/scientific-software/prism/		
Software, algorithm	DoubletFinder	https://github.com/chris-mcginnis-ucsf/DoubletFinder		

600
601

Methods

Cellular Isolation of VC trophoblast: Trophoblast were isolated from both floating and anchoring villi dissected from 2nd trimester human placentas. Trophoblast were isolated according to previously published protocols (Fisher et al., 1989; Kliman and Feinberg, 1990).

Cellular Isolation of SC trophoblast: Trophoblast were isolated according to previously published protocols (Garrido-Gomez et al., 2017).

Single Cell RNA-Sequencing and Analysis: To capture the transcriptome of individual cells we used the Chromium Single Cell 3' Reagent V3 Kit from 10X Genomics. For all samples 17,500 cells were loaded into one well of a Chip B kit for GEM generation. Library preparation including, reverse transcription, barcoding, cDNA amplification, and purification was performed according to Chromium 10x V3 protocols. Each sample was sequenced on a NovaSeq 6000 S4 to a depth of approximately 20,000-30,000 reads per cell. The gene expression matrices for each dataset was generated using the CellRanger software (v3.0.2 - 10x Genomics). All reads were aligned to GRCh38 using STAR. (https://support.10xgenomics.com/single-cell-gene-expression/software/pipelines/latest/advanced/references). The counts matrix was thresholded and analyzed in the package Seurat (v3.1.3). Cells with fewer than 500 or greater than 6000 unique genes, as well as all cells with greater than 15 percent mitochondrial counts, were excluded from all subsequent analyses. Doublet detection was performed for each sample using DoubletFinder and all doublets were excluded from analysis. For each sample, counts were scaled and normalized using ScaleData and NormalizeData, respectively, with default settings FindVariableFeatures used to identify the 2000 most variable genes as input for all future analyses. PCA was performed using RunPCA and significant PCs assessed using ElbowPlot and DimHeatmap. Dimensionality reduction and visualization using UMAP was performed by RunUMAP.

Integration of each timepoint into one dataset was performed using FindIntegrationAnchors and IntegrateData, both using 20 dimensions (after filtering each dataset for number of genes, mitochondrial counts, and normalizing as described above). Data scaling, PCA, selection of PCs, clustering and visualization proceeded as described above using 30 PCs and a resolution of 0.6.

To generate the trophoblast, stroma, and immune cell subsets, the respective clusters were subset from the integrated dataset using the function SubsetData based upon annotations from marker genes identified by FindAllMarkers. After subsetting, counts were scaled and normalized using ScaleData and NormalizeData, respectively, with default settings FindVariableFeatures used to identify the 2000 most variable genes. Differentially expressed genes for each integrated dataset were identified using FindAllMarkers.

scVelo: RNA velocity analysis was applied to the entire conglomerate dataset using Velocyto to generate spliced and unspliced reads for all cells. This dataset was then subset for the trophoblast dataset introduced in Figure 2. The scVelo dynamical model was run with default settings and subset by each timepoint.

Differential expression between regions: The number of trophoblast cells in each cluster were downsampled to 100 cells from each region of origin and performed differential expression using FindMarkers function and repeated this for 100 permutations. The number of genes found to be significantly differentially expressed (adj. P-value <0.05) from each permutation are plotted. Clusters having less that 100 cells from each region were excluded from this analysis.

Gene Ontology Analysis: Gene Ontology analysis was performed with ClusterProfiler enrichGO function. The simplify function within this package was used to consolidate hierarchically related terms using a cutoff of 0.5. Terms were considered significantly enriched with an adjusted P-value of less than 0.05.

Immunofluorescence Staining: Placental tissue for cryosectioning was fixed in 3% PFA at 4°C for 8hrs, washed in 1x PBS, then submerged in 30% sucrose overnight at 4°C prior to embedding in OCT medium. Placental tissue in OCT was sectioned at 5μm for all conditions. In brief, slides were washed in 1x PBST (1x PBS, 0.05% Tween-20), blocked for 1hr (1x PBS + 5% donkey or goat serum + 0.3% TritonX), incubated in primary antibody diluted for 3 hours at room temperature (or overnight at 4°C), washed in 1x PBST, incubated in secondary antibody (Alexafluor 488 or 594) for 1 hour at room temperature, incubated in DAPI for 10min at room temp, washed in 1x PBST, and mounted and sealed for imaging. Any antigen retrieval was performed prior to the blocking step by heating the slides in a 1x citrate buffer with 0.05% Tween-20 at 95C for 30 minutes. All antibodies and the dilutions are listed in the key resources table. All immunofluorescence staining was performed in n=3 biological replicates (3 distinct placentas) and representative image is shown.

Transwell Invasion Assay: 24-well plate transwell inserts with an 8 μM polycarbonate membrane (Corning Costar 3422) were coated with 10μL of Matrigel (growth factor-containing, Corning Corp, USA) diluted 1:1 in Serum Free Media (95% DME H-21 + Glutamine, 2% Nutridoma (mostly β-D xylopyranose), 1% Pen Strep, 1% Hepes, 0.1% Gentamycin). Cells from each region of the placenta were isolated as described above and then plated in the upper well of the transwell insert at a density of 250,000 cells in 250μL of Serum Free Media, with 1mL of Serum Free Media (or conditioned media) in the well below the insert. The cells were then cultured for 39 hours. The cells in the transwell were then fixed in 3% PFA for 10min at 4C, permeabilized in ice cold methanol for 10min at 4C, then washed in 1x PBS, incubated in Pan-CK primary antibody for 3 hours at 37C, washed in 1x PBS, incubated with secondary antibody (Alexafluor 594) and DAPI for 1 hour at 37C, washed in 1x PBS, then mounted and sealed for imaging.

Transwell Invasion Assay Quantification: Transwell membranes mounted on slides were imaged at 40x magnification and the number of Pan-CK positive projections through the membrane counted. Normalization for changes in cell density across fields of view and culture conditions was performed by quantifying the DAPI positive area of each image and the number of projections normalized to the median across comparisons.

Culture of placental cells and generation of conditioned media: Each well of a 24-well plate
was coated with 20µL of Matrigel undiluted (growth factor-containing, Corning Corp, USA) prior
to cell seeding. Cells from each region of the placenta were isolated as described above and
then plated into wells of a 24-well plate at a density of 1×10^6 cells in 1mL of Serum Free Media
(95% DME H-21 + Glutamine, 2% Nutridoma (mostly β-D xylopyranose), 1% Pen Strep, 1%
Hepes, 0.1% Gentamycin). The cells were cultured for 39 hours. After 39 hours in culture the
media was removed and centrifuged at 2000g for 10mins to remove cellular debris. The
supernatant was then removed, snap frozen in LN2, and stored at -80C. This media was then
thawed and used as conditioned media.

Data Availability

All sequencing data is available at the NCBI Gene Expression Omnibus GSE198373. Processed data are available as R objects at https://figshare.com/projects/Regionally_distinct_trophoblast_regulate_barrier_function_and_invasion_in_the_human_placenta/135191.

Acknowledgements

We thank the members of the University of California – San Francisco National Center of Translational Research in Reproduction and Infertility for helpful comments during the design, execution, and publication of this project. We would also like to thank all members of the Blelloch and Fisher Labs for their comments and support. We specifically thank Ali San and Nasim Zeighami for their research contributions. We would also like to acknowledge our funding sources, the NIH Eunice Kennedy Shriver National Institute for Child Health and Human Development P50 HD055764 and NIH R37 HD076253.

Competing Interests

Susan Fisher is a consultant for Novo Nordisk. All other authors declare no competing interests.

References

Baczyk, D., Drewlo, S., Proctor, L., Dunk, C., Lye, S., & Kingdom, J. (2009). Glial cell missing-1 transcription factor is required for the differentiation of the human trophoblast. *Cell Death and Differentiation*, *16*(5), 719–727. https://doi.org/10.1038/cdd.2009.1

Bailey, C. C., Zhong, G., Huang, I. C., & Farzan, M. (2014). IFITM-family proteins: The cell's first line of antiviral defense. *Annual Review of Virology*, *1*(1), 261–283. https://doi.org/10.1146/annurev-virology-031413-085537

Bartles, H., & Wang, T. (1983). Intercellular Junctions in the Human Fetal Membranes. *Anatomy and Embryology*, 103–120.

Benirschke, K., Kaufmann, P., & Baergen, R. (2006). Pathology of the Human Placenta. In *Pathology* (Vol. 38, Issue 2). https://doi.org/10.1016/s0031-3025(16)39684-2

Bergen, V., Lange, M., Peidli, S., Wolf, F. A., & Theis, F. J. (2020). Generalizing RNA velocity to transient cell states through dynamical modeling. *Nature Biotechnology*, *38*(12), 1408–1414. https://doi.org/10.1038/s41587-020-0591-3

Blaise, S., De Parseval, N., Bénit, L., & Heidmann, T. (2003). Genome wide screening for fusogenic human endogenous retrovirus envelopes identifies syncytin 2, a gene conserved on primate evolution. *Proceedings of the National Academy of Sciences of the United States of America*, *100*(22), 13013–13018. https://doi.org/10.1073/pnas.2132646100

Boyd, J. D., & Hamilton, W. J. (1967). Development and Structure of the Human Placenta From the End of the 3Rd Month of Gestation. *BJOG: An International Journal of Obstetrics & Gynaecology*, *74*(2), 161–226. https://doi.org/10.1111/j.1471-0528.1967.tb14864.x

Buchrieser, J., Degrelle, S. A., Couderc, T., Nevers, Q., Disson, O., Manet, C., Donahue, D. A., Porrot, F., Hillion, K. H., Perthame, E., Arroyo, M. V., Souquere, S., Ruigrok, K., Dupressoir, A., Heidmann, T., Montagutelli, X., Fournier, T., Lecuit, M., & Schwartz, O. (2019). IFITM proteins inhibit placental syncytiotrophoblast formation and promote fetal demise. *Science*, *365*(6449), 176–180. https://doi.org/10.1126/science.aaw7733

Del Peso, L., Castellanos, M. C., Temes, E., Martín-Puig, S., Cuevas, Y., Olmos, G., & Landázuri, M. O. (2003). The von Hippel Lindau/hypoxia-inducible factor (HIF) pathway regulates the transcription of the HIF-proline hydroxylase genes in response to low oxygen. *Journal of Biological Chemistry*, *278*(49), 48690–48695. https://doi.org/10.1074/jbc.M308862200

Dokras, A., Gardner, L. M. G., Seftor, E. A., & Hendrix, M. J. C. (2001). Regulation of human cytotrophoblast morphogenesis by hepatocyte growth factor/scatter factor. *Biology of Reproduction*, *65*(4), 1278–1288. https://doi.org/10.1095/biolreprod65.4.1278

Ferreira, L. M. R., Meissner, T. B., Mikkelsen, T. S., Mallard, W., O'Donnell, C. W., Tilburgs, T., Gomes, H. A. B., Camahort, R., Sherwood, R. I., Gifford, D. K., Rinn, J. L., Cowan, C. A., & Strominger, J. L. (2016). A distant trophoblast-specific enhancer controls HLA-G expression at the maternal-fetal interface. *Proceedings of the National Academy of*

Sciences of the United States of America, *113*(19), 5364–5369.
https://doi.org/10.1073/pnas.1602886113

Fisher, S. J., Cui, T., Zhang, L., Hartman, L., Grahl, K., Guo-Yang, Z., Tarpey, J., & Damsky,
C. H. (1989). Adhesive and degradative properties of human placental cytotrophoblast
cells in vitro. *Journal of Cell Biology*, *109*(2), 891–902.
https://doi.org/10.1083/jcb.109.2.891

Garrido-Gomez, T., Ona, K., Kapidzic, M., Gormley, M., Simón, C., Genbacev, O., & Fisher, S.
J. (2017). Severe pre-eclampsia is associated with alterations in cytotrophoblasts of the
smooth chorion. In *Development (Cambridge)* (Vol. 144, Issue 5, pp. 767–777).
https://doi.org/10.1242/dev.146100

Genbacev, O., Larocque, N., Ona, K., Prakobphol, A., Garrido-Gomez, T., Kapidzic, M.,
Bárcena, A., Gormley, M., & Fisher, S. J. (2016). Integrin α4-positive human trophoblast
progenitors: Functional characterization and transcriptional regulation. *Human
Reproduction*, *31*(6), 1300–1314. https://doi.org/10.1093/humrep/dew077

Genbačev, O., Vićovac, L., & Larocque, N. (2015). The role of chorionic cytotrophoblasts in the
smooth chorion fusion with parietal decidua. In *Placenta* (Vol. 36, Issue 7, pp. 716–722).
https://doi.org/10.1016/j.placenta.2015.05.002

Genbacev, O., Zhou, Y., Ludlow, J. W., & Fisher, S. J. (1997). Regulation of human placental
development by oxygen tension. *Science*, *277*(5332), 1669–1672.
https://doi.org/10.1126/science.277.5332.1669

Haider, S., Meinhardt, G., Saleh, L., Kunihs, V., Gamperl, M., Kaindl, U., Ellinger, A., Burkard,
T. R., Fiala, C., Pollheimer, J., Mendjan, S., Latos, P. A., & Knöfler, M. (2018). Self-
Renewing Trophoblast Organoids Recapitulate the Developmental Program of the Early
Human Placenta. *Stem Cell Reports*, *11*(2), 537–551.
https://doi.org/10.1016/j.stemcr.2018.07.004

Hamilton, W. J., & Boyd, J. D. (1960). Development of the human placenta in the first three
months of gestation. *Journal of Anatomy*, *94*, 297–328.
http://www.ncbi.nlm.nih.gov/pubmed/14399291%0Ahttp://www.pubmedcentral.nih.gov/arti
clerender.fcgi?artid=PMC1244370

Harris, L. K., Jones, C. J. P., & Aplin, J. D. (2009). Adhesion Molecules in Human Trophoblast
- A Review. II. Extravillous Trophoblast. *Placenta*, *30*(4), 299–304.
https://doi.org/10.1016/j.placenta.2008.12.003

Harvey, K. F., Zhang, X., & Thomas, D. M. (2013). The Hippo pathway and human cancer.
Nature Reviews Cancer, *13*(4), 246–257. https://doi.org/10.1038/nrc3458

Karantza, V. (2011). Keratins in health and cancer: More than mere epithelial cell markers.
Oncogene, *30*(2), 127–138. https://doi.org/10.1038/onc.2010.456

Kliman, H. J., & Feinbergt, R. F. (1990). Human trophoblast-extracellular matrix (ECM)
interactions in vitro: ECM thickness modulates morphology and proteolytic activity
(invasion/placenta/choriocarcinoma/Matrigel/cyclic AMP). *Proc. Nat. Acad. Sci. USA*,
87(April), 3057–3061.

Knöfler, M., Haider, S., Saleh, L., Pollheimer, J., Gamage, T. K. J. B., & James, J. (2019).
Human placenta and trophoblast development: key molecular mechanisms and model
systems. In *Cellular and Molecular Life Sciences*. https://doi.org/10.1007/s00018-019-
03104-6

Lala, P. K., & Graham, C. H. (1990). Mechanisms of trophoblast invasiveness and their control:
the role of proteases and protease inhibitors. *Cancer and Metastasis Review*, *9*(4), 369–
379. https://doi.org/10.1007/BF00049525

Lee, C. Q. E., Gardner, L., Turco, M., Zhao, N., Murray, M. J., Coleman, N., Rossant, J.,
Hemberger, M., & Moffett, A. (2016). What Is Trophoblast? A Combination of Criteria
Define Human First-Trimester Trophoblast. *Stem Cell Reports*, *6*(2), 257–272.
https://doi.org/10.1016/j.stemcr.2016.01.006

Liu, X.-Y., Chen, W., Wei, B., Shan, Y.-F., & Wang, C. (2011). IFN-Induced TPR Protein IFIT3
Potentiates Antiviral Signaling by Bridging MAVS and TBK1. *The Journal of Immunology*,
187(5), 2559–2568. https://doi.org/10.4049/jimmunol.1100963

Liu, Y., Fan, X., Wang, R., Lu, X., Dang, Y. L., Wang, H., Lin, H. Y., Zhu, C., Ge, H., Cross, J.
C., & Wang, H. (2018). Single-cell RNA-seq reveals the diversity of trophoblast subtypes
and patterns of differentiation in the human placenta. *Cell Research*.
https://doi.org/10.1038/s41422-018-0066-y

Lu, X., Wang, R., Zhu, C., Wang, H., Lin, H. Y., Gu, Y., Cross, J. C., & Wang, H. (2017). Fine-
Tuned and Cell-Cycle-Restricted Expression of Fusogenic Protein Syncytin-2 Maintains
Functional Placental Syncytia. *Cell Reports*, *21*(5), 1150–1159.
https://doi.org/10.1016/j.celrep.2017.10.019

Maxwell, P. H., Dachs, G. U., Gleadle, J. M., Nicholls, L. G., Harris, A. L., Stratford, I. J.,
Hankinson, O., Pugh, C. W., & Ratcliffe, P. J. (1997). Hypoxia-inducible factor-1
modulates gene expression in solid tumors and influences both angiogenesis and tumor
growth. *Proceedings of the National Academy of Sciences of the United States of*
America, *94*(15), 8104–8109. https://doi.org/10.1073/pnas.94.15.8104

McGinnis, C. S., Murrow, L. M., & Gartner, Z. J. (2019). DoubletFinder: Doublet Detection in
Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Systems*, *8*(4),
329-337.e4. https://doi.org/10.1016/j.cels.2019.03.003

McGowan, K. M., & Coulombe, P. A. (1998). Onset of keratin 17 expression coincides with the
definition of major epithelial lineages during skin development. *Journal of Cell Biology*,
143(2), 469–486. https://doi.org/10.1083/jcb.143.2.469

McMaster, M. T., Librach, C. L., Zhou, Y., Lim, K. H., Janatpour, M. J., DeMars, R., Kovats, S.,
Damsky, C., & Fisher, S. J. (1995). Human placental HLA-G expression is restricted to
differentiated cytotrophoblasts. *Journal of Immunology (Baltimore, Md. : 1950)*, *154*(8),
3771–3778. http://www.ncbi.nlm.nih.gov/pubmed/7706718

Mi, S. (2000). *Syncytin is a captive retrovirus*. *403*(February), 785–789. www.nature.com

Moll, R., Franke, W. W., Schiller, D. L., Geiger, B., & Krepler, R. (1982). The catalog of human
cytokeratins: Patterns of expression in normal epithelia, tumors and cultured cells. *Cell*,
31(1), 11–24. https://doi.org/10.1016/0092-8674(82)90400-7

Nelson, W. G., & Sun, T. T. (1983). The 50- and 58-kdalton keratin classes as molecular
markers for stratified squamous epithelia: Cell culture studies. *Journal of Cell Biology*,
97(1), 244–251. https://doi.org/10.1083/jcb.97.1.244

Okae, H., Toh, H., Sato, T., Hiura, H., Takahashi, S., Shirane, K., Kabayama, Y., Suyama, M.,
Sasaki, H., & Arima, T. (2018). Derivation of Human Trophoblast Stem Cells. *Cell Stem*
Cell, *22*(1), 50-63.e6. https://doi.org/10.1016/j.stem.2017.11.004

Otun, H. A., Lash, G. E., Innes, B. A., Bulmer, J. N., Naruse, K., Hannon, T., Searle, R. F., &
Robson, S. C. (2011). Effect of tumour necrosis factor-α in combination with interferon-γ
on first trimester extravillous trophoblast invasion. *Journal of Reproductive Immunology*,
88(1), 1–11. https://doi.org/10.1016/j.jri.2010.10.003

Parry, S., & Strauss, J. (1998). Premature rupture of the fetal membranes. *The New England*
Journal of Medicine, *338*(10), 663–670.

Red-Horse, K., Zhou, Y., Genbacev, O., Prakobphol, A., Foulk, R., McMaster, M., & Fisher, S.
J. (2004). Trophoblast differentiation during embryo implantation and formation of the
maternal-fetal interface. *The Journal of Clinical Investigation*, *114*(6), 744–754.
https://doi.org/10.1172/JCI22991

Rock, J. R., Gao, X., Xue, Y., Randell, S. H., Kong, Y. Y., & Hogan, B. L. M. (2011). Notch-
dependent differentiation of adult airway basal stem cells. *Cell Stem Cell*, *8*(6), 639–648.
https://doi.org/10.1016/j.stem.2011.04.003

Rock, J. R., Onaitis, M. W., Rawlins, E. L., Lu, Y., Clark, C. P., Xue, Y., Randell, S. H., &
Hogan, B. L. M. (2009). Basal cells as stem cells of the mouse trachea and human airway
epithelium. *Proceedings of the National Academy of Sciences of the United States of*
America, *106*(31), 12771–12775. https://doi.org/10.1073/pnas.0906850106

Romero, R., Dey, S. K., & Fisher, S. J. (2014). Preterm labor: One syndrome, many causes.
Science, *345*(6198), 760–765. https://doi.org/10.1126/science.1251816

Rotty, J. D., & Coulombe, P. A. (2012). A wound-induced keratin inhibits Src activity during
keratinocyte migration and tissue repair. *Journal of Cell Biology*, *197*(3), 381–389.
https://doi.org/10.1083/jcb.201107078

Segre, J. A., Bauer, C., & Fuchs, E. (1999). Klf4 is a transcription factor required for
establishing the barrier function of the skin. *Nature Genetics*, *22*(4), 356–360.
https://doi.org/10.1038/11926

Spence, J. S., He, R., Hoffmann, H. H., Das, T., Thinon, E., Rice, C. M., Peng, T., Chandran,
K., & Hang, H. C. (2019). IFITM3 directly engages and shuttles incoming virus particles to
lysosomes. *Nature Chemical Biology*, *15*(3), 259–268. https://doi.org/10.1038/s41589-
018-0213-2

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., Hao, Y.,
Stoeckius, M., Smibert, P., & Satija, R. (2019). Comprehensive Integration of Single-Cell
Data. *Cell*, *177*(7), 1888-1902.e21. https://doi.org/10.1016/j.cell.2019.05.031

Suryawanshi, H., Morozov, P., Straus, A., Sahasrabudhe, N., Max, K. E. A., Garzia, A.,
Kustagi, M., Tuschl, T., & Williams, Z. (2018). A single-cell survey of the human first-
trimester placenta and decidua. *Science Advances*.
https://doi.org/10.1126/sciadv.aau4788

Takahashi, K., Yan, B., Yamanishi, K., Imamura, S., & Coulombe, P. A. (1998). The two
functional keratin 6 genes of mouse are differentially regulated and evolved independently
from their human orthologs. *Genomics*, *53*(2), 170–183.
https://doi.org/10.1006/geno.1998.5476

Thurman, J. M., & Holers, V. M. (2006). The Central Role of the Alternative Complement
Pathway in Human Disease. *The Journal of Immunology*, *176*(3), 1305–1310.
https://doi.org/10.4049/jimmunol.176.3.1305

Tilburgs, T., Crespo, Â. C., Van Der Zwan, A., Rybalov, B., Raj, T., Stranger, B., Gardner, L.,
Moffett, A., & Strominger, J. L. (2015). Human HLA-G+ extravillous trophoblasts: Immune-
activating cells that interact with decidual leukocytes. *Proceedings of the National
Academy of Sciences of the United States of America*, *112*(23), 7219–7224.
https://doi.org/10.1073/pnas.1507977112

Tirosh, I., Izar, B., Prakadan, S. M., Wadsworth, M. H., Treacy, D., Trombetta, J. J., Rotem, A.,
Rodman, C., Lian, C., Murphy, G., Fallahi-Sichani, M., Dutton-Regester, K., Lin, J. R.,
Cohen, O., Shah, P., Lu, D., Genshaft, A. S., Hughes, T. K., Ziegler, C. G. K., …
Garraway, L. A. (2016). Dissecting the multicellular ecosystem of metastatic melanoma by
single-cell RNA-seq. *Science*, *352*(6282), 189–196.
https://doi.org/10.1126/science.aad0501

Vento-Tormo, R., Efremova, M., Botting, R. A., Turco, M. Y., Vento-Tormo, M., Meyer, K. B.,
Park, J. E., Stephenson, E., Polański, K., Goncalves, A., Gardner, L., Holmqvist, S.,
Henriksson, J., Zou, A., Sharkey, A. M., Millar, B., Innes, B., Wood, L., Wilbrey-Clark, A.,
… Teichmann, S. A. (2018). Single-cell reconstruction of the early maternal–fetal interface
in humans. *Nature*. https://doi.org/10.1038/s41586-018-0698-6

Wang, F., Chen, S., Liu, H. B., Parent, C. A., & Coulombe, P. A. (2018). Keratin 6 regulates
collective keratinocyte migration by altering cell–cell and cell–matrix adhesion. *Journal of
Cell Biology*, *217*(12), 4314–4330. https://doi.org/10.1083/jcb.201712130

Wong, P., & Coulombe, P. A. (2003). Loss of keratin 6 (K6) proteins reveals a function for
intermediate filaments during wound repair. *Journal of Cell Biology*, *163*(2), 327–337.
https://doi.org/10.1083/jcb.200305032

Xu, B., Nakhla, S., Makris, A., & Hennessy, A. (2011). TNF-α inhibits trophoblast integration
into endothelial cellular networks. *Placenta*, *32*(3), 241–246.
https://doi.org/10.1016/j.placenta.2010.12.005

Yabe, S., Alexenko, A. P., Amita, M., Yang, Y., Schust, D. J., Sadovsky, Y., Ezashi, T., &
Roberts, R. M. (2016). Comparison of syncytiotrophoblast generated from human

embryonic stem cells and from term placentas. *Proceedings of the National Academy of*
Sciences of the United States of America, *113*(19), E2598–E2607.
https://doi.org/10.1073/pnas.1601630113

Yang, L., Jin, L., Ke, Y., Fan, X., Zhang, T., Zhang, C., Bian, H., & Wang, G. (2018). E3 Ligase
Trim21 Ubiquitylates and Stabilizes Keratin 17 to Induce STAT3 Activation in Psoriasis.
Journal of Investigative Dermatology, *138*(12), 2568–2577.
https://doi.org/10.1016/j.jid.2018.05.016

Yeh, I. T., O'Connor, D. M., & Kurman, R. J. (1989). Vacuolated Cytotrophoblast: A
subpopulation of trophoblast in the chorion laeve. *Placenta*, *10*(5), 429–438.
https://doi.org/10.1016/0143-4004(89)90053-2

Yu, G., Wang, L. G., Han, Y., & He, Q. Y. (2012). ClusterProfiler: An R package for comparing
biological themes among gene clusters. *OMICS A Journal of Integrative Biology*, *16*(5),
284–287. https://doi.org/10.1089/omi.2011.0118

Yuan, B., Ohyama, K., Bessho, T., & Toyoda, H. (2006). Contribution of inducible nitric oxide
synthase and cyclooxygenase-2 to apoptosis induction in smooth chorion trophoblast cells
of human fetal membrane tissues. *Biochemical and Biophysical Research*
Communications, *341*(3), 822–827. https://doi.org/10.1016/j.bbrc.2006.01.042

Yuan, B., Ohyama, K., Bessho, T., Uchide, N., & Toyoda, H. (2008). Imbalance between ROS
production and elimination results in apoptosis induction in primary smooth chorion
trophoblast cells prepared from human fetal membrane tissues. *Life Sciences*, *82*(11–12),
623–630. https://doi.org/10.1016/j.lfs.2007.12.016

Yuan, B., Ohyama, K., Takeichi, M., & Toyoda, H. (2009). Direct contribution of inducible nitric
oxide synthase expression to apoptosis induction in primary smooth chorion trophoblast
cells of human fetal membrane tissues. *International Journal of Biochemistry and Cell*
Biology, *41*(5), 1062–1069. https://doi.org/10.1016/j.biocel.2008.09.031

Zhou, Y., McMaster, M., Woo, K., Janatpour, M., Perry, J., Karpanen, T., Alitalo, K., Damsky,
C., & Fisher, S. J. (2002). Vascular endothelial growth factor ligands and receptors that
regulate human cytotrophoblast survival are dysregulated in severe preeclampsia and
hemolysis, elevated liver enzymes, and low platelets syndrome. *American Journal of*
Pathology, *160*(4), 1405–1423. https://doi.org/10.1016/S0002-9440(10)62567-9

Zhu, J.-Y., Pang, Z.-J., & Yu, Y.-H. (2012). Regulation of trophoblast invasion: the role of
matrix metalloproteinases. *Reviews in Obstetrics & Gynecology*, *5*(3–4), e137-43.
https://doi.org/10.3909/riog0196

<u>Supplementary Tables</u>

Supplementary Table 1 - Number of cells captured from each region per cluster for the
trophoblast dataset.

Supplementary Figures
952
953

Figure 1-S1 – **Metrics of the integrated dataset. (a)** Violin plots of the number of unique genes (Left), number of UMI (Middle), and the percent of mitochondrial reads (Right) per cell for each broad cell type cluster. **(b)** The total number of cells (Left) and the percent of the total library (Right) in each broad cell type cluster from each placental sample are shown. **(c)** UMAP projections of the integrated dataset shown by each placental sample. Colors correspond to each broad cell type cluster in the legend at the right. The number of cells analyzed from each placental sample is listed beneath. **(d)** The number of cells in each broad cell type cluster from each placental region (VC – black; SC – blue).

Figure 1-S2 – Metrics and markers of the immune cell subset. (a) UMAP projection of all subclustered immune cells (n=14,805). Colors correspond to the clusters in the legend at the right. **(b)** Heatmap of selected marker genes of each immune cell cluster. Expression was displayed as the scaled mean expression in the cluster. **(c)** Violin plot of XIST expression in each immune cell cluster. **(d)** UMAP projection of immune cells shown by region of origin. **(e)** Stacked bar chart of the number of cells (left) or proportion of cells (right) in each cluster by region of origin.

Figure 1-S3 – Metrics and markers of the stromal cell subset. (a) UMAP projection of subclustered stromal cells (n=3,883). Colors correspond to the clusters in the legend at the right. **(b)** Heatmap of selected marker genes for each stromal cell cluster. Expression was displayed as the scaled mean expression in the cluster. **(c)** UMAP projection of stromal cells shown by region of origin. **(d)** Stacked bar chart of the number of cells (left) or percent of cells (right) in each cluster by region of origin. **(e)** Violin plot of XIST expression in each stromal cell cluster. **(f)** Violin plot of COL11A1 expression in each cluster shown by each region, showing expression in only SC cells **(g)** Violin plots of HGF, WNT2, and RSPO3 expression in each cluster and shown by each region, showing expression in both VC and SC cells.

Figure 2-S1 – Metrics of the trophoblast subset. (a) Violin plots of the number of unique genes (left), number of UMI (middle), and the percent of mitochondrial reads (right) per cell for each trophoblast cluster. **(b)** UMAP projections of the trophoblast dataset shown by each placental sample. Colors correspond to each trophoblast cluster shown in the legend at the right. The number of cells analyzed from each placental sample is listed beneath. **(c)** The number of cells (left) and the percentage (right) in each trophoblast cluster from each placental sample.

Figure 2-S2 – Metrics of the trophoblast subset. (a) Heatmap of the expression of the top 20 marker genes for each trophoblast population. **(b)** Expression of phasic transcripts MKI67 (top), PCNA (middle), and TOP2A (bottom) were projected in UMAP space. **(c)** Heatmap of the Pearson Correlation coefficients between each trophoblast population.

a CDH1

b
Villous Chorion   Smooth Chorion

c
Villous Chorion   Villous Chorion

d
KRT6A

KRT6B

KRT6C

Figure 2-S3 – KRT6 expression in the VC region. (a) Violin plot of *CDH1* transcript expression across all trophoblast clusters. **(b)** Immunofluorescence co-localization of CDH1 and PAGE4 in the VC (Left) and SC (Right). **(c)** Immunofluorescence co-localization of Pan-CK (marker of all trophoblast) and KRT6 in the VC. Arrowheads denote KRT6+ cells. **(d)** Violin plots of expression of each KRT6 isoform, showing expression of only KRT6A in trophoblasts. For all images nuclei were visualized by DAPI stain; Scale bar = 100µm. Abbreviations: AV = Anchoring Villi; Amn. = Amnion; SC = Smooth Chorion epithelium; Dec. = Decidua; FV = Floating Villi.

Figure 3-S1 – SC trophoblast display reduced expression of STB and increased expression of epithelial TFs. (a)
Violin plots of expression of markers of STB differentiation in each trophoblast cluster shown by region, demonstrating reduced expression all markers in STB Precursor and STB clusters. **(c)** Violin plots of expression of YAP1 and HES1, demonstrating increasing expression in CTB1-4 and greater expression in the SC compared to the VC.

Figure 4-S1 – Complete CTB Gene Ontology Analysis. The unabridged Gene Ontology results from each CTB cluster, displayed as adjusted p-values. Dark red corresponds to the lowest p-values and white represents p-values greater than 0.0005. Ontology categories are organized by hierarchical clustering along the y-axis. Marker genes for each cluster were used as inputs for the analysis.

Figure 4-S2 – Similarities and differences between CTB 1 in VC and SC. (a) Immunofluorescence co-localization of CDH1 and non-phosphorylated CTNNB1 in the VC (left) and SC (right), showing identical domains of expression in both regions. For all images nuclei were visualized by DAPI stain; Scale bar = 100μm. Abbreviations: FV = Floating Villi; MB = Maternal Blood space; SC = Smooth Chorion epithelium; Dec. = Decidua. **(b)** Gene ontology analysis for CTB 1 genes common to VC and SC, genes enriched in VC, and genes enriched in SC. **(c)** Violin plots of genes within the gene ontology categories indicated in (b) for common categories (top), VC enriched categories (middle), and SC enriched categories (bottom).

Figure 4-S3 – Cytokeratin expression in VC and SC trophoblast. (a) Violin plots of cytokeratins expressed in trophoblast in both VC and SC regions by cluster and region of origin. **(b)** Violin plots of cytokeratins expressed in SC trophoblast by cluster and region of origin.

Figure 4-S4 – IFITM3 expression in CTB populations. Violin plot of IFITM3 expression in trophoblast clusters by region, showing increased expression in SC compared to VC.

none

Figure 5-S1 – Functional annotation of EVT clusters. (a) Gene Ontology results from each EVT cluster displayed as adjusted p-values. Dark red corresponds to the lowest p-values and white represents p-values greater than 0.0005. Ontology categories are organized by hierarchical clustering along the y-axis. Marker genes for each cluster were used as inputs for the analysis. **(b)** Selected genes in the 'placenta development' gene ontology category (top). Violin plots of expression for selected genes in each EVT cluster by region. **(c)** Selected genes in the 'Extracellular Structure Organization' gene ontology category (top). Violin plots of expression for selected genes in each EVT cluster by region.

a

b

c

Pearson Correlation Between Regions	
Cluster	***r***
CTB 1	0.813
EVT 4	0.772
EVT 3	0.769
CTB S-phase	0.747
EVT 2	0.741
CTB G2M-phase	0.680
CTB 2	0.653
CTB 4	0.647
EVT 1	0.632
STB Precursor	0.596
CTB 3	0.497

Figure 5-S2 – Differential expression between EVT from VC and SC. (a) Volcano plots showing the differentially expressed genes the VC or SC for each EVT cluster. Only significantly differentially expressed genes are plotted (p-value <0.05). Genes with higher expression in VC compared to SC are shown in black. Genes with higher expression in SC compared to VC are shown in blue. Selected differentially expressed genes are labelled. **(b)** Violin plots of VC-specific genes in EVTs. **(c)** Pearson correlation coefficients between VC and SC for each cluster ranked from greatest to least similarity across regions. STB and EVT Precursor populations are excluded due to the low number of cells recovered in the SC.

Figure 6-S1 – CTB-EVT interactions in the VC or SC region. Colocalization of KRT6 (SC-CTB marker) and HLA-G (EVT marker) showing rare KRT6+ cells in the VC (top). These cells are few and do not interact with EVT in the same manner as was observed in the SC. Co-localization of non-phosphorylated CTNNB1 (CTB 1 marker) and HLA-G (EVT marker) in the SC (Bottom). Limited interactions between these populations was observed. The basal lamina separating the fetal stroma from the SC epithelium is marked by the white dashed line. For all images nuclei were visualized by DAPI stain; Scale bar = 100μm. Abbreviations: AV = Anchoring Villi; FV = Floating Villi; SC = Smooth Chorion epithelium; Dec. = Decidua.

Transswell Invasion Assay
Pan-CK Projections

Figure 6-S2 – Representative images of the transwell invasion assay. Representative images of trophoblast projections through the transwell filter are seen a white dots. For quantification, the number of projections (white bright dots) were counted. For each sample, the median DAPI area was quantified across the transwell membrane, and then the projections multiplied by a factor to normalize for cell density. Scale bar = 100μm.

Figure 6-S3 – Cell density is not correlated with culture in conditioned media. The DAPI+ area in each field of view was quantified and plotted as a measure of cell density. Each dot represents the percent area of the field of view which stained with DAPI. Measurements from each experiment are shown and the gestational ages of the cultured cells is on the y-axis at the left. Measurements of cells treated with control media (Serum Free Media) are shown in red, VC cell conditioned media are shown in black, and SC cell conditioned media are shown in open squares.