1   **Macrophages only sense infectious SARS-CoV-2 when they express sufficient ACE2 to**
2   **permit viral entry, where rapid cytokine responses then limit viral replication.**
3

4   Larisa I Labzin[1,2,3]*, Keng Yih Chew[4], Kathrin Eschke[1], Xiaohui Wang[1,2], Tyron Esposito[1,2],
5   Claudia J Stocks[1,2], James Rae[1,9], Ralph Patrick[1], Helen Mostafavi[1], Brittany Hill[1], Teodor E.
6   Yordanov[1], Caroline L Holley[1,2], Stefan Emming[1,2], Svenja Fritzlar[5], Francesca L. Mordant[5],
7   Daniel P. Steinfort[6,7], Kanta Subbarao[5,8], Christian M. Nefzger[1], Anne K Lagendijk[1], Emma
8   Gordon[1], Robert Parton[1,9], Kirsty R. Short[3,4], Sarah L. Londrigan[5, #], Kate Schroder[1,2,3,4, #].
9
10 [#]These authors contributed equally
11 * Correspondence should be addressed to l.labzin@uq.edu.au
12
[1]Institute for Molecular Bioscience (IMB), The University of Queensland, Brisbane, QLD,
Australia
[2]IMB Centre for Inflammation and Disease Research, The University of Queensland,
Brisbane, QLD, Australia
[3]Australian Infectious Diseases Research Centre, The University of Queensland, Brisbane,
QLD, Australia
[4]School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane,
QLD, Australia
[5]Dept of Microbiology and Immunology, The University of Melbourne at the Peter Doherty
Institute for Infection and Immunity, Melbourne, Victoria, Australia.
[6]Department of Medicine, The University of Melbourne, Parkville, Victoria, Australia
[7]Department of Respiratory Medicine, Royal Melbourne Hospital, Parkville, Victoria,
Australia
[8] WHO Collaborating Centre for Reference and Research on Influenza, Victorian Infectious
Diseases Reference Laboratory, at The Peter Doherty Institute for Infection and Immunity,
Melbourne, Victoria, Australia.
[9]Centre for Microscopy and Microanalysis, The University of Queensland, Brisbane, QLD,
Australia

31
32
33
34

Abstract

Macrophages are key cellular contributors to COVID-19 pathogenesis. Whether SARS-CoV-2 can enter macrophages, replicate and release new viral progeny remains controversial. Similarly, whether macrophages need to sense replicating virus to drive cytokine release is also unclear. Macrophages are heterogeneous cells poised to respond to their local microenvironment, and accordingly, the SARS-CoV-2 entry receptor ACE2 is only present on a subset of macrophages at sites of human infection. Here, we use in vitro approaches to investigate how SARS-CoV-2 interacts with ACE2-negative and ACE2-positive human macrophages and determine how these macrophage populations sense and respond to SARS-CoV-2. We show that SARS-CoV-2 does not replicate within ACE2-negative human macrophages and does not induce pro-inflammatory cytokine expression. By contrast, ACE2 expression in human macrophages permits SARS-CoV-2 entry, replication, and virion release. ACE2-expressing macrophages sense replicating virus to trigger pro-inflammatory and anti-viral programs that limit virus release. These combined findings resolve several controversies regarding macrophage-SARS-CoV-2 interactions and identify a signaling circuit by which macrophages sense SARS-CoV-2 cell entry and respond by restricting viral replication.

One sentence summary:

Lack of macrophage ACE2 expression precludes SARS-CoV-2 entry and sensing, while ACE2-expressing macrophages sense intramacrophage SARS-CoV-2 replication to induce rapid anti-viral responses that limit new virion release.

Main Text

INTRODUCTION

Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), belonging to the

Coronaviridae family, is an enveloped, single-stranded RNA virus with a positive-sense

genome. Many human respiratory viruses, including SARS-CoV-2, infect epithelial cells

lining the upper and lower airways, resulting in productive replication and the release of

newly synthesized infectious viral particles. The virus Spike (S) glycoprotein facilitates entry

into target epithelial cells by binding to the surface-expressed angiotensin-converting enzyme

2 (ACE2) receptor (*1*). The well-characterized ACE2-Spike interaction exposes a critical S

cleavage site (S2) that can be cleaved by the host serine protease TMPRSS2 (transmembrane

protease serine 2), also expressed on the plasma membrane. This allows the fusion of the

viral and cellular membranes, followed by the release of viral RNA directly into the

cytoplasm (reviewed in (*2*) and (*3*)). SARS-CoV-2, particularly omicron variants, can also

attach to ACE2 and enter cells via the endocytic pathway. Here, the S protein is cleaved by

endosomal proteases to allow fusion between the viral and endosomal membranes (*4*). Host

cell ribosomes immediately translate the infecting, positive sense RNA genome into two

large polyproteins. These polyproteins are proteolytically processed to generate individual

viral proteins required for replication. New virions are assembled in the endoplasmic

reticulum and Golgi of the host cell, before secretion from the cell via exocytosis or through a

lysosomal egress pathway (*3*). Our understanding of the SARS-CoV-2 replication cycle

comes from studies in primary epithelial cells or epithelial cell lines. How this may differ in

other potentially susceptible cell types, including distinct macrophage populations, remains

unclear.

Effective host defense against infection relies on accurate and timely immune detection.

Accumulating evidence suggests that severe COVID-19 results from a failure of early host-

interferon signaling to control SARS-CoV-2, followed by exacerbated pro-inflammatory

responses driving tissue damage. Airway epithelial cells, the primary target for SARS-CoV-2

infection (*5*), respond by releasing anti-viral and pro-inflammatory cytokines (*6*, *7*). Airway-

resident or newly recruited macrophages also appear to be a key source of pro-inflammatory

cytokines in severe COVID-19 (*8*, *9*), with macrophage-derived cytokines implicated in the

severe pathology observed in patients (*10–13*).

Macrophages are sentinel innate immune cells that defend the airways during respiratory

viral infection. Yet, and in contrast to epithelial cells, infection of human macrophages results

in an abortive replication cycle for many respiratory viruses (including seasonal influenza A

viruses (*14*) and rhinovirus (*15*)), despite macrophages supporting the early stages of

infection (entry and synthesis of new viral RNA and protein). Therefore, macrophages can

function as a viral 'dead end' to limit viral dissemination (*16*) and can sense infectious viral

particles, infected cells, and tissue damage to orchestrate anti-viral and pro-inflammatory

programs (*17–19*). Single-cell RNA sequencing has revealed viral RNA within macrophage

populations in human COVID-19 lungs (*8*, *20*) and *ex vivo* lung explants (*21*).

Similarly, immunostaining of COVID-19 autopsy lungs showed macrophages containing

viral antigens (Spike and the RNA-dependent RNA polymerase: RdRp) *in situ* (*22*). Whether

SARS-CoV-2 enters these macrophages via ACE2 remains unclear, as is the fate of

intramacrophage SARS-CoV-2. These in vivo studies did not ascertain if SARS-CoV-2

productively replicates within macrophages to produce new viral RNA and protein leading to

new virion assembly, nor whether macrophages release newly synthesized infectious virions.

Further, previous studies did not elucidate whether SARS-CoV-2 needs to enter and actively

replicate in macrophages to trigger macrophage anti-viral and pro-inflammatory responses.

In vitro models of SARS-CoV-2 interactions with macrophages have focused on human

monocyte-derived macrophages (HMDM). While there is consensus that HMDMs exposed to

SARS-CoV-2 do not release newly-synthesized infectious virions (*7*, *12*, *13*, *23–25*), some

studies report that HMDMs support the early stages of infection that include viral entry and

replication (i.e., viral RNA replication and protein synthesis) (*12*, *13*, *23*). In contrast, others

report that HMDM are refractory to SARS-CoV-2 entry (*7*, *25*). Accordingly, whether

SARS-CoV-2 exposure induces HMDM pro-inflammatory and anti-viral signaling or

whether this requires active viral entry and replication is also controversial. Multiple reports

suggest that macrophage challenge with SARS-CoV2 triggers pro-inflammatory responses

(*12*, *13*, *26*). In contrast, others report that SARS-CoV-2 challenge does not activate

macrophage inflammatory functions, consistent with a lack of viral entry and the early stages

of viral replication (*7*, *25*).

The current study investigates whether macrophage ACE2 expression dictates macrophage

susceptibility to SARS-CoV-2 entry and replication and how this shapes macrophage

inflammatory responses. We found that SARS-CoV-2 cannot enter or replicate within ACE2-
negative macrophages, and these cells do not produce pro-inflammatory cytokines in
response to viral challenge. In contrast, macrophages engineered to ectopically express ACE2
support SARS-CoV-2 entry, replication, and new virion release. ACE2-expressing
macrophages sense newly synthesized viral RNA and induce pro-inflammatory and anti-viral
mediator expression. This rapid induction of anti-viral programs in ACE2-expressing
macrophages limits ongoing virion release. Thus, we identify two blocks to SARS-CoV-2
infection and replication in ACE2-negative and -positive macrophages. ACE2-negative
macrophages are refractory to SARS-CoV-2 infectious entry and do not support the early
stages of replication. In contrast, ACE2-positive macrophages are susceptible to the early
stages of infection and replication but mount a second post-entry block that constrains the
production and release of newly synthesized infectious virions.

RESULTS

In vitro models of ACE2-positive and ACE2-negative macrophages

Given that ACE2 is the primary receptor for SARS-CoV-2, we anticipated that ACE2
expression would determine the susceptibility and permissivity of macrophages to SARS-
CoV-2 infection and replication. We reanalyzed published scRNA-seq lung data from three
studies (*20*, *27*, *28*), representing two sets of control and two sets of COVID-19-infected
individuals. We aligned the datasets to a common cell type reference to study *ACE2* mRNA
expression amongst myeloid cell types (e.g., infiltrating monocytes, monocyte-derived
macrophages, and resident alveolar macrophages). Less than 2% of all cell types expressed
ACE2 mRNA, including alveolar epithelial AT1 and AT2 cells targeted by SARS-CoV-2
(**fig. S1**). In control and COVID-19-infected lungs, a small proportion (<0.05%) of
monocyte-derived macrophages expressed *ACE2* mRNA. Very few tissue-resident
macrophages and transitioning monocyte-derived macrophages expressed *ACE2* mRNA in
control lungs, though this proportion increased in the COVID-19-infected lung. Monocytes
did not express *ACE2* mRNA in either condition (**fig. S1**). These results are consistent with
published scRNA sequencing analyses on *ACE2* mRNA expression across cell types (*20*, *27*–
29). A recent preprint reported ACE2-positive macrophages in lung explants infected *ex vivo*,
as determined by immunofluorescence (*30*). These data suggest that both ACE2-positive and
ACE2-negative macrophages are present in SARS-CoV-2-infected lungs.

We next assessed ACE2 expression in HMDM cultured from blood monocytes to determine

their suitability as an in vitro cell model for ACE2-positive or ACE2- negative macrophages.

Monitoring ACE2 mRNA and protein revealed low *ACE2* mRNA expression in HMDM, and

undetectable levels of ACE2 protein (**Fig. 1A, B**), even when HMDM were treated with

IFNβ (**Fig. 1B**). HMDM thus resemble ACE2-negative macrophages in COVID-19 lungs.

The lung-resident macrophage populations are immune sentinels of the airways, with distinct

origins and properties to HMDM (*31*). Along with airway epithelial cells, airway

macrophages are prime targets for infection by many respiratory viruses. We thus

investigated the ACE2 expression status of this lung-resident macrophage population. We

utilized airway macrophages from bronchoalveolar lavage (BAL) as representative resident

lung macrophages and observed that their ACE2 protein expression was below detection

levels (**Fig. 1C**). Together, this data indicates that HMDM and BAL macrophages cultured ex

vivo model ACE2-negative macrophage populations in vivo.

To model ACE2-positive cells, we used lentiviral transduction to ectopically express either

ACE2 (untagged) or a control protein (mScarlet: mSc) in the THP-1 monocytic cell line prior

to their differentiation into macrophage-like cells with phorbol-myristate-acetate (PMA). We

confirmed that ectopic ACE2 expression in THP-1 cells yielded readily detectable ACE2

mRNA and protein (**Fig. 1A, 1B**). The surface protease TMPRSS2, which is required for S

protein cleavage, was also readily detected at the protein level in HMDM, THP-1-ACE2, and

THP-1-mSc cells (**Fig. 1B**), indicating its presence in both ACE2-positive and ACE2-

negative macrophages**.**

ACE2-negative macrophages do not support SARS-CoV-2 entry and replication

Despite the small proportion of ACE2-positive macrophages at sites of human SARS-CoV-2

infection ((*20*, *27*), **fig S1**), many macrophages are positive for SARS-CoV-2 RNA or protein

in vivo (*8*, *20*, *22*), potentially reflecting an ACE2-independent mechanism of viral entry into

macrophages. We thus investigated whether SARS-CoV-2 can enter HMDMs, which do not

express ACE2 (**Fig. 1A, 1B**)**.** We left the viral inoculum on HMDM to maximize the uptake

of SARS-CoV-2 virions. For these experiments, we included SARS-CoV-2-permissive Calu-

3 epithelial cells as a positive control for viral infection and replication. Calu-3 cells were

susceptible to infection at high and low doses (MOI 5 and 0.5). We detected an increase in cell-associated viral RNA between two hours (indicative of input virus) and 24 h post-infection (p.i.), indicating newly synthesized viral RNA and active replication (**Fig. 2A**). Consistent with productive infection in Calu-3 cells, we also detected viral sub-genomic RNA (indicating new viral RNA synthesis; **Fig. S2A**) and newly synthesized viral nucleoprotein (NP, increased from two to 72 h; **Fig. 2B**). In contrast, cell-associated viral RNA levels did not rise over the same time course in HMDM infected with either MOI dosage (**Fig. 2C**). Viral sub-genomic RNA was barely detectable in HMDM (**Fig. S2B**), and while we could detect NP expression in HMDM at two hours p.i., NP levels were barely detectable 24 h p.i. (**Fig. 2D**). These data indicate that HMDM are not susceptible to SARS-CoV-2 infection and do not support the early stages of viral replication, including the production of newly synthesized viral RNA and protein.

We next tested whether BAL macrophages support ACE2-independent viral entry and replication by challenging these cells with SARS-CoV-2 at an MOI of 1. SARS-CoV-2-challenged BAL macrophages showed no increase in cell-associated viral RNA (indicative of the early stages of virus replication) between two and 48 h post-infection, in contrast to the SARS-CoV-2-susceptible control cell line, VERO E6 (**Fig. 2E**). An increase in viral RNA isolated from cell-free VERO E6 supernatants between two and 48 h post-infection indicated productive viral replication. However, we did not observe an increase in viral RNA in supernatants of SARS-CoV-2 infected BAL macrophages (**Fig. 2F**). As for HMDM, these data indicate that ACE2-negative BAL macrophages are not permissive to the early stages of SARS-CoV-2 infection and replication.

ACE2-positive macrophages support viral entry, replication, and new virion release

We next investigated whether ACE2-positive macrophages could support SARS-CoV-2 entry and replication. We challenged THP-1-ACE2 (**Fig. 1A, 1B**) with SARS-CoV-2 (MOI 0.5 and 5) and compared viral replication and release to THP-1-mSc (ACE2-negative) and Calu-3 control cells. Cell-associated viral RNA significantly increased from 0 h (input) to 24 h p.i. for both MOIs in THP-1-ACE2 cells before plateauing from 24 to 72 h p.i., indicating cell susceptibility and early-stage viral RNA replication (**Fig. 3A, 3B**). By contrast, viral RNA levels did not increase in the THP-1-mSc control cells at any time (**Fig. 3A, 3B**), similar to our observations in other ACE2-negative macrophages (HMDM and BAL; **Fig. 2C, 2E, 2F**).

Calu-3 cells support viral replication and virion release (*32*); accordingly, SARS-CoV-2

RNA levels continued to rise over time (**Fig. 3A, 3B**). Consistent with new viral protein

synthesis, we observed robust NP staining at 24 h post-infection in THP-1-ACE2 but not

THP-1-mSc cells (**Fig. 3C**). Only 16 % of THP-1-ACE2 cells were NP positive at 24 h p.i.

(**Fig. 3D**), indicating heterogeneity in the cellular response to SARS-CoV-2.

To determine whether the increase in cell-associated viral RNA levels in THP-1-ACE2 cells

was coupled to the release of infectious virus, we performed plaque assays on cell-free

supernatants between 0 h and 72 h p.i. At MOI 0.5, we observed a significant increase in

infectious viral particles in the THP-1-ACE2 cell-free supernatants at 24 h p.i., compared

with the levels of residual input virus (0 h p.i.), which is consistent with productive infection

(**Fig. 3E**). We observed a similar increase in THP-1- ACE2 cells at 24 h p.i. using MOI 5

(**Fig. 3F**). Control epithelial cells (Calu-3: **Fig. 3E, 3F**, and A549-ACE2 cells, **Fig. S3**)

supported productive virus release.   As expected, there was no evidence of productive

replication in control THP-1-mSc cells at either MOI (**Fig. 3E, 3F**), which is consistent with

a lack of susceptibility to infection (**Fig. 3A, 3B**). This shows that ACE2-expressing

macrophages support the initial stages of infection, new viral RNA synthesis, and assembly

and release of newly synthesized, infectious viral particles.

Macrophages can internalize SARS-CoV-2 independently of ACE2

Additional SARS-CoV-2 entry receptors beyond ACE2 are reported, including the C-type

lectin receptors (CLRs) (*26*) and CD169(*33*). Since our assays to measure viral protein and

RNA could not distinguish whether the virus is intracellular or extracellular (**Fig. 2C,2D**), we

next determined whether SARS-CoV-2 can still bind and enter HMDM, despite the lack of

ACE2 expression. We used transmission electron microscopy to determine the sub-cellular

location of incoming virions, using a high MOI of 20. HMDM (**Fig. 4A, B**) and THP-1-

ACE2 (**Fig. 4C**) displayed internalized, intact virions within the phagosomal system after one

hour of infection. The THP-1-ACE2 cells also showed virions bound to the plasma

membrane and possibly undergoing fusion (**Fig. 4D, E**). These virions were morphologically

similar to the new structures released from Calu-3 at 72 h p.i. (**Fig. 4F**). Since THP-1-ACE2

cells endogenously express TMPRSS2 (**Fig. 1B**), this suggests that THP-1-ACE2 cells

support viral fusion at the plasma membrane to deliver the viral genome and NP directly into

the cytoplasm. Together, these results indicate that while HMDM take up SARS-CoV-2 into

phagosomal compartments, low ACE2 expression will preclude SARS-CoV-2 S processing

and virus-cell membrane fusion, necessary steps for this virus to enter the cytoplasm.

ACE2-negative macrophages do not release pro-inflammatory cytokines or anti-viral

mediators upon SARS-CoV-2 exposure

Macrophages express a suite of pattern recognition receptors (PRRs) at strategic subcellular

locations to detect microbial challenges. PRRs can sense viral RNA and proteins incorporated

in incoming virions and RNA and proteins synthesized during viral replication (*34–36*). We

detected viral particles in HMDM phagosomal compartments at one-hour p.i. (**Fig. 4B**), and

so we next assessed whether ACE2-negative macrophages could sense incoming virions from

these compartments. We challenged HMDM with high dose SARS-CoV-2 (MOI 5) for 24 h

without removing the virus to allow maximal macrophage responses. SARS-CoV-2 did not

trigger cytokine (C-X-C motif chemokine ligand 10: CXCL10; Interleukin-6: IL-6; or Tumor

Necrosis Factor: TNF) release in vitro (**Fig. 5A**), despite reports of high circulating levels of

these cytokines during human SARS-CoV-2 infection (*37*, *38*). By contrast, HMDM showed

a robust secretory response to synthetic viral mimetics such as the toll-like receptor (TLR)

7/8 ligand, R848 (TNF, IL-6), and the melanoma differentiation-associated protein 5

(MDA5) ligand, transfected poly I:C (pIC) (CXCL10) (**Fig. 5A**). mRNA analyses revealed

that SARS-CoV-2 did not provoke HMDM expression of *IFNB1, IFNL1, CXCL10, IL6,*

TNF, or *IL1B* at two or 24 h p.i., at either low (0.5) or high (5) MOI (**Fig. 5B**). R848 and pIC

stimulation robustly induced these genes, indicating that the HMDM were signaling

competent (**Fig. 5B**). Thus, HMDMs do not respond to SARS-CoV-2 exposure in vitro, even

at the high MOI of 5 or when virions are present in the phagosomal system.

ACE2-positive cells sense replicating SARS-CoV-2 to trigger pro-inflammatory and

anti-viral responses

Since ectopic ACE2 expression in macrophages permits efficient SARS-CoV-2 entry and

early-stage viral replication, we examined whether THP-1-ACE2 cells produce pro-

inflammatory and anti-viral mediators upon SARS-CoV-2 challenge. THP-1-ACE2 cells

strongly upregulated *IFNB1, IFNL1, CXCL10,* and *IL6* mRNA expression after infection with

SARS-CoV-2 for 24 h (at MOI 5, **Fig. 6A**; or MOI 0.5, **fig. S4A**), correlating with increased

SARS-CoV-2 viral RNA levels (**Fig. 3A-B**). In contrast, THP-1-mSc cells did not respond to

SARS-CoV-2 challenge at either MOI (**Fig. 6A, fig. S4A)**, like HMDM (**Fig. 5A, 5B**).

SARS-CoV-2 infection downregulated *IL1B* mRNA levels in THP-1-ACE2 cells but not in

THP-1-mSc cells (**Fig. 6B, fig. S4A**), likely due to negative regulation of *IL1B* mRNA

expression by interferons (*39*). Viral sensing pathways were operational in THP-1-ACE2 and

THP-1-mSc cells, as both cell types responded equally to MDA5 stimulation via pIC

transfection, including robustly downregulating *IL1B* mRNA expression (**fig. S4B**). In Calu-3

cells, anti-viral (*IFNB1, IFNL1, CXCL10*) and pro-inflammatory (*TNF, IL6*) gene induction

peaked at 72 h p.i. (at MOI 5, **Fig. 6C**; MOI 0.5, **fig. S4C**), consistent with published

observations (*7*) and the rise in SARS-CoV-2 viral RNA expression (**Fig. 3A, B**) and

productive virus release (**Fig. 3E, F**). To determine whether THP-1-ACE2 cells responded to

incoming versus replicating virus, we treated cells with the RdRp inhibitor remdesivir (*40*).

Remdesivir ablated viral replication (**Fig. 6D**) and the ensuing induction of *IFNB1, IFNL1,*

and *IL6* (**Fig. 6E**). Intriguingly, remdesivir did not modulate *CXCL10* expression during

infection at MOI 5 (**Fig. 6E**). This suggests that incoming virions may also be sensed and

trigger distinct inflammatory responses compared with replicating virions.   Collectively,

these data indicate that ACE2-positive macrophages are poised to sense entry and early-stage

viral replication and respond by mounting potent anti-viral programs.

316
317   **Blocking cytokine signaling in ACE2-positive cells prolongs productive virion release**

318
Recent studies suggest that host cell death curbs the release of SARS-CoV-2 infectious

virions (*41*). We next explored whether the death of THP-1-ACE2 cells was responsible for

the decline in viral titers at 48h and 72h p.i. (**Fig. 3E, 3F**). Indeed, virus-challenged THP-1-

ACE2 cells showed a modest but significant decrease (~20%) in viability compared to THP-

1-mSc cells at 72 h p.i. (**Fig. 7A**). Thus, cell death may limit virus release from THP-1-ACE2

challenged with high viral loads.

325
We also noted that the rapid and robust induction of type I and III interferons in THP-1-

ACE2 cells peaked at 24 h p.i., much earlier than in Calu-3 cells (**Fig. 6A, 6B**). Interferons

provide autocrine and paracrine signals to activate an anti-viral state through the induction of

interferon-stimulated genes, which act as restriction factors to limit viral infection (*42*).

Indeed, type I IFN aborts influenza A virus infection in murine macrophages (*16*). We thus

hypothesized that interferon signaling in THP-1-ACE2 cells induces host factors that impede

viral replication, limiting the release of productive virions. In contrast, the delayed induction

of *IFNB1* and *IFNL1* in Calu-3 cells, potentially through viral antagonism, may contribute to

ongoing viral replication and release at 48 and 72 h p.i. (**Fig. 3E, 3F**). We tested this

hypothesis using a TBK1 inhibitor (BX-795) to block virus-induced interferon induction and

signaling (*43*) in THP-1-ACE2 cells (**Fig. 7B**). As expected, BX-795 suppressed pIC-induced

expression of *IFNB1*, *IFNL1,* and the interferon-stimulated gene *CXCL10* (**fig. S5**). We next

challenged THP-1-ACE2 and Calu-3 cells with SARS-CoV-2 for 72 h in the presence and

absence of BX-795. BX-795 suppressed SARS-CoV-2-induced CXCL10 production from

THP-1-ACE2 cells (**Fig. 7C**), significantly increasing viral release at 72h p.i. (**Fig. 7D**). BX-

795 did not boost the release of productive virions from Calu-3 cells (**Fig. 7E**), consistent

with published observations (*7*). These findings indicate that even though SARS-CoV-2 can

enter ACE2-positive macrophages, rapid anti-viral signaling limits viral replication and

release.

DISCUSSION

A key question in SARS-CoV-2 pathogenesis is which type of host cells efficiently sense

SARS-CoV-2 to trigger inflammatory cytokine and anti-viral mediator release. Since

macrophages are prime candidates for sensing and responding to SARS-CoV-2 (*11*), a second

unresolved question is whether SARS-CoV-2 can infect and productively replicate in human

macrophages. This study identifies ACE2 as a critical determinant of macrophage

susceptibility to infection and sensing of infectious SARS-CoV-2.

We demonstrate here that HMDM do not express ACE2 and are thus suitable in vitro models

for ACE2-negative macrophages. While HMDMs phagocytose virus, they do not permit

early-stage viral replication or protein synthesis and do not sense phagocytosed virus to

trigger inflammatory responses. Our observation that SARS-CoV-2 does not replicate in

HMDM or BAL is consistent with multiple reports of abortive SARS-CoV-2 infection of

macrophages in vitro (*12*, *13*, *23*, *24*). While these studies demonstrate that macrophage viral

challenge results in decreasing viral RNA (*12*, *13*, *23*) and viral protein (*24*) levels, and

macrophages fail to produce new infectious virus (*12*, *23*), these reports nevertheless observe

virus entry into macrophages via quantification of SARS-CoV-2 nucleoprotein-positive cells.

This contrasts with studies reporting no viral entry into HMDM (*7*, *25*). We observe virions

within phagosomes (**Fig. 4B**), indicating that SARS-CoV-2 virions can enter the macrophage

phagolysosomal system. Despite this, we propose that the virus does not enter the HMDM

cytoplasm, likely because, without ACE2, the Spike protein does not undergo the necessary

conformational changes for membrane fusion.

ACE2 expression analysis across cell types in healthy and COVID-19 lungs shows that a

small proportion of macrophages express ACE2, corroborating new reports that both ACE2-

positive and -negative macrophages are likely to be present at sites of infection in vivo (*20*,

27, *29*). Age, sex, and co-morbidities may influence cellular ACE2 expression (*44*), as well

as environmental cues such as hypoxia within tissues (*45*, *46*). How macrophage ACE2

expression is regulated has yet to be elucidated. Uncovering the macrophage-specific cues

and pathways regulating ACE2 expression and function will yield further insight into how

ACE2-positive macrophages can be detected and targeted.

In line with our findings herein, other non-epithelial cells that do not usually express ACE2,

such as endothelial cells, are rendered susceptible and permissive to productive SARS-CoV-2

infection by ectopic ACE2 expression (*32*). In THP-1 cells, ectopic ACE2 expression

facilitated SARS-CoV-2 entry and early-stage replication, but in response, these cells

initiated an interferon program that limits new virion release after 24h p.i. (**Fig. 3E, 3F, 6A**).

This limit could reflect the death of infected macrophages, as observed for antibody-

dependent monocyte infection(*47*), or restriction by macrophage anti-viral programs. While

we observe some virus-induced cell death in ACE2-positive macrophages (**Fig. 7A**), our

TBK1 inhibitor experiments (**Fig.s 7C-D**) indicate that a potent macrophage anti-viral

response interrupts the viral replication cycle. Future studies should elucidate the precise anti-

viral programs underpinning macrophage restriction of SARS-CoV-2. Nevertheless, our

findings suggest that ACE2-positive macrophages induce a sufficiently robust anti-viral

response to prevent them from acting as 'Trojan horses' to disseminate SARS-CoV-2 to

extra-respiratory tissues. If such anti-viral signaling programs are compromised, such as in

patients with inborn errors of immunity or circulating IFN autoantibodies(*48*, *49*), then

ACE2-positive macrophages may substantially contribute to increased viral loads and viral

dissemination.

The proportion of macrophages positive for viral RNA in COVID-19 lungs is much higher

than that of ACE2-positive macrophages (*20*, *50*, *51*). Our data suggest that this is not due to

ACE2-independent SARS-CoV-2 entry into and replication within macrophages, in contrast

to a recent study that reported CD169-dependent, ACE2-independent macrophage

infection(*33*). Instead, we expect macrophages positive for viral RNA may be ACE2-negative

macrophages with phagocytosed virions (**Fig. 4B**), extracellular viral RNA, or virus-infected

cells (*52*). Alternatively, ACE-negative macrophages might be infected via an antibody-

dependent uptake route, as recently shown for SARS-CoV-2-infected human monocytes (*47*).

Our data show that the small proportion of macrophages positive for ACE2 can be directly

infected and trigger a distinct inflammatory response to ACE2-negative macrophages

encountering SARS-CoV-2 virions. Importantly, for ACE2-positive macrophages, our data

suggest that the viral RNA present in macrophages in vivo also reflects limited productive

infection. A direct comparison of macrophage responses to these distinct routes of viral

uptake would elucidate whether targeting one subset could maintain protective anti-viral

responses while potentially limiting pathology.

The failure of primary HMDM to respond to SARS-CoV-2 exposure (**Fig. 5A, 5B**) contrasts

with previous reports that SARS-CoV-2 induces the expression of cytokine mRNAs

(including *IL6, CXCL10,* and *TNF*) in HMDM (*12*, *13*, *26*, *53*). Typically, a virus must enter

and replicate within a cell to activate cytosolic PRRs (so-called 'cell intrinsic' sensing),

concomitant with a high threat to that cell. Our data indicate that ACE2-negative

macrophages do not respond directly to SARS-CoV-2 exposure with cytokine production

(**Fig. 5A, 5B**) because lack of virus entry precludes 'cell intrinsic' sensing. Our results thus

agree with published observations that SARS-CoV-2 does not trigger cytokine production

from HMDM (*25*) and that SARS-CoV-2 also fails to trigger the interferon system in human

alveolar macrophages (*24*). In contrast, ACE2-positive macrophages detected actively

replicating SARS-CoV-2 to trigger potent *IFNB1* and *IFNL1* induction (**Fig. 6E**). We

hypothesize that the cytosolic viral sensors retinoic acid-inducible gene-1 (RIG-I) and

MDA5, which detect replicating SARS-CoV-2 RNA in epithelial cells (*7*, *54*, *55*), drive

macrophage anti-viral responses. Macrophage-derived interferon critically orchestrates the

anti-viral immune response to limit disease in other respiratory viral infections (e.g.,

Newcastle Disease Virus, Respiratory Syncytial Virus) (*17*, *18*). We speculate that interferon

produced by ACE2-positive macrophages via cell-intrinsic sensing may also contribute to

early control of SARS-CoV-2 infection in the lung. Alternatively, this same macrophage-

derived interferon could contribute to disease pathology (*56*, *57*).

In addition to cytosolic PRRs that detect serious threats, macrophages are also equipped with

PRRs at other subcellular locations (e.g., cell surface, endosomes) to detect moderate threats.

This 'cell extrinsic' sensing detects neighboring infected cells, viral pathogen-associated

molecular patterns (PAMPs), or host danger-associated molecular patterns (DAMPs) in the

extracellular milieu (*58*). We and others (*24*, *25*) find that HMDM do not sense SARS-CoV-

2, either at the cell surface or in endosomes, to trigger pro-inflammatory responses (**Fig. 5**),

which contradicts reports that SARS-CoV-2 selectively induces a pro-inflammatory response

(e.g., TNF, IL-6, CXCL10) from HMDM (*12*, *13*, *23*). Differences in viral stock preparation

or quantification could underpin these divergent observations. SARS-CoV-2 is usually

cultured in cell lines (e.g., VERO E6 or Calu-3 cells (*12*)) that inherently respond to infection

by producing cytokines or undergoing cell death. Indeed, virus is often harvested at a time

point when a visible cytopathic effect emerges. Individual viral preparations quantified for

intact virions by $TCID_{50}$ or plaque assays may also contain free viral RNA, viral proteins,

non-infectious virions, and parental cell line-derived DAMPs and cytokines (*59*). Thus,

published macrophage responses to viral preparations could represent an indirect response to

epithelial DAMPs or free viral PAMPs. Nevertheless, our viral stocks did not trigger cytokine

release from HMDM or THP-1 cells in the absence of ACE2, indicating that ACE-dependent

entry and early-stage replication are necessary for macrophage sensing (**Fig. 5A, 5B, Fig.**

6A). We anticipate this observation will also apply to ACE2-dependent SARS-CoV-2

variants, including Omicron (BA.4 and BA.5 lineages) and, indeed, any beta-coronaviruses

that utilize ACE2. Consistent with this, SARS-CoV-1, which also requires ACE2 for entry,

does not trigger macrophage cytokine responses (*60*).

This study reveals that human macrophages do not sense SARS-CoV-2 unless they express

ACE2 and support new viral synthesis. Our findings give new insight into SARS-CoV-2 cell

tropism and the influence therein of macrophage innate immune pathways. Such studies of

innate immune cell-intrinsic and -extrinsic SARS-CoV-2 recognition and response will help

reveal novel therapeutic targets for new drugs that dampen pathogenic pro-inflammatory

signaling in virulent viral infections without impeding host anti-viral defense.

MATERIALS AND METHODS

Re-analysis of human lung scRNA-seq

We reanalyzed three lung scRNA-seq datasets representing 1) samples from control or

COVID-19- infected individuals (*27*), 2) samples from COVID-19-infected individuals (*20*),

and 3) samples from control individuals (*28*).

To compare matched cell types across datasets, we used Seurat v 4.1.1(*61*) integration and

label transfer analysis to impose the 'fine level' cell identities defined by Melms et al. onto

the remaining datasets. We first applied the Seurat CCA integration pipeline to integrate the

individuals from Melms et al. before label transfer analysis. Functions *NormalizeData* and

FindVariableFeatures were run on each individual, and *SelectIntegrationFeatures* was run

with *nfeatures* set to 1000. *FindIntegrationAnchors* and *IntegrateData,* followed by

ScaleData, and *RunPCA,* were all run with default parameters. We then transferred the cell

type identities from Melms et al. to the remaining datasets using the *FindTransferAnchors*

and *TransferData* functions with dims = 1:50.

Reagents and inhibitors

Low molecular weight Poly I:C (Invivogen, tlrl-picw) was transfected into cells with

lipofectamine LTX (Thermofisther, A12621) at 0.5 μg pIC per well, while R848 Resiquimod

(Miltenyi Biotech, 130-109-376) was used at a final concentration of 250 ng/ml. The TBK1

inhibitor BX-795 (Sigma-Aldrich, SML0694) was used at a final concentration of 5 μM.

Remdesivir (Gilead) at a final concentration of 10 μM.

Cells

Peripheral blood mononuclear cells were isolated from buffy coats by density centrifugation

using Ficoll-Paque Plus (GE Healthcare). CD14+ monocytes were subsequently isolated

using magnetic-activated cell sorting (Miltenyi Biotech) according to the manufacturer's

instructions. Human macrophages were differentiated from human CD14+ monocytes as

previously described (*62*) and then used for experiments on day 7 of differentiation. Human

monocyte-derived macrophages (HMDMs) were cultured in media consisting of RPMI 1640

medium (Life Technologies) supplemented with 10% heat-inactivated fetal bovine serum

(FBS), 2 mM GlutaMAX (Life Technologies), 50 U/ml penicillin-streptomycin (Life

Technologies) and 150 ng/ml recombinant human macrophage colony-stimulating factor

(CSF-1, endotoxin-free, expressed and purified by the University of Queensland Protein

Expression Facility). HMDM were seeded 16 h prior to experiments at 500 000 cells per well

in 12-well plates or 200 000 cells per well in 24-well plates. Studies using primary human

cells were approved by the University of Queensland Human Medical Research Ethics

Committee. THP-1 cells (TIB-202; ATCC) were maintained in RPMI 1640 medium

supplemented with 10% heat-inactivated fetal bovine serum (FBS), 2 mM GlutaMAX (Life

Technologies) and 50 U/ml penicillin-streptomycin. For experiments, THP-1 cells were

seeded at 500 000 cells per well of a 12-well plate and differentiated for 48 h with 30 ng/ml

phorbol-myristate-acetate (PMA). Calu-3 cells purchased from ATCC (HTB-55) were

maintained in Minimal Essential Media (Invitrogen) containing 10% heat-inactivated fetal

bovine serum (Cytiva), 50 U/ml penicillin, and streptomycin (Life Technologies Australia),

and then seeded at 300 000 cells per well in 12-well plates 48 h prior to experiments. A549

cells (provided by Dr. Chris Macmillan, SCMB, UQ) were maintained in Dulbecco's

Modified Essential Medium (DMEM; Invitrogen) supplemented with 10% heat-inactivated

FBS and 50 U/ml penicillin-streptomycin.

Patient bronchoalveolar lavage (BAL) was obtained at the time of diagnostic bronchoscopy

as previously described (63). Briefly, the bronchoscope was wedged into a non-dependent

sub-segmental bronchus (64) of a radiologically normal segment of the lung, and 20 ml of

normal saline was instilled, retrieved, and then discarded to clear the bronchoscope of

bronchial secretions. A further 80-100 ml was instilled in 20 ml aliquots and retrieved via

hand aspiration of the syringe. Studies using primary human cells were approved by the

Royal Melbourne Hospital and the University of Melbourne Human Research Ethics

Committees. BAL was filtered, and cells were washed and seeded at $1x10^6$ cells per ml

overnight in 48-well tissue culture plates in RPMI 1640 medium supplemented with 10%

heat-inactivated fetal bovine serum (FBS), two mM GlutaMAX (Life Technologies) and 50

U/ml penicillin-streptomycin (Life Technologies). Non-adherent cells were removed via

media change four hours post-seeding, resulting in a >90% macrophage population, as

described previously (65).

Lentiviral transduction

A lentiviral construct containing human ACE2 (Addgene 155295) or mScarlet (Addgene

85044) was cloned into pLV-CMV-MCS-IRES-Puro-Sin (66) and packaged into lentivirus in

HEK-293T cells using third-generation lentiviral packaging plasmids (67). Lentivirus-

containing supernatant was harvested on days 2 and 3 after transfection and then concentrated

using a Lenti-X concentrator (Clontech, 631232). HEK-293T cells were transfected with the

expression vectors according to the manufacturer's protocol with PEI 2500 (BioScientific),

then transduced target THP-1 cells were selected with puromycin (1 µg/mL) after 24 h and

used for assays after 72 h.

Viruses and cell infections

SARS-CoV-2 isolate hCoV-19/Australia/QLD02/2020 was provided by Queensland Health

Forensic & Scientific Services, Queensland Department of Health. Virus was grown on Vero

E6 TMPRSS2 cells for 48 h in DMEM with 2% FBS, and cell debris was cleared by

centrifugation at 500G for 5 minutes at room temperature. Virus was titered as described

previously by plaque assay (*32*). Sanger sequencing was used to confirm that no mutations

occurred in the spike gene relative to the original clinical isolate. Cells (HMDM, THP-1,

Calu-3) in 12-well plates (500 000 cells/well) were challenged for 1 h at 37°C and 5% CO_2

with 2.5 x 10^6 plaque-forming units (PFUs) for MOI 5, or 2.5 x 10^5 PFUs for MOI 0.5. For

Calu-3 cells, virus was added to cells for a total volume of 500 μL of RPMI 1640 with 2%

FBS (HMDM and THP-1) or MEM with 2% FBS (Calu-3) per well. The viral inoculum was

removed, and the medium was replaced with DMEM (Invitrogen) or MEM (Invitrogen)

containing 2% FBS. Alternatively, virus was not removed from the cells. All studies with

SARS-CoV-2 were performed under physical containment 3 (PC3) conditions and were

approved by The University of Queensland Biosafety Committee (IBC/374B/ SCMB/2020,

IBC/518B/IMB/SCMB/2022) and the University of Melbourne Institutional Biosafety

Committee in consultation with the Doherty Institute High Containment Facility

Management Group.

For studies involving SARS-CoV-2 infection of BAL macrophages**,** the SARS-CoV-2 isolate

hCoV-19/Australia/VIC01/2020 (kindly provided by the Victorian Infectious Diseases

Reference Laboratory) was grown in Vero cells for 72 h in serum-free MEM with 1 μg/ml

TPCK trypsin. The median tissue culture infectious dose ($TCID_{50}$) was calculated using the

Reed-Muench method. BAL macrophages (approx. 2.5 x 10^5 cells) were seeded overnight in

48-well plates. Macrophages were infected with SARS-CoV-2 (MOI 1) in serum-free media

(RPMI supplemented as described above) for 1 h. The inoculum was removed, and cells were

washed before the media was replaced with 400 μl serum-free media for a further 2 to 48 h.

VERO control cells were infected at an MOI of 0.5, and maintenance media included 1 μg/ml

of TPCK trypsin. RNA from cell-free supernatant was collected and extracted using a

QIAamp Viral RNA Mini Kit (Qiagen), while RNA from cell monolayers was extracted

using the Rneasy Plus Mini Kit (Qiagen).

Cell Death

THP-1 were seeded at 50 000 cells per well of a black-walled, clear-bottomed 96-well plate (CLS3916, Corning). After 48 h of differentiation with PMA, cells were inoculated with virus at MOI 5 and incubated for one h before the media was replaced with 100 µl of fresh RPMI with 2% FCS. After 72 h, ~60 µl of media was removed per well, and 30 µl of ATPlite substrate solution (Perkin Elmer) was added to the remaining 30 µl. After 10 minutes of incubation at room temperature, plates were sealed with Optical Adhesive Film (Thermofisher), and luminescence was read on a Victor Nivo Plate Reader (Perkin Elmer).

RNA extraction and qPCR analysis

Cells were lysed in Buffer RLT plus β-mercaptoethanol, and the RNA was directly processed using a Rneasy Mini Kit (Qiagen) with on-column DNase digestion according to the manufacturer's instructions. RNA concentration was measured using a NanoDrop spectrophotometer, with equal starting concentrations of RNA for each sample used for reverse transcription. Reverse transcription was performed using Superscript III Reverse Transcriptase (ThermoFisher) with random hexamer priming. Quantitative PCR was performed using SYBR green reagent (Applied Biosystems) on a QuantStudio 7 Flex Real-Time PCR System (ThermoFisher) in 384-well plates (Applied Biosystems), and relative gene expression was determined using the change-in-threshold ($2^{-\Delta\Delta CT}$) method, using Hypoxanthine Phosphoribosyltransferase 1 (*HPRT*) as an endogenous control. Alternatively, gene expression was determined relative to a standard curve generated from plasmids containing SARS-CoV-2 Main Protease (Mpro). Primers are as follows: *HPRT* F*:* TCAGGCAGTATAATCCAAAGATGGT R: AGTCTGGCTTATATCCAACACTTCG; *ACE2* F: TCACGATTGTTGGGACTCTGC, R: TCGCTTCATCTCCCACCACT; *IL6* F: CTCAGCCCTGAGAAAGGAGACAT, R: TCAGCCATCTTTGGAAGGTTCA; *TNF* F: TGCCTGCTGCACTTTGGAGTGA, R: AGATGATCTGACTGCCTGGGCCAG; *IL1B* F: GAAGCTGATGGCCCTAAACA, R: AAGCCCTTGCTGTAGTGGTG, *IFNB1* F: CAGTCCTGGAAGAAAAACTGGAGA, R: TTGGCCTTCAGGTAATGCAGAA; *CXCL10* F: TGAAAGCAGTTAGCAAGGAAAGGT, R: AGCCTCTGTGTGGTCCATCC; *IFNL1* F: CGCCTTGGAAGAGTCACTCA, R: GAAGCCTCAGGTCCCAATTC; SARS-CoV-2 *Mpro* F: GAGACAGGTGGTTTCTCAATCG, R: ACGGCAATTCCAGTTTGAGC; SARS-CoV-2 *E* gene F: ACAGGTACGTTAATAGTTAATAGCGT, R:

ATATTGCAGCAGTACGCACACA. *Subgenomic E* gene: F:

AATATTGCAGCAGTACGCACACA, R: CGATCTCTTGTAGATCTGTTCTC

Copy number of the SARS-CoV-2 E gene was measured in BAL or VERO cellular RNA

(equal starting concentration of RNA) and in RNA from cell culture media (equal starting

volume). SensiFAST™ Probe Lo-ROX One-Step Kit with E-specific probe (FAM-

ACACTAGCCATCCTTACTGCGCTTCG-QQA) was used to detect the E gene, and a

plasmid-based standard curve was used to quantify the number of E copies. The following

primers were used to amplify the Mpro sequence from cDNA with the Phusion polymerase

kit (New England BioLabs). F:AATAAGGTACCAGTGGTTTTAGAAAAATGG,

R:TTATTGCGGCCGCTCATTGGAAAGTAACACC.

The Mpro expression vector was generated by cloning the Mpro PCR product into a

modified pEF6 plasmid, with an HA-tag N-terminal of the multiple cloning site, by standard

restriction digest cloning techniques. The vector and correct insertion of Mpro were verified

by Sanger sequencing.

Cytokine analysis

Cytokine titers were determined using an AlphaLISA Immunoassay kit (Perkin Elmer)

according to the manufacturer's instructions and analyzed on a Victor Nivo Plate Reader

(Perkin Elmer).

Immunofluorescence

Cells were cultured on 12 mm glass coverslips and washed with $PBS_{+/+}$ (PBS supplemented

with 1mM CaCl2 and 0.5mM MgCl2, fixed for 30 min in 4% paraformaldehyde (Sigma

Aldrich), and blocked and permeabilized for 30 min with 3% BSA, 0.3% Triton X-100

(Sigma Aldrich) at room temperature. Primary antibodies were incubated in 1.5% BSA for

1.5 h at room temperature, and secondary antibodies conjugated to Alexa fluorophores

(Invitrogen) were incubated in 1.5% BSA for 60 min in the dark at room temperature. Cells

were stained with Phalloidin (PHDN1-A; Jomar Life Research) and NP primary nanobody.

NP primary nanobody was kindly supplied by Dr. Ariel Isaacs, Watterson Laboratory, School

of Chemistry and Molecular Biology, UQ, QLD. After each incubation, cells were washed 3x

with $PBS_{+/+}$ and mounted with ProlongGold + DAPI solution (Cell Signaling Technologies)

onto glass slides. Z-stack image acquisition was performed on a confocal laser scanning

microscope (Zeiss LSM880) using a 40x NA 1.3 water immersion objective or 20x NA 0.8

objective. Raw images were processed and analyzed using Fiji (v1.53q). Images were batch

processed using a custom-made FIJI script (*68*). The image processing included background

subtraction and median filtering; then, the threshold was determined using the MinError

filter. The holes were filled in the binary image, and Analyze Particles was used to detect and

count nuclei and infected cells. Figures were prepared using Adobe Illustrator. For

quantification analysis, five random regions of interest (ROIs) per group were acquired.

Electron Microscopy

Electron microscopy samples were processed using a method adapted from a previous study

(*69*). Briefly, cells were fixed in 2.5% glutaraldehyde (Electron Microscopy Services) in PBS

for 24 h, then post-fixed in 1% osmium (ProSciTech) for one h and contrasted with 1%

aqueous uranyl acetate (Electron Microscopy Services) for one h. Samples were then serially

dehydrated in increasing percentages of ethanol before serial infiltration with LX-112 resin

(Ladd Research) in a Biowave microwave (Pelco). Ultrathin sections were obtained using an

ultramicrotome (UC6:Leica) and further contrasted using Reynold lead post-stain.

Micrographs were acquired using a JEOL 1011 transmission electron microscope at 80 kV,

with a Morada CCD camera (Olympus) utilizing iTEM software.

Immunoblotting

For total cell lysates, cells were washed once with PBS and lysed with RIPA buffer (50 mM

Tris, 150 mM NaCl, one mM EDTA, 1% Triton X-100, 0.1% SDS, 1% sodium deoxycholate,

protease inhibitor, pH 8.0). A Pierce BCA protein assay kit (Thermo Scientific) was used to

equalize protein amounts, wherein SDS sample buffer containing 100 mM DTT (Astral

Scientific) was added, and samples were boiled at 95°C for 10 min to denature proteins.

Proteins were then separated on 4-15% mini protean TGX precast gels (Bio-Rad) in running

buffer (200 mM glycine, 25 mM Tris, 0.1% SDS (pH8.6)), transferred to PVDF or

nitrocellulose membrane (Bio-Rad 1620112) in blot buffer (48 nM Tris, 39 nM glycine,

0.04% SDS, 20% MeOH) and subsequently blocked with 5% (w/v) milk powder or BSA in

Tris-buffered saline with Tween-20 (TBST) for 30 min. Primary antibodies were incubated

overnight at 4 deg C, followed by secondary antibodies linked to horseradish peroxidase

(HRP) (Cell Signalling) or AlexaFluor647 (Invitrogen), and after each step, immunoblots

were washed 3x with TBST. HRP signals were visualized by enhanced chemiluminescence

(ECL) (Bio-Rad) and imaged with a Vilber Fusion Imaging system (Vilber). The

fluorescence signal was detected using the AI600 imager (Amersham). The following

antibodies were used: SARS-CoV-2 Nucleoprotein (Sino Biological), Tubulin (9F3; Cell

Signalling Technology), ACE2 (AF933; RnD Systems), TMPRSS2 (Abcam) and Actin

(8H10D10; Cell Signalling Technology), ACE2 (MA532307, Invitrogen), Calnexin

(ab22595, Abcam).

Statistical Analysis:

Statistics were calculated using GraphPad Prism using tests as indicated in figure legends.

677
678   **Supplementary Materials**
679
Figure S1: Re-analysis of macrophage ACE2 expression in vivo

Figure S2: SARS-CoV-2 does not replicate in HMDM

Figure S3: SARS-CoV-2 replicates in ACE2-A549 cells

Figure S4: Macrophage ACE2 expression potentiates macrophage inflammatory responses to

SARS-CoV-2 at MOI 0.5

Figure S5: BX-795 blocks anti-viral cytokine induction in THP1-ACE2 cells.

REFERENCES
689

1.   M. Hoffmann, H. Kleine-Weber, S. Schroeder, M. A. Mü, C. Drosten, S. Pö, N. Krü, T.
Herrler, S. Erichsen, T. S. Schiergens, G. Herrler, N.-H. Wu, A. Nitsche, S. Pö Hlmann,
SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a
Clinically Proven Protease Inhibitor Article SARS-CoV-2 Cell Entry Depends on ACE2
and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell*. **181**, 1–
10 (2020).

2.   C. B. Jackson, M. Farzan, B. Chen, H. Choe, Mechanisms of SARS-CoV-2 entry into
cells. *Nat. Rev. Mol. Cell Biol.* **0123456789** (2021), doi:10.1038/s41580-021-00418-x.

3.   P. V'kovski, A. Kratzel, S. Steiner, H. Stalder, V. Thiel, Coronavirus biology and
replication: implications for SARS-CoV-2. *Nat. Rev. Microbiol.* **19**, 155–170 (2021).

4.   B. Meng, A. Abdullahi, I. A. T. M. Ferreira, N. Goonawardane, A. Saito, I. Kimura, D.
Yamasoba, P. P. Gerber, S. Fatihi, S. Rathore, S. K. Zepeda, G. Papa, S. A. Kemp, T.
Ikeda, M. Toyoda, T. S. Tan, J. Kuramochi, S. Mitsunaga, T. Ueno, K. Shirakawa, A.
Takaori-Kondo, T. Brevini, D. L. Mallery, O. J. Charles, CITIID-NIHR BioResource
COVID- Collaboration, Genotype to Phenotype Japan (GP-Japan) Consortium
members, Ecuador-COVID19 Consortium, J. E. Bowen, A. Joshi, A. C. Walls, L.
Jackson, D. Martin, K. G. C. Smith, J. Bradley, J. A. G. Briggs, J. Choi, E. Madissoon,
K. Meyer, P. Mlcochova, L. Ceron-Gutierrez, R. Doffinger, S. A. Teichmann, A. J.
Fisher, M. S. Pizzuto, A. de Marco, D. Corti, M. Hosmillo, J. H. Lee, L. C. James, L.
Thukral, D. Veesler, A. Sigal, F. Sampaziotis, I. G. Goodfellow, N. J. Matheson, K.
Sato, R. K. Gupta, Altered TMPRSS2 usage by SARS-CoV-2 Omicron impacts tropism
and fusogenicity. *Nature* (2022), doi:10.1038/s41586-022-04474-x.

5.   Y. J. Hou, K. Okuda, C. E. Edwards, D. R. Martinez, K. H. Dinnon, T. Kato, R. E. Lee,
B. L. Yount, T. M. Mascenik, G. Chen, K. N. Olivier, A. Ghio, V. Longping, S. R.
Leist, L. E. Gralinski, A. Schäfer, H. Dang, S. Nakano, L. Sun, M. L. Fulcher, A.
Livraghi-butrico, N. I. Nicely, M. Cameron, C. Cameron, D. J. Kelvin, A. De, D. M.
Margolis, A. Markmann, L. Bartelt, R. Zumwalt, J. Martinez, S. P. Salvatore, A.
Borczuk, P. R. Tata, A. Kimple, I. Jaspers, W. K. O. Neal, S. H. Randell, C. Richard, R.
S. Baric, SARS-CoV-2 Reverse Genetics Reveals a Variable Infection Gradient in the
Respiratory Tract. *Cell* (2020), doi:10.1016/j.cell.2020.05.042.

6.   D. Blanco-Melo, B. E. Nilsson-Payant, W.-C. Liu, S. Uhl, D. Hoagland, R. Møller, T.
X. Jordan, K. Oishi, M. Panis, D. Sachs, T. T. Wang, R. E. Schwartz, J. K. Lim, R. A.
Albrecht, B. R. Tenoever, Imbalanced host response to SARS-CoV-2 drives
development of COVID-19. *Cell*, 1–10 (2020).

7.   L. G. Thorne, A. Reuschl, L. Zuliani☐Alvarez, M. V. X. Whelan, J. Turner, M.
Noursadeghi, C. Jolly, G. J. Towers, SARS☐CoV☐2 sensing by RIG☐I and MDA5
links epithelial infection to macrophage inflammation. *EMBO J.*, 1–17 (2021).

8.   P. Bost, A. Giladi, Y. Liu, Y. Bendjelal, G. Xu, E. David, R. Blecher-Gonen, M. Cohen,
C. Medaglia, H. Li, A. Deczkowska, S. Zhang, B. Schwikowski, Z. Zhang, I. Amit,
Host-viral infection maps reveal signatures of severe COVID-19 patients. *Cell* (2020),
doi:10.1016/j.cell.2020.05.006.

9. R. L. Chua, S. Lukassen, S. Trump, B. P. Hennig, D. Wendisch, F. Pott, O. Debnath, L.
Thürmann, F. Kurth, M. T. Völker, J. Kazmierski, B. Timmermann, S. Twardziok, S.
Schneider, F. Machleidt, H. Müller-Redetzky, M. Maier, A. Krannich, S. Schmidt, F.
Balzer, J. Liebig, J. Loske, N. Suttorp, J. Eils, N. Ishaque, U. G. Liebert, C. von Kalle,
A. Hocke, M. Witzenrath, C. Goffinet, C. Drosten, S. Laudi, I. Lehmann, C. Conrad, L.
E. Sander, R. Eils, COVID-19 severity correlates with airway epithelium–immune cell
interactions identified by single-cell analysis. *Nat. Biotechnol.* **38**, 970–979 (2020).

10. R. Knoll, J. L. Schultze, J. Schulte-Schrepping, Monocytes and Macrophages in
COVID-19. *Front. Immunol.* **12**, 1–12 (2021).

11. M. Merad, J. C. Martin, Pathological inflammation in patients with COVID-19: a key
role for monocytes and macrophages. *Nat. Rev. Immunol.* **20**, 355–362 (2020).

12. J. Zheng, Y. Wang, K. Li, D. K. Meyerholz, C. Allamargot, S. Perlman, Severe Acute
Respiratory Syndrome Coronavirus 2-Induced Immune Activation and Death of
Monocyte-Derived Human Macrophages and Dendritic Cells. *J. Infect. Dis.* **223**, 785–
795 (2021).

13. D. Yang, H. Chu, Y. Hou, Y. Chai, H. Shuai, A. C. Y. Lee, X. Zhang, Y. Wang, B. Hu,
X. Huang, T. T. T. Yuen, J. P. Cai, J. Zhou, S. Yuan, A. J. Zhang, J. F. W. Chan, K. Y.
Yuen, Attenuated interferon and proinflammatory response in SARS-CoV-2-infected
human dendritic cells is associated with viral antagonism of STAT1 phosphorylation. *J.*
Infect. Dis. **222**, 734–745 (2020).

14. K. R. Short, A. G. Brooks, P. C. Reading, S. L. Londrigan, The fate of influenza A virus
after infection of human macrophages and dendritic cells. *J. Gen. Virol.* **93**, 2315–2325
(2012).

15. J. E. Gern, E. C. Dick, W. M. Lee, S. Murray, K. Meyer, Z. T. Handzel, W. W. Busse,
Rhinovirus enters but does not replicate inside monocytes and airway macrophages. *J.*
Immunol. Baltim. Md 1950. **156**, 621–7 (1996).

16. S. L. Londrigan, L. M. Wakim, J. Smith, A. J. Haverkate, A. G. Brooks, P. C. Reading,
IFITM3 and type I interferons are important for the control of influenza A virus
replication in murine macrophages. *Virology.* **540**, 17–22 (2020).

17. M. Goritzka, S. Makris, F. Kausar, L. R. Durant, C. Pereira, Y. Kumagai, F. J. Culley,
M. Mack, S. Akira, C. Johansson, Alveolar macrophage-derived type I interferons
orchestrate innate immunity to RSV through recruitment of antiviral monocytes. *J. Exp.*
Med. **212**, 699–714 (2015).

18. Y. Kumagai, O. Takeuchi, H. Kato, H. Kumar, K. Matsui, E. Morii, K. Aozasa, T.
Kawai, S. Akira, Alveolar Macrophages Are the Primary Interferon-α Producer in
Pulmonary Infection with RNA Viruses. *Immunity.* **27**, 240–252 (2007).

19. C. Schneider, S. P. Nobs, A. K. Heer, M. Kurrer, G. Klinke, N. van Rooijen, J. Vogel,
M. Kopf, Alveolar Macrophages Are Essential for Protection from Respiratory Failure
and Associated Morbidity following Influenza Virus Infection. *PLoS Pathog.* **10** (2014),
doi:10.1371/journal.ppat.1004053.

20.   T. M. Delorey, C. G. K. Ziegler, G. Heimberg, R. Normand, Y. Yang, Å. Segerstolpe,
D. Abbondanza, S. J. Fleming, A. Subramanian, D. T. Montoro, K. A. Jagadeesh, K. K.
Dey, P. Sen, M. Slyper, Y. H. Pita-Juárez, D. Phillips, J. Biermann, Z. Bloom-
Ackermann, N. Barkas, A. Ganna, J. Gomez, J. C. Melms, I. Katsyv, E. Normandin, P.
Naderi, Y. V. Popov, S. S. Raju, S. Niezen, L. T.-Y. Tsai, K. J. Siddle, M. Sud, V. M.
Tran, S. K. Vellarikkal, Y. Wang, L. Amir-Zilberstein, D. S. Atri, J. Beechem, O. R.
Brook, J. Chen, P. Divakar, P. Dorceus, J. M. Engreitz, A. Essene, D. M. Fitzgerald, R.
Fropf, S. Gazal, J. Gould, J. Grzyb, T. Harvey, J. Hecht, T. Hether, J. Jané-Valbuena,
M. Leney-Greene, H. Ma, C. McCabe, D. E. McLoughlin, E. M. Miller, C. Muus, M.
Niemi, R. Padera, L. Pan, D. Pant, C. Pe'er, J. Pfiffner-Borges, C. J. Pinto, J. Plaisted, J.
Reeves, M. Ross, M. Rudy, E. H. Rueckert, M. Siciliano, A. Sturm, E. Todres, A.
Waghray, S. Warren, S. Zhang, D. R. Zollinger, L. Cosimi, R. M. Gupta, N. Hacohen,
H. Hibshoosh, W. Hide, A. L. Price, J. Rajagopal, P. R. Tata, S. Riedel, G. Szabo, T. L.
Tickle, P. T. Ellinor, D. Hung, P. C. Sabeti, R. Novak, R. Rogers, D. E. Ingber, Z. G.
Jiang, D. Juric, M. Babadi, S. L. Farhi, B. Izar, J. R. Stone, I. S. Vlachos, I. H. Solomon,
O. Ashenberg, C. B. M. Porter, B. Li, A. K. Shalek, A.-C. Villani, O. Rozenblatt-Rosen,
A. Regev, COVID-19 tissue atlases reveal SARS-CoV-2 pathology and cellular targets.
Nature. **595**, 107–113 (2021).

21.   T. T.-H. Wu, K. J. Travaglini, A. Rustagi, D. Xu, Y. Zhang, S. K. Jang, A. Gillich, R.
Dehghannasiri, G. Martinez-Colon, A. Beck, A. J. Wilk, M. Morri, W. L. Trope, J. B.
Shrager, S. R. Quake, C. S. Kuo, J. Salzman, P. S. Kim, C. A. Blish, M. A. Krasnow,
bioRxiv, in press, doi:10.1101/2022.05.10.491266.

22.   E. Sefik, R. Qu, C. Junqueira, E. Kaffe, H. Mirza, J. Zhao, J. R. Brewer, A. Han, H. R.
Steach, B. Israelow, H. N. Blackburn, S. E. Velazquez, Y. G. Chen, S. Halene, A.
Iwasaki, E. Meffre, M. Nussenzweig, J. Lieberman, C. B. Wilen, Y. Kluger, R. A.
Flavell, Inflammasome activation in infected macrophages drives COVID-19 pathology.
Nature. **606**, 585–593 (2022).

23.   M. M. Abdelmoaty, P. Yeapuri, J. Machhi, K. E. Olson, F. Shahjin, V. Kumar, Y. Zhou,
J. Liang, K. Pandey, A. Acharya, S. N. Byrareddy, R. L. Mosley, H. E. Gendelman,
Defining the Innate Immune Responses for SARS-CoV-2-Human Macrophage
Interactions. *Front. Immunol.* **12**, 1–15 (2021).

24.   L. Dalskov, M. Møhlenberg, J. Thyrsted, J. Blay☐Cadanet, E. T. Poulsen, B. H.
Folkersen, S. H. Skaarup, D. Olagnier, L. Reinert, J. J. Enghild, H. J. Hoffmann, C. K.
Holm, R. Hartmann, SARS☐CoV☐2 evades immune detection in alveolar
macrophages. *EMBO Rep.* **21**, 1–10 (2020).

25.   M. A. Niles, P. Gogesch, S. Kronhart, S. Ortega Iannazzo, G. Kochs, Z. Waibler, M.
Anzaghe, Macrophages and Dendritic Cells Are Not the Major Source of Pro-
Inflammatory Cytokines Upon SARS-CoV-2 Infection. *Front. Immunol.* **12**, 1–12
(2021).

26.   Q. Lu, J. Liu, S. Zhao, M. F. Gomez Castro, M. Laurent-Rolle, J. Dong, X. Ran, P.
Damani-Yokota, H. Tang, T. Karakousi, J. Son, M. E. Kaczmarek, Z. Zhang, S. T.
Yeung, B. T. McCune, R. E. Chen, F. Tang, X. Ren, X. Chen, J. C. C. Hsu, M. Teplova,
B. Huang, H. Deng, Z. Long, T. Mudianto, S. Jin, P. Lin, J. Du, R. Zang, T. T. Su, A.
Herrera, M. Zhou, R. Yan, J. Cui, J. Zhu, Q. Zhou, T. Wang, J. Ma, S. B. Koralov, Z.
Zhang, I. Aifantis, L. N. Segal, M. S. Diamond, K. M. Khanna, K. A. Stapleford, P.

Cresswell, Y. Liu, S. Ding, Q. Xie, J. Wang, SARS-CoV-2 exacerbates
proinflammatory responses in myeloid cells through C-type lectin receptors and Tweety
family member 2. *Immunity*. **54**, 1304-1319.e9 (2021).

27. J. C. Melms, J. Biermann, H. Huang, Y. Wang, A. Nair, S. Tagore, I. Katsyv, A. F.
Rendeiro, A. D. Amin, D. Schapiro, C. J. Frangieh, A. M. Luoma, A. Filliol, Y. Fang,
H. Ravichandran, M. G. Clausi, G. A. Alba, M. Rogava, S. W. Chen, P. Ho, D. T.
Montoro, A. E. Kornberg, A. S. Han, M. F. Bakhoum, N. Anandasabapathy, M. Suárez-
Fariñas, S. F. Bakhoum, Y. Bram, A. Borczuk, X. V. Guo, J. H. Lefkowitch, C. Marboe,
S. M. Lagana, A. Del Portillo, E. J. Tsai, E. Zorn, G. S. Markowitz, R. F. Schwabe, R.
E. Schwartz, O. Elemento, A. Saqi, H. Hibshoosh, J. Que, B. Izar, A molecular single-
cell lung atlas of lethal COVID-19. *Nature*. **595**, 114–119 (2021).

28. E. Madissoon, A. Wilbrey-Clark, R. J. Miragaia, K. Saeb-Parsy, K. T. Mahbubani, N.
Georgakopoulos, P. Harding, K. Polanski, N. Huang, K. Nowicki-Osuch, R. C.
Fitzgerald, K. W. Loudon, J. R. Ferdinand, M. R. Clatworthy, A. Tsingene, S. van
Dongen, M. Dabrowska, M. Patel, M. J. T. Stubbington, S. A. Teichmann, O. Stegle, K.
B. Meyer, scRNA-seq assessment of the human lung, spleen, and esophagus tissue
stability after cold preservation. *Genome Biol.* **21**, 1 (2019).

29. C. Muus, M. D. Luecken, G. Eraslan, L. Sikkema, A. Waghray, G. Heimberg, Y.
Kobayashi, E. D. Vaishnav, A. Subramanian, C. Smillie, K. A. Jagadeesh, E. T. Duong,
E. Fiskin, E. Torlai Triglia, M. Ansari, P. Cai, B. Lin, J. Buchanan, S. Chen, J. Shu, A.
L. Haber, H. Chung, D. T. Montoro, T. Adams, H. Aliee, S. J. Allon, Z. Andrusivova, I.
Angelidis, O. Ashenberg, K. Bassler, C. Bécavin, I. Benhar, J. Bergenstråhle, L.
Bergenstråhle, L. Bolt, E. Braun, L. T. Bui, S. Callori, M. Chaffin, E. Chichelnitskiy, J.
Chiou, T. M. Conlon, M. S. Cuoco, A. S. E. Cuomo, M. Deprez, G. Duclos, D. Fine, D.
S. Fischer, S. Ghazanfar, A. Gillich, B. Giotti, J. Gould, M. Guo, A. J. Gutierrez, A. C.
Habermann, T. Harvey, P. He, X. Hou, L. Hu, Y. Hu, A. Jaiswal, L. Ji, P. Jiang, T. S.
Kapellos, C. S. Kuo, L. Larsson, M. A. Leney-Greene, K. Lim, M. Litviňuková, L. S.
Ludwig, S. Lukassen, W. Luo, H. Maatz, E. Madissoon, L. Mamanova, K.
Manakongtreecheep, S. Leroy, C. H. Mayr, I. M. Mbano, A. M. McAdams, A. N.
Nabhan, S. K. Nyquist, L. Penland, O. B. Poirion, S. Poli, C. Qi, R. Queen, D. Reichart,
I. Rosas, J. C. Schupp, C. V. Shea, X. Shi, R. Sinha, R. V. Sit, K. Slowikowski, M.
Slyper, N. P. Smith, A. Sountoulidis, M. Strunz, T. B. Sullivan, D. Sun, C. Talavera-
López, P. Tan, J. Tantivit, K. J. Travaglini, N. R. Tucker, K. A. Vernon, M. H.
Wadsworth, J. Waldman, X. Wang, K. Xu, W. Yan, W. Zhao, C. G. K. Ziegler, Single-
cell meta-analysis of SARS-CoV-2 entry genes across tissues and demographics. *Nat.*
Med. **27**, 546–559 (2021).

30. M. Magnen, R. You, A. A. Rao, R. T. Davis, L. Rodriguez, C. R. Simoneau, L. Hysenaj,
K. H. Hu, T. U. C. Consortium, C. Love, P. G. Woodruff, D. J. Erle, C. M.
Hendrickson, C. S. Calfee, M. A. Matthay, J. P. Roose, A. Sil, M. Ott, C. R. Langelier,
M. F. Krummel, M. R. Looney, Immediate myeloid depot for SARS-CoV-2 in the
human lung (2022), p. 2022.04.28.489942, , doi:10.1101/2022.04.28.489942.

31. C. Blériot, S. Chakarov, F. Ginhoux, Determinants of Resident Tissue Macrophage
Identity and Function. *Immunity*. **52**, 957–970 (2020).

32. L. Schimmel, K. Y. Chew, C. J. Stocks, T. E. Yordanov, P. Essebier, A. Kulasinghe, J.
Monkman, A. F. R. Santos Miggiolaro, C. Cooper, L. Noronha, K. Schroder, A. K.

Lagendijk, L. I. Labzin, K. R. Short, E. J. Gordon, Endothelial cells are not productively
infected by SARS☐CoV☐2. *Clin. Transl. Immunol.* **10**, 1–18 (2021).

33. S. Jalloh, J. Olejnik, J. Berrigan, A. Nisa, E. L. Suder, H. Akiyama, M. Lei, S.
Ramaswamy, S. Tyagi, Y. Bushkin, E. Mühlberger, S. Gummuluru, CD169-mediated
restrictive SARS-CoV-2 infection of macrophages induces pro-inflammatory responses.
PLOS Pathog. **18**, e1010479 (2022).

34. L. I. Labzin, M. Bottermann, P. Rodriguez-Silvestre, S. Foss, J. T. Andersen, M.
Vaysburd, D. Clift, L. C. James, Antibody and DNA sensing pathways converge to
activate the inflammasome during primary human macrophage infection. *EMBO J.* **38**,
e101365 (2019).

35. S. S. Diebold, T. Kaisho, H. Hemmi, S. Akira, C. Reis E Sousa, Innate Antiviral
Responses by Means of TLR7-Mediated Recognition of Single-Stranded RNA. *Science.*
303, 1529–1531 (2004).

36. M. Weber, A. Gawanbacht, M. Habjan, A. Rang, C. Borner, A. M. Schmidt, S.
Veitinger, R. Jacob, S. Devignot, G. Kochs, A. García-Sastre, F. Weber, Incoming RNA
virus nucleocapsids containing a 5′-triphosphorylated genome activate RIG-I and
antiviral signaling. *Cell Host Microbe.* **13**, 336–346 (2013).

37. C. Lucas, P. Wong, J. Klein, T. B. R. Castro, J. Silva, M. Sundaram, M. K. Ellingson, T.
Mao, J. E. Oh, B. Israelow, T. Takahashi, M. Tokuyama, P. Lu, A. Venkataraman, A.
Park, S. Mohanty, H. Wang, A. L. Wyllie, C. B. F. Vogels, R. Earnest, S. Lapidus, I. M.
Ott, A. J. Moore, M. C. Muenker, J. B. Fournier, M. Campbell, C. D. Odio, A.
Casanovas-Massana, A. Obaid, A. Lu-Culligan, A. Nelson, A. Brito, A. Nunez, A.
Martin, A. Watkins, B. Geng, C. Kalinich, C. Harden, C. Todeasa, C. Jensen, D. Kim,
D. McDonald, D. Shepard, E. Courchaine, E. B. White, E. Song, E. Silva, E. Kudo, G.
DeIuliis, H. Rahming, H. J. Park, I. Matos, J. Nouws, J. Valdez, J. Fauver, J. Lim, K. A.
Rose, K. Anastasio, K. Brower, L. Glick, L. Sharma, L. Sewanan, L. Knaggs, M.
Minasyan, M. Batsu, M. Petrone, M. Kuang, M. Nakahata, M. Campbell, M. Linehan,
M. H. Askenase, M. Simonov, M. Smolgovsky, N. Sonnert, N. Naushad, P.
Vijayakumar, R. Martinello, R. Datta, R. Handoko, S. Bermejo, S. Prophet, S.
Bickerton, S. Velazquez, T. Alpert, T. Rice, W. Khoury-Hanold, X. Peng, Y. Yang, Y.
Cao, Y. Strong, R. Herbst, A. C. Shaw, R. Medzhitov, W. L. Schulz, N. D. Grubaugh,
C. Dela Cruz, S. Farhadian, A. I. Ko, S. B. Omer, A. Iwasaki, Longitudinal analyses
reveal immunological misfiring in severe COVID-19. *Nature.* **584**, 463–469 (2020).

38. M. Koutsakos, L. C. Rowntree, L. Hensen, B. Y. Chua, C. E. van de Sandt, J. R. Habel,
W. Zhang, X. Jia, L. Kedzierski, T. M. Ashhurst, G. H. Putri, F. Marsh-Wakefield, M.
N. Read, D. N. Edwards, E. B. Clemens, C. Y. Wong, F. L. Mordant, J. A. Juno, F.
Amanat, J. Audsley, N. E. Holmes, C. L. Gordon, O. C. Smibert, J. A. Trubiano, C. M.
Hughes, M. Catton, J. T. Denholm, S. Y.-C. Tong, D. L. Doolan, T. C. Kotsimbos, D. C.
Jackson, F. Krammer, D. I. Godfrey, A. W. Chung, N. J.-C. King, S. R. Lewin, A. K.
Wheatley, S. J. Kent, K. Subbarao, J. McMahon, I. Thevarajan, T. H.-O. Nguyen, A. C.
Cheng, K. Kedzierska, Integrated immune dynamics define correlates of COVID-19
severity and antibody responses. *Cell Rep. Med.*, 100208 (2021).

39. L. I. Labzin, M. A. R. Lauterbach, E. Latz, Interferons and inflammasomes:
Cooperation and counterregulation in disease. *J. Allergy Clin. Immunol.* **138**, 37–46
(2016).

40. G. Kokic, H. S. Hillen, D. Tegunov, C. Dienemann, F. Seitz, J. Schmitzova, L. Farnung,
A. Siewert, C. Höbartner, P. Cramer, Mechanism of SARS-CoV-2 polymerase stalling
by remdesivir. *Nat. Commun.* **12**, 279 (2021).

41. C. Junqueira, Â. Crespo, S. Ranjbar, J. Ingber, B. Parry, S. Ravid, L. B. de Lacerda, M.
Lewandrowski, S. Clark, F. Ho, S. M. Vora, V. Leger, C. Beakes, J. Margolin, N.
Russell, L. Gehrke, U. Das Adhikari, L. Henderson, E. Janssen, D. Kwon, C. Sander, J.
Abraham, M. Filbin, M. B. Goldberg, H. Wu, G. Mehta, S. Bell, A. E. Goldfeld, J.
Lieberman, SARS-CoV-2 infects blood monocytes to activate NLRP3 and AIM2
inflammasomes, pyroptosis and cytokine release. *MedRxiv Prepr. Serv. Health Sci.*
(2021), doi:10.1101/2021.03.06.21252796.

42. J. W. Schoggins, Interferon-stimulated genes: Roles in viral pathogenesis. *Curr. Opin.*
Virol. **6**, 40–46 (2014).

43. K. Clark, L. Plater, M. Peggie, P. Cohen, Use of the pharmacological inhibitor BX795
to study the regulation and physiological roles of TBK1 and IκB Kinase ∈:A distinct
upstream kinase mediates ser-172 phosphorylation and activation. *J. Biol. Chem.* **284**,
14136–14146 (2009).

44. S. Beyerstedt, E. B. Casaro, É. B. Rangel, COVID-19: angiotensin-converting enzyme 2
(ACE2) expression and tissue susceptibility to SARS-CoV-2 infection. *Eur. J. Clin.*
Microbiol. Infect. Dis. **40**, 905–919 (2021).

45. S. Joshi, H. Wollenzien, E. Leclerc, Y. P. Jarajapu, Hypoxic regulation of angiotensin-
converting enzyme 2 and Mas receptor in human CD34+ cells. *J. Cell. Physiol.* **234**,
20420–20431 (2019).

46. T. H. Beacon, G. P. Delcuve, J. R. Davie, Epigenetic regulation of ACE2, the receptor
of the SARS-CoV-2 virus. *Genome*. **64**, 386–399 (2021).

47. C. Junqueira, Â. Crespo, S. Ranjbar, L. B. de Lacerda, M. Lewandrowski, J. Ingber, B.
Parry, S. Ravid, S. Clark, M. R. Schrimpf, F. Ho, C. Beakes, J. Margolin, N. Russell, K.
Kays, J. Boucau, U. Das Adhikari, S. M. Vora, V. Leger, L. Gehrke, L. A. Henderson,
E. Janssen, D. Kwon, C. Sander, J. Abraham, M. B. Goldberg, H. Wu, G. Mehta, S.
Bell, A. E. Goldfeld, M. R. Filbin, J. Lieberman, FcγR-mediated SARS-CoV-2 infection
of monocytes activates inflammation. *Nature*. **606** (2022), doi:10.1038/s41586-022-
04702-4.

48. Q. Zhang, P. Bastard, Z. Liu, J. Le Pen, M. Moncada-Velez, J. Chen, M. Ogishi, I. K. D.
Sabli, S. Hodeib, C. Korol, J. Rosain, K. Bilguvar, J. Ye, A. Bolze, B. Bigio, R. Yang,
A. A. Arias, Q. Zhou, Y. Zhang, F. Onodi, S. Korniotis, L. Karpf, Q. Philippot, M.
Chbihi, L. Bonnet-Madin, K. Dorgham, N. Smith, W. M. Schneider, B. S. Razooky, H.-
H. Hoffmann, E. Michailidis, L. Moens, J. E. Han, L. Lorenzo, L. Bizien, P. Meade, A.-
L. Neehus, A. C. Ugurbil, A. Corneau, G. Kerner, P. Zhang, F. Rapaport, Y.
Seeleuthner, J. Manry, C. Masson, Y. Schmitt, A. Schlüter, T. Le Voyer, T. Khan, J. Li,
J. Fellay, L. Roussel, M. Shahrooei, M. F. Alosaimi, D. Mansouri, H. Al-Saud, F. Al-

Mulla, F. Almourfi, S. Z. Al-Muhsen, F. Alsohime, S. Al Turki, R. Hasanato, D. van de
Beek, A. Biondi, L. R. Bettini, M. D'Angio, P. Bonfanti, L. Imberti, A. Sottini, S.
Paghera, E. Quiros-Roldan, C. Rossi, A. J. Oler, M. F. Tompkins, C. Alba, I.
Vandernoot, J.-C. Goffard, G. Smits, I. Migeotte, F. Haerynck, P. Soler-Palacin, A.
Martin-Nalda, R. Colobran, P.-E. Morange, S. Keles, F. Çölkesen, T. Ozcelik, K. K.
Yasar, S. Senoglu, Ş. N. Karabela, C. R. Gallego, G. Novelli, S. Hraiech, Y. Tandjaoui-
Lambiotte, X. Duval, C. Laouénan, A. L. Snow, C. L. Dalgard, J. Milner, D. C. Vinh, T.
H. Mogensen, N. Marr, A. N. Spaan, B. Boisson, S. Boisson-Dupuis, J. Bustamante, A.
Puel, M. Ciancanelli, I. Meyts, T. Maniatis, V. Soumelis, A. Amara, M. Nussenzweig,
A. García-Sastre, F. Krammer, A. Pujol, D. Duffy, R. Lifton, S.-Y. Zhang, G.
Gorochov, V. Béziat, E. Jouanguy, V. Sancho-Shimizu, C. M. Rice, L. Abel, L. D.
Notarangelo, A. Cobat, H. C. Su, J.-L. Casanova, Inborn errors of type I IFN immunity
in patients with life-threatening COVID-19. *Science*. **28**, eabd4570 (2020).

49. P. Bastard, A. Gervais, T. Le Voyer, J. Rosain, Q. Philippot, J. Manry, E. Michailidis,
H. Hoffmann, S. Eto, M. Garcia-Prat, L. Bizien, A. Parra-Martínez, R. Yang, L.
Haljasmägi, M. Migaud, K. Särekannu, J. Maslovskaja, N. de Prost, Y. Tandjaoui-
Lambiotte, C.-E. Luyt, B. Amador-Borrero, A. Gaudet, J. Poissy, P. Morel, P. Richard,
F. Cognasse, J. Troya, S. Trouillet-Assant, A. Belot, K. Saker, P. Garçon, J. G. Rivière,
J.-C. Lagier, S. Gentile, L. B. Rosen, E. Shaw, T. Morio, J. Tanaka, D. Dalmau, P.-L.
Tharaux, D. Sene, A. Stepanian, B. Megarbane, V. Triantafyllia, A. Fekkar, J. R. Heath,
J. L. Franco, J.-M. Anaya, J. Solé-Violán, L. Imberti, A. Biondi, P. Bonfanti, R.
Castagnoli, O. M. Delmonte, Y. Zhang, A. L. Snow, S. M. Holland, C. Biggs, M.
Moncada-Vélez, A. A. Arias, L. Lorenzo, S. Boucherit, B. Coulibaly, D. Anglicheau, A.
M. Planas, F. Haerynck, S. Duvlis, R. L. Nussbaum, T. Ozcelik, S. Keles, A. A.
Bousfiha, J. El Bakkouri, C. Ramirez-Santana, S. Paul, Q. Pan-Hammarström, L.
Hammarström, A. Dupont, A. Kurolap, C. N. Metz, A. Aiuti, G. Casari, V. Lampasona,
F. Ciceri, L. A. Barreiros, E. Dominguez-Garrido, M. Vidigal, M. Zatz, D. van de Beek,
S. Sahanic, I. Tancevski, Y. Stepanovskyy, O. Boyarchuk, Y. Nukui, M. Tsumura, L.
Vidaur, S. G. Tangye, S. Burrel, D. Duffy, L. Quintana-Murci, A. Klocperk, N. Y.
Kann, A. Shcherbina, Y.-L. Lau, D. Leung, M. Coulongeat, J. Marlet, R. Koning, L. F.
Reyes, A. Chauvineau-Grenier, F. Venet, G. Monneret, M. C. Nussenzweig, R.
Arrestier, I. Boudhabhay, H. Baris-Feldman, D. Hagin, J. Wauters, I. Meyts, A. H.
Dyer, S. P. Kennelly, N. M. Bourke, R. Halwani, N. S. Sharif-Askari, K. Dorgham, J.
Sallette, S. M. Sedkaoui, S. AlKhater, R. Rigo-Bonnin, F. Morandeira, L. Roussel, D. C.
Vinh, S. R. Ostrowski, A. Condino-Neto, C. Prando, A. Bonradenko, A. N. Spaan, L.
Gilardin, J. Fellay, S. Lyonnet, K. Bilguvar, R. P. Lifton, S. Mane, M. S. Anderson, B.
Boisson, V. Béziat, S.-Y. Zhang, E. Vandreakos, O. Hermine, A. Pujol, P. Peterson, T.
H. Mogensen, L. Rowen, J. Mond, S. Debette, X. de Lamballerie, X. Duval, F. Mentré,
M. Zins, P. Soler-Palacin, R. Colobran, G. Gorochov, X. Solanich, S. Susen, J.
Martinez-Picado, D. Raoult, M. Vasse, P. K. Gregersen, L. Piemonti, C. Rodríguez-
Gallego, L. D. Notarangelo, H. C. Su, K. Kisand, S. Okada, A. Puel, E. Jouanguy, C. M.
Rice, P. Tiberghien, Q. Zhang, A. Cobat, L. Abel, J.-L. Casanova, Autoantibodies
neutralizing type I IFNs are present in ~ 4% of uninfected individuals over 70 years old
and account for ~ 20% of COVID-19 deaths. *Sci. Immunol.* **6**, eabl4340 (2021).

50. R. A. Grant, L. Morales-Nebreda, N. S. Markov, S. Swaminathan, M. Querrey, E. R.
Guzman, D. A. Abbott, H. K. Donnelly, A. Donayre, I. A. Goldberg, Z. M. Klug, N.
Borkowski, Z. Lu, H. Kihshen, Y. Politanska, L. Sichizya, M. Kang, A. Shilatifard, C.
Qi, J. W. Lomasney, A. C. Argento, J. M. Kruser, E. S. Malsin, C. O. Pickens, S. B.

Smith, J. M. Walter, A. E. Pawlowski, D. Schneider, P. Nannapaneni, H. Abdala-
Valencia, A. Bharat, C. J. Gottardi, G. R. S. Budinger, A. V. Misharin, B. D. Singer, R.
G. Wunderink, R. A. Grant, L. Morales-Nebreda, N. S. Markov, S. Swaminathan, M.
Querrey, E. R. Guzman, D. A. Abbott, H. K. Donnelly, A. Donayre, I. A. Goldberg, Z.
M. Klug, N. Borkowski, Z. Lu, H. Kihshen, Y. Politanska, L. Sichizya, M. Kang, A.
Shilatifard, C. Qi, J. W. Lomasney, A. C. Argento, J. M. Kruser, E. S. Malsin, C. O.
Pickens, S. B. Smith, J. M. Walter, A. E. Pawlowski, D. Schneider, P. Nannapaneni, H.
Abdala-Valencia, A. Bharat, C. J. Gottardi, G. R. S. Budinger, A. V. Misharin, B. D.
Singer, R. G. Wunderink, A. A. Wagh, A. R. Hauser, A. R. Wolfe, A. Thakrar, A. V.
Yeldandi, A. A. Wang, A. R. Levenson, A. M. Joudi, B. Tran, C. A. Gao, C. Kurihara,
C. J. Schroedl, C. M. Horvath, D. Meza, D. D. Odell, D. W. Kamp, D. R. Winter, E. A.
Ozer, E. D. Shanes, E. T. Bartom, E. J. Rendleman, E. M. Leibenguth, F. Wehbe, G. Y.
Liu, G. T. Gadhvi, H. T. Navarro, J. I. Sznajder, J. E. Dematte, J. Le, J. M. Arnold, J. C.
Du, J. Coleman, J. I. Bailey, J. S. Deters, J. A. Fiala, J. Starren, K. M. Ridge, K.
Secunda, K. Aren, K. L. Gates, K. Todd, L. D. Gradone, L. N. Textor, L. F. Wolfe, L. L.
Pesce, L. A. Nunes Amaral, M. L. Rosenbaum, M. Kandpal, M. Jain, M. A. Sala, M.
Saine, M. Carns, M. J. Alexander, M. J. Cuttica, M. H. Prickett, N. H. Khan, N. S.
Chandel, N. D. Soulakis, O. R. Rivas, P. C. Seed, P. A. Reyfman, P. D. Go, P. H. S.
Sporn, P. R. Cooper, R. Tomic, R. Patel, R. Garza-Castillon, R. Kalhan, R. I. Morimoto,
R. J. Mylvaganam, S. S. Kim, S. W. M. Gatesy, S. Thakkar, S. Ben Maamar, S. H. Han,
S. R. Rosenberg, S. Nozick, S. J. Green, S. R. Russell, T. A. Poor, T. J. Zak, T. A.
Lombardo, T. Stoeger, T. Shamaly, Z. Ren, Circuits between infected macrophages and
T cells in SARS-CoV-2 pneumonia. *Nature*. **590**, 635–641 (2021).

51. P. Bost, A. Giladi, Y. Liu, Y. Bendjelal, G. Xu, E. David, R. Blecher-Gonen, M. Cohen,
C. Medaglia, H. Li, A. Deczkowska, S. Zhang, B. Schwikowski, Z. Zhang, I. Amit,
Host-viral infection maps reveal signatures of severe COVID-19 patients. *Cell*, 1–14
(2020).

52. A. C. Salina, D. dos-Santos, T. S. Rodrigues, M. Fortes-Rocha, E. G. Freitas-Filho, D.
L. Alzamora-Terrel, I. M. Castro, T. F. Fraga da Silva, M. H. de Lima, D. C.
Nascimento, C. M. Silva, J. E. Toller-Kawahisa, A. Becerra, S. Oliveira, D. B. Caetité,
L. Almeida, A. Y. Ishimoto, T. M. Lima, R. B. Martins, F. Veras, N. B. do Amaral, M.
C. Giannini, L. P. Bonjorno, M. I. Lopes, M. N. Benatti, S. S. Batah, R. C. Santana, F.
C. Vilar, M. A. Martins, R. L. Assad, S. C. de Almeida, F. R. de Oliveira, E. Arruda
Neto, T. M. Cunha, J. C. Alves-Filho, V. L. Bonato, F. Q. Cunha, A. T. Fabro, H. I.
Nakaya, D. S. Zamboni, P. Louzada-Junior, R. D. Oliveira, L. D. Cunha, Efferocytosis
of SARS-CoV-2-infected dying cells impairs macrophage anti-inflammatory functions
and clearance of apoptotic cells. *eLife*. **11**, e74443 (2022).

53. M. Zheng, R. Karki, E. P. Williams, D. Yang, E. Fitzpatrick, P. Vogel, C. B. Jonsson, T.
D. Kanneganti, TLR2 senses the SARS-CoV-2 envelope protein to produce
inflammatory cytokines. *Nat. Immunol.* **22**, 829–838 (2021).

54. A. Rebendenne, A. L. Chaves Valadão, M. Tauziet, G. Maarifi, B. Bonaventure, J.
McKellar, R. Planès, S. Nisole, M. Arnaud-Arnould, O. Moncorgé, C. Goujon, SARS-
CoV-2 Triggers an MDA-5-Dependent Interferon Response Which Is Unable To
Control Replication in Lung Epithelial Cells. *J. Virol.* **95**, e02415-20 (2021).

55.   N. G. Sampaio, L. Chauveau, J. Hertzog, A. Bridgeman, G. Fowler, J. P. Moonen, M.
1038   Dupont, R. A. Russell, M. Noerenberg, J. Rehwinkel, The RNA sensor MDA5 detects
1039   SARS-CoV-2 infection. *Sci. Rep.* **11**, 13638 (2021).

1040   56.   A. Broggi, S. Ghosh, B. Sposito, R. Spreafico, F. Balzarini, A. Lo Cascio, N. Clementi,
1041   M. de Santis, N. Mancini, F. Granucci, I. Zanoni, Type III interferons disrupt the lung
1042   epithelial barrier upon viral recognition. *Science*. **369**, 706–712 (2020).

1043   57.   J. Major, S. Crotta, M. Llorian, T. M. McCabe, H. H. Gad, S. L. Priestnall, R.
1044   Hartmann, A. Wack, Type I and III interferons disrupt lung epithelial repair during
1045   recovery from viral infection. *Science*. **369**, 712–717 (2020).

1046   58.   A. Iwasaki, P. S. Pillai, Innate immunity to influenza virus infection. *Nat. Rev. Immunol.*
1047   **14**, 315–328 (2014).

1048   59.   Q. Niu, L. Ma, S. Zhu, L. Li, Q. Zheng, J. Hou, H. Lian, L. Wu, X. Yan, Quantitative
1049   Assessment of the Physical Virus Titer and Purity by Ultrasensitive Flow Virometry.
1050   *Angew. Chem. - Int. Ed.* **60**, 9351–9356 (2021).

1051   60.   O. García-Nicolás, P. V'kovski, F. Zettl, G. Zimmer, V. Thiel, A. Summerfield, No
1052   Evidence for Human Monocyte-Derived Macrophage Infection and Antibody-Mediated
1053   Enhancement of SARS-CoV-2 Infection. *Front. Cell. Infect. Microbiol.* **11**, 1–10
1054   (2021).

1055   61.   Y. Hao, S. Hao, E. Andersen-Nissen, W. M. Mauck, S. Zheng, A. Butler, M. J. Lee, A.
1056   J. Wilk, C. Darby, M. Zager, P. Hoffman, M. Stoeckius, E. Papalexi, E. P. Mimitou, J.
1057   Jain, A. Srivastava, T. Stuart, L. M. Fleming, B. Yeung, A. J. Rogers, J. M. McElrath,
1058   C. A. Blish, R. Gottardo, P. Smibert, R. Satija, Integrated analysis of multimodal single-
1059   cell data. *Cell*. **184**, 3573-3587.e29 (2021).

1060   62.   R. C. Coll, J. R. Hill, C. J. Day, A. Zamoshnikova, D. Boucher, N. L. Massey, J. L.
1061   Chitty, J. A. Fraser, M. P. Jennings, A. A. B. Robertson, K. Schroder, MCC950 directly
1062   targets the NLRP3 ATP-hydrolysis motif for inflammasome inhibition. *Nat. Chem. Biol.*
1063   **15**, 556–559 (2019).

1064   63.   E. Tsantikos, M. Lau, C. M. N. Castelino, M. J. Maxwell, S. L. Passey, M. J. Hansen, N.
1065   E. McGregor, N. A. Sims, D. P. Steinfort, L. B. Irving, G. P. Anderson, M. L. Hibbs,
1066   Granulocyte-CSF links destructive inflammation and comorbidities in obstructive lung
1067   disease. *J. Clin. Invest.* **128**, 2406–2418 (2018).

1068   64.   N. Radhakrishna, M. Farmer, D. P. Steinfort, P. King, A Comparison of Techniques for
1069   Optimal Performance of Bronchoalveolar Lavage. *J. Bronchol. Interv. Pulmonol.* **22**,
1070   300–305 (2015).

1071   65.   J. Q. Davies, S. Gordon, "Isolation and Culture of Human Macrophages BT - Basic
1072   Cell Culture Protocols" in, C. D. Helgason, C. L. Miller, Eds. (Humana Press, Totowa,
1073   NJ, 2005), pp. 105–116.

1074   66.   L. Schimmel, M. van der Stoel, C. Rianna, A. M. van Stalborch, A. de Ligt, M.
1075   Hoogenboezem, S. Tol, J. van Rijssel, R. Szulcek, H. J. Bogaard, P. Hofmann, R. Boon,
1076   M. Radmacher, V. de Waard, S. Huveneers, J. D. van Buul, Stiffness-Induced

Endothelial DLC-1 Expression Forces Leukocyte Spreading through Stabilization of the
ICAM-1 Adhesome. *Cell Rep.* **24**, 3115–3124 (2018).

67. T. Dull, R. Zufferey, M. Kelly, R. J. Mandel, M. Nguyen, D. Trono, L. Naldini, A
Third-Generation Lentivirus Vector with a Conditional Packaging System. *J. Virol.* **72**,
8463–8471 (1998).

68. J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S.
Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D. J. White, V.
Hartenstein, K. Eliceiri, P. Tomancak, A. Cardona, Fiji: an open-source platform for
biological-image analysis. *Nat. Methods*. **9**, 676–682 (2012).

69. M. T. Howes, M. Kirkham, J. Riches, K. Cortese, P. J. Walser, F. Simpson, M. M. Hill,
A. Jones, R. Lundmark, M. R. Lindsay, D. J. Hernandez-Deviez, G. Hadzic, A.
McCluskey, R. Bashir, L. Liu, P. Pilch, H. McMahon, P. J. Robinson, J. F. Hancock, S.
Mayor, R. G. Parton, Clathrin-independent carriers form a high capacity endocytic
sorting system at the leading edge of migrating cells. *J. Cell Biol.* **190**, 675–691 (2010).

1091
1092

ACKNOWLEDGEMENTS

We gratefully acknowledge Dr. Lilian Schimmel for producing the ACE2 and mScarlet

expressing lentiviruses, Dr. Ariel Isaacs for producing the anti-N nanobody, and Dr. Fiona

Wylie for critical reading of this manuscript. This work was supported by National Health

and Medical Research Council (NHMRC) of Australia funding, including fellowships

(1124612 to LL; 1177174 to K. Subbarao; 1156489 to RGP; 2009075 to K. Schroder) and

grants (2010757 to LL and EG; 2013574 to CMN, 1140064 and 1150083 to RGP; 2007979

to KRS; 1184532 to SL; 2009677 to K. Schroder) and the Doherty Institute COVID-19

Agility Fund. The Melbourne WHO Collaborating Centre for Reference and Research on

Influenza is supported by the Australian Government Department of Health.

1103

COMPETING INTERESTS:

KRS is a consultant for Sanofi, Roche, and NovoNordisk. K. Schroder is a co-inventor on

patent applications for NLRP3 inhibitors licensed to Inflazome Ltd, a company

headquartered in Dublin, Ireland. Inflazome is developing drugs that target the NLRP3

inflammasome to address unmet clinical needs in inflammatory disease. K. Schroder served

on the Scientific Advisory Board of Inflazome in 2016–2017 and serves as a consultant to

Quench Bio, USA, and Novartis, Switzerland.

1111

AUTHOR CONTRIBUTIONS

Conception: LIL and SLL

Data acquisition and analysis: LIL, KYC, KE,XW, TE, CJS, JR, RP, HM, BH, TY, CLH, SE,

SF, FLM, DPS, CMN, AKL, EG, RGP, SLL

Funding acquisition: LIL, SLL, K.Schroder

Manuscript original draft: LIL, K.Schroder

Manuscript editing: LIL, K.Subbarao, KRS, SLL, K.Schroder

1119
1120   **DATA AND MATERIALS AVAILABILITY:** All data are available in the main text or the

1121   supplementary materials.

1122

  FIGURE LEGENDS

  Figure 1: ACE2 is not expressed on HMDM or BAL but can be ectopically expressed in

  THP-1 cells.

  A: HMDM and THP-1 cells were analyzed by qPCR for *ACE2* mRNA expression,

  with each data point showing an independent donor or experiment (n = 3). B: HMDM

  were stimulated with IFNβ (10 ng/ml) for six h, and protein extracts were analyzed by

  immunoblot alongside extracts from THP-1 cells (WT, THP-1-ACE2, THP-1-mSc).

  C: BAL macrophages from 3 donors were adhered overnight and lysed. The

  expression of ACE2 in BAL macrophages was analyzed by immunoblot relative to

  loading control (Calnexin). Lysate from A549 cells overexpressing ACE2 was used as

  a positive control.

  Figure 2: ACE2-negative HMDM and BAL do not support SARS-CoV-2 entry or early-

  stage replication.

  A-B: Calu-3 cells (A, B) or HMDM (C,D) were infected as indicated, and the virus

  was left on the cells. Viral RNA isolated from cells was measured by qPCR (A, C), or

  viral protein by immunoblot (B,D). Blots are representative of three independent

  experiments (Calu-3) or three independent donors (HMDM). E-F: BAL macrophages

  (MOI 1) or Vero cells (MOI 0.5) were infected with SARS-CoV-2 for one h before

  the virus was removed, and cell-associated viral RNA (E) or viral RNA released in

  cell-free supernatants (F) was analyzed by qPCR of SARS-CoV-2 E gene at 2- and

  48-h post-infection. Data are mean + SEM, with each point representing an individual

  donor (HMDM, BAL macrophages; n = 5) or independent experiments (Calu-3,

  VERO; n = 2-3). Asterisks indicate significance: $p \leq 0.05$ (*), $p \leq 0.001$ (**), $p \leq$

  0.0001 (***) (two-way ANOVA, Tukey's multiple comparison test).

  Figure 3: ACE2-positive macrophages support SARS-CoV-2 entry, early-stage

  replication and productive virion release.

  A-F: Cells were infected with SARS-CoV-2 at MOI 0.5 or MOI 5, as indicated. After

  one h, the virus inoculum was removed, cells were washed, and cells or supernatants

  were harvested at the indicated times (0 h = immediately after the virus inoculum was

  removed). Cellular viral mRNA was analyzed by qPCR (A-B). Viral N protein at 24 h

1156	was assessed by immunofluorescence staining, scale bar = 30 μm (C), and quantified
1157	relative to the total number of cells (D). Infectious virions released into cell
1158	supernatants were measured by plaque assay (E-F). Data show the mean + SEM of 3-
1159	8 independent experiments, with data points representing individual experiments.
1160	Asterisks indicate significance: $p \leq 0.05$ (*), $p \leq 0.001$ (**), $p \leq 0.0001$ (***) (A-B,
1161	E-F: two-way ANOVA, Tukey's multiple comparison test; D: unpaired t-test, Holm-
1162	Šídák test).
1163	
1164	**Figure 4: Macrophages can take up SARS-CoV-2 independently of ACE2**
1165	A-F: Transmission electron microscopy of indicated cells infected with SARS-CoV-2
1166	(MOI 20) at indicated time points. For low magnification images (A, D) scale bar =
1167	10 μm, and for all other images (B, C, E, F) scale bar = 500 nm.
1168	
1169	**Figure 5: SARS-CoV-2 does not trigger inflammatory responses in ACE2-negative**
1170	**HMDM**
1171	A: HMDM were infected with SARS-CoV-2 (MOI 5) or stimulated with R848 or
1172	transfected pIC for 24 h. Cytokines in cell supernatants were analyzed by alphaLISA.
1173	Each data point represents an individual donor (n = 6-15). Graphs show mean + SEM,
1174	and significance is indicated by asterisks (one-way ANOVA, Dunnett's multiple
1175	comparison test). B: HMDM were challenged with SARS-CoV-2 (MOI 0.5, 5) or
1176	transfected pI:C. Gene expression at 2 and 24 h was analyzed by qPCR. Graphs show
1177	mean + SEM, where each data point represents an individual donor (n = 3), and
1178	asterisks indicate significance: $p \leq 0.05$ (*), $p \leq 0.001$ (**), $p \leq 0.0001$ (***) (two-
1179	way ANOVA, Dunnett's multiple comparison test).
1180	
1181	**Figure 6: ACE2-positive macrophages sense entering or replicating SARS-CoV-2 to**
1182	**drive pro-inflammatory and anti-viral responses**
1183	A-C: THP-1-ACE-2 and THP-1-mSc cells (A,B) or Calu-3 cells (C) were infected
1184	with SARS-CoV-2 at MOI 5. Cells were harvested at the indicated times (0 h =
1185	immediately after the virus inoculum was removed), and gene expression was
1186	quantified by qPCR. Gene expression at each time point is presented relative to the
1187	mock control to show fold-gene induction. Data are mean + SEM of 4 independent
1188	experiments (indicated by individual data points), and significance is indicated by

asterisks: $p \leq 0.05$ (*), $p \leq 0.001$ (**), $p \leq 0.0001$ (***) (two-way ANOVA, Tukey's

multiple comparison test). (D-E). THP-1-ACE2 cells were infected with SARS-CoV-

2 at MOI 5 in the presence of 10 μm remdesivir, and viral RNA (C) or inflammatory

cytokines (D) were measured at 24 h p.i. Data show mean + SEM of 3 independent

experiments (indicated by individual data points), and asterisks indicate significance:

$p \leq 0.05$ (*), $p \leq 0.001$ (**), $p \leq 0.0001$ (***) (two-way ANOVA, Sidak's multiple

comparison test).

1196

1197   Figure 7: Blocking cytokine signaling in ACE2-positive macrophages prolongs

1198   infectious virion release

A: THP-1-ACE-2 and THP-1-mSc cells were infected with SARS-CoV-2 (MOI 5).

Cell death was analyzed by ATPlite assay at 72 h. Data are presented as cell viability

relative to mock infection and are mean + SEM of 3 independent experiments.

Asterisks indicate significance: $p \leq 0.05$ (*), $p \leq 0.001$ (**), $p \leq 0.0001$ (***) (two-

way ANOVA, Tukey's multiple comparison test). B: Schematic of TBK1 (BX-795)

inhibition. C-E: THP-1-ACE-2 or Calu-3 cells were stimulated with SARS-CoV-2 at

MOI 5. After one h, the viral inoculum was removed, and BX-795 was added.

Supernatants were harvested at 72 h, CXCL10 was analyzed by ELISA (C) and viral

titers were analyzed by plaque assay (D, E). Data show the mean + SEM of at least

three independent experiments, with each data point representing a different

experiment. Significance is indicated by asterisks: $p \leq 0.05$ (*), $p \leq 0.001$ (**), $p \leq$

0.0001 (***) (C: one-way ANOVA, Tukey's multiple comparison test; D, E: ratio-

paired t-test).

Figure 1

A

$Ace2/Hprt$

HMDM
Thp1 WT
Thp1 ACE2
Thp1 mSc

B

HMDM   THP-1

IFNβ   D1   D2   D3   D4   WT ACE2 mSc
       - +   - +   - +   - +

120
100   ACE2

50   TMPRSS2

50   β-TUBULIN

C

BAL MF   ACE2 - A549
D1 D2 D3

kDa
120   ACE2

70   Calnexin

Figure 2

Figure 3

Figure 4

HMDM 1h p.i.

HMDM 1h p.i.

THP-1 ACE2 1h p.i.

THP-1 ACE2 1h p.i.

THP-1 ACE2 1h p.i.

Calu3 72h p.i.

Figure 7