Title:

Density-dependent attributes of schooling in small pelagic fishes

Authors:

Sho Furuichi[1]*, Yasuhiro Kamimura[1], Masahiro Suzuki[1], Ryuji Yukami[1]

Affiliations:

[1]Fisheries Resources Institute, Japan Fisheries Research and Education Agency, Yokohama,

Japan

*Correspondence:

Name: Sho Furuichi

Email: sfuru@kyudai.jp

ORCID ID:

Sho Furuichi: 0000-0002-6262-4076

Yasuhiro Kamimura: 0000-0002-4236-2734

Abstract

Many small pelagic fishes obligately form schools; some of these schools attain a remarkable

size. Although the school is a fundamental and important ecological unit and is the site of

biological interactions such as competition and predation, information on schooling processes

in the field remains scarce. Here, we examined the quantitative relationships between

population density and school size, the number of schools, and other school characteristics

(i.e., packing density, volume, and cross-sectional area) in three species of small pelagic

fishes: Japanese anchovy *Engraulis japonicus*, Japanese sardine *Sardinops melanostictus*, and

chub mackerel *Scomber japonicus*. We found that school size increased almost linearly with

population density, whereas the number of schools and other characteristics increased

non-linearly with population density, whereby the rate of increase slowed radically as

population density increased. These results indicate that, at low population densities, an

increase in density causes an increase in both school size and the number of schools, whereas

at higher population densities, an increase in density triggers the formation of larger schools

rather than more schools. Furthermore, we found that the shapes of the relationships of all

school characteristics with population density differed among species. Our results indicate

that the schooling behaviour of small pelagic fishes is density-dependent, and responses to

changes in density are species-specific. Our results provide insight into how biological

interactions such as intra- and inter-specific competition and predator-prey interactions

mediate the density-dependent processes that underlie the population dynamics and

community structure of small pelagic fishes in marine ecosystems.

Keywords

pelagic fish, schooling, group living, density dependence, western North Pacific

Introduction

Group living is a widespread phenomenon across the animal kingdom (Krause and Ruxton 2002; Beauchamp 2014; Ward and Webster 2016), and is particularly important in fishes. Most fish species form groups at some stage of their life history, and fishes display more diverse grouping behaviour and a greater range of group sizes and shapes than other taxa (Pavlov and Kasumyan 2000; Vicsek and Zafeiris 2012).

Among fishes, the grouping behaviour of small pelagic fishes, such as sardines and anchovies, is especially distinctive and ecologically important. Many small pelagic fishes obligately aggregate to form schools, shoals, or swarms (hereafter, simply 'schools', although the terms are not strictly synonymous), and they sometimes form massive schools consisting of thousands or even millions of individuals (Misund 1993; Makris et al. 2006). Small pelagic fishes are also important components of marine ecosystems that link lower trophic-level plankton to upper trophic-level predators (Rice 1995; Cury et al. 2000; Bakun 2006), and the formation of schools by small pelagic fishes is likely to determine population dynamics, community structure, and energy flow in marine ecosystems through processes such as intra- and inter-specific competition and predator–prey interaction (Maury and Poggiale 2013; Maury 2017; Dalziel et al. 2021). Therefore, clarifying how and why school characteristics such as school size in small pelagic fishes vary is necessary to better understand marine ecosystem dynamics.

Population density is one of the most fundamental factors that affect schooling (Niwa 2004; Hensor et al. 2005). An increase in population density leads to a higher encounter rate between conspecific individuals or schools and hence an increased opportunity for school fusion. Therefore, mean school size is expected to increase with population density (Niwa 2004; Hensor et al. 2005). However, the formation of schools involves not only benefits (e.g., avoiding predation) but also costs (e.g., competition for food), meaning that there is likely to

be an optimum school size that allows individuals to maximize their net benefit (Krause and

Ruxton 2002; Ward and Webster 2016), and it has been suggested that individuals adjust their

schooling behaviour toward this optimum (Sibly 1983). Therefore, school sizes might

converge to a stable size or remain constant regardless of population density, rather than

simply increasing in direct proportion to population density (Sibly 1983).

The relationships between population density and schooling are not simple and need to

be verified by field or experimental observations. However, field or experimental studies on

the relationships between these factors in small pelagic fishes remain scarce. One of the few

relevant studies reported that in Atlantic herring (*Clupea harengus*), the spatial size (i.e.,

cross-sectional area) of schools is relatively stable, and only the number of schools increases

as population density increases (Jech and Stroman 2012). On the other hand, for several small

pelagic fishes in European areas, it has been reported that as population density increases, the

proportion of spatially large schools increases (Petitgas et al. 2001) as does the number of

schools (Brierley and Cox 2015). Therefore, in small pelagic fishes, the relationship between

population density and school size remains unclear, and quantitative relationships between

population density and school size and other schooling characteristics have not been well

established.

The purpose of this study is to identify quantitative relationships between population

density and school size, the number of schools, and other schooling characteristics in three

small pelagic fishes: Japanese anchovy *Engraulis japonicus*, Japanese sardine *Sardinops*

melanostictus, and chub mackerel *Scomber japonicus*. These three species inhabit the western

North Pacific around Japan and are important fisheries resources that account for the majority

of total fisheries catch by weight in Japan (Yatsu 2019). The stock abundance of the three

species has fluctuated widely in the last decades (Yatsu 2019), which allows us to observe a

wide range of population densities in the field.

First, to examine whether school characteristics (school size, number of schools, school

packing density, school volume, and school cross-sectional area) increase or remain constant

as population density increases and whether the shapes of the relationships are linear or

non-linear (e.g., saturating), we fitted power-law models to the relationships between school

characteristics and local population density at the feeding grounds of the three small pelagic

fishes. Second, to examine whether the relationships between population density and school

characteristics are different among species, we compared models with species-specific and

species-common relationships.

Materials and methods

Survey

The Japan Fisheries Research and Education Agency has conducted trawl-acoustic surveys to

examine the densities and distributions of small pelagic fishes every year since 2005 between

September and October in the western North Pacific off northern Japan and the Kuril Islands

(Fig. 1). The study area corresponds to the feeding grounds of the study species during the

months surveyed. In this study, we used data from 2007, when the survey design was

established, to 2021, the most recent year available. The specifications of the trawl net are as

follows: opening dimensions, 30 m × 30 m; cod-end mesh aperture, 17.5 mm; tow duration,

15–60 min; ship speed, 6.5–9.3 km h^{-1} (3.5–5.0 kn). We conducted three trawl hauls per day

at depths < 40 m. At each station, fish species were identified on board, and the catch was

weighed by species. Up to 380 individuals of each species were randomly selected from the

trawl catch and stored at –25 °C. Sampled fish were measured to the nearest 0.01 cm fork

length for chub mackerel, and standard length for Japanese sardine and Japanese anchovy in

the laboratory.

Acoustic data were collected during trawling using a Simrad EK60 scientific

echosounder (SIMRAD Kongsberg Maritime AS, Horten, Norway) equipped with a

hull-mounted transducer, at an operating frequency of 38 kHz. The echosounder was

calibrated in accordance with standard procedures before each survey (Foote 1987). During

operation, the pulse duration and ping interval were set to 1.024 ms and 0.7–1.3 s,

respectively.

Conductivity–temperature–depth (CTD) water column profiles were collected at each

station. The CTD profiles were used to determine sound speed (Mackenzie 1981) and

absorption coefficient (Francois and Garrison 1982) for processing of acoustic data.

To analyse schools dominated by one species rather than mixed-species schools, we

used only data corresponding to trawl catches where more than 80% of the trawl catch (by

weight) was composed of one species (Japanese anchovy, Japanese sardine, or chub mackerel),

and where catch exceeded 10 kg. Additionally, because the species of interest tend to disperse

in layers at night, making it very difficult to accurately delineate separate schools (Coetzee

2000), we used only data from daytime trawls.

Acoustic data processing

The analysis of acoustic data (volume backscattering strength [dB re 1 m^{-1}]) was performed

with Echoview (version 11.1; Echoview Software Pty Ltd, Hobart, Australia). To avoid

potential surface noise (e.g., wave generated bubbles) and to account for near-field effects,

acoustic data shallower than 7.5 m were removed. We also removed data deeper than 50 m

because schools of the species of interest rarely extend deeper than 50 m in the study area (the

centres of schools of the species were mainly shallower than 30 m [Table 1]). Additionally,

manual inspection was used to remove any remaining undesired data (e.g., reverberation,

bubble noise). To remove small spikes, two convolution kernels were applied, characterized

by a 3×3 median filter and a 5×5 dilation filter (Fernandes 2009; Aronica et al. 2019). A

minimum threshold of –60 dB was applied to pre-processed echograms to distinguish fishes

from other particulate targets such as plankton (Jech and Michaels 2006).

For subsequent analyses, we calculated the mean target strength (dB re 1 m^2) of each

school and at each station. Target strength indicates the acoustic reflectivity of the ensonified

target (e.g., a fish). The mean target strength was calculated from the mean total length of

catch samples using the following equations. For chub mackerel (Simmonds and MacLennan

2008), the equation is:

$$\overline{TS} = 20 \log_{10} \overline{L_t} - 67.6$$

and for Japanese sardine and Japanese anchovy (Zhao et al. 2008), the equation is:

$$\overline{TS} = 20 \log_{10} \overline{L_t} - \frac{20}{3} \log_{10} \left(1 + \frac{\bar{z}}{10}\right) - 70.9$$

where $\overline{L_t}$ is the mean total length (cm) of a catch sample at a given station. For the mean

target strength at a station, $\bar{z}$ is defined as the mean depth (m) of the distribution of Japanese

sardine or Japanese anchovy at the station, and for the mean target strength in a school, $\bar{z}$ is

defined as the mean depth of the school. The mean depth is the average of all sample depths at

the station or in the school weighted by volume-backscattering coefficient values. Because

body length was measured as fork length (L_f) for chub mackerel, and standard length (L_s) for

Japanese sardine and Japanese anchovy, measurements were converted to total length (L_t) by

using the following formulae (Furuichi et al. 2021): $L_t = 1.07 L_f - 0.136$ for chub

mackerel; $L_t = 1.13 L_s + 0.371$ for Japanese sardine; and $L_t = 1.12 L_s + 0.358$ for

Japanese anchovy.

We estimated local population density (ind. m^{-3}) at each station by using the following

formula: $D = 10^{(\overline{SV}_{station} - \overline{TS})/10}$, where D is the local population density at a station,

$\overline{SV}_{station}$ is the mean volume-backscattering strength of the water column (7.5–50.0 m)

during trawling at the station, and $\overline{TS}$ is the mean target strength at the station (Simmonds

and MacLennan 2008).

Fish schools were detected and isolated from the pre-processed echograms by using the

Shoals Analysis and Patch Estimation System (SHAPES) algorithm, implemented in

Echoview (Barange 1994; Coetzee 2000). In the SHAPES algorithm, schools are detected by

first identifying candidates, which are groups of adjacent pixels that meet minimum criteria

for length and thickness. These individual school candidates are then linked together to form a

larger school candidate if the horizontal and vertical distances between them are less than the

specified maximum linking distances. After all linking has been carried out, schools are

recognized if the final school candidates are larger than the defined minimum total school

length and thickness. The detection parameters we used were as follows: minimum total

school length, 4 m; minimum school height, 2 m; minimum candidate length, 2 m; minimum

candidate height, 1 m; maximum vertical linking distance, 5 m; and maximum horizontal

linking distance, 15 m. This set of parameters was selected based on previous studies and the

resolution of the acoustic data (Coetzee 2000; Fernandes 2009; D'Elia et al. 2014).

All detected schools were classified as Japanese anchovy, Japanese sardine, or chub

mackerel based on the trawl catch composition. We extracted school packing density (number

of individuals m^{-3}), school volume (m^3), and cross-sectional area (m^2) for each school. The

packing density of a school was calculated as $\rho = 10^{(\overline{SV}_{school} - \overline{TS})/10}$, where ρ is the packing

density of a school, $\overline{SV}_{school}$ is the mean volume-backscattering strength of the school, and

$\overline{TS}$ is the mean target strength in the school (Simmonds and MacLennan 2008). The volume

of a school was calculated assuming a cylindrical shape as $V = \sum_i \pi \cdot (T_i/2)^2 \cdot L_i$, where V

is volume, T_i is the thickness of the school at ping i, and L_i is the distance from ping i to

ping $i + 1$. School size (number of individuals) was calculated by multiplying the packing

density by school volume. In addition, we also calculated the number of schools at each

station.

Statistical analysis

We examined relationships between local population density and school characteristics

(school size, number of schools, packing density, school volume, cross-sectional area). To

consider the possible nonlinearity of the relationships, we applied power-law models:

$\mu = a \times x^b$

where μ is the mean value of a school characteristic (school size, number of schools per km,

packing density, school volume, or cross-sectional area), x is local population density, a is

a coefficient (>0), and b is a power exponent. The power exponent b determines the

direction (positive, constant, or negative) and shape (linear or nonlinear) of the relationship. A

negative power exponent ($b < 0$) means a negative relationship between local population

density and a school characteristic, an exponent of zero ($b = 0$) means that a school

characteristic is constant regardless of local population density, and a positive exponent

($b > 0$) means a positive relationship. Additionally, the power exponent also determines the

shape of the relationship when the exponent is greater than zero: an exponent greater than one

($1 < b$) generates a convex relationship (accelerating increase with increasing local

population density), an exponent equal to one ($b = 1$) generates a linear relationship

(proportionality), and an exponent lower than one ($0 < b < 1$) generates a concave

relationship (slowing increase with increasing local population density). The coefficient a

serves as a simple scaling factor, meaning that the shape of the relationship will not change

depending on the coefficient, but its placement will move up or down as the coefficient

increases or decreases, respectively.

For school size, packing density, school volume, and cross-sectional area, we assumed

that observed values follow a Gamma distribution because of their continuous, non-negative,

and right-skewed distributions. Therefore, the models were defined as follows:

$$y_{i,j,k} \sim \text{Gamma}\left(\alpha_i, \beta_{i,j}\right)$$

$$\beta_{i,j} = \alpha_i / \mu_{i,j}$$

$$\mu_{i,j} = a_i \times x_{i,j}^{b_i} \times r_l$$

where $y_{i,j}$ is the observed value of a school characteristic of species i at station j in school

k, α_i is a shape parameter (>0), $\beta_{i,j}$ is a rate parameter (>0), $\mu_{i,j}$ is the expected (mean)

value of the school characteristic, $x_{i,j}$ is local population density, a_i is a coefficient, b_i is a

power exponent, and r_l is a random effect of date l. Factors other than population density

have also been reported to affect schooling behaviour (Borner et al. 2015; Romenskyy et al.

2020; Zheng and Fu 2021), and these factors may cause spatiotemporal autocorrelation in

school characteristics. To consider this spatiotemporal autocorrelation, survey date was

incorporated into the model as a multiplicative random effect following a normal distribution

with mean 1 and standard deviation σ_r.

For the number of schools, we assumed that observed values followed a negative

binomial distribution because the data were count data characterized by a large variance.

Therefore, the model was defined as follows:

244            $y_{i,j} \sim \text{NegativeBinomial}(\lambda_{i,j}, \varphi_i)$

245            $\lambda_{i,j} = d_j \times \mu_{i,j}$

246            $\mu_{i,j} = a_i \times x_{i,j}^{b_i} \times r_l$

where $y_{i,j}$ is the observed number of schools of species i at station j, $\lambda_{i,j}$ is a location

parameter (>0), φ_i is a dispersion parameter (>0), d_j is sampling distance (km), $\mu_{i,j}$ is the

expected (mean) number of schools per km, $x_{i,j}$ is local population density, a_i is a

coefficient, b_i is a power exponent, and r_l is a random effect of date l. To consider any

spatiotemporal autocorrelation, survey date was incorporated into the model as a

multiplicative random effect following a normal distribution with mean 1 and standard

deviation σ_r.

Furthermore, we investigated whether the relationships between local population

density and school characteristics are different or common among species. In addition to the

above models assuming species-specific relationships, we also constructed models assuming

species-common relationships, where the coefficient and power exponent were same among

species, that is, $a_{anchovy} = a_{sardine} = a_{mackerel}$ and $b_{anchovy} = b_{sardine} = b_{mackerel}$. To

compare models, the widely applicable Bayesian information criterion (WBIC) was

calculated for each model (Watanabe 2013).

We fitted the models in a Bayesian framework using Markov chain Monte Carlo

methods via No-U-Turn sampling with the software STAN, which was called from R (R

Development Core Team 2020) using the package 'rstan' (Stan Development Team 2020), to

obtain posterior parameter distributions. We used weak informative priors (Cauchy or

half-Cauchy distribution with location parameter 0 and scale parameter 5) for all parameters

except σ_r. For σ_r, a normal distribution with mean 0 and standard deviation 0.2 was used as

the prior distribution because of the difficulty in converging. We used four chains, each of

which had 40,000 iterations including a burn-in of 10,000 iterations with a thinning interval of

10, resulting in 12,000 values for the posterior distribution of each parameter. Convergence of

Markov chain Monte Carlo algorithms was checked by using the potential scale reduction

statistic ($\hat{R}$) and effective sample sizes (Carpenter et al. 2017); for all parameters, the $\hat{R}$ was

less than 1.01 and effective sample sizes were greater than 400.

Results

In total, data from 88 stations and 518 schools were obtained (Table 1). A wide range of local

population density (ind. m^{-3}) was observed ranging from 1.18×10^{-4} to 0.41 for Japanese

anchovy, from 1.26×10^{-4} to 0.58 for Japanese sardine, and from 4.6×10^{-6} to 0.016 for chub

mackerel.

School size

For all species, school size tended to increase as local population density increased, and the

power exponents were greater than zero, indicating positive relationships (Figs 2a and 3, and

Table S1). In each case, mean school size increased almost linearly with local population

density (Fig. 3). The estimated power exponents were relatively close to one (although

slightly lower in chub mackerel than other species), but the credible intervals did not include

one (Fig. 2a and Table S1). This indicates that school size increased almost linearly with local

population density, but the rate of increase slowed slightly as local population size increased.

The model with species-specific relationships between school size and population density had

a lower WBIC than the model with species-common relationships (species-specific, 37.8;

species-common, 60.6), indicating that the relationships differed among species (Fig. S1a).

Number of schools

For all species, the number of schools per km tended to increase as local population density

increased, but the rate of increase slowed with increasing local population density (Fig. 4).

For all species, the power exponents were greater than zero (indicating positive relationships)

but much less than one (ranging from 0.31 to 0.43; Fig. 2b and Table S1), indicating that the

rate of increase of the number of schools slowed considerably with increasing local

population density (Fig. 4). The model with species-specific relationships between the number

of schools and population density had a lower WBIC than the model with species-common

relationships (species-specific, 222.4; species-common, 225.7), indicating that the

relationships differed among species (Fig. S1b).

School packing density

In all species, packing density tended to increase as local population density increased, but the

rate of increase slowed with increasing population density (Fig. 5). For all species, the power

exponents were greater than zero (indicating positive relationships) but much less than one

(ranging from 0.30 to 0.40; Fig. 2c and Table S1), indicating that the rate of increase of

packing density slowed considerably with increasing local population (Fig. 5). The model

with species-specific relationships between packing density and population density had a

lower WBIC than the model with species-common relationships (species-specific, 456.1;

species-common, 460.2), indicating that the relationships differed among species (Fig. S1c).

Spatial extent of a school

For all species, school volume tended to increase with increasing local population density, but

the rate of increase slowed as population density increased (Fig. 6). For all species, the power

exponents were greater than zero (indicating positive relationships) but much less than one

(ranging from 0.22 to 0.40; Fig. 2d and Table S1), indicating that the rate of increase of school

volume slowed considerably with increasing local population (Fig. 6). The model with

species-specific relationships between school volume and population density had a lower

WBIC than the model with species-common relationships (species-specific, 208.1;

species-common, 210.4), indicating that the relationships differed among species (Fig. S1d).

In school cross-sectional area, for all species, the power exponents were greater than

zero (indicating positive relationships) but much less than one (ranging from 0.13 to 0.26; Fig.

2e and Table S1), indicating that the rate of increase of cross-sectional area slowed

considerably with increasing local population (Fig. 7). The model with species-specific

relationships between cross-sectional area and population density had a lower WBIC than the

model with species-common relationships (species-specific, 690.7; species-common, 693.1),

indicating that the relationships differed among species (Fig. S1e).

Discussion

The effects of population density on school size and the number of schools is important for

understanding population dynamics, community structure, and energy flow in marine

ecosystems because the school is a fundamental ecological unit in which biological

interactions such as competition and predation take place (Maury and Poggiale 2013; Maury

2017). However, information on the relationships between population density and school

characteristics among small pelagic fishes in the field remains scarce, and quantitative

relationships are yet to be clarified. In the present study, we show that school size increases

almost linearly with population density, whereas the number of schools increases non-linearly

in such a way that the rate of increase slows considerably with increasing population density.

Our study provides important information on the quantitative relationships between

population density and school characteristics.

Our results are largely consistent with the predictions of a theoretical model based on

simple assumptions of encounter, merge, and split processes (Niwa 2004). In the model,

schools are assumed to move at random through their environment. Schools merge whenever

they encounter another school, and each school splits periodically into two schools. The

schooling process of the studied species may resemble this simple model. However, there is a

slight difference between our results and the predictions of the model. Whereas the model

predicts that mean school size will increase linearly with population density, our results show

that the rate of increase of mean school size slowed slightly with increasing local population

size (i.e., a near-linear relationship), implying the existence of an optimal (stable) school size.

The reason for this difference is likely that the theoretical model is purely mechanistic and

does not consider the functions (i.e., the ultimate factors) of schooling behaviour. The

formation of schools involves not only benefits but also costs, leading to the existence of an

optimal group size that allows individuals to maximize their net benefit (Krause and Ruxton

2002; Ward and Webster 2016), and it has been suggested that individuals adjust their

schooling behaviour to increase their net benefit (Sibly 1983). Accordingly, in our study

species, the probability of school merging and splitting likely changed with school size in a

way that served to increase this net benefit (Gueron and Levin 1995; Niwa 1998). A

combination of mechanistic and functional approaches will be needed in future studies to

better understand the dynamics of school size (Sumpter 2010; Griesser et al. 2011).

Our results show that whereas at low population densities, increasing density resulted in

both more and larger schools, at high population densities, increasing density resulted in

larger schools rather than more schools. This can be explained by the following mechanism:

at low population densities, the frequency of encounters with conspecific schools is low, and

hence many small schools are formed, whereas at high population densities, the frequency of

encounter with conspecific schools is high, and as many schools merge to form larger schools,

the number of schools becomes relatively stable. Similar trends have also been found in

freshwater cyprinid fishes and terrestrial mammals, which form much smaller groups than

marine small pelagic fishes (Southwell 1984; Pépin and Gerard 2008; Holubová et al. 2019).

This suggests that this pattern might be widespread in animals that form fluid groups.

Our results also show that school packing density increased with increasing population

density. This suggests that school size and packing density increase simultaneously as

population densities increase. In fact, school size and packing density were strongly correlated

(Japanese anchovy: $\rho = 0.79$, $p < 0.001$; Japanese sardine: $\rho = 0.82$, $p < 0.001$; chub mackerel:

$\rho = 0.75$, $p < 0.001$). For small to mid-sized schools, predation risk per individual is expected

to decrease as school size increases (Lima 1995; Krause and Ruxton 2002; Caro 2005). For

large schools, however, as school size increases, the risk of predator attacks also increases as

the school becomes more visible to predators (Pitcher and Parrish 1993; Maury 2017). Denser

schools can engage in highly coordinated antipredator reactions through the efficient

transmission of information within schools (Rieucau et al. 2014a, b), and therefore, the

formation of dense schools could be a response of individuals against the threat of predation.

Small pelagic fishes form especially massive schools, and individuals might cope with the

increased risk of predator attacks at these large school sizes by forming dense and compact

schools that allow them to efficiently obtain information from other schooling individuals.

In small pelagic fishes, a strong density-dependent decline in body growth is often

observed during times of increasing population density (Wada et al. 1995; Watanabe and

Yatsu 2004; Canales et al. 2020; Kamimura et al. 2021b). Density-dependent growth

reduction is mainly caused by intraspecific competition for food (Hansen et al. 1999; Jenkins

et al. 1999; Post et al. 1999; Kamimura et al. 2021b). Food competition can occur between

schools and within a school, but within-school competition is likely to be stronger and more

direct than between-school competition. Several studies in animals that form groups (although

none of the studies were conducted on fish) have reported that competition for food within

groups is more intense than competition between groups (Janson 1985, 1988). The increase in

school size and packing density with increasing population density observed in our study is

likely to create fairly strong competition for food. This might be one of the reasons for the

strong negative density-dependence of growth observed in small pelagic fishes.

In our study, the shapes of the relationships between population density and school

characteristics differed among small pelagic fishes (although the overall trends were similar

across species). In krill, which are similar to small pelagic fishes in that they often form

massive swarms and are preyed upon by a wide variety of pelagic megafauna, it has been

reported that increasing population density causes a linear increase in the number of schools,

but does not affect school size (Brierley and Cox 2015). This is different from the

relationships observed in small pelagic fishes in our study. Differences in selection pressures

(e.g., competition and predation) can lead to differences in optimal school size and hence to

differences in aggregation tendency (i.e., the frequency of school fission and fusion) (Rieucau

et al. 2015). This may account for the different shapes of the relationships between population

density and school characteristics among species and taxa.

Furthermore, relationships between population density and school characteristics could

also vary within a species depending on several factors. For example, seasonality (e.g.,

spawning and feeding seasons), body size, and age can affect optimal school sizes and

aggregation tendencies (Hoare et al. 2000; Croft et al. 2003; Makris et al. 2009). Our study

was conducted during the feeding season, and variations of body size and age in the study

area were small throughout the study period (2007–2021). The body lengths of most chub

mackerel and Japanese sardine corresponded to those of young of the year (Kamimura et al.

2021b, a). Further studies across seasons, locations, body sizes, ages, and species are needed

to identify general patterns in relationships between population density and school

characteristics.

Why small pelagic fishes form such massive schools remains an open question

(Rieucau et al. 2015). In general, reducing predation risk is thought to be the most

widely-applicable explanation for animal aggregation, although there are other explanations

such as facilitating encounters with reproductive partners, improving foraging efficiency, and

hydrodynamic gains (Pitcher and Parrish 1993; Krause and Ruxton 2002; Ward and Webster

2016). However, current theories on the function of schooling behaviour do little to explain

the formation of massive schools (Rieucau et al. 2015; Maury 2017). This is because for

massive schools, the benefits of schooling are likely to stay constant with increasing group

size while the costs increase continually, meaning that the benefit-to-cost ratio declines

rapidly with increasing group size and is quite small. The type of predator encountered has

been suggested to be an important factor when assessing the benefits of massive schools

(Rieucau et al. 2015), because the benefits of group size could be affected by predator hunting

strategies (e.g., solitary vs. social predators) and attack efficiencies (e.g., the number of prey

caught per strike) (Rieucau et al. 2015). In our results, the relationships between population

density and school size differed among species, and the observed school size tended to be

smaller for chub mackerel than for Japanese sardine and Japanese anchovy. The main

predators of chub mackerel could differ from those of Japanese sardine and Japanese anchovy.

A more detailed examination of these factors could help explain why small pelagic fishes

form such massive schools.

Our results will be useful for population-status assessment of these species. Population

density was positively correlated with school characteristics, indicating that school

characteristics could be used as an indicator of population status. However, the relationships

were highly nonlinear (except for that of school size). Therefore, this nonlinearity should be

taken into account when using school characteristics as an indicator of population status

(Erisman et al. 2011).

In the present study, we inferred schooling characteristics from echosounder data.

Echosounder signals provide snapshot information in two dimensions: vertically through the

water column and horizontally with successive acoustic signals along the vessel track

(Simmonds and MacLennan 2008). This means that we were able to sample only a small

fraction of each school we identified. In fact, several schools were estimated to be quite small.

A previous study has reported that there are no significant differences between schooling

characteristics obtained from echosounders and multibeam sonar, which can image an entire

school in three dimensions (Brierley and Cox 2015). Therefore, the overall trend in our results

is likely to be robust. However, the use of methods that can obtain three-dimensional and/or

continuous-monitoring data will enable more accurate and detailed analysis (Makris et al.

2006; Paramo et al. 2007) in the future and could provide new biological and ecological

insights.

In conclusion, our results show that school size increased almost linearly with

increasing population density, whereas the number of schools and other school characteristics

(school packing density, school volume, and school cross-sectional area) increased

non-linearly such that the rate of increase slowed considerably with increasing population

density. Furthermore, we found that the shapes of the relationships between population

density and school characteristics differed among species, although the overall trend was

similar. These results indicate that the schooling behaviour of small pelagic fish species is

density-dependent, and responses to changes in density differ among species. Our results will

help advance understanding of how biological interactions such as intra- and inter-specific

competition and predator–prey interactions mediate the density-dependent processes that

underlie the population dynamics and community structure of small pelagic fishes in marine

ecosystems (Maury 2017).

Acknowledgements

The dataset analysed in this study was made possible by enormous efforts undertaken by

numerous researchers, as well as the captains, officers, and crews of the training vessel

Hokuho-Maru. We thank Dr. K. Abe for providing information on methods for analysing

echosounder data. We also thank English-speaking professional editors from ELSS, Inc., for

English proofreading.

Author contribution

Sho Furuichi conceived of the study and performed data analysis. All authors contributed to

data collection. The first draft of the manuscript was written by Sho Furuichi and all authors

commented on previous versions of the manuscript.

Funding

This study was partially funded by the Japan Fisheries Research and Education Agency and

the Fisheries Agency of Japan.

Declarations

Conflict of interest

The authors declare no competing interests.

References

Aronica S, Fontana I, Giacalone G, et al (2019) Identifying small pelagic Mediterranean fish schools from acoustic and environmental data using optimized artificial neural networks. Ecol Inform 50:149–161. https://doi.org/10.1016/j.ecoinf.2018.12.007

Bakun A (2006) Wasp-waist populations and marine ecosystem dynamics: Navigating the "predator pit" topographies. Prog Oceanogr 68:271–288. https://doi.org/10.1016/j.pocean.2006.02.004

Barange M (1994) Acoustic identification, classification and structure of biological patchiness on the edge of the Agulhas Bank and its relation to frontal features. South African J Mar Sci 14:333–347. https://doi.org/10.2989/025776194784286969

Beauchamp G (2014) Social predation: How groupliving benefits predators and prey. Academic Press, London, UK

Borner KK, Krause S, Mehner T, et al (2015) Turbidity affects social dynamics in Trinidadian guppies. Behav Ecol Sociobiol 69:645–651. https://doi.org/10.1007/s00265-015-1875-3

Brierley AS, Cox MJ (2015) Fewer but not smaller schools in declining fish and krill populations. Curr Biol 25:75–79. https://doi.org/10.1016/j.cub.2014.10.062

Canales TM, Delius GW, Law R (2020) Regulation of fish stocks without stock–recruitment relationships: the case of small pelagic fish. Fish Fish 21:857–871. https://doi.org/10.1111/faf.12465

Caro T (2005) Antipredator defenses in birds and mammals. University of Chicago Press, Chicago, IL

Carpenter B, Gelman A, Hoffman MD, et al (2017) Stan: A probabilistic programming language. J Stat Softw 76:. https://doi.org/10.18637/jss.v076.i01

Coetzee J (2000) Use of a shoal analysis and patch estimation system (SHAPES) to

characterise sardine schools. Aquat Living Resour 13:1–10.

https://doi.org/10.1016/S0990-7440(00)00139-X

Croft DP, Krause J, Couzin ID, Pitcher TJ (2003) When fish shoals meet: outcomes for

evolution and fisheries. Fish Fish 4:138–146.

https://doi.org/10.1046/j.1467-2979.2003.00113.x

Cury P, Bakun A, Crawford RJM, et al (2000) Small pelagics in upwelling systems: Patterns

of interaction and structural changes in "wasp-waist" ecosystems. ICES J Mar Sci

57:603–618. https://doi.org/10.1006/jmsc.2000.0712

D'Elia M, Patti B, Bonanno A, et al (2014) Analysis of backscatter properties and application

of classification procedures for the identification of small pelagic fish species in the

Central Mediterranean. Fish Res 149:33–42.

https://doi.org/10.1016/j.fishres.2013.08.006

Dalziel BD, Novak M, Watson JR, Ellner SP (2021) Collective behaviour can stabilize

ecosystems. Nat Ecol Evol 5:1435–1440. https://doi.org/10.1038/s41559-021-01517-w

Erisman BE, Allen LG, Claisse JT, et al (2011) The illusion of plenty: Hyperstability masks

collapses in two recreational fisheries that target fish spawning aggregations. Can J Fish

Aquat Sci 68:1705–1716. https://doi.org/10.1139/f2011-090

Fernandes PG (2009) Classification trees for species identification of fish-school echotraces.

ICES J Mar Sci 66:1073–1080. https://doi.org/10.1093/icesjms/fsp060

Foote KG (1987) Fish target strengths for use in echo integrator surveys. J Acoust Soc Am

82:981–987. https://doi.org/10.1121/1.395298

Francois RE, Garrison GR (1982) Sound absorption based on ocean measurements. Part II:

Boric acid contribution and equation for total absorption. J Acoust Soc Am

72:1879–1890. https://doi.org/10.1121/1.388673

540 Furuichi S, Kamimura Y, Yukami R (2021) Length–length and length–weight relationships

541   for four dominant small pelagic fishes in the Kuroshio–Oyashio current system. Thalass

542   An Int J Mar Sci 37:651–657. https://doi.org/10.1007/S41208-021-00300-9

543 Griesser M, Ma Q, Webber S, et al (2011) Understanding animal group-size distributions.

544   PLoS One 6:e23438. https://doi.org/10.1371/journal.pone.0023438

545 Gueron S, Levin SA (1995) The dynamics of group formation. Math Biosci 128:243–264.

546   https://doi.org/10.1016/0025-5564(94)00074-A

547 Hansen TF, Stenseth NC, Henttonen H, Tast J (1999) Interspecific and intraspecific

548   competition as causes of direct and delayed density dependence in a fluctuating vole

549   population. Proc Natl Acad Sci U S A 96:986–991. https://doi.org/10.1073/pnas.96.3.986

550 Hensor E, Couzin ID, James R, Krause J (2005) Modelling density-dependent fish shoal

551   distributions in the laboratory and field. Oikos 110:344–352.

552   https://doi.org/10.1111/j.0030-1299.2005.13513.x

553 Hoare DJ, Krause J, Peuhkuri N, Godin J-GJ (2000) Body size and shoaling in fish. J Fish

554   Biol 57:1351–1366. https://doi.org/10.1111/j.1095-8649.2000.tb02217.x

555 Holubová M, Čech M, Vašek M, Peterka J (2019) Density dependent attributes of fish

556   aggregative behaviour. PeerJ 2019:e6378. https://doi.org/10.7717/peerj.6378

557 Janson C (1985) Aggresive competition and individual food consumption in wild brown

558   capuchin monkeys (*Cebus apella*). Behav Ecol Sociobiol 18:125–138.

559   https://doi.org/10.1007/BF00299041

560 Janson CH (1988) Food competition in brown capuchin monkeys (*Cebus apella*): quantitative

561   effects of group size and tree productivity. Behaviour 105:53–76.

562   https://doi.org/10.1163/156853988X00449

563 Jech JM, Michaels WL (2006) A multifrequency method to classify and evaluate fisheries

acoustics data. Can J Fish Aquat Sci 63:2225–2235. https://doi.org/10.1139/F06-126

Jech JM, Stroman F (2012) Aggregative patterns of pre-spawning Atlantic herring on Georges

Bank from 1999-2010. Aquat Living Resour 25:1–14

Jenkins TM, Diehl S, Kratz KW, Cooper SD (1999) Effects of population density on

individual growth of brown trout in streams. Ecology 80:941–956.

https://doi.org/10.1890/0012-9658(1999)080[0941:EOPDOI]2.0.CO;2

Kamimura Y, Tadokoro K, Furuichi S, Yukami R (2021a) Stronger density-dependent growth

of Japanese sardine with lower food availability: Comparison of growth and zooplankton

biomass between a historical and current stock-increase period in the western North

Pacific. bioRxiv 2021.12.26.474216. https://doi.org/10.1101/2021.12.26.474216

Kamimura Y, Taga M, Yukami R, et al (2021b) Intra- and inter-specific density dependence

of body condition, growth, and habitat temperature in chub mackerel (*Scomber*

japonicus). ICES J Mar Sci 78:3254–3264. https://doi.org/10.1093/ICESJMS/FSAB191

Krause J, Ruxton GD (2002) Living in groups. Oxford University Press, Oxford, UK

Lima SL (1995) Back to the basics of anti-predatory vigilance: the group-size effect. Anim

Behav 49:11–20. https://doi.org/10.1016/0003-3472(95)80149-9

Mackenzie K V. (1981) Nine-term equation for sound speed in the oceans. J Acoust Soc Am

70:807–812. https://doi.org/10.1121/1.386920

Makris NC, Ratilal P, Jagannathan S, et al (2009) Critical population density triggers rapid

formation of vast oceanic fish shoals. Science 323:1734–1737.

https://doi.org/10.1126/science.1169441

Makris NC, Ratilal P, Symonds DT, et al (2006) Fish population and behavior revealed by

instantaneous continental shelf-scale imaging. Science 311:660–663.

https://doi.org/10.1126/science.1121756

Maury O (2017) Can schooling regulate marine populations and ecosystems? Prog Oceanogr

156:91–103. https://doi.org/10.1016/j.pocean.2017.06.003

Maury O, Poggiale JC (2013) From individuals to populations to communities: A dynamic

energy budget model of marine ecosystem size-spectrum including life history diversity.

J Theor Biol 324:52–71. https://doi.org/10.1016/j.jtbi.2013.01.018

Misund O (1993) Dynamics of moving masses: variability in packing density, shape, and size

among herring, sprat, and saithe schools. ICES J Mar Sci 50:145–160.

https://doi.org/10.1006/jmsc.1993.1016

Niwa H-S (2004) Space-irrelevant scaling law for fish school sizes. J Theor Biol 228:347–357.

https://doi.org/10.1016/j.jtbi.2004.01.011

Niwa H-S (1998) School size statistics of fish. J Theor Biol 195:351–361.

https://doi.org/10.1006/jtbi.1998.0801

Paramo J, Bertrand S, Villalobos H, Gerlotto F (2007) A three-dimensional approach to

school typology using vertical scanning multibeam sonar. Fish Res 84:171–179.

https://doi.org/10.1016/j.fishres.2006.10.023

Pavlov DS, Kasumyan AO (2000) Patterns and mechanisms of schooling behavior in fish: a

review. J Ichthyol 40:S163

Pépin D, Gerard JF (2008) Group dynamics and local population density dependence of group

size in the Pyrenean chamois, *Rupicapra pyrenaica*. Anim Behav 75:361–369.

https://doi.org/10.1016/j.anbehav.2006.09.030

Petitgas P, Reid D, Carrera P, et al (2001) On the relation between schools, clusters of schools,

and abundance in pelagic fish stocks. ICES J Mar Sci 58:1150–1160.

https://doi.org/10.1006/jmsc.2001.1130

Pitcher TJ, Parrish JK (1993) Functions of shoaling behaviour in teleosts. In: Pitcher TJ (ed)

The behaviour of teleost fishes, 2nd edn. Chapman and Hall, London, UK, pp 263–439

Post JR, Parkinson EA, Johnston NT (1999) Density-dependent processes in structured fish populations: interaction strengths in whole-lake experiments. Ecol Monogr 69:155–175. https://doi.org/10.1890/0012-9615(1999)069[0155:DDPISF]2.0.CO;2

R Development Core Team (2020) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria

Rice J (1995) Food web theory, marine food webs, and what climate change may do to northern fish populations. Can Spec Publ Fish Aquat Sci 121:561–568

Rieucau G, Boswell KM, De Robertis A, et al (2014a) Experimental evidence of threat-sensitive collective avoidance responses in a large wild-caught herring school. PLoS One 9:e86726. https://doi.org/10.1371/journal.pone.0086726

Rieucau G, De Robertis A, Boswell KM, Handegard NO (2014b) School density affects the strength of collective avoidance responses in wild-caught Atlantic herring *Clupea harengus*: a simulated predator encounter experiment. J Fish Biol 85:1650–1664. https://doi.org/10.1111/jfb.12520

Rieucau G, Fernö A, Ioannou CC, Handegard NO (2015) Towards of a firmer explanation of large shoal formation, maintenance and collective reactions in marine fish. Rev Fish Biol Fish 25:21–37. https://doi.org/10.1007/s11160-014-9367-5

Romenskyy M, Herbert-Read JE, Ioannou CC, et al (2020) Quantifying the structure and dynamics of fish shoals under predation threat in three dimensions. Behav Ecol 31:311–321. https://doi.org/10.1093/beheco/arz197

Sibly RM (1983) Optimal group size is unstable. Anim Behav 31:947–948. https://doi.org/10.1016/S0003-3472(83)80250-4

Simmonds J, MacLennan DN (2008) Fisheries acoustics: theory and practice, second edi.

Blackwell Publishing, Oxford, UK

Southwell CJ (1984) Variability in grouping in the eastern grey kangaroo, *Macropus*

giganteus I. Group density and group size. Wildl Res 11:423–435.

https://doi.org/10.1071/WR9840423

Stan Development Team (2020) Stan: the R interface to Stan. R package version 2.19.2

Sumpter DJT (2010) Collective animal behavior. Princeton University Press, Princeton, NJ

Vicsek T, Zafeiris A (2012) Collective motion. Phys Rep 517:71–140.

https://doi.org/10.1016/j.physrep.2012.03.004

Wada T, Matsubara T, Matsumiya Y, Koizumi N (1995) Influence of environment of stock

fluctuations of Japanese sardine, Sardinops melanostictus. Can Spec Publ Fish Aquat Sci

121:387–394

Ward A, Webster M (2016) Sociality: The behaviour of group-living animals. Springer

International Publishing

Watanabe C, Yatsu A (2004) Effects of density-dependence and sea surface temperature on

interannual variation in length-at-age of chub mackerel (Scomber japonicus) in the

Kuroshio-Oyashio area during 1970-1997. Fish Bull 102:206

Watanabe S (2013) A widely applicable bayesian information criterion. J Mach Learn Res

14:867–897

Yatsu A (2019) Review of population dynamics and management of small pelagic fishes

around the Japanese Archipelago. Fish Sci 85:611–639.

https://doi.org/10.1007/s12562-019-01305-3

Zhao X, Wang Y, Dai F (2008) Depth-dependent target strength of anchovy (*Engraulis*

japonicus) measured in situ. ICES J Mar Sci 65:882–888.

https://doi.org/10.1093/icesjms/fsn055

Zheng YH, Fu SJ (2021) Effects of fasting on collective movement and fission–fusion

dynamics in both homogeneous and heterogeneous shoals of a group-living cyprinid fish

species. J Fish Biol 99:1640–1649. https://doi.org/10.1111/JFB.14872

Table 1. Characteristics of detected schools of Japanese anchovy, Japanese sardine, and chub

mackerel by year. L_t, total length; Q1, first quartile; Q3, third quartile.

Species	Year	Total number of stations	Total number of schools	School depth (m)		Mean L_t (by station; cm)	
				Median	Q1–Q3	Median	Q1–Q3
Japanese anchovy	2007	1	5	24.6	24.1–27.5	12.7	–
Japanese anchovy	2008	3	21	22.0	18.0–24.7	11.8	11.1–12.2
Japanese anchovy	2009	12	53	26.0	22.4–30.9	12.1	11.6–12.4
Japanese anchovy	2010	10	68	20.0	17.8–23.9	12.8	12.5–13.2
Japanese anchovy	2011	13	92	21.4	19.2–25.4	10.8	10.4–11.8
Japanese anchovy	2012	8	34	19.3	16.7–22.0	13.3	12.9–13.7
Japanese anchovy	2013	2	21	26.1	22.6–27.6	13.0	12.9–13.1
Japanese sardine	2014	1	3	25.2	24.6–25.9	15.3	–
Japanese sardine	2015	6	38	18.9	16.6–22.4	15.3	15.3–15.6
Japanese sardine	2017	1	29	20.8	14.9–22.5	14.4	–
Japanese sardine	2018	4	61	16.4	15.7–17.6	13.6	13.4–13.7
Japanese sardine	2019	2	3	20.7	19.5–23.9	14.5	14.2–14.8
Japanese sardine	2020	1	1	19.4	–	14.5	–
Japanese sardine	2021	10	38	23.9	19.8–29.2	14.0	13.6–14.5
Chub mackerel	2010	1	3	14.7	14.4–26.0	21.7	–
Chub mackerel	2013	4	25	19.2	15.0–20.7	19.6	19.2–19.9
Chub mackerel	2014	1	1	22.3	–	23.1	–
Chub mackerel	2015	1	1	19.8	–	19.4	–
Chub mackerel	2016	1	4	21.4	18.0–23.0	18.7	–
Chub mackerel	2018	2	2	18.4	14.9–21.9	24.8	20.6–29.0
Chub mackerel	2019	1	1	18.4	–	25.8	–
Chub mackerel	2020	3	14	25.6	21.9–32.1	19.2	18.6–19.7

Figure captions

Fig. 1   Sampling stations (translucent circles) used for trawl-acoustic surveys in the Pacific

Ocean off Japan and the Kuril Islands from 2007 to 2021.

Fig. 2   Estimated power exponents for the relationship between local population density and

(a) school size, (b) the number of schools, (c) school packing density, (d) school volume, and

(e) school cross-sectional area. Points indicate estimated values (posterior means), and error

bars indicate 90% credible intervals.

Fig. 3   Relationship between local population density and school size for (a) Japanese

anchovy, (b) Japanese sardine, and (c) chub mackerel. Translucent circles indicate observed

values of individual schools. Solid lines indicate estimated relationships (posterior means),

and grey bands indicate 90% credible intervals.

Fig. 4   Relationship between local population density and the number of schools for (a)

Japanese anchovy, (b) Japanese sardine, and (c) chub mackerel. Translucent circles indicate

observed values at individual stations. Solid lines indicate estimated relationships (posterior

means), and grey bands indicate 90% credible intervals.

Fig. 5   Relationship between local population density and school packing density for (a)

Japanese anchovy, (b) Japanese sardine, and (c) chub mackerel. Translucent circles indicate

observed values of individual schools. Solid lines indicate estimated relationships (posterior

means), and grey bands indicate 90% credible intervals.

Fig. 6   Relationship between local population density and school volume for (a) Japanese

anchovy, (b) Japanese sardine, and (c) chub mackerel. Translucent circles indicate observed

values of individual schools. Solid lines indicate estimated relationships (posterior means),

and grey bands indicate 90% credible intervals.

Fig. 7   Relationship between local population density and school cross-sectional area for (a)

Japanese anchovy, (b) Japanese sardine, and (c) chub mackerel. Translucent circles indicate

observed values of individual schools. Solid lines indicate estimated relationships (posterior

means), and grey bands indicate 90% credible intervals.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7