The effect of migration and variation on populations of *Escherichia coli* adapting to complex fluctuating environments

S Selveshwari[1], Kaustubh Chandrakant Khaire[1,2], Aditee Kadam[1,3], Sutirth Dey[1,*]

1 Population Biology Laboratory, Department of Biology, Indian Institute of Science Education and Research Pune, Dr Homi Bhabha Road, Pashan, Maharashtra, 411008, India.

2 *Current address*: School of Energy Science and Engineering, Indian Institute of Technology Guwahati, Guwahati 781039, Assam, India

3 *Current address*: Department of Molecular Cell Biology, Weizmann Institute of Science, Rehovot, 7610001 Israel

*Name and address of the corresponding author:

Sutirth Dey

Biology Division
Indian Institute of Science Education and Research Pune
Dr Homi Bhabha Road
Pune, Maharashtra 411 008, India.
Tel.: +91 20 25908054; fax: +91 (20) 2590 8186;
e-mail: s.dey@iiserpune.ac.in

Acknowledgements: We thank Gayathri Pananghat and Milind Watve for their valuable inputs. SS was supported by a Junior Research Fellowship initially sponsored by IISER Pune and then by Department of Biotechnology (DBT), Govt. of India. AK was supported through INSPIRE fellowship of Department of Science and Technology (DST), Government of India. This project was supported by a grant from Department of Biotechnology, Government of India (#BT/PR22328/BRB/10/1569/2016) and internal funding from IISER Pune. The authors declare no conflict of interest.

Title: The effect of migration and variation on populations of *Escherichia coli* adapting to complex fluctuating environments

Abstract

Migration, a critical evolutionary force, can have contrasting effects on adaptation. It can aid as well as impede adaptation. The effects of migration on microbial adaptation have been studied primarily in simple constant environments. Very little is known about the effects of migration on adaptation to complex, fluctuating environments. In our study, we subjected replicate populations of *Escherichia coli*, adapting to complex and unpredictably fluctuating environments to different proportions of clonal ancestral immigrants. Contrary to the results from simple/constant environments, the presence of clonal immigrants reduced all measured proxies of fitness. However, migration from a source population with a greater variance in fitness resulted in no change in fitness w.r.t the no-migration control, except at the highest level of migration. Thus, the presence of variation in the immigrants could counter the adverse effects of migration in complex and unpredictably fluctuating environments. Our study demonstrates that the effects of migration are strongly dependent on the nature of the destination environment and the genetic makeup of immigrants. These results enhance our understanding of the influences of migrating populations, which could help better predict the consequences of migration.

Keywords

Clonal immigrants
Variation in fitness
Escherichia coli
Experimental evolution
Sub-optimal environment
Unpredictable fluctuations

1. Introduction

Migration affects several ecological and evolutionary processes, such as a species' range (Kirkpatrick and Barton 1997, Barton 2001, Sexton, McIntyre et al. 2009), composition, and diversity in meta-populations and natural communities (Venail, MacLean et al. 2008, Albright and Martiny 2018) and evolution of traits like virulence (Boots and Sasaki 1999, Lively 1999) and antibiotic resistance (Perron, Gonzalez et al. 2007). Interestingly, when it comes to adaptation, migration can have contrasting effects. For example, migration can impede adaptation in coevolving host-phage systems (Morgan, Gandon et al. 2005, Vogwill, Fenton et al. 2011) and microbial communities subjected to warmer temperatures (Lawrence, Bell et al. 2016). Similarly, in viruses, migration can reduce the extent of specialization to different tissue types (Cuevas, Moya et al. 2003). One of the ways migration negatively affects adaptation is by swamping the destination environment with alleles that are beneficial or neutral at the source environment but maladaptive at the destination (Kawecki and Holt 2002, Kawecki and Ebert 2004, Yeaman and Guillaume 2009). At the same time, several studies have demonstrated that migration can promote adaptation. For example, asexual populations of *Chlamydomonas* exposed to herbicides (Lagator, Morgan et al. 2014) and yeast populations evolving in the presence of salt stress (Bell and Gonzalez 2011) adapted more rapidly in the presence of migration. Similarly, Φ6 phage populations adapted better to a novel host when there was migration from populations that had the ability to infect this novel host (Ching, Musheyev et al. 2013). However, when migrants came from control populations unable to infect the novel hosts, there was no effect on the absolute fitness (Ching, Musheyev et al. 2013). Migration can positively influence adaptation by increasing the supply of beneficial mutations, particularly when populations are mutation limited (Holt 2003, Sexton, McIntyre et al. 2009), such as in asexual microbes.

Interestingly, most empirical studies on how migration influences adaptation in microbes have been carried out only in constant environments, typically in the presence of a single selection pressure (Morgan, Gandon et al. 2005, Dennehy, Friedenberg et al. 2010, Ching, Musheyev et al. 2013, Lagator, Morgan et al. 2014, Lawrence, Bell et al. 2016). However, in nature, organisms are often faced with heterogeneous environments that contain multiple stressors simultaneously. To further complicate matters, the magnitudes of these stresses can fluctuate over time, either predictably or unpredictably. Adaptations in such spatially and/or temporally heterogeneous environments can differ from those in simple constant environments (Levins 1968, Reboud and Bell 1997, Cooper and Lenski 2010, Karve, Bhave et al. 2016). The effect of migration on adaptation under such complex and fluctuating environments has received less attention (however, see (Perron, Gonzalez et al. 2007)).

Here, we present the results of a study on the effects of different migration rates on adaptation in replicate populations of *Escherichia coli* that were subjected to complex environments undergoing unpredictable fluctuations. We also looked at the effects of migrants that were either clonal or carrying variation. When the immigrants were clonal, the recipient populations evolved reduced fitness compared to the no migration control. Interestingly, the magnitude of fitness reduction varied positively with the fraction of immigrants received. However, treatments that received immigrants with variation in fitness showed little or no change in fitness compared to the no migration control.

2. Materials and Methods

For experimental evolution, we used *Escherichia coli* MG1655 with a kanamycin resistance cassette. All cultures were maintained at 37°C and 150 RPM throughout the selection and assays, except where stated otherwise.

2.1 Immigrant Populations

This study consisted of two selection experiments. In each selection experiment, we used two populations: the immigrant and the native. The native populations evolved in the complex fluctuating environments and experienced the effects of immigration. The immigrant populations were non-evolving cultures freshly revived every day (see section 2.3 Selection protocol). The two selection experiments differed only in terms of the nature of the immigrant populations. In experiment 1, we used a **C**lonal immigrant population (henceforth C) derived from a single *E.coli* colony and grown in 150ml NB with kanamycin (NB^Kan) for 18 hours. Multiple 1 ml glycerol stocks (15%) of this culture were prepared and stored at -80°C. In the second experiment, we used the **V**ariation immigrant population (henceforth V), which was derived from the C population. V population was initiated by reviving 1ml glycerol stock of the C population in 10ml NB^Kan followed by inoculation of 1ml of this revived culture in 50ml NB^Kan. We sub-cultured (1/10th dilution) this population for the next 15 days into 50ml NB^Kan every 12 hours. After 15 days (i.e., 30 transfers), 50ml of the grown culture was added to 50ml fresh NB^Kan and incubated for another 12 hours. Multiple 1ml glycerol stocks (15%) were prepared and stored at -80°C. Over these 15 days (~100 generations) that the V populations spent in a benign environment, we expected many spontaneous mutations to arise in the culture. Since we maintained this population under optimal conditions, large culture volume (N_f: 50ml) and lenient bottlenecks (1/10), we expected most mutations arising in the population to accumulate. Thus, the V population

should harbor more genetic variation than the C population. To confirm this, we quantified the within-population variance in fitness in the C and V populations using the methodology of an earlier study (McDonald, Hsieh et al. 2012).

2.2 Quantification of within-population variance

To quantify the variation accumulated in the V population, we assayed the fitnesses of 72 individual colonies of both C and V populations in 6 environments. We used the within-population fitness variance as a proxy for genetic variation (McDonald, Hsieh et al. 2012). 1ml glycerol stock of the C or V population was revived in 10ml NB^{Kan}, overnight. The revived cultures were streaked on nutrient agar (NA) with kanamycin from where, we made single colony suspensions from 6 similar-sized colonies of both C and V populations. We then inoculated 10µL of the single colony suspension in 200µL of six assay environments, assayed in a single 96-well plate. The environments included sub-lethal concentrations of all stress combinations and NB^{Kan}. Environment I: pH 5+Salt 3.5g%, Environment II: pH 8.5+Salt 2.5g%, Environment III: pH 5+4µl 0.3% H_2O_2, Environment IV: pH 8.5+2µl 0.3% H_2O_2, Environment V: Salt 2.5g%+1.5µl 0.3% H_2O_2 and Environment VI: NB^{Kan}. H_2O_2 was added 2 hours after inoculation, where required. The populations were continuously monitored (OD_{600}) for 24 hours, from inoculation, using a plate reader (Synergy HT) at 37°C and continuous medium shaking. We repeated the entire procedure over 12 days for a total of 72 colonies from each of the C and V populations. Growth rate and yield were estimated using a custom python script; see section 2.4 for details. The coefficient of variation (CV) in fitness was computed between the six colonies assayed every day. CV was computed for fitness measured as both growth rate and yield. The CV of the C and V populations were compared in each of the six environments.

2.3 Selection Protocol

We initiated 48 replicate populations from both the C and the V ancestor populations. We then revived 1ml of the corresponding glycerol stocks overnight in 10ml NB^{Kan} and inoculated 20µL of the revived culture (OD_{600} 1.0 – 1.1) into 2 ml of the selection environment

Selection environment: The populations were subjected to selection for 30 days in complex environments (i.e., multiple stressors were present simultaneously) and fluctuated unpredictably. We used the same selection regimes as a previous study (Karve, Daniel et al. 2015). Briefly, selection involved pH, osmotic (NaCl), and oxidative (H_2O_2) stress. We chose the combinations of the stresses so that two of the three components were present at inhibitory concentrations and the third was present at a concentration as found in NB (i.e., pH=7, NaCl=0.5g% and H_2O_2=0). We tested several such combinations for their effect on the growth of the WT in a pilot experiment. We chose combinations that resulted in ~ 40 – 70% reduction in growth compared to NB growth. We chose a total of 28 combinations and a sequence of 30 environments from a uniform distribution with replacement (Table 1).

Migration treatment: We used four levels of migration, namely 0% (control), 10% (low migration), 50% (intermediate migration) and 90% (high migration). If we simply added the immigrants to the native population, the total population size would increases with the migration rate, thus resulting in large differences in population sizes across treatments. Previous studies have shown that population size can affect evolutionary outcomes in microbial populations (Chavhan, Ali et al. 2019). Therefore, to avoid the confounding effect of population size, we kept the inoculum size constant (~10^7 cells) and defined migration as the percentage of immigrants in the inoculum. For example, in the low (10%) migration treatment, 10% of the individuals in the subculture inoculum consisted of immigrants. The

remaining 90% were individuals from the native population, evolving under a complex fluctuating environment. The proportions were adjusted based on the OD_{600} values of the immigrant and native populations at the time of subculture. When $OD_{600} = 1$, the culture contained ~10^9 cells/ml of NB (S Selveshwari, personal observations).

Every day, we revived 1ml glycerol stock of the C or V population in 10ml NBKan. OD_{600} of the revived culture was adjusted to $1.0 - 1.1$ and used as immigrants. We also measured OD_{600} of the native populations and used an appropriate culture volume containing the required inoculum size for the sub-culture. When the OD_{600} of these populations was less than 0.3, the populations were considered extinct, and the native population was obtained from the previous non-extinct population, stored at 4°C. We stored the selected populations as glycerol stocks at the end of 30 days.

2.4 Fitness assays

We measured the fitnesses of the evolving populations during and after selection. We noted the OD_{600} in the selection environment every 24 hours (i.e., at carrying capacity, just before sub-culturing). We used the geometric mean of these values over the 30 days of selection to measure fitness under fluctuating stress (Orr 2009). Using the same data, we also estimated the probability of extinction ($OD_{600} < 0.3$) during selection.

We measured post-selection fitnesses as growth rate and yield of the populations in three representative environments: environment 1: pH 8.5+salt 4.5g%; environment 2: pH 5+0.5μL H_2O_2; environment 3: salt 2.5g%+0.7μL H_2O_2. All 48 replicate populations of each selection experiment were assayed twice in each assay environment. 4μL of glycerol stock was revived in 2ml NB^{Kan}, overnight. We measured the OD_{600} of the revived culture and inoculated a volume containing 10^7 cells (assuming 10^9 cells/2ml when $OD_{600} = 1$) in 2ml assay environments in 24-well tissue culture plates. The OD_{600} was measured every 2 hours for 24

hours at 37ºC and slow continuous shaking using a plate reader (Synergy HT BioTek, Winooski, VT, USA). We used population growth rate and yield as fitness measures (Karve, Daniel et al. 2015, Chavhan, Ali et al. 2019). The growth curve data were analyzed using a custom python script that fits overlapping straight lines over 6-hour windows. We estimated growth rate as the maximum slope of the curve and yield as the maximum OD_{600} reached in 24 hours.

2.5 Statistical analysis

All fitness comparisons were performed independently for the two selections. Population fitness during selection (geometric mean of growth and extinction probability) was compared across the migration treatments using separate one-way ANOVAs with migration treatment (0, 10, 50, 90) as a fixed factor. We compared post-selection fitnesses (growth rate and yield) for each assay environment, using separate two-way mixed model ANOVAs. We treated migration treatment (0, 10, 50, 90) as a fixed factor. The growth rate and yield from the two rounds of assays were considered measurement replicates. The biological replicates (12 levels) acted as random factors and were nested in migration treatment. To account for the inflation of family-wise error rate, the *P*-values of the main effect of migration were subjected to Holm–Šidák correction (Abdi 2010). We performed pairwise comparisons using Tukey's post hoc analysis when the corrected *P*-values was significant. We also computed the Cohen's *d* statistics (Cohen 2013) to measure the effect sizes (Sullivan and Feinn 2012). The biological significance of the differences between the treatments were interpreted as small, medium and large for $0.2 < d < 0.5$, $0.5 < d < 0.8$ and $d > 0.8$, respectively. We compared the within-population variance in the fitness of the C and V populations using paired t-Test in each environment. The inflation in family-wise error rate was controlled using Holm–Šidák correction.

We performed all the ANOVAs on STATISTICA v7.0 (Statsoft Inc.). Cohen's d statistics were estimated using the freeware Effect Size Generator v2.3.0 (Devilly 2004).

Table 1. Sequence of complex fluctuating environments used during selection.

Day	pH	NaCl (g/100ml)	H_2O_2 (µL)
1	5	4	-
2	4.5	3	-
3	5	4.5	-
4	-	2.5	0.8
5	5	3.5	-
6	5	-	0.6
7	5	3	-
8	-	4	0.5
9	-	3	0.5
10	5	3	-
11	-	4	0.5
12	-	3	0.5
13	5	-	0.7
14	-	3	0.5
15	8.5	-	0.5
16	-	3	0.7
17	5	-	0.5
18	5	3	-
19	9	-	0.5
20	5	5	-
21	-	3	0.5
22	9	-	0.5
23	8.5	5	-
24	4.5	3	-
25	8.5	4	-
26	-	2	0.8
27	5	3	-
28	8.5	4	-
29	5	4	-
30	8.5	4	-

3. Results

3.1. Clonal immigration impedes adaptation in complex and unpredictable environment

Figure 1. Effect of clonal migration during selection in complex, unpredictable environments. Fitness was measured as A) Geometric mean of growth during selection. B) Probability of extinction in the selection environments, during selection. Each box plot represents data from 12 replicate populations. Solid lines represent median, dotted lines denote mean, whiskers denote 10[th] and 90[th] percentiles, and dots denote 5[th] and 95[th] percentile. Box plots denoted by different letters are significantly different from each other (P < 0.05 in Tukey's posthoc analysis).

After 30 rounds of clonal migration and selection (corresponding to ~200 generations in the no migration control populations), we found that the control populations (i.e., those that did not receive any immigrants) adapted significantly more than populations that received immigrants (from clonal source, C). The geometric mean of growth during selection was significantly different between the migration treatments (Fig. 1A; $F_{3,44} = 28.76$, $P = 1.88$E-10). The no migration control populations had the highest GM of growth. However, this was only significantly higher than intermediate (Tukey's p = 0.01; Cohen's $d = 1.49$ (large)) and high (Tukey's p = 0.0002; Cohen's $d = 3.56$ (large)) migration treatments. The reduced GM

of growth of high migration treatment was also significant w.r.t low (Tukey's p = 0.0002; Cohen's d = 2.88 (large)) and intermediate (Tukey's p = 0.0002; Cohen's d = 2.42 (large)) migration treatments. This reduction in growth was accompanied by a significant effect in terms of extinction probability (Fig. 1B; $F_{3,44}$ = 13.48, P = 2.25E-06). Tukey's post hoc indicated that the high migration treatment had significantly greater extinction probability compared to all treatment population (control: p = 0.0002; Cohen's d = 2.11 (large); intermediate: p = 0.0002; Cohen's d = 2.13 (large); high: p = 0.006; Cohen's d = 1.66 (large)). Thus, during selection, the presence of migration reduced the populations' ability to survive and grow in the selection environment, resulting in a significant increase in extinction probability at high migration. We next tested how this reduced survivability and growth affected the overall adaptation.

3.2. Presence of variation in the immigrant pool counters the negative effect of migration.

Figure 2. Effect of clonal migration on post-selection fitness. Two fitness proxies, A) growth rate and B) growth yield, were measured in three complex environments. See methods for the composition of the three complex environments. Each box plot represents 24 values, i.e., 12 replicate populations, assayed twice. Solid lines represent median, dotted lines denote mean, whiskers denote 10th and 90th percentiles, and dots denote 5th and 95th percentile. Box plots denoted by different letters are significantly different from each other (P < 0.05 in Tukey's posthoc analysis).

After selection, we estimated the extent of adaptation as growth rate and yield in three complex environments. When the fitnesses across all three environments were analyzed together in a single ANOVA, we saw that the interaction between migration treatments and assay environment was significant (Fig. 2; Growth rate: $F_{6,144} = 2.43$, $P = 0.03$; Yield: $F_{6,144} = 4.62$, $P = 4.0E-04$). Therefore, the effect of clonal migration on fitness in the three complex environments was analyzed separately.

The results of ANOVAs, P-values and Cohen's d of all pairwise comparisons are summarized in Tables 2 and 3. Briefly, all migration treatments had significantly lower growth rates and yield than the no migration control populations in environment 1. The growth rate of the low migration treatment was significantly higher than both intermediate and high migration treatments, which were not significantly different from each other. Similarly, although the yield of the low migration treatment was higher, it was statistically significant w.r.t only the high migration treatment. However, both intermediate and high migration treatments were not significantly different. In environment 2, only two migration treatments (intermediate and high) had significantly lower growth rates and yield w.r.t the control. However, the growth rate of the low migration populations was only marginally insignificant (Tukey's p = 0.054) w.r.t control populations in environment 2. Both the growth rate and the yield of the low migration treatment were significantly higher than the intermediate and high migration treatments. However, there was no significant difference between growth rates and yields of intermediate and high migration treatments. In environment 3, the growth rate and the yield of only the high migration treatment were significantly lower than all other treatments. Thus, taken together, we saw that migration can have an overall negative effect on evolutionary outcomes under complex and unpredictable environments.

Table 2. Summary of the *P*-values for the growth rate of clonal migration post-selection. *P*-values of the main effect of migration was reported after Holm-Šidák correction for familywise error. Migration treatments were compared with each other using Tukey's post-hoc analysis. Pairwise effect sizes were computed as Cohen's d.

Growth rate					
	Holm Sidak Corrected *P*-values	Pairwise comparison	*P*-values (Tukey's post hoc)	Cohen's *d*	Interpretation
Envt. 1	0.001252	Cont. vs. Low	0.011	0.491	Small
		Cont. vs. Inter	1.67E-04	1.043	Large
		Cont. vs. High	1.67E-04	1.245	Large
		Low vs. Inter	0.023	0.702	Medium
		Low vs. High	0.002	0.997	Large
		Inter vs. High	0.814	0.348	Small
Envt. 2	5.68E-06	Cont. vs. Low	0.054	0.684	Medium
		Cont. vs. Inter	1.75E-04	1.321	Large
		Cont. vs. High	1.67E-04	2.330	Large
		Low vs. Inter	0.037	0.626	Medium
		Low vs. High	2.98E-04	1.302	Large
		Inter vs. High	0.255	0.486	Small
Envt. 3	5.07E-05	Cont. vs. Low	0.462	0.390	Small
		Cont. vs. Inter	0.330	0.473	Small
		Cont. vs. High	1.76E-04	1.915	Large
		Low vs. Inter	0.995	0.059	Small
		Low vs. High	0.002	1.184	Large
		Inter vs. High	0.003	1.177	Large

Table 3. Summary of the *P*-values for Yield of clonal migration, post-selection.
P-value of the main effect of migration was reported after Holm-Šidák correction for familywise error. Migration treatments were compared with each other using Tukey's post-hoc analysis. Pairwise effect sizes were computed as Cohen's d.

Yield					
	Holm Sidak Corrected *P*-values	Pairwise comparison	*P*-values (Tukey's post hoc)	Cohen's *d*	Interpretation
Envt. 1	0.000116	Cont. vs. Low	0.171	0.598	Medium
		Cont. vs. Inter	0.321	1.113	Large
		Cont. vs. High	1.68E-04	1.304	Large
		Low vs. Inter	0.984	0.642	Medium
		Low vs. High	0.002	0.894	Medium
		Inter vs. High	0.001	0.227	Small
Envt. 2	1.11E-08	Cont. vs. Low	0.532	0.350	Small
		Cont. vs. Inter	1.68E-04	1.763	Large
		Cont. vs. High	1.67E-04	2.851	Large
		Low vs. Inter	2.85E-04	1.088	Large
		Low vs. High	1.67E-04	1.668	Large
		Inter vs. High	0.441	0.465	Small
Envt. 3	1.5E-05	Cont. vs. Low	0.034	0.567	Medium
		Cont. vs. Inter	2.13E-04	0.479	Small
		Cont. vs. High	1.72E-04	2.122	Large
		Low vs. Inter	0.159	0.088	Small
		Low vs. High	0.044	1.137	Large
		Inter vs. High	0.934	1.271	Large

3.3 Variant (V) population has higher variation in fitness than the clonal (C) population

Figure 3. The average coefficient of variation of the clonal and variant populations. We measured the coefficient of variation (CV) of growth rate and yield of the two populations, assayed in 6 representative environments. Each bar represents the average CV over 72 individual colonies assayed over 12 days. Error bars are SE of the mean. * denote *P*-value < 0.05 in paired t-Test after Holm-Šidák correction.

Next, we investigated the effects of increased variation in the migrant population on the evolutionary outcome of the recipient populations. For this, we enriched the variation of the ancestral population (variation source, V). After ~100 generations of lenient (1/10) bottlenecks, in a benign environment, the V population showed a significant increase in variation in fitness, measured as coefficient of variation, in 4 out of 6 tested environments (Fig. 3 and Table 4). The increase in variation was seen w.r.t both growth rate and yield in environments II, III, V and NB. Thus, the V population had a larger within-population fitness variance than the C population and was used as the source of migrants in a second selection experiment.

Table 4. Summary of the *P*-values of paired T-test of coefficient of variation in fitness between C and V ancestors, after Holm-Šidák correction

	P-value after Holm-Šidák correction	
Environment	Growth rate	Yield
I	0.649	0.425
II	0.003	0.0001
III	0.018	0.005
IV	0.413	0.858
V	0.044	0.014
NB	0.006	0.009

3.4 Presence of variation in the immigrant pool counters the negative effect of migration.

Figure 4. The effect of migration, when the immigrants carry variation. Fitness was measured as A) Geometric mean of growth during selection. B) Probability of extinction in the selection environments, during selection. Each box plot represents data from 12 replicate populations. Solid lines represent median, dotted lines denote mean, whiskers denote 10th and 90th percentiles, and dots denote 5th and 95th percentile. Box plots denoted by different letters are significantly different from each other (P < 0.05 in Tukey's posthoc analysis).

When the migrant populations carried variation (V population), migration had an effect only at high levels of migration. A significant main effect was observed w.r.t geometric mean of growth (Fig 4A; $F_{3,44} = 34.14$, $P = 1.48E\text{-}11$). GM of growth was reduced at high levels of migration and this reduction was significantly different from all other treatments in Tukey's post-hoc analysis (control: p = 1.69E-04, Cohen's d = 2.25 (large); low: p = 1.69E-04, Cohen's d = 3.37 (large); intermediate: p = 1.69E-04, Cohen's d = 3.52 (large)). Subsequently, extinction probability also had a significant main effect of migration (Fig 4B; $F_{3,44} = 6.07$, p= 0.0015) and high migration treatment had an elevated probability of extinction. However, this increase in extinction was significantly different from only low and intermediate migration treatment (low: p = 9.32E-04, Cohen's d = 2.02 (large); intermediate: p = 0.03, Cohen's d = 0.90 (large)). Thus in contrast to migration from a clonal source, the negative effect of migration, during selection, was diminished when the migrants carry variation.

3.5. Presence of variation in the immigrant pool counters the negative effect of migration.

Figure 5. Effect of migrants with variation on post-selection fitness. Fitness was estimated as A) growth rate and B) yield in three complex environments. Each box plot represents data of 24 values, i.e., 12 replicate populations, assayed twice. Solid lines represent median, dotted lines denote mean, whiskers denote 10th and 90th percentiles, and dots denote 5th and 95th percentile. Box plots denoted by different letters are significantly different from each other (P < 0.05 in Tukey's posthoc analysis).

Following section 3.1, we analyzed the fitnesses of the populations receiving migrants with variation in the three complex environments using independent ANOVAs (Tables 5 and 6). In contrast to clonal migration, we found that the main effect of migration was either non-significant (growth rate in environments 2 and 3) or, when significant, the effect was limited to only the populations receiving a high level of migration. The growth rate and yield in environment 1 were significant only between the no-migration control and high migration

treatments. The main effect of growth rate in environment 2 was not significant ($F_{3,44} = 1.9$, $P = 0.144$). However, yield in environment 2 was significant, and Tukey's post hoc analysis showed that high migration treatment had significantly lower yield than control and low migration treatments (Table 6). Like environment 2, growth rate in environment 3 was also not significant ($F_{3,44} = 2.56$, $P = 0.13$). Similarly, yield in environment 3 was significant, and Tukey's post hoc analysis showed that high migration treatment has a significantly lower yield than only no-migration control (Table 6). These results illustrate that variation in the migrant pool can ameliorate the negative effects of migration.

Table 5. Summary of the *P*-values for the growth rate of populations receiving migrants with variation. *P*-values of the main effect of migration was reported after Holm-Šidák correction for family-wise error. Migration treatments were compared with each other using Tukey's post-hoc analysis. Pairwise effect sizes were computed as Cohen's d.

Growth rate					
	Holm Sidak Corrected *P*-values	Pairwise comparison	*P*-values (Tukey's post hoc)	Cohen's *d*	Interpretation
Envt. 1	0.02406	Cont. vs. Low	0.984	0.109	Small
		Cont. vs. Inter	0.612	0.395	Small
		Cont. vs. High	0.032	0.847	Large
		Low vs. Inter	0.819	0.283	Small
		Low vs. High	0.074	0.747	Medium
		Inter vs. High	0.377	0.510	Medium
Envt. 2	0.1435	NA	NA	NA	NA
Envt. 3	0.1296	NA	NA	NA	NA

Table 6. Summary of the *P*-values for yield of migration with variation. *P*-value of the main effect of migration was reported after Holm-Šidák correction for familywise error. Migration treatments were compared with each other using Tukey's post-hoc analysis. Pairwise effect sizes were computed as Cohen's d.

Yield					
	Holm Sidak Corrected *P*-values	Pairwise comparison	*P*-values (Tukey's post hoc)	Cohen's *d*	Interpretation
Envt. 1	1.9E-05	Cont. vs. Low	0.581	0.528	Medium
		Cont. vs. Inter	0.703	0.397	Small
		Cont. vs. High	0.010	1.070	Large
		Low vs. Inter	0.997	0.077	Small
		Low vs. High	0.200	0.681	Medium
		Inter vs. High	0.136	0.690	Medium
Envt. 2	0.007	Cont. vs. Low	1.000	0.046	Small
		Cont. vs. Inter	0.808	0.284	Small
		Cont. vs. High	0.009	0.934	Large
		Low vs. Inter	0.755	0.298	Small
		Low vs. High	0.007	0.922	Large
		Inter vs. High	0.084	0.555	Medium
Envt. 3	0.010	Cont. vs. Low	0.680	0.371	Small
		Cont. vs. Inter	0.328	0.587	Medium
		Cont. vs. High	0.003	1.057	Large
		Low vs. Inter	0.933	0.164	Small
		Low vs. High	0.061	0.627	Medium
		Inter vs. High	0.210	0.491	Small

4. Discussion

In this study, we investigated the effect of immigration on the adaptation of asexual populations in complex and unpredictably fluctuating environments. Immigration from a clonal and non-evolving source (ancestor) population resulted in a reduction in fitness during selection (Fig. 1) as well as post-selection (Fig. 2). During selection, as the proportion of immigrants increased, the geometric mean (GM) of growth decreased (Fig. 1A), and the extinction probability increased (Fig: 1B). After ~200 generations, the populations that received immigrants during selection had adapted less with a reduced growth rate (Fig. 2A) and yield (Fig. 2B). The fitness reduction increased with the fraction of immigrants in the evolving populations.

These results contradict several studies where migration promotes larger and/or faster adaptation in asexual organisms (Bell and Gonzalez 2011, Lagator, Morgan et al. 2014). In particular, it disagrees with a previous study (Perron, Gonzalez et al. 2007) that used a clonal source population as source of migration and showed that increasing immigration rates leads to rapid evolution. In their study, although the rates of adaptation were faster in benign (single antibiotic) environments than in harsh (two antibiotics) environments, the effect of migration was positive in all cases. In contrast, we saw increasing adverse effects of migration with increasing migration rates. One possible reason for the discrepancy in results could be the stress intensity used in the selection environments. While both experiments had multiple stress components, Perron *et al.* (Perron, Gonzalez et al. 2007) used lethal concentrations of stress, whereas we used sub-lethal concentrations. Lethal stresses create a sink environment where population size is expected to decline without sustained migration (as pointed out by the authors themselves) (Dias 1996, Holt 1997, Holt and Gomulkiewicz 1997). As in the case of the two experiments (Perron, Gonzalez et al. 2007) and this study),

immigration from a clonal source population can be expected to promote adaptation in terms of its demographic effect, i.e., changes in population sizes, which in turn can influence the adaptive dynamics of these populations. However, in non-lethal/ non-sink environments, like in our experiment, migration provided no demographic advantage as populations can persist here without immigration. Instead, the effect was largely negative as selection for locally fitter individuals can be diluted by increasing proportions of immigrants (Kawecki and Holt 2002, Lenormand 2002, Kawecki and Ebert 2004). The two studies taken together highlight that the nature of the environment faced by the evolving populations needs to be considered when studying the effect of migration on adaptation. The results can be very different when populations evolve in sub-optimal environments compared to those evolving in lethal environments.

Since clonal immigrants in sub-optimal environments did not provide any significant advantage to adaptation, we next investigated how variation in the immigrants influences adaptation to complex and unpredictable environments. To this end, we conducted a second selection experiment using a source population with a larger variation in fitness (Fig. 3). Immigration from this variant population had little or no effect on the evolving population, as seen from fitness measured during and post-selection (Figs. 4 and 5). Populations receiving low and intermediate levels of migration did not showed a reduction w.r.t any aspects of fitness. However, a reduction in fitness was observed when the populations were subjected to high migration.

It has been previously shown in bacteriophages that increased variation due to migration can promote adaptation (Dennehy, Friedenberg et al. 2010). However, in that study, the positive effect was limited to immigration from source populations grown in the same environment as the selection environment. Migration between similar environments could have promoted the

spread of beneficial variation between sub-populations (Kassen 2014). On the other hand, in our study, the source population was grown in a benign environment, unrelated to the selection environment. Migration from such a source is not limited to only beneficial variants.

Additionally, the distribution of fitness effects of new mutations is expected to vary greatly in a complex and fluctuating environment with rugged and shifting fitness landscapes (Van Cleve and Weissman 2015). Thus, in non-sink but complex unpredictable environments, we saw that the benefit of increased variation was only enough to counter the adverse effects of migration. These results agree with theoretical predictions that variation in the migrant pool can ameliorate the harmful effects of migration (Barton 2001). However, if the variation is too high, one can expect adverse effects on fitness, including extinctions (Barton 2001). Significant reduction in all aspects of fitness with high migration can indicate the existence of such a limit on the positive effects of increased variation via migration.

Populations receiving high levels of migrants, from both clonal and variant sources, relied on recurrent immigration and revival from previous time points for survival under complex, unpredictable conditions. This was a clear demonstration of the creation of pseudo-sinks where viable environments appear to have become sink environments due to high migration (Watkinson and Sutherland 1995). Repeated introduction of individuals into sub-optimal environments can limit adaptation and result in the populations being in a constant state of maladaptation, a phenomenon commonly observed at range margins (Kirkpatrick and Barton 1997).

5. Conclusions

Migrating individuals play multiple roles (demographic, variation) in the destination environment (Garant, Forde et al. 2007). The relative importance of these aspects of migration and their influence on adaptation is dependent on the quality of the environment. Maintaining a sustainable population might be more critical in a lethal sink environment. However, the supply of variation is more critical in sub-optimal environments, without which migration can create pseudo-sink environments. Since sub-optimal environments are likely more prevalent in nature, studies conducted under such conditions need to understand the effects of migration fully. Additionally, it becomes vital to consider the interactions between the role of migrating individuals and the environment into which they migrate.

6. References

Abdi, H. (2010). "Holm's sequential Bonferroni procedure." Encyclopedia of research design **1**(8): 1-8.

Albright, M. B. and J. B. Martiny (2018). "Dispersal alters bacterial diversity and composition in a natural community." The ISME journal **12**(1): 296-299.

Barton, N. (2001). Adaptation at the edge of a species' range. Integrating ecology and evolution in a spatial context. Cambridge, UK.

Bell, G. and A. Gonzalez (2011). "Adaptation and evolutionary rescue in metapopulations experiencing environmental deterioration." Science **332**(6035): 1327-1330.

Boots, M. and A. Sasaki (1999). "'Small worlds' and the evolution of virulence: infection occurs locally and at a distance." Proceedings of the Royal Society of London. Series B: Biological Sciences **266**(1432): 1933-1938.

Chavhan, Y. D., S. I. Ali and S. Dey (2019). "Larger numbers can impede adaptation in asexual populations despite entailing greater genetic variation." Evolutionary Biology **46**(1): 1-13.

Ching, J., S. A. Musheyev, D. Chowdhury, J. A. Kim, Y. Choi and J. J. Dennehy (2013). "Migration enhances adaptation in bacteriophage populations evolving in ecological sinks." Evolution **67**(1): 10-17.

Cohen, J. (2013). Statistical power analysis for the behavioral sciences. New York, Academic press.

Cooper, T. F. and R. E. Lenski (2010). "Experimental evolution with E. coli in diverse resource environments. I. Fluctuating environments promote divergence of replicate populations." BMC evolutionary biology **10**(1): 1-10.

Cuevas, J., A. Moya and S. Elena (2003). "Evolution of RNA virus in spatially structured heterogeneous environments." Journal of evolutionary biology **16**(3): 456-466.

Dennehy, J. J., N. A. Friedenberg, R. C. McBride, R. D. Holt and P. E. Turner (2010). "Experimental evidence that source genetic variation drives pathogen emergence." Proceedings of the Royal Society B: Biological Sciences **277**(1697): 3113-3121.

Devilly, G. (2004). "The effect size generator for windows: Version 2.3 (computer programme)." Centre for Neuropsychology, Swinburne University, Australia.

Dias, P. C. (1996). "Sources and sinks in population biology." Trends in Ecology & Evolution **11**(8): 326-330.

Garant, D., S. E. Forde and A. P. Hendry (2007). "The multifarious effects of dispersal and gene flow on contemporary adaptation." Functional Ecology **21**(3): 434-443.

Holt, R. D. (1997). "On the evolutionary stability of sink populations." Evolutionary Ecology **11**(6): 723-731.

Holt, R. D. (2003). "On the evolutionary ecology of species' ranges." Evolutionary ecology research **5**(2): 159-178.

Holt, R. D. and R. Gomulkiewicz (1997). "How does immigration influence local adaptation? A reexamination of a familiar paradigm." The American Naturalist **149**(3): 563-572.

Karve, S., S. Daniel, Y. Chavhan, A. Anand, S. Kharola and S. Dey (2015). "E scherichia coli populations in unpredictably fluctuating environments evolve to face novel stresses through enhanced efflux activity." Journal of evolutionary biology **28**(5): 1131-1143.

Karve, S. M., D. Bhave, D. Nevgi and S. Dey (2016). "Escherichia coli populations adapt to complex, unpredictable fluctuations by minimizing trade-offs across environments." Journal of evolutionary biology **29**(12): 2545-2555.

Kassen, R. (2014). Experimental evolution and the nature of biodiversity. Colorado, United States, Roberts.

Kawecki, T. J. and D. Ebert (2004). "Conceptual issues in local adaptation." Ecology letters **7**(12): 1225-1241.

Kawecki, T. J. and R. D. Holt (2002). "Evolutionary consequences of asymmetric dispersal rates." The American Naturalist **160**(3): 333-347.

Kirkpatrick, M. and N. H. Barton (1997). "Evolution of a species' range." The American Naturalist **150**(1): 1-23.

Lagator, M., A. Morgan, P. Neve and N. Colegrave (2014). "Role of sex and migration in adaptation to sink environments." Evolution **68**(8): 2296-2305.

Lawrence, D., T. Bell and T. G. Barraclough (2016). "The effect of immigration on the adaptation of microbial communities to warming." The American Naturalist **187**(2): 236-248.

Lenormand, T. (2002). "Gene flow and the limits to natural selection." Trends in Ecology & Evolution **17**(4): 183-189.

Levins, R. (1968). Evolution in changing environments. New Jersey, Princeton University Press.

Lively, C. (1999). "Migration, virulence, and the geographic mosaic of adaptation by parasites." The American Naturalist **153**(S5): S34-S47.

McDonald, M. J., Y.-Y. Hsieh, Y.-H. Yu, S.-L. Chang and J.-Y. Leu (2012). "The evolution of low mutation rates in experimental mutator populations of Saccharomyces cerevisiae." Current Biology **22**(13): 1235-1240.

Morgan, A. D., S. Gandon and A. Buckling (2005). "The effect of migration on local adaptation in a coevolving host–parasite system." Nature **437**(7056): 253-256.

Orr, H. A. (2009). "Fitness and its role in evolutionary genetics." Nature Reviews Genetics **10**(8): 531-539.

Perron, G. G., A. Gonzalez and A. Buckling (2007). "Source–sink dynamics shape the evolution of antibiotic resistance and its pleiotropic fitness cost." Proceedings of the Royal Society B: Biological Sciences **274**(1623): 2351-2356.

Reboud, X. and G. Bell (1997). "Experimental evolution in Chlamydomonas. III. Evolution of specialist and generalist types in environments that vary in space and time." Heredity **78**(5): 507-514.

Sexton, J. P., P. J. McIntyre, A. L. Angert and K. J. Rice (2009). "Evolution and ecology of species range limits." Annual Review of Ecology, Evolution, and Systematics **40**: 415-436.

Sullivan, G. M. and R. Feinn (2012). "Using effect size—or why the P value is not enough." Journal of graduate medical education **4**(3): 279-282.

Van Cleve, J. and D. B. Weissman (2015). "Measuring ruggedness in fitness landscapes." Proceedings of the National Academy of Sciences **112**(24): 7345-7346.

Venail, P. A., R. C. MacLean, T. Bouvier, M. A. Brockhurst, M. E. Hochberg and N. Mouquet (2008). "Diversity and productivity peak at intermediate dispersal rate in evolving metacommunities." Nature **452**(7184): 210-214.

Vogwill, T., A. Fenton and M. A. Brockhurst (2011). "Coevolving parasites enhance the diversity-decreasing effect of dispersal." Biology letters **7**(4): 578-580.

Watkinson, A. R. and W. J. Sutherland (1995). "Sources, sinks and pseudo-sinks." Journal of Animal Ecology: 126-130.

Yeaman, S. and F. Guillaume (2009). "Predicting adaptation under migration load: the role of genetic skew." Evolution **63**(11): 2926-2938.