1

7 Urbanization alters phenology of plant reproduction, foliar infection, and herbivory

9 Quinn N. Fox*, Mahal J. Bugay, Eleanor Grant, Olivia Shaw, Keiko Farah,

10 and Rachel M. Penczykowski

12 Department of Biology, Washington University in St. Louis, St. Louis, Missouri, 63130, USA

14 *Corresponding author: qfox@wustl.edu

Abstract

1. Urbanization is associated with numerous environmental changes that may affect the timing of plant reproduction and foliar damage by pathogens, herbivores, and human activities. Yet such effects have not previously been quantified simultaneously in plant populations across levels of urbanization. Thus, it remains unclear how urbanization impacts multiple fitness components of a single plant species. Moreover, it is unclear which effects are consistent between species.

2. We studied 22 populations of two co-occurring *Plantago* species in parks and nature areas spanning rural, suburban, and urban sections of the Greater St. Louis metropolitan area in the central United States. In monthly surveys of *Plantago lanceolata* and *P. rugelii* during summer–fall in two consecutive years, we quantified plant reproductive development and prevalence of powdery mildew infection. In the second year, we additionally quantified prevalence of insect herbivory and mowing damage and placed potted sentinel plants into the field populations to directly measure rates of infection and herbivory.

3. Urbanization was associated with earlier flowering and more seed production for *P. rugelii*, but less seed maturation for *P. lanceolata*. Mildew epidemics on *P. rugelii* started earlier and achieved greater infection prevalence in more urban sites. Correspondingly, sentinels only became infected in suburban and urban sites. By contrast, there was less mildew infection on *P. lanceolata*, including on sentinels, suggesting low availability of pathogen genotypes able to infect this species. Urbanization accelerated early-summer herbivory on both plant species, but did not affect herbivory of wild or sentinel plants later in summer. Finally, prevalence of mowing damage on leaves was generally greater in urban and suburban than rural sites.

4. Synthesis: We show that urbanization affects the timing of key reproductive events and multiple types of aboveground damage for wild herbaceous plants. The two plant species

responded differently to urbanization in terms of their reproductive phenology. However, in

months when significant effects of urbanization were detected, they were typically in the

direction of greater rates of pathogen infection, herbivory, and mowing damage. Overall, these

findings suggest multiple mechanisms by which urbanization may impact plant population

growth.

Key words: fungal pathogen; global change ecology; human disturbance; plant development and

life-history traits; plant phenology; plant-herbivore interactions; plant-pathogen interactions

Introduction

Life on Earth is becoming increasingly urbanized. As of 2018, 55% of the world's

population resides in urban areas, and that metric is anticipated to grow to 68% by 2050 (United

Nations, 2018). The process of urbanization involves dramatic changes to the environment,

typically including increases in habitat fragmentation, frequency of disturbance, coverage by

impervious surfaces, temperature, pollution, and introduction of invasive species (Johnson &

Munshi-South, 2017). These changes can alter ecological and evolutionary processes that impact

the fitness of organisms. For example, a meta-analysis of more than 100 published studies

revealed that wildlife face significantly greater risk of parasite infection in urban areas (Murray

et al., 2019). However, effects of urbanization on plant fitness are poorly understood (Johnson &

Munshi-South, 2017; Rivkin et al., 2019). Notably, only a single study has tested effects of

urbanization on tree diseases (van Dijk et al., 2022), and none have tested effects of urbanization

on diseases of wild herbs. Moreover, no studies have simultaneously quantified effects of

urbanization on timing of plant reproduction and damage from pathogens, herbivores, and human

activities.

One well documented consequence of urbanization is the heat island effect, in which

cities are warmer than surrounding environments due to retention of solar radiation by urban

materials and release of heat from high densities of vehicles and other machinery (Zhou et al.,

2017). These increased temperatures can impact plant growth and development. In a study of

herbarium records, 24% of herbaceous and woody plant species flowered earlier in urban

environments (Neil et al., 2010). However, whether urban heating advances plant phenology

depends on baseline climate in the region. For example, flowering date advances with city size

only in regions with cool or cold winters, where degree-days are limiting to development (Li et

al., 2019). Moreover, heat stress beyond a plant's thermal optimum can reduce shoot and root

growth (Desaint et al., 2021). Quantifying the impact of urban heating on plant growth and

development is necessary for understanding consequences of land use change. In addition,

studies conducted along spatial temperature gradients can provide insight into how plants will

respond to future climate warming (De Frenne et al., 2013).

Urban heating may increase or decrease prevalence of plant diseases, depending on

regional climate, magnitude of the urban heat island effect in a given city, and thermal sensitivity

of host and pathogen physiological traits. Plant pathogens include fungi, oomycetes, bacteria,

and viruses. Here, we focus on fungi because these are the most well studied in wild plant

populations (Alexander, 2010; Burdon & Laine, 2019). Across life history processes, plant-

associated fungi typically have temperature minima of 5-15 °C, optima of 18-28 °C, and maxima

of 26-38 °C (Chaloner et al., 2020). In regions where baseline climate is cooler than optimal for a

pathogen, urban heating may promote disease. Indeed, urban heat is linked to increased severity

of powdery mildew on English oak in Europe (van Dijk et al., 2022). However, in regions that

are already warmer than optimal for a pathogen, additional heat in cities may inhibit pathogen

growth. Effects of urban heat on disease can further depend on microclimate heterogeneity

(Penczykowski et al. 2018), temporal variation in climate (Egerer et al., 2020), and thermal

sensitivity of plant defense responses (Desaint et al., 2021; Wang et al., 2009).

In addition to being warmer, green spaces are often more fragmented in urban than rural

environments. This fragmentation may cause populations of some plant and pathogen species to

be more isolated in urban environments (Dubois & Cheptou, 2017). Alternatively, plant or

pathogen species that grow or disperse along roadways may be more highly connected in cities

with denser road networks. For example, fungal infection is more likely in populations of ribwort

plantain along roadsides, particularly at hubs of a road network, suggesting that vehicular traffic facilitates spore transport (Numminen & Laine, 2020). Similarly, roads enhance transmission of an oomycete pathogen between populations of Port Orford cedar (Jules et al., 2002). Thus, increasing road density and traffic volume from rural to urban environments could drive increases in fungal infection prevalence.

Rates of insect herbivory can also be altered by abiotic (e.g., temperature, light, and sound) and biotic (e.g., biodiversity of plants, insects, and predators) features of cities (Miles et al., 2019). For example, urban warming may accelerate emergence of insect herbivores in spring (Bale et al., 2002). If host plant phenology is similarly accelerated, this should lead to higher rates of herbivory earlier in the growing season. By contrast, if urban heating causes phenological asynchrony (e.g., insects emerge before their host plants, or after plants have developed morphological or chemical defenses), we would expect less herbivory (Bale et al., 2002). From the few published studies linking urbanization to herbivory, a variety of patterns have been documented (Miles et al., 2019). Positive relationships between urbanization and herbivory have been shown for wolf apple trees in Brazil (Cuevas-Reyes et al., 2013), for red maples in the southeastern United States (Just et al., 2019), and for English oaks in Europe (Moreira et al., 2019). Contrastingly, in an analysis of herbarium samples of three woody species and one understory herb collected over 112 years in the northeastern United States, urbanization was negatively correlated with insect abundance and herbivory (Meineke et al., 2019). Other evidence for changes in herbivory pressure with urbanization comes from clines in production of an antiherbivore chemical defense (hydrogen cyanide, HCN) in white clover (Johnson et al., 2018; Santangelo et al., 2022). Specifically, 75 of 160 cities around the world show clines in white clover HCN production along urban–rural transects, and most clines (62 of 160 cities) are

in the direction of decreasing HCN production with urbanization (Santangelo et al., 2022).

However, as for plant–pathogen interactions, so few studies have directly quantified rates of

herbivory on herbaceous plants across urbanization gradients that the generality of such trends

across plant species are unknown.

The frequency and intensity of disturbance from human activities including walking and

lawn mowing may also increase with urbanization. For herbs growing along paths, roadways, or

lawns, human activities may damage or remove leaves or reproductive tissues directly, or affect

the probability of damage by pathogens or herbivores. For example, human footsteps increased

severity of fungal anthracnose infections on annual bluegrass on a golf course (Roberts &

Murphy, 2014). Height of mower blades has also been shown to influence disease severity on

golf course grass (Inguagiato et al., 2009; Williams et al., 2001). Yet effects of walking and

mowing on plant populations across levels of urbanization remain to be explored.

Here, we tested effects of urbanization on phenology of plant reproduction and three

types of leaf damage (fungal infection, insect herbivory, and lawn mowing) across a 56-km land

use gradient in the Greater St. Louis metropolitan area in Missouri, United States (Fig. 1, Table

S1). We did this through monthly field surveys of *Plantago lanceolata* (ribwort plantain) and *P.*

rugelii (blackseed plantain) populations co-occurring in 22 sites in summer–fall 2019 and 2020.

In a separate 2020 field experiment, we placed groups of healthy, pathogen-free "sentinel plants"

of these two species into a subset of sites to directly measure rates of infection and herbivory

across the urbanization gradient. We hypothesized that urbanization would be associated with:

(1) earlier flowering and seed production, (2) earlier and larger powdery mildew epidemics, (3)

earlier and overall more prevalent insect herbivory, and (4) more mowing damage on leaves. We

further hypothesized that rates of infection and herbivory on sentinels would positively correlate

with prevalence of infection and herbivory in the field populations where they were placed.

Materials and Methods

Study system

This study focused on two co-occurring herbs, *Plantago lanceolata* (ribwort plantain) and

Plantago rugelii (blackseed plantain). These short-lived perennials grow as rosettes in pastures

or human-disturbed landscapes (e.g., lawns and roadsides), and they are tractable model

organisms for studies of plant–pathogen and plant–herbivore interactions across land use

gradients (Penczykowski & Sieg, 2021). *Plantago lanceolata* is native to Eurasia but

cosmopolitan in its distribution, and *P. rugelii* is endemic to eastern North America. We did not

include the similarly weedy and globally abundant *Plantago major* (common plantain) in our

study, because it is rare in our focal geographic region (Penczykowski & Sieg, 2021;

Yatskievych & Steyermark, 1999).

Both species produce flower stems bearing spikes of many small, wind-pollinated

flowers. Individuals can flower repeatedly over the growing season. The flowers are typically

monecious, though *P. lanceolata* are sometimes gynodioecious (Ross 1973). Fertilization of the

flowers can only occur via outcrossing for *P. lanceolata* (i.e., it is self-incompatible), while *P.*

rugelii can successfully self-pollinate (Hawthorn, 1974; Ross, 1973; Sagar & Harper, 1964).

Fruits then develop within seed capsules that can be visually assessed for maturity.

The most globally and locally common foliar fungal pathogens of *Plantago* are the

specialist powdery mildews *Podosphaera plantaginis* (Castagne; U. Braun and S. Takamatsu)

and *Golovinomyces sordidus* (L. Junell) V.P. Heluta (Braun & Cook, 2012). Powdery mildews –

obligate fungal pathogens in the order Erysiphales – grow on the surface of leaves and extract

nutrients from their host's epidermal tissue (Bushnell, 2002). Chains of asexual spores produced

from mycelia on the leaf surface give infected leaves a white, dusty appearance. The spores are

passively transmitted via wind, and more than 90% land within 2 m of the source plant (Tack et

al., 2014). However, occasional long-distance spore transport allows for pathogen persistence at

the regional scale (Ovaskainen & Laine, 2006). Powdery mildews overwinter via sexual resting

structures that release spores when conditions are favorable in spring (Tack & Laine, 2014).

Plantago are also host to numerous invertebrate herbivores (Penczykowski & Sieg,

2021). Here, we focus on two types of distinctive insect herbivory: leaf mines and chewing

damage. Leaf mines on *Plantago* are created by larvae of the leaf miner fly *Phytomyza*

plantaginis (Penczykowski & Sieg, 2021). Chewing damage on leaf edges or between leaf veins

could be due to a variety of arthropod taxa, particularly larval lepidopterans. In the region of our

study, these include specialist common buckeye caterpillars (*Junonia coenia*) as well as

generalist species (Penczykowski & Sieg, 2021).

Focal populations

We investigated the relationships between urbanization and plant phenology, powdery

mildew infection, and insect herbivory in a total of 22 sites with co-occurring *P. lanceolata* and

P. rugelii. These sites were parks and nature areas extending from downtown St. Louis City to

the southwest along the I-44 interstate highway corridor (Fig. 1a, Table S1). In a few large parks

or nature areas, we surveyed two sites separated by at least 0.45 km. The study began with 19

sites in July 2019. We added two sites at Shaw Nature Reserve in August and September 2019,

and we added a site at Forest Park in June 2020 (Table S1).

We classified the eight sites within St. Louis City as "urban", eight sites in St. Louis

County east of Missouri Route 141 as "suburban", and six sites west of Missouri Route 141 as

"rural". These classifications were based on a principal component analysis (PCA) of

environmental and spatial variables performed with the 'prcomp' function in R version 3.6.2 (R

Core Team, 2019). The PCA included average temperature from June through October (Temp),

nighttime radiance (NightLight), percent impervious area (%Imp), percent tree cover (%Tree),

and estimated intersection density of walkable roads (IntDen; walkable roads defined as having

speed limits between 8-55 miles per hour) (Fig. 1b). The temperature data represented long-term

(1970-2000) averages at these sites during the focal months of our study (WorldClim 2; Fick &

Hijmans, 2017). Nighttime radiance data came from the NASA Black Marble data product for

December 2020 (NASA Worldview).

For most sites, percent impervious area, percent tree cover, and intersection density were

obtained through the EnviroAtlas Community dataset for St. Louis (US EPA, 2021a,b,c).

However, some rural sites were located outside the boundaries of that urban-centered dataset.

Thus, for rural sites #1, 2, and 3 (Table S1), we estimated percent impervious area and percent

tree cover from the National Land Cover Database (MRLC, 2019). Specifically, we used the

Summarize Categorical Raster tool in ArcGIS Pro to extract land use pixel count data from the

NLCD layer for the census blocks containing the sites. We calculated percent impervious area as

percent of pixels in low, medium, or high intensity development classes. Similarly, we calculated

percent tree cover as the percent of pixels in deciduous forest, evergreen forest, mixed forest,

shrub scrub, or woody wetlands classes. Intersection density data was not available through

EnviroAtlas for rural sites #1, 2, 3, 5, or 6 (Table S1). For these, we estimated intersection

density using the streets layer from the ESRI Streetmap Premium dataset as a base (ESRI, 2020).

After limiting to streets with speed limits between 6-55 mph, we manually digitized all

intersections within a 750-m radius of the focal sites, and used the Kernel Density tool to

generate an intersection density layer. From that layer, we extracted values of intersection

density at the study sites, following methods in the EnviroAtlas metadata (US EPA, 2021c).

Field survey methods

We performed monthly surveys in the focal populations from July to October 2019 (four

surveys) and from June to October/November 2020 (five surveys). This time frame was chosen

to represent the majority of the growing season of the plants, their fungal pathogens, and their

insect herbivores. In each survey, we collected data for 50 individuals each of *P. lanceolata* and

P. rugelii. On a meandering walk through each population, we stopped every few steps and

selected the nearest plant of either species that was at least 1 m from the previously surveyed

individual of that same species. The average sizes of the surveyed areas were 1004, 1714, and

2699 m^2 for rural, suburban, and urban sites, respectively (Fig. S1; sizes estimated from

polygons in Google Earth). Rural survey areas tended to be smaller because those sites were

typically clearings bounded by tall grassland or woodlands where *Plantago* do not grow. By

contrast, urban and suburban sites typically had larger total areas with *Plantago* present, and our

surveys covered more ground.

For each selected plant, we recorded the status of the most mature flower stem on the

plant, if present, using the following ordered categories: (0) no flower stem or stem present but

flowers/seeds removed by mowing, (1) budding, (2) flowering, (3) immature seeds, (4) mature

seeds, or (5) seeds dispersed. Seeds were categorized as immature if the seed capsules were

green and plump, and mature if they were brown and papery. We recorded the powdery mildew

infection status for each plant as the presence or absence of conspicuous white mycelia and/or

conidia on the leaf surface. In 2020, to assess whether the infection statuses of the selected plants

were representative of nearby conspecifics, we additionally recorded the numbers of infected

conspecifics within a 1.5-m radius of each focal individual, using the following categories: (0) 0,

(1) 1-10, (2) 11-50, (3) 51-100, and (4) 101 or more plants. In 2020, we also recorded the

presence or absence of two main types of herbivory damage (chewing and leaf mines on leaves)

and mowing damage (straight cuts across leaves).

Sentinel experiment

In July 2020, we placed healthy, greenhouse-grown "sentinel" plants into 15 of our field

survey sites to directly test rates of powdery mildew infection and herbivory across the

urbanization gradient. These sites (five each of urban, suburban, and rural) were chosen from the

full set of 22 sites based on presence of mildew on at least one of the *Plantago* species in 2019.

We selected sites based on that criterion to avoid biasing the outcome of the experiment towards

rejection of the null hypothesis that infection prevalence would be the same across site types.

Sentinel plants were grown from *P. lanceolata* and *P. rugelii* seeds collected from wild,

uninfected plants in St. Louis City and County during July–October 2019. To capture variation in

plant genetic background, we planted seeds from six maternal lines of *P. lanceolata*, and five of

P. rugelii. Each maternal line consisted of a single seed spike from a mother plant. Seeds were

sown in BM6 All-Purpose soil (Berger) on 16 May 2020 in the Jeanette Goldfarb Plant Growth

Facility at Washington University in St. Louis. All six *P. lanceolata* lines but only three of the *P.*

rugelii lines successfully germinated. On 1 June, we moved the seedlings to a hoop house at

Tyson Research Center (Eureka, Missouri) and transplanted each into a 4.5-in diameter pot. To

ensure that sentinels remained uninfected before deployment into the field, we covered each with

an autoclaved, spore-proof pollination bag (PBS International, model 3D.55), which we secured

around the pot with a silicone band. Three times per week, we placed the sentinels in a shallow

pool of water for 10 min, allowing water to soak up through the base of the pot.

Sentinels were deployed into field sites between 7-9 July 2020. Due to much higher

germination success for *P. lanceolata* than *P. rugelii*, we were able to place 20-23 *P. lanceolata*

into each of 15 sites (n = 322) but only 3-4 *P. rugelii* into each of 9 sites (n = 36). Within sites,

sentinels were randomly divided between three seedling trays at the base of a single tree (i.e., all

sentinels were at one shaded location per site). We removed the pollination bags and collected

initial data on the number of leaves, length and width of largest leaf, and flower maturity stage.

We confirmed that no sentinels were infected with powdery mildew, and recorded any existing

damage on leaves (e.g., from greenhouse or hoop house insects encountered prior to enclosure

with pollination bags). We also recorded the distance from the trays of sentinels to the nearest

infected *Plantago* in the surrounding wild population. We placed a temperature datalogger

(HOBO MX2201) just below the soil surface in one pot per tray to monitor temperature

differences between sites. The trays were filled with an inch of water at the time of deployment

and once the following week. It did not rain during the week of the experiment, so this watering

scheme achieved healthy moisture levels in the pots.

Between 14-16 July 2020, sentinels were retrieved from the field sites in the same order

they were deployed. Before removal from the field, we recorded the number of leaves, length

and width of largest leaf, and presence/absence of chewing and leaf miner herbivory for each

leaf. Then we re-covered each sentinel with a pollination bag and moved them to a common

location 5 km from the nearest field site to allow any powdery mildew infections to develop for

another week. During that week, we watered plants twice (in shallow pools for 10 min), and at

the end of the week we recorded infection status of each leaf.

Statistical analyses

All statistics were performed in R version 3.6.2 (R Core Team, 2019). Field survey data

were analyzed separately for each species and year. In models testing effects of survey month

(ordered factor), site type (urban, suburban, or rural), and their interaction on a given response

variable, we included site (population) identity as a random effect. We performed post-hoc

Tukey's tests of contrasts between site types within each month if there was at least a marginally

significant ($P < 0.10$) site type x month interaction, or between site types averaged across months

if there was a significant ($P < 0.05$) main effect of site type but no interaction (package

'emmeans'; Lenth, 2022). We report all site type contrasts with $P < 0.10$, reserving the word

"significant" for $P < 0.05$.

We analyzed plant reproductive development stage as an ordered categorical response

variable (levels 0–5, as explained above in *Field survey methods*) in cumulative link mixed

models (package 'ordinal'; Christensen, 2019). Prevalences of powdery mildew infection, leaf

mines, and mowing were logit-transformed and analyzed in linear mixed effects models

(package 'nlme'; Pinheiro et al., 2012) with first order autocorrelation structure. For prevalence

values of zero or one, we added (zeros) or subtracted (ones) 0.0001 before logit-transforming.

We detected no temporal autocorrelation in prevalence of chewing herbivory, so this was

analyzed in a generalized linear mixed model (GLMM, package 'lme4'; Bates et al., 2015) with

binomial error distribution and logit link function.

To test if the infection statuses of the surveyed plants were representative of nearby

conspecifics, we used binomial generalized linear models (logit link). Specifically, we modelled

binary infection status of surveyed plants in response to the categorical abundance of infected

conspecifics in the surrounding 1.5 m radius, with site identity as a random effect.

In the sentinel experiment, we analyzed the proportion of leaves on each plant with

powdery mildew, leaf mines, and chewing damage (i.e., "severity" of each damage type) using

binomial GLMMs (logit link) with random effects of tray identity nested within site identity

(because there were three trays of sentinels in each field site). In addition to testing effects of site

type on infection and herbivory severity, we included covariates reflecting the initial size of

sentinels. Specifically, these were scores on the first two principal components axes from a PCA

of number of leaves and length and width of longest leaf at the start of the experiment. For *P.*

lanceolata, the first principal component axis (PC1) explained 65.06% of variation in initial

plant size data and the second (PC2) explained 34.72%; for *P. rugelii*, PC1 explained 75.36%

and PC2 explained 21.92% of variation. For analysis of infection severity, we included distance

to nearest infected wild plant as a covariate. For analyses of severity of leaf mines and chewing,

we additionally included the initial severity (chewing) or presence/absence (mines) of these

damage types as covariates, to account for any herbivory that occurred prior to placement in the

field. For both infection and herbivory, we tested whether prevalence on sentinels was correlated

with prevalence in the wild populations where they had been placed.

Results

Field site characteristics

In the PCA of environmental variables at the study sites, the first two principal

components (PC1 and PC2) together explained 89.85% of the total variation in the data (Fig. 1b).

The pre-classified site types (rural, suburban, and urban) were primarily separated along PC1,

which was associated with greater temperature, nighttime radiance, intersection density, and

percent impervious area, and with lower percent tree cover (Fig. 1b). Variation within site types

occurred primarily along PC2, which was associated with greater temperature, nighttime

radiance, and percent tree cover, and with lower percent impervious area (Fig. 1b).

Field surveys: plant reproductive phenology

In both 2019 and 2020, *P. lanceolata* populations had begun flowering in all site types

before our first survey, the frequency of flowering peaked in July, and frequency of mature

and/or dispersed seeds peaked in August (Fig. 2a,c). For *P. lanceolata* in 2019, parameters could

not be uniquely determined for the model of reproductive development through time when

October data were included. Excluding October 2019 from the analysis, the effect of site type on

P. lanceolata reproductive development (as an ordered categorical variable) varied through time

in both years (site type x month interaction in 2019: Likelihood Ratio [LR] χ^2 = 20.40, df = 4, P

= 0.0004; in 2020: LR χ^2 = 51.52, df = 8, P < 0.0001). Post-hoc Tukey's tests showed that, in

2019, maturity was marginally lower in urban and/or suburban than rural sites in July (urban-

rural: P = 0.050, suburban-rural: P = 0.068) and August (suburban-rural: P = 0.060), with no

other differences between site types within a given month (P > 0.10 for all unreported pairwise

contrasts; Fig. 2a). In 2020, *P. lanceolata* reproductive development was marginally lower in

urban than rural sites in August (P = 0.079) and significantly lower in urban than either suburban

or rural sites in September (urban-rural: P = 0.036, urban-suburban: P = 0.008; Fig. 2c).

In both years, flowering continued at a higher rate into August for *P. rugelii* compared to

P. lanceolata, and overall rates of seed production and maturation were also much higher for *P.*

rugelii (Fig. 2). The effect of site type on *P. rugelii* reproductive development varied through

time in both years (site type x month interaction in 2019: LR χ^2 = 115.72, df = 6, P < 0.0001; in

2020: LR χ^2 = 128.98, df = 8, P < 0.0001; Fig. 2b,d). Post-hoc tests revealed that *P. rugelii*

development was significantly lower in suburban than either rural (P = 0.0092) or urban (P =

0.0002) sites in September 2019, but increased with level of urbanization in October 2019

(suburban-rural: P = 0.0001, urban-rural: P < 0.0001, urban-suburban: P = 0.009; Fig. 2b). Our

June 2020 survey captured significantly earlier *P. rugelii* development with increasing levels of

urbanization (suburban-rural: P = 0.018, urban-rural: P < 0.0001, urban-suburban: P = 0.0011;

Fig. 2d). Maturity of *P. rugelii* was greater in urban than rural sites in both August and

September 2020 (P = 0.025 and P = 0.043, respectively), and greater in both suburban and urban

than rural sites in October/November 2020 (P < 0.0001 for both contrasts).

Field surveys: powdery mildew infection

There was very little powdery mildew infection on *P. lanceolata* in 2019, and generally

small epidemics in 2020 (Fig. 3a,c). For this species, there were no significant site type x month

effects on infection prevalence (in 2019: LR χ^2 = 4.00, df = 6, P = 0.68; in 2020: LR χ^2 = 10.34,

df = 8, P = 0.24), nor significant main effects of site type (in 2019: LR χ^2 = 2.43, df = 2, P =

0.30; in 2020: LR χ^2 = 3.36, df = 2, P = 0.19). There was a significant main effect of month in

2020 only (in 2019: LR χ^2 = 3.92, df = 3, P = 0.27; in 2020: LR χ^2 = 47.64, df = 4, P < 0.0001).

There were large powdery mildew epidemics on *P. rugelii* in both years, and epidemics

started earlier in suburban and urban populations (Fig. 3b,d). In both years, *P. rugelii* infection

prevalence was significantly affected by site type (in 2019: LR χ^2 = 8.89, df = 2, P = 0.012; in

2020: LR χ^2 = 22.00, df = 4, P < 0.0001) and month (in 2019: LR χ^2 = 22.57, df = 3, P < 0.001;

in 2020: LR χ^2 = 70.42, df = 4, P < 0.001), but there were no site type x month interactions (in

2019: LR χ^2 = 4.39, df = 6, P = 0.62; in 2020: LR χ^2 = 3.90, df = 6, P = 0.87). For 2019, post-hoc

testing revealed significantly greater infection prevalence in urban than rural sites (suburban-

rural: P = 0.067, urban-rural: P = 0.037; Fig. 3b). In 2020, there was significantly greater

infection prevalence in both suburban and urban relative to rural sites (suburban-rural: P =

0.0061, urban-rural: P = 0.0007; Fig. 3d).

The infection status of surveyed plant individuals was consistent with that of nearby

plants in the population. Specifically, presence/absence of powdery mildew infection on

surveyed plants was significantly positively related to the abundance of infected conspecifics

within a 1.5-m radius (*P. lanceolata:* χ^2 = 262.81, df = 3, P < 0.0001; *P. rugelii:* LR χ^2 = 1097.6,

df = 3, P < 0.0001; Fig. S2).

Field surveys: herbivory

For both species, prevalence of leaf mines depended on the interaction of site type and

month (*P. lanceolata*: χ^2 = 16.37, df = 8, P = 0.037; *P. rugelii*: χ^2 = 55.84, df = 8, P < 0.0001;

Fig. 4a,b), and the only significant differences between site types occurred in June. For *P.*

lanceolata, there was significantly greater prevalence of leaf mines in urban than suburban sites

in June (urban-rural: P = 0.092, urban-suburban: P = 0.0083). For *P. rugelii*, there was greater

prevalence of leaf mines in urban than either suburban or rural sites in June (urban-rural: P =

0.0017, urban-suburban: P = 0.0001).

Chewing-type herbivory also depended on the interaction between site type and month

(*P. lanceolata*: χ^2 = 66.17, df = 8, P < 0.0001; *P. rugelii*: χ^2 =88.51, df = 8, P < 0.0001; Fig.

4c,d). For both species, there was greater prevalence of chewing damage in urban than suburban

or rural sites in June (*P. lanceolata*: urban-rural: P < 0.0001, urban-suburban: P < 0.0001; *P.*

rugelii: urban-rural: P < 0.0001, urban-suburban: P < 0.0001).

Field surveys: mowing

For *P. lanceolata*, prevalence of mowing damage on leaves depended on site type ($\chi^2 =$

7.50, df = 2, P = 0.024) and month ($\chi^2 = 43.10$, df = 4, P < 0.0001), but not their interaction ($\chi^2 =$

10.28, df = 8, P = 0.25). For this species, there was marginally greater mowing prevalence in

suburban and urban than rural sites (suburban-rural: P = 0.049, urban-rural: P = 0.083; Fig. 5a).

For *P. rugelii*, there was a marginally significant site type x month effect on prevalence

of mowing damage (site type x month: $\chi^2 = 13.79$, df = 8, P = 0.087; site type: $\chi^2 = 8.56$, df = 2,

P = 0.014; month: $\chi^2 = 23.44$, df = 4, P = 0.0001; Fig. 5b). This weak interactive effect reflected

marginally greater mowing prevalence in suburban and urban relative to rural sites in June

(suburban-rural: P = 0.034, urban-rural: P = 0.062) and significantly greater mowing prevalence

in suburban and urban relative to rural sites in the final October/November survey (suburban-

rural: P = 0.0020, urban-rural: P = 0.0025; Fig. 5b).

Sentinel experiment: powdery mildew infection and herbivory

All 322 *P. lanceolata* sentinels survived to the end of the experiment, and only one

became infected with powdery mildew (Fig. 6a). This infected sentinel happened to be placed in

Buder South Park within 1.5 m of the only wild infected *P. lanceolata* individual we found after

a thorough search at that site (Fig. S3; note that the one wild infected plant was not among the 50

plants surveyed at that site in July 2020, so infection prevalence of wild plants was recorded as

zero for that site-date; Fig. 6a). No *P. lanceolata* sentinels became infected in Tilles Park, which

did have a small number of infected wild plants at that time (Fig. 6a), likely because the sentinels

were located almost 6 m from the nearest wild infected plant (Fig. S3).

Many *P. lanceolata* sentinels experienced herbivory during their week in the field;

however, there were no differences between site types for severity (proportion of leaves

damaged) of either leaf mines ($\chi^2 = 4.17$, df = 2, P = 0.12) or chewing ($\chi^2 = 3.82$, df = 2, P =

0.15). Prevalences of leaf mines and chewing (proportion of plants affected) were not correlated

between sentinels and wild *P. lanceolata* in the same sites during July 2020 (leaf mines: r = -

0.03, P = 0.92, Fig. 6c; chewing: r = -0.14, P = 0.62, Fig. 6e).

All 36 *P. rugelii* sentinels survived to the end of the experiment. Eight of these became

infected with powdery mildew in urban sites (Clifton Park and Tilles Park), three became

infected in a single suburban site (Kirkwood Park) and none became infected in rural sites (Fig.

6b). Although this trend was consistent with the greater prevalence of infection observed in more

urban populations of wild *P. rugelii* in 2020 (Fig. 3d), there was no significant correlation

between prevalence of infection on sentinels and wild plants surveyed in those sites during July

(P = 0.95, Fig. 6b). Instead, the proportion of infected *P. rugelii* sentinels increased with

proximity to wild infected plants within the site (P = 0.048, Fig. S3).

Only a single *P. rugelii* sentinel acquired a leaf mine during the field exposure. *Plantago*

rugelii sentinels experienced chewing-type herbivory in most field sites (Fig. 6f), and severity of

chewing damage did not differ between site types ($\chi^2 = 1.25$, df = 2, P = 0.53). Prevalence of

herbivory on *P. rugelii* sentinels was not correlated with that on wild conspecifics in those sites

during July 2020 (leaf mines: r = 0.15, P = 0.71, Fig. 6d; chewing: r = -0.52, P = 0.15, Fig. 6f).

Discussion

We found that urbanization affected the timing of key reproductive events and seasonal prevalence of multiple types of foliar damage for wild herbaceous plants. The co-occurring focal plant species responded differently in terms of their reproductive phenology. Specifically, urbanization was associated with earlier flowering and more seed production for *P. rugelii*, but less seed maturation for *P. lanceolata*. These species-specific effects on reproductive phenology were broadly consistent between the two years. Another consistent trend between years was that powdery mildew epidemics on *P. rugelii* started earlier and achieved greater infection prevalence in more urban sites. Correspondingly, *P. rugelii* sentinels only became infected in suburban and urban sites. By contrast, there was less mildew on *P. lanceolata*, including for sentinels placed into the field sites. There was no significant effect of urbanization on infection prevalence for *P. lanceolata*. Urbanization accelerated early-summer herbivory on both plant species, but did not affect herbivory of wild or sentinel plants later in summer. Finally, prevalence of mowing damage on leaves was generally greater in urban and suburban than rural sites.

Urbanization accelerated reproductive development of *P. rugelii*, consistent with trends for many other species (e.g., Neil et al., 2010; Li et al. 2019). However, there were neutral-to-negative effects of urbanization on reproductive development of *P. lanceolata*. One possible explanation for this difference involves the differential impact of mowing on flower stems of these species. *Plantago rugelii* flowers are produced along most of the length of the flower stem, starting below the height of mower blades. If the upper portion of the flower spike had been removed by mowing, we recorded the maturity of the flowers or seeds that remained. By contrast, *P. lanceolata* produces taller stems with flowers produced only at the tip, where they are highly susceptible to complete removal by lawn mowing. Indeed, stems with completely

removed flower spikes were more common for *P. lanceolata* than *P. rugelii* (Fig. 2). Therefore,

more frequent mowing in suburban and urban sites may have reduced the abundance of flower

and seed spikes on *P. lanceolata* but not affected the proportion of *P. rugelii* individuals with

flowers or seeds. How variation in frequency and/or height of mowing impacts plant

reproductive phenology across urbanization gradients is an open question. Given the consistently

shorter length of the flowering season of *P. lanceolata* compared to *P. rugelii*, their different

responses to urbanization may also be explained by differences in their responses to

environmental cues including temperature and light. Experimental tests of these possible

underlying mechanisms are needed to clarify why urbanization has species-specific effects on

reproduction. Moreover, for some plants, later flowering and senescence in urban sites is

adaptive (Lambrecht et al., 2016). Thus, future experiments testing the genetic basis for

differences in phenology between *Plantago* populations will contribute to our understanding of

how plant reproductive traits evolve in cities.

Urbanization was associated with earlier and overall larger powdery mildew epidemics

on *P. rugelii*. There are several non-mutually exclusive possible reasons for that trend. Two of

those possible reasons involve temperature. First, mildew survival is decreased by exposure to

freezing air temperatures (Penczykowski et al., 2015). Therefore, the urban heat island effect

may have increased mildew overwinter survival. Second, warmer temperatures during the

growing season can promote mildew growth and transmission. Because all the field sites were

parks and nature areas, the magnitude of the urban heat island effect experienced by our focal

plants might have been tempered by the presence of trees and other vegetation. However, soil

temperatures in the pots of sentinel plants increased from rural to suburban to urban sites (Fig.

S4), consistent with long-term trends in this region (Fig. 1b). Thus, as for powdery mildew on

oaks (van Dijk et al., 2022), warmer temperatures may have contributed to the increased

infection prevalence of *P. rugelii* in suburban and urban populations.

Our observation of increased mildew prevalence with urbanization could also be due to

greater rates of spore arrival to more highly connected urban sites. Dispersal along roadside

populations could allow mildew to spread between sites over the course of the growing season

(Numminen & Laine, 2020). Vehicular traffic can also facilitate transport of particles including

seeds (Lippe et al., 2013) and spores (Numminen & Laine, 2020). Hence, denser road networks

surrounding urban parks may bring more mildew spores. However, arriving spores are only

successful at infecting if they land on susceptible plant genotypes, and *Plantago* populations can

vary in their susceptibility to arriving mildew strains (Höckerstedt et al., 2018; Jousimo et al.,

2014; Laine, 2004). Thus, greater resistance of rural *P. rugelii* to mildew could also have

decreased their prevalence of infection, even if rates of spore arrival were equally high across the

land use gradient. The fact that none of the *P. rugelii* sentinel plants became infected in rural

sites is consistent with low spore arrival in those sites, though we note that sample sizes of *P.

rugelii* sentinels were small. At the same time, there were lower rates of infection on wild *P.

lanceolata* compared to *P. rugelii* growing in the same sites, as well as near-zero prevalence of

infection among the >300 *P. lanceolata* sentinels placed across the urbanization gradient. These

results suggest species-specificity of the pathogen strains, and lower regional abundance of

spores able to infect *P. lanceolata*. Aerial spore sampling and genotyping of mildew infections

collected from both *Plantago* species across the urbanization gradient will provide further

insights into the connectivity of pathogen populations within and between land use types.

We hypothesized that early-summer prevalence of insect herbivory would be elevated in

urban populations, due to urban heat island effects accelerating insect emergence and foraging

activity in spring. Indeed, both types of herbivory damage were greatest on urban *Plantago* in

our earliest survey month (June). However, we further hypothesized that urban sites would

continue to experience greater herbivory over the course of summer, and that herbivory damage

on sentinel plants would similarly increase with urbanization. These hypotheses for later-summer

patterns are not supported by our data. Instead, prevalence of leaf mines became similar across

site types, peaking at a moderate level in July-August. Meanwhile, prevalence of chewing

damage was sustained at a high level across site types for the remainder of the growing season.

Rates of herbivory on sentinels in July neither followed a trend with urbanization nor correlated

with prevalence of herbivory in the wild populations at that time. In general, our results are

consistent with herbarium evidence that increases in winter temperature are a strong driver of

increases in insect herbivory (Meineke et al., 2019). Yet additional experiments will be needed to

tease apart effects of winter and spring climate variables, artificial light at night, and other

abiotic and biotic facets of urbanization as contemporary drivers of increased early-summer

herbivory on *Plantago* hosts.

As urban areas continue to grow in size and prominence worldwide, there is a societal

need for understanding effects of urban environmental factors on plants (United Nations, 2018).

Urban vegetation provides ecosystem services that mitigate heat, air pollution, and flooding

(Heidt & Neef, 2008). Access to urban green space is also linked to mental and physical health

benefits for humans, including decreased heart rate, increased physical activity, and improved

mood (Kondo et al., 2018). The abundance, health, and connectivity of native and introduced

plant species are also key in supporting diversity of urban fauna (Johnson and Munshi-South,

2017). Given these important roles of urban plants, it is critical to resolve the mechanisms by

which urban environments impact their fitness. In this study, we showed how plant reproductive

development and prevalence of disease, herbivory, and mowing damage varied through time

across an urbanization gradient. Our study is the first to test for differences in fungal disease

prevalence in wild herbaceous plant populations across levels of urbanization. More importantly,

because we quantified effects of urbanization on multiple plant traits and sources of damage

through the growing season, our findings point to several mechanisms by which changes in land

use may impact plant population growth.

Author Contributions

QNF, MJB, and RMP conceived of the study design. QNF, MJB, EG, OSS, and KNF

performed field surveys. QNF, OSS, and KNF performed the sentinel experiment. RMP provided

logistical support for the field surveys and sentinel experiment. QNF and RMP performed

statistical analyses. QNF wrote the first draft, and all authors contributed to subsequent drafts of

the manuscript.

Data Availability

Data and code for analyses will be archived at Dryad upon acceptance of the article.

Acknowledgements

We thank Whitney Anthonysamy, Beth Biro, and Solny Adalsteinsson for help selecting

field sites and Michelle Pollowitz, Armando Sánchez-Conde, Philippa Tanford, and Shayna

Rosenbloom for assistance with field work. Carrie Goodson provided greenhouse and lab

support. Mike Dyer and Kim Medley provided greenhouse and hoop house support on the

Washington University in St. Louis campus and at Tyson Research Center (TRC), respectively.

Susan Flowers provided undergraduate mentoring support through TRC. Bill Winston assisted

with acquisition of land use data. We thank staff at St. Louis City Department of Parks,

Recreation, and Forestry; Forest Park Forever; Tower Grove Park; Webster Groves Parks and

Recreation Department; St. Louis County Parks and Recreation; Kirkwood Parks and Recreation

Department; Missouri Department of Conservation; TRC; Missouri State Parks; and Missouri

Botanical Garden's Shaw Nature Reserve for access to field sites.

References

Alexander, H. M. (2010). Disease in Natural Plant Populations, Communities, and Ecosystems:

Insights into Ecological and Evolutionary Processes. *Plant Disease*, *94*(5), 492–503.

https://doi.org/10.1094/PDIS-94-5-0492

Bale, J. S., Masters, G. J., Hodkinson, I. D., Awmack, C., Bezemer, T. M., Brown, V. K.,

Butterfield, J., Buse, A., Coulson, J. C., Farrar, J., Good, J. E. G., Harrington, R., Hartley,

S., Jones, T. H., Lindroth, R. L., Press, M. C., Symrnioudis, I., Watt, A. D., & Whittaker,

J. B. (2002). Herbivory in global climate change research: Direct effects of rising

temperature on insect herbivores. *Global Change Biology*, *8*(1), 1–16.

https://doi.org/10.1046/j.1365-2486.2002.00451.x

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. *Journal of Statistical Software*, *67*, 1–48.

https://doi.org/10.18637/jss.v067.i01

Braun, U., & Cook, R. T. A. (2012). *Taxonomic Manual of the Erysiphales (powdery Mildews)*.

CBS-KNAW Fungal Biodiversity Centre.

Burdon, J. J., & Laine, A.-L. (2019). *Evolutionary Dynamics of Plant-Pathogen Interactions*.

Cambridge University Press; Cambridge Core. https://doi.org/10.1017/9781108625517

Bushnell, W. R. (2002). The role of powdery mildew research in understanding host-parasite

interaction: Past, present, and future. *The Powdery Mildews: A Comprehensive Treatise*,

1–12.

Chaloner, T. M., Gurr, S. J., & Bebber, D. P. (2020). Geometry and evolution of the ecological

niche in plant-associated microbes. *Nature Communications*, *11*(1), 2955.

https://doi.org/10.1038/s41467-020-16778-5

Christensen, R. H. B. (2019). *ordinal: Regression Models for Ordinal Data* (2019.12-10)

[Computer software]. https://CRAN.R-project.org/package=ordinal

Cuevas-Reyes, P., Gilberti, L., González-Rodríguez, A., & Fernandes, G. W. (2013). Patterns of

herbivory and fluctuating asymmetry in Solanum lycocarpum St. Hill (Solanaceae) along

an urban gradient in Brazil. *Ecological Indicators*, *24*, 557–561.

https://doi.org/10.1016/j.ecolind.2012.08.011

De Frenne, P., Graae, B. J., Rodríguez-Sánchez, F., Kolb, A., Chabrerie, O., Decocq, G., De

Kort, H., De Schrijver, A., Diekmann, M., Eriksson, O., Gruwez, R., Hermy, M., Lenoir,

J., Plue, J., Coomes, D. A., & Verheyen, K. (2013). Latitudinal gradients as natural

laboratories to infer species' responses to temperature. *Journal of Ecology*, *101*(3), 784–

795. https://doi.org/10.1111/1365-2745.12074

Desaint, H., Aoun, N., Deslandes, L., Vailleau, F., Roux, F., & Berthomé, R. (2021). Fight hard

or die trying: When plants face pathogens under heat stress. *New Phytologist*, *229*(2),

712–734. https://doi.org/10.1111/nph.16965

Dubois, J., & Cheptou, P.-O. (2017). Effects of fragmentation on plant adaptation to urban

environments. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

372(1712), 20160038. https://doi.org/10.1098/rstb.2016.0038

Egerer, M., Liere, H., Lucatero, A., & Philpott, S. M. (2020). Plant damage in urban

agroecosystems varies with local and landscape factors. *Ecosphere*, *11*(3), e03074.

https://doi.org/10.1002/ecs2.3074

ESRI, Environmental Systems Research Institute (2020). ArcGIS Streetmap Premium "Streets"

Dataset, v. 2020.3. Retrieved: July 28, 2021.

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces

for global land areas. *International Journal of Climatology*, *37*(12), 4302–4315.

https://doi.org/10.1002/joc.5086

Hawthorn, W. R. (1974). THE BIOLOGY OF CANADIAN WEEDS. 4. *PLANTAGO MAJOR*

AND *P. RUGELII*. *Canadian Journal of Plant Science*, *54*(2), 383–396.

https://doi.org/10.4141/cjps74-059

Heidt, V., & Neef, M. (2008). Benefits of Urban Green Space for Improving Urban Climate. In

M. M. Carreiro, Y.-C. Song, & J. Wu (Eds.), *Ecology, Planning, and Management of*

Urban Forests: International Perspectives (pp. 84–96). Springer.

https://doi.org/10.1007/978-0-387-71425-7_6

Höckerstedt, L. M., Siren, J. P., & Laine, A.-L. (2018). Effect of spatial connectivity on host

resistance in a highly fragmented natural pathosystem. *Journal of Evolutionary Biology*,

31(6), 844–852. https://doi.org/10.1111/jeb.13268

Inguagiato, J. C., Murphy, J. A., & Clarke, B. B. (2009). Anthracnose Disease and Annual

Bluegrass Putting Green Performance Affected by Mowing Practices and Lightweight

Rolling. *Crop Science*, *49*(4), 1454–1462. https://doi.org/10.2135/cropsci2008.07.0435

Johnson, M. T. J., & Munshi-South, J. (2017). Evolution of life in urban environments. *Science*,

358(6363). https://doi.org/10.1126/science.aam8327

Johnson, M. T. J., Prashad, C. M., Lavoignat, M., & Saini, H. S. (2018). Contrasting the effects

of natural selection, genetic drift and gene flow on urban evolution in white clover (

Trifolium repens). *Proceedings of the Royal Society B: Biological Sciences*, *285*(1883),

20181019. https://doi.org/10.1098/rspb.2018.1019

Jousimo, J., Tack, A. J. M., Ovaskainen, O., Mononen, T., Susi, H., Tollenaere, C., & Laine, A.-

L. (2014). Ecological and evolutionary effects of fragmentation on infectious disease

dynamics. *Science*, *344*(6189), 1289–1293. https://doi.org/10.1126/science.1253621

Jules, E. S., Kauffman, M. J., Ritts, W. D., & Carroll, A. L. (2002). Spread of an Invasive

Pathogen Over a Variable Landscape: A Nonnative Root Rot on Port Orford Cedar.

Ecology, *83*(11), 3167–3181. https://doi.org/10.1890/0012-

9658(2002)083[3167:SOAIPO]2.0.CO;2

Just, M. G., Dale, A. G., Long, L. C., & Frank, S. D. (2019). Urbanization drives unique

latitudinal patterns of insect herbivory and tree condition. *Oikos*, *128*(7), 984–993.

https://doi.org/10.1111/oik.05874

Kondo, M. C., Fluehr, J. M., McKeon, T., & Branas, C. C. (2018). Urban Green Space and Its

Impact on Human Health. *International Journal of Environmental Research and Public*

Health, *15*(3), 445. https://doi.org/10.3390/ijerph15030445

Laine, A.-L. (2004). Resistance variation within and among host populations in a plant-pathogen

metapopulation: Implications for regional pathogen dynamics. *Journal of Ecology*, *92*(6),

990–1000. https://doi.org/10.1111/j.0022-0477.2004.00925.x

Lambrecht, S. C., Mahieu, S., & Cheptou, P.-O. (2016). Natural selection on plant physiological

traits in an urban environment. *Acta Oecologica*, *77*, 67–74.

https://doi.org/10.1016/j.actao.2016.09.002

Lenth, R. V. (2022). *emmeans: Estimated Marginal Means, aka Least-Squares Means*.

https://CRAN.R-project.org/package=emmean

Li, D., Stucky, B., Deck, J., Baiser, B., & Robert, G. (2019). The effect of urbanization on plant

phenology depends on regional temperature. *Nature Ecology & Evolution*, *3*, 1–7.

https://doi.org/10.1038/s41559-019-1004-1

Lippe, M. von der, Bullock, J. M., Kowarik, I., Knopp, T., & Wichmann, M. (2013). Human-

Mediated Dispersal of Seeds by the Airflow of Vehicles. *PLOS ONE*, *8*(1), e52733.

https://doi.org/10.1371/journal.pone.0052733

Meineke, E. K., Classen, A. T., Sanders, N. J., & Davies, T. J. (2019). Herbarium specimens

reveal increasing herbivory over the past century. *Journal of Ecology*, *107*(1), 105–117.

https://doi.org/10.1111/1365-2745.13057

Miles, L. S., Breitbart, S. T., Wagner, H. H., & Johnson, M. T. J. (2019). Urbanization Shapes

the Ecology and Evolution of Plant-Arthropod Herbivore Interactions. *Frontiers in*

Ecology and Evolution, *7*. https://doi.org/10.3389/fevo.2019.00310

Moreira, X., Abdala-Roberts, L., Teran, J. C. B. M. y, Covelo, F., Mata, R. de la, Francisco, M.,

Hardwick, B., Pires, R. M., Roslin, T., Schigel, D. S., Hoopen, J. P. J. G. ten,

Timmermans, B. G. H., Dijk, L. J. A. van, Castagneyrol, B., & Tack, A. J. M. (2019).

Impacts of urbanization on insect herbivory and plant defences in oak trees. *Oikos*,

128(1), 113–123. https://doi.org/10.1111/oik.05497

MRLC, Multi-Resolution Land Characteristics Consortium (2019). National Land Cover

Dataset. Retrieved: July 28, 2021 from https://www.mrlc.gov/

Murray, M. H., Sánchez, C. A., Becker, D. J., Byers, K. A., Worsley-Tonks, K. E., & Craft, M.

E. (2019). City sicker? A meta-analysis of wildlife health and urbanization. *Frontiers in*

Ecology and the Environment, *17*(10), 575–583. https://doi.org/10.1002/fee.2126

Neil, K. L., Landrum, L., & Wu, J. (2010). Effects of urbanization on flowering phenology in the

metropolitan phoenix region of USA: Findings from herbarium records. *Journal of Arid*

Environments, *74*(4), 440–444. https://doi.org/10.1016/j.jaridenv.2009.10.010

Numminen, E., & Laine, A.-L. (2020). The spread of a wild plant pathogen is driven by the road

network. *PLOS Computational Biology*, *16*(3), e1007703.

https://doi.org/10.1371/journal.pcbi.1007703

Ovaskainen, O., & Laine, A.-L. (2006). Inferring Evolutionary Signals from Ecological Data in a

Plant–Pathogen Metapopulation. *Ecology*, *87*(4), 880–891. https://doi.org/10.1890/0012-

9658(2006)87[880:IESFED]2.0.CO;2

Penczykowski, R. M., Parratt, S. R., Barres, B., Sallinen, S. K., & Laine, A.-L. (2018).

Manipulating host resistance structure reveals impact of pathogen dispersal and

environmental heterogeneity on epidemics. *Ecology*, *99*(12), 2853–2863.

https://doi.org/10.1002/ecy.2526

Penczykowski, R. M., & Sieg, R. D. (2021). Plantago spp. As Models for Studying the Ecology

and Evolution of Species Interactions across Environmental Gradients*. *The American*

Naturalist. https://doi.org/10.1086/714589

Penczykowski, R. M., Walker, E., Soubeyrand, S., & Laine, A.-L. (2015). Linking winter

684   conditions to regional disease dynamics in a wild plant-pathogen metapopulation. *New*

685   *Phytologist*, *205*(3), 1142–1152. https://doi.org/10.1111/nph.13145

686   Pinheiro, J. C., Bates, D. J., DebRoy, S., & Sakar, D. (2012). The Nlme Package: Linear and

687   Nonlinear Mixed Effects Models, R Version 3. In *R package version* (Vol. 6).

688   Rivkin, L. R., Santangelo, J. S., Alberti, M., Aronson, M. F. J., Keyzer, C. W. de, Diamond, S.

689   E., Fortin, M.-J., Frazee, L. J., Gorton, A. J., Hendry, A. P., Liu, Y., Losos, J. B.,

690   MacIvor, J. S., Martin, R. A., McDonnell, M. J., Miles, L. S., Munshi-South, J., Ness, R.

691   W., Newman, A. E. M., … Johnson, M. T. J. (2019). A roadmap for urban evolutionary

692   ecology. *Evolutionary Applications*, *12*(3), 384–398. https://doi.org/10.1111/eva.12734

693   Roberts, J. A., & Murphy, J. A. (2014). Anthracnose Disease on Annual Bluegrass as Affected

694   by Foot Traffic and Sand Topdressing. *Plant Disease*, *98*(10), 1321–1325.

695   https://doi.org/10.1094/PDIS-08-13-0877-RE

696   Ross, M. D. (1973). Inheritance of self-incompatibility in Plantago lanceolata. *Heredity*, *30*(2),

697   169–176. https://doi.org/10.1038/hdy.1973.19

698   Sagar, G. R., & Harper, J. L. (1964). Plantago Major L., P. Media L. and P. Lanceolata L.

699   *Journal of Ecology*, *52*(1), 189–221. https://doi.org/10.2307/2257792

700   Santangelo, J. S., Ness, R. W., Cohan, B., Fitzpatrick, C. R., Innes, S. G., Koch, S., Miles, L. S.,

701   Munim, S., Peres-Neto, P. R., Prashad, C., Tong, A. T., Aguirre, W. E., Akinwole, P. O.,

702   Alberti, M., Álvarez, J., Anderson, J. T., Anderson, J. J., Ando, Y., Andrew, N. R., …

703   Johnson, M. T. J. (2022). Global urban environmental change drives adaptation in white

704   clover. *Science*, *375*(6586), 1275–1281. https://doi.org/10.1126/science.abk0989

Tack, A. J. M., Hakala, J., Petäjä, T., Kulmala, M., & Laine, A.-L. (2014). Genotype and spatial

structure shape pathogen dispersal and disease dynamics at small spatial scales. *Ecology*,

95(3), 703–714. https://doi.org/10.1890/13-0518.1

Tack, A. J. M., & Laine, A.-L. (2014). Ecological and evolutionary implications of spatial

heterogeneity during the off-season for a wild plant pathogen. *New Phytologist*, *202*(1),

297–308. https://doi.org/10.1111/nph.12646

US EPA, United States Environmental Protection Agency. (2021a). EnviroAtlas. Percent

impervious area [census block group]. Retrieved: June 19, 2021 from

epa.gov/enviroatlas

US EPA, United States Environmental Protection Agency. (2021b). EnviroAtlas. Percent tree

cover. Retrieved: June 19, 2021 from epa.gov/enviroatlas

US EPA, United States Environmental Protection Agency. (2021c). EnviroAtlas. Estimated

intersection density of walkable roads. Retrieved: June 19, 2021 from

epa.gov/enviroatlas

United Nations, Department of Economic and Social Affairs, Population Division (2018). World

Urbanization Prospects 2018 – Highligts (ST/ESA/ SER.A/392).

van Dijk, L. J. A., Moreira, X., Barr, A. E., Abdala-Roberts, L., Castagneyrol, B., Faticov, M.,

Hardwick, B., ten Hoopen, J. P. J. G., de la Mata, R., Pires, R. M., Roslin, T., Schigel, D.

S., Timmermans, B. G. H., & Tack, A. J. M. (2022). Urbanization affects oak–pathogen

interactions across spatial scales. *Ecography*, *2022*(1).

https://doi.org/10.1111/ecog.06091

Wang, Y., Bao, Z., Zhu, Y., & Hua, J. (2009). Analysis of Temperature Modulation of Plant

Defense Against Biotrophic Microbes. *Molecular Plant-Microbe Interactions®*, *22*(5),

498–506. https://doi.org/10.1094/MPMI-22-5-0498

Williams, D. W., Burrus, P. B., & Vincelli, P. (2001). Severity of Gray Leaf Spot in Perennial

Ryegrass as Influenced by Mowing Height and Nitrogen Level. *Crop Science*, *41*(4),

1207–1211. https://doi.org/10.2135/cropsci2001.4141207x

Yatskievych, G. A., & Steyermark, J. A. (1999). *Steyermark's flora of Missouri* (Vol. 3).

Missouri Department of Conservation.

Zhou, B., Rybski, D., & Kropp, J. P. (2017). The role of city size and urban form in the surface

urban heat island. *Scientific Reports*, *7*(1), 4791. https://doi.org/10.1038/s41598-017-

04242-2

Figure Legends

Fig. 1. (a) Google Earth satellite view of the focal urbanization gradient into St. Louis, Missouri, USA. Urban sites are plotted as purple diamonds, suburban as green squares, and rural as yellow circles. (b) Biplot of a principal component analysis of environmental variables at the study sites. Black arrows represent loadings of percent impervious area (%Imp), percent tree cover (%Tree), estimated intersection density of walkable roads (IntDen), average temperature from June–October (Temp), and nighttime radiance (NightLight). Sites are numbered from west to east (see Table S1 for additional site information).

Fig. 2. Relative frequency of each stage of plant reproductive development for each site type and survey month in 2019 and 2020 (Rur = rural, Sub = suburban, Urb = urban).

Fig. 3. Proportion of plants infected with powdery mildew in each population. Box-and-whisker plots are grouped by site type and survey month in 2019 and 2020.

Fig. 4. Proportion of plants with chewing damage or leaf mines in each population. Box-and-whisker plots are grouped by site type and survey month in 2020.

Fig. 5. Proportion of plants with leaf damage from lawn mower blades in each population. Box-and-whisker plots are grouped by site type and survey month in 2020.

Fig. 6. Proportion of sentinels with (a, b) powdery mildew infection, (c, d) leaf mines, and (e, f) chewing damage in relation to the proportion of wild conspecifics with the same type of damage in each site in July 2020. There are 15 sites plotted in each panel, but 12 points with no mildew on either wild or sentinel plants (i.e., overlapping at the origin) in panel (a).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6