Introgression between highly divergent sea squirt genomes: an adaptive breakthrough?

Christelle Fraïsse[1,2,*], Alan Le Moan[3,4], Camille Roux[1], Guillaume Dubois[3], Claire Daguin-Thiébaut[3], Pierre-Alexandre Gagnaire[2], Frédérique Viard[2,3,°], Nicolas Bierne[2,°]

[1] CNRS, Univ. Lille, UMR 8198 – Evo-Eco-Paleo, F-59000 Lille, France.

[2] ISEM, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

[3] CNRS UMR 7144 - Sorbonne University, 29680 Roscoff, France.

[4] Department of Marine Sciences, Tjärnö Marine Laboratory, University of Gothenburg, 452 96 Strömstad, Sweden.

[*]Corresponding author: E-mail: christelle.fraisse@univ-lille.fr

[°]Co-last authors

Keywords: ascidians, trio-based genome phasing, anthropogenic hybridization, introgression hotspot, cytochrome P450.

Short title: Adaptive breakthrough in sea squirts

Abstract

Human-mediated introductions are reshuffling species distribution on a global scale. Consequently, an increasing number of allopatric taxa are now brought into contact, promoting introgressive hybridization between incompletely isolated species and new adaptive gene transfer. The broadcast spawning marine species, *Ciona robusta*, has been recently introduced in the native range of its sister taxa, *Ciona intestinalis*, in the English Channel and North-East Atlantic. These sea squirts are highly divergent, yet hybridization has been reported by crossing experiments and genetic studies in the wild. Here, we examined the consequences of secondary contact between *C. intestinalis* and *C. robusta* in the English Channel. We produced genomes phased by transmission to infer the history of divergence and gene flow, and analyzed introgressed genomic tracts. Demographic inference revealed a history of secondary contact with a low overall rate of introgression. Introgressed tracts were short, segregating at low frequency, and scattered throughout the genome, suggesting traces of past contacts during the last 30 ky. However, we also uncovered a hotspot of introgression on chromosome 5, characterized by several hundred Kb-long *C. robusta* haplotypes segregating in *C. intestinalis*, that introgressed during contemporary times the last 75 years. Although locally more frequent than the baseline level of introgression, *C. robusta* alleles are not fixed, even in the core region of the introgression hotspot. Still, LD patterns and haplotype-based tests suggest this genomic region is under recent positive selection. We further detected in the hotspot an over-representation of candidate SNPs lying on a cytochrome P450 gene with a high copy number of tandem repeats in the introgressed alleles. Cytochromes P450 are a superfamily of enzymes involved in detoxifying exogenous compounds, constituting a promising avenue for functional studies. These findings support that introgression of an adaptive allele is possible between very divergent genomes and that anthropogenic hybridization can provide the raw material for adaptation of native lineages in the Anthropocene.

Author summary

Introgression, the transfer of genetic material by hybridization between taxa, is increasingly recognized to sometimes persist for long during species divergence. However, the evolutionary consequences of human-induced introgression remain largely unknown, especially in the marine realm. While some argue it poses a threat to the genome integrity of native species, others consider it has a great potential to fuel adaptation. In this work, we quantify the magnitude and genomic distribution of introgression after secondary contact between a native sea squirt and its divergently related sister species recently introduced in the English Channel. The genome-wide pattern suggests introgression is mostly impeded between these two incompatible genomes that only exchanged a few very short chromosome tracts sparsely distributed in the native genome. We nonetheless found a hotspot of long tracts that recently introgressed in a single region of the genome, with a clear footprint of recent positive selection. In the center of the hotspot, we further detected a promising candidate gene for adaptive introgression: a cytochrome P450 detoxifying enzyme with a high copy number in the introgressed allele. Therefore, our results support that adaptive introgression can remain possible between very divergent genomes and that anthropogenic hybridization can provide the raw material for the adaptation of native lineages in the Anthropocene.

Introduction

Human-mediated introductions often result in interlineage introgression (North et al. 2021; Ottenburghs 2021). Pervasive introgression implies that most co-occurring introduced and native genomes of sister species are still to some extent permeable to interspecific gene flow, with various outcomes from genome-wide genetic swamping to adaptive introgression at few specific genomic regions (McFarlane and Pemberton 2019). In the marine realm, harbors, docks and piers are prime locations for such hybridization events between non-native and native lineages, sometimes resulting in singular outcomes. For example, (Simon et al. 2020) identified a unique ecotype of marine mussels in these artificial habitats ("docks mussels"), resulting from a recent admixture between two closely-related European mussel species. These anthropogenic hybridizations can also promote secondary contact between divergent genomes with long histories of allopatric divergence (Viard et al. 2020). They provide unique opportunities to investigate the outcomes of hybridization between co-occurring genetic lineages at a late stage of the speciation continuum.

Sea squirts are among the most critical invasive marine organisms forming a significant component of the non-indigenous community in artificial marine habitats (Shenkar and Swalla 2011; Zhan et al. 2015). For this reason, they were among the first marine taxa to be studied to test the hypothesis of the relationship between climate change and biological invasions (Stachowicz et al. 2002). *Ciona robusta* is a sea squirt species native to the Northwest Pacific introduced in the early 2000s to the English Channel in the native range of *Ciona intestinalis* (Bouchemousse, Bishop, et al. 2016). The two species are found in sympatry in these regions (Nydam and Harrison 2011). However, their relative abundance varies locally over seasons (Bouchemousse, Lévêque, et al. 2016), and they display contrasting genetic diversity patterns, with low mitochondrial diversity in *C. robusta* supporting its recent introduction (Bouchemousse, Bishop, et al. 2016). *C. robusta* and *C. intestinalis* represent a pair of species at the end of the speciation continuum with 14% of net synonymous

divergence, which is well above the ~2% suggested to delineate the end of the grey zone of speciation

in most animals (Roux et al. 2016). Despite this high molecular divergence, first and second-

generation crosses between the two species show successful hybridization in the laboratory

(Bouchemousse, Lévêque, et al. 2016; Malfant et al. 2018). Moreover, the two species produce

gametes synchronously in the wild, with juveniles recruiting simultaneously (Bouchemousse,

Lévêque, et al. 2016). However, the use of >300 ancestry-informative markers on 450 individuals

showed limited evidence for recent hybridization in the wild, with only one F1 and no later generation

hybrids found in the sympatric range (Bouchemousse, Liautard-Haag, et al. 2016). Therefore, efficient

reproductive barriers seem to restrict hybridization in nature.

Despite the paucity of first-generation hybrids, Le Moan et al. (2021) found compelling

evidence of contemporary introgression between *C. robusta* and *C. intestinalis* in the sympatric range

(Bay of Biscay, Iroise Sea and the English Channel) from RADseq-derived SNPs. Instead of genome-

wide admixture, Le Moan et al. (2021) detected a single genomic hotspot (~1.5 Mb) of long

introgressed *C. robusta* tracts into its native congener on chromosome 5. The absence of such

introgression tracts in allopatric populations suggests introgression occurred after the recent

introduction of *C. robusta*. At a fine spatial scale within the sympatric range, the introgressed tracts

displayed chaotic frequencies across sympatric localities, which has been attributed to human--

mediated transport among harbors (Hudson et al. 2016). The features of the introgression hotspot

identified between the two species, namely being 1) unidirectional, 2) localized in a single genome

region, and 3) made-up of long tracts, are reminiscent of the footprint of positive selection. Therefore,

Le Moan et al.'s work (2021) provides a seminal example of a contemporary introgression

breakthrough between two species at a late stage of the speciation continuum. It underlines the need to

densely scan genomes with genome-wide markers, notably when considering divergent genomes that

may only show very localized introgression hotspots (Ravinet et al. 2018; Maxwell et al. 2019;

Stankowski et al. 2020; Yamasaki et al. 2020).

Here, we extend Le Moan et al.'s study (2021) using whole-genome sequences fully phased by

transmission in both *C. robusta* and *C. intestinalis* taken from their sympatric range (English Channel),

to *i)* specifically delineate the core region of the genomic breakthrough, *ii)* test for the footprint of

selection, and *iii)* identify candidate loci driving the putatively adaptive introgression. We also

examined a non-introgressed *Ciona roulei* population in the Mediterranean Sea, used as a control.

Based on experimental crosses and genome-wide analyses, recent studies showed that *C. roulei* is a

"Mediterranean lineage" of the accepted species *C. intestinalis*, and thus its species status needs to be

revised (Malfant et al. 2018; Le Moan et al. 2021). However, we will continue to name it "*C. roulei*" in

this study. Based on whole genomes, we recovered the observation of Le Moan et al. (2021) for a

genomically localized introgression breakthrough on chromosome 5 from *C. robusta* to the sympatric

C. intestinalis, absent in the *C. roulei* population of the Mediterranean Sea. We also inferred the

divergence history of the two species and confirmed that they have hybridized in the past, far before

their introduction in Europe (Roux et al. 2013). However, when including chromosome 5, we

recovered a signal of contemporary introgression. Next, we inferred the haplotype ancestry of the *C.*

intestinalis genomes and delineated migrant genomic tracts. In sharp contrast with a genomic

background interspersed with small and sparse introgressed tracts attributed to past admixture, we

found a distinct pattern of very long introgressed tracts segregating at intermediate frequencies at the

introgression hotspot on chromosome 5. Finally, using haplotype-based tests, we provided evidence

that the high linkage disequilibrium (LD) observed in the genomic hotspot is due to some sort of

positive selection. Inspecting annotated genes at the core of the introgression breakthrough, our best

candidate for selection was a cytochrome P450 gene, on which differentiated SNPs were over-

represented, and that showed a high copy number tandem repeat in the *C. robusta* introgressed

haplotype.

Results

Genome-wide analysis of population structure

A principal component analysis on genome-wide unlinked SNPs (**Figure 1B**) showed, as expected, that the *C. robusta* individuals are clearly distinguished from the *C. intestinalis* individuals sampled in the sympatric populations of the English Channel (Brest and Aber Wrac'h, named 'Aber' in the following text) and from the non-introgressed population of the Mediterranean Sea (Banyuls, *C. roulei*). This primary component of genetic variation was carried by the first PCA axis (21.2% of explained variance). In comparison, the second axis (6.7%) revealed a slight genetic differentiation between *C. intestinalis* and *C. roulei*, validating previous findings with RADseq (Le Moan et al. 2021). The intraspecific F1 individuals produced in the lab (**Table S1**) fall within the genetic variance of their species, while the interspecific F1s fall halfway between the two species along the first axis (**Figure 1B**), validating their F1 hybrid status. The intraspecific variance along the first axis was substantial within *C. intestinalis*. At the same time, this was not the case of *C. robusta* and *C. roulei*, suggesting that interspecific introgression affects specifically *C. intestinalis* individuals in the English Channel.

The SNPs contributing to species divergence on the first axis are distributed genome-wide (**Figure 1C**). Still, a decline of SNP contribution on the first axis at the start of chromosome 5 indicated a reduction of the divergence between *C. robusta* and the sympatric *C. intestinalis* populations locally in the genome. This pattern is supported by the observation of a consistently high divergence across the genome between *C. robusta* and *C. intestinalis* (the maximal F_{ST} calculated in non-overlapping 10 Kb windows is equal to one), except at the start of chromosome 5, where the maximal F_{ST} value declined from 1 to 0.69 (**Figure S1A**). This striking decline, located between 700 Kb and 1.5 Mb, was not observed between *C. robusta* and *C. roulei* (**Figure S1B**). Moreover, in *C. intestinalis,* it did not correlate with a reduction of diversity (π), suggesting it is likely due to interspecific introgression rather than intraspecific selective sweeps. This pattern is very different from

what was observed for the averaged F_{ST} (**Figure S1**) that strongly varied across the genome. It was

notably higher at the beginning or in the middle of the chromosomes in regions of low intraspecific

genetic diversities. These large-scale averaged variations were observed in all species, indicating that

they may be due to the long-term effect of linked selection acting on a shared recombination landscape

(note that all chromosomes in *C. intestinalis* are metacentric, except chromosome 2, 7 and 8, which are

submetacentric (Shoguchi et al. 2006).

Genome-wide analysis of introgression

We calculated the Patterson's *D* statistic using *C. edwardsi* as an outgroup to test for genome-wide

admixture between the two species (**Figure S4**). We found evidence for an excess shared ancestry of

C. robusta with the sympatric *C. intestinalis* relative to *C. roulei* across all chromosomes (**Figure S4**).

To locate introgressed genomic regions, the fraction of the genome that has been shared between

species (*fd*) was then calculated in non-overlapping windows. *fd* varied around zero along each

chromosome, but we observed an outlying increase on chromosome 5 between 700 Kb and 1.5 Mb

showing a high admixture level between *C. robusta* and *C. intestinalis* (**Figure S4**). This *fd* increase

had its maximum (25% of admixture level) centered on the introgression hotspot of chromosome 5 and

was only present in the sympatric range. None of the other chromosomes showed outlying genomic

regions, neither with *C. intestinalis*, nor with *C. roulei*. Furthermore, the admixture proportion was

weakly negatively correlated with chromosome length, a known proxy for the recombination rate

(Kaback 1996), but chromosome 5 was a clear outlier (it has a higher *fd* value than expected given its

size).

We detected introgression tracts in *C. intestinalis* genomes using local ancestry inference on

640,044 phased SNPs and considering *C. robusta* and *C. roulei* as the parental populations. The

inferred tracts showed similar introgression patterns to the raw haplotypes obtained from SNPs fixed

between *C. robusta* and *C. roulei* (**Figure S2**). This suggests that local ancestry inferences indeed

detect the introgressed genomic regions while being less noisy than when considering raw haplotypes.

The proportion of *C. robusta* ancestry inferred was low (0.1% on average per individual), suggesting

that the introgression rate at the genome level is low. Furthermore, there was no significant correlation

in *C. robusta* ancestry between chromosomes (except in 5 pairs) among the *C. intestinalis* individuals

(**Table S2**). Introgressed tracts were short (median size of 380 bp) and widespread across the genome

(**Figure 2**). These short tracts had a bimodal frequency distribution with a majority segregating at low

frequency and a minority fixed in *C. intestinalis*. They likely have originated from past admixture

events between the two species and then progressively been chopped down by recombination over

time while they drifted towards loss or fixation. The introgression hotspot on chromosome 5

immediately appears as an outlier on the chromosome map (**Figure 2**). Its underlying tracts were much

longer (maximal size of 156 Kb) than the tracts outside of the hotspot, and they segregated at

intermediate frequencies (none of the long tracts on the hotspot was fixed), in line with a recent

introgression event.

We then analyzed the coding sequences inside and outside the genomic tracts identified as

being introgressed (**Figure S3**). Chromosome 5 carries by far the largest number of introgressed CDS:

65 of 69 were located on this chromosome, while the four other introgressed CDS were located on

three different chromosomes (3, 8 and 13). Among all CDS on chromosome 5, 6% were detected as

being on introgressed tracts, demonstrating that the hotspot does contain introgressed genes. As

introgression has not reached fixation in *C. intestinalis,* we would expect an increase of diversity

within *C. intestinalis* (π) and a decrease of interspecies divergence (d_{XY}) for the CDS on introgressed

tracts compared to the rest of the genome. However, this is not what was observed (**Figure S3A**),

probably because the *C. robusta* introgressed alleles segregate at an intermediate frequency that

negligibly impacts diversity. Therefore, we computed the G_{min} statistic, defined as the ratio of the

minimum d_{XY} to the average d_{XY}, which is better suited to capture the effect of recent introgression

events (Geneva et al. 2015). As expected if introgressed tracts originate from recent introgression

events, we found that G_{min} was significantly lower in the introgressed CDS than in the rest of the
genome (**Figure S3B**).

<u>The history of divergence and gene flow between *C. robusta* and *C. intestinalis*</u>

In order to address whether short and long *C. robusta* tracts introgressed in the *C. intestinalis* genomes
could result from different introgression events, we reconstructed the divergence history between the
two species based on their joint site frequency spectrum. Divergence models in which the history of
gene flow can take different forms were tested. The possibilities of having a heterogeneity of effective
population sizes and effective migration rates to model the effects of linked selection and species
barriers were also included in the models. This is because previous work showed that when these
features were not considered, the inferences led to ambiguous results in sea squirts (Roux et al. 2016).
We first excluded chromosome 5 from the inferences to capture the prominent history between the two
species (**Figure 3** and **Table S3** for details). Divergence with periodic connectivity and the effects of
linked selection was the best model, closely followed by a secondary contact model. The divergence
between the two species started with gene flow (during ~400 Ky), then it was followed by a ~1.5 My
period of isolation. Only in the 30,000 last years, *C. robusta* and *C. intestinalis* came into secondary
contact. This long period of introgression could explain the presence of the short introgressed tracts in
C. intestinalis. In line with this scenario, the estimates of migration rates show that introgression is
highly asymmetrical from *C. robusta* toward *C. intestinalis*. Furthermore, we observed a ten-fold
lower effective population size in *C. robusta* than *C. intestinalis*, which matches the difference in
nucleotide diversities between the two species and can be explained by the recent introduction of *C.*
robusta in Europe (**Figure S1**). Repeating the demographic analyses with chromosome 5 in the dataset
led to very similar parameter estimates, except for the divergence times (**Figure S5** and **Table S4** for
details). Indeed, the best model was now a secondary contact, where a long period of isolation (~2
My) was followed by a contemporary period of introgression (in the last 200 years), which may
capture the signal left by the long introgressed tracts on chromosome 5.

We used a neutral recombination clock to refine the time estimate since admixture at the

introgression hotspot on chromosome 5. The average length of the introgressed tracts can be estimated

using the formula $\bar{L} = [(1 - f) * r * (t - 1)]^{-1}$, where r is the local recombination rate (crossovers

per base pair per generation), f is the admixture proportion, and t is the time since the admixture event

in generations (Racimo et al. 2015). Given that the average length of introgressed tracts at the hotspot

is 19,898 bp, the mean frequency of introgression is 0.106, and the recombination rate is 3.82e-07

M/bp (Duret, pers. comm.), we found that the contemporary admixture between *C. robusta* and *C.*

intestinalis occurred about 75 years ago (assuming two generations per year; (Bouchemousse et al.

2017). Note that due to small regions lacking sufficient ancestry signal (**Figure S7**), some introgressed

tracts could be a bit longer than measured, indicating an even more contemporary introgression.

<u>The introgression hotspot on chromosome 5</u>

We have shown that maximal F_{ST} values between *C. robusta* and *C. intestinalis* form a valley at the

start of chromosome 5 (**Figure 4A**). This pattern is due to long *C. robusta* tracts segregating in the

sympatric populations of *C. intestinalis* (**Figure 4B**). The introgression tracts were variable in size.

They shared ancestral recombination breakpoints, clearly visible in the linkage disequilibrium (LD)

heatmap between pairs of diagnostic SNPs along chromosome 5 (**Figure 4D**). The hotspot region

between 700 Kb and 1.5 Mb exhibited stronger LD (r^2 median of 0.3) than the rest of chromosome 5

(r^2 median of 0.007). Introgression was maximal on either side of the "missing data region" from

1,009,000 to 1,055,000 bp (a region of significantly increased read depth: 100x in average inside the

region *vs* 25x outside). But we found no evidence of introgressed tracts in the hotspot that have

completely swept to fixation in *C. intestinalis* (**Figure 4C**).

To explicitly test if some sort of selection could explain this pattern on chromosome 5, we

used various methods. We first sought the footprint of a classic selective sweep, where a *de novo*

beneficial mutation arises on a *C. robusta* haplotype and quickly sweeps toward fixation, reducing

diversity and creating a signal of long-range LD around it. This signal can be captured by the extended

haplotype homozygosity (EHH), which measures the decay of identity-by-descent between haplotypes

as a function of the distance from a focal SNP. Taking as targets the SNPs with the highest *C. robusta*

frequency to the left and right of the "missing data region", we observed a slower EHH decay on the

C. robusta haplotypes compared to other haplotypes in the sympatric *C. intestinalis* populations, but

not in the *C. roulei* population (**Figure 4E**). To test for significance, the absolute normalized integrated

haplotype score (iHS) was then calculated in 50-Kb windows along chromosome 5, and we estimated

the proportion of SNPs in each window associated with outlying values of iHS. This proportion was

the highest in the core region of the introgression hotspot in *C. intestinalis* (8%) and *C. robusta* (20%),

but not in *C. roulei* (0%). This result indicates a low haplotype diversity over an extended region in

both the donor *C. robusta* and the introgressed alleles of the recipient *C. intestinalis* populations. The

genealogies of the 50-kb windows framing the "missing data region" further support a reduced

diversity of the *C. robusta* clade (**Figure S8**). Moreover, the alleles sampled in the introgressed *C.*

intestinalis genomes cluster within the star-like *C. robusta* clade, suggesting that a recent selective

sweep happened in *C. robusta* and a single beneficial haplotype introgressed into *C. intestinalis*.

Finally, we used a complementary approach (VolcanoFinder) to directly test for adaptive

introgression using all SNPs from the *C. intestinalis* recipient species only. Again, the method is

suitable to detect an adaptively introgressed allele that has swept to fixation in the recipient species,

producing intermediate-frequency polymorphism in its flanking regions. Although introgression was

incomplete in our case (generating a soft sweep pattern, which may lead to a decrease in power), we

nonetheless observed a signal of adaptive introgression on the hotspot of chromosome 5 (**Figure S6**).

Several other regions in the genome showed extreme values of the log-likelihood ratio test. However,

contrary to the introgression hotspot, these regions also displayed signals of *de novo* selective sweeps

within *C. intestinalis* (detected with SweepFinder) that globally correlated with genomic regions of

reduced diversity (**Figure S1**). In contrast, a signal of *de novo* selective sweep within *C. robusta* was

detected in the introgression hotspot, supporting the view that beneficial alleles in this species recently

swept to fixation and were adaptively introgressed into the sympatric *C. intestinalis*.

<u>Copy number variation at the introgression hotspot</u>

We then annotated the introgression hotspot region of chromosome 5 (700 Kb - 1.5 Mb) to identify

putative candidate genes under selection. To overcome the difficulty posed by the high coverage of the

"missing data region" at the center of the hotspot, we relied on the variant allele fraction (VAF)

calculated from read depth to find candidate SNPs. Because the reference genome used throughout this

paper is from *C. robusta*, the variant allele represents the alternate allele in the *C. robusta* genome.

Candidate SNPs were defined as being differentiated between *C. robusta* and *C. roulei*, therefore

having a low VAF in the former and a high VAF in the latter, and being exclusively introgressed in the

sympatric *C. intestinalis* (VAF below 50%). Using a lenient threshold of VAF higher than 85% in *C.*

roulei and below 15% in *C. robusta*, we found 28 candidate SNPs in the 800-Kb region of the hotspot

distributed across six different protein-coding genes and two non-coding loci (**Figure S9**). Only

variants in the "missing data region" (20 of 28 SNPs) showed a coverage pattern in line with multi-

copy genes. Notably, 16 of these SNPs were located on the cytochrome P450 family 2 subfamily U

gene. Three other cytochromes from family 2 were found in the "missing data region" (subfamilies J/

D/R; **Figure S10**), but none contained candidate SNPs.

We did not find candidate SNPs where the *C. robusta* allele had swept to fixation in *C.*

intestinalis. The SNP showing the highest introgression frequency (0.85, i.e., only two non-

introgressed *C. intestinalis* individuals out of 13 sampled) was located in a single-copy non-coding

locus at position 1,067,404 bp. Nevertheless, this pattern should be interpreted with caution as many

individuals had a shallow read depth at this SNP. Considering the multi-copy genes, the 16 variants on

the cytochrome P450 all exhibited the same pattern of a high copy number of the introgressed *C.*

robusta allele, while this was not the case of the other multi-copy genes (**Figure S9**). Two candidate

SNPs on the cytochrome P450 are represented in **Figure 5**. They showed that *C. robusta* individuals

carried from five to twenty copies of the reference allele, while *C. roulei* individuals had one or two

copies of the alternate allele. As for the introgressed *C. intestinalis*, they were heterozygous with one

copy of the *C. intestinalis* allele and at least ten copies of the *C. robusta* allele, while the non-

introgressed *C. intestinalis* individuals were like *C. roulei*. This pattern suggests the presence of

multiple copies in tandem repeats of the *C. robusta* allele on cytochrome P450, which might play a

critical role in adaptation, and have favored its introgression into *C. intestinalis*.

Discussion

We used phased genomes from whole-genome trio sequencing to document the fine-scale genomic

consequences of the human-mediated contact between the invasive *C. robusta* and the native *C.*

intestinalis sea squirt species in Europe. A Mediterranean *C. roulei* population was also whole-genome

sequenced to be used as a non-introgressed control. Despite their high divergence, we have

demonstrated that the introduced and native species still hybridize in their sympatric range, showing a

localized introgression hotspot in the native species. We provided several lines of evidence for a

selective sweep of an adaptive *C. robusta* allele introgressed into *C. intestinalis* at the hotspot and

identified a tandem repeat variation at the cytochrome P450 locus to be a promising candidate.

Introgression between highly divergent sea squirt genomes

Introgression between highly divergent lineages has been rarely reported, partly because there is a bias

against studying the end of the speciation continuum (Kulmuni et al. 2020). Indeed, the few cases that

documented introgression between divergent species consistently showed that it was rare and localized

to small genomic regions, suggesting that most introgression events were deleterious in the recipient

genome. Moreover, introgression occurred more often in regions depleted in conserved elements and

regions with high recombination rates, consistent with the idea that introgressed tracts escape the

effect of species barriers through recombination (Martin and Jiggins 2017). Examples include

drosophila flies (Turissini and Matute 2017), coccidioides fungi (Maxwell et al. 2019), nine-spined

sticklebacks (Yamasaki et al. 2020), sea snails (Stankowski et al. 2020) or aspen trees (Shang et al.

2020). In line with these previous findings, we found that gene flow was limited between the two

divergent sea squirt species, and that most introgressed tracts segregating within *C. intestinalis* were

short.

Based on coding sequences, previous work inferring the divergence history between *C.*

robusta and *C. intestinalis* found contrasting results. With sampling locations situated in the sympatric

and allopatric ranges of the two species, Roux et al. (2013) found evidence for introgression after

secondary contact. However, when they used only allopatric populations in a later work (Roux et al.

2016), and included the effect of linked selection by introducing within genome variations in effective

population size, they found unsupported evidence for ongoing gene flow. These differences illustrate

the effect of the sampling locations on the demographic inferences and demonstrate the importance of

geography to understand the complex history of divergence in non-indigenous species. Furthermore,

the two studies by Roux et al. differ in the set of models compared. Failing to account accurately for

genomic heterogeneity in gene flow or effective population size could result in falsely supporting a

wrong model (Roux et al. 2016).

Here, we tested whether the presence of many short and a few very long *C. robusta*

introgressed tracts in the genome of the sympatric *C. intestinalis* species could be explained by a

complex history of gene flow between the two species. Therefore, we fitted models that could include

genomic heterogeneities in effective population sizes and migration rates as well as periodic

connectivity between the two species. Despite a firm species boundary with about two-thirds of the

genome linked to species barriers, we found signals of past introgression (in the last ~30 Ky), far

preceding their contemporary contact in Europe. This is in line with the low rates of natural

hybridization between the two species (Bouchemousse, Liautard-Haag, et al. 2016). The pattern of

high differentiation we observed along the genomes also suggests highly polygenic barriers that

maintain the species boundaries between *C. robusta* and *C. intestinalis*. As the species diverged for

~1.5 to 2 million years in strict isolation, they had time to accumulate many barriers in their genomes,

contributing to selection against introgression upon secondary contact.

These inferences were made excluding chromosome 5 to capture the prominent history

between the two species. When including this chromosome, and so the long introgressed tracts in the

introgression hotspot, we found evidence for a much more recent introgression event dated 200 years

ago. This estimate was then refined using a recombination clock and the introgressed tract length

distribution. We found that the contemporary introgression event may have occurred about 75 years

ago, consistent with the human-induced introduction of *C. robusta* in the English Channel (Nydam and

Harrison 2011; Bouchemousse, Bishop, et al. 2016).

<u>Is there an adaptive breakthrough on chromosome 5?</u>

In line with the studies mentioned above on divergent species, we observed limited introgression

between *C. robusta* and *C. intestinalis* in the sympatric range. On top of the many short introgressed

tracts (average length of 2.6 Kb) widespread in the *C. intestinalis* genome and segregating at a low

frequency, we observed a very localized introgression signal between 700 Kb and 1.5 Mb on

chromosome 5. This hotspot of introgression harbored very long introgression tracts (maximal length

of 156 Kb) that were more frequent than the baseline introgression level. This pattern contrasts with

the tract length distribution observed in a secondary contact between two divergent *Drosophila* fly

species that diverged 3 My ago (Turissini and Matute 2017). Introgression produced mostly small

tracts (1 to 2.5 Kb on average), but the longest tracts were only 7.5 to 10 Kb long, ten times smaller

than what was observed in the sea squirt hotspot. The situation in sea squirts resembles more to the

introgression pattern between two fungi species that diverged 5 My ago (Maxwell et al. 2019). Most

introgression tracts were 3 to 4 Kb long on average and segregated at low frequency, but there was a

long tail of longer tracts (maximal length of 100 Kb), some of them being found in high frequency

within species.

Adaptive introgressed alleles are expected to increase in frequency in the recipient population.

However, alleles might also increase in frequency simply due to allele surfing at the front wave of a

range expansion (Klopfstein et al. 2006). In our study, we only sampled populations in the English

Channel (Aber and Brest), but Le Moan et al. (2021) demonstrated that the introgression hotspot was

present in multiple localities (10 of 18) across the contact zone (Bay of Biscay, Iroise Sea and the

English Channel). The populations we sampled in Aber and Brest were among the most introgressed,

together with populations in the western UK coastline. However, there was no evidence for a wave of

introgression in line with geography: the distribution of introgressed tracts was a geographic mosaic,

likely due to human-mediated transportation (Le Moan et al. 2021).

Furthermore, the introgression of genomic tracts across a species barrier is highly random at

short time scales. Therefore, one expects a large variance in the tract length distribution under neutral

admixture (Sachdeva and Barton 2018). Observing long haplotypes at intermediate frequency could

thus be explained with purely neutral processes. Still, the singularity of such a region found in the

genome of sympatric *C. intestinalis* individuals seems difficult to explain without invoking some sort

of selection. We identified signals of selection based on haplotype variations in the flanking regions of

the most introgressed alleles (Sabeti et al. 2002; Staubach et al. 2012). Indeed, the introgression

hotspot is characterized by unusually long-range LD in the introgressed *C. intestinalis* population. The

genealogy at the hotspot shows that the haplotypes sampled in C. intestinalis cluster together with the

start-like clade of the *C. robusta* haplotypes. This indicates that a recent selective sweep occurred in

the *C. robusta* population, leading to the fixation of a beneficial allele, which then introgressed into the

sympatric *C. intestinalis* populations. This scenario was supported using an independent method based

on polarized SNPs (Setter et al. 2020; Szpiech et al. 2021).

Nevertheless, we cannot claim yet that the hotspot on chromosome 5 contains alleles that were

adaptively introgressed *sensu stricto*. Indeed, the introgression is not fixed in the studied *C.*

intestinalis population (maximal frequency of 0.31), nor in other distant localities of the contact zone

included in (Le Moan et al. 2021). Le Moan et al. suggested that the maintenance of polymorphism at

these alleles could be explained with some sort of balancing selection: if the introgressed tracts are

under overdominance or frequency-dependent selection, and suffer a fitness reduction when frequent

and homozygous in a foreign genetic background. Therefore, an incomplete sweep aligns with

balancing selection acting on the introgressed alleles (e.g., humans and neanderthals: Sams et al.

2016). In addition, this pattern is also expected if admixture is very recent, typically when it has been

human-mediated, as then allele replacement may still be ongoing in the recipient population. For

example, this may be the case in honeybees where a haplotype of European ancestry, implicated in

reproductive traits and foraging, was found at high frequency (but not fixed) in Africanized honeybees

(Nelson et al. 2017, confirmed in Calfee et al. 2020). Incomplete introgression at a single region has

also been documented in cotton bollworm, where an insecticide resistance allele at a cytochrome P450

gene increased in frequency after introducing an invasive congener carrying the adaptation (Valencia-

Montoya et al. 2020).

<u>A usual suspect: cytochrome P450</u>

In the middle of the introgression hotspot, we identified a region with high coverage that we could not

analyze using called genotypes (from 1,009,000 to 1,055,000 bp). Therefore, we examined the read

depth at candidate SNPs in this genomic region to identify further variants introgressing at a high

frequency. This analysis pinpointed 28 candidate SNPs, of which one in a non-coding region was at a

high frequency (0.85), but its overall low read depth calls for caution. The second most introgressed

SNPs (*n*=16) were located on the cytochrome P450 family 2 subfamily U gene, and all showed the

same introgression pattern with a frequency of 0.35. Strikingly, the *C. robusta* alleles had a read depth

pattern consistent with them being multi-copy (5 to 20 copies), while this was not the case for the *C.*

intestinalis alleles sampled in the non-introgressed individuals.

These candidate variants could potentially be involved in adaptation. Notably, the cytochrome

P450 gene is an exciting candidate. It belongs to a large gene class of oxidase enzymes responsible for

the biotransformation of small endogenous molecules, detoxifying exogenous compounds, and it is involved in regulating the circadian rhythm. Cytochrome P450 family 2 is the largest and most diverse CYP family in vertebrates, and the U and R subfamilies were present in the vertebrate ancestor (Nelson 1998). A recent study experimentally showed that the candidate gene we identified here (cytochrome P450 2U) is involved in the inflammatory response in *C. robusta* (Vizzini et al. 2021). Although this phenotype indicates resistance toward toxic substances, future functional study of potential fitness differences between the tandem repeat *C. robusta* allele and the single copy *C. intestinalis* allele will be needed to determine what adaptive role these alleles play. Note, however, that if the tandem-repeat variant provides adaptation to pollution in harbors, this would result in local selection and explain the absence of fixation (the native alleles being fitter in wild habitats), as discussed above.

At a larger phylogenetic scale, resistance genes were identified as gene families enriched in adaptive introgressions (Moran et al. 2021). Notably, human-induced selection such as insecticide exposure drives strong and rapid development of resistance. In that context, gene amplification of detoxification enzymes is a crucial feature for adaptation as it increases the number of functional enzymes and/or allows neofunctionalization of the new copies. There are many examples of such processes involving cytochromes P450 in insects. Insecticide resistance is due to gene amplification that produces over-expression of the cytochrome P450 gene in the aphid *Myzus persicae* (neonicotinoids resistance, Puinean et al. 2010), *Drosophila melanogaster* (DDT resistance, Schmidt et al. 2010), *Anopheles funestus* (pyrethroid resistance, Wondji et al. 2009), and *Anopheles coluzzii* (ITN resistance, Main et al. 2018). Neonicotinoids resistance due to neofunctionalization of a duplicated cytochrome P450 was demonstrated in the brown planthopper, *Nilaparvata lugens* (Zimmer et al. 2018). Another example of high copy numbers of cytochrome P450 conferring insecticide resistance was found in the moth *Spodoptera frugiperda* (Yainna et al. 2021). In contrast, resistance against pyrethroid in the moth *Helicoverpa armigera* and introgressed *Helicoverpa zea* was due to a

chimeric cytochrome P450 gene resulting from recombination between two copies in tandem

(Valencia-Montoya et al. 2020).

Even though we are not yet at the step of functionally characterizing the cytochrome P450

candidate gene, we highlighted in this work the critical role of biological invasions for driving

adaptive introgression across species boundaries. Our work also illustrates that phased genomes offer

the opportunity to detect introgression signals between divergent species, even when they are rare and

localized in the genome. Genomically localized introgression breakthroughs are still an understudied

pattern that recent genomic surveys have only begun to unravel.

Materials and Methods

Sampling and whole-genome sequencing

Sixteen parent-offspring trios (six interspecific, six within *Ciona intestinalis* and four within *Ciona*

robusta) were generated by crossing wild-caught parents in the laboratory at Roscoff (**Table S1**). For

C. intestinalis, seven of the parents used were sampled in the marina of the Aber Wrac'h (Finistère,

France), and nine others in the marina of Moulin Blanc, Brest (Finistère, France). For *C. robusta*, the

ten parents used were also sampled in Moulin Blanc. The two parents and one randomly selected

descendant for each trio were fixed in absolute ethanol, and their whole genomic DNA was extracted

using a CTAB protocol. Five individuals were sampled in Banyuls-Sur-Mer (Méditerranée, France)

belonging to *Ciona roulei*. Based on crossing experiments and genetic analyses, the species status of

C. roulei has been repeatedly questioned (Nydam and Harrison 2010; Malfant et al. 2018; Le Moan et

al. 2021). In particular, recent genetic analyses clearly showed that *C. roulei* is a distinct lineage of *C.*

intestinalis, specific to the Mediterranean Sea (Le Moan et al. 2021). Therefore, we used these

individuals as a positive control for a non-introgressed population of *C. intestinalis*. For *C. roulei*

samples, genomic DNA was extracted using a Nucleospin Tissue kit (Macherey-Nagel). After quality

control, DNA extracts were sent to the LIGAN genomics platform (Lille, France) where whole-

genome sequencing libraries were prepared separately for each of the 48 individuals, and were sequenced on an Illumina Hi-Seq 2000 instrument using 100 bp PE reads. Three parents that were poorly sequenced (ad2, ad18 and ad31; **Table S1**) were excluded from analyses.

Furthermore, four *Ciona edwardsi* individuals were sampled in Banyuls-Sur-Mer. *C. edwardsi* is reproductively isolated from the other taxa included in this study, and it was used as an outgroup (Malfant et al. 2018). These individuals were fixed in RNAlater, and their DNA was extracted using a Nucleospin Tissue kit (Macherey-Nagel). Libraries were prepared separately for each of the four individuals, and were sequenced on an Illumina Hi-Seq 4000 instrument using 150 bp PE reads at FASTERIS (Plan-les-Ouates, Switzerland).

Genotyping and haplotyping pipeline

We followed the GATK best practice pipeline (Van der Auwera et al. 2013) including haplotype phasing-by-transmission, as applied in Duranton et al. (2018). All scripts used in the pipeline are available in the **Supplementary Scripts**. We generated seven different datasets with various levels of filtering, and with or without haplome phasing, that are described in the **Supplementary Data**.

All analyses were made using the newly available *C. robusta* assembly as the reference genome (GCA_009617815.1; Satou et al. 2019). As a cautionary note, analyses in Le Moan et al. (2021) were made using the previous *C. robusta* reference genome published in 2011 (GCA_000224145.1), therefore coordinates do not correspond between the two studies. Reads were aligned to the *C. robusta* reference genome using BWA-mem v0.7.5a (Li and Durbin 2009), and duplicates were marked using Picard v1.119. The individual bam files of the introgression hotspot were used as dataset **#7**. The mean read depth was 21x across all samples (**Table S1**).

A series of steps were then performed using GATK v3.4-0 (McKenna et al. 2010), including: *i*) local realignment around indels, *ii*) individual variant calling in gVCF format using the HaplotypeCaller (options: dontUseSoftClippedBases, heterozygosity=0.01, minimum base quality score=30), *iii*) joint genotyping using GenotypeGVCFs (heterozygosity=0.01), *iv*) genotype

refinement based on family priors. Hard-filtering was then applied on the SNPs and indels to produce

a database of high-confidence variants. The database was then used to recalibrate variant quality

scores with the VQSR algorithm. After recalibration, a second round of genotype refinement based on

family priors was applied.

We then introduced a step of genotype verification (and correction where required) based on

the individual variant allele fraction (VAF) at each site, i.e., the ratio of the alternate allele depth to the

total (alternate+reference) depth. A distribution across sites was plotted for the three possible

genotypes (homozygous reference: 0/0, heterozygous: 0/1, homozygous alternate: 1/1). While the

distributions for the homozygous genotypes were shaped as expected (i.e. 99% of the sites had a VAF

< 0.1 for 0/0 and VAF > 0.9 for 1/1), the distribution of heterozygous genotypes was normally

distributed around VAF=0.5, but showed additional peaks near 0 (and near 1 to a lesser extent).

Therefore, we corrected the miscalled 0/1 genotypes to 0/0 when the variant allele depth was below

the 99th quantile of the 0/0 distribution and to 1/1 when it was above the 1th quantile of the 1/1

distribution. In addition, heterozygous genotypes with a VAF < ⅓ or > ⅔ were assigned as missing

data (excluding the ones we corrected near VAF=0 or 1). We also considered as missing data the

genotypes with a total depth below ten reads or above the 99th quantile of the depth distribution (to

exclude repeated regions). Finally, low-quality variants were excluded from the VCF (QUAL<30), and

we applied a stringent filter on individual genotype quality (GQ<30). Different missing data thresholds

were then applied to produce datasets **#2** (five missing genotypes), **#5** (no missing genotypes allowed)

and **#6** (three missing genotypes).

Phased genomes were obtained using the tool PhaseByTransmission of GATK v3.4-0. All trios

were phased given parents and offspring genotype likelihoods, setting a *de novo* mutation prior to 1e-

8. Only sites where Mendelian transmission could be determined unambiguously were phased. The

non-missing phased SNPs were then used as a reference panel for BEAGLE v4.0 (Browning and

Browning 2007). BEAGLE was run without imputing genotypes (impute=false) on the filtered VCF,

with all variants being unphased. The parent-offspring relationships in the reference panel were

specified to inform phasing-by-transmission with BEAGLE, except for the five *C. roulei* samples,

which were not included in a trio and were statistically phased. Datasets **#1**, **#3** and **#4** are based on

this phased VCF.

A genomic region with high coverage failed to be genotyped in the introgression hotspot

(defined between 700 Kb and 1.5 Mb). This region was set as the "missing data region" and defined

from 1,009,000 to 1,055,000 bp.

<u>Analyses of population structure</u>

We used a Principal Component Analysis (PCA) to assess the partition of genetic variation in our

sample of 45 individuals (i.e., all individuals except the three poorly sequenced parents, and the *C.*

edwardsi individuals). SNPs were LD-pruned with PLINK v1.9 (Purcell et al. 2007) using a window

size (WD) of 20 SNPs, a window step size (CT) of 5 SNPs and a linkage threshold (r^2) of 0.1. PLINK

was then used to run a PCA on the unlinked SNPs. We recorded the amount of genotypic variance

explained by each principal component (PC) and the SNP weights on each PC. Only the first two PCs

were relevant to visualize population structure and were plotted using the R package tidyverse.

We used VCFtools v0.1.15 (Danecek et al. 2011) on all SNPs to calculate the per-site

nucleotide diversity (site-pi) in each population and the per-site F_{ST} (weir-fst-pop, Weir and

Cockerham 1984) between populations. We then calculated the average and maximum of these

statistics for each chromosome in non-overlapping windows of 10 Kb. Windows with less than 10

SNPs were excluded. The linkage disequilibrium on chromosome 5 (where an introgression hotspot

was detected) in the *C. intestinalis* individuals was estimated with the function "hap-r2" of VCFtools.

It was based on the calculation of the r^2 among all fixed SNPs (phased) between *C. robusta* and *C.*

roulei.

Detection of introgression with summary statistics

To evaluate the extent of genome-wide admixture, we computed the D-statistic (Green et al. 2010;

Patterson et al. 2012) from a polarized set of SNPs using the outgroup species, *C. edwardsi*. The

following topology was applied: (((P1 = *C. roulei* ; P2 = *C. intestinalis*) ; P3 = *C. robusta*) ; O = *C.*

edwardsi). Therefore, a positive value of D indicates an excess of ABBA sites, and so an excess of

shared ancestry of *C. robusta* with *C. intestinalis* over that shared with *C. roulei*. We also estimated the

fraction of the genome introgressed with the *fd* statistic (Martin et al. 2015), calculated in non-

overlapping windows of 100 SNPs.

Detection of introgression with local ancestry inference

We used Chromopainter (available in fineSTRUCTURE v2.0.7) to perform local ancestry inference

based on the phased dataset. *C. intestinalis* was considered as the recipient population, while *C.*

robusta and *C. roulei* (the latter being a non-introgressed population of *C. intestinalis*) were the donor

populations. We used ten iterations of the expectation-maximization algorithm to estimate the

probability of each position along each *C. intestinalis* haplotype to come from *C. robusta* or *C. roulei*.

We then determined the boundaries of each ancestry tract. A given position was considered originating

from *C. robusta* if this probability was >0.95. To define the tracts, an extension from this focal position

was then made as long as this probability was above 0.5 at the surrounding positions (Duranton et al.

2018).

Various statistics were then calculated focusing on the introgressed tracts originating from *C.*

robusta (i.e. those found in *C. intestinalis* haplotypes, but with a *C. robusta* ancestry): *i*) the *C. robusta*

ancestry fraction per individual, *ii*) the tract length, and *iii*) the frequency of the alleles lying on the

tracts. No filter on the minimal tract length was applied, and missing data were not allowed for the

allele frequency calculation.

We performed additional analyses on the coding sequences (CDS). They were obtained by

extracting the biallelic SNPs from the phased VCF. Then, the VCF was converted into a fasta file, and

exons were extracted with bedtools v2.25.0 based on the annotation file (HT.Gene.gff3) of the

reference genome. The CDS were classified as introgressed or not using the bounds inferred from

Chromopainter. The following statistics were calculated for the CDS: *i*) the pairwise nucleotide

diversity (π, Tajima 1983), *ii*) the raw divergence between *C. robusta* and *C. intestinalis* (d_{XY}, Nei and

Li 1979), and *iii*) the G_{min} measured as minimum(d_{XY})/average(d_{XY}) (Geneva et al. 2015).

<u>Testing for selection</u>

Selection for an adaptive variant is expected to reduce haplotype variation in flanking regions,

producing unusually long haplotypes (Sabeti et al. 2002). To capture such a signal, we measured the

extended haplotype homozygosity (EHH) score from the phased dataset using SelScan v2.0.0 (Szpiech

2021). Target SNPs were identified as the *C. robusta* alleles with the highest frequency to the left

(959,519 bp) and right (1,061,854 bp) of the "missing data region" on chromosome 5. The maximum

extension from the target SNP for a single EHH computation was 100 Kb. Then, we calculated with

SelScan the (absolute) integrated haplotype score (iHS). Values were normalized using the norm

v1.3.0 utility with 100 frequency bins over 50-Kb non-overlapping windows. Finally, we estimated the

proportion of SNPs in each window associated with extreme iHS values (iHS>3, which refers to the

99th quantile of the iHS distribution).

We also tested for the footprint of selective sweeps using SweepFinder v2.0 (DeGiorgio et al.

2016) and adaptive introgression using VolcanoFinder v1.0 (Setter et al. 2020). These methods are

based on polarized SNPs (using the outgroup species *C. edwardsi*) and do not use phase information.

Chromosomes were scanned with the two methods, and a log-ratio test for selection was performed at

test sites spaced by 1Kb.

Finally, SplitsTree4 V4.17.0 (Huson and Bryant 2006) was used on the phased dataset to

produce neighbor-joining trees from 50-Kb windows framing the "missing data region" on

chromosome 5.

<u>Analyses of copy number variation</u>

To overcome the absence of genotyping in the "missing data region" (due to our filtering of repeated

regions), we analyzed the read depth of the variants directly from the unfiltered bam files. Counts of

the reference and alternate alleles were collected with GATK (CollectAllelicCounts) from the bam

files, excluding duplicate reads and positions with a base quality (BQ) < 20. Candidate SNPs were

defined based on their variant allele fraction (VAF = alternate read depth / total read depth). The

following criteria were applied to identify variants differentiated between *C. robusta* and *C. roulei* (the

latter is used as a non-introgressed *C. intestinalis* population), and introgressed into *C. intestinalis*:

VAF <= 50% in *C. intestinalis*, VAF >= 85% (or 90%) in *C. roulei* and VAF <= 15% (or 10%) in *C.*

robusta. The copy number at each candidate SNP was then calculated as its allele read depth

normalized by the per-site read depth averaged across all sites (excluding sites with less than ten

reads) for each individual. Variants were annotated using the HT.Gene.gff3 file of the reference

genome.

<u>Demographic inferences</u>

We reconstructed the divergence history of *C. robusta* and *C. intestinalis* from the folded joint site

frequency spectrum (jSFS) using moments (Jouganous et al. 2017). No missing data was allowed, and

the SNPs were LD-pruned with PLINK v1.9 using a window size (WD) of 10 SNPs, a window step

size (CT) of 10 SNPs and a linkage threshold (r^2) of 0.5. We defined five demographic scenarios,

following (Fraïsse et al. 2018): SI = strict isolation, IM = isolation with continuous migration, SC =

secondary contact, AM = ancient migration, PER = periodic connectivity with both an ancient and a

current period of gene flow. Different versions of these scenarios were tested, following (Fraïsse et al.

2021): bbN = genomic heterogeneity of the effective population sizes (to capture the effect of

background selection), bbM = genomic heterogeneity of the effective migration rates (to capture the

effect of interspecies barriers to gene flow), 2N2M = combining both types of heterogeneities, "" = no

heterogeneities. Parameters were as follows: T = times in years (assuming two generations per year in

European waters), Ne = effective population sizes in numbers of individuals, m = migration rates

(independently estimated in both directions), %Barriers = fraction of the genome experiencing null

migration (i.e. species barriers and their associated loci), %Ne$_{reduced}$ = fraction of the genome

experiencing reduced Ne due to background selection, HRF = factor by which Ne is reduced. See

Tables S3 and **S4** for details. The scripts used to define the demographic models and run the

inferences are available in the **Supplementary Scripts**.

Each demographic model was then fitted to the observed jSFS, with singletons masked. We

ran five independent runs from randomized starting parameter values for each model. Likelihood

optimization was performed using a "dual annealing" algorithm (optimize_dual_anneal). It consists of

a series of global optimizations, each followed by a local optimization ("L-BFGS-B" method).

Settings of the global optimizations were as follows: maximum number of search iterations = 100,

initial temperature = 50, acceptance parameter = 1, and visit parameter = 1.01. The maximum number

of search iterations for the local optimization was set to 100. Model comparisons were made using the

Akaike information criterion (AIC), calculated as $2*k$ - $2*$ML, where k is the number of parameters in

the model, and ML its maximum log-likelihood value across the five runs.

Data Availability

634 Sequence reads have been deposited in NCBI Sequence Read Archive (SRA) under the accession

635 number PRJNA813009. Supplementary Data is available from Zenodo: XXX. Supplementary Figures,

636 Tables and Scripts can be found in the Supporting Information.

Acknowledgments

The authors are grateful to the divers of the Marine Operations department (*Service Mer &*

Observation) at the Roscoff Biological Station for the *Ciona* sampling in Brittany, and to Marine

Malfant, Sebastien Darras and divers of the laboratory Arago in Banyuls-Sur-Mer for providing the

Ciona edwardsi and *C. roulei* samples, and Charlotte Roby for DNA extractions. The authors are very

thankful to Jerome Coudret and Sarah Bouchemousse for their help for carrying the laboratory crosses

at Roscoff. We thank Véronique Dhennin from the LIGAN genomics platform (Lille, France), and

Nicolas González from the FASTERIS platform (Plan-les-Ouates, Switzerland). This work benefited

from the Montpellier Bioinformatics Biodiversity platform supported by the LabEx CeMEB, an ANR

"Investissements d'avenir" program (ANR-10-LABX-04-01). This work benefited from funding of the

French National Research Agency (ANR) with regards the ANR Project HYSEA (no. ANR-12-BSV7-

0011). It also benefited from the MarEEE project funded through the French National Research

Agency (ANR) under the "Investissements d'Avenir" programme with the reference ANR-16-IDEX-

0006 (i-site MUSE). The funders had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript.

References

Bouchemousse S, Bishop JDD, Viard F. 2016. Contrasting global genetic patterns in two biologically

similar, widespread and invasive Ciona species (Tunicata, Ascidiacea). *Sci. Rep.* 6:24875.

Bouchemousse S, Lévêque L, Dubois G, Viard F. 2016. Co-occurrence and reproductive synchrony do

not ensure hybridization between an alien tunicate and its interfertile native congener. *Evol. Ecol.*

30:69–87.

Bouchemousse S, Lévêque L, Viard F. 2017. Do settlement dynamics influence competitive

interactions between an alien tunicate and its native congener? *Ecol. Evol.* 7:200–213.

Bouchemousse S, Liautard-Haag C, Bierne N, Viard F. 2016. Distinguishing contemporary

hybridization from past introgression with postgenomic ancestry-informative SNPs in strongly

differentiated Ciona species. *Mol. Ecol.* 25:5527–5542.

Browning SR, Browning BL. 2007. Rapid and accurate haplotype phasing and missing-data inference

for whole-genome association studies by use of localized haplotype clustering. *Am. J. Hum.*

Genet. 81:1084–1097.

Calfee E, Agra MN, Palacio MA, Ramírez SR, Coop G. 2020. Selection and hybridization shaped the

rapid spread of African honey bee ancestry in the Americas. *PLoS Genet.* 16:e1009038.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth

GT, Sherry ST, et al. 2011. The variant call format and VCFtools. *Bioinformatics* 27:2156–2158.

DeGiorgio M, Huber CD, Hubisz MJ, Hellmann I, Nielsen R. 2016. SweepFinder2: increased

sensitivity, robustness and flexibility. *Bioinformatics* 32:1895–1897.

Duranton M, Allal F, Fraïsse C, Bierne N, Bonhomme F, Gagnaire P-A. 2018. The origin and

remolding of genomic islands of differentiation in the European sea bass. *Nature Comm.* 9:2518.

Fraïsse C, Popovic I, Mazoyer C, Spataro B, Delmotte S, Romiguier J, Loire É, Simon A, Galtier N,

Duret L, et al. 2021. DILS: Demographic inferences with linked selection by using ABC. *Mol.*

Ecol. Resour. 21:2629–2644.

Fraïsse C, Roux C, Gagnaire P-A, Romiguier J, Faivre N, Welch JJ, Bierne N. 2018. The divergence

history of European blue mussel species reconstructed from Approximate Bayesian Computation:

the effects of sequencing techniques and sampling strategies. *PeerJ* 6:e5198.

Geneva AJ, Muirhead CA, Kingan SB, Garrigan D. 2015. A New Method to Scan Genomes for

Introgression in a Secondary Contact Model. *PLoS One* 10:e0118621.

Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai W, Fritz

MH-Y, et al. 2010. A draft sequence of the Neandertal genome. *Science* 328:710–722.

Hudson J, Viard F, Roby C, Rius M. 2016. Anthropogenic transport of species across native ranges:

unpredictable genetic and evolutionary consequences. *Biol. Lett.* 12:20160620.

Huson DH, Bryant D. 2006. Application of phylogenetic networks in evolutionary studies. *Mol. Biol.*

Evol. 23:254–267.

Jouganous J, Long W, Ragsdale AP, Gravel S. 2017. Inferring the Joint Demographic History of

Multiple Populations: Beyond the Diffusion Approximation. *Genetics* 206:1549–1567.

Kaback DB. 1996. Chromosome-size dependent control of meiotic recombination in humans. *Nat.*

Genet. 13:20–21.

Klopfstein S, Currat M, Excoffier L. 2006. The fate of mutations surfing on the wave of a range

expansion. *Mol. Biol. Evol.* 23:482–490.

Kulmuni J, Butlin RK, Lucek K, Savolainen V, Westram AM. 2020. Towards the completion of

speciation: the evolution of reproductive isolation beyond the first barriers. *Philos. Trans. R. Soc.*

Lond. B Biol. Sci. 375:20190528.

Le Moan A, Roby C, Fraïsse C, Daguin-Thiébaut C, Bierne N, Viard F. 2021. An introgression

breakthrough left by an anthropogenic contact between two ascidians. *Mol. Ecol.* 30:6718–6732.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics 25:1754–1760.

Main BJ, Everitt A, Cornel AJ, Hormozdiari F, Lanzaro GC. 2018. Genetic variation associated with

increased insecticide resistance in the malaria mosquito, Anopheles coluzzii. *Parasit. Vectors*

11:225.

Malfant M, Darras S, Viard F. 2018. Coupling molecular data and experimental crosses sheds light

about species delineation: a case study with the genus Ciona. *Sci. Rep.* 8:1480.

Martin SH, Davey JW, Jiggins CD. 2015. Evaluating the Use of ABBA–BABA Statistics to Locate

Introgressed Loci. *Mol. Biol. Evol.* 32:244–257.

Martin SH, Jiggins CD. 2017. Interpreting the genomic landscape of introgression. *Curr. Opin. Genet.*

Dev. 47:69–74.

Maxwell CS, Mattox K, Turissini DA, Teixeira MM, Barker BM, Matute DR. 2019. Gene exchange

between two divergent species of the fungal human pathogen, Coccidioides. *Evolution* 73:42–58.

McFarlane SE, Pemberton JM. 2019. Detecting the True Extent of Introgression during Anthropogenic

Hybridization. *Trends Ecol. Evol.* 34:315–326.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D,

Gabriel S, Daly M, et al. 2010. The Genome Analysis Toolkit: a MapReduce framework for

analyzing next-generation DNA sequencing data. *Genome Res.* 20:1297–1303.

Moran BM, Payne C, Langdon Q, Powell DL, Brandvain Y, Schumer M. 2021. The genomic

consequences of hybridization. *Elife* 10:e69016.

Nei M, Li WH. 1979. Mathematical model for studying genetic variation in terms of restriction

endonucleases. *Proc. Natl. Acad. Sci. U. S. A.* 76:5269–5273.

Nelson DR. 1998. Metazoan cytochrome P450 evolution. *Comp. Biochem. Physiol. C Pharmacol.*

Toxicol. Endocrinol. 121:15–22.

Nelson RM, Wallberg A, Simões ZLP, Lawson DJ, Webster MT. 2017. Genomewide analysis of

admixture and adaptation in the Africanized honeybee. *Mol. Ecol.* 26:3603–3617.

Noé L, Kucherov G. 2005. YASS: enhancing the sensitivity of DNA similarity search. *Nucleic Acids*

Res. 33:W540–W543.

North HL, McGaughran A, Jiggins CD. 2021. Insights into invasive species from whole-genome

resequencing. *Mol. Ecol.* 30:6289–6308.

Nydam ML, Harrison RG. 2010. Polymorphism and divergence within the ascidian genus Ciona. *Mol.*

Phylogenet. Evol. 56:718–726.

Nydam ML, Harrison RG. 2011. Introgression despite substantial divergence in a broadcast spawning

marine invertebrate. *Evolution* 65:429–442.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de Bakker

PIW, Daly MJ, Sham PC. 2007. PLINK: a toolset for whole-genome association and population-

based linkage analysis. *Am. J. Hum. Genet.* 81.

Ottenburghs J. 2021. The genic view of hybridization in the Anthropocene. *Evol. Appl.* 14:2342–2360.

Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y, Genschoreck T, Webster T, Reich D.

2012. Ancient admixture in human history. *Genetics* 192:1065–1093.

Puinean AM, Foster SP, Oliphant L, Denholm I, Field LM, Millar NS, Williamson MS, Bass C. 2010.

Amplification of a cytochrome P450 gene is associated with resistance to neonicotinoid

insecticides in the aphid Myzus persicae. *PLoS Genet.* 6:e1000999.

Racimo F, Sankararaman S, Nielsen R, Huerta-Sánchez E. 2015. Evidence for archaic adaptive

introgression in humans. *Nat. Rev. Genet.* 16:359–371.

Ravinet M, Yoshida K, Shigenobu S, Toyoda A, Fujiyama A, Kitano J. 2018. The genomic landscape

at a late stage of stickleback speciation: High genomic divergence interspersed by small localized

regions of introgression. *PLoS Genet.* 14:e1007358.

Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP. 2011.

Integrative genomics viewer. *Nat. Biotechnol.* 29:24–26.

Roux C, Fraïsse C, Romiguier J, Anciaux Y, Galtier N, Bierne N. 2016. Shedding Light on the Grey

Zone of Speciation along a Continuum of Genomic Divergence. *PLoS Biol.* 14:e2000234.

Roux C, Tsagkogeorga G, Bierne N, Galtier N. 2013. Crossing the species barrier: genomic hotspots of

introgression between two highly divergent Ciona intestinalis species. *Mol. Biol. Evol.* 30:1574–

1587.

Sabeti PC, Reich DE, Higgins JM, Levine HZP, Richter DJ, Schaffner SF, Gabriel SB, Platko JV,

Patterson NJ, McDonald GJ, et al. 2002. Detecting recent positive selection in the human genome

from haplotype structure. *Nature* 419:832–837.

Sachdeva H, Barton NH. 2018. Introgression of a Block of Genome Under Infinitesimal Selection.

Genetics 209:1279–1303.

Sams AJ, Dumaine A, Nédélec Y, Yotova V, Alfieri C, Tanner JE, Messer PW, Barreiro LB. 2016.

Adaptively introgressed Neandertal haplotype at the OAS locus functionally impacts innate

immune responses in humans. *Genome Biol.* 17:246.

Satou Y, Nakmura R, Deli Y, Yoshida R, Hamada M, Fujie M, Hisata K, Takeda H, Satoh N. 2019. A

nearly-complete genome of Ciona intestinalis type A (C. robusta) reveals the contribution of

inversion to chromosomal evolution in the genus Ciona. *Genome Biol. Evol.* 11:3144–3157.

Schmidt JM, Good RT, Appleton B, Sherrard J, Raymant GC, Bogwitz MR, Martin J, Daborn PJ,

Goddard ME, Batterham P, et al. 2010. Copy number variation and transposable elements feature

in recent, ongoing adaptation at the Cyp6g1 locus. *PLoS Genet.* 6:e1000998.

Setter D, Mousset S, Cheng X, Nielsen R, DeGiorgio M, Hermisson J. 2020. VolcanoFinder: Genomic

scans for adaptive introgression. *PLoS Genet.* 16:e1008867.

Shang H, Hess J, Pickup M, Field DL, Ingvarsson PK, Liu J, Lexer C. 2020. Evolution of strong

reproductive isolation in plants: broad-scale patterns and lessons from a perennial model group.

Philos. Trans. R. Soc. Lond. B Biol. Sci. 375:20190544.

Shenkar N, Swalla BJ. 2011. Global diversity of Ascidiacea. *PLoS One* 6:e20657.

Shoguchi E, Kawashima T, Satou Y, Hamaguchi M, Sin-I T, Kohara Y, Putnam N, Rokhsar DS, Satoh

N. 2006. Chromosomal mapping of 170 BAC clones in the ascidian Ciona intestinalis. *Genome*

Res. 16:297–303.

Simon A, Arbiol C, Nielsen EE, Couteau J, Sussarellu R, Burgeot T, Bernard I, Coolen JWP, Lamy J-

B, Robert S, et al. 2020. Replicated anthropogenic hybridisations reveal parallel patterns of

admixture in marine mussels. *Evol. Appl.* 13:575–599.

Stachowicz JJ, Terwin JR, Whitlatch RB, Osman RW. 2002. Linking climate change and biological

invasions: Ocean warming facilitates nonindigenous species invasions. *Proc. Natl. Acad. Sci. U.*

S. A. 99:15497–15500.

Stankowski S, Westram AM, Zagrodzka ZB, Eyres I, Broquet T, Johannesson K, Butlin RK. 2020. The

evolution of strong reproductive isolation between sympatric intertidal snails. *Philos. Trans. R.*

Soc. Lond. B Biol. Sci. 375:20190545.

Staubach F, Lorenc A, Messer PW, Tang K, Petrov DA, Tautz D. 2012. Genome patterns of selection

and introgression of haplotypes in natural populations of the house mouse (Mus musculus). *PLoS*

Genet. 8:e1002891.

Szpiech ZA. 2021. Selscan 2.0: scanning for sweeps in unphased data. *biorxiv*. 2021.10.22.465497.

Tajima F. 1983. Evolutionary relationship of DNA sequences in finite populations. *Genetics* 105:437–

460.

Turissini DA, Matute DR. 2017. Fine scale mapping of genomic introgressions within the Drosophila

yakuba clade. *PLoS Genet.* 13:e1006971.

Valencia-Montoya WA, Elfekih S, North HL, Meier JI, Warren IA, Tay WT, Gordon KHJ, Specht A,

Paula-Moraes SV, Rane R, et al. 2020. Adaptive Introgression across Semipermeable Species

Boundaries between Local Helicoverpa zea and Invasive Helicoverpa armigera Moths. *Mol. Biol.*

Evol. 37:2568–2583.

Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A, Jordan T,

Shakir K, Roazen D, Thibault J, et al. 2013. From FastQ data to high confidence variant calls: the

Genome Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinformatics* 43:11.10.1–

11.10.33.

Viard F, Riginos C, Bierne N. 2020. Anthropogenic hybridization at sea: three evolutionary questions

relevant to invasive species management. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 375:20190547.

Vizzini A, Bonura A, La Paglia L, Fiannaca A, La Rosa M, Urso A, Mauro M, Vazzana M, Arizza V.

2021. Transcriptomic Analyses Reveal 2 and 4 Family Members of Cytochromes P450 (CYP)

Involved in LPS Inflammatory Response in Pharynx of *Ciona robusta*. *Int. J. Mol. Sci.* 22:11141.

Weir BS, Cockerham CC. 1984. Estimating f-statistics for the analysis of population structure.

Evolution 38:1358–1370.

Wondji CS, Irving H, Morgan J, Lobo NF, Collins FH, Hunt RH, Coetzee M, Hemingway J, Ranson

H. 2009. Two duplicated P450 genes are associated with pyrethroid resistance in Anopheles

funestus, a major malaria vector. *Genome Research* 19:452–459.

Yainna S, Nègre N, Silvie PJ, Brévault T, Tay WT, Gordon K, dAlençon E, Walsh T, Nam K. 2021.

Geographic Monitoring of Insecticide Resistance Mutations in Native and Invasive Populations

of the Fall Armyworm. *Insects* 12:468.

Yamasaki YY, Kakioka R, Takahashi H, Toyoda A, Nagano AJ, Machida Y, Møller PR, Kitano J. 2020.

Genome-wide patterns of divergence and introgression after secondary contact between Pungitius

sticklebacks. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 375:20190548.

Zhan A, Briski E, Bock DG, Ghabooli S, MacIsaac HJ. 2015. Ascidians as models for studying

invasion success. *Mar. Biol.* 162:2449–2470.

Zimmer CT, Garrood WT, Singh KS, Randall E, Lueke B, Gutbrod O, Matthiesen S, Kohler M, Nauen

R, Davies TGE, et al. 2018. Neofunctionalization of Duplicated P450 Genes Drives the Evolution

of Insecticide Resistance in the Brown Planthopper. *Curr. Biol.* 28:268–274.e5.

Figure 1 Genetic population structure. **A.** Geographical location of the samples in the English Channel and

Iroise Sea (*C. robusta* and *C. intestinalis*) and the Mediterranean Sea (*C. roulei*). Numbers in brackets refer to the

sample size of each population. "Lab F1s" indicates the intraspecific and interspecific offspring produced in the

laboratory. Further information on samples is provided in **Table S1**. The color code (*C. robusta* in red, *C.*

intestinalis in blue and *C. roulei* in green) is used throughout the manuscript. **B.** Principal Component Analysis

of 45 individuals genotyped at 194,742 unlinked SNPs (pruning threshold: $r^2 > 0.1$). Numbers in brackets refer to

the proportion of variance explained by each axis. The F1s were considered as supplementary individuals in the

PCA, and three poorly sequenced parents were removed from the analysis (see **Table S1**). **C.** SNP weights to the

first axis of the PCA (after removing the SNPs contributing less than the 75th quantile of the weight

distribution). The introgression hotspot on chromosome 5 is highlighted with an arrow. Dataset **#1** was used.

Figure 2 Local ancestry patterns of the *C.*
intestinalis genomes sampled in the English
Channel and Iroise Sea using 640,044 phased
SNPs. *C. robusta* and *C. roulei* were used as the
donor populations. **A.** Physical mapping across
the 14 chromosomes of the frequency of the *C.*
robusta tracts introgressed into *C. intestinalis*.
The color gradient (blue to yellow to red)
follows the gradient in allele frequency from
low to high; regions in white correspond to null
introgression. The direction of the arrow
indicates the coordinate direction from top
(start) to bottom (end). The R package
RIdeograms was used for the plotting. **B.**
Length distribution of the *C. robusta*
introgressed tracts (*n*=1,143 tracts). The
maximum tract length is 156 Kb, the average is
2.6 Kb, and the median is 0.38 Kb. A blue
dashed line depicts the 95th quantile of the
length distribution (8.3 Kb), and it was used as
a threshold to delineate long tracts. A total of 38
of 57 long tracts were detected on chromosome
5 (red arrows). **C.** Allele frequency of the SNPs
lying on the *C. robusta* introgressed tracts
(*n*=621,249 variants). The maximum frequency

is 1, the average is 0.14, and the median is 0.08. On the right, the frequency of variants lying on short tracts
(upper panel) or long tracts (lower panel) is depicted. Red arrows indicate the tracts on chromosome 5. Allele
frequency was calculated, excluding any position with missing data. Dataset **#3a** was used.

Figure 3 Inference of the divergence history between *C. robusta* and *C. intestinalis* with moments. **A.** AIC value of the best run for each model. **B.** Observed site frequency spectrum (SFS), modeled SFS and residuals of the best model. Maximum likelihood values of the parameters are provided. **C.** Same as in **B** but for the second-best model. Analyses were based on the folded SFS after LD-pruning the SNPs. Five demographic scenarios were modeled: SI = strict isolation, IM = isolation with continuous migration, SC = secondary contact, AM = ancient migration, PER = periodic connectivity. Different versions of these scenarios were tested: bbN = genomic heterogeneity of the effective population sizes, bbM = genomic heterogeneity of the effective migration rates, 2N2M = both types of heterogeneities, "" = no heterogeneities. Five replicates were run for each model. Parameters are as follows: T = times in years, assuming two generations per year in European waters (the "GF" label refers to gene flow), Ne = effective population sizes in numbers of individuals, m = migration rates (direction given by the arrow), %Barriers = proportion of the genome with null migration, %Ne$_{reduced}$ = fraction of the genome experiencing reduced Ne, HRF = factor by which Ne is reduced. Full details are provided in **Table S3**. Dataset **#5** was used, excluding chromosome 5.

Best model: PER2N2M ; AIC : 3,342 (2nd best model: SC2N2M ; AIC : 3,350)

SI   SIbbN   AM   AMbbN   AMbbM   AM2N2M   IM   IMbbN   IMbbM   IM2N2M   SC   SCbbN   SCbbM   SC2N2M   PER   PERbbN   PERbbM   PER2N2M

B

DATA — MODEL
C. intestinalis vs *C. robusta*
RESIDUALS

Best model PER2N2M

T_{1_GF} = 399 Ky ; $T_{2\_NO\_GF}$ = 1.5 My ; T_{3_GF} = 30 Ky

Ne_{int} = 13.1 M ; Ne_{rob} = 1.3 M ; Ne_{rob} = 9.9% Ne_{int}

$m_{rob \to int}$ >>> $m_{int \to rob}$; %Barriers = 61%

HRF = 0.038 ; %Ne$_{reduced}$ = 69%

C

DATA — MODEL
C. intestinalis vs *C. robusta*
RESIDUALS

2nd best model SC2N2M

$T_{1\_NO\_GF}$ = 1.8 My ; T_{2_GF} = 0.2 Ky

Ne_{int} = 12.6 M ; Ne_{rob} = 1.2 M ; Ne_{rob} = 9.5% Ne_{int}

$m_{rob \to int}$ >>> $m_{int \to rob}$; %Barriers = 74%

HRF = 0.039 ; %Ne$_{reduced}$ = 67%

Figure 4 Analyses of the introgression hotspot on chromosome 5. **A.** Maximum F_{ST} between *C. robusta* and *C.*

intestinalis was calculated in non-overlapping 10 Kb windows along chromosome 5. Windows with less than 10

SNPs were excluded. The x-axis is in bp. **B.** Haplotypes of the *C. intestinalis* individuals in the two sampled

localities (sample IDs are depicted on the right, see **Table S1**). Each individual displays two haplotypes

delimited by horizontal lines. The *C. robusta* introgressed tracts are shown as red bars. The white background

represents the non-introgressed tracts and missing data. The tract boundaries were determined based on the

ancestry probability of each position, as shown in **Figure S7**. **C.** Frequency of the *C. robusta* alleles lying on the

introgressed tracts along chromosome 5. Allele frequency was calculated, excluding any position with missing

data (e.g. the nearly fixed SNP at position 28,801 bp on panel **B** was excluded and designated with the first cross

on panel **C**). The grey horizontal band running through all panels refers to the "missing data region" (due to high

coverage) in the core region of the hotspot (from 1,009,000 to 1,055,000 bp). **D.** Linkage disequilibrium pattern

between the 111,951 SNPs fixed between *C. robusta* and *C. roulei*. The color scale indicates the level of LD

from blue (low) to red (high). **E.** Haplotype-based selection test using *SelScan*. EHH is shown for the *C. robusta*

haplotype (red) and the other haplotypes (black) in *C. intestinalis* (upper panel) and *C. roulei* (lower panel) using

a 100 Kb maximal extension. A separate analysis was done on the left and right of the "missing data region"

(grey band) using the most frequent *C. robusta* allele closest to the grey band as target SNP. Absolute iHS was

calculated based on the EHH results and normalized in windows of 50 Kb. The threshold value of the normalized

iHS was set to 3 (which refers to the 99th quantile). Dataset **#2** was used for the F_{ST}, **#3a** for Chromopainter

haplotypes and **#4** for the LD triangle.

Figure 5 Copy number variation at two candidate SNPs on the cytochrome P450 family 2 subfamily U gene. The two SNPs (labeled with their position in bp) lie in the "missing data region" of the introgression hotspot on chromosome 5. Candidates were defined as having a variant allele fraction, VAF <= 50% in *C. intestinalis*, a VAF >= 90% in *C. roulei* and a VAF <= 10% in *C. robusta*. No candidates were found in the other direction (i.e. with the minor VAF in *C. roulei*). Copy number at each SNP was calculated as its allele read depth normalized by the per-site read depth averaged across all sites (excluding sites with less than ten reads) for each individual (labeled on the left). A copy number of one (vertical dashed line) means that the SNP lies on a single-copy locus. Values for the *C. robusta* allele (red) and the *C. intestinalis* allele (blue) are separately shown. Read depth was obtained from the bam files. Horizontal dashed lines separate the different species, and *C. intestinalis* individuals introgressed at the hotspot (see **Figure 4**) were labeled as "introgressed". Dataset **#7** was used.