Validation of Reference Genes for RT-qPCR Relative Expression Analysis in Pre-Adult Stages of *Taenia solium*

David Castaneda-Carpio[1][¶], Jose Maravi[1][¶], Renzo Gutierrez-Loli[1]*, Valeria Villar[1] Juan Blume La Torre[1], Segundo W. Del Aguila[1] & Cristina Guerra-Giraldez[1]*[†].

[1] Laboratorio de Proliferación y Regeneración Celular. Laboratorios de Investigación y Desarrollo, Facultad de Ciencias e Ingeniería. Universidad Peruana Cayetano Heredia. Lima, Peru.

* Correspondence: renzo.gutierrez@upch.pe; cristina.guerra@upch.pe

¶ These authors contributed equally to this work.

† Lead contact.

SUMMARY

The larvae-to-adult development of the zoonotic parasitic tapeworm *Taenia solium* involves significant but often clinically overlooked events crucial in cestode biology. The early-adult stages can be studied in vitro, providing a valuable model to examine scolex evagination, strobilation, and worm development. Without a stage-specific transcriptome, postgenomic data exploration followed by single-gene relative expression analysis using RT-qPCR (reverse transcription-quantitative PCR) are effective strategies to study gene regulation during parasite development. However, achieving accurate comparisons with this approach requires the validation of an endogenous reference gene (RG).

To address this, we analyzed the expression stability of 17 candidate reference genes (RGs), representing various biological processes, in the context of the in vitro-induced early adult stages of *T. solium* larvae (cysts). RT-qPCR of the candidate RGs was performed on different stages, defined by distinct morphology in culture, and gene expression stability was comprehensively analyzed using the RefFinder tool. Genes *pgk1*, *bact1*, *mapk3*, *tbp*, *rpl13*, and *cox1* were ranked as the most stable and were used to normalize the expression of *h2b* and *wnt11a*, which are involved in proliferation and strobilation processes in parasitic tapeworms. This study represents the first attempt to identify reliable normalization standards for transcript analysis in the genus *Taenia*.

INTRODUCTION

The genus *Taenia* includes over a hundred species of tapeworms, parasites of wildlife and livestock. Three among these, *T. solium*, *T. saginata*, and *T. asiatica,* cause zoonotic infections, leading to taeniasis and cysticercosis. These helminthiases are part of the neglected tropical diseases prioritized by the WHO[1].

Cysticercosis can cause life-threatening conditions, particularly in poor, remote areas of developing countries in Africa, Asia, and Latin America, where it impacts the health and livelihoods of rural farming communities. The latest WHO report identified *T. solium* as a leading cause of death from food-borne diseases, resulting in a total loss of 2.8 million disability-adjusted life-years (DALYs)[2,3].

T. solium has a complex life cycle involving an intermediate host (pig) that harbors the larvae in its tissues (cysticercosis), and a definitive host (human) that carries the adult

tapeworm in its intestines (taeniasis). The tapeworm carrier spreads thousands of
microscopic parasite eggs through feces containing infective hexacanth embryos or
oncospheres. Upon ingestion by free-roaming pigs or other humans, active embryos are
released from their eggshells and cross the intestinal epithelium, spreading through the
bloodstream to skeletal muscle, heart, and nervous system, where they lodge and develop
into vesicular larvae (cysts)[4]. Poorly cooked pig meat may thus contain viable cysts. The
gastric and bile acids in the human digestive tube activate these, and the already formed
scolex inside the vesicle evaginates to attach to the intestinal wall. There, the parasite
grows into an adult tapeworm by producing a strobilus of continuous proglottids that will
carry new eggs when mature[4,5].

Due to the complexity of the host-parasite interactions, the complete life cycle of *T. solium*
has yet to be reproduced in the laboratory. However, observations in experimental or
naturally infected animals have produced valuable knowledge about the oncosphere-to-
cyst development and cyst-associated immunopathology in several tissues[6,7]. There is
much less information related to cyst-to-adult development, partly because taeniasis has
generally mild or non-specific symptoms[4,8]. Nonetheless, studying the stages of taeniasis
is necessary to understand endemicity's critical factors. Notably, a tapeworm carrier is the
leading risk factor for acquiring cysticercosis, spreading the infective stage and
perpetuating the parasite's life cycle.

Cyst-to-adult development involves several processes essential in cestode biology,
associated with the scolex's evagination and further strobilation[9]. The early adult stage in
T. solium can be addressed in vitro, starting from chemically[10] or enzymatically treated
cysts[11].

With the publication of the *T. solium* genome[12], the first partial cyst transcriptomes
obtained by RNA-seq[13,14], and the recent transcriptome of the abnormal form of the cyst
found in human subarachnoid neurocysticercosis[15], a significant amount of data is
currently available to study stage-specific events in the parasite's development[16].
Moreover, whole transcriptome analyses in other tapeworms (including families Taeniidae
– genera *Taenia* and *Echinococcus*[17-21], Hymenolepididae[22], and Mesocestoididae[23]) have
led to the identification of conserved pathways among cestodes that complement previous
datasets.

RT-qPCR helps analyze single-gene expression and compare experimental conditions due
to its high sensitivity and specificity. Validating the expression stability of candidate
reference genes (RGs) is thus critical to normalize data when studying any particular
species or stages with RT-qPCR[24]. Although a few analyses for expression stability of RGs
have been done in other cestodes[25,26], a similar validation for the pre-adult stages of *T.*
solium is lacking. Previous RT-qPCR comparisons used canonical housekeeping gene
targets such as *gapdh* (for the enzyme Glyceraldehyde-3-Phosphate Dehydrogenase)[27].

In this study, a representative set of 17 transcripts involved in key biological processes –
such as gene transcription, translation and protein synthesis, protein folding and quality
control, signal transduction, cytoskeleton and cell structure, energy metabolism, and
mitochondrial respiration – were selected as candidate RGs for the in vitro-induced pre-
adult stages of *T. solium*. The expression stability of these genes was analyzed in

cultivated cysts, classified into three morphological stages corresponding to different
moments in the process of scolex evagination. The RefFinder integrative tool, which
combines the statistical algorithms geNorm, NormFinder, BestKeeper, and the deltaCT
method, was used for the analysis.

Finally, to validate the reliability of the top-ranked candidates, these were used to
normalize the expression of two informative genes: *h2b*, involved in cell proliferation during
scolex evagination in tapeworms, and *wnt11a*, a morphogen associated with posterior
development in flatworms that acts as an effector in strobilation.

METHODS

Selection of candidate RGs and primer design

Primers were designed using Primer3[28] and BLAST® software based on the 17 mRNA
sequences of *T. solium* candidate RGs retrieved from GeneDB databases. The basic
parameters used for primer design were as follows: primer length between 18-21 bp, GC
content between 45-55%, melting temperature between 55-60 °C, and amplicon length
between 60-200 bp. Every primer was 100% complementary to the expected primer
binding sites, except for the COX1 primers, which had less (97%) complementarity to the
used mRNA sequence. The details of each selected candidate gene and the biological
features of each primer set are shown in Table 1.

Parasite collection and *in vitro* evagination

Taenia solium cysts were collected from the muscular tissue of a naturally infected pig, as
determined by a positive tongue examination, from a farm at Huancayo, Junin, Peru. Only
undamaged and complete cysts were selected and placed in phosphate buffered-saline
(PBS; 0.1 M, pH 7.4) supplemented with gentamicin (20 µg/mL) and benzylpenicillin (10
µg/mL) for transport.

The evagination of scolices was induced with 0.1% taurocholic acid (TA) (Sigma, St. Louis,
MO) diluted in RPMI 1640 medium, supplemented with 2 g/L $NaHCO_3$ (Merck®,
Burlington, MA). Parasites were cultured in 12-well plates (Corning, Corning, NY) with 2
mL of the described medium at 37°C in a 5% CO_2 atmosphere for up to 5 days. Scolex
evagination was monitored every 12 hours by direct observation to track intermediate
stages until full evagination occurred. Consistent collection criteria were established to
categorize parasites into three distinct stages: cysts with invaginated scolex ("non-
evaginated cysts," or PRE), cysts with early scolex exposure ("recently evaginated cysts,"
or EV), and cysts showing clear strobila ("fully evaginated cysts," or POST).

After culture, all parasites were washed with PBS to remove residual media. The
percentage of evagination, defined as the proportion of parasites with a visibly evaginated
scolex, was recorded for each culture condition and parasite group.

RNA extraction and DNAse treatment

Evaginated and non-evaginated parasites were embedded independently in 1 mL of
TRIzol (Invitrogen, Carlsbad, CA) and admixed with a tissue homogenizer tip (Omni

International, Kennesaw, GA) on ice. RNA was isolated following the manufacturer's
protocol, considering an extra washing step with 75% ethanol. After the solubilization,
whole RNA was treated with 2 U RNAse-free DNAse I (Invitrogen, Carlsbad, CA) for 30
min at 37° C. The DNAse was inactivated with TRIzol, and the RNA was re-purified. RNA
yield was measured by spectrophotometry with NanoDrop® (Thermofisher Scientific,
Waltham, MA), and approximately 100 ng of RNA were loaded in a 1.5% agarose gel
electrophoresis to test its integrity.

Two-step RT-qPCR reaction

cDNA was generated from 100 ng of RNA of every parasite by reverse transcription with
the High-Capacity Reverse Transcription Kit (Thermofisher Scientific, Waltham, MA) in a
final volume of 20 µL. The reaction mix was incubated 15 min at 25° C, 2 h at 37° C and 5
min at 85°C, and stored at 4° C until required.

qPCR reactions were done with the PowerUp® SYBR Green Master Mix (Thermofisher
Scientific, Waltham, MA). For every 10-µL reaction, forward and reverse primers were
added at a final concentration of 500 nM each, and 1 µL of cDNA was used as the
template. The cycling conditions were set in the PikoReal™ Real-Time PCR System
(Thermofisher Scientific, Waltham, MA) as follows: 2 min at 50°C, 2 min at 95°C; and 40
cycles of 15 s at 95°C and 1 min at 60°C. Every sample was run in triplicates, and a
melting curve step was done at the end of every experiment ranging from 60°C to 95°C
with increments of 0.2°C. Additionally, the size and specificity of amplification products
were confirmed with a 1.5% agarose gel electrophoresis. RG datasets with Cq values >35
were excluded for having low abundance.

Amplification efficiency and analytical performance

To determine the PCR amplification efficiency for every candidate RG, 5-point calibration
curves of cDNA were prepared using five 2-fold, 5-fold, or 10-fold serial dilutions. The
linear regression, slopes, and y-axis intercepts for every standard curve were calculated
using Graphpad Prism® software version 6.01 (Graphpad®, San Diego, CA). Every
dilution was tested in triplicate, and amplification efficiency was determined based on the
slope of the log-linear portion of the calibration curve ($10^{-1/slope} - 1$), following the MIQE
guidelines[29].

Data analysis for gene expression stability

The gene expression stability of the candidate RGs was compared and ranked with
different normalization algorithms integrated in the web-based tool RefFinder (available at:
https://www.ciidirsinaloa.com.mx/RefFinder-master/). This program integrates the analysis
performed by the extensively used computational algorithms geNorm[30], Normfinder[31],
BestKeeper[32], and the comparative DeltaCT method[33]. The selection of the most stable
RG was based on the recommended comprehensive ranking calculated by the geometric
mean of the stability scores obtained from each method.

Validation of RG in relative quantification

To validate the reliability of the selected RG, the relative expression of *hb2* and *wnt11a*
was normalized by the most and the least stable genes obtained by the normalization
algorithms. Indeed, both *h2b* and *wnt11a* genes have been reported previously as

differentially expressed upon scolex evagination in parasitic tapeworms[34, 35]. Ten biological
samples per condition and three technical replicates per sample were set up for this
analysis. Fold-change for *h2b*n and *wnt11a* at experimental conditions was analyzed using
the $2^{-\Delta\Delta Cq}$ method[36]. The sequence and other technical details for *h2b* and *wnt11a* primer
sets are shown in Table 1.

RESULTS

In vitro parasite evagination and definition of pre-adult stages

We confirmed that taurocholic acid (TA) induces scolex evagination. This was
demonstrated through direct and macroscopic observations and by quantifying the
percentage of cysts with evaginated scolices. The results showed that after two days in
culture, 0.1% TA in the medium led to twice as many cysts with evaginated scolices as the
untreated ones (Fig 1B).

Fig 1. Taurocholic acid (TA) induces in vitro evagination of *T. solium* cysts. (A)
Overview of the study design and analysis workflow. (B) Quantification of the percentage
of cyst evagination over 120 hours. Each point represents the mean of three independent
experiments, with error bars showing the maximum and minimum values. (C)
Representative macroscopic images of non-evaginated PRE (left) and POST (right)
parasites after 5 days of culture following 0.1% TA induction. Scale bar: 5 mm. These
samples were used for subsequent single-gene expression analysis comparisons.

To study the early development of the pre-adult stages, we examined several conditions,
including uncultured controls and cysts maintained in vitro with 0.1% TA. Our goal was to
identify culture conditions that achieve complete evagination of the scolices, allowing us to
establish three *bona fide* categories for comparing single-gene expression against the
controls. By five days post-TA induction, the parasites had fully evaginated the scolex,
neck, and early strobila from the vesicle (POST). We also included cysts with fully
invaginated scolices collected at approximately 4-8 hours in culture (PRE) and cysts
collected during the 24-48 hour period (EV), when the parasites had recently evaginated
the scolex (Fig 1C).

RT-qPCR amplification efficiency and product specificity

Primers of 17 candidate RG were used to amplify cDNA templates by qPCR. Agarose gel
electrophoresis revealed a single-band amplicon by end-point PCR for each reaction,
consistent with the expected product size (Fig 2A). Similarly, all graphs of the negative first
derivative of the melting curve showed a single peak, indicating that the primers had
strong specificity for the target region and no non-specific amplification occurred (Fig 2D).

Fig 2. Comprehensive Analysis of qPCR Performance for Candidate Reference Genes.
(A) Primer pair specificity for candidate reference genes: End-point PCR products for all
candidate RG were observed on gel electrophoresis and arranged according to product size.
MW: 100bp DNA Molecular Weight Ladder. (B) Raw Cq values across parasite samples:
Comparison of raw Cq values from qPCR amplification across all parasite samples,

showing gene expression for each candidate RG. (C) Cq Value distribution based on
treatment and morphology: The distribution of Cq values for each candidate RG, grouped
by TA treatment (left) and morphological condition of the parasite (right). (D) Ridgeline
plot of melting curves for candidate RG: Simplified representation of the melting curves
(dF/dT vs. T) obtained from qPCR runs. A single peak was observed for all candidate
genes, confirming specific amplification without contamination.

The amplification efficiency for all reactions performed with each primer set reached
values above 90%. However, genes *rpbii*, *etif4aiii*, *atub*, *ube3a*, and *cypa* were excluded
from downstream analysis because their efficiency values were above 100%, which could
lead to inaccurate measurements (Table 1).

Table 1. Primer sequences and RT-qPCR amplification features of 17 candidate RG and both
target genes.

Gene	Wormbase accession no.	Primer sequence 5'→ 3' (forward / reverse)	Product size (bp)	Tm (° C)	%E
atub	TsM_000816800	ACTCCGCGATAATGGGTCTT/ GGAGCTCTGGGAAGGGAATT	183	59.3	133.56
bact1	TsM_001199400	ATCGTAGCACCACCAGAACG/ CCCGATTCGTCGTACTCCTG	1113	60.0	94.13
bact2	TsM_001001400	GTGCGAGACATCAAGGAGAAG/ GGAAGCGTTCATTGCCAACT	142	60.4	98.31
gapdh1	TsM_000056400	TCCAAGAGATGAATGCCAATGC/ CAGAAGGAGCCGAGATGATGA	147	61.1	93.89
gapdh2	TsM_001083000	CAAACTTGCCAAACCTGCCA/ GCGTCGAAAGTGGATGATGC	153	58.7	97.87
pgk1	TsM_000796500	GACACCGCTACTTGTTGTGC/ TCCGAAAGAGCTGATACGCC	125	59.0	105.09
mdh	TsM_000048200	TAAGGTGCTGGTTGTTGGCA/ CGACCTGATAGATGGCACGG	131	59.8	91.18
cox1	TsM_000113400	CCGTTAGGAGGTGGTGATCC/ ACCCATAAAAGCCAAAAGCA	154	60.1	101.69
rpb2	TsM_000754600	CACCACGAAGACACCACAAG/ TAGGAATGGCGTGAGGATTG	149	58.6	95.31
tbp	TsM_000071900	GCCAAAGTGCGAGATGAGAT/ CCCGATTTGTCTGAATCCAA	89	56.8	96.374
eif4a	TsM_000855500	CTGCATTGGTACGCTTCAAC/ AGTGCTGTGACGACCTGATG	111	58.7	116.65
rpl13	TsM_000621100	TCCAATGCTATCGGGTGAAT/ CGCCTCGTGGATCGTATATT	74	56.0	103.09
ef1a	TsM_000233800	TTGCTGCTGGTACTGGTGAG/ CGCTATAATCAACCGCATCC	138	58.6	89.79
cypa	TsM_000676800	TCGCAGTTCTTCATCACCAC/ TCTTGCCACTACTGCTACCG	127	58.1	129.71
ub3a	TsM_000891500	TACGCTTGTGCTGGTGGTAG/ GGGAAGAAGGAGATCCAAGG	87	58.5	93.42
ub2	TsM_001232600	TTGTCCCACTTATCCCAAGG/ ACGGCCTAAGCACACTGAAC	100	59.1	110.49
mapk3	TsM_000447200	TCAAGCCGGACTACTTACCG/ CACGGTTGCTGTACATCTCG	140	58.6	101.56
h2b	TsM_000989300	GCTCCTAAAGTAGTGTCAGGCA/ ATGGCGTAGCTCTCCTTCCT	104	63.2	100.38

| *wnt11a* | TsM_000806100 | GGCAGGCGAGGTTCTGTAAT/G GCACAGTAGCGACCGATTT | 104 | 60.2 | 101.85 |

RT-qPCR performance and specificity and RG expression profile

To determine the dispersion of Cq values for the selected candidate RG under conditions
representing *T. solium* early-adult stages, initial qPCR runs were performed using cDNA
from controls and PRE, EV, and POST parasites. A small Cq value indicated higher gene
expression abundance, and vice versa. Mean Cq values for all RG were below 35 cycles,
suggesting that transcript abundances were not underrepresented.

A comprehensive heat map of the raw Cq values helped identify parasite samples
contributing to variability within their groups. Overall, the samples showed consistent
expression levels. Most candidate genes were upregulated due to culture conditions,
regardless of AT treatment. Notably, the *ube2* gene, encoding a ubiquitin-conjugating
enzyme involved in protein degradation, was specifically upregulated during early culture
only with AT treatment (Fig 2B). Similarly, genes *bact2* (beta-actin isoform 2, a cytoskeletal
protein) and *ef1a* (Eukaryotic Elongation Factor 1-alpha, crucial for protein synthesis)
exhibited distinct AT-dependent differential expression. Interestingly, *gapdh1*, involved in
glycolysis and other cellular processes, showed the lowest expression across all
conditions and treatments (Fig 2C).
236

Comprehensive stability analysis of candidate RG among conditions

RefFinder, a free web tool, combines the results from BestKeeper, geNorm, NormFinder,
and the comparative DeltaCT method, using raw Cq values as input to provide a geometric
mean ranking of RG stability.

Bestkeeper ranks the candidate RG based on two key indicators: (1) the standard
deviation (SD) of the Cq values (genes with SD > 1 are considered unstable), and (2) the
Pearson correlation coefficient (r), comparing each gene's expression to a BestKeeper
index derived from the geometric mean of all other genes. Higher correlation (r values
close to 1) indicates better stability. geNorm calculates a stability score (M) for each gene,
representing the average pairwise variation between that gene and all others being tested.
Genes with the highest M values (least stability) are eliminated in a stepwise manner. A
lower M value indicates higher stability. A cut-off of 0.15 is used to determine if additional
RG are needed. Normfinder ranks RG by calculating their expression stability considering
both intra- and inter-group variation. It identifies the genes with the least variation across
different conditions or treatments. A stability value below 0.15 is generally considered
suitable for normalization. Finally, the deltaCT method compares the relative expression
levels of candidate RG pairs across all samples. Gene pairs with the smallest variability
(smallest difference in their Cq values) are considered more stable.

The BestKeeper analysis identified *gapdh2*, *cox1*, and *tbp* as the most stable genes.
geNorm found *pgk1* and *bact1* to be equally stable, followed by *rpl13* and *mapk3*.
NormFinder ranked *mapk3*, *tbp*, and *pgk1* as the most stable, while the deltaCT method
highlighted *pgk1*, *rpl13*, and *bact1* as the top stable genes (Fig 3, left). The comprehensive
ranking, based on the geometric mean of all analyses, identified *pgk1*, *bact1*, *mapk3*, *tbp*,
rpl13, *cox1*, and *gapdh2* (in descending order) as the most stable genes across all

comparisons (Fig 3, right). These genes are essential for key cellular processes, including
metabolism, structural integrity, and signal transduction.

Figure 3. Comprehensive Gene Stability Ranking by RefFinder. The stability scores for
candidate RG were analyzed using geNorm, NormFinder, BestKeeper, and the deltaCT
method. These results were combined into a comprehensive stability ranking, calculated as
the geometric mean of the scores from all four methods. Lower values indicate higher gene
stability, reflecting reduced variability.

Validation of the top-ranked RG

To validate the stability of the selected RG, *h2b* and *wnt11a* were used as target genes
under the PRE, EV, and POST early-adult conditions. As previously reported in other
parasitic tapeworms, these genes are expected to be upregulated during scolex
evagination, as both cell proliferation and strobilation are inherent to this morphogenic
process.

After calculating the fold-change values using the $2^{-\Delta\Delta Cq}$ method, the proliferative gene *h2b*
showed a 2- to 6-fold increase in expression throughout the early-adult stages compared
to the uncultured control. Interestingly, when *h2b* expression was normalized using the
mapk3 and *tbp* RG, the uncultured control groups formed a more consistent cluster with
reduced variability. Similarly, *wnt11a* expression was upregulated in cultured parasites,
showing a 5- to 10-fold increase when using the different selected RG. The fold-change
increase remained within a consistent range for both *h2b* (no values exceeding an 8-fold
increase) and *wnt11a* (no values exceeding a 30-fold increase), suggesting that the
selected RG exhibit a similar expression pattern, supporting their suitability for
normalization (Fig 4A).

**Figure 4. Relative Quantification of *h2b* and *wnt11a* expression using the most and
least stable reference genes.** The relative expression of *h2b* and *wnt11a*, involved in cell
proliferation and strobilation, was calculated using the $2^{-\Delta\Delta Cq}$ method. Gene expression was
normalized with (A) the top-ranked and (B) the lowest-ranked reference genes identified
through the stability analysis.

In contrast, when the least stable candidates (*bact2*, *ub2*, *gapdh1*, and *ef1a*) were used for
normalization, the expression levels of *h2b* and *wnt11a* varied significantly across
treatments. The *h2b* fold-change fluctuated widely, ranging from 20- to 1000-fold,
depending on the gene used as reference. Similarly, *wnt11a* showed highly variable fold-
changes, ranging from 10- to 6000-fold, indicating that these unstable genes lead to
inconsistent normalization.

Notably, a common feature of the fold-change values obtained for *h2b* and *wnt11a* across
the different stages is their inconsistency, displaying a bimodal distribution in gene
expression. The values range from below the detection limit (<1, indicating
downregulation) to significant increases in expression, as seen when *bact1* or *ef1a* are
used for normalization (Fig 4B).

DISCUSSION

Like in other neglected infectious diseases, the biology of *T. solium* is largely unexplored.
This field holds crucial information for better understanding interactions with the hosts,
developmental changes through life cycles, and the associated disease endemicity. In the
absence of stage-specific transcriptomes, single-gene expression analysis is a powerful
approach to exploring the roles of genes of interest under well-defined biological
conditions. The controlled in vitro evagination of cysts in the presence ofTA provides a
valuable model for studying early-adult development. The normalization of gene
expression data often relies on canonical RG that have not been properly validated for this
organism or its (yet not described) developmental stages.

In our analysis, we selected a group of genes involved in essential eukaryotic
housekeeping processes. Some of them have previously been analyzed for stability in
closely related cestodes like *Echinococcus sp.* They are also widely used for normalization
in relative quantification studies across various organisms. The comprehensive ranking of
gene stability highlighted *pgk1*, *bact1*, *mapk3*, *tbp, rpl13, cox1*, and *gapdh2*.

PGK1 is a glycolytic enzyme. Due to the stability of glycolytic processes across many cell
types, it is frequently selected as a housekeeping gene for normalization in qPCR and
gene expression analyses in cell lines or animal tissues[37]. However, no similar analyses
have been performed in free-living or parasitic worms, suggesting that PGK1 expression
may be upregulated (and maintained) in glucose-rich environments, such as those
provided by culture media.

While genes involved in signal transduction, such as *mapk3*, are typically considered
highly variable, a small study on *H. microstoma* found that cAMP, a key second
messenger in signaling pathways, was consistently expressed along the anteroposterior
gradient of the adult body[38]. This suggests that the cellular signaling pathways may be
stably activated in response to the culture conditions.

Although *gapdh* is commonly used to normalize gene expression, it plays several roles
beyond its well-known function in the glycolytic pathway. For instance, it helps maintain
cellular iron homeostasis[39] and acts as a chaperone[40]. Furthermore, its expression levels
in human cells are not stable[41] and show greater variability across different tissues
compared to other genes[42,43]. Consequently, previous studies have raised concerns about
using GAPDH as a reference gene in qPCR experiments[44].

Moreover, gapdh has undergone gene duplication events throughout evolution, resulting in
different isoforms that may have distinct roles. As a result, the expression of gapdh
isoforms can vary depending on the tissue[45].

Our comprehensive analysis included the isoforms *gapdh1* and *gapdh2*, but neither
demonstrated high stability. Although the Bestkeeper algorithm identified *gapdh2* as the
most stable candidate, likely due to its low SD across all samples and conditions, other
algorithms ranked gapdh2 lower. Additionally, gapdh exhibited the highest raw Cq values
among the seventeen initial reference genes, suggesting low expression levels in parasite
tissues, making it unsuitable for relative expression analyses. Notably, when gapdh1 was
used to normalize the expression of h2b and wnt11a, the resulting fold change values

were significantly overestimated compared to those obtained with the top-ranked reference
genes.

Some of the top-ranked genes in our analysis, such as *tbp*, *rpl13*, and *bact1*, were also
previously reported as among the most stable in a similar study on *Echinococcus sp.*
protoscoleces that were activated in vitro with pepsin[26]. This suggests that, despite being
based on a limited number of representative elements, the expression profiles may share
common features across species within the phylum Platyhelminthes.

In contrast, one of our top-ranked genes, *tbp*, was identified in a recent study focused on
reference gene selection in the free-living echiuran worm *Urechis unicinctus*. Based on
transcriptome data, this study suggested that *tbp* is among the least stable genes[46].
However, the genera *Urechis* and *Taenia* occupy distant evolutionary branches within the
animal kingdom. Echiurans belong to the Phyllum Annelida, significantly divergent from the
Platyhelminths.

Finally, our in vitro design provides a versatile tool for exploring and analyzing the
expression of genes of interest throughout the early adult stages during chemically
induced evagination of *T. solium*. However, a significant limitation is that our
determinations were done on whole-cyst RNA preparations. As seen in other parasitic
cestodes, several genes exhibit discrete expression patterns along the adult worm's body.
Techniques such as in situ hybridization have revealed the presence of regional
expression gradients, which may be restricted to or extend across anatomical regions like
the scolex, the "transition zone" between the neck and the nascent strobila, as well as the
immature and mature proglottids[34,35]. Therefore, depending on the specific goals, a more
refined set of criteria should be established, such as preparing RNA from carefully
selected anatomical regions, to improve the accuracy of the analysis. Nevertheless, this
would be difficult on cysts with invaginated scolices (our PRE parasites).

Taken together, the results of this work validated that *pgk1*, *bact1*, *mapk3*, *tbp*, *rpl13*, and
cox1 could be suitable RG for normalizing single-gene relative expression qRT-PCR data
in the context of *T. solium* cyst evagination. This study provides a basis for future studies
of gene function in *T. solium* early-adult development.

ACKNOWLEDGMENTS

We thank Luz Moyano and Ricardo Gamboa (Global Health Tumbes, UPCH) for providing viable *T. solium* cysts; David Duran (Instituto de Medicina Tropical "Alexander von Humboldt", UPCH) for his kind assistance with gene expression equipment; Vanessa Arevalo (Laboratorio de Ecotoxicología, UPCH) for her assistance with macro photography of cultured parasites, and Francisco Villafuerte, Daniela Bermúdez and Rómulo J. Figueroa (Laboratorio de Fisiología Comparada, UPCH) for sharing their facilities for parasite culture. RGL and JBLT were supported by a training grant from NIH-Fogarty, USA (2D43TW007120-11A1). DC, JM, and VV were supported by a grant from the Peruvian Programa Nacional de Investigación Científica y Estudios Avanzados, Prociencia (PE501079376-2022), which also funded the completion of this work.

AUTHOR CONTRIBUTIONS

RGL and JBLT conceptualized and designed this work. DC, JM, VV, JBLT, and SDA performed experiments. RGL, DC, and JBLT performed all data analysis. RGL wrote the manuscript with input from all authors, and CGG finished the critical revision of the article.

DECLARATION OF INTERESTS

The authors declare that the research was conducted without any commercial or financial relationships that could be construed as a potential conflict of interest.

REFERENCES

1. World Health Organization. Ending the neglect to attain the Sustainable Development Goals: A road map for neglected tropical diseases 2021–2030. (2020). Available at: https://www.who.int/publications/i/item/9789240010352. (Accessed: 15th February 2021)

2. World Health Organization. WHO estimates of the global burden of foodborne diseases. (2015). Available at: https://www.who.int/foodsafety/publications/foodborne_disease/fergreport/en/. (Accessed: 15th February 2021)

3. World Health Organization. Taeniasis/cysticercosis. (2022). Available at: https://www.who.int/news-room/fact-sheets/detail/taeniasis-cysticercosis. (Accessed: 15th February 2021)

4. Garcia, H. H., Gonzalez, A. E. & Gilman, R. H. Taenia solium cysticercosis and its impact in neurological disease. *Clinical Microbiology Reviews* **33**, (2020).

5. Flisser, A. *et al.* Taenia solium: Current understanding of laboratory animal models of taeniosis. *Parasitology* **137**, 347–357 (2010).

6. Arora, N. *et al.* Recent advancements and new perspectives in animal models for Neurocysticercosis immunopathogenesis. *Parasite Immunology* **39**, (2017).

7. Palma, S. *et al.* In vitro model of postoncosphere development, and in vivo infection abilities of Taenia solium and Taenia saginata. *PLoS Negl. Trop. Dis.* **13**, (2018).

8. Ávila, G. *et al.* Laboratory animal models for human Taenia solium. in *Parasitology International* **55**, (Parasitol Int, 2006).

9. Koziol, U., Domínguez, M. F., Marín, M., Kun, A. & Castillo, E. Stem cell proliferation during in vitro development of the model cestode Mesocestoides corti from larva to adult worm. *Front. Zool.* **7**, 22 (2010).

10. Rabiela, M. T., Hornelas, Y., Garcia-Allan, C., Rodriguez-del-Rosal, E. & Flisser, A. Evagination of Taenia solium cysticerci: A histologic and electron microscopy study. *Arch. Med. Res.* **31**, 605–607 (2000).

11. Paredes, A. *et al.* In vitro analysis of albendazole sulfoxide enantiomers shows that (+)-(R)-albendazole sulfoxide is the active enantiomer against Taenia solium. *Antimicrob. Agents Chemother.* **57**, 944–949 (2013).

12. Tsai, I. J. *et al.* The genomes of four tapeworm species reveal adaptations to parasitism. *Nature* **496**, 57–63 (2013).

13. Almeida, C. R. *et al.* Transcriptome analysis of Taenia solium cysticerci using open reading frame ESTs (ORESTES). *Parasites and Vectors* **2**, (2009).

14. Lundström, J., Salazar-Anton, F., Sherwood, E., Andersson, B. & Lindh, J. Analyses of an expressed sequence tag library from Taenia solium, cysticerca. *PLoS Negl. Trop. Dis.* **4**, 1–8 (2010).

15. Orrego, M. A. *et al.* Transcriptomic analysis of subarachnoid cysts of Taenia solium reveals mechanisms for uncontrolled proliferation and adaptations to the microenvironment. *Sci. Rep.* **14**(1):11833 (2024).

16. Paludo, G. P. *et al.* Cestode strobilation: Prediction of developmental genes and pathways. *BMC Genomics* **21**, (2020).

17. Yang, D. *et al.* Annotation of the transcriptome from taenia pisiformis and its comparative analysis with three taeniidae species. *PLoS One* **7**, (2012).

18. Parkinson, J. *et al.* Correction: A Transcriptomic Analysis of Echinococcus granulosus Larval Stages: Implications for Parasite Biology and Host Adaptation. *PLoS Negl. Trop. Dis.* **6**, (2012).

19. Wu, X. *et al.* Detailed Transcriptome Description of the Neglected Cestode Taenia multiceps. *PLoS One* **7**, (2012).

20. Zhang, S. Comparative transcriptomic analysis of the larval and adult stages of Taenia pisiformis. *Genes (Basel).* **10**, (2019).

21. García-Montoya, G. M. *et al.* Transcriptome profiling of the cysticercus stage of the laboratory model Taenia crassiceps, strain ORF. *Acta Trop.* **154**, 50–62 (2016).

22. Olson, P. D. *et al.* Genome-wide transcriptome profiling and spatial expression analyses identify signals and switches of development in tapeworms. *Evodevo* **9**, (2018).

23. Basika, T. *et al.* Transcriptomic profile of two developmental stages of the cestode parasite Mesocestoides corti. *Mol. Biochem. Parasitol.* **229**, 35–46 (2019).

24. Huggett, J., Dheda, K., Bustin, S. & Zumla, A. Real-time RT-PCR normalisation; strategies and considerations. *Genes and Immunity* **6**, 279–284 (2005).

25. Pouchkina-Stantcheva, N. N., Cunningham, L. J. & Olson, P. D. Spatial and temporal consistency of putative reference genes for real-time PCR in a model tapeworm. *Mol.*

453   *Biochem. Parasitol.* **180**, 120–122 (2011).

454   26.   Espínola, S. M., Ferreira, H. B. & Zaha, A. Validation of suitable reference genes for
455   expression normalization in Echinococcus spp. larval stages. *PLoS One* **9**, (2014).

456   27.   Hou, J. *et al.* Sequence analysis and molecular characterization of Wnt4 gene in
457   metacestodes of Taenia solium. *Korean J. Parasitol.* **52**, 163–168 (2014).

458   28.   Untergasser, A. *et al.* Primer3-new capabilities and interfaces. *Nucleic Acids Res.* **40**,
459   (2012).

460   29.   Bustin, S. A. *et al.* The MIQE guidelines: minimum information for publication of
461   quantitative real-time PCR experiments. *Clin. Chem.* **55**, 611–22 (2009).

462   30.   Vandesompele, J. *et al.* Accurate normalization of real-time quantitative RT-PCR data
463   by geometric averaging of multiple internal control genes. *Genome Biol.* **3**,
464   RESEARCH0034 (2002).

465   31.   Andersen, C. L., Jensen, J. L. & Ørntoft, T. F. Normalization of real-time quantitative
466   reverse transcription-PCR data: A model-based variance estimation approach to
467   identify genes suited for normalization, applied to bladder and colon cancer data sets.
468   *Cancer Res.* **64**, 5245–5250 (2004).

469   32.   Pfaffl, M. W., Tichopad, A., Prgomet, C. & Neuvians, T. P. Determination of stable
470   housekeeping genes, differentially regulated target genes and sample integrity:
471   BestKeeper - Excel-based tool using pair-wise correlations. *Biotechnol Lett.* **26**, 509–
472   515 (2004).

473   33.   Silver, N., Best, S., Jiang, J. & Thein, S. L. Selection of housekeeping genes
474   for gene expression studies in human reticulocytes using real-time PCR. *BMC Mol*
475   *Biol.* **7**, 33 (2006).

476   34.   Jarero, F. *et al.* Muscular remodeling and anteroposterior patterning during
477   tapeworm segmentation. *Dev Dyn.* 1–26 (2024)

478   35.   Rozario, T., Quinn, E.B., Wang, J., Davis, R.E. & Newmark, P.A. Region-
479   specific regulation of stem cell-driven regeneration in tapeworms. *Elife.* **8**:e48958
480   (2019).

481   36.   Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-
482   time quantitative PCR and the $2\text{-}\Delta\Delta CT$ method. *Methods.* **25**, 402–408 (2001).

483   37.   Schwarz, A. P. *et al.* Reference gene expression stability within the rat brain under
484   mild intermittent ketosis induced by supplementation with medium-chain triglycerides.
485   *PloS one*, **18**(2), e0273224 (2023).

486   38.   Pouchkina-Stantcheva, N. N., Cunningham, L. J., & Olson, P. D. Spatial and temporal
487   consistency of putative reference genes for real-time PCR in a model tapeworm. *Mol*
488   *Biochem Parasitol.* 180(2), 120–122 (2011).

489   39.   Boradia, V. M., Raje, M., & Raje, C. I Protein moonlighting in iron metabolism:
490   glyceraldehyde-3-phosphate dehydrogenase (GAPDH). *Biochem Soc Trans.* **42**(6),
491   1796–1801 (2014).

492   40.   Sweeny, E. A. *et al.* Glyceraldehyde-3-phosphate dehydrogenase is a chaperone that
493   allocates labile heme in cells. *J Biol Chem.* 293(37), 14557–14568 (2018).

41. Zhu, G., Chang, Y., Zuo, J., Dong, X., Zhang, M., Hu, G., & Fang, F. Fudenine, a C-
terminal truncated rat homologue of mouse prominin, is blood glucose-regulated and
can up-regulate the expression of GAPDH. *Biochem Biophys Res Commun.* **281**(4),
951–956 (2001).

42. Barber, R. D., Harmer, D. W., Coleman, R. A., & Clark, B. J. GAPDH as a
housekeeping gene: analysis of GAPDH mRNA expression in a panel of 72 human
tissues. *Physiol Genomics.* **21**(3), 389–395 (2005).

43. Radonić, A., Thulke, S., Mackay, I. M., Landt, O., Siegert, W., & Nitsche, A. Guideline
to reference gene selection for quantitative real-time PCR. *Biochem Biophys Res
Commun.* **313**(4), 856–862 (2004).

44. Ke, L. D., Chen, Z., & Yung, W. K. A reliability test of standard-based quantitative
PCR: exogenous vs endogenous standards. *Mol Cell Probes.* **14**(2), 127–135 (2000).

45. Manchado, M., Infante, C., Asensio, E., & Cañavate, J. P. Differential gene
expression and dependence on thyroid hormones of two glyceraldehyde-3-phosphate
dehydrogenases in the flatfish Senegalese sole (Solea senegalensis Kaup). *Gene.*
400(1-2), 1–8 (2007).

46. Chen, J., Wang, Y., Yang, Z., Liu, D., Jin, Y., Li, X., Deng, Y., Wang, B., Zhang, Z., &
Ma, Y. Identification and validation of the reference genes in the echiuran worm
Urechis unicinctus based on transcriptome data. *BMC genomics.* **24**(1), 248 (2023).

Figure 1

Fig 1. Taurocholic acid (TA) induces in vitro evagination of T. solium cysts. (A) Overview of the study design and analysis workflow. (B) Quantification of the percentage of cyst evagination over 120 hours. Each point represents the mean of three independent experiments, with error bars showing the maximum and minimum values. (C) Representative macroscopic images of non-evaginated PRE (left) and POST (right) parasites after 5 days of culture following 0.1% TA induction. Scale bar: 5 mm. These samples were used for subsequent single-gene expression analysis comparisons.

Figure 2

Fig 2. Comprehensive Analysis of qPCR Performance for Candidate Reference Genes. (A) Primer pair specificity for candidate reference genes: End-point PCR products for all candidate RG were observed on gel electrophoresis and arranged according to product size. MW: 100bp DNA Molecular Weight Ladder. (B) Raw Cq values across parasite samples: Comparison of raw Cq values from qPCR amplification across all parasite samples, showing gene expression for each candidate RG. (C) Cq Value distribution based on treatment and morphology: The distribution of Cq values for each candidate RG, grouped by TA treatment (left) and morphological condition of the parasite (right). (D) Ridgeline plot of melting curves for candidate RG: Simplified representation of the melting curves (dF/dT vs. T) obtained from qPCR runs. A single peak was observed for all candidate genes, confirming specific amplification without contamination.

Figure 3

Figure 3. Comprehensive Gene Stability Ranking by RefFinder. The stability scores for candidate RG were analyzed using geNorm, NormFinder, BestKeeper, and the deltaCT method. These results were combined into a comprehensive stability ranking, calculated as the geometric mean of the scores from all four methods. Lower values indicate higher gene stability, reflecting reduced variability.

Figure 4

Figure 4. Relative Quantification of h2b and wnt11a expression using the most and least stable reference genes. The relative expression of h2b and wnt11a, involved in cell proliferation and strobilation, was calculated using the 2−ΔΔCq method. Gene expression was normalized with (A) the top-ranked and (B) the lowest-ranked reference genes identified through the stability analysis.