Two mosquito odorant receptors with reciprocal specificity mediated by a single amino acid residue

Flávia P. Franco[1], Pingxi Xu[1], Brandon J. Harris[2], Vladimir Yarov-Yarovoy[2,3], and Walter S. Leal[1]*

[1]Department of Molecular and Cellular Biology, [2]Department of Physiology and Membrane Biology, and [3]Department of Anesthesiology and Pain Medicine, University of California, Davis, Davis, CA 95616 USA

*Walter S. Leal, wsleal@ucdavis.edu

Keywords: *Culex quinquefasciatus, CquiOR2, CquiOR10, RosettaFold, RosettaLigand*

Dedicated to the late Professor James H. Tumlinson III

ABSTRACT

The southern house mosquito, *Culex quinquefasciatus,* utilizes two odorant receptors, CquiOR10 and CquiOR2, narrowly tuned to oviposition attractants and well conserved among mosquito species. They detect skatole and indole, respectively, with reciprocal specificity. We swapped the seven transmembrane (TM) domains of CquiOR10 and CquiOR2 and identified TM2 as a specificity determinant. With additional mutations, we showed that CquiOR10A73L behaved like CquiOR2. Conversely, CquiOR2L74A recapitulated CquiOR10 specificity. Next, we generated structural models of CquiOR10 and CquiOR10A73L using RoseTTAFold and AlphaFold and docked skatole and indole using RosettaLigand. These modeling studies suggested space-filling constraints around A73. Consistent with this hypothesis, CquiOR10 mutants with a bulkier residue (Ile, Val) were insensitive to skatole and indole, whereas CquiOR10A73G retained the specificity to skatole and showed a more robust response than the wildtype receptor CquiOR10. On the other hand, Leu to Gly mutation of the indole receptor CquiOR2 reverted the specificity to skatole. Lastly, CquiOR10A73L, CquiOR2, and CquiOR2L74I were insensitive to 3-ethylindole, whereas CquiOR2L74A and CquiOR2L74G gained activity. Additionally, CquiOR10A73G gave more robust responses to 3-ethylindole than CquiOR10. Thus, we suggest the specificity of these receptors is mediated by a single amino acid substitution, leading to finely tuned volumetric space to accommodate specific oviposition attractants.

INTRODUCTION

Insects perceive the world with a sophisticated olfactory system essential for survival and reproduction. Their antennae are biosensors *par excellence,* allowing detection of a plethora of compounds, some with extraordinary sensitivity and selectivity (Kaissling, 2014). The insect olfactory system is comprised mainly of odorant-binding proteins, odorant-degrading enzymes, ionotropic receptors, and the ultimate gatekeepers of selectivity (Leal, 2013, 2016; Leal, 2020) – the odorant receptors (ORs) (Clyne et al., 1999; Gao & Chess, 1999; Vosshall et al., 1999). The ORs are the binding units in functional heteromeric cation channels (Neuhaus et al., 2005) formed with an odorant receptor coreceptor (Orco) (Larsson et al., 2004). Unlike mammalian olfactory receptors, insect ORs and Orco have inverse topologies compared to G-protein coupled receptors (GPCRs), with a cytosolic N-terminus and an extracellular C-terminus (Benton et al., 2006; Lundin et al., 2007). One of the major breakthroughs in the fields of insect olfaction in the last two decades since the discovery of ORs (Leal, 2020) was the determination of the cryo-electron microscopy (cryo-EM) structure of the Orco homomer from the parasitic fig wasp*, Apocrypta bakeri,* AbakOrco (Butterwick et al., 2018). Subsequently, the structure for a promiscuous OR from the evolutionarily primitive (Apterygota, wingless) jumping bristletail, *Machilis harabi*, MhOR5, was solved (Del Marmol et al., 2021). Although structures of ORs from winged insects (Pterygota) have not been solved to date, amino acid residues critical for OR specificity have been reported (Auer et al., 2020; Cao et al., 2021; Hughes et al., 2014; Leary et al., 2012; Pellegrino et al., 2011; Yang et al., 2017; Yuvaraj et al., 2021). These studies focused on one-way alteration of specificity, but did not examine how an insect detects two odorants with reverse specicity.

The genome of the southern house mosquito, *Culex quinquefasciatus* (Arensburger et al., 2010)*,* has the most extensive repertoire of *OR* genes (Leal et al., 2013) among mosquito species. However, the number of odorants in the

environment far exceeds the number of OR genes, even in *Cx. quinquefasciatus*.

Therefore, it is unsurprising that many insect odorant receptors are promiscuous

(Leal, 2013; Pask, 2020). However, ORs detecting behaviorally critical

compounds may be narrowly tuned to pheromones and other semiochemicals

(Hughes et al., 2010; Nakagawa et al., 2005; Stensmyr et al., 2012).

Odorant receptors narrowly tuned to 3-methylindole (=skatole) and indole have

been found in mosquito species in the subfamilies Culicinae and Anophelinae.

Specifically, OR10 and OR2 have been de-orphanized in the southern house

mosquito (Hughes et al., 2010; Pelletier et al., 2010)*,* the yellow fever mosquito

(Bohbot et al., 2011), and the malaria mosquito (Carey et al., 2010; Wang et al.,

2010)*.* Recently, these so-called indolergic receptors (Bohbot & Pitts, 2015) have

also been found in the housefly, *Musca domestica* (Pitts et al., 2021)*.*

Remarkably, in all these species, OR10s are narrowly tuned to skatole and

respond with lower sensitivity to indole. In contrast, indole is the most potent

ligand for OR2s, which give lower responses to skatole. In *Cx. quinquefasciatus,*

these receptor genes were initially named *CquiOR10* and *CquiOR2*, respectively,

renamed *CquiOR21* and *CquiOR121*, but a proposition to restore the original

names is under consideration. Here, we refer to these receptors from *Cx.*

quinquefasciatus as CquiOR10 (VectorBase and GenBank I.D.s, CPIJ002479

and GU945397, respectively) and CquiOR2 (CPIJ014392, GU945396.1). These

oviposition-attractant-detecting receptors in *Cx. quinquefasciatus*, CquiOR10 and

CquiOR2 provide a suitable model to identify specificity determinants and

examine the long-standing question of reverse specificity.

To identify the specificity determinants of these receptors, we swapped

transmembrane (TM) domains and tested the chimeric receptors using the

Xenopus oocyte recording system. Given that CquiOR10 is the second most

sensitive *Cx. quinquefasciatus* OR (second only to CquiOR36 (Choo et al.,

2018)), we replaced the predicted (Reynolds et al., 2008) transmembrane

domains from CquiOR2 into CquiOR10 and measured the specificity of the

chimeric receptors. With this approach, we identified TM2 as a specificity

determinant. Next, we tested mutations of chimeric CquiOR10 and identified a

single amino acid residue in TM2 (Ala-73) that determines the receptor's

specificity. We then directly mutated the wildtype receptor and observed that

CquiOR10A73L is specific to indole as CquiOR2. Additionally, CquiOR2L74A

emulated the response profile of CquiOR10. To better understand the structural

basis of this single-point specificity determinant, we generated structural models

of CquiOR10 and CquiOR10A73L using RoseTTAFold (Baek et al., 2021) and

AlphaFold (Jumper et al., 2021). We identified the binding poses for skatole and

indole using RosettaLigand molecular docking and observed a finely tuned

volumetric space to accommodate specific oviposition attractants.

113
114
RESULTS

Chimeric OR with reversed specificity. We envisioned that studying a pair of

odorant receptors with reverse specificities, like CquiOR10 and CquiOR2, could

lead us to specificity determinants, which may have implications for the evolution

of other insect behaviors. CquiOR10 is activated by the oviposition attractant

skatole (Blackwell et al., 1993; Mboera et al., 2000; Millar et al., 1992) with high

specificity (*Figure 1A*), whereas CquiOR2 is specific to indole (*Figure 1B*). Our

approach was designed to swap TM domains using the more sensitive receptor,

CquiOR10, as the acceptor. Specifically, we generated chimeric receptors by

replacing CquiOR10 TM domains with related domains from CquiOR2 (*Figure*

1—figure supplement 1; see Appendix 1). We referred to these chimeric

receptors as CquiOR10^{Mx}, where Mx refers to TMx from CquiOR2. Then, we

performed functional assays of these CquiOR10^{Mx} receptors using the *Xenopus*

oocyte recording system. This long-term project could require as many as 127

possible chimeric receptors. We optimized the workflow to reduce the number of

tested mutants before finding a chimeric receptor with reversed specificity.

Figure 1. Concentration-response analysis for activation of wildtype ORs by

skatole and indole. (A) CquiOR10 and (B) CquiOR2. Lines were obtained with

nonlinear fit. Bars represent SEM. *n* = 4-5.

Figure supplement 1. Sequence alignment of CquiOR10 and CquiOR2.

Figure supplement 2. Concentration-response analysis for wildtype and

chimeric ORs.

Figure supplement 3. Overlap of CquiOR10 predicted TM domains and

AbakOrco cryo-EM structure.

We tested 36 chimeric receptors (see *Appendix 1* and *Figure 1—figure*

supplement 2). Fourteen chimeric receptors did not respond to skatole or indole,

and 21 receptors retained the specificity to skatole (*Extended—Table 1* and

Figure 1—figure supplement 2). Lastly, CquiOR10^{M2,7}/CquiOrco-expressing

oocytes responded to both skatole and indole with a reverse profile (*Figure 2A*).

This dataset shows that CquiOR10^{M2,7} emulated the profile of the indole receptor
CquiOR2 (*Figure 1B*).
150

151
Figure 2. Concentration-response curves obtained with chimeric ORs stimulated
with skatole and indole. (A) CquiOR10^{M2,7}; (B) CquiOR10^{M2,5,6,7}; (C)
CquiOR10^{M2,5,6,7}_Outer; (D) CquiOR10^{M2,5,6,7}_Mid;Inner; (E)
CquiOR10^{M2,5,6,7}_Inner; (F) CquiOR10^{M2,5,6,7}T78I; (G) CquiOR10^{M2,5,6,7}L73A; (H)
CquiOR10^{M7}A73L; (I) CquiOR10A73L; (J) CquiOR2L74A. Lines were obtained

with nonlinear fit. Bars represent SEM. The number of replicates (*n*) were 7, 4, 5,

5, 4, 3, 9, 7, 6, and 5, respectively.

Figure supplement 1. Partial sequences of CquiOR10 and chimeric ORs

highlighting the transmembrane domain-2 (TM2).

Figure supplement 2. Schematic view of the workflow.

A single-point mutation that reverses the specificity of the skatole and

indole receptors. We proceeded to identify the amino acid residues in the

swapped domains of CquiOR10^{M2,7}, directly affecting the specificity of the

chimeric and wildtype receptors. Given the observation that, by and large,

chimeric receptors with TM5 and TM6 from CquiOR2 gave stronger responses

(*Figure 1—figure supplement 2*), we asked whether CquiOR10^{M2,7} responses

with these two additional TM domains swapped would give more robust

responses while keeping the same specificity to indole.

CquiOR10^{M2,5,6,7}/CquiOrco-expressing oocytes were indeed more sensitive while

maintaining the selectivity to indole (*Figure 2B*). We then used CquiOR10^{M2,5,6,7}

and designed various mutants to rescue single or multiple residues in TM2 at a

time (*Figure 2—figure supplement 1*). We focused on TM2 because swapping

TM7 did not affect the specificity of the receptor (*Figure 1—figure supplement*

2E). We aimed to identify mutations in the chimeric receptor that reverts the

specificity to skatole as in the wildtype receptor. We divided TM2 into outer,

middle, and inner segments based on the topology predicted by OCTOPUS

(Viklund & Elofsson, 2008). It has been postulated that the extracellular halves of

TM domains form an odorant-binding pocket (Guo & Kim, 2010), thus we first

examined a mutant (CquiOR10^{M2,5,6,7}_Outer) having the residues in the outer

segment at 59, 60, 63, and 64 restored to Glu, Val, Asn, and Ala, respectively, as

in the wildtype receptor (*Figure 2—figure supplement 1*).

CquiOR10^{M2,5,6,7}_Outer/CquiOrco-expressing oocytes retained the specificity of

CquiOR10^{M2,5,6,7} with a more robust response to indole than skatole (*Figure 2C*).

These findings indicated that the residues in the outer segment of TM2 are not

specificity determinants. After that, we tested a chimeric receptor with the

residues in the middle and inner part of the TM2 domain rescued to match those

in the wildtype receptor (*Figure 2—figure supplement 1*).

CquiOR10^{M2,5,6,7}_Mid;Inner restored the skatole-specific profile of CquiOR10

(*Figure 2D*), thus suggesting that amino acid residues in these segments are

specificity determinants. Then, we tested CquiOR10^{M2,5,6,7}_Inner

(=CquiOR10^{M2,5,6,7}L73A;T78I), which had only residues at 73 and 78 rescued to

the wildtype Ala and Ile, respectively. Oocytes co-expressing

CquiOR10^{M2,5,6,7}_Inner and CquiOrco reverted the specificity to skatole (*Figure*

2E). As a result, we concluded that residues in the predicted inner part of TM2

are critical for the chimeric receptor's specificity. We further probed the chimeric

receptor CquiOR10^{M2,5,6,7} with single-point mutations to identify the residue(s)

determining specificity. CquiOR10^{M2,5,6,7}T78I/CquiOrco-expressing oocytes

showed the same specificity as the chimeric receptor CquiOR10^{M2,5,6,7} (*Figure*

2F). Specifically, CquiOR10^{M2,5,6,7}T78I gave a more robust response to indole

than skatole, suggesting that rescuing the residue at 78 did not affect

CquiOR10^{M2,5,6,7} specificity. By contrast, CquiOR10^{M2,5,6,7}L73A/CquiOrco-

expressing oocytes reverted the specificity to skatole (*Figure 2G*), thus behaving

like the wildtype receptor CquiOR10 (*Figure 1A*). To further examine the role of

Ala-73 as a specificity determinant residue, we obtained a single-point mutation

of CquiOR10^{M7}, which is specific to skatole (*Figure 1—figure supplement 2E*).

The responses recorded from CquiOR10^{M7}A73L/CquiOrco-expressing oocytes

showed a reverse, indole-specific profile (*Figure 2H*), similar to the CquiOR2

profile (*Figure 1B*). Having identified a single amino acid residue in the chimeric

receptor that switches the skatole/indole specificity, we tested the effect of single

point mutation on the specificity of the wildtype receptor CquiOR10 (EC50:

skatole, 3.6 µM; indole 29.9 µM). CquiOR10A73L showed a reverse specificity,

with dose-dependent responses to indole (*Figure 2I*) (EC50: indole, 3.4 µM;

skatole 53.7 µM). Collectively, these findings suggest that a single amino acid

residue in CquiOR10 determines the specificity of this receptor. Additionally, we

obtained an equivalent single-point mutation in the indole-specific CquiOR2

(*Figure 1B*) (EC50: indole, 7.7 µM; skatole 16.4 µM). Thus,

CquiOR2L74A/CquiOrco-expressing oocytes gave robust and specific responses

to skatole (Figure 2J) (EC50: skatole, 8.5 µM; indole 27.6 µM).

Lastly, we recorded the response of these ORs to other phenolic ligands that

activate indolic receptors, albeit generating small currents. While CquiOR10 and

CquiOR2 responded to phenol, 3,5-dimethylphenol activated only CquiOR10

(*Figure 3*). A single-point mutation in CquiOR10 rendered the chimeric receptor

insensitive to 3,5-dimethylphenol. By contrast, an equivalent mutation in

CquiOR2 recapitulated the profile of CquiOR10 (*Figure 3*).

Figure 3. Quantification of wildtype and chimeric receptors to phenol and 2,3-

dimethylphenol. Each receptor was co-expressed with CquiORco in *Xenopus*

oocytes and stimulated with the phenolic compounds at 1 mM. Bars represent

SEM. *n* = 3-5.

Additionally, we recorded responses elicited by methylindoles. Specifically, we

challenged oocytes with 1-methylindole, 2-methylindole, 4-methylindole, 5-

methylindole, 6-methylindole, and 7-methylindole. In these analyses, we did not

stimulate the oocyte preparations with 3-methylindole to avoid possible

desensitization. CquiOR10/CquiOrco-expressing oocytes elicited stronger

responses when challenged with 1-methylindole and 5-methylindole than when

stimulated with the other methylindoles (*Figure 4A*). By contrast,

CquiOR2/CquiOrco-expressing oocytes elicited similarly lower responses when

stimulated with methylindoles (*Figure 4B*). CquiOR2L74A with a single-point

mutation to mimic OR10 receptor recapitulated CquiOR10 response profile

(*Figure 4C*). These data suggest that a single amino acid residue determines a

receptor's specificity toward ligands eliciting robust or small responses.

Figure 4. Quantification of responses to methylindoles presented at 100 µM. (A)

CquiOR10/CquiOrco-, (B) CquiOR2/CquiOrco-, and (C)-CquiOR2L74A-

expressing oocytes. Bars represent SEM. *n* = 9-11.

As summarized in a graphical representation (*Figure 2—figure supplement 2*),

these findings demonstrate that these two mosquito odorant receptors,

CquiOR10 and CquiOR2, have reciprocal specificity mediated by a single amino

acid residue, Ala-73 and Leu-74, respectively.

CquiOR10 computational modeling suggests space-filling constraints for

indole-based odorants around A73. To structurally hypothesize the above-

described reciprocal specificity, we generated structural models of CquiOr10,

CquiOR2, CquiOR10A73L and CquiOR2L74A using RoseTTAFold (Baek et al.,

2021) and a structural model of CquiOR10 using AlphaFold (Jumper et al., 2021)

(*Figure 5*).

Figure 5. Structural models of CquiOR10 (A,B), CquiOR10A73L

(C),CquiOR2(D), and CquiOR2L74A(E) with AlphaFold (A) and RoseTTAFold (B-

E) structure prediction methods. Superposition of all RoseTTAFold models (F)

resulted in transmembrane helix root mean square deviation (RMSD) of 0.8 Å

when aligned with RoseTTAFold CquiOR10. (G) The transmembrane helix

RMSD of CquiOR10 RoseTTAFold (rainbow) vs AlphaFold (grey) was 1.7 Å.

Loops were not included in RMSD calculation due to inherent flexibility during

structure prediction.

Figure supplement 1. Representative model of docked eugenol with MhraOR5.

Figure supplement 2. Additional clusters of eugenol docked to MhraOR5.

CquiOR10 models of one homotetramer subunit were generated with AlphaFold

and RoseTTAFold, each producing five models. RoseTTAFold models of

CquiOR10, CquiOR10A73L, CquiOR2, and CquiOR2A73L produced a

transmembrane helix RMSD between α-Carbon atoms less than 1 Å across all

twenty models (five models per odorant receptor); this suggests that the

homologous CquiOR10 and CquiOR2 are structurally similar and that single-

point mutations should not cause great structural deviation from wild type.

Comparing CquiOR10 models, RoseTTAFold and AlphaFold produced a

transmembrane helix RMSD of 1.7 Å, with the transmembrane helices at the

extracellular membrane face having the largest structural deviation. Considering

there is little structural knowledge of insect odorant receptors, their binding

mecanisms, and their conformational changes, we suspected that homologous

odorant receptors would have similar binding modes. Using transmembrane 7b

(TM7b), we superimposed the top ranking CquiOR10 RoseTTAFold and

AlphaFold models with an experimentally resolved structure, *M. hrabei*

(MhraOR5) in complex with eugenol (PDB ID: 7LID, (Del Marmol et al., 2021)) to

identify which of our models resembled an odorant-bound conformation. With this

selection criteria, we proceeded with RoseTTAFold models of the odorant

receptors for Rosetta-based small-molecule docking method RosettaLigand

(Davis & Baker, 2009; DeLuca et al., 2015), as the structural similarity around the

hypothesized binding pocket was greater than the AlphaFold models of the

odorant receptors when compared to the MhraOR5 structure. We chose to select conformationally similar models over modeling and docking an apo structure into a bound conformation because it is a more cautious approach when there is little structural information. We perceived modeling an apo structure into a bound conformation to potentially yield more biologically implausible conformations than docking of a structurally comparative model.

RosettaLigand (37, 38) is a competitive protein docking method (Smith & Meiler, 2020) that has been used previously to predict the binding position of ligands within protein pores (Nguyen et al., 2017; Pressly et al., 2022). To verify that RosettaLigand could effectively sample odorants in receptors homologous to CquiOR10 and CquiOR2, we used the structure of eugenol in complex with the insect odorant receptor OR5 from MhraOR5 (PDB ID:7LID) as a control (Del Marmol et al., 2021). Rosetta protein-ligand docking employs energy-based analyses, such as the interface energy between protein and ligand, to select the representative models (Bidula et al., 2022). With this selection method, the RMSD of our MhraOR5-eugenol models relative to the experimental structure ranged from 0.5 – 5.0 Å (*Extended—Table 2*). RMSD values equal or below 2.0 Å are considered an appropriate range for validation (Park et al., 2021). After using hdbscan cluster analysis (McInnes et al., 2017) to group structurally similar models, the largest cluster had a RMSD of 0.75 Å while the lowest interface-energy model had a RMSD of 2.4 Å. Collectively, these data demonstrate that RosettaLigand paired with the hdbscan clustering method can recapitulate the MhraOR5 structure and blindly select a near-native model, thus is suitable for structural predictions of odorants with CquiOR10A73L and CquiOR2L74A (*Figure 5—figure supplements 1 and 2*).

For each receptor-ligand complex (CquiOR10-skatole, CquiOR10A73L-skatole, CquiOR10-indole, and CquiOR10A73L-indole), we generated 100,000 docking models using RosettaLigand, clustered the 10,000 lowest interface-energy models, and selected the lowest interface-energy model from the 10 largest

clusters, resulting in ten models per receptor-ligand complex from which to draw
structural hypotheses (*Extended—Tables 3 and 4*). Our modeling suggests that
both indole and skatole can readily reorient themselves in a similar pore depth
near residue 73, regardless of mutant or wildtype receptor. Comparing the lowest
interface-scoring model from each receptor-ligand complex, indole and skatole
are positioned in the membrane-embedded pore, flanked by transmembrane
helices S2, S4, S5, and S6, and show positional overlap in both and CquiOR10-
A73L CquiOR10 (Figure 6; *Figure 6—figure supplemens 1 and 2*).

Figure 6. Representative models of docked skatole and indole in complex with CquiOR10 and CquiOR10A73L using RosettaLigand. Each model shown is the lowest interface-energy model from the 10 largest clusters of each docking study. CquiOR10 – skatole (forest green), CquiOR10 – indole (brown), CquiOR10A73L – skatole (light blue), and CquiOR10A73L – indole (purple). Atoms that are not indole/skatole carbon atoms are color-coded by atom type: carbon (grey), nitrogen (dark blue), and oxygen (red). Ala-73 and Leu-73 indicated with space-filling representation.

Figure supplement 1. Clusters from RosettaLigand docking of skatole or indole to CquiOR10 or CquiOR10A73L.

Figure supplement 2. Zoom out view of CquiOR10 or CquiOR10A73L compexed to skatole or indole.

Figure supplement 3. Representative models of docked skatole or indole to CquiOR10 or CquiOR10A73L with PLIC analysis.

Figure supplement 4. Superimposition of all OR-skatole and OR-indole clusters

In most models, skatole and indole form contacts with CquiOR10 and CquiOr10A73L in a similar plane about a center of rotation. These observations are supported by skatole and indole not containing rotatable bonds, thus relying on rigid translational movements and rotation to form favorable contacts with the rotatable and repackable receptor residues. Additionally, our models position indole and skatole within a series of nonpolar, polar-uncharged, and aromatic amino acids. Protein-ligand interaction profiler (PLIP) analysis (Adasme et al., 2021) suggests that the bulk of favorable interactions are nonpolar, occasional hydrogen bonding with the odorant NH group, and occasional parallel pi stacking (*Figure 6—figure supplement 3*).

The ligand-binding pocket is formed by TMs 2, 4, 5, and 6. The most frequent hydrophobic contacts were formed by Tyr-185 (38/40 models; TM4) and Ile-69 (32/40 models; TM2) (*Extended—Tables 5-7*). Phe-261 (TM5) also formed

hydrophobic interactions in all four receptor-ligand complex sets of models, but less frequently (22/40 models). Asn-72 (TM2), Tyr-287 (TM6), and Ile-291 (TM6) formed hydrophobic contacts more frequently with skatole structures compared to indole structures (Asn-72: 12/20 vs 9/20, Tyr-287: 14/20 vs 5/20, Ile-291: 8/20 vs 3/20). Notably, Leu-73 (TM2) in CquiOR10A73L formed hydrophobic contacts with indole (2/10) and skatole (5/10), but Ala-73 (TM2) did not form hydrophobic contacts with indole or skatole. Tyr-138 (TM3) formed hydrophobic contacts with skatole in CquiOR10A73L (2/10), but not in CquiOR10. Gln-294 (TM6) formed hydrophobic contacts with skatole and indole in these structures, except CquiOR10A73L-skatole; however, Gln-294 does form a hydrogen bond in one to two models for each structure. Of note, the range of PLIP-reported hydrophobic contacts among the ten representative models for each structure changed from an average of 6.3±0.5 contacts (Mean±SEM) in CquiOR10 to 5.2±0.3 contacts in CquiOR10A73L (p = 0.0354, Mann-Whitney test). Considering the number of unique residues forming hydrophobic contacts, the averages were 4.9±0.4 and 4.9±0.2 contacts, respectively (p = 0.97, Mann-Whitney test).

There were at most two models in each structure with hydrogen bonds (*Extended—Tables 8-10*). For CquiOR10-skatole, two models formed hydrogen bonds with Gln-294. For CquiOR10A73L-skatole, one model formed hydrogen bonds with Asn-72, and one other with Gln-294 (*Extended—Tables 10*). For CquiOr10-indole, one model formed a hydrogen bond with the backbone of Ile-69, another model with Tyr-138, another with Tyr-287, and a fourth model with Gln-294. For CquiOR10A73L-indole, only one model formed a hydrogen bond at Gln-294.

Parallel pi-stacking interactions (*Extended—Tables 11-13*) were identified for CquiOR10A73L-skatole, CquiOR10-indole, and CquiOR10A73L-indole. For CquiOR10A73L-skatole, two models each formed a parallel pi-stacking interaction with Tyr-185 (*Extended—Table 13*). For CquiOR10-indole, a model formed a parallel pi-stacking interaction with Tyr-138 and another model formed

with Tyr-152. For CquiOR10A73L-indole, eight parallel pi-stacking interactions

with Tyr-185 were identified across seven representative models.

From the PLIP analysis of our docking study, we find of most importance skatole

and indole not forming contacts with Ala-73 in CquiOR10 models. By contrast,

skatole formed hydrophobic contacts with Leu-73 in five of the ten representative

models, while indole formed contacts with Leu-73 in two representative models.

This suggests that Ala-73 may indirectly affect specificity by modulating the

volume of the binding pocket.

Structurally aligning CquiOR10 and CquiOR10A73L receptors by TM7b

(*Extended—Table 14*), demonstrates approximately a 1 Å α-Carbon outward shift

of A73L (*Extended—Table 15*), suggesting a tightly constrained space in

CquiOR10 and an expanded space in CquiOR10A73L relative to the protein

backbone (*Figure 7*).

Figure 7. Comparison of CquiOR10 and CquiOR10A73L models. An

approximate 1 Å α-Carbon outward shift of Leu-73 (forest green) in CquiOR10

model relative to Ala-73 (light blue) in CquiOR10A73L model. Models were

superimposed using the TM7b region. Residue 73 amino nitrogen is colored in

dark blue, and carboxyl oxygen is colored in red in each model.

Figure supplement 1. Effect of single-point mutations around A73.

Figure supplement 2. Quantification of single-point mutations around A73.

This difference in binding pocket volume is not ligand-induced, but rather

independent of skatole or indole binding.

Consistent with our structural hypothesis of space constraints, mutation of Ala-73

in CquiOR10 to Ile or Val negatively affected receptor function, whereas

CquiOR10A73G retained specificity and showed higher sensitivity. Specifically,

CquiOR10A73I/CquiOrco- and CquiOR10A73V/CquiOrco-expressing oocytes did

not respond to skatole or indole (*Figure 7—figure supplement 1A,B*). We

concluded that mutations with these bulkier residues caused loss of binding to

indole or skatole, given that these receptors were functional, as indicated by the

potent responses elicited by OLC12. On the other hand,

CquiOR10A73G/CquiOrco-expressing oocytes showed the same skatole

specificity as the wildtype receptor (*Figure 7—figure supplement 1C*).

Interestingly, skatole elicited stronger currents recorded from CquiOR10A73G

than the wildtype receptor CquiOR10 (*Figure 7—figure supplement 2*), further

implying a higher affinity for skatole when there is a reduced constraint, or

increased volume, of the binding pocket.

Since CquiOR2 is homologous to CquiOR10, we also propose that our space

constraint structural hypothesis can be applied to CquiOR2. While we did not

perform CquiOR2 docking, we indirectly examined space constraints in CquiOR2

by mutating Leu-74 to Ile, Val, or Gly. We then tested oocytes co-expressing

CquiOrco with CquiOR2L74I, CquiOR2L74V, or CquiOR2L74G (*Figure 7—figure*

supplement 1D-F). Consistent with the relaxation of the space constraints,

CquiOR2L74I retained the specificity of the wildtype receptor to indole, whereas

CquiOR2L74G showed a reverse specificity profile (*Figure 7—figure supplement*

1F), i.e. more robust response to skatole than indole. Interestingly,

CquiOR2L74V/CquiOrco-expressing oocytes generated nearly equal, albeit

small, currents when stimulated with the two oviposition attractants (skatole,

8.7±2.7 nA; indole, 9.0±2.5 nA at 100 µM; *n* = 3, P>0.9999, Wilcoxon matched-

pairs signed-rank test).

Figure 8. Concentration-dependent responses elicited by 3-ethylindole in

oocytes co-expressing CquiOrco with CquiOR10, CquiOR2, or single-point

mutants. Bars represent SEM (n = 4-10).

Next, we tested the space constraints hypothesis with a bulkier ligand, 3-

ethylindole. A CquiOR10 mutant with a less space-filling residue,

CquiOR10A73G, elicited responses to 3-ethylindole (526±110 nA) higher than

the responses to indole (205±81 nA, at 100 µM), although less robust than the

skatole responses (1939±142 nA; all at 100 µM) (*Figure 7—figure supplement 3*).

By contrast, receptors with a bulkier residue (CquiOR2WT, CquiO10A73L, and

CquiOR2L74I) did not respond to 3-ethylindole (Figure 8). However, CquiOR2

mutant with less space-filling residues, CquiOR2L74A and CquiOR2L74G,

responded to 3-ethylindole in a dose-dependent manner (Figure 8). Additionally,

CquiOR10A73G elicited dose-dependent strong responses to 3-ethylindole than

CquiOR10 (Figure 8). In summary, the findings that a bulkier, non-natural ligand,

elicited more robust responses when CquiOR10 and CquiOR2 residues at 73

and 74, respectively, were mutated into Gly are consistent with the space

constraints hypothesis.

DISCUSSION

Our mutation studies demonstrated that a single amino acid residue substitution

in two narrowly tuned odorant receptors from the southern house mosquito can

revert their specificity to the oviposition attractants skatole and indole.

RoseTTAFold modeling and RosettaLigand docking studies suggest that the

specificity determinant amino acid residue (Ala-73) did not form direct contacts

with ligands but rather provided a finely-tuned, sensitive volumetric space to

accommodate odorants. Interpreting CquiOR10-A73L as an expanded space is

counterintuitive when compared to our oocyte recordings; the wild-type

CquiOR10 receptor responded better to the bulkier skatole, (*Figure 1A*), while

CquiOR10-A73L responded better to the smaller indole (*Figure 2I*).

Considering both experimental and modeling data, the reason for differing

receptor responses at residue 73 could be due to the repacking of this position to

accommodate more carbon sidechain atoms that are reducing the binding pocket

volume. Leu-73 contains three additional carbon sidechain atoms: γ-Carbon, δ1-

Carbon, and δ2-Carbon. Adding these atoms increases the sidechain rotatable

bond count from 0 (for Ala-73) to 2 (for Leu-73), causing an increase in

repackable states for the binding pocket. From our models, the distance of Leu-

73 α-Carbon to its δ1-Carbon or δ2-Carbon is approximately 2.8 to 3.9 Å. The

Ala-73 α-Carbon to its β-Carbon is approximately 1.5 Å. Since the Ala-73 α-

Carbon is approximately 0.9 to 1.1 Å inward relative to Leu-73 α-Carbon, Leu-73

C-δ atoms could be further inward by approximately 0.2-1.5 Å relative to Ala-73

C-β. This could cause repacking of surrounding side chains while reducing the

volume of the binding pocket, or shift key, nearby residues required for receptor

response. Further, from our models, the Carbon-Carbon length of skatole's

methyl group is approximately 1.5 Å. This distance discrepancy, the fact that

indole is a rigid molecule, and that the methyl group of skatole is the only

rotamer, could also suggest that the binding pocket around position 73 is tightly

regulated. The 1.5 Å increase from skatole's methyl group could prevent skatole

from occupying the appropriate configuration for receptor response with Leu-73.

Conversely, the additional inward 0.2-1.5 Å increase from Leu-73's sidechain

could provide indole more contacts or a more favorable configuration with

surrounding CquiOR10 sidechains to prompt increased receptor response. This

hypothesis is supported by seven of ten representative models of

CquiOR10A73L-indole having eight total parallel pi-stacking interactions with Tyr-

185; this is a considerable increase compared to two representative interactions

of CquiOR10-indole having two total parallel pi-stacking interactions with Tyr-138

and Tyr-152. Combined, our results suggest that the residue at 73 provide a

finely tuned volumetric space to accommodate specific oviposition attractants.

While modeling generally has a degree of uncertainty, RosettaFold and

AlphaFold are highly accurate (Baek & Baker, 2022). Given the challenge of

solving insect OR cryo-EM structures, the new accurate modeling methods are

currently the best way to advance the field and generate hypothesis for future

empirical tests.

513
514

Materials and methods

Construction of chimeric receptors. We used a previously obtained pGEMHE-

517 CquiOR10 (Hughes et al., 2010) plasmid to amplify the sequences of the desired

chimeric receptors. To obtain the full length of chimeric *OR* genes, we used

specific primers, with overlapping sequences at each end to target sequences

(adaptor underlined) for cloning into pGEMHE vector, which was linearized at

restriction sites *XmaI* (5'-CCCGGG-3') and *XbaI* (5'-TCTAGA-3'):CqOR10_Fw 5'-

GATCAATTCCCCGGGACCATGACCGCGGCACCCATTTT-3' and CqOR10_Rv

5'-CAAGCTTGCTCTAGATCAATTATAAACGCGTCTCAGCAGGGT-3'. To

generate a chimeric OR by PCR amplification, we designed specific primers for

the desired CquiOR2 TM domain. Simultaneously, we prepared the recipient

fragments of CquiOR10 to receive the CquiOR2 TM domains. The fragments

were then assembled using In-Fusion HD cloning kit (Clontech) to obtain the

desired chimeric OR. Each CquiOR2 TM domain was divided into two parts,

each part was synthesized with a specific primer, except for CquiOR2 TM7

domain, which used only one primer with the full domain. These primers

contained an overlap with CquiOR10 sequence followed by one part of the

CquiOR2 TM sequence and an overlap with the other part of the CquiOR2 TM

domain sequence, which was synthesized in another PCR. The underlined

sequences represent CquiOR2 TM domains and, unless otherwise specified,

sequences in italic are overlaps for CquiOR2 TM domains. CqOR10^{M1}_Fw: *5'-*

*GGTGACCGTGCTGA*ACGTGTTCCAGTTTATGAACCTGTTTCGAGCCTGGGG

CAACATC-3'; CqOR10^{M1}_Rv: 5'-

*TCAGCACGGTCACC*GGAATGCAGCCGAGGAGGTAACTGAGGACGTTGCCT

ACTGTGATCTCAAGG-3'; CqOR10^{M2}_Fw: 5'-

*CCGTGCTGTACTT*CAACCTTGTGTTGAGAACCACGTTTATACTGTGCAATCG

TCAGGATTATGAGG-3'; CqOR10^{M2}_Rv: 5'-

*AAGTACAGCACGG*TGAAATATCCGTCGATGATGATTTTGTCGATGTTGCCCC

AGGCT-3'; CqOR10^{M3}_Fw: 5'-

*GTTCATCAGTGCGT*GCTTCGTGACGTATCCGCTTTTTTCACCGACACGTAGC

CTCCCGTACG-3'; CqOR10^{M3}_Rv: 5'-

*ACGCACTGATGAAC*GCTCCCAGCCAGAGGTTCGATTTGGACAGCAGTCGG

GCACGTTTGGTGA-3'; CqOR10^{M4}_Fw: 5'-

*CACGTTTCCGGCGT*GCTGCATGTACATTCCGTTTACCAGCTTCTTCGCCACG

ACTACTTTG-3'; CqOR10^{M4}_Rv: 5'-

*ACGCCGGAAACGTG*AGGTACACTTGCAGAAAAACACAACCTGGTACAGGG

GCGTC-3'; CqOR10^{M5}_Fw: 5'-

*GCTATGCGCCTTGC*TGTTTCTACTTAGCACCAGCAATAATCCCGCGCAAATT

ATCATCGTGG-3'; CqOR10^{M5}_Rv: 5'-

*GCAAGGCGCATAGC*ATCATCCCAAACGATAGCAACTCAATCAGACAGATGT

AGGTCACCAGCG-3'; CqOR10^{M6}_Fw: 5'-

*TCTTTATGATTCT*GTCCCAGATGTACGCCCTGTACTGGCACGCCAACGAGCT

GCG-3'; CqOR10^{M6}_Rv: 5'-

*AGAATCATAAAGA*TGTACGATCCGATCATCACCATCTGCGCGGGATTTTCGA

TAATGTTCAGC-3'; CqOR10^{M7}_Rv: 5'-

*CAAGCTTGCTCTAGA*TCAATTATAAACGCGTCTCAGCAACGTAAAGTACGAA

TACGAGGCATTGATCAACTTTTGAAACATTTCCAAGGTCATCGGATAGACGT

TGCCTACTGTGATCTCAAGG-3' (here the italic represents the pGEMHE

adaptor. The two fragments were amplified using a combination of CqOR10_Fw

with CqOR10^{Mx}_Rv primers and CqOR10_Rv with CqOR10^{Mx}_Fw primers. Then,

we cloned the two fragments amplified by PCR into pGEMHE vector using the In

Fusion system as described below. Chimeric plasmid with a single TM swapped

was used as a template to generate chimeric OR with two TM swapped and

subsequently chimeric ORs with multiple CquiOR2 TM domains.

Chimeric OR cloning and subcloning into pGEMHE. The fragments

amplification was performed using Platinum Taq DNA Polymerase High Fidelity

(Thermo Fisher Scientific) with the following conditions: 95°C for 5 min, followed

by 30 cycles of 95°C for 20 s, 57ºC 30 s for annealing and 68°C for 1.5 min, and

extension at 68°C for 5 min. PCR products were purified by QIAquick gel

extraction kit (Qiagen). The target pGEMHE plasmid was cut by *XmaI* and *XbaI*

in separate reactions. Ligation was done with In Fusion (Takara Bio USA) system

and the transformation was performed using Stellar competent cells (Takara Bio

USA) in heat-shock. After selecting the cells, plasmids were extracted with

QIAprep Spin Miniprep kit (Qiagen). The cloned gene was verified by DNA

sequencing (Berkeley Sequencing Facility).

Site-directed point mutagenesis and fragment replacement. Phusion™ Site-

Directed Mutagenesis Kit (Thermo Scientific, West Palm Beach, FL) was used to

generate point mutations and TM fragment replacements. Mutations were

created with mismatched 5′-phosphorylated mutagenic primers and PCR

amplification. The chimeric sequences (CquiOR10^{M2,5,6,7}) in pGEMHE vector

were used as templates for rescues, whereas the wildtype sequences in

pGEMHE vector served as templates for point-mutations. Rescue primers:

CqOR10M2_M2567OUTERup: 5'-

GATGATGACCTCGTCGATGTTGCCCCAGGC-3';

CqOr10M2_M2567OUTERdn: 5'-

AACGCATATTTCACCGTGCTGTACTTCAACC-3'; CqOr10M2_M2567INNERup:

5'-CGCAGCACCGCGTTGAAGTACAGCACGGTG-3';

CqOr10M2_M2567INNERdn: 5'-AACAATTTTCATACTGTGCAATCGTCAGGA-

3'; CqOR10M2_M2567MID-INNERup: 5'-

CATCGACGGCTACTTTGCGATGATTTTCTTCAACGCG-3';

CqOR10M2_M2567MID-INNERdn: 5'-

ATGATTTTGTCGATGTTGCCCCAGGCTC-3'; CqOr10M2_M2567_L73Aup: 5'-

CGTTGAAGTACAGCACGGTGAAATAT-3'; CqOr10M2_M2567_L73Adn: 5'-

CTGTGTTGAGAACCACGTTTATACTGT-3'; CqOr10M2_M2567_T78Iup: 5'-

ATGGTTCTCAACACAAGGTTGAAGTA-3'; CqOr10M2_M2567_T78Idn: 5'-

CTTTATACTGTGCAATCGTCAGGATTATG-3'; Point mutation primers:

CqOR10A73up: 5'-GTTGAAGAAAATCATCGCAAAGTA-3'; CqOR10A73Ldn: 5'-

CTGGTGCTGCGAACAATTTTC-3'; CqOR10A73Adn: 5'-

GCGGTGCTGCGAACAATTTTC-3'; CqOR10A73Gdn: 5'-

GGGGTGCTGCGAACAATTTTC-3'; CqOR10A73Vdn: 5'-

GTGGTGCTGCGAACAATTTTC-3'; CqOR10A73Idn: 5'-

ATCGTGCTGCGAACAATTTTC-3'; CqOR2L74up: 5'-

GTTGAAGTACAGCACGGTGAAATA-3'; CqOr2L74Adn: 5'-

GCTGTGTTGAGAACCACGTTTATAC-3'; CqOr2L74Idn: 5'-

ATCGTGTTGAGAACCACGTTTATAC-3'; CqOr2L74Gdn: 5'-

GGGGTGTTGAGAACCACGTTTATAC-3'; CqOr2L74Vdn: 5'-

GTGGTGTTGAGAACCACGTTTATAC-3'. The amplified linear PCR products

containing the desired modification were ligated and transformed into Stellar™

Competent Cells (Takara Bio USA). All sequences were confirmed by DNA

sequencing (UC Berkeley DNA Sequencing Facility).

In vitro transcription, oocyte microinjection and two-electrode voltage-

clamp assay (TEVC). Capped OR cRNA was prepared using mMESSAGE

mMACHINE T7 Kit (Ambion) as previously described (Xu et al., 2019). Purified

OR cRNA was resuspended in nuclease-free water at 200 ng/µL and

microinjected into *Xenopus laevis* oocytes on stage V or VI (EcoCyte Bioscience)

along with the same amount of CquiOrco. Injected oocytes were incubated at 18

°C for 3–7 d in a modified Barth's solution [88 mM NaCl, 1 mM KCl, 2.4 mM

NaHCO$_3$, 0.82 mM MgSO$_4$, 0.33 mM Ca(NO$_3$)$_2$, 0.41 mM CaCl$_2$, and 10 mM

HEPES at pH 7.4] supplemented with 10 µg/mL gentamycin, 10 µg/mL

streptomycin and 1.8 mM sodium pyruvate. Two-electrode voltage-clamp

technique (TEVC) was used to measure odorant-induced currents at a holding

potential of −80 mV. Signals were amplified with an OC-725C amplifier (Warner

Instruments), low-pass filtered at 50 Hz, and digitized at 1 kHz. Data acquisition

and analysis were performed with Digidata 1440A and software pCLAMP 10

(Molecular Devices). Skatole (98%, CAS# 83-34-1), indole (98%, CAS# 120-72-

9), and 3-ethylindole (97%, CAS# 1484-19-1) were provided by Sigma-Aldrich

(Milwaukee, WI), ACROS Organics (Geel, Belgium), and AmBeed (Arlington Hts,

IL), respectively. Phenol (99.5%, CAS# 108-95-2), 3,5-dimethylphenol (99%,

CAS# 108-68-9), 1-methylindole (97%, CAS# 603-76-9), 2-methylindole (98%,

CAS# 95-20-5), 5-methylindole (99%, CAS# 614-96-0), 6-methylindole (97%,

CAS# 3420-02-8) and 7-methylindole (97%, CAS# 933-67-5) were acquired from Sigma-Aldrich, and 4-methylindole (99%, CAS# 16096-32-5) was provided by ACROS.

Rosetta structural modeling and docking. We used RoseTTAFold (Baek et al., 2021) and AlphaFold (Jumper et al., 2021) to generate predicted structures of CquiOR10, CquiOR2, CquiOR10A73L, and CquiOR2L74A. The OpenEye Omega toolkit (Hawkins et al., 2010) was used to generate conformer libraries. Molecular docking was performed using RosettaLigand (Davis & Baker, 2009; DeLuca et al., 2015). The top models were visually analyzed using UCSF Chimera (Pettersen et al., 2004).

To verify RosettaLigand could effectively sample homologous receptors in complex with odorants, we used the structure of eugenol in complex with the insect olfactory receptor OR5 from *Machilis hrabei* (MhOR5) as a control test case (Del Marmol et al., 2021). After verifying RosettaLigand could recapitulate the MhraOR5 — eugenol complex from structural studies, we docked indole and skatole to the RoseTTAFold structure models of CquiOR10 and CquiOR10A73L receptors with the same method. Briefly, the initial placement of all ligands into respective structures was guided by the position of eugenol in a complex with MhOR5 as a basis since it is the only homologous structure with an odorant. For an unbiased receptor sampling, an initial transformation was performed on the ligand relative to the receptor site using Rosetta's Transform mover. Within a 7 Å sphere, the ligand underwent a Monte Carlo simulation, whereby the ligand was allowed to translate up to 0.2 Å and rotate up to 20 degrees. This was performed 500 times on the ligand, with the lowest scoring pose used as the starting pose for docking. This form of docking is termed local docking and is commonplace when there is experimental evidence and homologous structures as templates. physiologically relevant binding modes can be identified with local, high-

resolution ligand docking (Kaufmann et al., 2009). For additional methods, see

Appendix 2.

Appendix 1

Preliminary screening of chimeric receptors. During the life of this project, the

structure of an odorant receptor coreceptor AbakOrco was reported (Butterwick

et al., 2018). We then compared the Cryo-EM structure (Butterwick et al., 2018)

with the predicted topology for AbakOrco using the same OCTOPUS method

(Viklund & Elofsson, 2008) we used to identify CquiOR10 and CquiOR2 TMs

(*Figure 1—figure supplement 1*) (Viklund & Elofsson, 2008). The almost perfect

overlap between OCTOPUS prediction and the Cryo-EM structure (*Figure 1—*

figure supplement 3) validated not only our TM predictions (*Figure 1—figure*

supplement 1), but also the 21 chimeric ORs already tested when the structure of

the coreceptor AbakOrco was reported (Hughes et al., 2017; Luetje et al., 2013).

CquiOR10[M1,2,3,4,5,6,7]/CquiOrco-expressing oocytes did not respond to skatole or

indole. Likewise, the chimeric receptor with six transmembrane domains

replaced, CquiOR10[M1,2,3,4,5,6] was also silent. Next, we replaced one TM at a

time. CquiOR10[M1] and CquiOR10[M2] were silent, whereas CquiOR10[M3] (*Figure*

1—figure supplement 2A), CquiOR10[M4] (*Figure 1—figure supplement 2B*),

CquiOR10[M5] (*Figure 1—figure supplement 2C*), CquiOR10[M6] (*Figure 1—figure*

supplement 2D), and CquiOR10[M7] (*Figure 1—figure supplement 2E*) responded

to both skatole and indole with specificity similar to that of the wildtype receptor.

Although these chimeric receptors differed from CquiOR10 in terms of sensitivity,

they showed the same specificity to skatole as CquiOR10 (*Figure 1A*). While

CquiOR10[M3] and CquiOR10[M4] were less sensitive than CquiOR10, CquiOR10[M7]

showed comparable sensitivity to CquiOR10, and CquiOR10[M5] and CquiOR10[M6]

were more sensitive to skatole than the wildtype receptor. We remained focused

on specificity and did not attempt to quantify the different responses.

696 We tested a chimeric receptor with the other five TM domains swapped.

697 CquiOR10^{M3,4,5,6,7}/CquiOrco-expressing oocytes mirrored the profiles obtained

698 with CquiOR10 in terms of sensitivity and specificity (*Figure 1—figure*

699 *supplement 2F*). We then kept the wildtype TM7, in addition to TM1 and TM2, in

700 the next chimeric ORs tested. Albeit less sensitive than CquiOR10,

701 CquiOR10^{M3,4,5,6} showed the same specificity of the wildtype receptor (*Figure 1—*

702 *figure supplement 2G*). Following, we tested chimeric receptors with three TM

703 domains swapped. CquiOR10^{M4,5,6} (*Figure 1—figure supplement 2H*),

704 CquiOR10^{M3,5,6} (*Figure 1—figure supplement 2I*), CquiOR10^{M3,4,6} (*Figure 1—*

705 *figure supplement 2J*), and CquiOR10^{M3,4,5} (*Figure 1—figure supplement 2K*) did

706 not differ from CquiOR10 in terms of specificity or sensitivity. Next, we focused

707 on chimeric receptors with two TMs swapped. Thus, we tested CquiOR10^{M3,4}

708 (*Figure 1—figure supplement 2L*), CquiOR10^{M3,5} (*Figure 1—figure supplement*

709 *2M*), CquiOR10^{M3,6} (*Figure 1—figure supplement 2N*), CquiOR10^{M3,7} (*Figure 1—*

710 *figure supplement 2O*), CquiOR10^{M4,5} (*Figure 1—figure supplement 2P*),

711 CquiOR10^{M4,6} (*Figure 1—figure supplement 2Q*), CquiOR10^{M4,7} (*Figure 1—figure*

712 *supplement 2R*), CquiOR10^{M5,6} (*Figure 1—figure supplement 2S*), CquiOR10^{M5,7}

713 (*Figure 1—figure supplement 2T*), and CquiOR10^{M6,7} (*Figure 1—figure*

714 *supplement 2U*). They all showed skatole-specific profiles, most of them with

715 comparable sensitivity, whereas CquiOR10^{M4,5}, CquiOR10^{M5,7}, and CquiOR10^{M6,7}

716 were more sensitive and CquiOR10^{M3,6} less sensitive than CquiOR10.

717 Interestingly, none of the chimeric receptors having TM1 as one of the two

718 domains swapped was functional. Specifically, oocytes co-expressing CquiOrco

719 with either CquiOR10^{M1,2}, CquiOR10^{M1,3}, CquiOR10^{M1,4}, CquiOR10^{M1,5},

720 CquiOR10^{M1,6}, CquiOR10^{M1,7} did not respond to skatole or indole (*Extended—*

721 *Table 1*). Likewise, CquiOR10^{M2,3}, CquiOR10^{M2,4}, CquiOR10^{M2,5}, and

722 CquiOR10^{M2,6} did not respond to these oviposition attractants (*Extended—Table*

723 *1*).

Appendix 2

Ligand conformer library generation. The Protein DataBank file (.pdb) of eugenol was extracted from the MhOR5 structure (PDB ID: 7LID), while indole and skatole were extracted as Structure-Data Files files (.sdf) from PubChem (Kim et al., 2021). Each ligand structure was bond-corrected, protonated at pH 7.4, energy minimized using the Merck Molecular Force Field (Halgren, 1996), and saved as Tripos Mol2 files (.mol2) with the Avogadro software (Hanwell et al., 2012). Next, partial charge, atom, and bond type assignment were performed with Antechamber AM1BCC correction using Amber Tools (Case et al., 2021; Salomon-Ferrer et al., 2013); AM1BCC partial-charge correction is common in Rosetta ligand docking protocols (Park et al., 2021) and has been shown to be similarly ranked with other Rosetta-based partial-charge methods for RosettaLigand (Smith & Meiler, 2020). An in-house script using the OpenEye Omega toolkit and Rosetta was used to generate the conformer library and associated Rosetta-readable ligand parameters file.

Protein receptor preparation. The MhOR5 structure was relaxed with backbone constraints using the RosettaRelax protocol (Nivon et al., 2013) prior to RosettaLigand docking. This is commonplace to allow the repacking of protein sidechains and minimization of the structure into the Rosetta score function for comparison between structures. For RoseTTAFold-generated structures, the RosettaRelax protocol was not performed since the Rosetta score function was used to generate the structures.

RosettaLigand docking. RosettaLigand docking was performed using RosettaScripts protocols described previously (Davis & Baker, 2009; DeLuca et al., 2015). Briefly, initial placement of all ligands into respective structures was guided by the position of eugenol in complex with MhOR5 as a basis. For an unbiased sampling of the receptor, an initial transformation was performed on the

ligand relative to the receptor site using Rosetta's Transform mover. Within a 7 Å

sphere, the ligand underwent a Monte Carlo simulation, whereby the ligand was

allowed to translate up to 0.2 Å and rotate up to 20 degrees. This was performed

500 times on the ligand, with the lowest scoring pose used as the starting pose

for docking. Six cycles of high-resolution docking were performed with repacking

of sidechains every third iteration using Rosetta's HighResDocker mover. Lastly,

the protein/ligand complex was minimized using Rosetta's FinalMinimizer mover

and interface scores reported using the InterfaceScoreCalculator.

For every test case, 100,000 docking poses were generated. The top 10,000

models were selected by Rosetta interface energy (Interface_delta_X) metric,

followed by ligand clustering using Biomol2Clust (D. et al., 2021), the hdbscan

algorithm (McInnes et al., 2017), and a minimum cluster threshold of 50 poses.

The 10 most frequently sampled clusters were visually analyzed using UCSF

Chimera (Pettersen et al., 2004) by using the lowest interface energy pose in

each cluster as the representative for that cluster. The Protein Ligand Interaction

Profiler (Salentin et al., 2015) was performed on these representative poses to

identify potential hydrophobic interactions, hydrogen bonds, π-Stacking, and salt

bridges. Hydrogen bond and pi-stacking interactions were filtered by previously

reported bond distances (Bissantz et al., 2010) to present a conservative

estimate of potential contacts. Atom types from PLIP analysis are reported using

IDATM nomenclature (Meng & Lewis, 1991).

Acknowledgments. We thank Drs. David K. Wilson, Enoch Baldwin (UC

Davis), and members of our research groups for enlightening discussions. The

Orco ligand candidate OLC12 was provided by the Vanderbilt Institute of

Chemical Biology, Chemical Synthesis Core, Vanderbilt University. Research

reported in this publication was supported by the National Institute of Allergy and

Infectious Diseases (NIAID) of the National Institutes of Health under award

number R01AI095514. The content is solely the responsibility of the authors and
does not necessarily represent the official views of the NIH.

Competing Interest Statement: The authors declare no competing interest.

Author Contributions: W.S.L., P.X., and V.Y-Y. designed research; F.P.F., P.X.
B.J.H, V.Y-Y., and W.S.L. performed research and analyzed the data; W.S.L.
and B.J.H wrote the paper, and F.P.F., P.X., and V.Y-Y. edited the manuscript.

Data Availability. All data generated during this study are included in the
manuscript and supporting files.

References

Adasme, M. F., Linnemann, K. L., Bolz, S. N., Kaiser, F., Salentin, S., Haupt,
V J., & Schroeder, M. (2021). PLIP 2021: expanding the scope of the
protein–ligand interaction profiler to DNA and RNA. *Nucleic Acids*
Research, 49(W1), W530-W534. https://doi.org/10.1093/nar/gkab294

Arensburger, P., Megy, K., Waterhouse, R. M., Abrudan, J., Amedeo, P., Antelo,
B., Bartholomay, L., Bidwell, S., Caler, E., Camara, F., Campbell, C. L.,
Campbell, K. S., Casola, C., Castro, M. T., Chandramouliswaran, I.,
Chapman, S. B., Christley, S., Costas, J., Eisenstadt, E., Feschotte, C.,
Fraser-Liggett, C., Guigo, R., Haas, B., Hammond, M., Hansson, B. S.,
Hemingway, J., Hill, S. R., Howarth, C., Ignell, R., Kennedy, R. C., Kodira,
C. D., Lobo, N. F., Mao, C., Mayhew, G., Michel, K., Mori, A., Liu, N.,
Naveira, H., Nene, V., Nguyen, N., Pearson, M. D., Pritham, E. J., Puiu,
D., Qi, Y., Ranson, H., Ribeiro, J. M., Roberston, H. M., Severson, D. W.,
Shumway, M., Stanke, M., Strausberg, R. L., Sun, C., Sutton, G., Tu, Z. J.,
Tubio, J. M., Unger, M. F., Vanlandingham, D. L., Vilella, A. J., White, O.,
White, J. R., Wondji, C. S., Wortman, J., Zdobnov, E. M., Birren, B.,
Christensen, B. M., Collins, F. H., Cornel, A., Dimopoulos, G., Hannick, L.
I., Higgs, S., Lanzaro, G. C., Lawson, D., Lee, N. H., Muskavitch, M. A.,
Raikhel, A. S., & Atkinson, P. W. (2010, Oct 1). Sequencing of Culex
quinquefasciatus establishes a platform for mosquito comparative
genomics. *Science, 330*(6000), 86-88.
https://doi.org/10.1126/science.1191864

Auer, T. O., Khallaf, M. A., Silbering, A. F., Zappia, G., Ellis, K., Alvarez-Ocana,
823   R., Arguello, J. R., Hansson, B. S., Jefferis, G., Caron, S. J. C., Knaden,
824   M., & Benton, R. (2020, Mar). Olfactory receptor and circuit evolution
825   promote host specialization. *Nature, 579*(7799), 402-408.
826   https://doi.org/10.1038/s41586-020-2073-7

828 Baek, M., & Baker, D. (2022, Jan). Deep learning and protein structure modeling.
829   *Nat Methods, 19*(1), 13-14. https://doi.org/10.1038/s41592-021-01360-8

831 Baek, M., DiMaio, F., Anishchenko, I., Dauparas, J., Ovchinnikov, S., Lee, G. R.,
832   Wang, J., Cong, Q., Kinch, L. N., Schaeffer, R. D., Millan, C., Park, H.,
833   Adams, C., Glassman, C. R., DeGiovanni, A., Pereira, J. H., Rodrigues, A.
834   V., van Dijk, A. A., Ebrecht, A. C., Opperman, D. J., Sagmeister, T.,
835   Buhlheller, C., Pavkov-Keller, T., Rathinaswamy, M. K., Dalwadi, U., Yip,
836   C. K., Burke, J. E., Garcia, K. C., Grishin, N. V., Adams, P. D., Read, R.
837   J., & Baker, D. (2021, Aug 20). Accurate prediction of protein structures
838   and interactions using a three-track neural network. *Science, 373*(6557),
839   871-876. https://doi.org/10.1126/science.abj8754

841 Benton, R., Sachse, S., Michnick, S. W., & Vosshall, L. B. (2006, Feb). Atypical
842   membrane topology and heteromeric function of Drosophila odorant
843   receptors in vivo. *PLoS Biol, 4*(2), e20.
844   https://doi.org/10.1371/journal.pbio.0040020

846 Bidula, S., Nadzirin, I. B., Cominetti, M., Hickey, H., Cullum, S. A., Searcey, M.,
847   Schmid, R., & Fountain, S. J. (2022, Jan). Structural basis of the negative
848   allosteric modulation of 5-BDBD at human P2X4 receptors. *Mol
849   Pharmacol, 101*(1), 33-44. https://doi.org/10.1124/molpharm.121.000402

851 Bissantz, C., Kuhn, B., & Stahl, M. (2010, Jul 22). A medicinal chemist's guide to
852   molecular interactions. *J Med Chem, 53*(14), 5061-5084.
853   https://doi.org/10.1021/jm100112j

855 Blackwell, A., Mordue, A. J., Hansson, B. S., Wadhams, L. J., & Pickett, J. A.
856   (1993, Dec). A behavioral and electrophysiological study of oviposition
857   cues for Culex quinquefasciatus. *Physiological Entomology, 18*(4), 343-
858   348. <Go to ISI>://WOS:A1993MP27600003

860 Bohbot, J. D., Jones, P. L., Wang, G., Pitts, R. J., Pask, G. M., & Zwiebel, L. J.
861 (2011, Jan). Conservation of indole responsive odorant receptors in
862 mosquitoes reveals an ancient olfactory trait. *Chemical Senses, 36*(2),
863 149-160. https://doi.org/10.1093/chemse/bjq105

864
865 Bohbot, J. D., & Pitts, R. J. (2015, 2015-April-10). The narrowing olfactory
866 landscape of insect odorant receptors [Hypothesis and Theory]. *Frontiers*
867 *in Ecology and Evolution, 3*. https://doi.org/10.3389/fevo.2015.00039

868
869 Butterwick, J. A., Del Marmol, J., Kim, K. H., Kahlson, M. A., Rogow, J. A., Walz,
870 T., & Ruta, V. (2018, Aug). Cryo-EM structure of the insect olfactory
871 receptor Orco. *Nature, 560*(7719), 447-452.
872 https://doi.org/10.1038/s41586-018-0420-8

873
874 Cao, S., Liu, Y., Wang, B., & Wang, G. R. (2021, Sep 24). A single point mutation
875 causes one-way alteration of pheromone receptor function in two Heliothis
876 species. *Iscience, 24*(9). https://doi.org/ARTN 102981

877 10.1016/j.isci.2021.102981

878
879 Carey, A. F., Wang, G., Su, C. Y., Zwiebel, L. J., & Carlson, J. R. (2010, Mar 04).
880 Odorant reception in the malaria mosquito Anopheles gambiae. *Nature,*
881 *464*(7285), 66-71. https://doi.org/10.1038/nature08834

882
883 Case, D. A., Aktulga, H. M., Belfon, K., Ben-Shalom, I. Y., Brozell, S. R., Cerutti,
884 D. S., Cheatham III, T. E., Cisneros, G. A., Cruzeiro, V. W. D., Darden, T.
885 A., & Duke, R. E. (2021). *Amber 2021.*
886 https://ambermd.org/doc12/Amber21.pdf

887
888 Choo, Y. M., Xu, P., Hwang, J. K., Zeng, F., Tan, K., Bhagavathy, G., Chauhan,
889 K., & Leal, W. S. (2018). Reverse chemical ecology approach for the
890 identification of an oviposition attractant for Culex quinquefasciatus. *Proc*
891 *Natl Acad Sci U S A, 115*(4), 714-719.

892
Clyne, P. J., Warr, C. G., Freeman, M. R., Lessing, D., Kim, J., & Carlson, J. R.
(1999, Feb). A novel family of divergent seven-transmembrane proteins:
candidate odorant receptors in Drosophila. *Neuron, 22*(2), 327-338.
https://doi.org/10.1016/s0896-6273(00)81093-4

D., T., Y., S., V., Š., & D., S. (2021). *Biomol2Clust*. In
https://biokinet.belozersky.msu.ru/Biomol2Clust

900
901   Davis, I. W., & Baker, D. (2009, Jan 16). RosettaLigand docking with full ligand
902   and receptor flexibility. *J Mol Biol, 385*(2), 381-392.
903   https://doi.org/10.1016/j.jmb.2008.11.010

904
905   Del Marmol, J., Yedlin, M. A., & Ruta, V. (2021, Sep). The structural basis of
906   odorant recognition in insect olfactory receptors. *Nature, 597*(7874), 126-
907   131. https://doi.org/10.1038/s41586-021-03794-8

908
909   DeLuca, S., Khar, K., & Meiler, J. (2015). Fully flexible docking of medium sized
910   ligand libraries with RosettaLigand. *PLOS ONE, 10*(7), e0132508.
911   https://doi.org/10.1371/journal.pone.0132508

912
913   Gao, Q., & Chess, A. (1999, Aug 15). Identification of candidate Drosophila
914   olfactory receptors from genomic DNA sequence. *Genomics, 60*(1), 31-39.
915   https://doi.org/10.1006/geno.1999.5894

916
917   Guo, S., & Kim, J. (2010). Dissecting the molecular mechanism of drosophila
918   odorant receptors through activity modeling and comparative analysis.
919   *Proteins: Structure, Function, and Bioinformatics, 78*(2), 381-399.
920   https://doi.org/https://doi.org/10.1002/prot.22556

921
922   Halgren, T. A. (1996, Apr). Merck molecular force field .1. Basis, form, scope,
923   parameterization, and performance of MMFF94. *Journal of Computational*
924   *Chemistry, 17*(5-6), 490-519. https://doi.org/Doi 10.1002/(Sici)1096-
925   987x(199604)17:5/6<490::Aid-Jcc1>3.0.Co;2-P

926
927   Hanwell, M. D., Curtis, D. E., Lonie, D. C., Vandermeersch, T., Zurek, E., &
928   Hutchison, G. R. (2012, Aug 13). Avogadro: an advanced semantic
929   chemical editor, visualization, and analysis platform. *Journal of*
930   *Cheminformatics, 4*. https://doi.org/Artn 17

931   10.1186/1758-2946-4-17

932
933   Hawkins, P. C., Skillman, A. G., Warren, G. L., Ellingson, B. A., & Stahl, M. T.
934   (2010, Apr 26). Conformer generation with OMEGA: algorithm and
935   validation using high quality structures from the Protein Databank and

Cambridge Structural Database. *J Chem Inf Model, 50*(4), 572-584. https://doi.org/10.1021/ci100031x

Hughes, D. T., Pelletier, J., Luetje, C. W., & Leal, W. S. (2010, Aug). Odorant receptor from the southern house mosquito narrowly tuned to the oviposition attractant skatole. *J Chem Ecol, 36*(8), 797-800. https://doi.org/10.1007/s10886-010-9828-9

Hughes, D. T., Pelletier, J., Rahman, S., Chen, S., Leal, W. S., & Luetje, C. W. (2017, May 01). Functional and nonfunctional forms of CquiOR91, an odorant selectivity subunit of Culex quinquefasciatus. *Chemical Senses, 42*(4), 333-341. https://doi.org/10.1093/chemse/bjx011

Hughes, D. T., Wang, G., Zwiebel, L. J., & Luetje, C. W. (2014, Nov). A determinant of odorant specificity is located at the extracellular loop 2-transmembrane domain 4 interface of an Anopheles gambiae odorant receptor subunit. *Chemical Senses, 39*(9), 761-769. https://doi.org/10.1093/chemse/bju048

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A. W., Kavukcuoglu, K., Kohli, P., & Hassabis, D. (2021, Aug). Highly accurate protein structure prediction with AlphaFold. *Nature, 596*(7873), 583-589. https://doi.org/10.1038/s41586-021-03819-2

Kaissling, K. E. (2014). Pheromone reception in Insects: The example of silk moths. In C. Mucignat-Caretta (Ed.), *Neurobiology of Chemical Communication*. https://www.ncbi.nlm.nih.gov/pubmed/24830037

Kaufmann, K. W., Dawson, E. S., Henry, L. K., Field, J. R., Blakely, R. D., & Meiler, J. (2009, Feb 15). Structural determinants of species-selective substrate recognition in human and Drosophila serotonin transporters revealed through computational docking studies. *Proteins, 74*(3), 630-642. https://doi.org/10.1002/prot.22178

Kim, S., Chen, J., Cheng, T., Gindulyte, A., He, J., He, S., Li, Q., Shoemaker, B. A., Thiessen, P. A., Yu, B., Zaslavsky, L., Zhang, J., & Bolton, E. E. (2021, Jan 8). PubChem in 2021: new data content and improved web interfaces. *Nucleic Acids Res, 49*(D1), D1388-D1395. https://doi.org/10.1093/nar/gkaa971

Larsson, M. C., Domingos, A. I., Jones, W. D., Chiappe, M. E., Amrein, H., & Vosshall, L. B. (2004, Sep 2). Or83b encodes a broadly expressed odorant receptor essential for Drosophila olfaction. *Neuron, 43*(5), 703-714. https://doi.org/10.1016/j.neuron.2004.08.019

Leal, W. S. (2013). Odorant reception in insects: roles of receptors, binding proteins, and degrading enzymes. *Annu Rev Entomol, 58*, 373-391. https://doi.org/10.1146/annurev-ento-120811-153635

Leal, W. S. (2016). Molecular biology of reception. *Pheromone Communication in Moths: Evolution, Behavior, and Application*, 127-138. <Go to ISI>://WOS:000388302800009

Leal, W. S. (2020). Mechanism of action of insect pheromones and other semiochemicals. In H.-W. Liu & T. P. Begley (Eds.), *Comprehensive Natural Products III* (pp. 222-236). Elsevier. https://www.sciencedirect.com/science/article/pii/B9780124095472147353

Leal, W. S., Choo, Y. M., Xu, P., da Silva, C. S., & Ueira-Vieira, C. (2013, Nov 12). Differential expression of olfactory genes in the southern house mosquito and insights into unique odorant receptor gene isoforms. *Proc Natl Acad Sci U S A, 110*(46), 18704-18709. https://doi.org/10.1073/pnas.1316059110

Leary, G. P., Allen, J. E., Bunger, P. L., Luginbill, J. B., Linn, C. E., Jr., Macallister, I. E., Kavanaugh, M. P., & Wanner, K. W. (2012, Aug 28). Single mutation to a sex pheromone receptor provides adaptive specificity between closely related moth species. *Proc Natl Acad Sci U S A, 109*(35), 14081-14086. https://doi.org/10.1073/pnas.1204661109

Luetje, C. W., Nichols, A. S., Castro, A., & Sherman, B. L. (2013). Functional assay of mammalian and Insect olfactory receptors using Xenopus oocytes. In C. J. Crasto (Ed.), *Olfactory Receptors: Methods and*

1014   *Protocols* (pp. 187-202). Humana Press. https://doi.org/10.1007/978-1-
1015   62703-377-0_14

1016
1017   Lundin, C., Kall, L., Kreher, S. A., Kapp, K., Sonnhammer, E. L., Carlson, J. R.,
1018   Heijne, G., & Nilsson, I. (2007, Dec 11). Membrane topology of the
1019   Drosophila OR83b odorant receptor. *FEBS Lett, 581*(29), 5601-5604.
1020   https://doi.org/10.1016/j.febslet.2007.11.007

1021
1022   Mboera, L. E. G., Takken, W., Mdira, K. Y., Chuwa, G. J., & Pickett, J. A. (2000,
1023   May). Oviposition and behavioral responses of Culex quinquefasciatus to
1024   skatole and synthetic oviposition pheromone in Tanzania. *Journal of
1025   Chemical Ecology, 26*(5), 1193-1203. https://doi.org/Doi
1026   10.1023/A:1005432010721

1027
1028   McInnes, L., Healy, J., & Astels, S. (2017). hdbscan: Hierarchical density based
1029   clustering. *J. Open Source Softw., 2*, 205.

1030
1031   Meng, E. C., & Lewis, R. A. (1991, Sep). Determination of Molecular Topology
1032   and Atomic Hybridization States from Heavy-Atom Coordinates. *Journal of
1033   Computational Chemistry, 12*(7), 891-898. https://doi.org/DOI
1034   10.1002/jcc.540120716

1035
1036   Millar, J. G., Chaney, J. D., & Mulla, M. S. (1992, Mar). Identification of
1037   oviposition attractants for Culex quinquefasciatus from fermented
1038   Bermuda grass infusions. *Journal of the American Mosquito Control
1039   Association, 8*(1), 11-17. <Go to ISI>://WOS:A1992HM11400002

1040
1041   Nakagawa, T., Sakurai, T., Nishioka, T., & Touhara, K. (2005, Mar 11). Insect
1042   sex-pheromone signals mediated by specific combinations of olfactory
1043   receptors. *Science, 307*(5715), 1638-1642.
1044   https://doi.org/10.1126/science.1106267

1045
1046   Neuhaus, E. M., Gisselmann, G., Zhang, W., Dooley, R., Stortkuhl, K., & Hatt, H.
1047   (2005, Jan). Odorant receptor heterodimerization in the olfactory system
1048   of Drosophila melanogaster. *Nat Neurosci, 8*(1), 15-17.
1049   https://doi.org/10.1038/nn1371

1050
1051   Nguyen, H. M., Singh, V., Pressly, B., Jenkins, D. P., Wulff, H., & Yarov-Yarovoy,
1052   V. (2017, Apr). Structural insights into the atomistic mechanisms of action

of small molecule inhibitors targeting the KCa3.1 channel pore. *Mol Pharmacol, 91*(4), 392-402. https://doi.org/10.1124/mol.116.108068

Nivon, L. G., Moretti, R., & Baker, D. (2013). A Pareto-optimal refinement method for protein design scaffolds. *PLOS ONE, 8*(4), e59004. https://doi.org/10.1371/journal.pone.0059004

Park, H., Zhou, G., Baek, M., Baker, D., & DiMaio, F. (2021, Mar 9). Force field optimization guided by small molecule crystal lattice data enables consistent sub-Angstrom protein-ligand docking. *J Chem Theory Comput, 17*(3), 2000-2010. https://doi.org/10.1021/acs.jctc.0c01184

Pask, G. M. (2020). Invertebrate olfactory receptors. In B. Fritzsch (Ed.), *The Senses: A Comprehensive Reference (Second Edition)* (pp. 453-465). Elsevier. https://doi.org/doi.org/10.1016/B978-0-12-809324-5.23814-5

Pellegrino, M., Steinbach, N., Stensmyr, M. C., Hansson, B. S., & Vosshall, L. B. (2011, Oct 27). A natural polymorphism alters odour and DEET sensitivity in an insect odorant receptor. *Nature, 478*(7370), 511-514. https://doi.org/10.1038/nature10438

Pelletier, J., Hughes, D. T., Luetje, C. W., & Leal, W. S. (2010). An odorant receptor from the southern house mosquito Culex pipiens quinquefasciatus sensitive to oviposition attractants. *PLOS ONE, 5*(4), e10090. https://doi.org/10.1371/journal.pone.0010090

Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C., & Ferrin, T. E. (2004, Oct). UCSF Chimera--a visualization system for exploratory research and analysis. *Journal of Computational Chemistry, 25*(13), 1605-1612. https://doi.org/10.1002/jcc.20084

Pitts, R. J., Huff, R. M., Shih, S. J., & Bohbot, J. D. (2021, Dec). Identification and functional characterization of olfactory indolergic receptors in Musca domestica. *Insect Biochem Mol Biol, 139*, 103653. https://doi.org/10.1016/j.ibmb.2021.103653

Pressly, B., Lee, R. D., Singh, V., Pessah, I. N., & Wulff, H. (2022, May). The seizure-inducing plastic explosive RDX inhibits the alpha1beta2gamma2

GABAA receptor. *Ann Clin Transl Neurol, 9*(5), 600-609.
https://doi.org/10.1002/acn3.51536

Reynolds, S. M., Kall, L., Riffle, M. E., Bilmes, J. A., & Noble, W. S. (2008, Nov).
Transmembrane topology and signal peptide prediction using dynamic
bayesian networks. *PLoS Comput Biol, 4*(11), e1000213.
https://doi.org/10.1371/journal.pcbi.1000213

Salentin, S., Schreiber, S., Haupt, V. J., Adasme, M. F., & Schroeder, M. (2015,
Jul 1). PLIP: fully automated protein-ligand interaction profiler. *Nucleic
Acids Res, 43*(W1), W443-447. https://doi.org/10.1093/nar/gkv315

Salomon-Ferrer, R., Case, D. A., & Walker, R. C. (2013, Mar-Apr). An overview
of the Amber biomolecular simulation package. *Wiley Interdisciplinary
Reviews-Computational Molecular Science, 3*(2), 198-210.
https://doi.org/10.1002/wcms.1121

Smith, S. T., & Meiler, J. (2020). Assessing multiple score functions in Rosetta
for drug discovery. *PLOS ONE, 15*(10), e0240450.
https://doi.org/10.1371/journal.pone.0240450

Stensmyr, M. C., Dweck, H. K., Farhan, A., Ibba, I., Strutz, A., Mukunda, L., Linz,
J., Grabe, V., Steck, K., Lavista-Llanos, S., Wicher, D., Sachse, S.,
Knaden, M., Becher, P. G., Seki, Y., & Hansson, B. S. (2012, Dec 7). A
conserved dedicated olfactory circuit for detecting harmful microbes in
Drosophila. *Cell, 151*(6), 1345-1357.
https://doi.org/10.1016/j.cell.2012.09.046

Viklund, H., & Elofsson, A. (2008). OCTOPUS: improving topology prediction by
two-track ANN-based preference scores and an extended topological
grammar. *Bioinformatics, 24*(15), 1662-1668.
https://doi.org/10.1093/bioinformatics/btn221

Vosshall, L. B., Amrein, H., Morozov, P. S., Rzhetsky, A., & Axel, R. (1999, Mar
5). A spatial map of olfactory receptor expression in the Drosophila
antenna. *Cell, 96*(5), 725-736. https://doi.org/10.1016/s0092-
8674(00)80582-6

1129 Wang, G., Carey, A. F., Carlson, J. R., & Zwiebel, L. J. (2010, Mar 02). Molecular
1130   basis of odor coding in the malaria vector mosquito Anopheles gambiae.
1131   *Proc Natl Acad Sci U S A, 107*(9), 4418-4423.
1132   https://doi.org/10.1073/pnas.0913392107

1133

1134 Xu, P., Choo, Y. M., Chen, Z., Zeng, F., Tan, K., Chen, T. Y., Cornel, A. J., Liu,
1135   N., & Leal, W. S. (2019, Jul 12). Odorant inhibition in mosquito olfaction.
1136   *Iscience, 19*, 25-38. https://doi.org/10.1016/j.isci.2019.07.008

1137

1138 Yang, K., Huang, L. Q., Ning, C., & Wang, C. Z. (2017, Oct 24). Two single-point
1139   mutations shift the ligand selectivity of a pheromone receptor between two
1140   closely related moth species. *Elife, 6*. https://doi.org/10.7554/eLife.29100

1141

1142 Yuvaraj, J. K., Roberts, R. E., Sonntag, Y., Hou, X. Q., Grosse-Wilde, E.,
1143   Machara, A., Zhang, D. D., Hansson, B. S., Johanson, U., Lofstedt, C., &
1144   Andersson, M. N. (2021, Jan 26). Putative ligand binding sites of two
1145   functionally characterized bark beetle odorant receptors. *BMC Biol, 19*(1),
1146   16. https://doi.org/10.1186/s12915-020-00946-6

1147

1148