1 **3D Reconstruction of Murine Mitochondria Exhibits Changes in Structure Across Aging**
2 **Linked to the MICOS Complex**
3

Zer Vue[*], Edgar Garza-Lopez[*], Kit Neikirk[*], Prasanna Katti[*], Larry Vang, Heather Beasley,
Jianqiang Shao, Andrea G. Marshall, Amber Crabtree, Alexandria C. Murphy, Brenita C.
Jenkins, Praveena Prasad, Chantell Evans, Brittany Taylor, Margaret Mungai, Mason Killion,
Dominique Stephens, Trace A. Christensen, Jacob Lam, Benjamin Rodriguez, Mark A. Phillips,
Nastaran Daneshgar, Ho-Jin Koh, Alice Koh, Jamaine Davis, Nina Devine, Saleem Muhammod,
Estevão Scudese, Kenneth Ryan Arnold, Valeria Vanessa Chavarin, Ryan Daniel Robinson,
Moumita Chakraborty, Jennifer A. Gaddy, Mariya Sweetwyne, Genesis Wilson, Elma Zaganjor,
James Kezos, Cristiana Dondi, Anilkumar K. Reddy, Brian Glancy, Annet Kirabo, Anita M.
Quintana, Dao-Fu Dai, Karen Ocorr, Sandra A. Murray, Steven M. Damo, Vernat Exil, Blake
Riggs, Bret C. Mobley, Jose A. Gomez, Melanie R. McReynolds[#], Antentor Hinton, Jr[#]
14
Affiliations:
Z.V., K.N., D.S., L.V., M.M.; M.K.; H.B., A.G.M., E.Z.; G.W.; J.A.G., A.H.J.;: Department of
Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN, 37232, USA
E.G., J.L., B.R.: Department of Internal Medicine, University of Iowa, Iowa City, IA, 52242,
USA
J.S.; Central Microscopy Research Facility, University of Iowa, Iowa City, IA, 52242, USA.
C.E.: Department of Cell Biology, Duke University School of Medicine, Durham, NC, 27708,
USA
B.T.; J. Crayton Pruitt Family Department of Biomedical Engineering, University of Florida,
Gainesville, FL, 32611, USA
T.A.C.: Microscopy and Cell Analysis Core Facility, Mayo Clinic, Rochester, MN, 55905, USA
M.A.P.; N.D.; M.C.: Department of Integrative Biology, Oregon State University, Corvallis, OR,
97331, USA
P.K.; B.G.: National Heart, Lung and Blood Institute, National Institutes of Health, 9000
Rockville Pike, Bethesda, MD 20892, USA.
B.G.: National Institute of Arthritis and Musculoskeletal and Skin Diseases, National Institutes
of Health, 9000 Rockville Pike, Bethesda, MD 20892, USA.
J.D.: Department of Biochemistry, Cancer Biology, Neuroscience, and Pharmacology, Meharry
Medical College, Nashville, TN 37208, USA
A.M.Q.: Department of Biological Sciences, Border Biomedical Research Center, University of
Texas at El Paso, El Paso, TX, 79968, USA
S.M.D.: Department of Life and Physical Sciences, Fisk University, Nashville, Tennessee,
37208, U.S.A.; Center for Structural Biology, Vanderbilt University, Nashville, Tennessee,
37232, U.S.A.
V.G.: Department of Cell and Developmental Biology, Vanderbilt University, Nashville, TN,
37232, USA
M.S.: Department of Laboratory Medicine and Pathology, University of Washington, Seattle,
WA, 98195, USA
E.S.: Laboratory of Biosciences of Human Motricity (LABIMH) of the Federal University of
State of Rio de Janeiro (UNIRIO), Rio de Janeiro, Brazil; Sport Sciences and Exercise
Laboratory (LaCEE), Catholic University of Petrópolis (UCP), Brazil

S.A.M: Department of Cell Biology, School of Medicine, University of Pittsburgh, Pittsburgh,
PA, 15260, USA
V.E.: Department of Pediatrics, Carver College of Medicine, University of Iowa, Iowa City, IA.
52242, USA
A.K.R.: Department of Medicine, Baylor College of Medicine, Houston, TX, 77001
V.E.: Department of Pediatrics, Div. of Cardiology, St. Louis University School of Medicine, St.
Louis, MO, 63104, USA
B.M.: Department of Pathology, Vanderbilt University Medical Center, Nashville, TN, 37232
USA
J.A.Gaddy: Department of Medicine, Vanderbilt University Medical Center, Nashville,
Tennessee, 37232, U.S.A.; Department of Medicine Health and Society, Vanderbilt University,
Nashville, Tennessee, 37232, U.S.A.; Department of Pathology, Microbiology and Immunology,
Vanderbilt University Medical Center, Nashville, 37232, Tennessee, U.S.A.; Department of
Veterans Affairs, Tennessee Valley Healthcare Systems, Nashville, Tennessee, 37212, U.S.A.
J.A.Gomez.; A.K.: Department of Medicine, Vanderbilt University Medical Center, Nashville,
TN, 37232, USA
D.D.: Department of Pathology, University of John Hopkins, Baltimore, MD, USA.
H.K.; A.K.; Department of Biological Sciences, Tennessee State University, Nashville, TN
37209, USA
M.R.M.;A.C.M.; B.C.J.; P.P.: Department of Biochemistry and Molecular Biology, The Huck
Institute of the Life Sciences, Pennsylvania State University, State College, PA 16801
J.K.; C.D.; K.O.: Sanford Burnham Prebys Medical Discovery Institute, La Jolla, CA 92037,
USA.
K.R.A.; V.V.C.; R.D.R.: Department of Ecology and Evolutionary Biology, University of
California at Irvine, Irvine, CA 92697
B.R.: Department of Biology at San Francisco State University, San Francisco, CA 94133, USA.
72
73 *These authors share co-first authorship.
74 #These authors share senior authorship.
75
76 Corresponding Author:
77 Antentor Hinton
78 Department of Molecular Physiology and Biophysics
79 Vanderbilt University
80 antentor.o.hinton.jr@vanderbilt.edu, 319-383-3095

81
82 **Graphical Abstract:**
83
ABSTRACT

Background: During aging, muscle gradually undergoes loss of function including sarcopenia,
losing mass, strength, endurance, and oxidative capacity. While mitochondrial aging is
associated with decreased mitochondrial capacity, the genes associated with morphological
changes in mitochondria during aging still require further elucidation. Furthermore, it is not
completely understood how 3D mitochondrial structures are altered during aging in skeletal
muscle and cardiac tissues.

Methods: We measured changes in mitochondrial morphology and mitochondrial complexity
during the aging of murine gastrocnemius, soleus, and cardiac tissues using serial block face-
scanning electron microscopy and 3D reconstruction. Lipidomic and metabolomic analysis
elucidated concomitant changes associated with aging. We also used qPCR, transmission
electron microscopy quantification, Seahorse Analyzer, and metabolomics to evaluate changes in
mitochondria morphology and function upon loss of the MICOS complex.
Results: We identified significant changes in 3D mitochondrial size and network configuration
in murine gastrocnemius, soleus, and cardiac tissue during aging. These changes were
concomitant with loss of mitochondria contact site and cristae organizing system (MICOS) gene
expression during aging. Mitochondrial morphology was similar between aged mice and young
mice. We show an age-related loss of the MICOS complex (Chchd3, chchd6, and Mitofilin)
while their knockout results in alterations in mitochondrial morphology. Given the critical role of
mitochondria in maintaining cellular metabolism, we perform cellular metabolic profiling of
young and aged tissues. Metabolomics and lipidomics showed profound alterations, including in
membrane integrity, that support our observations of age-related changes in these muscle tissues.
Discussion: In tandem, our data suggest a relationship between the MICOS complex and aging,
which could be linked to disease states with further 3D reconstruction studies. Our study
highlights the importance of understanding tissue-dependent 3D mitochondrial phenotypical
changes which occur across aging with evolutionary conservation between *Drosophila* and
murine models.
111
112

113 **Keywords**: MICOS, aging, mitochondria, 3D morphometry; mitochondrial disease;
114 mitochondrion; reconstruction; reticulum; serial block-face SEM; skeletal muscle, *Drosophila*.

115
116
117
INTRODUCTION
Sarcopenia is the loss of muscle mass associated with aging, which impacts the quality of life.
Sarcopenia predominately affects type II muscle fibers, but also affects type I fibers.
Mitochondrial dysfunction is a hallmark of aging[1]. With age, sarcopenia may occur in skeletal
muscle, leading to a body mass-independent loss of skeletal function[2]. Decreased expression of
genes associated with mitochondrial dynamics and loss of function is known to contribute to
sarcopenia and other age-related diseases [3]. A hallmark of aging is alterations in mitochondria
structure [1]. Thus, mitochondria are a key target for the development of potential therapeutics for
age-related pathologies [1,3]. Mitochondria maintain finely regulated structures, through various
mechanisms, including fission and fusion [4]. Because their structure changes dynamically as
mitochondria shift from fission to fusion, many important mitochondrial functions depend on
their structure[5–7]; therefore, it is important to understand mitochondrial structural changes over
time. Essential mitochondrial functions are associated with the cristae, the inner folds of the
mitochondrial membrane, which house the oxidative phosphorylation machinery[6] and various
transporters. Our objective was to determine how mitochondrial structures change during aging.
We hypothesized that age-related alterations in metabolism and lipids cause increased

mitochondrial fragmentation over time with concomitant losses in mitochondrial cristae
integrity.
The ultrastructure and phenotypical variation of mitochondria play important roles in
mitochondrial function. Disruption of optic atrophy 1 (OPA-1), an inner membrane protein that
regulates mitochondrial fusion, causes mitochondrial fragmentation and affects the dimensions,
shapes, and sizes of the cristae[6]. Furthermore, disruption of dynamin-related protein-1 (DRP1), a
protein associated with mitochondrial fission, causes elongated mitochondria and resistance to
cristae remodeling[8,9]. Nanotunnels, or "mitochondria-on-a-string", are thin, double-membrane
protrusions that allow mitochondria to communicate across distances. Nanotunnels may increase
in mitochondrial disease[10,11] and may be associated with mitochondrial dysfunction during aging.
Thus, the alterations in mitochondrial structure and bioenergetics may simultaneously drive
pathologies.
Mutations in genes that regulate the morphology of cristae are associated with aging
cardiomyocytes[12]. These proteins are located at the crista junctions in the inner membrane and
are a part of the mitochondrial contact site and cristae organizing system (MICOS) complex,
which is important for maintaining mitochondrial shape and size[13]. DRP1 or OPA-1 loss can
similarly affect mitochondria morphology[14,15]. Cristae membranes contain the electron transport
chain complexes and ATP synthases for oxidative phosphorylation[16–18]. Since mitochondrial
morphology affects function, altering the structure by knocking out MICOS-associated genes or
OPA-1, a GTPase, could alter the metabolism and function of mitochondria during aging[16–18].
We sought to better understand how the MICOS complex affects gross mitochondrial structure
beyond only cristae, as well as how mitochondrial structure changes in aging. Critically, we
predicted that MICOS-associated gene products are lost during aging; therefore, loss of MICOS-
associated genes would mimic the change in mitochondria morphology observed in aging.
Mitochondria have a tissue-dependent response to the environment [19], which may be due
to heterogeneity in mitochondrial DNA (mtDNA) quality check mechanisms across tissues [20].
Using manual contour tracings, 3D reconstruction is an important tool to understand specific
phenotypes of mitochondria as well as how they may differ across tissue types. Therefore, we
sought to better understand the alterations of mitochondrial structure in aging; we compared size,
shape, quantity, complexity, and branching of mitochondria using 3D reconstructions of aged
gastrocnemius, soleus, and cardiac tissue in 3-month and 2-year mice. We observed an age-
related loss of the MICOS complex in both qPCR and western blotting. Beyond this, we used
CRISPR/Cas9 on myotubes to knockout (KO) three genes of the MICOS complex: *Chchd3*
(Mic19), *Chchd6* (Mic25), and *Mitofilin* (Mic60) to understand how the MICOS complex may
present phenotypically similar to the aging process through modulation of mitochondrial size,
morphology, and oxygen consumption rate. To further understand factors that may affect
mitochondrial structure in aging, multivariate analysis was used to identify changes in
metabolites and characterize the lipidome across aging to understand commonalities to metabolic
changes with loss of the MICOS complex. Finally, a *Drosophila* model was used to further
explore the role of the MICOS complex beyond murine and to elucidate changes in heart
function and MICOS gene expression in aging.
175
RESULTS
Aging Results in Smaller Mitochondria in Murine Gastrocnemius, Soleus, and Cardiac
Tissue

The gastrocnemius is a mixed muscle with both oxidative fibers, containing many
mitochondria, and glycolytic fibers, with few mitochondria [21]. This heterogeneity makes the
gastrocnemius ideal to study changes in mitochondrial dynamics. In contrast, soleus tissue is
made up predominately of slow-twitch muscle fibers, reliant on oxidative metabolism dependent
on the relatively increased mitochondrial content [22]. Mitochondria in cardiac tissue, as we have
previously studied [23], further have differential roles more reliant on efficient energy transfer to
myofibrils and constant ATP production to maintain contractile function [24]. Since many
mitochondrial functions depend on structure [5–7], it is important to examine how mitochondrial
structure changes over time. We hypothesized that increasing mitochondrial fragmentation over
time is concomitant with losses in mitochondrial crista integrity.
Our objective was to determine how aging alters mitochondrial networks and individual
mitochondrial structures. We imaged gastrocnemius, soleus, and cardiac biopsies from
adolescent (3-month-old) and aged (2-year-old) mice by serial block-face scanning electron
microscopy (SBF-SEM) with resolutions of 10 nm for the x- and y- planes and 50 nm for the z-
plane, which allows visualization of the electron connectome. Approximately 50
intermyofibrillar (IMF) mitochondria were segmented from each image stack (Figure 1A-F) and
a 3D surface view was generated (Figure 1A'-F'). We analyzed IMF, as opposed to other
mitochondrial subpopulations such as subsarcolemmal, as IMF mitochondria are larger and
displayed more significant age-related changes. We analyzed mitochondria sub-network volumes
from four ROIs with an average of 175 mitochondria for each mouse (n = 3), for a total of 500
mitochondria. Mitochondrial networks in aged skeletal muscle tissue mice showed largely
interconnected mitochondria that were not composed of a single reticulum (Figures 1A''–D'').
As in our previous study, we found cardiac tissue mitochondria remained relatively clumped
with no apparent distribution changes (Figure 1E"-F"). We found that across all tissue types,
volume, area, and perimeter of mitochondria of samples from 2-year-old mice were significantly
lower than those from 3-month-old mice (Figures 1G-X). Here mitochondria volume is a
measure of their total capacity (Figure 1Y), while area represents a measure analogous to surface
area (Figure 1Z) and perimeter represents boundary pixel count (Figure 1AA).
While there was some variability across the three mice for each age cohort
(Supplemental Figure 1), this heterogeneity is more pronounced in the gastrocnemius muscle
(Figure 1G-H). Gastrocnemius also showed a greater reduction in mitochondrial size and surface
area, demonstrating much smaller mitochondria no longer presenting as hyperbranched. Soleus
tissue similarly demonstrated heterogeneity between the samples with similar reductions in
mitochondrial volume across aging (Figure 1M-R). In contrast to skeletal muscle, cardiac tissue
shows a much more homogenous mitochondrial population (Figure 1S-X). In general, the trend
showed a downward trajectory emblematic of increased fragmentation and smaller mitochondria.
This showed that the size and length of mitochondria change with age, so we sought to further
elucidate the complexity of mitochondria, which is implicated in mitochondria communication.
217
218 **Aging Causes Altered Mitochondrial Phenotypes with Poorly Connected Mitochondria**
219 **with Decreased Branching in Murine Gastrocnemius, Soleus, and Cardiac Tissue**
220
Next, we measured mitochondrial complexity to better understand changes in mitochondrial
shape during aging. We hypothesized that fewer networks and simpler shapes would occur as
aging and dysfunction continued. Since we observed in 3D reconstructions that mitochondrial
populations are heterogeneous and diverse, we used mito-otyping, which is the karyotyping-like

method for arranging mitochondria[25], to capture the diversity of IMF mitochondria (Figure 2A-
C). This analysis suggested, at every volumetric measurement, that mitochondria were smaller
and less complex with age. Soleus and gastrocnemius tissue both showed branched distributions
in the adolescent populations while aged populations appeared much less branched, in addition to
having smaller volumes. Notably, skeletal muscle showed large phenotypical changes with aging
while cardiac tissue does not show the same changes. To validate these changes, we utilized
previously established methods were used to determine 3D mitochondrial complexity from 3D
form factor measurements[25,26]. We measured mitochondrial complexity index (MCI) and
sphericity to further understand changes in complexity (Figure 2D-O). MCI and sphericity are
analogous measures to understand the general roundness of mitochondria (Figure 2P-Q). In
gastrocnemius tissue, there was an increase in MCI concomitant with an increase in sphericity
with aging (Figure 2D-G). This shows that, contrary to the appearance of mito-otyping,
gastrocnemius exhibits increased sphericity of mitochondria. When examining the specific three
male mice sampled for each age cohort, some variation exists for both metrics. Soleus tissue,
which shows a similar heterogeneity, is exposed to be less complex and more spherical across
aging (Figure 2H-K) although this is not as significant of a difference as that exhibited in
gastrocnemius. Finally, cardiac tissue only shows a significant increase in MCI (Figure 2L-O).
Together, these data suggest that complexity is altered with age, but the specific way in which
complexity changes is dependent on tissue type. These findings caused us to look at the MICOS
complex to elucidate the role it has on mitochondrial structure and function, as a potential
effector of age-related changes in mitochondrial structure.
246
Age-Related Changes in the MICOS Complex Expression
The MICOS complex and OPA-1 are key players in mitochondrial biogenesis [13,17,27], but
how their interactions regulate aging and mitochondrial structures is poorly understood. Since
we observed mitochondrial dynamics change with aging, we measured transcripts for *Opa1*,
Chchd3 (*Mic19*), *Chchd6* (*Mic25*), and *Mitofilin* (*Mic60*), the genes for the principal MICOS
complex subunits. We used reverse-transcriptase qPCR (RT-qPCR) to determine the effect of
aging on transcription. Aging causes mitochondrial fragmentation [1], which is associated with the
loss of *Opa1* and MICOS complex proteins [17,28,29]. We hypothesized that aging would also lead
to decreases in the associated MICOS complex proteins. We also sought to confirm prior studies
that found *Opa1* loss across aging, by using *Opa1* as a positive control [17,30–32]. We found a
significant loss of both *Opa1* and MICOS complex subunits, as measured by loss of transcripts
for the genes in skeletal muscle when comparing 3-month and 2-year samples (Figure 3A–D).
When looking at cardiac tissue (Figure 3E-H) we noticed a similar age-related loss of all
MICOS genes products. These data suggest a correlation between mitochondrial morphological
changes and decreases in expression of *Opa1* and MICOS subunits *Chchd3*, *Chchd6*, and
Mitofilin during aging; however, they are not sufficient to demonstrate causation, therefore we
did additional measurements to test if reduced protein expression levels may be responsible for
the morphological changes. When looking at the MTCO1 (Cox-1) or Sam50 levels, there were
not significant alterations across the aging process (Figure 3I-K). However, both gastrocnemius
and skeletal tissue showed reduced protein levels of Mitofilin and OPA1 (Figure 3I-J); notably,
however, there was a differential aging response with no age-related changes in protein levels in
cardiac tissue (Figure 3K).
From there we sought to validate these results in another model. Using qPCR, we looked
at differences in gene expression for MICOS complex as well as mitochondrial endoplasmic

reticulum contact (MERC) genes between experimentally evolved Drosophila populations
subjected to hundreds of generations for accelerated aging and their controls [33]. We looked at
MERC genes as the relationship between the MICOS complex and MERCs remains unclear, and
we believed there may be concomitant loss. Specifically, we compared patterns between 21-day-
old flies, with the accelerated aging flies displaying phenotypes of old age, where past studies
have shown there are large differences in age-specific mortality, gene expression profiles, and
metabolomic profiles [33–35]. Flight muscle tissues, ATF-4, Drp, Marf, Opa1, MINOS1 (mic10),
CHCHD3/6(mic19), QIL1(mic13), APOO (mic16/27), IP3R/Itpr, VDAC/Porin, Bip(grp78),
GRP-75/Hsc70-5, and Ire1 all showed downregulated expression in indirect flight muscle of
aged flies when compared with controls (Table 1; Figure 3L). However, only the expression
changes associated with Drp, Marf, Opa1, CHCHD3/6, QIL1, APOO, IP3R, VDAC, and Bip
were significant (p-value < 0.05; Unpaired T-test) (Table 1; Figure 3L). We saw a slight
upregulated expression of ATF-6 and Mitofilin in the aged flies, however, these changes were
not significant. ATF-4, Opa1, MINOS1, and Mitofilin all showed downregulated expression in
the cardiac tissues of the aged flies when compared with controls, however, only the change in
ATF-4 and Mitofilin expression were significant (Table 2; Figure 3M). In contrast, there was a
significant increase in FGF-21 expression in the aged flies (Table 2). These validate that
mitochondrial, MICOS, and MERC proteins are differentially impacted in cardiac and skeletal
muscle tissue. Interestingly, together these also suggest that in some models MICOS complex
has a greater dependence on the aging process.
291
2D and 3D Structural Changes in Cristae and Mitochondria Exhibited Upon Loss of
MICOS Complex and *Opa1*
To determine the role of OPA-1 and the MICOS complex in mitochondrial structure and
networking, we ablated the genes for *Opa1* and the MICOS complex proteins in isolated primary
skeletal muscle cells from 3-month-old mice. We isolated primary satellite cells, then
differentiated myoblasts into myotubes. Using CRISPR/Cas9 method and a control plasmid, we
knocked out the genes for MICOS complex components and *Opa1* from skeletal muscle cells.
We measured 1250 mitochondria across 10 cells, with loss of *Opa1* as a positive control
for mitochondrial morphological changes, since it is well understood that *in vitro* deletion of
Opa1 alters mitochondrial morphology [14,36,37]. Although *Opa1* expression decreases with age[38],
how loss of the MICOS complex affects mitochondria 3D morphology is poorly elucidated. Past
studies have indicated knockout of MICOS subunit *Chchd3* results in fragmented mitochondria
as the cristae lose their normal structure[39]. Similarly, *Chchd6* is important in maintaining crista
structure and its downregulation results in hollow cristae that lack an electron-dense matrix,
thereby inhibiting ATP production and cell growth[40–42]. Using transmission electron microscopy
(TEM), we compared mitochondria and cristae in myotubes from wild-type (WT) and knockouts
of *Opa1-*, *Mitofilin-, Chchd3-*, and *Chchd6*-knockout myotubes, which are essential for the
organization of mitochondrial cristae[41,43] (Figures 4A–E). Mitochondrial average area decreased
for *Opa1-*, *Mitofilin-, Chchd3-*, and *Chchd6*-knockout myotubes (Figure 4F), while the
mitochondrial circularity index (the roundness and symmetry of mitochondria) and the number
of mitochondria, once normalized, increased (Figures 4G–H). This suggests that mitochondria
become less complex, smaller, and more abundant upon loss of the MICOS complex. For *Opa1-*,
Chchd3-, *Mitofilin-*, and *Chchd6*-knockouts, the number of cristae per mitochondrion decreased,
as did the cristae score and cristae surface area compared with the WT (Figures 4I–K). Here
cristae score is defined by the following:

0: No sharply defined cristae are visible.
1: Over 50% of the mitochondrial area lacks cristae.
2: Over 25% of the mitochondrial area lacks cristae.
3: Irregularly shaped cristae are visible, less than 25% of mitochondria lack cristae. over
75% of the mitochondrial area.
4: Many regular-shaped cristae are visible, less than 25% of mitochondria lack cristae.
While the loss of the MICOS complex shows similar changes across all of these
knockouts, in general, the least significant change was seen in the *Chchd3* knockout. Together,
these data show quantitative and structural changes in both mitochondria and cristae upon loss of
MICOS proteins.
TEM provides cristae detail, but not 3D morphology; therefore, we used SBF-SEM to
look at the 3D structure of the mitochondria. Using myotubes with ablated genes for *Opa1* and
MICOS complex subunits, as described above, we measured a total of 200 mitochondria across
10 cells. We compared mitochondria in WT, *Opa1*-, *Mitofilin-, Chchd3*-, and *Chchd6*-knockout
myotubes (Figures 4L-P). We found that compared with the elongated mitochondria in the WT,
the 3D length was much shorter in *Opa1* and MICOS protein knockouts (Figure 4Q). Similarly,
the volume of mitochondria was also reduced in *Chchd3-*, *Chchd6-*, *Mitofilin-*, and *Opa1-*
knockouts compared with the control (Figure 4R). The 3D reconstruction data, in combination
with the prior TEM results, show how mitochondrial dynamics change with the loss of MICOS
subunits and mimic the mitochondrial phenotypes observed across aging.
We also wanted to consider how loss of the MICOS complex and other mitochondrial
genes impacted mitochondrial structure in a *Drosophila* model. Since QIL1 (Mic13) and
CHCHD3/6 (mic19) were all specifically downregulated across the aging process in skeletal
tissue, we sought to understand their role in *Drosophila* mitochondrial structure. We also
considered how the loss of Mitofilin, which was slightly upregulated in aging, changed
mitochondrial structure. Finally, we also looked at standard mitochondrial fusion proteins of
OPA1 and MARF and the fission protein DRP1. Mitochondrial-actin straining showed
differences in mitochondrial organization and relative Myofibrillar density (Figure 4S). From
there, we utilized TEM to understand how *in vivo* flight muscle mitochondrial morphology was
altered upon these KDs. As expected, loss of DRP1 caused increases in the mitochondrial area
while inversely loss of MARF and OPA1 both reduced mitochondrial area (Figure 4T). Loss of
all MICOS complex proteins (Mic13, Mic19, and Mic60) all showed reductions in mitochondrial
volume, although this reduction was less severe in Mic13 (Figure 4T). Looking at circularity to
understand mitochondrial complexity, we notice that interestingly loss of the MICOS complex
paralleled the knockout of DRP1 marked by a decrease in circularity (Figure 4U). In contrast,
loss of mitochondrial fusion proteins increased circularity of mitochondria. Finally, for Mic13,
Mic19, and Mic60 we notably show that while the loss of Mic13 and Mic19 reduces
mitochondrial quantity, yet loss of Mic60 increases mitochondrial quantity (Figure 4V). This
suggests that loss of gene expressions of the MICOS complex and related mitochondrial
structure may be evolutionarily conserved to some degree yet there may be certain organismal-
dependent alterations in associated mitochondrial dynamics and age-related gene expression
reduction.
359
360
Changes in Oxygen Respiration Rate and Metabolite Changes Upon Loss of MICOS
Complex and *Opa1*

To understand the functional impact of these structural remodeling of mitochondria upon
loss of MICOS complex, we used an XF24 seahorse analyzer to measure oxygen consumption
rate (OCR). Past studies have indicated that loss of *Opa1* induces bioenergetic stress and
decreased electron transport chain function[37], and ablation of the MICOS complex alters
mitochondrial capacity[44,45]. We found that loss of *Opa1* or *Mitofilin* in myotubes decreased basal
OCR (Figure 5A–B) and decreased ATP-linked, maximum, and reserve capacity OCR (Figure
5C–E). Although *Opa1*-knockout myotubes exhibited a decrease in proton leak, which represents
protons that go from the mitochondria to the matrix without producing ATP, *Mitofilin* knockouts
showed no significant difference (Figure 5F). To determine the global effects of loss of *Opa1* or
the MICOS complex in skeletal muscle myotubes, we analyzed the metabolome to identify
changes in metabolites that occurred with changes in mitochondria and cristae. Principal
component analysis (PCA) revealed distinct populations in the control and the *Mitofilin*-
knockout strains, which suggested that their genotypes contributed to the clustering (Figure 5G).
To identify metabolites with the most robust ability to discriminate between true positives and
false positives, we constructed a model using analysis of variance (ANOVA) to determine
several key metabolites that were statistically significant (Figure 5H). This unique metabolite
signature revealed that *Mitofilin* plays a critical role in regulating amino acid metabolism and
steroidogenesis (Figure 5I–J). Upregulation of steroidogenesis pathways may result from the
increased fluidity of membranes caused by *Mitofilin*[46,47].
In *Opa1*-, *Chchd3*-, and *Chchd6*-knockouts, there was a decrease in basal, ATP-linked,
maximum, and reserve capacity OCR compared with the control (Figures 5K-O). Although
proton leak OCR decreased in *Opa1*- and *Chchd3*-knockout myotubes (Figure 5P), there was no
significant difference between the control and *Chchd6*. The decrease in OCR may be attributed
to smaller and fragmented mitochondria; mitochondrial density decreases as fragmentation
targets them for autophagy[38,48]. Together, these results show that MICOS and *Opa1* are essential
for the normal respiration of muscle tissue. We also measured the effect of ablating the genes for
Chchd3 and *Chchd6* in skeletal muscle myotubes on bioenergetic metabolism. PCA revealed
distinct populations in the control and the *Chchd3*- and *Chchd6*-knockouts, which showed a
similar profile to that we observed in *Mitofilin* (Figure 5Q). We constructed a model using
ANOVA to determine which metabolite changes in *Chchd3* and *Chchd6* knockouts were
statistically significant (Figure 5R). There was a loss of protein synthesis and changes in
carbohydrate metabolism (Figure 5S–T). Loss of *Opa1* typically favors fatty acid synthesis, so
the results showing increased carbohydrate metabolism differ from previous *Opa1*-knockout
responses[49–51]. This atypical response was evident in the increase in lactose and starch synthesis,
but there was poor protein turnover, as seen in methionine metabolism (Figure 5T). Given that
we saw a loss of MICOS complex proteins caused a change in metabolism, we sought to see if
this change in metabolism parallels the age-related change in gastrocnemius, soleus, and cardiac
metabolism.
401
Metabolomics/Lipidomic Show that Across Aging, Gastrocnemius, Soleus, and Cardiac
Tissue Exhibit Altered Metabolism Across Aging
Together, our results up to this point suggested an age-related change in the MICOS
complex with implications for structural and functional alterations in the MICOS complex.
Given we saw a loss of the MICOS complex was implicated in altered steroidogenesis and
metabolism, we examined if similar pathway alterations are noted in aging to explicate future
areas of study.

To investigate the factors influencing the observed changes in aged tissues, we conducted
metabolomics and lipidomics analyses on both young and aged muscle tissues (Figure 6A-F).
Our analysis revealed significant metabolic shifts in all three tissue types. These changes
encompassed various processes, including NAD^+ metabolism, linolenic metabolism, porphyrin
synthesis, heme biosynthesis, and glycine and lysine metabolism (SFigure 3D-I). Notably, we
observed an accumulation of cholic acid in aged soleus and gastrocnemius muscles (SFigure 3A,
B, D, and E), a known inducer of muscle atrophy [52]. Amino acid metabolism was dysregulated
across all muscle tissues.
417
We conducted lipidomic profiling to investigate the role of dysregulated lipid metabolism
in muscle tissues during aging (Figure 6 B, D, F; STables 1-4). Our analysis revealed notable
changes in both lipid classes (SFigure 3 J-L) and lipid chain lengths (SFigure 3 M-O) with age.
Among the three muscle types, the gastrocnemius and cardiac tissues exhibited the most
pronounced alterations in lipid classes, while the soleus showed minimal dysregulation (SFigure
3 J-L). Additionally, the cardiac muscle displayed a higher degree of disrupted lipid chain
lengths compared to the soleus and gastrocnemius muscles (SFigure 3M-O). These findings
highlight the importance of altered lipid chain lengths in cellular and physiological processes, as
they can affect membrane integrity, fluidity, and functionality, thereby influencing a wide range
of cellular functions.
428
The gastrocnemius muscle exhibited dynamic changes in lipids with age (Figure 6B). These
alterations affected various lipid classes, including triglycerides oligomers (TGO), triglycerides
(TG), phosphatidylserine (PS), phosphatidylglycerol (PG), ceramide (Cer), and acylcarnitine
(CAR). In the gastrocnemius, the levels of PS, the most abundant negatively charged
phospholipid in eukaryotic membranes and enriched at cell-cell contact regions [53], decreased
with age (SFigure 3J). Acylcarnitine, which plays a crucial role in transporting acyl groups from
the cytosol into the mitochondrial matrix for beta-oxidation [54], was significantly reduced in aged
gastrocnemius tissues (SFigure 3J). Conversely, ceramide levels increased with age in these
tissues (SFigure 3J). This observation is supported by previous reports of elevated ceramide
levels in replicative senescent cells, with ceramide proposed to contribute to senescence by
inducing cell cycle arrest [55].
440
In the aged soleus tissues, we observed alterations in only four lipid classes (SFigure 3K). These
classes included triglycerides (TG), sulfatide hexosylceramide (SHexCer), monoradylglycerol
(MG), and hexosylceramide (HexCer). Notably, both HexCer and SHexCer exhibited depletion
with age in the soleus (SFigure 3K). These lipids belong to the sphingolipid class, suggesting a
novel role for sphingolipids in the aging process of the soleus. Additionally, MGs were found to
accumulate in the aged soleus (SFigure 3K). These lipids have been linked to lipotoxicity, which
triggers immunesenescence [56]. Lastly, TGs, which are crucial for storing and transporting fatty
acids within cells and in the circulation, showed a decrease in the aged soleus (SFigure 3K). This
finding aligns with previous studies reporting decreased TG levels in plasma and increased levels
of fatty acids during aging [57].
451
The cardiac muscle displayed age-related alterations in several lipid classes, including
triglycerides oligo (TGO), phophatidyl glycerol (PG), lysophosphatidylcholines (LPC),
cardiolipin (CL), cholesteryl esters (CE), acyl carnitine (CAR), and

bis(monoacylglycerol)phosphate (BMP) (SFigure 3L). Notably, cardiolipins, which play a
crucial role in regulating mitochondrial proteins and maintaining mitochondrial structures such
as cristae and contact sites [58], exhibited a decrease in aged cardiac muscles (SFigure 3L). This
lipid class has been implicated in age-related alterations in mitochondrial bioenergetics [58–60].
Furthermore, we observed an accumulation of TGO in the heart, which has been linked to
inflammation, endothelial dysfunction, oxidative stress, atherosclerotic plaques, and ultimately
cardiovascular disease (SFigure 3L) [61]. TGOs are closely associated with the aging process [57].
462
Returning to metabolic changes, we detected an accumulation of Nicotinic Acid Riboside
(NaR) in the soleus, cardiac, and gastrocnemius tissues (Figure 6G), indicating potential
compensatory mechanisms to maintain NAD^+ levels, which are known to decline with age in
muscle tissues [62]. Furthermore, the levels of linoleic acid, an essential fatty acid crucial for cell
membrane integrity and the synthesis of inflammation-related eicosanoids, increased with age in
these three tissues (Figure 6H). In our metabolic profiling, aminolevulinic acid stood out as a
depleted metabolite in the cardiac, soleus, and gastrocnemius tissues of aged mice (Figure 6I).
Aminolevulinic acid is essential for heme production, a process vital for mitochondrial oxygen
and energy production [63]. Collectively, these profound metabolic alterations support our
observations of age-related changes in these muscle tissues.
473
474   In summary, our metabolic analysis and lipid profiling unveiled significant metabolic
475   alterations in cardiac, soleus, and gastrocnemius muscles with age, some of which may parallel
476   the metabolic alterations seen with the loss of the MICOS complex.
477
478
DISCUSSION
We demonstrated that either aging or loss of MICOS proteins in skeletal muscle resulted
in suboptimal mitochondria morphology in a tissue-dependent manner, suggesting a potential
correlation between aging and MICOS protein expression. Previous studies used 3D focused ion
beam-scanning electron microscopy (FIB-SEM) to characterize the networking of the
mitochondria in human[64] and mouse skeletal muscle[65]. Another report did quantitative 3D
reconstructions using SBF-SEM to define the morphological differences in the skeletal muscles
of humans versus mice, and they compared patients with primary mitochondrial DNA diseases
with healthy controls[25]. To the best of our knowledge, our current study is the first to use a 3D
reconstruction method as an approach to studying mitochondria phenotypes in skeletal muscle
aging. We prioritized using manual contour tracing as opposed to machine learning techniques to
ensure the accuracy of these highly varied mitochondrial phenotypes. Future research may
consider looking further at human skeletal muscle, to better compare if human gastrocnemius
and soleus muscles have a similar phenotype to murine skeletal muscle. While the murine soleus
is understood to have a similar transcriptome to that of humans [66], further elucidation of these
similarities are necessary. Furthermore, considering that oxidative muscle types vary in function
and structure, future studies may further seek to better discern them.
Skeletal muscle is highly mitochondrial-dependent, as well as mitochondria enriched,
which comprise ~6% of the cell volume and changes with aging[67]. Gastrocnemius muscle has
both type I slow-twitch muscle fibers and type II fast-twitch muscle fibers; type I fibers are more
effective for endurance, while type II fibers better support short bursts of muscle activity[21,67,68].
Of relevance, sarcopenia is characterized by both types reducing in frequency and size while

type II fibers transition to type I [69]. In contrast, other forms of muscle degeneration such as
disuse atrophy does not see a change in fiber number and an inverse shift from type I fibers to
type II [69]. This suggests it is possible that age-related changes in mitochondrial shape may be due
to sarcopenia-dependent alterations in fiber frequency, which may have different mitochondrial
phenotypes. However, our 3D morphologic data did not permit discrimination of the two fiber
types, although we observed many variable muscle fibers within a sample and between all
differently aged cohorts. Past studies in human skeletal muscle separated these by taking two
portions of samples and assessing fiber type via SDS-PAGE[70]. Importantly, past studies have
found rather large differences in mitochondrial content and morphology in chicken muscle fiber
subtypes[71]. In a chicken model, past literature has found that type I does not contain lipid
droplets, and used the presence of lipid droplets to identify type II fibers [71,72]. However, to our
knowledge, this technique has not yet been applied to the 3D reconstruction of aged murine
skeletal muscle models.
Using 3D reconstructions, we found that all parameters of the mitochondria changed
during aging, including smaller volume, area, and perimeter (Figure 1A-X), and the
mitochondria became less interconnected in gastrocnemius while other tissue types saw a less
significant decrease in volume. This increased fragmentation suggests a decreased mitochondrial
fusion, in association with age-dependent decreases of OPA-1, which is a regulator of
mitochondrial fusion. We also saw a decrease in the MCI during aging, suggesting that
mitochondria decrease in networking, but their shape may not radically change as they increase
in sphericity as they age (Figure 2).
MICOS proteins play key regulatory roles in mitochondrial structure and function[17,27,48].
We determined by TEM 3D reconstructions that the loss of *Mitofilin*, *Chchd3*, and *Chchd6*
resulted in fragmentation, disrupted cristae, and smaller mitochondria (Figure 4), similar to the
loss of *Opa1*, which is known to cause changes in oxidative phosphorylation [17,30,37]. Overall,
mitochondria lacking the MICOS genes had characteristics similar to those of aged mouse
skeletal muscle (Figures 1–2). This similarity in phenotype suggests there may be an association.
Thus, changes in mitochondrial morphology due to aging may be caused by a lack of MICOS
protein expression. This is supported by the RT-qPCR data that showed decreased *Chchd3*,
Chchd6, *Mitofilin*, and *Opa1* transcription with aging (Figure 3). Notably, however, while both
tissue types exhibit diminished mRNA transcripts, only skeletal muscle shows diminished
protein levels of the MICOS complex and OPA1 (Figure 3). This may be a cell-specific
response, exhibiting down-regulated proteins in cardiomyocytes but more abundantly expressed
proteins in fibroblasts [73]. This may also represent a compensatory mechanism, where the half-life
of the protein is much longer in cardiac tissue to buffer against increased loss of mRNA
transcripts. We also note that there are differential metabolomics and lipidomics across different
tissue types, which suggests that this MICOS-complex-dependent response to aging may differ
across cell types due to altered metabolomics. In any case, the tissue-specific loss of the MICOS
complex must be further elucidated.
In *Drosophila*, since *Mitofilin* is slightly upregulated (Figure 3L), its expression may act
as a counteractive mechanism relative to the loss of other MICOS complex proteins across the
aging process. Notably, in flies, we also show that *Mitofilin* loss increases mitochondrial quantity
conversely to the loss of other MICOS complex components. This suggests that the roles of
Mitofilin may be differential in Drosophila and murine models, highlighting the need for
additional studies on *Mitofilin*. Furthermore, the differential roles of the MICOS complex
proteins need further elucidation. While there are commonalities in phenotypes between all

MICOS complex components when knocked down, we see that loss of *Mitofilin* and *Chchd6* do
not have the same reduction in proton leak as *Chchd3* KD (Figure 5F, P). While past studies
have indicated that all three of these components are necessary for the stabilization of the
MICOS and SAM complex, and consequently cristae architecture [27,39–41,74], our findings suggest
greater research is necessary to understand if compensation for specific components of the
complex avoids dysfunctional phenotypes upon loss of certain MICOS complex components.
Although there is a link between aging and loss of *Opa1* [31,38], little is known about the
role of the MICOS complex in aging. Changes in mitochondrial architecture and loss of integrity
may be caused by decreased MICOS proteins; thus, it will be important to develop methods to
restore MICOS proteins and *Opa1* lost during aging to decrease the deleterious effects of
mitochondrial dysfunction. Although studies have examined the role of knockouts on
mitochondrial dynamics, few studies have examined how the loss of MICOS proteins may be
restored to mitochondria[30,75]. To better elucidate potential future studies, we sought to consider
the similarities in metabolomics and lipidomics in loss of the MICOS complex and the aging
process.
We noticed differential changes in metabolomics and lipids both in comparison between
skeletal and cardiac tissue, but even between soleus and gastrocnemius (Figure 6). This
highlights the importance of considering tissue-dependent differences, which may arise due to
the tissue-dependent mitochondrial structures we observed. Metabolomics showed that several
pathways acted to cause muscular dystrophy while other pathways such as Nicotinic Acid
Riboside acted to rescue these pathways. Cardiac tissue shows the largest change in lipid
alterations across aging, which suggests that lipid accumulation may play a larger role in cardiac,
which matches prior literature which shows that lipid metabolism is closely linked with
pathology [76]. Conversely, soleus shows a Sphingolipid decrease, which may show that the soleus
age due to lipo-toxicity. Sphingolipid also is associated with MERC regions during and prior to
apoptosis [77]. This highlights the importance of future studies examining mitochondria-lipid
droplet associations, or interactions with ER [78] to see if these contact sites change across the
aging process to protect against lipotoxicity, as well as experiments employing mass
spectrometry imaging to better understand the spatial distribution of these metabolites across
aging [79]. In gastrocnemius tissue, we observe numerous phospholipids decrease (Figure 6). We
also saw a loss of linoleic acid, which is important for membrane integrity [80]. This suggests
altered mitochondrial membrane viscosity across the aging process. Past findings have also
observed age-dependent increases in viscosity which affect oxidative phosphorylation through
modulation of supercomplexes performance [81]. Notably, decreased membrane viscosity occurs in
age-related pathologies including Alzheimer's Disease, highlighting that these changes may have
implications in pathomechanisms [82]. One potential target is the MICOS complex, as past studies
highlight that lipids, including cardiolipin, can interact with cristae [83]. While we only see a
change in cardiolipins in cardiac tissue, it is possible that since this region has the least drastic
change in mitochondrial structure, cardiolipins protect against MICOS-dependent loss in
structure. It is possible that when you lose cristae morphology, you may increase in cardiolipins.
In conclusion, we can conclude that aging may also be due to alterations in membranes mediated
by lipids, affecting the MICOS complex and modulating mitochondrial structure and function.
Some of these age-related lipidomic and metabolomic changes may be due age-
dependent alterations in the MICOS complex. Many studies have analyzed the mitochondrial
metabolome using mouse skeletal muscles[67,84–88]. We found that loss of *Mitofilin* affected cristae
morphology (Figures 4I–K), decreased oxidative phosphorylation (Figure 5A), and may have

increased lipid and steroid synthesis, which may be important for MERCs regulation and cristae
formation. We found an increase in tryptophan and methyl histidine metabolism (Figure 5J) and
an increase in taurine metabolism and hypotaurine, a key sulfur-containing amino acid for fat
metabolism. Loss of *Opa1* also changes amino acid and lipid metabolism, similar to the loss of
Mitofilin[49–51]. Steroidogenesis, which makes the membrane less rigid, was increased. Since the
loss of *Mitofilin*, *Chchd6*, or *Chchd3* caused a loss of oxidative capacity (Figure 5A-F; 5K-P),
increased steroid synthesis may allow the cell to recover bioenergetic functions, as steroid
compounds decrease membrane viscosity, with the incorporation of estrogen[46]. In the absence of
Mitofilin, cristae junctions and contact sites fall apart[89]; thus, *Mitofilin* is critical for maintaining
cristae[90,91]. Cells lacking *Mitofilin* may make steroids to help the membrane to reconfigure
broken cristae. Since the loss of *Opa1* causes more MERCs[92], loss of *Mitofilin* may increase
phospholipids (Figure 5J) because of increased smooth MERCs, which are associated with lipid
changes[93]. This is supported by the fact that biosynthesis of phosphatidylethanolamine and
phosphatidylcholine, and metabolism of arachidonic acid and sphingolipid, increased with loss
of *Mitofilin* (Figure 5J). Since these phospholipids aggregate around MERCs and may shuffle
into the ER, *Mitofilin* may be a critical gene for regulating cristae morphology, with a key, novel
role in regulating mitochondrial metabolism.
Mitofilin may be an important target in the future to restore energy production. Loss of
Mitofilin may lead to ER stress, which, via ATF4, activates amino acid transporters[94] that further
activate mTORC1. ER stress activates mTORC as a result of a decrease in glucose[88]. Critically,
mTORC1 affects glucose homeostasis[95], which may lead to inefficient energy use. This can
result in changes in autophagy. Therefore, as a downstream effect of *Mitofilin* loss increasing
mTORC1, this may explain why deletion of MICOS in *Drosophila* increases autophagy[48].
Similarly, loss of *Opa1* increases ER stress[37], and loss of *Mitofilin* may cause a similar increase
in amino acid catabolism. If ER stress activates amino acid transporters, branched-chain amino
acids could increase ER stress, resulting in a positive feedback loop that affects the health of the
cell, cellular energy, metabolism, and antioxidants. ER stress may also be responsible for the
poor performance and fragmentation of mitochondria (Figure 4-5). Loss of *Mitofilin* may result
in the breakdown of protein pathways that regulate ER stress. Other amino acid pathways, such
as homocysteine (Figure 5J), are involved in triglyceride uptake and increased intracellular
cholesterol, suggesting that proteins like ATF4[88] and the MICOS complex[13,27] are important for
aging. In particular, the MICOS components may prevent mitochondrial fragmentation by
blocking ER stress pathways in aging. We notably show that several MERC proteins are
differentially regulated concomitantly with MICOS complex proteins across the aging process in
Drosophila (Tables 1-2). Further studies are needed to better understand the role of MICOS in
MERC formation and the relationship between smooth MERC and lipid synthesis.
Although *Mitofilin* is a key component of the MICOS complex, other components are
likely also important. The loss of *Chchd3* or *Chchd6* leads to a decrease in and disassembly of all
Mitofilin subcomplex components in mammals, with abnormal cristae morphology and growth
defects[28,29,40,74,96–98]. Downregulation of *Chchd3* is linked to type 2 diabetes[99]. In our
metabolomics enrichment dataset (Figure 6T), loss of *Chchd3* or *Chchd6* in mouse myotubes
resulted in a preference for alternative energy sources, such as lactate, lactose, and starches.
Supplementation of healthy myotubes with galactose leads to a 30% increase in oxidative
capacity (i.e., OCR) due to an increase in AMPK phosphorylation and cytochrome c oxidase
(COX) activity, thereby forcing cells to become more oxidative to maintain ATP levels[100]. In our
samples, as oxidative metabolism decreased, anaerobic metabolism and lactate levels increased,

forcing cells to produce ATP by anaerobic glycolysis. However, long and high-level exposure to
D-galactose generates free radicals, which alter MERCs, causing mitochondrial dysfunction and
inducing aging[101,102]. This is the likely explanation for mitochondrial fragmentation in aged
samples and loss of the MICOS complex, which should be investigated further.
In conclusion, we present a quantitative evaluation of mitochondrial morphology in
mouse skeletal muscle and cardiac tissue using 3D reconstructions, while TEM was utilized in
cell lines to understand factors including cristae architecture. We found structural changes,
including in gross 3D mitochondrial structure, which conferred functional differences upon loss
of MICOS proteins. Similar changes in mitochondrial morphology were observed in aging
muscles and for loss of MICOS proteins in mouse skeletal muscle, and MICOS proteins
decreased with age. We also found that metabolomics and lipidomics are heavily altered across
the aging process, pointing towards the potential roles of the MICOS complex in membrane
integrity across the aging process which must be further investigated. *In vivo* Drosophila models
also highlight the importance of understanding the tissue-dependent response to aging, roles of
individual components in the MICOS complex, and potential MICOS complex-MERC pathway
interactions that may regulate mitochondria and structure. This suggests a relationship between
the MICOS complex and aging, and further studies using 3D reconstruction could elucidate the
link between sarcopenia, the MICOS complex, and disease states in mitochondria.
657
EXPERIMENTAL PROCEDURES
Animal Care and Maintenance
All procedures for the care of mice were in accordance with humane and ethical protocols
approved by the University of Iowa Animal Care and Use Committee (IACUC) following the
National Institute of Health (NIH) Guide for the Care and Use of Laboratory Animals as
described previously[37]. Therefore, all studies are performed in accordance with the ethical
standards laid down in the 1964 Declaration of Helsinki and its later amendments. All
experiments used WT male C57Bl/6J mice housed at 22 °C on a 12-hour light, 12-hour dark
cycle with free access to water and standard chow. Mice were anesthetized with 5%
isoflurane/95% oxygen.
668
RNA Extraction and RT-qPCR
Total RNA was extracted from tissue using TRIzol reagent (Invitrogen, cat #), purified with the
RNeasy kit (Qiagen Inc, cat #), and quantitated by the absorbance at 260 nm and 280 nm using a
NanoDrop 1000 (NanoDrop products, Wilmington, DE, USA) spectrophotometer. Total RNA
(~1 µg) was reverse transcribed using a High-Capacity cDNA Reverse Transcription Kit,
(Applied Biosciences, Carlsbad CA, cat #) followed by real-time quantitative PCR (qPCR)
reactions using SYBR Green (Life Technologies, Carlsbad, CA, cat #)[103]. Triplicate samples for
qPCR (~50 ng) in a 384-well plate were placed into ABI Prism 7900HT instrument (Applied
Biosystems) programmed as follows: 1 cycle at 95°C for 10 min; 40 cycles of 95°C for 15 s;
59°C for 15 s, 72°C for 30 s, and 78°C for 10 s; 1 cycle of 95°C for 15 s; 1 cycle of 60°C for
15 s; and one cycle of 95°C for 15 s. Data were normalized to glyceraldehyde-3-phosphate
dehydrogenase (*Gapdh*), and results are shown as fold changes. qPCR primers were designed
using Primer-Blast or were previously published sequences[37] as shown in Table 1.
682
683 Table 1: qPCR Primers Used

Gene	Primers	

Opa1	Forward	5'-ACCAGGAGACTGTGTCAA-3'
	Reverse	5'-TCTTCAAATAAACGCAGAGGTG-3'
Chchd3	Forward	5'-GAAAAGAATCCAGGCCCTTCCACGCGC-3'
	Reverse	5'-CAGTGCCTAGCACTTGGCACAACCAGGAA-3'
Chchd6	Forward	5'-CTCAGCATGGACCTGGTAGGCACTGGGC-3'
	Reverse	5'-GCCTCAATTCCCACATGGAGAAAGTGGC-3'
Mitofilin	Forward	5'-CCTCCGGCAGTGTTCACCTAGTAACCCCTT-3'
	Reverse	5'-TCGCCCGTCGACCTTCAGCACTGAAAACCTAT-3'

684
685
686
687   *Experimentally evolved Drosophila populations, RNA extraction, qPCR*
688
For this work, we used groups of experimentally *Drosophila melanogaster* where hundreds of
generations of selection on reproductive timing has produced populations with markedly
different aging an longevity patterns. In the control populations, termed CO_{1-5}, genetically
diverse populations (census size ~2000 per replicate) are maintained on a 28-day generation
cycle. From these CO populations, a set of population maintained on a 9-day cycle were created
– ACO_{1-5} (See [104] for more details on how populations were created and still maintained). As a
result of this difference and hundreds of generations, the ACO populations have evolved to
optimize for early reproduction and now develop more quickly and die at much younger ages
than their controls [34]. We also know that the accelerated aging seen in the ACO flies is underlain
by significant genetic differentiation [105], differences in patterns of gene expression [35], and
metabolomic differentiation [33]. The ACO populations are also referred to in this paper as "aged
flies", whereas the CO populations are referred to as "control flies". As such, we believe
comparisons between them can yield insights into the factor that broadly shape differences in
rates of senescence between individuals.
703
Here we use qPCR to compare gene expression for the following genes (Table 2). Specifically,
we looked at differences in cardiac tissue from 21-day old flies from the ACO and CO
populations. We chose day 21 based on demographic data [34] and whole-body transcriptomic
data[35] showing that cohorts are especially differentiated at this age. For each group, we collected
heart tissue using the following protocols. On day 21 from egg, female fruit flies from each of
the 10 ACO and CO populations were anesthetized using Fly Nap (Carolina, NC, USA), a
triethylamine-based anesthetic, for about 1-minute or until no movement was detected. Flies
were dissected in oxygenated artificial hemolymph to expose the cardiac tubes. Fly abdomens
were cut open in order to remove guts/intestines, fat, and ovaries. After removal, hearts were
exposed and additional fat and pericardial cells were carefully suctioned away from the cardiac
tube [106]. It is not possible to fully remove all fat and pericardial cells from the cardiac tube
without damaging the cardiac tube itself. Approximately 18-20 hearts from adults were pooled
for each biological replicate. Three biological replicates were collected for each ACO and CO
population. Total RNA was extracted using Qiazol (Qiagen) and miRNeasy Mini Kit (Qiagen,
catalog no. 217004). An optional on-column DNase digestion was also performed since we
expected our samples to contain less than 1 ug total RNA (Qiagen RNase-Free DNase set,
cataolog no. 79254). RNA was reverse transcribed to cDNA using a QuantiTect Reverse
Transcription Kit (Qiagen).

All cDNA samples were diluted to a standard concentration of 1ng/µl prior to data collection.

Three replicates from each population (ACO and CO) were arbitrarily selected after filtering for

DNA concentration and purity (260/280 ratio >1.8). Samples were run in duplicate on a CFX96

Touch platform (Bio Rad Laboratories, Hercules, CA). Each well contained 7.5 uL of 2X iTaq

SYBR Green Supermix (Bio Rad Laboratories, Hercules, CA), 3 uL of ddH2O, 0.75 uL of each

primer, and 3 uL of template cDNA. The protocol for each primer pair was as follows: 1 cycle at

95ºC for 2 minutes, 40 cycles of 95ºC for 30 seconds, 60ºC for 30 seconds after which a melt

curve analysis was conducted for quality control. Genes dMic60, Bip, and Ire1 were run at an

annealing temperature of 62ºC. Data was analyzed using the ΔΔCt method and normalized to the

Drosophila gene rp49/RpL32. Primers were designed using Primer Blast or previously published

sequences as shown in Table 3.

Table 2: Drosophila qPCR primer sequences

Gene	Primers
ATF-4	F: 5′-TCGCAAAAGTTGGTTAAACG-3′
	R: 5′-TCCGTAGGATTCAACTGCTG-3′
Drp1	F: 5′-TCCACAATCTTCTCGTGCAG-3′
	R: 5′-CATTCACGAGGAGATGCAGC-3′
Marf	F: 5′-GTATGTCCATGAGACGACCA-3′
	R: 5′-CTTGTACACATAGCTTTCGA-3′
Opa1	F: 5′-GACTCTGACCGGGAATACGA-3′
	R: 5′-CTAGAACCATGCGTTGCAGA-3′
MINOS1 (Mic10)	F: 5' GTCCTTTTAACGTTGTTTTGGCA 3'
	R: 5' AGCGATGCCCATTCCAAATC 3'
QIL1 (Mic13)	F: 5' AGACTCTGACCAGACGGACA 3'
	R: 5' AGGGCAGCATGTGGATGAAA 3'
CHCHD3/6 (mic19)	F: 5' GCTAGAGGAACTTCAAAGATGG 3'
	R: 5' GGGATAGGAGGATACTTTCGG 3'
Dmic60 (mitofilin)	F: 5' TCCAAAACATAAATACTCTGGGAAG 3'
	R: 5' ACAAAGCTTGCCAATTTCAGC 3'
APOO (mic26/27)	F: 5' CGGTCTGGCTGGTTTCATCT 3'
	R: 5' GGCACTACGGGAACATCCTC 3'
VDAC/Porin	F: 5'-ATCTGAAGACTAAGACCTCGTCG-3'
	R: 5'-AGACCTTTCCAGACTCCTGGT-3'
IP3R/Itpr	F: 5'- ATGGGCGACAATATAATTGGCTC- 3'
	R: 5'-GTGCTCAAGAAACCGCAAACG-3'
FGF21/ bln	F: 5'- TGTCGCCCGCTGACAATAAT-3'

	R: 5' TTGCTGATGGGCGTGTTACT-3'
Ire1	F: 5' GAACGCGAGTGCGAAGAAAA 3'
	R: 5' CTGATGCAATAAGCCCGCTG 3'
Bip(grp78)/ Hsc 70-3	F: 5' CGCATCGAAATTGAATCCTT 3'
	R: 5' TTCAGGGTGGAACGGAATAG 3'
ATF-6	F: 5'-TGAGCCTAATTCGTCTCCCAC-3'
	R: 5'-TAGACCGCCTCTTCGTTAGAA-3'
RP49	F: 5' AGGCCCAAGATCGTGAAGAA 3'
	R: 5' TCGATACCCTTGGGCTTGC 3'
GRP75/Hsc70-5	F: 5'-CGCGTACCCAAGTTTCTGC-3'
	R: 5'-CGGAACATGCTAGAAGCTCC-3'

Drosophila strains and genetics

Genetic crosses were conducted using yeast corn medium at a consistent temperature of 25□°C,
unless otherwise specified. The Mef2-Gal4 was employed as a control within their respective
genetic backgrounds. Both male and female subjects were used and analyzed collectively, as no
discernible sex differences in mitochondrial morphology were observed in wild-type muscles.
Mef2-Gal4 (III) was utilized for muscle-specific knockdown of MICOS genes. The
mitochondrial network was visualized using UAS-mito-GFP, located on the second chromosome
(BS# 8442).To achieve muscle-specific knockdown MICOS genes, UAS-RNAi trip lines were
used for UAS- *Chchd3* RNAi (BS#51157), UAS-*Mitofilin* RNAi (BS# 63994), UAS-QIL1
RNAi (BS# 44634), UAS-Drp1 RNAi (BS# 51483), UAS-Marf RNAi (BS# 55189), and UAS-
Opa1 RNAi (BS#32358). All stocks were acquired from the Bloomington Drosophila stock
center, denoted by the Bloomington Stock Number (BS#). All chromosome designations and
gene symbols are mentioned in FlyBase ((http://flybase.org)).

Mitochondrial staining
Adult Drosophila thoraces, aged 2-3 days, were dissected in 4% paraformaldehyde (PF, Sigma)
using fine scissors and processed as mentioned in Katti et al[1]. The dissection process isolated
including Indirect Flight Muscles (IFMs). Muscles were then fixed in 4% PF for a specific
duration: IFMs for 1.5 hours. This fixation process was performed using a rotor, followed by a
washing sequence in PBSTx (PBS□+□0.3% TritonX100) thrice, each time for 15 minutes.
Mitochondria were visualized using mitochondrial staining dyes, either Mito Tracker Red
(M22425, Thermofisher, USA) or DMef2-Gal4 driven UAS-mito-GFP. Actin staining was
conducted by incubating the muscles in 2.5□µg/ml of Phalloidin in PBS (Sigma, 1□mg/ml stock
of Phalloidin TRITC) at 25□°C. there incubated muscles for 40 minutes at room temperature.
Following staining, tissues were mounted on a glass slide using Prolong Glass Antifade
Mountant with NucBlue stain (P36985, Thermofisher, USA). Images were subsequently
captured using a Zeiss 780 confocal microscope.

Mitochondrial measurements
Measurements were carried out following procedures previously established[1,2]. Images were
loaded into the ImageJ software (https://imagej.net), a freely available image processing

program. Individual mitochondria were outlined using the freehand tool on 2D light microscopic images provided by the software. The area and aspect ratio (the ratio of the major axis to the minor axis) of each mitochondrion were then calculated using the same ImageJ software. For each data set, three animals (n=3 animals) were analyzed, and these analyses were part of three independent experiments conducted to gather quantifiable data. For further analysis, the number of mitochondria was counted across every three-sarcomere segment.

Isolation of Satellite Cells
Satellite cell differentiation was performed as previously described[14,37].

Gastrocnemius muscles were dissected from 8–10 week-old WT mice and washed twice with 1× PBS supplemented with 1% penicillin-streptomycin and Fungizone (300 mL/100 mL). DMEM-F12 medium with collagenase II (2 mg/mL), 1% penicillin-streptomycin, and Fungizone (300 mL/100 mL) was added to the muscle which was agitated for 90 min at 37°C. For the second wash, collagenase II was changed to 0.5 mg/mL, and the muscle was agitated for 30 min at 37°C. The tissue was cut, passed through a 70 mm cell strainer, and after centrifugation satellite cells were plated on BD Matrigel-coated dishes. To differentiate cells into myoblasts, a mixture of DMEM-F12, 20% fetal bovine serum (FBS), 40 ng/mL basic fibroblast growth factor (bFGF, R and D Systems, 233-FB/CF), 1× non-essential amino acids, 0.14 mM β-mercaptoethanol, and 1× penicillin/streptomycin, and Fungizone was used. Myoblasts were maintained with 10 ng/mL bFGF and when cells reached 80% confluence, myoblasts were differentiated in DMEM-F12, 2% FBS, 1× insulin–transferrin–selenium medium. Cells were cultured at 37°C, 5% CO_2 Dulbecco's modified Eagle's medium (DMEM; GIBCO) supplemented with 10% FBS (Atlanta Bio selected), and 1% penicillin-streptomycin (Gibco, Waltham, MA, USA).

CRISPR-Cas9 Knockouts
After three days, myotubes were incubated with CRISPR/Cas9 mixtures to produce the following knockouts—control CRISPR/Cas9 (sc-418922), *Chchd6* (Mic25) CRISPR (sc-425817), *Chchd3* (Mic19) CRISPR (sc-425804), and *Mitofilin* (Mic60) CRISPR (sc-429376) (Santa Cruz Biotechnology, California, US), with the use of guide RNA (Table 2). We incubated 2.5% relevant CRISPR, 2.5% RNAiMax (ThermoFisher Scientific; cat # 13778075), and 95% Opti-MEM (Gibco; cat #31985070) in a tube for 20 minutes. Cells were washed twice with PBS after removal of the medium, then 800 µL of OPT-MEM and 200 µL of the CRISPR mixture were added to each well and ran in triplicates. Cells were incubated for 4 hours at 37 C, 1.0 mL of DMEM medium was added, cells were incubated overnight. The myotubes were then washed with PBS and the medium was replaced. Experiments were performed between 3 and 7 days after knockout for a total of 6 days of differentiation.

Table 3: Guide RNA and Plasmids Used

Gene Name	Type of Plasmid	CAS Number
Mitofilin	CRISPR/Cas9 KO (m)	sc-429376
Chchd6	CRISPR/Cas9 KO (m)	sc-425817
Chchd3	CRISPR/Cas9 KO (m)	sc-425804
Control	CRISPR/Cas9 KO (m)	sc-418922

Serial Block-Face Scanning Electron Microscope (SBF-SEM) Processing of Mouse Muscle
Fibers
SBF-SEM preparation was performed as described previously[15,36,107]. Male mice were
anesthetized with 5% isoflurane. Once the hair and skin were removed, the hindlimbs were
incubated in 2% glutaraldehyde with 100 mM phosphate buffer for 30 min. Gastrocnemius
muscles were dissected, cut into 1-mm^3 cubes, and incubated in 2.5% glutaraldehyde, 1%
paraformaldehyde, 120 mM sodium cacodylate solution for 1 hour. Tissues were washed three
times with 100 mM cacodylate buffer at room temperature before immersion in 3% potassium
ferrocyanide and 2% osmium tetroxide for 1 hour at 4°C, then treated with 0.1%
thiocarbohydrazide, 2% filtered osmium tetroxide for 30 min, and left overnight in 1% uranyl
acetate at 4°C. Between each step, three de-ionized water washes were performed. The following
day, samples were immersed in 0.6% lead aspartate solution for 30 min at 60°C and dehydrated
in graded concentrations of acetone. Dehydration was for five min each in 20%, 50%, 70%, 90%,
95%, and 100% acetone. Tissues were impregnated in Epoxy Taab 812 hard resin, then
embedded in fresh resin, and polymerized at 60°C for 36–48 hours. Once polymerization was
complete, blocks were sectioned for TEM to identify areas of interest, trimmed to 0.5 mm × 0.5
mm, and glued to aluminum pins. Afterward, pins were placed in an FEI/Thermo Scientific
Volumescope 2 SEM, a state-of-the-art SBF imaging system. Running on a FEI/Thermo
Scientific Volumescope 2 SEM, a state-of-the-art SBF imaging system, we obtained 300–400 10
µm by 10 µm ultrathin (90 nm) serial sections per previous techniques [15]. All sections were
collected onto formvar-coated slot grids (Pella, Redding CA), stained, and imaged as previously
described[15,36,107].
832
Quantification of TEM Micrographs and Parameters Using ImageJ
Quantification of TEM images was performed as described previously using the
NIH *ImageJ* software[14,36]. Cells were divided into four quadrants and two quadrants were
selected randomly for complete analysis. Individuals blinded to the experimental design had a
minimum of 10 cells to measure with three analyses to obtain accurate and reproducible values.
If variability occurred, the number of cells was expanded to 30 cells per individual to reduce the
variability.
840
Measurement of OCR Using Seahorse
To measure cellular respiration, the XF24 extracellular flux (XF) bioanalyzer (Agilent
Technologies/Seahorse Bioscience, North Billerica, MA, USA) was used. Cells were plated at a
density of 2×10^4 per well and differentiated. After 3 days of differentiation, *Opa-1*, *CHCHD3*,
CHCHD6, or *Mitofilin* genes were knocked out as described above. Three days after knockdown,
the medium was changed to XF-DMEM, and cells were kept in a non-CO_2 incubator for 60 min.
The basal OCR was measured in XF-DMEM. Oxygen consumption was measured after the
addition of oligomycin (1 µg/mL); carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone
(FCCP; 1 µM); rotenone (1 µM) and antimycin A (10 µM) [37,108]. Cells were then switched to
glucose-free XF-DMEM and kept in a non-CO_2 incubator for 60 min for the glycolysis stress
test. Seahorse experimental data are for triplicate Seahorse plates. Three independent
experiments were performed with four to six replicates for each time and for each condition and
representative data from the replicates are shown.
854
Segmentation and Quantification of 3D SBF-SEM Images Using Amira

Intermyofibrillar (IMF) mitochondria are located between myofibrils, arranged in pairs at the z-
band of each sarcomere, with 2D elongated tubular shapes[109], and these were the mitochondria
analyzed. For each region of interest across the three age groups, we analyzed 300 slices at 50
µm intervals in transverse intervals. For 3D reconstruction, SBF-SEM images were manually
segmented in Amira as described previously[15,36]. All serial sections (300–400 slices) were loaded
onto Amira and structural features were traced manually on sequential slices of micrograph
blocks. Structures in mice were collected from 30–50 serial sections that were then stacked,
aligned, and visualized using Amira to make videos and quantify volumetric structures. An
average of 500 total mitochondria across four ROIs from three mice were collected for
quantification. For 3D reconstruction of myotubes, approximately 20 mitochondria from a
minimum of 10 cells were collected. Quantification of SBF-SEM images was performed as
described previously[15] using the Amira software (Thermo Scientific).
868
Western-Blotting
Tissues from adult (3 months) and aged (2 years) mice were lysed with RIPA lysis buffer (1%
NP40, 150 mM NaCl, 25 mM Tris base, 0.5% sodium deoxycholate, 0.1% SDS, 1% phosphatase
inhibitor cocktails #2 (Sigma P5726-1ML) and #3 (Sigma P0044-1ML), 1 cOmplete protease
inhibitor tablet (Sigma 04693159001)). Protein content was quantified using a BCA Assay
(Thermo Scientific VLBL00GD2) and equal protein was run on 4-20% Tris-Glycine Gels
(Invitrogen WXP42012BOX). Protein was transferred to a nitrocellulose membrane (Li-Cor 926-
31092). Membranes were incubated overnight with primary antibodies overnight at 4 $^{\circ}$C:
MTOC1 (Invitrogen PA5-26688), phospho S406 ATGL (Abcam ab135093), DRP1 (CST
8570S), pDRP1 (CST 6319S), OPA1 (BD Biosciences 612306), Mic60/mitofilin (Abcam
ab110329), SLC25A46 (Abcam ab237760), SAM50 (Proteintech 20824-1-AP), tubulin (Novus
NB100-690). Secondary antibodies used were diluted to 1:10000 and incubated at room
temperature for 1 hour: Donkey anti-Mouse IgG (H+L) (Invitrogen A32789), Donkey anti-
Rabbit IgG (H+L) (Invitrogen A32802). Blots were imaged with the Li-Cor Odyssey
CLx infrared imaging system. Raw blots shown in SFigure 7.
884
Gas Chromatography-Mass Spectrometry (GC-MS) for MICOS KO
Samples were extracted for metabolites and prepared as previously designed[33,110].
Samples were extracted for metabolites in −80°C 2:2:1 methanol/acetonitrile/water that
contained a mixture of nine internal standards (d_4-citric acid, $^{13}C_5$-glutamine, $^{13}C_5$-glutamic
acid, $^{13}C_6$-lysine, $^{13}C_5$-methionine, $^{13}C_3$-serine, d_4-succinic acid, $^{13}C_{11}$-tryptophan, d_8-valine;
Cambridge Isotope Laboratories) at a concentration of 1 µg/mL each, at a ratio of 18:1
(extraction solvent:sample volume). Cells were lyophilized overnight before extraction and
homogenized with a ceramic bead mill homogenizer after the addition of extraction buffer.
Samples were incubated for one hour at −20°C and centrifuged at maximum speed for 10 min.
All supernatants were transferred to fresh tubes and pooled quality control (QC) samples were
prepared by adding an equal volume of each sample to a fresh 1.5 mL microcentrifuge tube. A
speed vac was used to evaporate the pooled QCs, samples, and processing blanks, which were
made by adding extraction solvent to microcentrifuge tubes. Derivatives of the dried products
were obtained using methoxamine hydrochloride (MOX) and *N*,O-
bis(trimethylsilyl)trifluoroacetamide (TMS). Products were rehydrated in 30 µL of 11.4 mg/mL
Molybdenum Carbide (MOC) in anhydrous pyridine (VWR), vortexed for 10 minutes, and
incubated at 60°C for 1 hour. Then 20 µL of TMS was added, vortexing was repeated for one

min, and samples were heated for an hour at 60°C. Samples of 1 µL were analyzed by GC-MS
using a Thermo Trace 1300 GC with a TraceGold TG-5SilMS column for GC chromatographic
separation. The GC temperature was set at 250°C for the inlet, with the oven temperature at a
gradient with 3 min at 80 °C, increasing by 20°C a minute to a final 280°C for the last 8 minutes.
The settings for the GC machine were 20:1 split ratio, split flow; 24 µL/min, purge flow; 5
mL/min, carrier mode; constant flow, carrier flow rate: 1.2 mL/min. Between each injection
sample, the column was washed three times with pyridine. Metabolites were detected using the
Thermo ISQ single quadrupole mass spectrometer, with data acquired from 3.90 to 21.00 min in
the EI mode (70 eV) by single-ion monitoring. The profiling of the metabolites was performed
using TraceFinder 4.1 with standard verified peaks and retention times. TraceFinder was used to
compare metabolite peaks in each sample against an in-house library of standards. For these
standards, retention times and fragment ions for each were analyzed, with fragment ions for both
the target peak and two confirming ions. For the samples, we identified metabolites that matched
both retention times and the three fragment ions. TraceFinder was also used for GC-MS peak
integration to obtain peak areas for each metabolite. The profiling of the metabolites was
performed using TraceFinder 4.1 with standard verified peaks and retention times. TraceFinder
was used to compare metabolite peaks in each sample against an in-house library of standards.
TraceFinder was also used for GC-MS peak integration to obtain peak areas for each metabolite.
After this analysis, we used previously described protocols[111] to correct for drift over time by
using QC samples run at both the beginning and end of the sequence. The data was then
normalized to an internal standard to control for extraction, derivatization, and/or loading effects.
923
Liquid Chromatography-Mass Spectrometry (LC-MS) for MICOS KO
Myotubes were dried, rehydrated in 40 µL acetonitrile:water (1:1), and vortexed. For LC-MS, 2
µL of the sample was used with a Thermo Q Exactive hybrid quadrupole Orbitrap mass
spectrometer with a Vanquish Flex UHPLC system and a Millipore SeQuant ZIC-pHILIC
column (length area = 2.1×150 mm, 5 µm particle size) with a ZIC-pHILIC guard column
(length area = 20×2.1 mm). The mobile phase comprised solvent A (20 mM ammonium
carbonate [$(NH_4)_2CO_3$] and 0.1% ammonium hydroxide [NH_4OH]) and solvent B (acetonitrile).
The mobile phase gradient started at 80% solvent B, decreased to 20% solvent B over 20 min,
returned to 80% solvent B in 0.5 min, and was held at 80% for 7 min [112]. From there, the mass
spectrometer was operated in the full-scan, polarity-switching mode for 1 to 20 min, spray
voltage set to 3.0 kV, capillary heated at 275°C, and HESI probe heated at 350°C. The sheath gas
flow, auxiliary gas flow, and sweep gas flow were 40 units, 15 units, and 1 unit, respectively. We
examined an *m/z* range of 70–1000, the resolution was set at 70,000, the AGC target at 1×10^6,
and the maximum injection time was set to 200 ms [111]. TraceFinder 4.1 software was used for
analysis and metabolites were identified based on an in-house library. Drift was corrected for as
described above[111]. Data were normalized and further visualization and analysis were performed
on MetaboAnalyst 5.0[113].
941
Analyzing Metabolomic Data for MICOS KO
Metabolomic analysis was performed as described previously[33] using the web service
MetaboAnalyst 5.0 (https://www.metaboanalyst.ca/MetaboAnalyst/ModuleView.xhtml, last
accessed on 8 February 2022) that combines machine learning methods and statistics to group
data using PCA, heat mapping, metabolite set enrichment analysis, and statistical analysis. One-
way ANOVA and Fisher's LSD multiple comparison test were also used. PCA uses score plots

to provide an overview of variance for the principal components. Heatmaps separate hierarchical
clusters leading to progressively larger clusters. Clusters are based on similarity using Euclidean
distance and Ward's linkage to minimize the clustering needed. Metabolite Set Enrichment
Analysis (MSEA), which determines whether a set of functionally related metabolites is altered,
can be used to identify consistent changes across many metabolites with similar roles.
Overrepresentation analysis determines whether a group of compounds is overrepresented in
comparison to pure chance and whether a group of metabolites have similar changes. In this
analysis, the fold enrichment was calculated by dividing the observed hits by the expected
metabolites. Expected number of hits are calculated by MetaboAnalyst 5.0. GraphPad Prism
software (La Jolla, CA, USA) was used for statistical analysis with data expressed as mean ±
standard deviation, and one-tailed p-values ≤ 0.01 were considered significant.
959
Metabolomics for Aged Samples
Methods used previously verified techniques [114–116]. Frozen tissues were weighed, ground with a
liquid nitrogen in a cryomill (Retsch) at 25 Hz for 45 seconds, before extracting tissues 40:40:20
acetonitrile: methanol: water +0.5% FA +15% NH4HCO3 [116] with a volume of 40µL solvent per
1mg of tissue, vortexed for 15 seconds, and incubated on dry ice for 10 minutes. Tissue samples
were then centrifuged at 16,000 g for 30 minutes. The supernatants were transferred to new
Eppendorf tubes and then centrifuged again at 16,000 g for 25 minutes to remove and residual
debris before analysis.
968
Extracts were analyzed within 24 hours by liquid chromatography coupled to a mass
spectrometer (LC-MS). The LC–MS method was based on hydrophilic interaction
chromatography (HILIC) coupled to the Orbitrap Exploris 240 mass spectrometer (Thermo
Scientific) [114]. The LC separation was performed on a XBridge BEH Amide column (2.1 x 150
mm, 3.5 µm particle size, Waters, Milford, MA). Solvent A is 95%: 5% H2O: acetonitrile with
20 mM ammonium acetate and 20mM ammonium hydroxide, and solvent B is 90%: 10%
acetonitrile: H2O with 20 mM ammonium acetate and 20mM ammonium hydroxide. The
gradient was 0 min, 90% B; 2 min, 90% B; 3 min, 75% B; 5 min, 75% B; 6 min, 75% B; 7 min,
75% B; 8 min, 70% B; 9 min, 70% B; 10 min, 50% B; 12 min, 50% B; 13 min, 25% B; 14min,
25% B; 16 min, 0% B; 18 min, 0% B; 20 min, 0% B; 21 min, 90% B; 25 min, 90% B. The
following parameters were maintained during the LC analysis: flow rate 150 mL/min, column
temperature 25 °C, injection volume 5 µL and autosampler temperature was 5 °C. For the
detection of metabolites, the mass spectrometer was operated in both negative and positive ion
mode. The following parameters were maintained during the MS analysis: resolution of 180,000
at m/z 200, automatic gain control (AGC) target at 3e6, maximum injection time of 30 ms and
scan range of m/z 70-1000. Raw LC/MS data were converted to mzXML format using the
command line "msconvert" utility [115]. Data were analyzed via the EL-MAVEN software version
12.
987
Lipidomics for Aged Samples
Tissue homogenization and extraction for lipids:
Tissues were homogenized using a Retsch CryoMill. The homogenate was mixed with 1 mL of
Extraction Buffer containing IPA/H2O/Ethyl Acetate (30:10:60, v/v/v) and Avanti Lipidomix
Internal Standard (diluted 1:1000) (Avanti Polar Lipids, Inc. Alabaster, AL). Samples were
vortexed and transferred to bead mill tubes for homogenization using a VWR Bead Mill at 6000

g for 30 seconds, repeated twice. The samples were then sonicated for 5 minutes and centrifuged
at 15,000 g for 5 minutes at 4°C. The upper phase was transferred to a new tube and kept at 4°C.
To re-extract the tissues, another 1 mL of Extraction Buffer (30:10:60, v/v/v) was added to the
tissue pellet-containing tube. The samples were vortexed, homogenized, sonicated, and
centrifuged as described earlier. The supernatants from both extractions were combined, and the
organic phase was dried under liquid nitrogen gas.

1000

Sample reconstitution for lipids:
The dried samples were reconstituted in 300 µL of Solvent A (IPA/ACN/H2O, 45:35:20, v/v/v).
After brief vortexing, the samples were sonicated for 7 minutes and centrifuged at 15,000 g for
10 minutes at 4°C. The supernatants were transferred to clean tubes and centrifuged again for 5
minutes at 15,000 g at 4°C to remove any remaining particulates. For LC-MS lipidomic analysis,
60 µL of the sample extracts were transferred to mass spectrometry vials.

1007

LC-MS analysis for lipids:
Sample analysis was performed within 36 hours after extraction using a Vanquish UHPLC
system coupled with an Orbitrap Exploris 240™ mass spectrometer equipped with a H-ESI™
ion source (all Thermo Fisher Scientific). A Waters (Milford, MA) CSH C18 column (1.0×150
mm $\times$ 1.7 µm particle size) was used. Solvent A consisted of ACN:H2O (60:40; v/v) with 10
mM Ammonium formate and 0.1% formic acid, while solvent B contained IPA:ACN (95:5; v/v)
with 10 mM Ammonium formate and 0.1% formic acid. The mobile phase flow rate was set at
0.11 mL/min, and the column temperature was maintained at 65 °C. The gradient for solvent B
was as follows: 0 min 15% (B), 0–2 min 30% (B), 2–2.5 min 48% (B), 2.5–11 min 82% (B), 11–
11.01 min 99% (B), 11.01–12.95 min 99% (B), 12.95–13 min 15% (B), and 13–15 min 15% (B).
Ion source spray voltages were set at 4,000 V and 3,000 V in positive and negative mode,
respectively. Full scan mass spectrometry was conducted with a scan range from 200 to 1000
m/z, and AcquireX mode was utilized with a stepped collision energy of 30% with a 5% spread
for fragment ion MS/MS scan.

1022

Lipidomics Analysis:
After normalization, all data was analyzed in R using the *lipidr* package (Mohamed et al.
2020). All code to analyze data and generate figures can be found here:
https://github.com/mphillips67/Lipidomic-Analysis-Young-and-Aged-Mouse-Tissue. Data sets
associated with each tissue type we looked at were analyzes independently. Prior to running any
analysis, data was log transformed and further processed. For lipids where there were multiple
readings across replicates, sets of readings with the highest values were identified and all others
were discarded. Next there were several instances (2 to 3) per tissue type were all samples had
identical measurements. As this was mostly likely due to technical errors, these entries were also
discarded from the data set. Lastly, to be compatible with *lipidr*, lipid names had to be modified
to fit a standard "CLS xx:x/yy:y" naming scheme where CLS refers to the abbreviated lipid class
and xx:x and yy:y refer to the first and second chains (note: code to generate a conversion key
are available through the previously mentioned GitHub repository).
After processing, lipid composition between young and old samples were compared using
the "de_analysis" function from *lipidr* with default settings. Here *lipidr* uses moderated t-tests to
identify significantly differentiated lipids between samples types within a given tissue type.
Significantly differentiated lipids are those with adjusted p-values < 0.05 (note: p-values were

adjusted to correct for multiple comparisons using and FDR procedure) and log fold changes greater than 1 or less than -1. These results were then used to perform a lipid set enrichment analysis using the "lsea" function where entries were ranked by fold change, only classes with at least 4 associated lipids were considered, and 100000 permutations were run. Here the method *lipdr* used is based on the commonly used gene set enrichment analysis approach outlined in Subramanian et al. (2005). Briefly, lipid class and chain length categories are determined from annotations extracted from lipid names in the data set and lipids are ranked by fold change. A permutation algorithm is then used to calculate enrichment scores and p-values for each lipid set. Sets with adjusted p-values < 0.05 were defined as significantly enriched. Lastly, heatmaps were generated for significantly enriched lipid classes using the "plot_heatmap".

Data Analysis
All SBF-SEM and TEM data were presented as the mean of at least three independent experiments with similar outcomes. Results were presented as mean ± standard error with individual data points shown. Data with only two groups were analyzed using an unpaired t-test. For nanotunnel quantification, a Mann-Whitney test (unpaired, nonparametric) t-test was performed between two groups. If more than two groups were compared, one-way ANOVA was performed, and significance was assessed using Fisher's protected least significant difference (LSD) test. GraphPad Prism software package was used for t-tests and ANOVA analyses (La Jolla, CA, USA). For all statistical analyses, $p < 0.05$ indicated a significant difference. Higher degrees of statistical significance (**, ***, ****) were defined as $p < 0.01$, $p < 0.001$, and $p < 0.0001$, respectively.

ACKNOWLEDGEMENTS
This project was funded by the National Institute of Health (NIH) NIDDK T-32, number DK007563 entitled Multidisciplinary Training in Molecular Endocrinology to Z.V.; National Institute of Health (NIH) NIDDK T-32, number DK007563 entitled Multidisciplinary Training in Molecular Endocrinology to A.C.; Integrated Training in Engineering and Diabetes", Grant Number T32 DK101003; Burroughs Wellcome Fund Postdoctoral Enrichment Program #1022355 to D.S.; The UNCF/ Bristol-Myers Squibb (UNCF/BMS)- E.E. Just Postgraduate Fellowship in Life sciences Fellowship and Burroughs Wellcome Fund/ PDEP #1022376 to H.K.B.; NSF MCB #2011577I to S.A.M.; NSF EES2112556, NSF EES1817282, NSF MCB1955975, and CZI Science Diversity Leadership grant number 2022-253614 from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation to S.D.; The UNCF/Bristol-Myers Squibb E.E. Just Faculty Fund, Career Award at the Scientific Interface (CASI Award) from Burroughs Welcome Fund (BWF) ID # 1021868.01, BWF Ad-hoc Award, NIH Small Research Pilot Subaward to 5R25HL106365-12 from the National Institutes of Health PRIDE Program, DK020593, Vanderbilt Diabetes and Research Training Center for DRTC Alzheimer's Disease Pilot & Feasibility Program. CZI Science Diversity Leadership grant number 2022- 253529 from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation to A.H.J.; and National Institutes of Health grant HD090061 and the Department of Veterans Affairs Office of Research award I01 BX005352 (to J.G.). Howard Hughes Medical Institute Hanna H. Gray Fellows Program Faculty Phase (Grant# GT15655 awarded to M.R.M); and Burroughs Wellcome Fund PDEP Transition to Faculty (Grant# 1022604 awarded to M.R.M). Additional support was provided by the Vanderbilt Institute for

Clinical and Translational Research program supported by the National Center for Research
Resources, Grant UL1 RR024975–01, and the National Center for Advancing Translational
Sciences, Grant 2 UL1 TR000445–06 and the Cell Imaging Shared Resource. Its contents are
solely the responsibility of the authors and do not necessarily represent the official view of the
NIH. The funders had no role in study design, data collection and analysis, decision to publish,
or preparation of the manuscript. The co-authors would like to acknowledge the Huck
Metabolomics Core Facility for use of the UHPLC coupled to the OE240 Mass Spectrometer and
staff Drs. Imhoi Koo, Ashley Shay, and Sergei Koshkin for helpful discussions on sample
preparation for lipidomics. We would like to thank Mariya Sweetwyne for her advice as an aging
expert. We would like to thank Anastasia Berlynn Casper for her contributions to qPCR.

1096
1097

CONFLICT OF INTEREST
1099 All authors declare that they have no conflict of interest.

1100

ETHICAL STANDARDS:
1102 The manuscript does not contain clinical studies or patient data.

1103

CONSENT FOR PUBLICATION
1105 All authors have agreed to the final version of this manuscript.

1106

DATA AVAILABILITY STATEMENT
1108 The data that support the findings of this study are available from the corresponding author upon
1109 reasonable request.

1110

AUTHOR CONTRIBUTIONS
Z.V., E.G., L.V., J.S., H.K.B., S.A.M., M.A.P., M.R.M., A.H.J., J.A.G., and D.D. conceived and

designed research; A.G.M., A.C., L.V., Z.V., T.A.C, B.C.M., J.L., H.K.B., B.R, C.E., D.D.,

A.C.M., B.C.J., P.P, M.R.M, A.H.J., and J.A.G. performed experiments; J.D., K.N, J.S., E.G.,

Z.V., J.L., B.R., T.A.C., A.K.R., A.M.Q., V.E., E.G., D.D., A.C.M., B.C.J., P.P, M.R.M., J.A.G.,

and A.H.J. analyzed data; B.T, K.N, J.S., E.G., Z.V., S.A.M., A.M.Q., V.E., H.K.B., A.C.,

A.G.M., J.D., M.A.P., M.R.M., D.D., J.A.G., and A.H.J.. interpreted results of experiments and

prepared figures; K.N, E.G., Z.V., J.S., S.A.M., L.V., A.G.M., M.A.P., A.K.R., B.C.M., B.T.,

C.E., A.C., H.K.B., M.R.M., D.D., J.A.G., and A.H.J. drafted manuscript, edited, and revised

manuscript; M.R.M., A.H.J., D.D., and J.A.G. approved final version of manuscript.

1121

REFERENCES

1123

1. Haas RH. Mitochondrial Dysfunction in Aging and Diseases of Aging. *Biology*. 2019;8:48.

2. Miller BF, Baehr LM, Musci RV, Reid JJ, Peelor III FF, Hamilton KL, Bodine SC. Muscle-
specific changes in protein synthesis with aging and reloading after disuse atrophy. *Journal*
of Cachexia, Sarcopenia and Muscle. 2019;10:1195–1209.

3. Coen PM, Musci RV, Hinkley JM, Miller BF. Mitochondria as a Target for Mitigating
Sarcopenia. *Front Physiol*. 2019;9:1883.

4. Anand R, Wai T, Baker MJ, Kladt N, Schauss AC, Rugarli E, Langer T. The i-AAA protease
YME1L and OMA1 cleave OPA1 to balance mitochondrial fusion and fission. *J Cell Biol*.
2014;204:919–929.

5. Renken CW. The structure of mitochondria. 2004.

6. Cogliati S, Enriquez JA, Scorrano L. Mitochondrial cristae: where beauty meets
functionality. *Trends in biochemical sciences*. 2016;41:261–273.

7. Kühlbrandt W. Structure and function of mitochondrial membrane protein complexes. *BMC*
Biology. 2015;13:89.

8. Otera H, Ishihara N, Mihara K. New insights into the function and regulation of
mitochondrial fission. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research*.
2013;1833:1256–1268.

9. Favaro G, Romanello V, Varanita T, Desbats MA, Morbidoni V, Tezze C, Albiero M,
Canato M, Gherardi G, De Stefani D. DRP1-mediated mitochondrial shape controls calcium
homeostasis and muscle mass. *Nature communications*. 2019;10:1–17.

10. Vincent AE, Turnbull DM, Eisner V, Hajnóczky G, Picard M. Mitochondrial Nanotunnels.
Trends in Cell Biology. 2017;27:787–799.

11. Zhang L, Trushin S, Christensen TA, Bachmeier BV, Gateno B, Schroeder A, Yao J, Itoh K,
Sesaki H, Poon WW. Altered brain energetics induces mitochondrial fission arrest in
Alzheimer's Disease. *Scientific reports*. 2016;6:1–12.

12. Zhang J, He Z, Fedorova J, Logan C, Bates L, Davitt K, Le V, Murphy J, Li M, Wang M,
Lakatta EG, Ren D, Li J. Alterations in mitochondrial dynamics with age-related
Sirtuin1/Sirtuin3 deficiency impair cardiomyocyte contractility. *Aging Cell*. 2021;20:e13419.

13. Kozjak-Pavlovic V. The MICOS complex of human mitochondria. *Cell and tissue research*.
2017;367:83–93.

14. Lam J, Katti P, Biete M, Mungai M, AshShareef S, Neikirk K, Garza Lopez E, Vue Z,
Christensen TA, Beasley HK, Rodman TA, Murray SA, Salisbury JL, Glancy B, Shao J,
Pereira RO, Abel ED, Hinton A. A Universal Approach to Analyzing Transmission Electron
Microscopy with ImageJ. *Cells*. 2021;10:2177.

15. Garza-Lopez E, Vue Z, Katti P, Neikirk K, Biete M, Lam J, Beasley HK, Marshall AG,
Rodman TA, Christensen TA, Salisbury JL, Vang L, Mungai M, AshShareef S, Murray SA,
Shao J, Streeter J, Glancy B, Pereira RO, Abel ED, Hinton A. Protocols for Generating
Surfaces and Measuring 3D Organelle Morphology Using Amira. *Cells*. 2022;11:65.

16. Rampelt H, Zerbes RM, van der Laan M, Pfanner N. Role of the mitochondrial contact site
and cristae organizing system in membrane architecture and dynamics. *Biochimica et
Biophysica Acta (BBA)-Molecular Cell Research*. 2017;1864:737–746.

17. Hu C, Shu L, Huang X, Yu J, Li L, Gong L, Yang M, Wu Z, Gao Z, Zhao Y, Chen L, Song
Z. OPA1 and MICOS Regulate mitochondrial crista dynamics and formation. *Cell Death
Dis*. 2020;11:1–17.

18. Friedman JR, Mourier A, Yamada J, McCaffery JM, Nunnari J. MICOS coordinates with
respiratory complexes and lipids to establish mitochondrial inner membrane architecture.
Elife. 2015;4:e07739.

19. Holmström MH, Iglesias-Gutierrez E, Zierath JR, Garcia-Roves PM. Tissue-specific control
of mitochondrial respiration in obesity-related insulin resistance and diabetes. *American
Journal of Physiology-Endocrinology and Metabolism*. 2012;302:E731–E739.

20. Herbers E, Kekäläinen NJ, Hangas A, Pohjoismäki JL, Goffart S. Tissue specific differences
in mitochondrial DNA maintenance and expression. *Mitochondrion*. 2019;44:85–92.

21. Mukund K, Subramaniam S. Skeletal muscle: A review of molecular structure and function,
in health and disease. *Wiley Interdisciplinary Reviews: Systems Biology and Medicine*.
2020;12:e1462.

22. Crupi AN, Nunnelee JS, Taylor DJ, Thomas A, Vit J-P, Riera CE, Gottlieb RA, Goodridge
HS. Oxidative muscles have better mitochondrial homeostasis than glycolytic muscles
throughout life and maintain mitochondrial function during aging. *Aging (Albany NY)*.
2018;10:3327–3352.

23. Vue Z, Neikirk K, Vang L, Garza-Lopez E, Christensen TA, Shao J, Lam J, Beasley HK,
Marshall AG, Crabtree A, Anudokem J, Rodriguez B, Kirk B, Bacevac S, Barongan T, Shao
B, Evans CS, Taylor B, Reddy AK, Murray SA, Mobley BC, Gomez JA, Phillips MA, Exil
V, Dai D-F, Hinton A. Tri-dimensional Mitochondria Reconstructions of Cardiac Muscle
Changes in Size Across Aging due to the MICOS Complex. 2022;:2022.04.23.489291.

24. Chaudhary KR, El-Sikhry H, Seubert JM. Mitochondria and the aging heart. *J Geriatr
Cardiol*. 2011;8:159–167.

25. Vincent AE, White K, Davey T, Philips J, Ogden RT, Lawless C, Warren C, Hall MG, Ng
YS, Falkous G, Holden T, Deehan D, Taylor RW, Turnbull DM, Picard M. Quantitative 3D
Mapping of the Human Skeletal Muscle Mitochondrial Network. *Cell Reports*. 2019;26:996-
1009.e4.

26. Koopman WJH, Visch H-J, Verkaart S, van den Heuvel LWPJ, Smeitink JAM, Willems
PHGM. Mitochondrial network complexity and pathological decrease in complex I activity
are tightly correlated in isolated human complex I deficiency. *Am J Physiol Cell Physiol*.
2005;289:C881-890.

27. Li H, Ruan Y, Zhang K, Jian F, Hu C, Miao L, Gong L, Sun L, Zhang X, Chen S.
Mic60/Mitofilin determines MICOS assembly essential for mitochondrial dynamics and
mtDNA nucleoid organization. *Cell Death & Differentiation*. 2016;23:380–392.

28. Genin EC, Plutino M, Bannwarth S, Villa E, Cisneros☐Barroso E, Roy M, Ortega☐Vila B,
Fragaki K, Lespinasse F, Pinero☐Martos E. CHCHD 10 mutations promote loss of
mitochondrial cristae junctions with impaired mitochondrial genome maintenance and
inhibition of apoptosis. *EMBO molecular medicine*. 2016;8:58–72.

29. Genin EC, Bannwarth S, Lespinasse F, Ortega-Vila B, Fragaki K, Itoh K, Villa E, Lacas-
Gervais S, Jokela M, Auranen M, Ylikallio E, Mauri-Crouzet A, Tyynismaa H, Vihola A,
Augé G, Cochaud C, Sesaki H, Ricci J-E, Udd B, Vives-Bauza C, Paquis-Flucklinger V.
Loss of MICOS complex integrity and mitochondrial damage, but not TDP-43 mitochondrial
localisation, are likely associated with severity of CHCHD10-related diseases. *Neurobiol
Dis*. 2018;119:159–171.

30. Zheng J, Croteau DL, Bohr VA, Akbari M. Diminished OPA1 expression and impaired
mitochondrial morphology and homeostasis in Aprataxin-deficient cells. *Nucleic acids
research*. 2019;47:4086–4110.

31. Varanita T, Soriano ME, Romanello V, Zaglia T, Quintana-Cabrera R, Semenzato M,
Menabò R, Costa V, Civiletto G, Pesce P. The OPA1-dependent mitochondrial cristae
remodeling pathway controls atrophic, apoptotic, and ischemic tissue damage. *Cell
metabolism*. 2015;21:834–844.

32. Khin PP, Hong Y, Yeon M, Lee DH, Lee JH, Jun H-S. Dulaglutide improves muscle
function by attenuating inflammation through OPA-1-TLR-9 signaling in aged mice. *Aging
(Albany NY)*. 2021;13:21962–21974.

33. Phillips MA, Arnold KR, Vue Z, Beasley HK, Garza-Lopez E, Marshall AG, Morton DJ,
McReynolds MR, Barter TT, Hinton A. Combining Metabolomics and Experimental
Evolution Reveals Key Mechanisms Underlying Longevity Differences in Laboratory
Evolved Drosophila melanogaster Populations. *International Journal of Molecular Sciences*.
2022;23:1067.

34. Burke MK, Barter TT, Cabral LG, Kezos JN, Phillips MA, Rutledge GA, Phung KH, Chen
RH, Nguyen HD, Mueller LD, Rose MR. Rapid divergence and convergence of life-history
in experimentally evolved Drosophila melanogaster. *Evolution*. 2016;70:2085–2098.

35. Barter TT, Greenspan ZS, Phillips MA, Mueller LD, Rose MR, Ranz JM. Drosophila
transcriptomics with and without ageing. *Biogerontology*. 2019;20:699–710.

36. Hinton A, Katti P, Christensen TA, Mungai M, Shao J, Zhang L, Trushin S, Alghanem A,
Jaspersen A, Geroux RE. A comprehensive approach for artifact-free sample preparation and
assessment of mitochondrial morphology in tissue and cultured cells. *bioRxiv*. 2021.

37. Pereira RO, Tadinada SM, Zasadny FM, Oliveira KJ, Pires KMP, Olvera A, Jeffers J,
Souvenir R, Mcglauflin R, Seei A. OPA 1 deficiency promotes secretion of FGF 21 from
muscle that prevents obesity and insulin resistance. *The EMBO journal*. 2017;36:2126–2145.

38. Tezze C, Romanello V, Desbats MA, Fadini GP, Albiero M, Favaro G, Ciciliot S, Soriano
ME, Morbidoni V, Cerqua C, Loefler S, Kern H, Franceschi C, Salvioli S, Conte M, Blaauw
B, Zampieri S, Salviati L, Scorrano L, Sandri M. Age-Associated Loss of OPA1 in Muscle
Impacts Muscle Mass, Metabolic Homeostasis, Systemic Inflammation, and Epithelial
Senescence. *Cell Metab*. 2017;25:1374-1389.e6.

39. Darshi M, Taylor SS. Mitochondrial ChChD3 acts as a Scaffold for Mitofilin, Sam50 and
PKA. 2008.

40. An J, Shi J, He Q, Lui K, Liu Y, Huang Y, Sheikh MS. CHCM1/CHCHD6, Novel
Mitochondrial Protein Linked to Regulation of Mitofilin and Mitochondrial Cristae
Morphology *. *Journal of Biological Chemistry*. 2012;287:7411–7426.

41. Ding C, Wu Z, Huang L, Wang Y, Xue J, Chen S, Deng Z, Wang L, Song Z, Chen S.
Mitofilin and CHCHD6 physically interact with Sam50 to sustain cristae structure. *Scientific
reports*. 2015;5:1–11.

42. Ott C, Dorsch E, Fraunholz M, Straub S, Kozjak-Pavlovic V. Detailed analysis of the human
mitochondrial contact site complex indicate a hierarchy of subunits. *PloS one*.
2015;10:e0120213.

43. John GB, Shang Y, Li L, Renken C, Mannella CA, Selker JM, Rangell L, Bennett MJ, Zha J.
The mitochondrial inner membrane protein mitofilin controls cristae morphology. *Molecular
biology of the cell*. 2005;16:1543–1554.

44. Kondadi AK, Anand R, Reichert AS. Functional Interplay between Cristae Biogenesis,
Mitochondrial Dynamics and Mitochondrial DNA Integrity. *Int J Mol Sci*. 2019;20:4311.

45. Stephan T, Brüser C, Deckers M, Steyer AM, Balzarotti F, Barbot M, Behr TS, Heim G,
Hübner W, Ilgen P. MICOS assembly controls mitochondrial inner membrane remodeling
and crista junction redistribution to mediate cristae formation. *The EMBO journal*.
2020;39:e104105.

46. Torres MJ, Kew KA, Ryan TE, Pennington ER, Lin C-T, Buddo KA, Fix AM, Smith CA,
Gilliam LA, Karvinen S, Lowe DA, Spangenburg EE, Zeczycki TN, Shaikh SR, Neufer PD.
17β-Estradiol Directly Lowers Mitochondrial Membrane Microviscosity and Improves
Bioenergetic Function in Skeletal Muscle. *Cell Metabolism*. 2018;27:167-179.e7.

47. Kitajima Y, Ono Y. Estrogens maintain skeletal muscle and satellite cell functions. *Journal
of Endocrinology*. 2016;229:267–275.

48. Wang L, Hsu T, Lin H, Fu C. Drosophila MICOS knockdown impairs mitochondrial
structure and function and promotes mitophagy in muscle tissue. *Biology Open*.
2020;9:bio054262.

49. Wasilewski M, Semenzato M, Rafelski SM, Robbins J, Bakardjiev AI, Scorrano L. Optic
Atrophy 1-Dependent Mitochondrial Remodeling Controls Steroidogenesis in Trophoblasts.
Current Biology. 2012;22:1228–1234.

50. Sarzi E, Seveno M, Angebault C, Milea D, Rönnbäck C, Quilès M, Adrian M, Grenier J,
Caignard A, Lacroux A, Lavergne C, Reynier P, Larsen M, Hamel CP, Delettre C, Lenaers
G, Müller A. Increased steroidogenesis promotes early-onset and severe vision loss in
females with OPA1 dominant optic atrophy. *Human Molecular Genetics*. 2016;25:2539–
2551.

51. Chao de la Barca JM, Fogazza M, Rugolo M, Chupin S, Del Dotto V, Ghelli AM, Carelli V,
Simard G, Procaccio V, Bonneau D, Lenaers G, Reynier P, Zanna C. Metabolomics
hallmarks OPA1 variants correlating with their in vitro phenotype and predicting clinical
severity. *Hum Mol Genet*. 2020;29:1319–1329.

52. Abrigo J, Gonzalez F, Aguirre F, Tacchi F, Gonzalez A, Meza MP, Simon F, Cabrera D,
Arrese M, Karpen S, Cabello-Verrugio C. Cholic acid and deoxycholic acid induce skeletal
muscle atrophy through a mechanism dependent on TGR5 receptor. *Journal of Cellular
Physiology*. 2021;236:260–272.

53. Jeong J, Conboy IM. Phosphatidylserine directly and positively regulates fusion of myoblasts
into myotubes. *Biochem Biophys Res Commun*. 2011;414:9–13.

54. Dambrova M, Makrecka-Kuka M, Kuka J, Vilskersts R, Nordberg D, Attwood MM, Smesny
S, Sen ZD, Guo AC, Oler E, Tian S, Zheng J, Wishart DS, Liepinsh E, Schiöth HB.
Acylcarnitines: Nomenclature, Biomarkers, Therapeutic Potential, Drug Targets, and
Clinical Trials. *Pharmacol Rev*. 2022;74:506–551.

55. Stith JL, Velazquez FN, Obeid LM. Advances in determining signaling mechanisms of
ceramide and role in disease. *J Lipid Res*. 2019;60:913–918.

56. Feng Q, Xia W, Dai G, Lv J, Yang J, Liu D, Zhang G. The Aging Features of Thyrotoxicosis
Mice: Malnutrition, Immunosenescence and Lipotoxicity. *Front Immunol*. 2022;13:864929.

57. Johnson AA, Stolzing A. The role of lipid metabolism in aging, lifespan regulation, and
age□related disease. *Aging Cell*. 2019;18:e13048.

58. Paradies G, Paradies V, Ruggiero FM, Petrosillo G. Role of Cardiolipin in Mitochondrial
Function and Dynamics in Health and Disease: Molecular and Pharmacological Aspects.
Cells. 2019;8:728.

59. Shen Z, Ye C, McCain K, Greenberg ML. The Role of Cardiolipin in Cardiovascular Health.
Biomed Res Int. 2015;2015:891707.

60. Semba RD, Moaddel R, Zhang P, Ramsden CE, Ferrucci L. Tetra-linoleoyl cardiolipin
depletion plays a major role in the pathogenesis of sarcopenia. *Med Hypotheses*.
2019;127:142–149.

61. Singh AK, Singh R. Triglyceride and cardiovascular risk: A critical appraisal. *Indian J*
Endocrinol Metab. 2016;20:418–428.

62. McReynolds MR, Chellappa K, Chiles E, Jankowski C, Shen Y, Chen L, Descamps HC,
Mukherjee S, Bhat YR, Lingala SR, Chu Q, Botolin P, Hayat F, Doke T, Susztak K, Thaiss
CA, Lu W, Migaud ME, Su X, Rabinowitz JD, Baur JA. NAD+ flux is maintained in aged
mice despite lower tissue concentrations. *Cell Syst*. 2021;12:1160-1172.e4.

63. Sawicki KT, Chang H-C, Ardehali H. Role of heme in cardiovascular physiology and
disease. *J Am Heart Assoc*. 2015;4:e001138.

64. Dahl R, Larsen S, Dohlmann TL, Qvortrup K, Helge JW, Dela F, Prats C. Three-dimensional
reconstruction of the human skeletal muscle mitochondrial network as a tool to assess
mitochondrial content and structural organization. *Acta Physiol (Oxf)*. 2015;213:145–155.

65. Glancy B, Hartnell LM, Malide D, Yu Z-X, Combs CA, Connelly PS, Subramaniam S,
Balaban RS. Mitochondrial reticulum for cellular energy distribution in muscle. *Nature*.
2015;523:617–620.

66. Kho AT, Kang PB, Kohane IS, Kunkel LM. Transcriptome-scale similarities between mouse
and human skeletal muscles with normal and myopathic phenotypes. *BMC Musculoskelet*
Disord. 2006;7:23.

67. Garnier A, Fortin D, Delomenie C, Momken I, Veksler V, Ventura□Clapier R. Depressed
mitochondrial transcription factors and oxidative capacity in rat failing cardiac and skeletal
muscles. *The Journal of physiology*. 2003;551:491–501.

68. Lin I-H, Chang J-L, Hua K, Huang W-C, Hsu M-T, Chen Y-F. Skeletal muscle in aged mice
reveals extensive transformation of muscle gene expression. *BMC genetics*. 2018;19:1–13.

69. Romanick M, Thompson LV, Brown-Borg HM. Murine models of atrophy, cachexia, and
sarcopenia in skeletal muscle. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of*
Disease. 2013;1832:1410–1420.

70. Galpin AJ, Raue U, Jemiolo B, Trappe TA, Harber MP, Minchev K, Trappe S. Human
skeletal muscle fiber type specific protein content. *Anal Biochem*. 2012;425:175–182.

71. Hosotani M, Kametani K, Ohno N, Hiramatsu K, Kawasaki T, Hasegawa Y, Iwasaki T,
Watanabe T. The unique physiological features of the broiler pectoralis major muscle as
suggested by the three-dimensional ultrastructural study of mitochondria in type IIb muscle
fibers. *Journal of Veterinary Medical Science*. 2021;:21–0408.

72. Makida S, Kametani K, Hosotani M, Takahashi N, Iwasaki T, Hasegawa Y, Takaya T, Ueda
H, Watanabe T. Three-dimensional structural analysis of mitochondria composing each

subtype of fast-twitch muscle fibers in chicken. *Journal of Veterinary Medical Science*. 2022;advpub:22–0080.

73. Beasley HK, Wanjalla CN, Kirabo A, Hinton A. β2ARs: double edge sword in heart function. *Trends Mol Med*. 2023;:S1471-4914(23)00063–1.

74. Darshi M, Mendiola VL, Mackey MR, Murphy AN, Koller A, Perkins GA, Ellisman MH, Taylor SS. ChChd3, an Inner Mitochondrial Membrane Protein, Is Essential for Maintaining Crista Integrity and Mitochondrial Function *. *Journal of Biological Chemistry*. 2011;286:2918–2932.

75. Liu T, Woo JA, Bukhari MZ, LePochat P, Chacko A, Selenica MB, Yan Y, Kotsiviras P, Buosi SC, Zhao X. CHCHD10⎕regulated OPA1⎕mitofilin complex mediates TDP⎕43⎕induced mitochondrial phenotypes associated with frontotemporal dementia. *The FASEB Journal*. 2020;34:8493–8509.

76. Chung KW. Advances in Understanding of the Role of Lipid Metabolism in Aging. *Cells*. 2021;10:880.

77. Mignard V, Dubois N, Lanoé D, Joalland M-P, Oliver L, Pecqueur C, Heymann D, Paris F, Vallette FM, Lalier L. Sphingolipid distribution at mitochondria-associated membranes (MAMs) upon induction of apoptosis [S]. *Journal of Lipid Research*. 2020;61:1025–1037.

78. Ilacqua N, Anastasia I, Raimondi A, Lemieux P, de Aguiar Vallim TQ, Toth K, Koonin EV, Pellegrini L. A three-organelle complex made by wrappER contacts with peroxisomes and mitochondria responds to liver lipid flux changes. *Journal of Cell Science*. 2021;135:jcs259091.

79. Hogan KA, Zeidler JD, Beasley HK, Alsaadi AI, Alshaheeb AA, Chang Y-C, Tian H, Hinton AO, McReynolds MR. Using mass spectrometry imaging to visualize age-related subcellular disruption. *Frontiers in Molecular Biosciences*. 2023;10. Available at https://www.frontiersin.org/articles/10.3389/fmolb.2023.906606. Accessed April 8, 2023.

80. Cury-Boaventura MF, Pompéia C, Curi R. Comparative toxicity of oleic acid and linoleic acid on Jurkat cells. *Clinical Nutrition*. 2004;23:721–732.

81. Dencher NA, Frenzel M, Reifschneider NH, Sugawa M, Krause F. Proteome Alterations in Rat Mitochondria Caused by Aging. *Annals of the New York Academy of Sciences*. 2007;1100:291–298.

82. Kuter K, Kratochwil M, Berghauzen-Maciejewska K, Głowacka U, Sugawa MD, Ossowska K, Dencher NA. Adaptation within mitochondrial oxidative phosphorylation supercomplexes and membrane viscosity during degeneration of dopaminergic neurons in an animal model of early Parkinson's disease. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*. 2016;1862:741–753.

83. Ikon N, Ryan RO. Cardiolipin and mitochondrial cristae organization. *Biochimica et Biophysica Acta (BBA) - Biomembranes*. 2017;1859:1156–1163.

84. de La Barca JMC, Arrázola MS, Bocca C, Arnauné-Pelloquin L, Iuliano O, Tcherkez G,
Lenaers G, Simard G, Belenguer P, Reynier P. The metabolomic signature of Opa1
deficiency in rat primary cortical neurons shows aspartate/glutamate depletion and
phospholipids remodeling. *Scientific reports*. 2019;9:1–8.

85. Bocca C, Nzoughet JK, Leruez S, Amati-Bonneau P, Ferré M, Kane M-S, Veyrat-Durebex
C, de La Barca JMC, Chevrollier A, Homedan C. A plasma metabolomic signature involving
purine metabolism in human optic atrophy 1 (OPA1)-related disorders. *Investigative*
ophthalmology & visual science. 2018;59:185–195.

86. de La Barca JMC, Simard G, Sarzi E, Chaumette T, Rousseau G, Chupin S, Gadras C,
Tessier L, Ferré M, Chevrollier A. Targeted metabolomics reveals early dominant optic
atrophy signature in optic nerves of Opa1delTTAG/+ mice. *Investigative ophthalmology &*
visual science. 2017;58:812–820.

87. Garcia-Cazarin ML, Snider NN, Andrade FH. Mitochondrial isolation from skeletal muscle.
JoVE (Journal of Visualized Experiments). 2011;:e2452.

88. Wortel IMN, van der Meer LT, Kilberg MS, van Leeuwen FN. Surviving Stress: Modulation
of ATF4-Mediated Stress Responses in Normal and Malignant Cells. *Trends Endocrinol*
Metab. 2017;28:794–806.

89. Kondadi AK, Anand R, Hänsch S, Urbach J, Zobel T, Wolf DM, Segawa M, Liesa M,
Shirihai OS, Weidtkamp☐Peters S, Reichert AS. Cristae undergo continuous cycles of
membrane remodelling in a MICOS☐dependent manner. *EMBO Rep*. 2020;21:e49776.

90. Hessenberger M, Zerbes RM, Rampelt H, Kunz S, Xavier AH, Purfürst B, Lilie H, Pfanner
N, van der Laan M, Daumke O. Regulated membrane remodeling by Mic60 controls
formation of mitochondrial crista junctions. *Nat Commun*. 2017;8:15258.

91. Tarasenko D, Barbot M, Jans DC, Kroppen B, Sadowski B, Heim G, Möbius W, Jakobs S,
Meinecke M. The MICOS component Mic60 displays a conserved membrane-bending
activity that is necessary for normal cristae morphology. *J Cell Biol*. 2017;216:889–899.

92. Rowland AA, Voeltz GK. Endoplasmic reticulum–mitochondria contacts: function of the
junction. *Nat Rev Mol Cell Biol*. 2012;13:607–615.

93. Rieusset J. The role of endoplasmic reticulum-mitochondria contact sites in the control of
glucose homeostasis: an update. *Cell death & disease*. 2018;9:1–12.

94. Han J, Back SH, Hur J, Lin Y-H, Gildersleeve R, Shan J, Yuan CL, Krokowski D, Wang S,
Hatzoglou M. ER-stress-induced transcriptional regulation increases protein synthesis
leading to cell death. *Nature cell biology*. 2013;15:481–490.

95. Zhang X, Wang X, Yuan Z, Radford SJ, Liu C, Libutti SK, Zheng XFS. Amino acids-
Rab1A-mTORC1 signaling controls whole-body glucose homeostasis. *Cell Reports*.
2021;34:108830.

96. Piñero-Martos E, Ortega-Vila B, Pol-Fuster J, Cisneros-Barroso E, Ruiz-Guerra L, Medina-
Dols A, Heine-Suñer D, Lladó J, Olmos G, Vives-Bauzà C. Disrupted in schizophrenia 1
(DISC1) is a constituent of the mammalian mitochondrial contact site and cristae organizing
system (MICOS) complex, and is essential for oxidative phosphorylation. *Human Molecular*
Genetics. 2016;25:4157–4169.

97. Park Y-U, Jeong J, Lee H, Mun JY, Kim J-H, Lee JS, Nguyen MD, Han SS, Suh P-G, Park
SK. Disrupted-in-schizophrenia 1 (DISC1) plays essential roles in mitochondria in
collaboration with Mitofilin. *Proceedings of the National Academy of Sciences*.
2010;107:17785–17790.

98. Bannwarth S, Ait-El-Mkadem S, Chaussenot A, Genin EC, Lacas-Gervais S, Fragaki K,
Berg-Alonso L, Kageyama Y, Serre V, Moore DG, Verschueren A, Rouzier C, Le Ber I,
Augé G, Cochaud C, Lespinasse F, N'Guyen K, de Septenville A, Brice A, Yu-Wai-Man P,
Sesaki H, Pouget J, Paquis-Flucklinger V. A mitochondrial origin for frontotemporal
dementia and amyotrophic lateral sclerosis through CHCHD10 involvement. *Brain*.
2014;137:2329–2345.

99. Eramo MJ, Lisnyak V, Formosa LE, Ryan MT. The 'mitochondrial contact site and cristae
organising system' (MICOS) in health and human disease. *The Journal of Biochemistry*.
2020;167:243–255.

100.   Martin CR, Preedy VR, Rajendram R. Assessments, Treatments and Modeling in Aging
and Neurological Disease: The Neuroscience of Aging. Academic Press; 2021: 1-608.

101.   Barja G. The mitochondrial free radical theory of aging. *Progress in molecular biology*
and translational science. 2014;127:1–27.

102.   Kandlur A, Satyamoorthy K, Gangadharan G. Oxidative Stress in Cognitive and
Epigenetic Aging: A Retrospective Glance. *Frontiers in Molecular Neuroscience*. 2020;13.
Available at https://www.frontiersin.org/article/10.3389/fnmol.2020.00041. Accessed March
17, 2022.

103.   Boudina S, Sena S, Theobald H, Sheng X, Wright JJ, Hu XX, Aziz S, Johnson JI, Bugger
H, Zaha VG, Abel ED. Mitochondrial energetics in the heart in obesity-related diabetes:
direct evidence for increased uncoupled respiration and activation of uncoupling proteins.
Diabetes. 2007;56:2457–2466.

104.   Chippindale AK, Alipaz JA, Chen H-W, Rose MR. EXPERIMENTAL EVOLUTION OF
ACCELERATED DEVELOPMENT IN DROSOPHILA. 1. DEVELOPMENTAL SPEED
AND LARVAL SURVIVAL. *Evolution*. 1997;51:1536–1551.

105.   Graves JL Jr, Hertweck KL, Phillips MA, Han MV, Cabral LG, Barter TT, Greer LF,
Burke MK, Mueller LD, Rose MR. Genomics of Parallel Experimental Evolution in
Drosophila. *Molecular Biology and Evolution*. 2017;34:831–842.

106.   Vogler G, Ocorr K. Visualizing the Beating Heart in Drosophila. *J Vis Exp*. 2009;:1425.

107.   Neikirk K, Vue Z, Katti P, Shao J, Christensen TA, Lopez EG, Marshall A, Palavicino-
Maggio C, Ponce J, Alghanem A. Systematic Transmission Electron Microscopy-Based
Identification of Cellular Degradation Machinery. *bioRxiv*. 2021.

108.   Wende AR, O'Neill BT, Bugger H, Riehle C, Tuinei J, Buchanan J, Tsushima K, Wang
L, Caro P, Aili G, Sloan C, Kim BJ, Wang X, Pereira RO, McCrory MA, Nye BG,
Benavides GA, Darley-Usmar VM, Shioi T, Weimer BC, Abela ED. Enhanced cardiac
Akt/protein kinase B signaling contributes to pathological cardiac hypertrophy in part by
impairing mitochondrial function via transcriptional repression of mitochondrion-targeted
nuclear genes. *Molecular and Cellular Biology*. 2015;35:831–846.

109.   Vendelin M, Béraud N, Guerrero K, Andrienko T, Kuznetsov AV, Olivares J, Kay L,
Saks VA. Mitochondrial regular arrangement in muscle cells: a "crystal-like" pattern.
American Journal of Physiology-Cell Physiology. 2005;288:C757–C767.

110.   Yoon J, Daneshgar N, Chu Y, Chen B, Hefti M, Vikram A, Irani K, Song L, Brenner C,
Abel ED, London B, Dai D. Metabolic rescue ameliorates mitochondrial
encephalo□cardiomyopathy in murine and human iPSC models of Leigh syndrome. *Clin
Transl Med*. 2022;12:e954.

111.   Li B, Tang J, Yang Q, Li S, Cui X, Li Y, Chen Y, Xue W, Li X, Zhu F. NOREVA:
normalization and evaluation of MS-based metabolomics data. *Nucleic acids research*.
2017;45:W162–W170.

112.   Cantor JR, Abu-Remaileh M, Kanarek N, Freinkman E, Gao X, Louissaint Jr A, Lewis
CA, Sabatini DM. Physiologic medium rewires cellular metabolism and reveals uric acid as
an endogenous inhibitor of UMP synthase. *Cell*. 2017;169:258–272.

113.   Chong J, Soufan O, Li C, Caraus I, Li S, Bourque G, Wishart DS, Xia J. MetaboAnalyst
4.0: towards more transparent and integrative metabolomics analysis. *Nucleic acids
research*. 2018;46:W486–W494.

114.   Wang L, Xing X, Chen L, Yang L, Su X, Rabitz H, Lu W, Rabinowitz JD. Peak
Annotation and Verification Engine for Untargeted LC–MS Metabolomics. *Anal Chem*.
2019;91:1838–1846.

115.   Adusumilli R, Mallick P. Data Conversion with ProteoWizard msConvert. In: Comai L,
Katz JE, Mallick P, eds. Proteomics: Methods and Protocols. New York, NY: Springer;
2017: 339–368.

116.   Lu W, Wang L, Chen L, Hui S, Rabinowitz JD. Extraction and Quantitation of
Nicotinamide Adenine Dinucleotide Redox Cofactors. *Antioxidants & Redox Signaling*.
2018;28:167–179.

1484
1485
1486
1487 (**13**) **SUPPORTING INFORMATION**

1488   All data may be requested from the corresponding author.
1489
1490
1491   **(15) FIGURE LEGEND**
1492
Figure 1: Decreased mitochondrial size and volume in gastrocnemius, soleus, and cardiac
muscle of aged mice in SBF-SEM 3D reconstructions.
(**A-B**) Representative SBF-SEM orthoslice for male murine gastrocnemius, (**C-D**) soleus, and
(**E-F**) cardiac tissue. (**A'-B'**) 3D reconstructions of mitochondria (various colors) in
gastrocnemius, (**C'-D'**) soleus, and (**E'-F'**) cardiac tissue of three-month and 2-year mice
overlaid on ortho slices. (**A''-B''**) Pseudo-colored individual mitochondria in gastrocnemius,
(**C''-D''**) soleus, and (**E''-F''**) cardiac tissue differentiate micro-level changes. (**G-X**)
Quantification of 3D reconstructions, with each dot representing the overall average of all
mitochondria quantified for each mouse. (**G**) Mitochondrial volume in gastrocnemius muscle of
3-months and 2-year murine samples, and (**H**) mitochondrial volume distributed on basis of
percent of total mitochondria to visualize relative heterogeneity. (**I**) Mitochondrial 3D area in
gastrocnemius muscle of 3-months and 2-year murine samples, and (**J**) mitochondrial area
distributed on basis of percent of total mitochondria to visualize relative heterogeneity. (**K**)
Mitochondrial perimeter in gastrocnemius muscle of 3-months and 2-year murine samples, and
(**L**) mitochondrial perimeter distributed on basis of percent of total mitochondria to visualize
relative heterogeneity. These same quantifications are also displayed in (**M-R**) soleus and (**S-X**)
cardiac tissues. Cartoon representations of metrics to calculate (**Y**) mitochondrial volume, (**Z**)
perimeter, and (**AA**) perimeter. Approximately 550 mitochondria analyzed for each tissue type
and age cohort (n=3 mice per age cohort). Significance values **** represents $p \leq 0.0001$.
1512
1513
Figure 2: SBF-SEM 3D reconstruction in gastrocnemius, soleus, and cardiac muscle of
aged mice showing altered mitochondrial networks
Representative examples of 3D reconstruction of mitochondria in (**A**) gastrocnemius, (**B**) soleus,
and (**C**) cardiac tissue of three-month and 2-year mice ages organized by volume to show
phenotypes of mitochondria. (**D**) Mitochondrial complexity index (MCI), a measure analogous to
sphericity, in gastrocnemius muscle of 3-months and 2-year murine samples, and (**E**) MCI
distributed on basis of percent of total mitochondria to visualize relative heterogeneity. (**F**)
Sphericity in gastrocnemius muscle of 3-months and 2-year murine samples, and (**G**)
mitochondrial sphericity distributed on basis of percent of total mitochondria to visualize relative
heterogeneity. These same quantifications are also displayed in (**H-K**) soleus and (**L-O**) cardiac
tissues. Cartoon representations of metrics to calculate (**P**) MCI and (**Q**) sphericity.
Approximately 550 mitochondria analyzed for each tissue type and age cohort (n=3 mice per age
cohort). Significance values * $p \leq 0.05$, **** $p \leq 0.0001$.
1527
1528
Figure 3: Changes in Expression of *Opa1* and MICOS genes in Aging Mouse Tissue and
Drosophila
(**A-D**) Fold changes in the amount of *Opa1* and MICOS gene transcripts in mitochondria of
skeletal muscle of 3-month-old and 2-year-old mice as measured by RT-qPCR. (**A**) *Opa1*, (**B**)
Mitofilin, (**C**) *Chchd3*, and (**D**) *Chchd6* transcripts in skeletal muscle. (**E-H**) qPCR analyzing the

gene transcript fold changes of Opa-1 and MICOS across aging in cardiac tissue. (**E**) *Opa1*, (**F**)
Mitofilin, (**G**) *Chchd3*, (**H**) and *Chchd6* transcripts. (**I**) Altered protein expressions of MTCO1,
SAM50, Mitofilin, OPA1, and tubulin in gastrocnemius, (**J**) soleus, and (**K**) cardiac tissue. (**L-**
M) Altered *Drosophila* mitochondrial genes with age. (**L**) qPCR of *OPA1* and MICOS complex
components in aging Drosophila in thorax and (**M**) cardiac tissue. Significance values * $p \leq 0.05$,
** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$. For all qPCR experiments, n=6.
1540
Figure 4. Knockout of *Opa1*, *Mitofilin*, *Chchd3*, or *Chchd6* in myotubes results in changes
in structural changes of mitochondria and cristae in TEM and 3D reconstruction.
(**A-E**) Representative images of mitochondria and cristae from myotubes of *Opa1*, *Mitofilin*,
Chchd3, and *Chchd6*-knockout mice compared to WT. (**F**) Mitochondrial area in myotubes of
Opa1, *Mitofilin, Chchd3*, and *Chchd6*-knockout mice compared to WT. (**G**) Circularity index,
measuring the roundness and symmetry of mitochondria, in myotubes of *Opa1*, *Mitofilin,*
Chchd3, and *Chchd6*-knockout mice compared to WT. (**H**) The number of mitochondria in
myotubes of *Opa1*, *Mitofilin, Chchd3*, and *Chchd6*-knockout mice compared to WT. (**I**)
Quantification of individual cristae quantity in myotubes of *Opa1*, *Mitofilin, Chchd3*, and
Chchd6-knockout mice compared to WT. (**J**) Cristae scores measuring the uniformity and
idealness of cristae in myotubes of *Opa1*, *Mitofilin, Chchd3*, and *Chchd6*-knockout mice
compared to WT. (**K**) Surface area of the average cristae in myotubes of *Opa1*, *Mitofilin,*
Chchd3, and *Chchd6*-knockout mice compared to WT. (**L-P**) Representative images showing 3D
reconstructions of mitochondria in myotubes of *Opa1*, *Mitofilin*, *Chchd3*, and *Chchd6* knockout
mice compared to WT. (**Q**) Mitochondrial 3D length in myotubes of *Opa1*, *Mitofilin*, *Chchd3*,
and *Chchd6* knockout mice compared to WT. (**R**) Mitochondrial volume on log scale in
myotubes of *Opa1*, *Mitofilin, Chchd3*, and *Chchd6* knockout mice compared to WT.
(**S-V**) Altered *Drosophila* mitochondrial structure with loss of the MICOS complex and
mitochondrial proteins. (**S**) Actin-Mitochondria staining for *Drosophila* flight tissue across
knockout of MICOS complex and mitochondrial proteins. (**T**) Transmission electron microscopy
(TEM) quantification of changes in *Drosophila* flight tissue the mitochondrial area, (**U**)
circularity, and (**V**) quantity per sarcomere upon knockout of MICOS complex and
mitochondrial proteins. Significance values * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, ****
$p \leq 0.0001$. Dots represent number of mitochondria quantified.
1565
1566
Figure 5: Knockout of the MICOS complex in myotubes results in changes in oxygen
consumption rates and metabolomics.
(**A**) OCR in myotubes of *Opa1* and *Mitofilin*-knockout mice compared to WT. (**B**) Basal OCR,
the net respiration rate once non-mitochondrial respiration has been removed, in myotubes of
Opa1 and *Mitofilin*-knockout mice compared to WT. (**C**) ATP-linked respiration, shown from
intervals 4−7 in the OCR graphs, was determined by the addition of oligomycin (an inhibitor of
respiration), thus representing the respiration dependent on ATP, in myotubes of *Opa1* and
Mitofilin-knockout mice compared to WT. (**D**) Maximum OCR, represented by the peak from
intervals 7−11 once non-mitochondrial respiration was deducted, in myotubes of *Opa1* and
Mitofilin-knockout mice compared to WT. (**E**) The reserve capacity, which is represented by the
difference between basal OCR and maximum OCR, in myotubes of *Opa1* and *Mitofilin*-
knockout mice compared to WT. (**F**) Proton leak, representing non-phosphorylating electron
transfer, in myotubes of *Opa1* and *Mitofilin*-knockout mice compared to WT. (**G-J**)

Metabolomic analysis upon loss of *Mitofilin* is also shown. (**G**) Metabolite PCA and (**H**) T-test
comparing myotubes for control to *Mitofilin*-knockout mice. (**I**) Heatmap showing the relative
abundance of ions and (**J**) enrichment analysis of metabolites, which links together several
similarly functioning metabolites, with the relative abundance for *Mitofilin*-knockout.
(**K**) OCR in myotubes of *Chchd3, Chchd6*, and *Opa1* knockout mice compared to WT. (**L**) Basal
OCR in myotubes of *Chchd3*, *Chchd6*, and *Opa1* knockout mice compared to WT. (**M**) ATP-
linked respiration in myotubes of *Chchd3*, *Cchchd6*, and *Opa1* knockout mice compared to WT.
(**N**) Maximum OCR in myotubes of *Chchd3*, *Chchd6*, and *Opa1* knockout mice compared to
WT. (**O**) The reserve capacity in myotubes of *Chchd3*, *Chchd6*, and *Opa1* knockout mice
compared to WT. (**P**) Proton leak in myotubes of *Opa1*, *Chchd3*, and *Chchd6*, knockout mice
compared to WT. (**Q-T**) Metabolomic analysis upon loss of *Chchd3* or *Chchd6* is also shown.
(**Q**) Metabolite PCA and (**R**) ANOVA test comparing control to myotubes of *Chchd3* and
Chchd6 knockout mice (**S**) Heatmap showing the relative abundance of ions for control and (**T**)
enrichment analysis metabolite for *Chchd3* and *Chchd6* knockout mice. Significance values *
$p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$. For seahorse n=6 plates for experimental
knockouts, while for control n=16.

1596
Figure 6: Metabolomics analysis and lipidomic profiling reveals metabolic dysregulation
and disruptions in lipid classes with age in gastrocnemius, soleus, and cardiac muscles. (**A**)
Metabolic heatmap representing relative abundance of metabolites and (**B**) heatmaps
representing the lipidome in young and aged gastrocnemius, (**C-D**) soleus, and (**E-F**) cardiac
samples. For each tissue and metabolite in the heatmaps, aged are normalized to the median of
young, and then log2 transformed. Significantly differentiated lipid classes are those with
adjusted p-values < 0.05 (note: p-values were adjusted to correct for multiple comparisons using
and FDR procedure) and log fold changes greater than 1 or less than -1. Young, n=4; aged, n=4.
For all panels, *error bars* indicate S.E.M. ***, $p < 0.001$; **, $p < 0.01$; *, $p < 0.05$; §, $p < 0.1$,
calculated with Student's *t*-test.

1607
1608
1609 **Supplementary:**

1610
SFigure 1: Distribution of mitochondria quantifications among mice samples.
Mitochondrial volume in (**A**) gastrocnemius, (**B**) soleus, and (**C**) cardiac tissue of 3-months and
2-year murine samples distribution in each individual mouse to represent intra-individual
heterogeneity. This is also done for (**D-F**) mitochondrial area, (**G-I**) mitochondrial perimeter, (**J-**
L) mitochondrial complexity index, and (**M-O**) mitochondrial sphericity in gastrocnemius,
soleus, and cardiac tissue of 3-months and 2-year murine samples.

1617
1618 **SFigure 2**: Raw Western blots for gastrocnemius, soleus, and cardiac tissue of tubulin, OPA-1,
1619 and mitofilin.

1620
SFigure 3: Metabolomics and lipidomic analysis reveal metabolic dysregulation and
disruptions in lipid classes and chain lengths with age in gastrocnemius, soleus, and cardiac
muscles. (**A**) Volcano plot of metabolites that were differentially regulated in aged compared to
young gastrocnemius, (**B**) soleus, and (**C**) cardiac samples. For our volcano plots, the x-axis
represents the median and the y-axis represents the adjusted FDR. (**D**) Enrichment analysis

displaying enriched metabolites in aged gastrocnemius, (**E**) soleus, and (**F**) cardiac muscles. (**G**)
PCA plot for young and aged gastrocnemius, (**H**) soleus, and (**I**) cardiac samples. (**J**) Lipid class
fold-change ratios between young and old tissues in gastrocnemius, (**K**) soleus, and (**L**) cardiac
samples. Significantly differentiated lipid classes are those with adjusted p-values < 0.05 (note:
p-values were adjusted to correct for multiple comparisons using and FDR procedure) and log
fold changes greater than 1 or less than -1. (**M**) Lipid chain-length fold-change ratios between
young and old tissues in gastrocnemius, (**N**) soleus, and (**O**) cardiac samples. Significantly
differentiated lipid chain lengths are those with adjusted p-values < 0.05 (note: p-values were
adjusted to correct for multiple comparisons using and FDR procedure) and log fold changes
greater than 1 or less than -1.
1636
1637
1638
1639 **Tables:**
1640

Gene	ΔΔCt	Fold change
ATF-4	-0.781	**1.718**
Drp	-0.781	**2.980**
Marf	-1.764	**3.397**
Opa1	-2.388	**5.235**
MINOS1 (Mic10)	-0.385	**1.306**
QIL1 (Mic13)	-1.549	**2.926**
CHCHD3/6 (mic19)	-2.279	**4.853**
Dmic60 (mitofilin)	0.244	0.844
APOO (mic26/27)	-1.053	**2.075**
VDAC/Porin	-3.265	**9.612**
IP3R/Itpr	-0.997	**1.996**
FGF21/ bln	-0.140	1.102
Ire1	-1.219	**2.328**
Bip(grp78)/ Hsc 70-3	-2.259	**4.786**
ATF-6	0.072	0.951
GRP75/Hsc70-5	-1.203	**2.301**

1641
Table 1:
Drosophila qPCR flight tissue results of altered MICOS, mitochondrial, and MERCs proteins in
relative fold change across the aging process.
Fold change represents change in gene expression when comparing population A with population
C. Fold change values of >1.0 indicate that the gene is expressed more in the C populations (28-
day cycle) than in the A populations (9-day cycle) at day 21. Fold change values of <1.0 indicate

that the gene is expressed more in the A populations than in the C populations at day 21. Values
of ~1.0 indicate no difference between the populations. Values deviating from 1.0 by more than
0.20 are highlighted in yellow. This distinction is arbitrary and based on the researchers'
personal interpretation of significance.
1652
1653

Gene	ΔΔCt	Fold change
ATF-4	-0.595	1.510
Drp	0.010	0.993
Marf	0.032	0.978
Opa1	-0.428	1.346
MINOS1 (Mic10)	-0.452	1.368
QIL1 (Mic13)	0.012	0.992
CHCHD3/6 (mic19)	0.123	0.918
Dmic60 (mitofilin)	-1.791	3.460
APOO (mic26/27)	-0.012	1.008
VDAC/Porin	0.290	0.818
IP3R/Itpr	0.243	0.845
FGF21/ bln	0.927	0.526
Ire1	0.221	0.858
Bip(grp78)/ Hsc 70-3	-0.223	1.167
ATF-6	0.141	0.907
GRP75/Hsc70-5	-0.224	1.168

1654
Table 2: qPCR results from *Drosophila* cardiac tissue
Fold change represents change in gene expression when comparing population A with population
C. Fold change values of >1.0 indicate that the gene is expressed more in the C populations (28-
day cycle) than in the A populations (9-day cycle) at day 21. Fold change values of <1.0 indicate
that the gene is expressed more in the A populations than in the C populations at day 21. Values
of ~1.0 indicate no difference between the populations. Values deviating from 1.0 by more than
0.20 are highlighted in yellow. This distinction is arbitrary and based on the researchers'
personal interpretation of significance.
1663
1664
1665
1666 **Supplementary Tables:**
1667
1668 Supplementary Table 1. Lipid classes present in our dataset and associated abbreviations.

1669
1670   Supplementary Table 2. Results for set enrichment analysis, lipid class and chain length, based
1671   on lipidomic differentiation between young and old gastrocnemius tissue.
1672
1673   Supplementary Table 3. Results for set enrichment analysis, lipid class and chain length, based
1674   on lipidomic differentiation between young and old soleus tissue.
1675
1676   Supplementary Table 4. Results for set enrichment analysis, lipid class and chain length, based
1677   on lipidomic differentiation between young and old heart tissue.

Highest volume → Lowest volume

$$\text{Mitochondria Complexity Index} = \frac{Surface\ Area^3}{16\pi^2 Volume^2}$$

$$\text{Sphericity} = \frac{\pi^{1/3}(6*Volume)^{2/3}}{Surface\ Area}$$

Mice

A
Opa-1 mRNA fold change

C
Cdcdh3 mRNA fold change

C
Mitofilin mRNA Fold Change

3 Month   2 Month   2 year
** **

H
Chchd6 mRNA Fold Change

3 Month   3 Month   2 year
** ***

I

MTCO1 (56 kDa)
SAM50 (52)
Mic60/mitofilin (84)
OPA1 (80-100)
tubulin (50-55)

K
MTCO1 (56 kDa)

J
MTCO1 (56 kDa)

Thorax

Cardiac

L

Drosophila

Opa1 mRNA Fold Change
ACO   CO
**

Mic60 mRNA Fold Change
ACO   CO
ns

M

Opa1 mRNA Fold Change
ACO   CO
ns

Mic60 mRNA Fold Change
ACO   CO
*

Mic19 mRNA Fold Change
ACO   CO
***

Mic10 mRNA Fold Change
ACO   CO
ns

Mic13 mRNA Fold Change
ACO   CO
*

Mic19 mRNA Fold Change
ACO   CO
ns

Mic10 mRNA Fold Change
ACO   CO
ns

Mic13 mRNA Fold Change
ACO   CO
ns

Myotubes

Drosophila