The genomics and physiology of abiotic stressors associated with global elevation gradients

in *Arabidopsis thaliana*

Short Title: Global elevational clines in Arabidopsis

Diana Gamba[1*], Claire Lorts[1], Asnake Haile[1,2], Seema Sahay[3], Lua Lopez[1,4], Tian Xia[1],

Margarita Takou[1], Evelyn Kulesza[1,5], Dinakaran Elango[5,6], Jeffrey Kerby[7], Mistire Yifru[2],

Collins E. Bulafu[8], Tigist Wondimu[2], Katarzyna Glowacka[3*], Jesse R. Lasky[1]

[1] Department of Biology, Pennsylvania State University, University Park, PA 16802, USA

[2] Department of Plant Biology and Biodiversity Management, Addis Ababa University, Addis

Ababa, Ethiopia

[3] Department of Biochemistry and Center for Plant Science Innovation, University of Nebraska-

Lincoln, Lincoln, NE 68588, USA

[4] Department of Biology, California State University San Bernardino, San Bernardino, CA

92407, USA

[5] Department of Plant Science, Pennsylvania State University, University Park, PA 16802, USA

[6] Department of Agronomy, Iowa State University, Ames, IA 50011, USA

[7] Aarhus Institute for Advanced Studies, 1630, Høegh-Guldbergs Gade 6B, Aarhus, Denmark

8000

[8] Department of Plant Sciences, Microbiology, and Biotechnology, Makarere University,

Kampala, Uganda

*Corresponding authors: dxg5484@psu.edu, kglowacka2@unl.edu

Abstract

Phenotypic and genetic diversity in *Arabidopsis thaliana* may be associated with adaptation

along its wide elevational range. We took a multi-regional view of elevational adaptation and in

a diverse panel of ecotypes measured plant responses to high elevation stressors: low partial CO_2

pressure, high light, and night freezing. We conducted genome-wide association studies (GWAS)

and found evidence of contrasting locally adaptive clines between regions. Western

Mediterranean ecotypes showed low $\delta^{13}C$/early flowering at low elevations to high $\delta^{13}C$/late

flowering at high elevations, while Asian ecotypes showed the opposite pattern. We mapped

different candidate genes for each region, and trait-associated SNPs often showed elevational

clines likely maintained by selection. Antioxidants and pigmentation showed regional

differentiation but rarely elevational clines. GWAS for antioxidants identified an ascorbate

transporter *PHT4;4* (AT4G00370), which we show alters non-photochemical quenching kinetics

under high light and may be involved in local adaptation to Moroccan mountains. The low-

antioxidant *PHT4;4* GWAS allele was associated with lower *PHT4;4* expression and this

haplotype was characterized by binding sites of a transcription factor family, DOF, involved in

light response. Our results highlight how physiological and genomic elevational clines in

different regions can be unique, underlining the complexity of local adaptation in widely

distributed species.

Introduction

Changes in environmental conditions with changing elevation are some of the most iconic

natural gradients (Caldas, 1966). Correspondingly, changes in plant growth forms are dramatic

over hundreds or thousands of meters of elevation (Caldas, 1966; Hedberg and Hedberg, 1979;

Körner, 2021). Elevational gradients are among the key factor driving local adaptation, and

hence diversity, within species that have broad elevational ranges (Clausen et al., 1940; Kooyers

et al., 2015). At high elevations in particular, the combination of cold and high light during

vegetative periods poses a major challenge. Additional environmental changes from sea level to

alpine environments include wider fluctuations in daily temperature, decreased partial CO_2

pressure (pCO_2), altered (i.e., orographic) precipitation and clear sky radiation, and shorter

growing seasons (Körner, 2003, 2007). While local adaptation to high elevation conditions has

been detected in reciprocal transplants and common gardens (Villemereuil et al., 2018; Wos et

al., 2022), its physiology and genetic basis is less clear.

76   Photosynthesis sits at the nexus of multiple physiological challenges for high

elevation plants. For example, pCO_2 declines approximately linearly with elevation, at 4000 m

being only ~65% of sea level pCO_2 (Körner, 2021). Species from high elevations may exhibit

higher fitness and better regulation of physiological processes under low atmospheric pCO_2

(Ward and Strain, 1997; Zhu et al., 2010). However, pCO_2 limitation is likely counteracted by

elevational reductions in pO_2 and photorespiration (Wang et al., 2017). Additionally, clear sky

radiation increases with elevation, but cold temperatures slow the dark reactions of

photosynthesis, leading to photo-oxidative damage if excess energy from photons is not properly

handled (Wise, 1995). One mechanism to counteract photo-oxidative damage is non-

photochemical quenching (NPQ) in which excess absorbed photons dissipate via the xanthophyll

cycle (García-Plazaola et al., 2012; Muller et al., 2001). When light intensity rapidly increases

(e.g., due to clearing cloud cover), rapid acclimation of NPQ (fast NPQ kinetics) may be

adaptive (Rungrat et al., 2019). Another mechanism is the increased production of antioxidants

(e.g., carotenoids and flavonoids), which can scavenge reactive-oxygen-species (Wise, 1995).

Alpine environments can also experience extreme temperature fluctuations, often reaching below

freezing temperatures every night, which not only exacerbates oxidative damage but can also

physically damage cells (Suzuki and Mittler, 2006).

The model plant *Arabidopsis thaliana* (hereafter Arabidopsis) has a broad geographical, environmental, and elevational range, from below sea level to ~4400 m, making it an ideal system for studying multi-regional elevational adaptation. Several studies have addressed elevational adaptation in Eurasia, though in single mountain systems. For example, in the Pyrenees low elevation/coastal sites experience early summer drought, potentially selecting for a drought-escape strategy (Montesinos⧠Navarro et al., 2011, 2012; Picó, 2012; Vidigal et al., 2016; Wolfe and Tonsor, 2014). In the Alps, flowering time and its gene expression may contribute to adaptation to seasonal freezing variation along elevation (Suter et al., 2014), but physiological clines are unclear (Günther et al., 2016; Lampei et al., 2019; Luo et al., 2015a). In the western Himalayas flowering time can be plastic in response to temperature, especially in ecotypes from high elevation (Singh and Roy, 2017). Little is known about elevational clines in African Arabidopsis, where populations are older than most Eurasian ones, have higher genetic diversity, and grow under the highest recorded elevations (up to ~4400 m) in the unique Afroalpine ecosystem (Brennan et al., 2014; Brochmann et al., 2021; Durvasula et al., 2017).

One challenge in deriving general conclusions about adaptive responses to elevation for widespread species is that elevational changes in environment often differ from region to region, potentially leading to region-specific local adaptations. For example, while temperature consistently decreases with elevation, different regions may show different elevational clines in cloudiness, resulting in fluctuating light intensity that complicates elevational light patterns in different geographical regions (Körner, 2021). Even when gradients are consistent, local adaptation to isolated mountain ranges can occur via shared or distinct genetic and phenotypic mechanisms (Bohutínská et al., 2021). Furthermore, plant adaptation to extreme conditions often comprises complex multivariate responses where homeostatic regulation during photosynthesis is achieved in combination with changes in phenology (Fernández⧠Marín et al., 2020; Kooyers et al., 2015; Smith and Knapp, 1990).

Here, we took a multi-regional view of Arabidopsis genomes, life history, and physiology across elevational gradients. We asked whether putatively adaptive physiological and genomic elevational clines were consistent among regions and hypothesized that ecotypes from higher elevations have robust performance in the face of high elevation stressors (i.e., better growth and resource use, lower oxidative stress) using a diverse set of naturally inbred lines (ecotypes). We integrate large-scale experiments and genome-wide association studies (GWAS)

with detailed physiological panels, revealing the complexity of elevational adaptation across the

range of Arabidopsis. For a GWAS candidate gene (*PHT4;4*), we explore the physiological

effects of natural and mutant allelic variants in response to high light. With our regional

approach in a widespread model plant, we show that different mountains exhibit different

strategies of local adaptation along elevation.

Results

Population genetic structure and regional patterns in climate

To investigate natural phenotypic and genetic variation across elevational gradients in different

geographical regions, we studied a total of 271 ecotypes, including 262 from native regions in

Eurasia and Africa (Figure 1 A, Supplemental Dataset S1). We defined five geographical regions

based on location and genetic cluster (Supplemental Figure S1): Asia (46 ecotypes), central

Europe/Caucasus (59 ecotypes), northwestern Europe (26 ecotypes), western Mediterranean (112

ecotypes), and eastern Africa (19 ecotypes, including 9 sequenced). The last region encompasses

the poorly studied genetically diverse Afroalpine Arabidopsis populations (Fulgione and

Hancock, 2018).

140   We examined population genetic structure in our diversity panel using a published

variant call format (VCF) file (Durvasula et al., 2017) that we combined with 16 ecotypes (eight

phenotyped) from eastern Africa that we sequenced here (Supplemental Methods S1). We

calculated pairwise genetic distances of 261 ecotypes (Supplemental Dataset S1, 'sequenced'

column) from 562,194 whole genome resequencing single nucleotide polymorphisms (SNPs)

filtered for minor allele frequency (MAF<0.05) and linkage-disequilibrium (LD). A neighbor-

joining tree showed that population genetic structure largely corresponded to our geographical

regions and not elevation. Ecotypes from low and high elevations on the same mountain range

were more closely related than ecotypes of similar elevations on different mountains. In the

Caucasus, however, ecotypes from two of our regions overlapped (Asia and central

Europe/Caucasus), highlighting the genetic heterogeneity of this region (Figure 1 B).

151   A principal components analysis (PCA) of 85 climatic variables from the native

range of 257 studied ecotypes with accurate provenance/climate data (Supplemental Dataset S2)

showed that ecotypes from different regions occupied distinct climates and climatic extremes

(Figure 1 C). Asia had prominent temperature seasonality, with short growing seasons and cold

winters. Central Europe/Caucasus had milder temperature seasonality and warm growing

seasons. Northwestern Europe was cold with mild dry seasons. In the western Mediterranean

temperature variation was wide, with dry seasons at high elevation ranging from hot, dry, and

bright (Atlas Mountains) to temperate and moist (Pyrenees). East Africa occupied the most

distinct climate of long growing seasons with sharp day/night temperature changes. Elevational

range and extent differed among regions, but environmental gradients were generally similar,

with lower night temperatures and shorter growing seasons at higher elevations (Figure 1 D, E).

Figure 1 Provenance, genetic relationships, and climates of studied native ecotypes colored by
region. Symbols represent elevation below (closed circles) and above (open circles) 1000 m for
all regions, except for eastern Africa where the threshold is 4000 m. **A** Geographic location of
ecotypes. **B** Neighbor-joining tree based on 261 ecotypes and 562,194 resequencing SNPs
filtered for LD. Tips with no symbol are ecotypes outside of the native range. **C** PC1 and PC2 of
climate space representing conditions in 257 ecotypes based on 85 climatic variables. Arrows
and text summarize loadings more strongly correlated with PC1 and PC2, indicating the direction
of the relationship (not proportional to eigenvectors). PAR=photosynthetically active radiation,

PET=potential evapotranspiration, AUS=Austria, ETH=Ethiopia, MAR=Morocco, RUS=Russia,
SWE=Sweden, TAN=Tanzania, UGA=Uganda. **D**, **E** Linear regressions of climatic variables on
elevation of origin. Selected variables describe the growing season. Significant relationships ($p <$
0.01) are depicted by their R^2.
175
Trait variation under low pCO2, high light and night-time freezing conditions

To characterize genetic variation in physiology, resource use, and performance under alpine-

associated stressors (*i.e.*, trait differences among ecotypes in our diversity panel), we performed

three separate experiments (12 h photoperiod) with subsets (111–254) of the 271 studied

ecotypes (Supplemental Dataset S1). Subsets were different based on seed and space availability

and germination success, with a total of 101 ecotypes overlapping between experiments. Our

first experiment imposed low partial CO_2 pressure (pCO_2; with cool and bright conditions) on

253 ecotypes (Supplemental Dataset S3). Our next two experiments imposed contrasting light

levels (250 vs 600 μmol m^{-2} s^{-1}) on 111–114 ecotypes (performed simultaneously under cool

conditions) and contrasting night-time temperatures (–4 vs 4ºC) on 170–172 ecotypes (performed

sequentially under bright conditions) (Supplemental Datasets S4–S5).

In our first experiment we measured δ^{13}C and δ^{15}N, two stable isotope ratios

associated with carbon/water and nitrogen use, respectively, and integrated across the plant life

span, root diameter, rosette diameter, and rosette compactness. We examined coordinated life

history variation (*i.e.*, in life cycle and physiology) with a PCA of these traits along with

published flowering time data (Alonso-Blanco et al., 2016) (N=206 ecotypes). Asian ecotypes

tended to have smaller and more compact rosettes with smaller roots and higher δ^{13}C and δ^{15}N,

in contrast to western Mediterranean ecotypes which showed greater phenotypic variation,

including larger and less compact rosettes with larger roots lower δ^{13}C and δ^{15}N (Figure 2 A).

For flowering time, we found high variation within regions, with generally later flowering in

northwestern Europe (likely due to strict vernalization requirements), and earlier flowering in

central Europe/Caucasus (likely due to weak flowering time cold response genes), while Asia

and the western Mediterranean were not different to each other (Supplemental Figure S2 A).

This high within-region diversity was also apparent at the level of some major known flowering

time genes (Supplemental Figure S2 B).

In our next experiments, we found significant trait changes between treatments

and trait differences among regions (ANOVA, $p<8E–10$; Figure 2) in fresh aboveground

biomass, total antioxidant activity, and the multidimensional color variation of leaves (high light

experiment) and their extracts (night-time freezing experiment) based on a PCA of RGB color

values (Figure 2 B, C, Supplemental Results S1). Color may indicate phenotypes connected to

elevational adaptation because pigments can be adaptive as they act as antioxidants under high

light and/or cold (Havaux and Kloppstech, 2001).

Under cool temperatures and high light, plants were smaller, had higher

antioxidant activity, and dark purple (vs. green) leaves (lower values of $PC1_{LeafColor}$/higher values

of $PC2_{LeafColor}$) than when grown under low light (Figure 2 B, D–F). Survival rates were slightly

but significantly lower under high than low light (93 vs 96%, $p=0.02$), and positively correlated

with rosette biomass (Pearson correlation coefficient (cor)=0.23, $p=0.01$) under high light.

Furthermore, ANOVA showed significant differences in antioxidant activity at high light among

regions ($p=0.004$), with western Mediterranean ecotypes having the lowest values (Figure 2 E),

suggesting lower sensitivity to high light.

Under night-time freezing, plants were larger, had higher antioxidant activity, and

more orange (as opposed to yellow) leaf extracts (lower values of $PC1_{ExtractColor}$) than when

grown with night-time temperatures just above freezing (Figure 2 C, G–I). Survival was

significantly lower under night-time freezing than under no freezing (76 vs 100%, $p<0.0001$),

and in the western Mediterranean survival was negatively correlated with higher antioxidants

under night freezing (*i.e.*, antioxidant under night freezing - antioxidant under no freezing; cor=

–0.38, $p=0.001$), suggesting maladaptive plasticity. Additionally, ANOVA showed significant

differences in $PC1_{ExtractColor}$ under night-time freezing between regions (i.e., orange in extracts,

$p=0.007$), with eastern Africa ecotypes (all from >2775 m) having yellower extracts (Figure 2 I).

In eastern Africa more orange extracts (lower values of $PC1_{ExtractColor}$) were associated with

higher survival rates (cor= –0.92, $p=0.001$, df=6), suggesting selection associated with pigments.

Taken together, ecotypes showed great variation in flowering time and in

response to high elevation experimental conditions. Ecotypes from different geographical

regions showed distinct morphology under low pCO_2 (bright and cool conditions) and strong

treatment effects on all phenotypes measured under high vs. low light, and under night-time

freezing vs. night-time above freezing. We detected some differences in phenotypes among

regions, suggesting decreased sensitivity in antioxidants to high light in western Mediterranean

ecotypes (Figure 2 E), and decreased content in leaf extract greenness under night-time freezing

in eastern African ecotypes (Figure 2 I).

Figure 2 Ecotype responses to high elevation experimental conditions colored by region. **A** Eigenvector plot of the loadings of 8 variables (d: diameter, c: compactness) onto PC1 and PC2 of phenotypes under low pCO_2 along with flowering time (FT at 10ºC: FT10 and at 16ºC: FT16) and flowering time plasticity (FTp: FT10–FT16) in 206 ecotypes (5–6 replicates). **B–C** PC1 and PC2 of RGB colors in experiment 2 (from rosette photos of 110 ecotypes-5 replicates under high light: HL vs. low light: LL) and experiment 3 (from leaf-extract photos of 163 ecotypes-6 pooled replicates under night-time freezing: Frz vs. no night-time freezing: no Frz), respectively. **D–I** Significant phenotypic plasticity under contrasting light (center: Asia N=23, C Europe/Caucasus N=17, NW Europe N=7, W Mediterranean N=63) and night-time temperatures (bottom: Asia N= 27, C Europe/Caucasus N=38, NW Europe N= 10, E Africa N=13, W Mediterranean N=75). Boxplots are proportional to sample size and depict the median and interquartile range, whiskers cover the data extent. **D, G** Fresh aboveground biomass (mg). **E, H** Antioxidant activity (ug per leaf mg). **F, I** PC1 of RGB color from leaves and leaf extracts, respectively.

Genetic variation in NPQ kinetics under cold and drought

Non-photochemical quenching (NPQ) kinetics in response to fluctuating light (e.g., due to cloud changes) can be an important mechanism for limiting photooxidative damage at high elevations (Kumar et al., 2021). To capture diverse strategies, we selected a global subset of 11 ecotypes representing the lowest (22–254 m) and highest (1576–4078 m) elevational range of Arabidopsis (Supplemental Dataset S1). We performed detailed measurements of NPQ kinetics under fluctuating light-dark cycles in response to two potential high elevation stressors: cold (22 vs. 4°C under well-watered conditions for 17 h) and drought (daily-watering vs. 5-day-drought under 22°C). We fitted exponential equations to individual NPQ time series to obtain parameters that describe NPQ induction and relaxation (Sahay et al., 2023, Supplemental Figure S3).

We found that cold and drought significantly affected NPQ induction and relaxation, but this effect significantly varied between ecotypes (ANOVA, Supplemental Table S1, Supplemental Figures S4–S5). For example, CYR from low and seasonally dry France showed a slow NPQ response to cold but a fast response to drought via rapid NPQ induction (Figure 3 A–B). Conversely, Dja-1 from high and cold Kyrgyzstan showed no changes in NPQ kinetics in response to drought but a strong response to cold via sustained high NPQ after fluctuating light treatments (2nd and 3rd cycles; Figure 3 C–D), which could be important to avoid high light stress under cold. Sij-1 from high and cold Uzbekistan showed a similar pattern (Figure 3 G–H). In IP-Trs-0 from low and mesic Iberia, cold and drought had similarly minor effects on NPQ kinetics (Figure 3 E–F), suggesting reduced sensitivity to these abiotic stressors. Interestingly, under control conditions, IP-Trs-0 and JL011-2-1, an ecotype with Afroalpine origin where nights are subject to below freezing temperatures all-year-round, had the lowest fresh weight and the highest dry matter content (>60%), indicating low water content (Supplemental Figure S6), which could aid in drought-avoidance or in freezing-tolerance. Additionally, under control conditions, JL011-2-1 had the lowest chlorophyll and total carotenoid content, but the highest chlorophyll ($a + b$) to carotenoids ratio (Supplemental Table S2).

Taken together, while there was no clear pattern relative to elevation of origin across this global sample of ecotypes, responses to drought or cold in NPQ kinetics greatly varied among ecotypes, with responses potentially corresponding to the environmental pressures at their sites of origin.

Figure 3 Four representative examples of non-photochemical quenching curves during three sequential light-dark cycles under cold (left panels) and drought (right panels) with corresponding control. Chosen ecotypes represent the lowest and highest elevations of the Arabidopsis range: **A–B** CYR (52 m). **C–D**. Dja-1 (2995 m). **E–F** IP-Trs-0 (254 m). **G–H** Sij-1 (2459 m). Bars are means ± SE (5–10 replicates).

Regional elevational clines in phenotype and the genomic basis of phenotypic variation

To understand how different elevational gradients shape phenotypic (Figure 4) and genomic variation, we tested for region-specific trait- and SNP-elevation clines with linear mixed models that controlled for genome-wide similarity among ecotypes (kinship), which when significant suggest that selection is driving clines (Josephs et al., 2019). GWAS that controlled for kinship (Supplemental Datasets S6–S9) were performed on our global dataset of 19 phenotypes (N=110–253 ecotypes, 2.7–2.8M SNPs), on regional datasets for six phenotypes that significantly varied with elevation (N=44–112, 1.9–2.8M SNPs), and on regional datasets for flowering time at 10 and 16ºC and flowering time plasticity (*i.e.*, flowering time at 10ºC minus flowering time at 16 ºC; N=67–398, 2–2.6M SNPs). We report candidate protein coding genes whose annotated function relates to phenotype function (Table 1) and highlight regional QTL for which top SNP significantly changed with elevation in mixed models that controlled for kinship.

Figure 4 Linear regressions on elevation of origin of flowering time (FT) and phenotypes under low pCO$_2$, high light, and night-time freezing. Ecotypes and fitted lines are colored by region: Asia N=20–60, central Europe/Caucasus N=17–264, eastern Africa N=5–11, northwestern Europe N=7–331, and western Mediterranean N=64–174. Significant relationships from linear

304 mixed effects kinship models (p<0.05) are depicted by their likelihood-ratio based pseudo-R^2. **A**
305 Days to flower at 10ºC (FT10) and 16ºC (FT16) obtained from the 1001 Genomes Consortium.
306 **B–F** Phenotypes measured in a low pCO_2 growth chamber. **G–I** Selected phenotypes measured
307 in a high light growth chamber. J–L. Selected phenotypes measured in a night-time freezing
308 growth chamber.
309
310

Evidence of adaptive elevational clines in Asia. Using mixed models that accounted for kinship

among ecotypes we found evidence that several traits showed elevational clines maintained by

selection. Flowering time (at 10 and 16ºC), $\delta^{13}C$, $\delta^{15}N$, rosette compactness, and plasticity in leaf

extract color in response to night-time freezing (*i.e.*, the Euclidean distance in PC 1-2 space

between contrasting conditions for a given ecotype, Figure 2 C), decreased with elevation (all

p<0.03). Conversely, flowering time plasticity, rosette diameter, stomatal density, biomass under

high and low light, and survival rates under high light and night-time freezing, increased with

elevation (all *p*<0.025; Figure 4 A–G). We found corroborating evidence in our growth chamber

experiments. Under high light stress, flowering time (at 10 and 16ºC) was negatively correlated

with both biomass and survival (cor ranged from –0.45 to –0.59, all *p*<0.04), while $\delta^{13}C$ was

negatively correlated with survival (cor= –0.46, *p*=0.03). Thus, high elevation ecotypes from this

region seemingly showed a life history strategy of rapid phenology but inefficient water and

nitrogen use, where accelerated reproduction and low water use efficiency (WUE) under high

light stress may enhance biomass and survival.

GWAS mapped quantitative trait loci (QTL) for $\delta^{13}C$, rosette compactness, and

rosette diameter, that showed elevational variation potentially maintained by selection (*i.e.*, after

controlling for kinship, p<0.05), although for $\delta^{13}C$ and rosette diameter no SNP passed

FDR<0.05 in the GWAS (Figure 5 A, D, G). For $\delta^{13}C$, the 25[th] top QTL was *FLOWERING*

LOCUS M (*FLM*; AT1G77080), a negative regulator of flowering in response to cold, where

loss-of-function removes the response to cool (≤10ºC) temperatures and results in earlier

flowering (Jin et al., 2022). For rosette compactness, the top QTL was a GDSL-like protein

(AT5G41890) which is poorly studied and belongs to a family of lipolytic enzymes involved in

morphogenesis and development (Lai et al., 2017). For rosette diameter, the 3[rd] top QTL was

downstream of a secreted peptide (*STMP5*; AT1G65490) that functions in plant growth and

potentially disease defense (Yu et al., 2019).

Figure 5 Genome-wide association studies (GWAS), geographic distribution, and variation of GWAS SNP alleles for selected traits and associated SNPs that showed elevational variation in Asia: $\delta^{13}C$ (top panels), rosette compactness (middle panels), and rosette diameter (bottom panels). **A**, **D**, **G** Manhattan plots over the five chromosomes of the Arabidopsis nuclear genome with hit SNPs marked in red and dotted lines corresponding to a genome-wide Bonferroni threshold. Right panels are zoomed-in on candidate gene QTL (surrounded by dashed lines), showing their orientation (+: forward, –: reverse) and associated SNP marked in red. SNPs were filtered for MAF=0.05, N=45 ecotypes and 1,984,688 SNPs for $\delta^{13}C$ and N=46 ecotypes and 2,002,562 SNPs for rosette compactness and diameter. **B**, **E**, **H** Map of GWAS SNP alleles in Asia. **C**, **F**, **I** Variation of GWAS SNP alleles (0: alternate allele, 2: reference allele) relative to traits of interest (left, with GWAS FDR), and elevation (right, with *p*-value from a kinship mixed model). Boxplot width is proportional to allele frequency and depict the median and interquartile range, whiskers extend to the data extremes.

Allelic variants in the aforementioned candidate genes involved in high $\delta^{13}C$, and compact/small rosettes, were restricted to low elevations (linear-mixed kinship model *p*=0.000003 for chr. 1: 28957836 bp, *p*<0.000001 for chr. 5: 16766371 bp, *p*=0.02 for chr. 1: 24356159 bp, respectively), suggesting that changing selection with elevation on *FLM*, the

GDSL-like enzyme, and *STMP5* expression contributes to elevational clines in $\delta^{13}C$ via

flowering time variation in response to cold and in rosette morphology via growth and

development. Although these SNPs were not found among the top 100 in our flowering time

GWAS, the high $\delta^{13}C$/rosette compactness variants were significantly associated with delayed

flowering at 10ºC (Wilcox-tests $p=0.002$ for chr. 1: 28957836 bp and $p=0.003$ for chr. 5:

16766371 bp).

Evidence of adaptive elevational clines in the western Mediterranean. We found evidence for

clines maintained by selection in flowering time (at 10 and 16ºC), $\delta^{13}C$, $\delta^{15}N$, root diameter,

rosette compactness, and biomass under high and low light, which all increased with elevation

(all $p\leq0.03$; Figure 4 A–G). Additionally, stomatal density was positively correlated with $\delta^{13}C$

(cor=0.39, $p=0.02$) and flowering time (at 10 and 16ºC: cor=0.49 and 0.45, $p=0.007$ and 0.02,

respectively), suggesting that stomatal patterning varies with life history and water use strategies.

Thus, high elevation ecotypes from this region apparently displayed a strategy of delayed

phenology and efficient water and nitrogen use, directed toward root and aboveground biomass

accumulation.

We found elevational variation in GWAS QTL for flowering time and $\delta^{13}C$ (all

FDR<0.05; Figure 6). For flowering time, the top QTL (at 16ºC) and the 2nd top QTL (at 10ºC)

was *DELAY OF GERMINATION 1* (*DOG1*; AT5G45830), an extensively studied gene involved

in regulating seed dormancy and flowering time (Alonso-Blanco et al., 2016; Graeber et al.,

2014; Huo et al., 2016; Martínez-Berdeja et al., 2020). For $\delta^{13}C$, the top and the 7th top QTL

were adjacent (downstream and upstream, respectively) of two genes involved in regulation of

abscisic acid (ABA) in response to osmotic stress. The first is *NICOTINAMIDASE 3* (*NIC3*;

AT5G23220), which encodes an enzyme that prevents the accumulation of intracellular

nicotinamide by converting it into nicotinic acid in the NAD+ pathway, and which is expressed

via Repressor of Silencing 1 (ROS1) DNA demethylation in response to ABA (Kim et al., 2019).

The other is *MAP3 kinase* (*MAP3k*; AT5G11850), which is involved in the phosphorylation of

SnRK2 kinases that in turn trigger osmotic-stress and ABA responses (Takahashi et al., 2020).

Figure 6 Genome-wide association studies (GWAS), geographic distribution, and variation of GWAS SNP alleles for selected traits and associated SNPs that showed elevational variation in the western Mediterranean: flowering time (top panels) and δ^{13}C (middle and bottom panels). **A**, **D** Manhattan plots over the five chromosomes of the Arabidopsis nuclear genome with hit SNPs marked in red and dotted lines corresponding to a genome-wide Bonferroni threshold. Right panels are zoomed-in on candidate gene QTL (surrounded by dashed lines), showing their orientation (+: forward, −: reverse) and associated SNP marked in red. SNPs were filtered for MAF=0.05, N=178 ecotypes and 2,663,439 SNPs for flowering time (at 16℃) and N=110 ecotypes and 2,831,054 SNPs for δ^{13}C. **B**, **E**, **G** Map of GWAS SNP alleles in the western Mediterranean. **C**, **F**, **H** Variation of GWAS SNP alleles (0: alternate allele, 2: reference allele) relative to traits of interest (left, with GWAS FDR), and elevation (right, with *p*-value from a kinship mixed model). Boxplot width is proportional to allele frequency and depict the median and interquartile range, whiskers extend to the data extremes.

SNP alleles involved in late flowering and high δ^{13}C were restricted to high elevation (linear-mixed kinship model *p*=0.000007 for chr5: 18590591 bp and *p*=0.02 for chr5: 7818730 bp, although *p*=0.07 for chr5: 3821491 bp, respectively), suggesting changing selection with elevation on *DOG1*, *NIC3* and *MAP3k* contributes to elevational clines in flowering time via seed dormancy variation and in δ^{13}C via the ABA pathway and stomatal dynamics. In Iberia,

403   allelic variants associated with low δ^{13}C/rapid flowering occurred in coastal and lowland areas

404   with late-spring heat and drought (Figure 6), where ecotypes appear to display a drought-escape

405   strategy (Montesinos-Navarro et al., 2011; Wolfe and Tonsor, 2014).

406   The *DOG1* SNP (chr5: 18590591 bp) was previously identified by Martínez-

Berdeja et al. (2020) because of its association with seed dormancy in response to cold. The late

flowering allele tags some (15 of 52 in Iberia and 15 of 68 in Morocco) of the ecotypes carrying

the widespread ancestral haplotype of the *DOG1* self-binding domain (ECCY), but this SNP did

not segregate in ecotypes with other haplotypes, nor in ECCY ecotypes from Asia (which only

carry the other GWAS SNP allele). Delayed flowering was more frequent in ecotypes carrying

the ECCY haplotype in all regions except in Asia (Kruskal-Wallis *p*=0.8 in Asia and *p*<0.0001

for all other regions; Supplemental Figure S7). Ecotypes carrying ECCY behave as winter

annuals: seeds germinate soon after dispersal in the late summer–early fall (if not, extended cold

induces secondary dormancy), plants overwinter as small rosettes, and flower over the spring

(i.e., delayed flowering after vernalization) (Martínez-Berdeja et al., 2020). Our finding indicates

that in Asia, delayed flowering is not exclusive to ecotypes carrying the ECCY overwintering

haplotype, but also occurs in ecotypes carrying the D-SY haplotype where seeds only germinate

after cold exposure (Martínez-Berdeja et al., 2020).

420   Martínez-Berdeja (2020) noted global climatic associations between *DOG1*

haplotypes, including elevation, but did not characterize the direction of elevational clines. Here

we found that *DOG1* haplotype frequency significantly changes with elevation in all regions

(Kruskal-Wallis *p*<0.01), except in central Europe/Caucasus (*p*=0.3; Supplemental Figure S7).

Relative to other haplotypes, ECCY was associated with higher elevations in the western

Mediterranean, but with lower elevations in Asia and northwestern Europe. In Asia, the more

frequent D-SY *DOG1* haplotype (Supplemental Figure S7) or mutations at other genes could

also confer delayed flowering, which is the low elevation strategy in this region (opposite the

western Mediterranean). Thus, the role of *DOG1* in local adaptation along elevation likely differs

from region to region due to alternate life-history clines among regions.

Latitudinal but not elevational clines detected in northwestern Europe. We found no

significant trait-elevational clines (all FDR>0.05). However, the elevational gradient in this

group was narrow, occupying the lowest part of the range: 0–500 m, with only one ecotype at

434 1060 m. Similar to the latter two regions, flowering time (at 10 and 16ºC) was positively

435 correlated with δ^{13}C and δ^{15}N (cor=0.52–0.68, p=0.01–0.0006), while negatively correlated with

436 rosette diameter (cor= –0.50 and –0.45, p= 0.02 and 0.04, respectively). We examined changes in

437 phenotype along latitude, given that winters become longer and colder at higher latitudes, which

438 may in turn relate to cold stress response. We found that delayed flowering and high δ^{13}C were

439 associated with higher latitudes (p<0.0001) in mixed models controlling for kinship among

440 ecotypes, suggesting that overwintering at higher latitudes is adaptive via delayed flowering and

441 high-water use efficiency. We did not perform GWAS on measured phenotypes for this region,

442 given their small sample size (up to 26), but report flowering time QTL (see Supplemental

443 Results S2).

445 ***Potentially adaptive elevational clines in central Europe/Caucasus but no clear QTL.*** Mixed

446 models that accounted for kinship among ecotypes showed that flowering time (at 10 and 16ºC)

447 and rosette compactness increased with elevation as in the western Mediterranean (all p<0.05;

448 Figure 4 A, D). Stable isotopes did not vary with elevation in this region, but as in other regions

449 both δ^{13}C and δ^{15}N were positively correlated to flowering time at 10 and 16ºC (cor=0.31–0.41,

450 p=0.01–0.001). Although flowering time was not correlated to leaf color traits under stress,

451 plasticity in leaf color in response to high light was lower (i.e., lower photosensitivity) at higher

452 elevations (linear-mixed model, p=0.0001, Figure 4 I). We did not detect QTL mapped with

453 GWAS that seemed functionally relevant for rosette compactness, and report one flowering time

454 QTL that was, however, not associated with elevation (see Supplemental Results S2).

456 ***Elevational clines in eastern Africa.*** Afroalpine ecotypes were only included in the night-time

457 freezing experiment and represented the highest part of the Arabidopsis range (mostly 3691–

458 4374 m, with one ecotype at 2775 m). Although based on a limited sample size of seven

459 ecotypes, unlike other regions, here a mixed model that accounted for kinship found that

460 antioxidant activity under night-time freezing increases with elevation (p=0.005; Figure 4 K).

462 ***Genotypes and phenotypes of montane Moroccan ecotypes under high light.*** In our global

463 dataset, we mapped two QTL associated with antioxidant activity under high light stress (Figure

464 7 A). The top tagged *Ferric Reduction Oxidase 7* (*FRO7*; AT5G49740) and was upstream (in the

putative promoter region) of *FRO6* (AT5G49730), both involved in iron chloroplast uptake and expressed in aerial green tissue (Jain et al., 2014; Jeong et al., 2008, 2009). *FRO6* expression is light-dependent, with several light-responsive elements in the promoter region (Feng et al., 2006). The second top QTL was downstream of a phosphate transporter (*PHT4;4*; AT4G00370) that carries ascorbate into the chloroplast (Miyaji et al., 2015). *PHT4;4* expression is higher in the chloroplast envelope membrane and increases under high light stress, with knockouts having lower levels of reduced ascorbate (an antioxidant) in the leaves, and lower content of ascorbate-dependent xanthophylls and ß-carotene after high light exposure (Miyaji et al., 2015).

Figure 7 Genome-wide association study (GWAS), geographic distribution, and variation of GWAS SNP alleles associated with antioxidant activity under high light. **A** Manhattan plots over the five chromosomes of the Arabidopsis nuclear genome with hit SNPs marked in red and dotted lines corresponding to a genome-wide Bonferroni threshold. Bottom panels are zoomed-in on candidate gene QTL (surrounded by dashed lines), showing their orientation (+: forward, –: reverse) and associated SNP marked in red. SNPs were filtered for MAF=0.05, N=110 ecotypes and 2,865,792 SNPs. **B**, **D** Map of GWAS SNP alleles in the western Mediterranean. **C**, **E** Variation of GWAS SNP alleles (0: alternate allele, 2: reference allele) in the western Mediterranean relative to antioxidant activity under high light (left, with GWAS FDR), elevation (center, with *p*-value from a kinship mixed model), and spring photosynthetically active radiation (PAR) (left, with *p*-value from a kinship mixed model). Boxplot width is proportional to allele frequency and depict the median and interquartile range, whiskers extend to the data extremes.

To account for potential under-fitting of the genomic background effect (kinship random effects) for the genetically distinct Moroccan ecotypes (Atlas Mountains in Figure 1 B) that also had low antioxidant activity (Figure 2 E, 4 H), we performed another GWAS on this

490   trait incorporating a covariate that indicated if the accession was from Morocco. P-values on this

491   GWAS were higher (FDR>0.9) and previously mapped QTL were no longer mapped. The

492   original GWAS results were thus driven by the low antioxidant activity of most Moroccan

493   ecotypes.

494   Variants in top SNPs (chr5: 20206030 bp for *FRO6/FRO7* and chr4: 160735 bp

for *PHT4;4*) associated with low antioxidant activity were globally uncommon (allele frequency

AF=0.09 and 0.08, respectively), but dominant (AF=0.81 and 0.84) in montane (>1000 m)

Moroccan ecotypes, and infrequent (AF=0.13 for chr5: 20206030 bp) to rare (AF=0.03 for chr4:

160735 bp) in Iberian ecotypes. Furthermore, these SNPs were strongly linked in Morocco

(cor=0.36, *p*=0.004), but not elsewhere (cor= –0.01, *p*=0.8). In the western Mediterranean, the

respective low-antioxidant GWAS SNP alleles were more frequent in ecotypes from sites with

higher photosynthetically active radiation during the growing season (Kruskal-Wallis *p*=0.0001

for chr4: 160735 bp and Kruskal-Wallis *p*=0.003 for chr5: 20206030 bp), but only for chr4:

160735 bp the low-antioxidant allele was dominant at higher elevations (Kruskal-Wallis *p*=

0.004), but these patterns were not significant in mixed-linear kinship models (Figure 7 C, E).

505   To further assess evidence of selection on these loci, we calculated F_{ST} values for

506   all SNPs with MAF>0.05 in chromosome 4 (where *PHT4;4* is located) and 5 (where

507   *FRO6/FRO7* are located), comparing Moroccan ecotypes to the global panel. We found F_{ST}=0.19

508   for chr4: 160735 bp and was in the 0.005 tail for the SNPs on the chromosome, and Chr 5:

509   20206030 had a similar pattern: F_{ST}=0.14 and was in the 0.01 tail for the SNPs on the

510   chromosome, suggesting these variants might be locally adaptive in Morocco.

511   Using published transcriptome data on 727 diverse ecotypes under typical low

light laboratory conditions (Klepikova et al., 2016), we found that *PHT4;4* expression was

weakly associated with allelic variation at chr4: 160735 bp in the western Mediterranean, where

the low-antioxidant GWAS SNP allele was associated with lower expression (cor= –0.18,

p=0.02), but not globally (cor<0.01). Likewise, *FRO7* expression (but not *FRO6*) was weakly

associated with allelic variation in chr5: 20206030 bp in the western Mediterranean (cor= –0.19,

p=0.01), but not globally (cor=0.07).

518   The *PHT4;4* antioxidant SNP covaried with our color phenotypes in a way

consistent with Miyagi et al. (2015)'s findings on xanthophylls and ß-carotene. The variant

associated with low antioxidant activity was significantly associated with lighter leaves and leaf

extracts (i.e., higher values of $PC1_{LeafColor}$ and $PC1_{ExtractColor}$) in our high light (Wilcox-tests

$p=0.004$), night-freezing ($p=0.02$) and no night-freezing experiments ($p=0.007$), suggesting the

low-antioxidant *PHT4;4* variant is associated with low carotenoids. We did not find this

association in the low light treatment ($p=0.12$), consistent with *PHT4;4* expression being

upregulated under high light (Miyagi et al., 2015).

Physiological effects of PHT4;4 mutations. We further investigated the phenotypic effects of

PHT4;4 mutations in response to high light (vs. low light) under cold using *PHT4;4* insertion

lines (CS444342, Salk_082875, and CS26443) in two different genetic backgrounds (*i.e.*, wild-

type: Col-0 for the first two mutants and Landsberg erecta (LE; for the third) (Supplemental

Table S3) and a subset of western Mediterranean ecotypes that segregate at the *PHT4;4* GWAS

SNP (Supplemental Dataset S1).

We found that the speed of NPQ induction was significantly different among

ecotypes (ANOVA, $p<0.001$), being on average 12% slower in ecotypes with the low antioxidant

GWAS allele compared to ecotypes with the other allele ($p=0.006$; Figure 8 A and B). In

agreement with Miyagi et al. (2015)'s results on *PHT4;4* knockouts, our results were conditioned

on high light exposure: the slower NPQ induction in ecotypes with the low-antioxidant allele was

only detected when plants were exposed for 2 days to high light but not when plants

continuously grew under low light (Figure 8 A). Accordingly, ecotypes with the low-antioxidant

GWAS allele had a 27% average reduction in *PHT4;4* expression under high light compared to

ecotypes with the other allele ($p=0.001$; Figure 8 B).

Furthermore, a prolongation of high light exposure to 4 days combined with cold

(day and night at 6°C) revealed that ecotypes with the low-antioxidant GWAS allele resembled

PHT4;4 insertion lines in their NPQ kinetics. For instance, the steady state of NPQ under light

(*NPQasymptote$_{light}$*) and the range of NPQ change under dark (*NPQamplitude$_{dark}$*) were 15%

lower ($p=0.01$) in ecotypes with the low-antioxidant allele vs. ecotypes with the other allele, and

9% ($p=0.02$) and 16% ($p=0.008$) lower, respectively, in the *PHT4;4* knockout CS26443 vs. its

LE wild-type (Figure 8 C).

Figure 8 Effects of *PHT4;4* variation in NPQ kinetics and gene expression. **A** Speed of NPQ induction (*time constant*$_{light}$; time needed to achieve 63% of the maximum capacity of NPQ) for plants grown under high light (HL: 550 µmol m^{-2} s^{-1}) vs. low light (LL; 200 µmol m^{-2} s^{-1}) at room temperature (5–6 replicates per ecotype), with ANOVA *p*-values indicating significant differences among ecotypes (left panel) and between ecotype groups with different GWAS SNP alleles (middle panel). **B** Significant downregulation in normalized *PHT4;4* expression in ecotypes with the GWAS SNP allele vs. ecotypes with the other allele when grown under HL (5–6 replicates per ecotype). **C** NPQ response to light and dark cycles for plants grown under HL and cold (6ºC). *PHT4;4* knockouts CS26443, Salk_082875, and CS444342, are preceded by corresponding wild-type LE and Col-0. Genotype means are depicted with standard errors based on 4–6 replicates. Asterisks indicate significant differences between ecotypes with different GWAS SNP alleles or between mutants with their corresponding wild-type based on a Dunnett's test adjusted to multiple comparisons (*p*≤0.05).

In a separate experiment of plants exposed for 7 days to high vs. low light (day and night at 4ºC), the phenotypic effects of different *PHT4;4* variants were not as congruent

between natural variation and insertion lines (Supplemental Results S3). Under high light, all

western Mediterranean ecotypes had the lowest antioxidant activity, brightest/most transparent

leaf extracts, and lowest chlorophyll content (Supplemental Figure S8 B–D). *PHT4;4* insertion

lines showed lower chlorophyll content conditioned on high light exposure relative to their

respective wild-type, consistent with our observation that in western Mediterranean ecotypes

with the low-antioxidant GWAS allele, chlorophyll content was generally lowest (Figure S8).

However, insertion lines and wild-type did not differ in antioxidant levels (ANOVA *p*=0.3–0.5)

while neither did western Mediterranean ecotypes with different GWAS SNP alleles (ANOVA

p=0.7).

575   Taken together, ecotypes with the low-antioxidant GWAS allele behaved

similarly to *PHT4;4* insertion lines under high light, in that they had low *PHT4;4* gene

expression, slow NPQ kinetics, and lower chlorophyll content. However, *PHT4;4* insertions did

not affect total antioxidants, which were higher compared to the low antioxidant activity in our

subset of western Mediterranean ecotypes. This inconsistency could be due to a *cis*-regulatory

natural variant (among other explanations) of *PHT4;4* having distinct phenotypes from *PHT4;4*

knockouts.

Potential cis-regulatory variants in the Moroccan PHT4;4 haplotype. To look for potential

causal *cis*-regulatory variants, we scanned for enriched sequence motifs in the 3.5 kbp upstream

and 3.5 kbp downstream region of *PHT4;4* in 63 ecotypes (61 western Mediterranean, 1 from

Japan, and 1 from Czech Republic) with the low-antioxidant GWAS SNP allele/haplotype

(Supplemental Figure S9 A), in contrast to 53 closely related ecotypes with the high-antioxidant

allele (Supplemental Dataset 1). We discovered 54 significantly enriched motifs (p<0.05) in the

low-antioxidant allele ecotypes, one of which was lacking in the high-antioxidant allele group,

six of which were fixed, and 46 of which were present in more than 90% of the ecotypes with the

low antioxidant allele. When we cross referenced the discovered motif sequences with the plant

JASPAR (Castro-Mondragón et al., 2022) database of experimentally defined plant transcription

factor binding sites (TFBS), the 54 discovered motifs in the low-antioxidant allele group had a

significant overlap with 339 known TFBS. The motif lacking in the high-antioxidant allele group

overlapped with four known TFBS in the HD-ZIP family (MA0990.1, MA1212.1, MA1369.1,

MA1375.1). Members of this family function in regulating stress response and plant growth

(Sharif et al., 2021). Amongst the most common TFBS across the discovered motifs were

members of the DOF family, with DOF1.7, DOF5.8 and DOF6.3 being present in 20%, 18.5%

and 15.4% of the total discovered motifs respectively (Table 2, Supplemental Figure S9 B, C).

Members of this family function in regulating growth, development, and biotic and abiotic stress

response (Zou and Sun, 2023). Lastly, motifs in the MIKC TFBS family were common in more

than 10% of the discovered motifs (Table 2). Members of this family function in regulating

cyclical developmental processes in response to cold (Castelán-Muñoz et al., 2019).

Discussion

Arabidopsis exhibits natural genetic variation in abiotic stress response, which may be linked to

phenological and physiological adaptation along elevational gradients. Germination and

flowering time show evidence of local adaptation in response to seasonal cold at high elevations

and to early drought at low elevations (Méndez-Vigo et al., 2011; Suter et al., 2014; Vidigal et

al., 2016). Water use efficiency (WUE) and its genetic basis may be associated with phenology

(Kenney et al., 2014; Lovell et al., 2013), though changes in WUE with elevation have been less

studied (DeLeo et al., 2020; Des Marais et al., 2014; Dittberner et al., 2018; Mckay et al., 2003;

Wolfe and Tonsor, 2014). Elevational clines related to dwarfism evolution at high elevations

have been identified (Luo et al., 2015a, b; Singh and Roy, 2017; Tripathi et al., 2019; Tyagi et

al., 2016). However, past studies have typically focused on individual geographic regions in

Eurasia, precluding a broader understanding of adaptation to global elevational gradients.

617   Here we examined the range-wide genomics and physiology of adaptation to

elevation of Arabidopsis using a multi-regional approach informed by phylogeography. We

uncovered elevational clines indicating changing selection for several traits and genomic

variants. Most clines were region-specific, despite similarity between regions in elevational

environmental gradients, where differences in phenology are likely part of a coordinated

phenology-physiology strategy that differs between regions. Antioxidant activity, pigmentation,

and NPQ kinetics showed substantial genetic variation. These traits may be particularly

important to local adaptation to Moroccan mountains. We showed that even when selective

environmental gradients are consistent among mountain ranges, patterns of multivariate

adaptation to elevation and their genetic basis can be unique to each mountain range (see

Tusiime et al., 2020; Wos, et al. 2022). This could be due to phenotypic and genetic

heterogeneity (sensu López-Arboleda et al., 2021) across different regions affected by subtle

differences in selective gradients and phenotypic clines despite similar environmental clines, or

by the historical contingency of each region. Our regional approach in a widespread model plant

highlights that a better integration of plant adaptation at a range-wide scale will help improve

evolutionary predictions (*e.g.*, in globally important crops or in invasive species) under the

current scenario of accelerated anthropogenic global change.

Coordinated life history and physiology along elevation In Asia, high elevation ecotypes

showed a putatively adaptive strategy of rapid and plastic phenology (*i.e.*, more rapid at 16ºC

than at 10ºC) and low water and nitrogen use. High elevation genotypes with low $\delta^{13}C$ tended to

harbor mutations in *FLM*, a flowering time regulator, suggesting a potential pleiotropic effect of

flowering time genetic control on water use efficiency in Asia. Previous studies have found

genotypic correlations between low WUE and earlier flowering both due to variation across the

genome (Kenney et al., 2014) and specifically to pleiotropic effects of a flowering time regulator

(*FRI*) (Lovell et al., 2013). Furthermore, studies in the western Himalayas suggest that increasing

cold and light intensity with elevation influences life history and genomic variation (Singh and

Roy, 2017; Tripathi et al., 2019; Tyagi et al., 2016). Here, two ecotypes from high elevation

central Asia showed a strong increase of NPQ kinetics after cold exposure. High light and low

pCO_2 induces drought stress in leaves via ABA signaling and stomatal dynamics (Kuromori et

al., 2022; Moustakas et al., 2022), while high light and cold increases the risk of photo-oxidation

(Wise, 1995). We hypothesize that in Asia high elevation ecotypes exhibit a multivariate stress-

escape strategy involving rapid NPQ and a fast 'spring annual' life cycle.

In the western Mediterranean, high elevation ecotypes showed a putatively

adaptive strategy of delayed flowering and high water and nitrogen use efficiency, suggesting

drought avoidance. By contrast, low elevation ecotypes had open rosettes associated with a

rapid-cycling strategy (early flowering) potentially allowing drought escape (Ludlow, 1989). The

genetic basis of this cline is likely partly caused by allelic variation in the seed

dormancy/flowering time *DOG1* gene (Graeber et al., 2014; Huo et al., 2016), and in two

candidate genes related to abiotic stress response in the ABA pathway, *MAP3k* and *NIC3* (Kim et

al., 2019; Takahashi et al., 2020). Previous studies in the Pyrenees have shown that earlier

flowering relates to drought-escape in the lowlands (Montesinos□Navarro et al., 2009, 2011;

Wolfe and Tonsor, 2014), and our results indicate that this strategy may involve evolution of

ABA signaling. Ecotypes from high elevation have alleles that allow them to counteract water

loss under stress, which in turn is adjusted with a 'winter-annual' life cycle, but not those from

low elevations which are probably less commonly exposed to environmental stress because of

their rapid cycle during wet periods (Picó 2012).

Traits associated with protection from photo-oxidation Ecotypes showed shared increases in

antioxidants and changes in leaf color in response to bright light and night-time freezing, but

little associations with elevation of origin. Photo-oxidation is also a threat at lower elevations

due to drought, potentially resulting in no clear change in selection (Nakabayashi et al., 2014).

Furthermore, while changes in antioxidants and pigments mediate high light acclimation (Aarabi

et al., 2023), they might not indicate cold acclimation (Distelbarth et al., 2013), but could be

related to freezing tolerance via flavonoid accumulation (Hannah et al., 2016). The combination

of cold and high light can thus comprise diverse metabolic and genomic pathways, that could be

region-specific (see next section).

Afroalpine populations occupy an outlier environment with respect to clear sky

radiation and night-time cold throughout the year. Here, antioxidant activity appeared to increase

with elevation under night-time freezing. Afroalpine ecotypes also had yellower leaf extracts

compared to dark orange and deep red extracts in other regions, potentially indicating less

xanthophylls used in NPQ, a hypothetical reduced sensitivity to cold (Havaux and Kloppstech,

2001). More orange extracts and higher plasticity in extracts color were associated with higher

survival under night-time freezing, suggesting potential selection on pigment content.

Additionally, a high elevation Afroalpine ecotype was very small, had little total carotenoid

content, with slower but greater NPQ under cold, and distinctly high leaf dry matter content

under controlled conditions. These characteristics could be adaptive if they enhance freezing

resistance (Gorsuch et al., 2010).

We found significant genetic variation in NPQ kinetics parameters in response to

cold and drought (i.e., ecotypes were significantly different in their response), but this variation

was not clearly related to elevation of origin. Cold resulted in slower NPQ induction and

relaxation across genotypes, while drought resulted in highly variable NPQ induction and

relaxation among ecotypes. NPQ can remove over 75% of absorbed light energy, thus its fast

induction in response to stress can play an essential role for rapid defense against damage from

photoinhibition (Malnoë, 2018). Slow relaxation of NPQ (particularly evident in two Asian

highland ecotypes) could limit NPQ and plant growth, but provide better protection to cold

(Kromdijk et al., 2016). We hypothesize that the generally slower NPQ speed in cold vs. drought

is due to temperature-dependent changes in the activity of both NPQ-related enzymes,

violaxanthin de-epoxidase and *zeaxanthin* epoxidase (Nilkens et al., 2010).

Natural and knockout *PHT4;4* variants differ in NPQ induction and gene expression under

high light Under bright and cool conditions, western Mediterranean ecotypes (mostly from

Morocco) had the lowest antioxidant activity, chlorophyll content, and slow NPQ kinetics. These

ecotypes have almost exclusively alleles near a transporter gene for ascorbate into the chloroplast

(*PHT4;4*) associated with low antioxidant activity under high light, and from the subset we

tested under high light, they showed a significant downregulation of *PHT4;4* expression relative

to ecotypes with the other allele. Previous studies (Miyaji et al., 2015) showed that the level of

PHT4;4 expression increases with light exposure. *PHT4;4* knockouts under high light had slower

NPQ induction, lower levels of reduced ascorbate in leaves, and lower levels of xanthophyll and

beta-carotene (Miyaji et al., 2015). Here we show that under high light and cold, *PHT4;4*

knockouts and western Mediterranean ecotypes with the low-antioxidant GWAS allele had lower

NPQ and chlorophyll content. Ascorbate is an antioxidant and cofactor used in the xanthophyll

cycle which in turn aids induction of NPQ. The western Mediterranean ecotypes carrying the

distinct *PHT4;4* haplotype linked to our GWAS SNP, might be tolerating high light and cold

conditions by maintaining a low but steady NPQ speed involving downregulated *PHT4;*4, while

reducing oxidative risk by producing less chlorophyll by, for example, reducing light-harvesting

antenna pigments of Photosystem II (Havaux and Kloppstech, 2001), which could incur a cost

under lower light conditions. This strategy might allow to limit photo-oxidative damage

independently from antioxidant activity in the bright and seasonally cold Atlas Mountains.

716   Furthermore, we showed that the western Mediterranean low-antioxidant GWAS

PHT4;4 haplotype (compared to related ecotypes lacking this haplotype) has putative *cis*-

regulatory motifs enriched primarily for transcription factor binding sites (TFBS) in the DOF

zinc-finger family. This family is involved in hormonal response and photoperiodic regulation

during plant development and under stress (Zou and Suri, 2023). DOF response elements are

present in the promoter region of several cold-induced genes which expression is exclusively

upregulated in response to light exposure under cold (Soitamo et al., 2008). We also identified

motifs associated with TFBS in the MIKC family, which are involved in floral development and

are key components of gene regulatory networks involved in stress response and developmental-

phenological plasticity in response to seasonal cold (Castelán-Muñoz et al., 2019). Furthermore,

our analysis revealed a motif only present in the western Mediterranean low-antioxidant GWAS

PHT4;4 haplotype, that overlapped with four known TFBS in the HD-ZIP family. Genes

involved with this family are critical for plant growth and development under stressful

environmental conditions (Sharif et al., 2021). We believe the evidence may indicate a *cis-*

regulatory variant at *PHT4;4* that reduces its expression under high light causing slower NPQ

kinetics. This phenotype might be part of a more conservative growth strategy that does not

require rapid NPQ kinetics, possibly due to lower chlorophyll levels in ecotypes from the region.

Finally, *PHT4;4* knockouts might not show antioxidant phenotypes due to

epistasis with other genes in western Mediterranean ecotypes, like the top GWAS QTL *FRO6/7*,

an iron transporter with similar allele frequency to the *PHT4;4* SNP. Nam et al. (2021) found

prevention of downregulation of photosynthesis related genes via ascorbate accumulation by

PHT4;4 induced by transcription factors responsive to iron deficiency. We hypothesize that the

interaction between *PHT4;4* and *FRO6/7* transcription factors in response to high light and cold

could influence the low antioxidant and chlorophyll content of western Mediterranean ecotypes

with downregulated *PHT4;4*. Indeed further validation studies with mapping populations or

transgenics targeting the specific natural variation found in the western Mediterranean are

needed. Our findings highlight the complexity of investigating the underpinnings of the

functional genomic and phenotypic variation in natural systems.

Materials and Methods

Plant material We selected a diverse set of 271 ecotypes (i.e., naturally inbred genotypes;

Supplemental Dataset S1) which we collectively call our diversity panel: 232 from the 1001

Genomes Project (ABRC stocks), 21 from Morocco and Tanzania (NASC stocks), and 18 newly

collected lines (8 of them here sequenced; Supplemental Methods S1) along elevational gradients

in Ethiopia (in 2017) and Uganda (in 2018) ranging from 3691–4374 m. We specifically targeted

several regional elevational gradients with this diversity panel, which we subsequently classified

into **five regions** following genetic cluster (according to Alonso-Blanco et al., 2016 and

Durvasula et al., 2017, see Supplemental Figure S1) and geographic origin:

1) Asia (46 ecotypes; 45 in the Asia genetic cluster + 1 admixed from Kazakhstan).

2) central Europe/Caucasus (59 ecotypes; 16 in the Central Europe cluster + 37 in the

Italy-Balkan-Caucasus cluster + 6 admixed from Germany, Italy, Lebanon, Switzerland,

and Turkey).

3) northwestern Europe (26 ecotypes; 6 in the Germany genetic cluster + 3 in the North

Sweden cluster + 9 in the South Sweden cluster + 4 in the Western Europe cluster + 4

admixed from the Netherlands and the United Kingdom).

4) western Mediterranean (112 ecotypes; all from Iberia and Morocco: 13 in the Relict

genetic cluster + 53 in the Spain cluster + 9 in the Western Europe cluster + 22 in the

Moroccan clusters + 15 admixed from Spain).

5) eastern Africa (19 ecotypes, only 9 sequenced; 16 from Ethiopia + 2 from Uganda + 1

from Tanzania).

The remaining accessions were from North America and Japan, thus designated as unassigned,

along with one accession from Cape Verde given how different this location is from all other

regions (9 ecotypes). With subsets of our diversity panel, we conducted three large-scale growth

chamber experiments (Experiment 1–3 below, Supplemental Dataset S1), and a detailed study on

NPQ kinetics in response to cold and drought. When trait measurements (explained below) could

be averaged across biological replicates, we obtained breeding values of phenotypes from the

fixed effects of linear mixed models that estimated the trait mean per ecotype with 'tray' as a

random factor in the R lme4 package. All phenotypes were measured in adult pre-flowering

plants.

Seed bulking and initial growing conditions Seeds from 271 ecotypes were bulked after dark-

stratification in distilled water for 4 d at 4ºC, in a growth chamber at 25ºC day/15ºC night with a

16 h day/8 h night photoperiod in a reach-in growth chamber (Model PGC-40L2, Percival

Scientific, IA, USA). One 21 d old plant per pot was kept and vernalized for 10 wk. at 4ºC with

an 8 h day/16 h night photoperiod, then returned to initial growth conditions until harvesting.

Harvested seeds were dark-stratified as in the bulking before performing experiments in the

Percival, then grown in cone-tainers (1.5-inch diameter, 164 ml) in a RL98 rack with nutrient

rich soil (Supplemental Methods S2). After the first true leaves expanded (ca. 23 d old plants),

pots were thinned to one plant per pot. We used a 12 h day/night photoperiod for all large-scale

experiments, randomized pots withing trays, and periodically rotated trays and moved them

around the growth chamber (or within shelves) to prevent positional effects.

Experiment 1: Low pCO_2 in 253 ecotypes We randomly selected 140 ecotypes for 5 biological

replicates (individual plants) and 113 ecotypes for 4 biological replicates, for a total of 1152

plants. The difference in number of replicates was due to space constraints in the growth

chamber. We imposed a treatment of low pCO_2 by scrubbing CO_2 from the Percival air using a

soda lime absorbent material (SODASORB HP, Divers Supply Inc., LA, USA). We set the

growth chamber to 200 ppm CO_2, which at our elevation of ~350 m corresponds to ~19.5 Pa, the

approximate pre-industrial partial pressure of CO_2 at 3000 m, compared to a pre-industrial partial

pressure of ~28 Pa at sea level. We sought to mimic high elevation conditions by growing plants

under high light (~462–616 μmol m^{-2} s^{-1}) and cool conditions (~3–13ºC; Supplemental Dataset

S3) with moderate soil drying. We terminated the experiment 156 d after planting and

subsequently completed harvesting and phenotyping within 5 d.

Phenotypes in Experiment 1 We recorded phenotypic variation in six traits on 250–253

ecotypes: $\delta^{13}C$, $\delta^{15}N$, root diameter, rosette diameter, and rosette compactness. Just prior to

harvest, we photographed overhead rosettes of each plant and used ImageJ (Schneider et al.,

2012) to measure rosette diameter (as the average between the longest and the shortest distance

between opposite leaf tips) and rosette compactness (by assigning a rating from 1–5 where 5

corresponded to more compact rosettes with shorter petioles and total leaf overlap, thus less soil

exposure and 1 corresponded to less compact rosettes with longer petioles and minimal leaf

overlap, thus more soil exposure). After harvesting, root diameter for each plant was measured 3

mm below the hypocotyl/root junction with a caliper. We then used 2–4 young healthy leaves per

plant for estimating $\delta^{13}C$ and $\delta^{15}N$. Leaves of each ecotype were pooled, dried at 27°C for 24 h,

and homogenized to a fine powder. We sent 1.5–2.5 mg of fine powder per ecotype to the

University of California Davis Stable Isotope Facility where $\delta^{13}C$ and $\delta^{15}N$ were quantified

following international standards using an elemental analyzer interfaced to a continuous flow

isotope ratio mass spectrometer (Sharp, 2005). We measured stomatal density on a subset of 80

widely distributed ecotypes (Supplemental Dataset S1): 2–4 young healthy leaves were collected

from 2–3 plants per ecotype and stored in 75% ethanol at 4°C. Epidermal peels from the abaxial

surface of each leaf were softened in distilled water for 3–7 min, immediately placed into 0.02%

Toluidine Blue stain for 3–5 min, rinsed with distilled water for 3 s, and put onto a slide with a

small drop of distilled water for imaging. Images were taken at 20X magnification using a Zeiss

compound microscope and stomatal density (stomata/mm^2) was quantified in ImageJ, using

distance-calibrated images of one representative area of each epidermal peel. To assess

coordinated life history variation among regions, we summarized variation in phenotypes from

Experiment 1 and days to flower (in plants grown at 10 and 16 ºC; published data in Alonso-

Blanco et al. 2016) with a principal component analysis (PCA; R 'prcomp' function).

Experiment 2: Low vs. High Light in 111–114 ecotypes We conducted Experiment 2 using

identical growth parameters as above except for light intensity (Supplemental Dataset S4). Five

biological replicates of 111–114 ecotypes, for a total of 1125 plants, were grown under two light

levels under cool conditions (~11ºC day/3ºC night). Plants grown on the lower level of the

growth chamber received the "high light" treatment with 100% maximum light output (~616

μmol m^{-2} s^{-1}), and plants grown on the upper level of the chamber received the "low light"

treatment with 57% maximum light output (~350 μmol m^{-2} s^{-1}). We distributed plants this way,

as the upper level of the chamber tends to get warmer, balancing out the heat from bright lights

in the bottom shelf. Soil temperature after 43 d taken with a FLIR ONE (Gen 2) thermal camera

for iOS (FLIR Systems, USA) showed that differences in soil temperature were on average only

within 1.3ºC, with warmer pots under the high light treatment (12.4°C vs. 11.1°C).

Experiment 3: Night-time above vs. night-time below freezing in 170–172 ecotypes

Experiment 3 consisted of two consecutive experiments using identical growth parameters as

Experiment 1, except for night-time temperatures (Supplemental Dataset S5). Six replicates of

170–172 ecotypes were grown under night-time below freezing temperature (850 plants) and

consecutively under night-time above freezing temperature (860 plants). Both experiments

maintained high light (~462–616 μmol m^{-2} s^{-1}) and cool conditions throughout (~15ºC day/4ºC

night), but in the night-time freezing experiment we introduced below freezing temperatures

(from –4 to –2ºC) after 40 d for 30 d. We monitored soil temperature with probes every 30 min,

which did not reveal soil frost during the day.

Phenotypes in Experiment 2 and 3 We recorded phenotypic variation in three traits related to

photosynthesis: fresh aboveground biomass, antioxidant activity, and leaf color. Ecotypes were

harvested 82–89 d (Experiment 2) and 72–75 d (Experiment 3) after planting. Biological

replicates from each ecotype were pooled together (due to their small size) and cleaned with

distilled water for assessing collective fresh aboveground mass and antioxidant activity with 1,1-

diphenyl-2-picrylhydrazyl (DPPH), an assay that estimates free radical scavenging activity of

antioxidants (protocol in Seyoum et al., 2006). Leaf color was estimated with ImageJ, excluding

late germinants. In Experiment 2, rosette leaf color was measured from the pixels selected at a

representative location within a mid-aged leaf on rosette photographs of each plant taken just

prior to harvest. We used the option "Color Histogram" to obtain the variation and calculate

percent of red, green, and blue pixels (RGB values). In Experiment 3, we obtained variation in

RGB values from images of the pooled leaves extracts per ecotype in test tubes used for DPPH.

Images were white-balanced and zoomed into the test tube avoiding any discolored, dark, or

bright areas. We note that upon visual examination of photographs, rosette colors are probably

more indicative of anthocyanins, while extract colors may also inform on other pigments like

carotenoids, including xanthophylls. We then used PCA to summarize variation in color assessed

from RGB values in contrasting conditions. Based on the color space in PC1 and PC2, we

calculated plasticity in color: the Euclidean distance between ecotype values of the two

contrasting conditions using the Pythagorean theorem. We also calculated plasticity in

antioxidant activity by finding the difference between experimental conditions (stressful minus

control condition). Survival rates per ecotype were obtained from the number of replicates that

survived at the end of each experiment divided by the number of replicates that germinated

within treatments. We used survival as a proxy of fitness and analyzed how other traits affect it

with Pearson correlations.

Assessing region-specific climatic and trait variation Based on ecotypes with accurate

coordinates and elevation of origin, we examined how study regions varied climatically and

differed in phenotypes. We first evaluated regional variation of environmental variables on

elevation using linear models (R 'lm' function) and performed a PCA on 85 environmental

variables (Supplemental Dataset S2). We then used ANOVA to test for differences in trait

variation among regions, followed by Tukey's honest significant difference post-hoc tests to

detect pairwise regional differences. We also used ANOVA to test for significant differences in

trait variation between conditions (plasticity) in Experiments 2 and 3.

Assessing region-specific putatively adaptive trait-elevation clines For each trait of interest,

including the first two PCs of Experiment 1–3 described above, we tested for associations with

elevation within regions using linear mixed effects kinship models with the function 'lmekin' in

the kinship2 R package (Therneau, 2012). These models accounted for population genetic

structure by incorporating a kinship matrix of ecotypes as a random effect. When significant,

they thus provide evidence of selection maintaining trait-elevation clines. A kinship matrix of

sequenced ecotypes was obtained from a published VCF of genotypes (Durvasula et al., 2017),

which we concatenated with the here sequenced eastern Africa ecotypes (Supplemental Methods

1), for a total of 1233 ecotypes (261 in our study) in that VCF. First, this VCF was converted to

gds format for manipulation in the SNPRelate R package (Zheng et al., 2012). We then

visualized population genetic structure with a neighbor-joining tree with a subset of 562,194

unlinked SNPs obtained with 'snpgdsLDpruning' in SNPRelate with a linkage-disequilibrium

threshold of 0.5 and a 10 kb window. With this set of SNPs, we calculated a distance matrix

based on genetic dissimilarity between ecotypes and used the 'nj' and 'plotnj' function in the R

package ape (Paradis et al., 2004). We obtained a kinship matrix with the snpgdsIBS function

(*i.e.*, based on the fraction of identity by state for each pair of samples) and the same set of

unlinked SNPs.

Assessing genome-wide phenotypic associations and genomic clines We imputed the above

VCF based on LD with Beagle v.4.1 (Browning et al., 2018), resulting in 22,619,232 SNPs of

which 11,500,298 were polymorphic. We used univariate linear mixed effects models in gemma

v.0.98.3 (Zhou and Stephens, 2012) to perform genome wide association studies (GWAS).

Because traits generally differed in the number of analyzed individuals, we filtered out loci with

maf<0.05 directly in gemma. To account for multiple testing, we used a false-discovery-rate

(FDR) threshold of 0.05 to detect significantly associated SNPs with traits of interest. We ran

gemma on the global dataset for all traits, as well as by region for those traits that showed a

significant association with elevation. For flowering time regional GWAS, we used all the

ecotypes with available data: 60 for Asia, 289 for central Europe/Caucasus, 398 for northwestern

Europe, and 183 for western Mediterranean. We identified QTL by assigning the top SNPs (i.e.,

those with the lowest 100 p-values for each trait) to candidate genes using the Araport11

reannotation (Cheng et al., 2017) and a 5 kb window. To examine if selection on elevation

maintained allelic variation in top SNPs in putative QTL, we evaluated the association of allelic

variation with elevation while accounting for population genetic structure with 'lmekin'.

Assessing NPQ kinetics in a global panel of 9–11 ecotypes We performed a detailed study on

NPQ kinetics based on ecotypes with a global distribution representing the lowest (22–254 m)

and highest (1576–4078 m) elevational range of Arabidopsis, that also showed great variation in

antioxidant activity. With this panel we aimed to capture diverse strategies, but given the limited

number of ecotypes, not necessarily make conclusive cline tests.

Seeds were dark-stratified at 4ºC for 4 d in water (or in 800 PPM gibberellic acid

for the JL011-2-1 ecotype), sown in 8.89×8.89 cm pots with soil-less BM2 potting mix under

moist conditions, and grown in a reach-in Percival chamber (model AR-66L2). Growing

conditions were set at 21ºC day/18ºC night with an 8 h day/16 h night photoperiod (200 μmol m^{-}

^{2} s^{-1}) and 60% RH, 5 d old plants thinned to one per pot, well-watered, and trays repositioned at

random locations in the chamber.

Two weeks after germination, NPQ kinetics was measured on seedlings in two

separate experiments (5–10 biological replicates per ecotype per treatment) using a fluorescence

imager (Closed FluorCam 7 FC 800-C, Photon System Instruments, Czech Republic). Half of the

seedlings were subjected to drought (5 d of no watering; 2 ecotypes excluded due to flowering)

or to 17 h of overnight chilling at 6°C, and the other half were maintained under normal

conditions and used as control. Plants were subjected to three cycles of 3 min 1000 μmol m^{-2} s^{-1}

followed by 2 min of dark. NPQ of chlorophyll fluorescence was determined assuming the Stern-

Volmer quenching model (NPQ $= F_m/F_m{}' - 1$; Bilger and Bjorkman, 1994). NPQ was then

calculated at following intervals of 10s, 10s, 20s, 20s, 60s and 60s in light and 10s, 20s, 30s and

60s in dark. To this end, raw images were processed manually using a FluorCam 7 by drawing

three circles per plant of 170–180 pixels on three young fully developed leaves. The NPQ data

were fitted to exponential equations to parametrize the NPQ kinetics of its induction in light and

relaxation in dark of three subsequent light-dark cycles (Sahay et al., 2023, Supplemental Figure

S3) using a custom-made script in MATLAB v. R2018b. After 30 d since germination, we

harvested rosettes from control conditions to estimate fresh and dry aboveground biomass, and

photosynthesis related pigments. For each ecotype, we harvested 4–6 biological replicates, which

were immediately weighted to obtain fresh weight. To estimate dry weight, fresh rosettes were

dried to constant weight at 70°C. On the remaining 4–6 biological replicates, we quantified

photosynthesis related pigments from 2.32 cm^2 leaf tissue from the interveinal region following a

100% methanol-based extraction (Lichtenthaler, 1987). To understand variation among ecotypes

in NPQ kinetics, biomass, and pigments, we performed ANOVA and Tukey HSD post-hoc tests

(α=0.05) in SAS v.9.4.

Assessing the physiological effect of allelic variation and gene expression of *PHT4;4* in

western Mediterranean ecotypes We investigated the effect of allelic variation in *PHT4;4*

(chr4: 160735) on western Mediterranean ecotypes, using seven ecotypes with the low-

antioxidant allele and six closely related ecotypes with the high-antioxidant allele (Supplemental

Dataset S1), along with three *PHT4;4* knockouts. We obtained a Ds transposon tagged mutant in

a Ler genetic background (CS26443) and two T-DNA insertional mutants in a Col-0 genetic

background (Salk_082875 and CS444342) from the Cold Spring Harbor Laboratory and the

Arabidopsis Biological Resource Center (Ohio State University), respectively. Lines were selfed

until a homozygous progeny for knockouts was confirmed by PCR (Supplemental Table S3).

Two weeks after germination under control conditions, half of the seedlings were subjected to

two days of high light (HL; 550 μmol m^{-2} s^{-1} at 24°C/22°C day /night with an 8 h day/16 h night

photoperiod) or to four days of HL under cold (550 μmol m^{-2} s^{-1} at 6°C with a 16 h day/8 h night

photoperiod) in a walk-in growth chamber (Conviron model GR48, Controlled Environments,

Manitoba, Canada). The other half were maintained under normal conditions and used as

controls. NPQ kinetics was investigated during 10 min of light and 10 min of dark, and

parametrized as in the global NPQ experiment, except that saturating flashes were spaced at the

following intervals (in s): 15, 30, 30, 60, 60, 60, 60, 60, 60, 60, 60, 60, 60, 9, 15, 30, 60,180, and

300.

To quantify *PHT4;4* expression under high light, four leaf discs (2.32 cm^2) from a

fully expanded leaf of 5–6 biological replicates were collected 2 h after the photoperiod started

in ecotypes treated for 2 days with HL at RT (first HL experiment above). Total mRNA was

extracted from leaf tissue with a NucleoSpin RNA/Protein kit (REF740933, Macherey-Nagel

GmbH & Co.), then treated with DNase before transcription into cDNA using a Superscript III

First-Strand Synthesis System (18080-051, Thermo Fisher Scientific). The cDNA was amplified

by quantitative reverse transcription PCR (RT-qPCR) to quantify the *PHT4;4* transcript (5'-

GCATATAGCTCTCCCAAGGATG-3' and 5'- TAGAGACCCGACTGAGAGAATG-3')

relative to the elongation factor (AtEF1α; 5'-TGAGCACGCTCTTCTTGCTTTCA-3' and 5'-

GGTGGTGGCATCCATCTTGTTACA-3'). The RT-qPCR was performed using a

SsoAdvanced Universal SYBR Green Supermix (172-5270, BioRad). We used ANOVA

followed by a Dunnett's test to investigate differences in physiological traits and *PHT4;4* gene

expression between ecotypes with different GWAS alleles and between knockouts and

corresponding wild-type.

We performed an additional experiment to quantify antioxidant activity in

PHT4;4 knockouts and their respective wild-type, along with five western Mediterranean

ecotypes that differ in the GWAS allele (Supplemental Dataset S1). Seeds were dark stratified at

4ºC in gibberellic acid (1000 PPM) for 7 d, then sown in 4×4 cm pots with BMX potting mix

and kept moist. We grew plants for 20 d in a walk-in chamber (Conviron model CMP4030,

Controlled Environments, ND, USA) with a 16 h day/8 h night photoperiod at ~22ºC day/18ºC

night, and subsequently transferred them to a cold room at 4–8ºC day and night, where 10–16

randomized biological replicates of each genotype were placed under high (~600 μmol m^{-2} s^{-1})

or low light (~150 μmol m^{-2} s^{-1}). We started with an 8 h day/ 16 h night photoperiod, and

gradually increased it to 12 h over the course of 1 wk. After 10 d, 2 h after the photoperiod

started, we measured chlorophyll content with a SPAD meter (Spectrum Technologies, IL, USA)

based on an average of three points along a fully expanded young leaf, then harvested rosettes to

perform the same DPPH assay as in our GWAS high light experiment. We took photographs

from leaf extracts and quantified RGB colors and brightness with ImageJ in white-balanced

photos. We then performed a PCA on RGB values across genotypes and used PC1 and PC2 as

color traits.

Assessing potential *cis*-regulatory variation of *PHT4;4* in Moroccan ecotypes We used

samtools v1.10 (Li et al. 2009) to extract the 3.5kbp upstream and downstream regions of

PHT4;4. Specifically, we extracted the 3kbp upstream and 0.5kbp downstream of the gene's start

codon for the upstream regulatory coding region, and the 0.5kbp upstream and 3kbp downstream

of the gene's end codon for the downstream regulatory coding region. We then used the standard

functions of the program STREME (Bailey, 2021) to calculate significantly enriched motifs

within 63 western Mediterranean ecotypes carrying the low-antioxidant GWAS haplotype

(Supplemental Figure S9) vs. 53 closely related ones with the other allele (based on genome-

wide genetic distance, see the tree in Supplemental Figure S1). We then searched for overlaps

between the discovered enriched motifs and the JASPAR 2022 CORE (Castro-Mondragón et al.,

2022) plant transcription factor binding sites database with MEME suite's online tool TOMTOM

(Gupta et al., 2007).

1010

1011 **Table 1** Candidate genes mapped with GWAS which associated SNP follows a putatively
1012 adaptive elevational cline. CHR: chromosome, Coord: genomic coordinates in bp, Rosette D:
1013 rosette diameter, Rosette Comp: rosette compactness, W Med: western Mediterranean.
1014

Locus	Gene	Function	CHR	Coord (bp)	Strand	SNP (bp)	Phenotype	Region
AT1G77080	*FLOWERING LOCUS M*	Flowering response to cold	1	28955522–28960111	Forward	28957836	δ^{13}C	Asia
AT1G65490	*SECRETED PEPTIDE STMP5*	Plant growth and pathogen defense	1	24354646–24355490	Forward	24356159	Rosette D	Asia
AT5G41890	*GDSL*-motif	Biotic and abiotic stress responses	5	16764211–16766702	Reverse	16766371	Rosette Comp	Asia
AT5G45830	*DELAY OF GERMINATION1*	Seed dormancy response to cold	5	18589438–18591506	Reverse	18590591	FT10 & FT16	W Med
AT5G23220	*NICOTINAMID-ASE 3*	Abscisic acid signaling	5	7819608–7820760	Reverse	7818730	δ^{13}C	W Med
AT5G11850	*MAP3* kinase	Abscisic acid signaling	5	3816080–3821170	Reverse	3821491	δ^{13}C	W Med
AT4G00370	Phosphate transporter (*PHT4;4*)	Ascorbate transporter	4	162897–166357	Reverse	160735	HL antioxidants	Global
AT5G49730	*FERRIC REDUCTION OXIDASE 6*	Iron transporter	5	20201037–20204595	Reverse	20206030	HL antioxidants	Global
AT5G49740	*FERRIC REDUCTION OXIDASE 7*	Iron transporter	5	20205302–20208777	Reverse	20206030	HL antioxidants	Global

1015

1016 **Table 2** Most frequent transcription factor binding sites (TFBS) found within 54 discovered
1017 motifs 3 kbp around *PHT4;4* in 63 western Mediterranean lines carrying the low-antioxidant
1018 *PHT4;4* haplotype. For each known TFBS, we report the number of overlapping motifs (N).

TFBS id	TFBS name	TFBS family	N motifs
MA1277.1	DOF1.7	DOF	13
MA1267.1	DOF5.8	DOF	12
MA1274.1	DOF6.3	DOF	10
MA1184.1	RVE1	Myb-related	9
MA1191.1	RVE7L	Myb-related	9
MA1268.1	CDF5	DOF	9
MA1273.1	DOF4.2	DOF	9
MA1279.1	DOF1.5	DOF	9
MA0563.1	SEP3	MIKC	8
MA1012.1	AGL27	MIKC	7
MA1204.1	AGL13	MIKC	7
MA1205.1	AGL6	MIKC	7
MA1281.1	DOF5.1	DOF	7

1019

Accession numbers

1021 Sequence data (bam files), a VCF with newly sequenced eastern Africa ecotypes, and

1022 phenotypes from this article are available in the Figshare public repository with DOI:

1023 10.6084/m9.figshare.23960727

1024

Acknowledgments

Permits: Plant material was exported from Uganda with permission of The Ministry of

Agriculture, Animal Industry and Fisheries' Plant Quarantine and Inspection Services, permit

UQIS 4414/93/PC (E). Material was exported from Ethiopia with permission of the Ethiopian

Biodiversity Institute, Ref. no. EBE71/7065/2018. Material was imported to the USA under

USDA APHIS permits P37-17-01651 and P37-18-00230. Chloee McLaughlin, Jeremy

Sutherland, Tim Gilpatrick, Connor Campana, Victoria Meagher, and Jaden Hill assisted in

experiment setup and rosette harvests. Shawn Berghard advised in pest and disease control.

Alexander Batelaan, Darla Brennan, Rachel Gerdes, Bailey McLean, and Annie Nelson helped in

performing NPQ kinetics assays. Funding was provided by NSF DEB-1927009 and NIH

1R35GM138300-01 awards to JRL, and by NSF IOS-2142993 to KG.

1036

1037

1038 **Author contributions**

1039 DG, CL, SS, KG, and JRL designed the research; DG, CL, AH, SS, LL, TX, MT, EK, DE, JK,

1040 MY, CEB, and TW performed research; DG, CL, SS, LL, TX, MT, EK analyzed data; DG, CL,

1041 SS, KG, and JRL wrote the paper with contributions of all other authors.

1042

References

Aarabi, et al. (2023). Genome-wide association study unveils ascorbate regulation by *PAS/LOV*

PROTEIN during high light acclimation. Plant Physiology **kiad323.**

Alonso-Blanco, C. et al. (2016). 1,135 Genomes Reveal the Global Pattern of Polymorphism in

Arabidopsis thaliana. Cell **166**: 481–491.

Bailey, T.L. (2021). STREME: accurate and versatile sequence motif discovery. Bioinformatics

37: 2834–2840.

Bilger, W. and Bjorkman, O. (1994). Relationships among violaxanthin deepoxidation,

thylakoid membrane conformation, and nonphotochemical chlorophyll fluorescence

quenching in leaves of cotton (Gossypium hirsutum L.). Planta **193**.

Bohutínská, M., Vlček, J., Yair, S., Laenen, B., Konečná, V., Fracassetti, M., Slotte, T., and

Kolář, F. (2021). Genomic basis of parallel adaptation varies with divergence in *Arabidopsis*

and its relatives. Proc Natl Acad Sci USA **118**: e2022713118.

Brennan, A.C., Méndez-Vigo, B., Haddioui, A., Martínez-Zapater, J.M., Picó, F.X., and

Alonso-Blanco, C. (2014). The genetic structure of *Arabidopsis thaliana* in the south-

western Mediterranean range reveals a shared history between North Africa and southern

Europe. BMC Plant Biol **14**: 17.

Brochmann, C., Gizaw, A., Chala, D., Kandziora, M., Eilu, G., Popp, M., Pirie, M.D., and

Gehrke, B. (2021). History and evolution of the afroalpine flora: in the footsteps of Olov

Hedberg. Alpine Botany.

Browning, B.L., Zhou, Y., Browning, S.R. (2018). A one-penny imputed genome from next-

generation reference panels. Am J Hum Genet **103**: 338–348.

Caicedo, A., Stinchcombe, J.R., Olsen, K.M., Schmitt, J., and Purugganan, M. (2004).

Epistatic interaction between Arabidopsis *FRI* and *FLC* flowering time genes generates a

latitudinal cline in a life history trait. Proc Nat Acad Sci **101**: 15670–15675.

Caldas, F.J. (1966). Obras completas de Francisco José de Caldas: publicadas por la

Universidad Nacional de Colombia como homenaje con motivo del sesquicentenario de su

muerte 1816 - Octubre 29 - 1966 (Imprenta Nacional: Bogotá, Colombia).

Castelán-Muñoz, Herrera, J., Cajero-Sánchez, W., Arrizubieta, M., Trejo, C., García-

Ponce, B., Sánchez, M.P., Álvarez-Buylla, E.R., and Garay-Arroyo, A. (2019). MADS-

box genes are key components of genetic regulatory networks involved in abiotic stress and

plastic developmental responses in plants. Front Plant Sci **10**: 853.

Castro-Mondragón J.A., et al. (2022) JASPAR 2022: the 9th release of the open-access

database of transcription factor binding profiles. Nucleic Acids Res. **50**: D165-D173.

Cheng, C.-Y., Krishnakumar, V., Chan, A.P., Thibaud☐Nissen, F., Schobel, S., and Town,

C.D. (2017). Araport11: a complete reannotation of the Arabidopsis thaliana reference

genome. The Plant Journal **89**: 789–804.

Clausen, J., Keck, D.D., and Hiesey, W.M. (1940). Experimental studies on the nature of

species. I. Effect of varied environments on western North American plants. Experimental

studies on the nature of species. I. Effect of varied environments on western North American

plants.

DeLeo, V.L., Menge, D.N.L., Hanks, E.M., Juenger, T.E., and Lasky, J.R. (2020). Effects of

two centuries of global environmental variation on phenology and physiology of *Arabidopsis*

thaliana. Glob Change Biol **26**: 523–538.

Des Marais, D.L., Auchincloss, L.C., Sukamtoh, E., McKay, J.K., Logan, T., Richards,

J.H., and Juenger, T.E. (2014). Variation in MPK12 affects water use efficiency in

Arabidopsis and reveals a pleiotropic link between guard cell size and ABA response. Proc

Natl Acad Sci USA **111**: 2836–2841.

Distelbarth, H., Nägele, T., and Heyer, A.G. (2013). Responses of antioxidant enzymes to cold

and high light are not correlated to freezing tolerance in natural accessions of *Arabidopsis*

thaliana. Plant Biol J **15**: 982–990.

Dittberner, H., Korte, A., Mettler-Altmann, T., Weber, A.P.M., Monroe, G., and de Meaux,

J. (2018). Natural variation in stomata size contributes to the local adaptation of water-use

efficiency in *Arabidopsis thaliana*. Mol Ecol **27**: 4052–4065.

Durvasula, A., Fulgione, A., Gutaker, R.M., Alacakaptan, S.I., Flood, P.J., Neto, C.,

Tsuchimatsu, T., Burbano, H.A., Picó, F.X., Alonso-Blanco, C., and Hancock, A.M.

(2017). African genomes illuminate the early history and transition to selfing in *Arabidopsis thaliana*. Proc Natl Acad Sci USA **114**: 5213–5218.

Feng, H., An, F., Zhang, S., Ji, Z., Ling, H.-L., and Zuo, J. (2006). Light-regulated, tissue-specific, and cell differentiation-specific expression of the *Arabidopsis* Fe(III)-chelate reductase gene AtFRO6. **140**: 1345–1354.

Fernández☐Marín, B., Gulías, J., Figueroa, C.M., Iñiguez, C., Clemente☐Moreno, M.J., Nunes☐Nesi, A., Fernie, A.R., Cavieres, L.A., Bravo, L.A., García☐Plazaola, J.I., and Gago, J. (2020). How do vascular plants perform photosynthesis in extreme environments? An integrative ecophysiological and biochemical story. Plant J **101**: 979–1000.

Fulgione, A. and Hancock, A.M. (2018). Archaic lineages broaden our view on the history of *Arabidopsis thaliana*. New Phytol **219**: 1194–1198.

García-Plazaola, J. I., Esteban, R., Fernández-Marín, B., Kranner, I., and Porcar-Castell, A. (2012). Thermal energy dissipation and xanthophyll cycles beyond the Arabidopsis model. Photosynthesis Research **113**: 89–103.

Gorsuch, P. A., Pandey, S., and Atkin, O. K. (2010). Thermal de-acclimation: how permanent are leaf phenotypes when cold-acclimated plants experience warming? Plant, Cell & Environment **33**: 1124–1137.

Graeber, K. et al. (2014). *DELAY OF GERMINATION 1* mediates a conserved coat-dormancy mechanism for the temperature- and gibberellin-dependent control of seed germination. Proc Natl Acad Sci USA **111**: E3571–E3580.

Günther, T., Lampei, C., Barilar, I., and Schmid, K.J. (2016). Genomic and phenotypic differentiation of *Arabidopsis thaliana* along altitudinal gradients in the North Italian Alps. Mol Ecol **25**: 3574–3592.

Gupta, S., Stamatoyannopoulos, J. A., Bailey, T. L., Noble, W. S. (2007). Quantifying similarity between motifs. Genome Biol. **8**: R24.

Hannah, M.A., Wiese, D., Freund, S., Fiehn, O., Heyer, A.G., and Hincha, D.K. (2006). Natural genetic variation of freezing tolerance in *Arabidopsis*. Plant Physiology **142**: 98–112.

Havaux, M. and Kloppstech, K. (2001). The protective functions of carotenoid and flavonoid pigments against excess visible radiation at chilling temperature investigated in *Arabidopsis* npq and tt mutants. Planta **213**: 953–966.

Hedberg, I. and Hedberg, O. (1979). Tropical-alpine life-forms of vascular plants. OIKOS **33**:
297–307.

Hepworth, J., et al. (2018). Absence of warmth permits epigenetic memory of winter in
Arabidopsis. Nature Communications **9**: 639.

Hepworth, J., et al. (2020). Natural variation in autumn expression is the major adaptive
determinant distinguishing Arabidopsis *FLC* haplotypes. eLife **9**: e57671.

Huo, H., Wei, S., and Bradford, K.J. (2016). *DELAY OF GERMINATION1* (*DOG1*) regulates
both seed dormancy and flowering time through microRNA pathways. Proc Natl Acad Sci
USA **113**: E2199–E2206.

Jain, A., Wilson, G. T., Connolly, E. L. (2014). The diverse roles of *FRO* family
metalloreductases in iron and copper homeostasis. Front in Plant Sci **5**: 100.

Jeong, J., Cohu, C., Kerkeb, L., Pilon, M., Connolly, E.L., and Guerinot, M.L. (2008).
Chloroplast Fe (III) chelate reductase activity is essential for seedling viability under iron
limiting conditions. Proc Nat Acad Sci **105**: 10619–10624.

Jeong, J. and Connolly, E.L. (2009). Iron uptake mechanisms in plants: Functions of the *FRO*
family of ferric reductases. Plant Science **176**: 709–714.

Jin, S., Kim, S.Y., Susila, H., Nasim, Z., Youn, G., and Ahn, J. H. (2022). *FLOWERING*
LOCUS M isoforms differentially affect the subcellular localization and stability of *SHORT*
VEGETATIVE PHASE to regulate temperature-responsive flowering in *Arabidopsis*.
Molecular Plant **15**: 1696–1709.

Josephs, E., Berg, J.J., Ross-Ibarra, J., and Coop, G. (2019). Detecting adaptive
differentiation in structured populations with genomic data and common gardens. Genetics
211: 989–1004.

1152

1153 **Kenney, A.M., McKay, J.K., Richards, J.H., and Juenger, T.E.** (2014). Direct and indirect
1154   selection on flowering time, water□use efficiency (WUE, δ^{13}C), and WUE plasticity to
1155   drought in *Arabidopsis thaliana*. Ecol Evol **4**: 4505–4521.

1156 **Kim, D.-H., and Sung, S.** (2013). Coordination of the vernalization response through a VIN3
1157   and *FLC* gene family regulatory network in *Arabidopsis*. Plant Cell **25**: 454–69.

Kim, J.-S., Lim, J.Y., Shin, H., Kim, B.-G., Yoo, S.-D., Kim, W.T., and Huh, J.H. (2019). ROS1-Dependent DNA Demethylation Is Required for ABA-Inducible *NIC3* Expression. Plant Physiol. **179**: 1810–1821.

Klepikova, A., Kasianov, A.S., Gerasimov, E.S., Logacheva, M.D., and Penin, A.A. (2016). A high-resolution map of the *Arabidopsis thaliana* developmental transcriptome based on RNA-seq profiling. The Plant J **88**: 1058–1070.

Kooyers, N.J., Greenlee, A.B., Colicchio, J.M., Oh, M., and Blackman, B.K. (2015). Replicate altitudinal clines reveal that evolutionary flexibility underlies adaptation to drought stress in annual *Mimulus guttatus*. New Phytol **206**: 152–165.

Körner, C. (2021). Alpine plant life: functional plant ecology of high mountain ecosystems (Springer International Publishing: Cham).

Körner, C. (2003). Ecological impacts of atmospheric CO_2 enrichment on terrestrial ecosystems. Philosophical Transactions of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences **361**: 2023–2041.

Körner, C. (2007). The use of 'altitude'in ecological research. Trends in Ecology & Evolution **22**: 569–574.

Kromdijk, J., Głowacka, K., Leonelli, L., Gabilly, S.T., Iwai, M., Niyogi, K.K., and Long, S.P. (2016). Improving photosynthesis and crop productivity by accelerating recovery from photoprotection. Science **354**: 857–861.

Kumar, M., Patel, M.K., Kumar, N., Bajpai, A.B., and Siddique, K.H.M. (2021). Metabolomics and molecular approaches reveal drought stress tolerance in plants. Int J Mol Sci **22**: 9108.

Kuromori, T., Fujita, M., Takahashi, F., Yamaguchi-Shinozaki, K., and Shinozaki, K. (2022). Inter-tissue and inter-organ signaling in drought stress response and phenotyping of drought tolerance. The Plant Journal **109**: 342–358.

Lai, C.P., Huang, L.-M., Chen, L.-F. O., Chan, M.-T., Shaw, J.-F. (2017). Genome-wide analysis of GDSL-type esterases/lipases in *Arabidopsis*. Plant Molecular Biology **95**: 181–197.

Lampei, C., Wunder, J., Wilhalm, T., and Schmid, K.J. (2019). Microclimate predicts frost hardiness of alpine *Arabidopsis thaliana* populations better than elevation. Ecol Evol **9**: 13017–13029.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis,

G., Durbin, R., 1000 Genome Project Data Processing Subgroup. (2009). The Sequence

Alignment/Map format and SAMtools. Bioinformatics **25**: 2078–2079.

Lichtenthaler, H.K. (1987). 34. Chlorophylls and carotenoids: Pigments of photosynthetic

biomembranes. In Methods in Enzymology (Academic Press), pp. 350–382.

López-Arboleda, W.A., Reinert, S., and Nordborg, M. (2021). Global genetic heterogeneity in

adaptive traits. Molecular Biology and Evolution **38**: 4822–4831.

Lovell, J.T., Juenger, T.E., Michaels, S.D., Lasky, J.R., Platt, A., Richards, J.H., Yu, X.,

Easlon, H.M., Sen, S., and McKay, J.K. (2013). Pleiotropy of *FRIGIDA* enhances the

potential for multivariate adaptation. Proc. R. Soc. B. **280**: 20131043.

Ludlow, M.M. (1989). Strategies of response to water stress. In Structural and Functional

Responses to Environmental Stresses, K.H. Kreeb, H. Richter, and T.M. Hinckley, eds (SPB

Academic), pp. 269–281.

Luo, Y., Widmer, A. and Karrenberg, S. (2015a). The roles of genetic drift and natural

selection in quantitative trait divergence along an altitudinal gradient in Arabidopsis thaliana.

Heredity **114**: 220–228.

Luo, Y., Dong, X., Yu, T., Shi, X., Li, Z., Yang, W., Widmer, A., and Karrenberg, S.

(2015b). A single nucleotide deletion in *Gibberellin20-oxidase1* causes alpine dwarfism in

Arabidopsis. Plant Physiol. **168**: 930–937.

Malnoë, A. (2018). Photoinhibition or photoprotection of photosynthesis? Update on the (newly

termed) sustained quenching component qH. Environmental and Experimental Botany **154**:

123–133.

Martínez-Berdeja, A., Stitzer, M.C., Taylor, M.A., Okada, M., Ezcurra, E., Runcie, D.E.,

and Schmitt, J. (2020). Functional variants of *DOG1* control seed chilling responses and

variation in seasonal life-history strategies in *Arabidopsis thaliana*. Proc Natl Acad Sci USA

117: 2526–2534.

Mckay, J.K., Richards, J.H., and Mitchell Olds, T. (2003). Genetics of drought adaptation in

Arabidopsis thaliana: I. Pleiotropy contributes to genetic correlations among ecological

traits. Molecular Ecology **12**: 1137–1151.

Méndez-Vigo, B., Picó, F.X., Ramiro, M., Martínez-Zapater, J.M., and Alonso-Blanco, C. (2011). Altitudinal and climatic adaptation is mediated by flowering traits and *FRI*, *FLC*, and *PHYC* genes in *Arabidopsis*. Plant Physiology **157**: 1942–1955.

Miyaji, T., Kuromori, T., Takeuchi, Y., Yamaji, N., Yokosho, K., Shimazawa, A., Sugimoto, E., Omote, H., Ma, J.F., Shinozaki, K., and Moriyama, Y. (2015). AtPHT4;4 is a chloroplast-localized ascorbate transporter in Arabidopsis. Nat Commun **6**: 5928.

Montesinos-Navarro, A., Tonsor, S.J., Alonso-Blanco, C., and Picó, F.X. (2009). Demographic and genetic patterns of variation among populations of *Arabidopsis thaliana* from contrasting native environments. PLoS ONE **4**: e7213.

Montesinos-Navarro, A., Picó, F.X., and Tonsor, S.J. (2012). Clinal variation in seed traits influencing life cycle timing in *Arabidopsis thaliana*. Evolution **66**: 3417–3431.

Montesinos☐Navarro, A., Wig, J., Xavier Pico, F., and Tonsor, S.J. (2011). *Arabidopsis thaliana* populations show clinal variation in a climatic gradient associated with altitude. New Phytologist **189**: 282–294.

Moustakas, M., Sperdouli, I., and Moustaka, J. (2022). Early drought stress warning in plants: color pictures of photosystem II photochemistry. Climate **10**: 179.

Muller, P., Li, X.-P., and Niyogi, K.K. (2001). Non-photochemical quenching. A response to excess light energy. Plant physiology **125**: 1558–1566.

Nam, H.-I., Shahzad, Z., Dorone, Y., Clowez, S., Zhao, K., Bouain, N., Lay-Pruitt, K.S., Cho, H., Rhee, S.Y., and Rouached, H. (2021). Interdependent iron and phosphorus availability controls photosynthesis through retrograde signaling. Nat Commun **12**: 7211.

Nakabayashi, R. et al. (2014). Enhancement of oxidative and drought tolerance in *Arabidopsis* by overaccumulation of antioxidant flavonoids. Plant J **77**: 367–379.

Nilkens, M., Kress, E., Lambrev, P., Miloslavina, Y., Müller, M., Holzwarth, A.R., and Jahns, P. (2010). Identification of a slowly inducible zeaxanthin-dependent component of non- photochemical quenching of chlorophyll fluorescence generated under steady-state conditions in Arabidopsis. Biochimica et Biophysica Acta (BBA) Bioenergetics **1797**: 466–475.

Paradis, E., Claude, J., and Strimmer, K. (2004). APE: analyses of phylogenetics and evolution in R language. Bioinformatics **20**: 289–290.

Picó, F.X. (2012). Demographic fate of *Arabidopsis thaliana* cohorts of autumn- and spring-
germinated plants along an altitudinal gradient: Cohort dynamics of *Arabidopsis thaliana*.
Journal of Ecology **100**: 1009–1018.

Rungrat, T., Almonte, A.A., Cheng, R., Gollan, P.J., Stuart, T., Aro, E., Borevitz, J.O.,
Pogson, B., and Wilson, P.B. (2019). A genome☐wide association study of
non☐photochemical quenching in response to local seasonal climates in *Arabidopsis*
thaliana. Plant Direct **3**.

Sahay, S., Grzybowski, J., Schnable, J.C., and Glowacka, K. (2023). Genetic control of
photoprotection and photosystem II operating efficiency in plants. New Phytol **239**: 1068–
1082.

Seyoum, A., Asres, K., and El-Fiky, F.K. (2006). Structure-radical scavenging activity
relationships of flavonoids. Phytochemistry **67**: 2058–2070.

Sharif, R., Raza, A., Chen, P., Li, Y., El-Ballat, E. M., Rauf, A., Hano, C., and El-Esawi M,
A. (2021). HD-ZIP gene family: potential roles in improving plant growth and regulating
stress-responsive mechanisms in plants. Genes (Basel) **12**: 1256.

Sharp, Z. (2005). Principles of stable isotope geochemistry. (Prentice Hall, USA).

Singh, A. and Roy, S. (2017). High altitude population of *Arabidopsis thaliana* is more plastic
and adaptive under common garden than controlled condition. BMC Ecol **17**: 39.

Smith, W.K. and Knapp, A.K. (1990). Ecophysiology of high elevation forests. In Plant
Biology of the Basin and Range, Ecological Studies. (Springer Berlin Heidelberg: Berlin,
Heidelberg), pp. 87–142.

Soitamo, A.J., Plippo, M., Allahverdiyeva, Y., Battchikova, N., and Aro, E.-M. (2008). Light
has a specific role in modulating *Arabidopsis* gene expression at low temperature. BMC
Plant Biol **8**: 13.

Stinchcombe, J.R., Caicedo, A.L., Hopkins, R., Mays, C., Boyd, E.W., Purugganan, M.D.,
and Schmitt, J. (2005). Vernalization sensitivity in *Arabidopsis thaliana* (Brassicaceae): the
effects of latitude and *FLC* variation. American Journal of Botany **92**: 1701–1707.

Suter, L., Rüegg, M., Zemp, N., Hennig, L., and Widmer, A. (2014). Gene regulatory
variation mediates flowering responses to vernalization along an altitudinal gradient in
Arabidopsis. Plant Physiol. **166**: 1928–1942.

Takahashi, Y., Zhang, J., Hsu, P.-K., Ceciliato, P.H.O., Zhang, L., Dubeaux, G.,

Munemasa, S., Ge, C., Zhao, Y., Hauser, F., and Schroeder, J.I. (2020). MAP3Kinase-

dependent SnRK2-kinase activation is required for abscisic acid signal transduction and rapid

osmotic stress response. Nat Commun **11**: 12.

Therneau, T. (2012). The lmekin function. Rochester, MN: Mayo Clinic.

Tripathi, A.M., Singh, A., Singh, R., Verma, A.K., and Roy, S. (2019). Modulation of

miRNA expression in natural populations of A. thaliana along a wide altitudinal gradient of

Indian Himalayas. Sci Rep **9**: 441.

Tusiime, F.M., Gizaw, A., Gussarova, G., Nemomissa, S., Popp, M., Masao, C.A.,

Wondimu, T., Abdi, A.A., Mirré, V., Muwanika, V., Eilu, G., and Brochmann, C.

(2020). Afro-alpine flagships revisited: Parallel adaptation, intermountain admixture, and

shallow genetic structuring in the giant senecios (*Dendrosenecio*). PLoS ONE **15**: e0228979.

Tyagi, A., Yadav, A., Tripathi, A.M., and Roy, S. (2016). High light intensity plays a major

role in emergence of population level variation in Arabidopsis thaliana along an altitudinal

gradient. Sci Rep **6**: 26160.

Vasseur, F., Exposito-Alonso, M., Ayala-Garay, O.J., Wang, G., Enquist, B.J., Vile, D.,

Violle, C., and Weigel, D. (2018). Adaptive diversification of growth allometry in the plant

Arabidopsis thaliana. Proc Natl Acad Sci USA **115**: 3416–3421.

Vidigal, D.S., Marques, A.C.S.S., Willems, L.A.J., Buijs, G., Méndez-Vigo, B., Hilhorst,

H.W.M., Bentsink, L., Picó, F.X., and Alonso-Blanco, C. (2016). Altitudinal and climatic

associations of seed dormancy and flowering traits evidence adaptation of annual life cycle

timing in *Arabidopsis thaliana*: Climatic adaptation of annual life cycles. Plant, Cell &

Environment **39**: 1737–1748.

Wang, H., Prentice, I.C., Davis, T.W., Keenan, T.F., Wright, I.J., and Peng, C. (2017).

Photosynthetic responses to altitude: an explanation based on optimality principles. New

Phytol **213**: 976–982.

Ward, J.K. and Strain, B.R. (1997). Effects of low and elevated CO2 partial pressure on

growth and reproduction of Arabidopsis thaliana from different elevations. Plant Cell

Environ **20**: 254–260.

Wise, R.R. (1995). Chilling-enhanced photooxidation: The production, action and study of

reactive oxygen species produced during chilling in the light. Photosynth Res **45**: 79–97.

Wolfe, M.D. and Tonsor, S.J. (2014). Adaptation to spring heat and drought in northeastern
Spanish *Arabidopsis thaliana.* New Phytol **201**: 323–334.

Wos, et al. (2022). Parallel local adaptation to an alpine environment in *Arabidopsis arenosa.*
Journal of Ecology **110**: 2448– 2461.

Yu, Y., Wang, S., Wang, Z., Gao, R., Lee, J. (2023). *Arabidopsis thaliana*: a powerful model
organism to explore histone modifications and their upstream regulations. EPIGENETICS
18: 2211362.

Yu, Z., Xu, Y., Zhu, L., Zhang, L., Liu, L., Zhang, D., Li, D., Wu, C., Huang, J., Yang, G.,
Yan, K., Zhang, S., Zheng, C. (2019). The Brassicaceae-specific secreted peptides, STMPs,
function in plant growth and pathogen defense. J Integr Plant Biol **12817**.

Zhang, Z., Luo, X., and He, Y. (2023). Cold induction of nuclear *FRIGIDA* condensation in
Arabidopsis. Nature **619**: E27–E32.

Zheng, X., Levine, D., Shen, J., Gogarten, S.M., Laurie, C., and Weir, B. (2012). A high-
performance computing toolset for relatedness and principal component analysis of SNP
data. Bioinformatics **28**: 3326–3328.

Zhou, X. and Stephens, M. (2012). Genome-wide efficient mixed-model analysis for
association studies. Nat Genet **44**: 821-824.

Zhu, Y., Siegwolf, R.T., Durka, W., and Körner, C. (2010). Phylogenetically balanced
evidence for structural and carbon isotope responses in plants along elevational gradients.
Oecologia **162**: 853–863.

Zou, X. and Sun, H. (2023). DOF transcription factors: Specific regulators of plant biological
processes. Front Plant Sci. **14**: 1044918.

1331

1332

1333

1334

1335

1336

1337

1338

1339

List of Supplemental Data

Supplemental Methods

S1. Sequencing and Variant Calling.

S2. Analysis of RGB color data.

Supplemental Results

S1. Analysis of RGB color data.

S2. Flowering time QTL in northwestern Europe.

S3. Physiological effects under high light of *PHT4;4* variants.

Supplemental Figures

S1. NJ tree of the global population genetic structure of Arabidopsis with 1233 ecotypes.

S2. NJ gene trees of vernalization-pathway gene sequences.

S3. Representative NPQ kinetics curve and parameters.

S4. Cold effect on NPQ kinetics for three subsequent light-dark cycles in 11 ecotypes.

S5. Drought effect on NPQ kinetics for three subsequent light-dark cycles in 9 ecotypes.

S6. Aboveground biomass of 11 ecotypes grown under control conditions.

S7. Regional *DOG1* haplotype variation relative to elevation and flowering time.

S8. Antioxidant-related phenotypes under HL vs LL (cold) in *PHT4;4* variants.

S9. NJ tree and motifs linked to the low antioxidant *PHT4;4* haplotype.

Supplemental Tables

S1. ANOVA on NPQ kinetics in response to cold and drought.

S2. Quantification of photosynthesis related pigments in 8 ecotypes.

S3. Information of insertional mutants in *PHT4;4*.

Supplemental Datasets

S1. Coordinates, elevation, and region of ecotypes used in all experiments.

S2. List of environmental variables, source, and definition used in the PCA of Figure 1.

S3–S5. Percival programs and HOBO data for conditions in Experiments 1–3.

S6. Dataset, top SNPs, and functional gene annotation in Flowering Time GWAS.

S7. Top SNPs and gene annotation for the low pCO_2 GWAS.

S8. Top SNPs and gene annotation for the HL vs. LL GWAS.

S9. Top SNPs and gene annotation for the night-freezing vs. no freezing GWAS.