1 Phosphoprotein Phosphatase Activity Positively Regulates Oligomeric Pyrin to Trigger
2 Inflammasome Assembly in Response to Bacterial Effectors and Toxins that Inactivate RhoA in
3 Macrophages
4
5
6

7 Haleema S. Malik[1], Nicole A. Loeven[1], Jose M. Delgado[2], Arminja N. Kettenbach[1,2] and James B.
8 Bliska[1]
9
10 [1]Department of Microbiology and Immunology, and [2]Department of Biochemistry and Cell Biology,
11 Geisel School of Medicine at Dartmouth, Hanover NH, USA
12
13 [2]Norris Cotton Cancer Center, Lebanon, NH, USA.
14
15
16 Correspondence:
17 James B. Bliska
18 Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth, 66
19 College Street, Hanover NH, 03755 USA
20 Email: james.bliska@dartmouth.edu

21
22

23
24
25   ABSTRACT:
26

Pyrin is a pattern-recognition receptor in phagocytes that triggers capase-1 inflammasome
assembly in response to bacterial toxins and effectors that inactivate RhoA. Pyrin contains coiled
coil and B-box domains that promote oligomerization. Pyrin is negatively regulated by
phosphorylation of two linker domain residues, S205 and S241 (murine) or S208 and S242
(human), via the kinase PKN, which is in turn activated by RhoA. Inactivation of RhoA by the TcdB
toxin of *Clostridioides difficile*, the YopE and YopT effectors of *Yersinia,* or the TecA effector of
Burkholderia cenocepacia triggers pyrin inflammasome assembly in macrophages. Inactivation
of RhoA removes the negative regulation but a phosphoprotein phosphatase (PPP) is needed to
positively regulate pyrin by dephosphorylation. No PPP that dephosphorylates pyrin has been
identified, the oligomeric state of murine pyrin has not been determined, and the phosphorylation
status of pyrin oligomers is unknown. We used primary murine macrophages and human THP-1
cells combined with the use of inhibitors, native PAGE, immunoblotting with phospho-specific
pyrin antibodies and siRNA knockdowns to determine if a PPP positively regulates oligomeric
pyrin. Results obtained with inhibitor calyculin A indicate that PPP activity is required to
dephosphorylate S205 or S242 in murine or human pyrin, respectively, in response to inactivation
of RhoA by TcdB, YopE/T or TecA. Findings from native PAGE show that murine pyrin exists as
dimers and higher order oligomers that are phosphorylated on S205 prior to RhoA inactivation.
Calyculin A treatment causes reduced mobility of murine pyrin on native PAGE and
hyperphosphorylation of S242 in human pyrin, suggesting a PPP constitutively counterbalances
PKN to keep the second linker site hypophosphorylated. Data from low dose okadaic acid and
siRNA knockdown experiments implicates PP2A in dephosphorylation of S205 and positive
regulation of pyrin in response to RhoA inactivation.

INTRODUCTION

Numerous Gram-negative bacterial pathogens encode type III secretion systems (T3SSs) that are essential for virulence (1). T3SSs translocate effectors into or across the plasma membrane and into the eukaryotic cytosol during bacterial-host cell contact. Translocated effectors can inhibit innate immune responses to promote pathogenesis (1, 2). Host cells can sense perturbations caused by T3SSs and/or effectors and produce compensatory innate immune responses to counteract infection by pathogens (3, 4). Perturbations induced by T3SSs and/or effectors in macrophages infected with bacterial pathogens can result in the assembly of caspase-1 inflammasomes (5, 6). Inflammasomes are multi-oligomeric complexes that are assembled in response to various danger signals in the cell cytosol and serve as a molecular platform for the recruitment and maturation of caspase-1 into its active form (5, 6). Active caspase-1 cleaves gasdermin-D (GSDMD), pro-IL-1β and pro-IL-18 (7-9). The N-terminal domain of GSDMD forms pores in the plasma membrane which allow for release of mature IL-1β and IL-18 (7-9). GSDMD pores can also promote cell death, termed pyroptosis, resulting in the release of additional pro-inflammatory molecules and intracellular bacteria (10). Pyroptosis amplifies the immune response by restricting growth of bacteria and facilitating bacterial clearance by neutrophils (7). Virulent bacterial pathogens can inhibit caspase-1 inflammasomes in host cells to counteract protective immune responses mediated by IL-1β, IL-18 and pyroptosis (11).

Yersinia species are Gram-negative bacterial pathogens that cause invasive human infections ranging in severity from plague (*Y. pestis*) to mesenteric lymphadenitis (*Y. pseudotuberculosis, Yptb*). These bacteria replicate in lymphoid tissues as extracellular microcolonies and require a plasmid-encoded T3SS for virulence (12). The T3SS delivers seven effectors called *Yersinia* outer proteins (Yops) into the cytosol of phagocytes that are in contact with *Yersinia* (12). The effectors inhibit key host cell processes, including phagocytosis and proinflammatory cytokine production (13). For example, YopE and YopT inhibit phagocytosis by inactivating Rho GTPases, including RhoA (13). YopE mimics a Rho GTPase activating protein (GAP) to accelerate GTP hydrolysis to GDP in RhoA (14), while YopT is a cysteine protease which cleaves RhoA at its C-terminus so that it is released from the membrane (15). Two effectors that are essential virulence factors inhibit inflammasome activation: YopM, a leucine-rich repeat (LRR)-containing protein (16-18) and YopK (19, 20). YopM inhibits the pyrin inflammasome (17, 18). YopK interacts with *Yersinia* translocon components to limit activation of NLRP3 and NLRC4 inflammasomes (19).

Inactivation of RhoA by YopE or YopT in macrophages triggers assembly of the pyrin
inflammasome as a compensatory protective innate immune response (17, 18, 21, 22). Pyrin is a
cytosolic pattern recognition receptor (PRR) which specifically senses pathogen induced
modifications to RhoA (or RhoB or RhoC) to assemble an inflammasome (23). RhoA, RhoB and
RhoC appear to be functionally redundant in regulating pyrin activity (23) and will be referred to
as RhoA from here on. Inactivation of RhoA by the TcdB toxin of *Clostridioides difficile* or the
TecA effector of *Burkholderia cenocepacia* also triggers pyrin inflammasome assembly in
macrophages (23, 24). Pyrin is encoded by the *MEFV* gene (*Mefv* in mice) whose expression is
restricted to phagocytes such as neutrophils, monocytes and activated macrophages (25). Pyrin
is a TRIM protein and like other members of this family (26) human pyrin forms dimers via coiled
coil (CC) domains (27). Additionally, human pyrin has been shown to form trimers (28). Pyrin
potentially forms higher order oligomers (i.e. dimers of dimers), via B-box domains (21). It is not
known if murine pyrin forms oligomers. Structurally, murine and human pyrin are similar except
for their C-terminus. Murine pyrin lacks the B30.2 domain present in human pyrin (29). However,
the mechanism of pyrin regulation appears to be conserved in both species (30).

Negative regulation of pyrin occurs in a manner similar to the guard hypothesis observed in plants,
whereby a plant disease resistance PRR recognizes modifications to host proteins made by
pathogens and induces an innate response (31). More specifically, pyrin follows an indirect guard
hypothesis in which it does not directly interact with but is negatively regulated by RhoA (23).
Hence, pyrin acts as the guard of RhoA (guardee) to protect host cells from bacterial effectors or
toxins which target it. RhoA functions through a switch like mechanism between an "on" (Rho-
GTP) and "off" (Rho-GDP) state. RhoA has numerous roles in phagocytosis, cell cycle and
migration and is targeted by multiple bacterial pathogens (23, 32). Pyrin is negatively regulated
by serine phosphorylation, via the kinase PKN (also known as PRK), which is in turn is activated
by RhoA. PKN phosphorylates murine pyrin on serines 205 and 241 (S208 and S242 in human
pyrin) within 14-3-3-binding sites located in the linker region to negatively regulate pyrin (30, 33).
Mutational analysis of S205 and S241 by alanine substitution in ectopically expressed murine
pyrin indicates that loss of either site results in partial inflammasome activation and when both
sites are disrupted there is an additive effect (30). Similar studies of S208 and S242 in human
pyrin show that the inflammasome is activated when either site is substituted (33, 34). Moreover,
an S242R mutation in MEFV causes an autoinflammatory disease, PAAND, which leads to a
partially active form of pyrin and further underscores the importance of linker phosphorylation in
pyrin regulation (35). When pyrin is phosphorylated, the N-terminal pyrin domain (PYD) may be

sequestered, preventing interaction with ASC (21). RhoA inactivation removes the negative
regulation axis of pyrin, presumably resulting in dephosphorylation of S205/S208 and S241/S242
(21, 22, 30, 33). These findings lead to a model wherein when both serines in the pyrin linker are
dephosphorylated, 14-3-3 detaches, the PYD is exposed, and ASC and caspase-1 are recruited
to assemble inflammasomes (21, 30, 33). A limitation of this model is that no phosphoprotein
phosphatase (PPP) that dephosphorylates pyrin has been identified and the phosphorylation
status of pyrin oligomers is unknown.

Assembly of the pyrin inflammasome in phagocytes infected with WT *Yersinia* is blocked due to
the action of YopM (17, 18). YopM binds to and activates the protein PKN and a second kinase,
RSK (36). YopM hijacks PKN and RSK to keep pyrin phosphorylated (17, 21, 34). A *Yptb ΔyopM*
mutant is fully virulent in Mefv[-/-] mice, demonstrating that YopM promotes virulence by inhibiting
pyrin (17). The importance of pyrin in pathogenesis extends beyond *Yersinia, C. difficile* and *B.*
cenocepacia as numerous other bacteria including *Clostridium botulinum*, *Vibrio*
parahaemolyticus and *Histophilus somni* secrete toxins that trigger assembly of this
inflammasome in phagocytes (23).

Codon changes in the *MEFV* gene that result in gain of function pyrin variants are responsible for
the monogenic human autoinflammatory disease Familial Mediterranean Fever (FMF) (25, 34).
FMF is characterized by recurrent episodes of fever and serositis and primarily effects eastern
Mediterranean populations (37). Magnotti et al demonstrated that pyrin dephosphorylation was
sufficient to promote inflammasome activation using PKN inhibitors on monocytes from FMF
patients, which implies that these patients have a lower threshold for pyrin activation than healthy
patients (38). It has been hypothesized that the high carrier frequency of FMF in Mediterranean
and Middle Eastern populations is the result of a selective advantage in resistance to an unknown
infection (34). Population genetic evidence is consistent with the idea that plague selected for
FMF mutations in Middle Eastern populations (34). In addition, infection assays with human cells
and mice provided evidence that pyrin FMF variants promote resistance to *Y. pestis* (34). These
finding underscore the importance of the pyrin inflammasome for human health. Therefore, it is
important to understand the dual role of pyrin in promoting immune responses against pathogens
and FMF and a deeper understanding of pyrin inflammasome regulation will aid in this goal.

Here we used primary murine macrophages and human THP-1 cells in conjunction with PPP
inhibitors, native PAGE, immunoblotting with phospho-specific pyrin antibodies and siRNA
knockdowns to determine if PPP activity is required to dephosphorylate oligomeric pyrin and

positively regulate inflammasome assembly in response to bacterial effectors and toxins (YopE,

YopT, TecA or TcdB) that inactivate RhoA.

RESULTS

PPP activity is required for pyrin dephosphorylation and positive regulation of

inflammasome assembly

Inactivation of RhoA removes negative regulation by PKN, however, a PPP is needed to positively

regulate pyrin by serine dephosphorylation. The PPP family consists of PP1, PP2A, PP2B (also

known as calcineurin or PP3), PP4, PP5, PP6, and PP7 (39). To test the role of PPPs in

dephosphorylation of pyrin LPS-primed bone marrow-derived macrophages (BMDMs) were

pretreated for 15 min with DMSO as a vehicle control or one the following inhibitors: calyculin A

(CA) at 10 nM and okadaic acid (OA) at 1000 nM, which at these concentrations inhibit PP1,

PP2A, PP4, PP5, and PP6 but not PP2B; OA at 100 nM to more specifically inhibit PP2A; and

1000nM of cyclosporin A (CSA) which inhibits PP2B. After the pretreatment the BMDMs were

maintained in the presence of DMSO or the inhibitors during intoxication for 90 min with *C. difficile*

TcdB, which glucosylates and inactivates RhoA (40), to trigger pyrin inflammasome assembly

(23). BMDM lysates were analyzed by immunoblotting with commercially available monoclonal

antibodies specific for total pyrin or phosphoserine 205 (PS205) (30) or actin as loading control.

We were unable to monitor PS241 in murine pyrin because the antibody reported to recognize

this modification (30) did not work in our hands for this purpose but did recognize human pyrin

PS242 (see Fig.S3 below). We found that treatment of BMDMs with CA or 1000 nM OA prevented

pyrin S205 dephosphorylation in response to TcdB intoxication (Fig. S1A). Results of ELISA on

BMDM supernatants showed that CA or 1000 nM OA also significantly reduced release of mature

IL-1β during TcdB intoxication (Fig. S1B). CSA treatment did not inhibit pyrin S205

dephosphorylation or IL-1β release upon TcdB intoxication (Fig.S1) which indicates that PP2B

does not regulate this inflammasome. 100nM OA treatment was also ineffective (Fig. S1) but

previous publications (41) and subsequent experiments (see Fig. 4 below) indicate that at this

concentration longer pretreatments are needed for the inhibitor to reach effective concentrations

in cells. We noted that CA and 1000nM OA in the absence or presence of TcdB caused an upshift

in the positions of pyrin bands on the immunoblots, but not a corresponding increase in PS205

signal (Fig.S1A). This observation suggests that S205 is uniformly phosphorylated and a PPP

constitutively dephosphorylates pyrin at an additional site(s), possibly S241, and when this

phosphatase is inhibited, hyperphosphorylation by PKN results in reduced mobility of pyrin by

SDS-PAGE. In summary, these results suggest that a PPP other than PP2B is required to

dephosphorylate pyrin S205 and trigger assembly of the inflammasome in response to TcdB. In
addition, a PPP may act constitutively to dephosphorylate pyrin at another site(s).

We used CA treatment to determine if a PPP is important for YopE and YopT to trigger pyrin S205
dephosphorylation and inflammasome assembly in BMDMs infected with *Yptb.* BMDMs treated
with DMSO or CA as above were infected with a Δ*yopM* mutant to activate the pyrin
inflammasome. We also infected DMSO or CA-treated BMDMs with a Δ*yopK* mutant, which
triggers assembly of the NLRP3 inflammasome (and to a lesser degree the NLRC4
inflammasome) (19, 20), since there is evidence that PP2A is important for dephosphorylation
and activation of NLRP3 (42). Immunoblot and ELISA analysis of BMDMs pre-treated with CA
followed by Δ*yopM* infection showed that pyrin remained phosphorylated on S205 and there was
a significant decrease in IL-1β release (Fig. 1). Results of infection with Δ*yopK* also showed
reduced IL-1β release when the BMDMs were treated with CA, although the decrease was not
significant (Fig. 1B). Pyrin was not dephosphorylated on S205 in response to Δ*yopK* infection of
untreated BMDMs as expected (Fig. 1A). The finding that CA had a stronger inhibitory effect on
IL-1β secretion with Δ*yopM* as compared to Δ*yopK* infection (Fig. 1B) might reflect residual
NLRC4 inflammasome assembly triggered by the latter strain (19).

Similar results were obtained when CA was used to treat BMDMs infected with Δ*yopM Yptb*
strains expressing only active YopE (*yopT*^{C139A}) or YopT (*yopE*^{R144A}) (Fig. S2). In addition,
immunoblotting indicated that CA did not decrease steady state levels of pro-IL-1β (Fig. S2A).
199
B. cenocepacia infection of naive THP-1 cells has been shown to trigger assembly of the pyrin
inflammasome (43). To determine if PPP inhibition prevents dephosphorylation of human pyrin
we infected THP-1 cells with *B. cenocepacia* WT or Δ*tecA* strains in the absence or presence of
DMSO or CA*.* Lysates were immunoblotted with antibodies that recognize total pyrin, PS242,
pro-IL-1β or actin as loading control. *B. cenocepacia* infection triggered dephosphorylation of
S242 in a TecA-dependent manner in DMSO-treated THP-1 cells (Fig.S3). Interestingly, when
THP-1 cells were pretreated and maintained in the presence of CA S242 became
hyperphosphorylated in both uninfected and infected conditions (Fig.S3), which was evident from
comparing the short and long exposures of the immunoblots. The hyperphosphorylation of pyrin
in the presence of CA in uninfected THP-1 cells suggests that a PPP is constitutively
counterbalancing PKN to keep S242 hypophosphorylated. CA treatment prevented infection-
induced production of pro-IL-1β as determined by immunoblotting (Fig.S3) and as a result the

impact of the inhibitor on release of mature IL-1β from the THP-1 cells could not be measured.

These data indicate that a PPP dephosphorylates pyrin S242 in response to inactivation of RhoA

by TecA in THP-1 cells, however, it also appears that a PPP is acting constitutively to keep most

of the pyrin in the cell dephosphorylated at this site prior to infection.

PPP activity regulates phosphorylation of oligomeric pyrin

Although human pyrin has been shown to form trimers (28) the oligomeric status of murine pyrin

has not been investigated.   In addition, it is not known if pyrin oligomers are differentially

phosphorylated or dephosphorylated.   We used blue native PAGE (BN-PAGE) and

immunoblotting (44, 45) to examine the impact RhoA inactivation and PPP inhibition on

phosphorylation of oligomeric pyrin. To optimize the procedure, we first tested different non-ionic

detergent extraction conditions on untreated uninfected WT or *Mefv*[-/-] BMDMs. As shown in

Fig.S4, when extracts of untreated BMDMs were separated by BN-PAGE and immunoblotted for

total (A) or PS205 (B) pyrin, we detected oligomers of pyrin in WT but not *Mefv*[-/-] BMDMs.

Monomeric pyrin (~90 kDa) was not detected and based on the apparent molecular weight of the

oligomers we tentatively assigned them as dimer (~242 kDa) and higher-order oligomers (~480-

720 kDa). Thus, most of the pyrin in BMDMs is in a higher oligomeric form and phosphorylated

on S205. Recovery of total and PS205 pyrin was optimal using 0.5% or 1% digitonin (Fig.S4) and

1% was used going forward.

We next analyzed the phosphorylation and oligomeric status of pyrin in BMDMS infected with WT

or Δ*yopM Yptb* in the absence of CA using BN-PAGE and immunoblotting. As shown in Fig.2, the

total (A) and PS205 (B) pyrin oligomer signals decreased in BMDMs infected with Δ*yopM*, due

increased insolubility of the dephosphorylated form of pyrin in non-ionic detergent (E) (22). This

insolubility issue prevented us from determining if the oligomeric state of pyrin is altered upon

dephosphorylation. In BMDMs treated with CA there was an upshift in the positions of the dimeric

and higher oligomeric forms under all conditions, most evident with the PS205 signal, and there

was no insolubility or dephosphorylation of pyrin in response to infection with Δ*yopM* (Fig.2CD).

In addition, in the presence of CA a very large mystery oligomer at ~1048 kDa was detected with

the PS205 signal under all conditions (Fig.2D). These results confirm the upshift in mobility due

to hyperphosphorylation of pyrin in the presence of CA seen in some SDS-PAGE immunoblots

(Fig.S1A). These data further suggest that hyperphosphorylation of pyrin increases

oligomerization.

YopM interacts with oligomeric pyrin

YopM can interact with pyrin as determined by co-immunoprecipitation (17, 18, 34). We wondered if YopM interacts with the different oligomeric forms of YopM and if this interaction is sensitive to PPP inhibition. To determine if YopM is associated with pyrin oligomers, samples from untreated WT BMDMs left uninfected or infected with WT or Δ*yopM Yptb* were separated by BN-PAGE and immunoblotted with antibody to YopM. We detected a YopM signal at the dimer and higher-order oligomer positions in the WT but not in the Δ*yopM Yptb* samples (Fig.3A, lanes 2 and 3). We then repeated the experiments using WT BMDMs untreated or treated with CA or untreated *Mefv⁻ᐟ⁻* BMDMs. In WT BMDMs there was a YopM signal corresponding to dimeric pyrin in the DMSO and CA treated WT *Yptb* infected samples (Fig.3B, lanes 2 and 5). The higher order oligomeric YopM signal around ~480kDa seen with DMSO treatment was not present in the CA treated sample (Fig.3B, lane 5). In addition, there was a faint YopM signal around ~480kDa in the *Mefv⁻ᐟ⁻* BMDMs infected with WT *Yptb* (Fig.3B, lane 8). Two dimensional BN-PAGE immunoblot analysis of the DMSO treated and WT *Yptb* infected BMDMs sample showed a good correspondence in the molecular weights of the YopM and oligomeric pyrin signals (Fig. 3C). These results strongly indicate that YopM interacts with dimeric and higher order oligomers of pyrin. Possible explanations for why YopM interaction with the higher order pyrin oligomer is sensitive to PPP inhibition, and the presence of a weak YopM signal at this position in Mefv-/- BMDMs are discussed below (see Discussion).

The PP1 inhibitor tautomycetin does not prevent pyrin dephosphorylation or inflammasome assembly

To further delineate which PPP activities positively regulate pyrin, we used tautomycetin (TTN), a more specific inhibitor of PP1 (46, 47). WT BMDMs were pretreated for 15 min and maintained in DMSO or 1 or 2μM of TTN during infection with Δ*yopM Yptb*. We found that treatment of BMDMs with TTN did not prevent pyrin PS205 dephosphorylation or IL-1β release in response to infection with Δ*yopM Yptb* (Fig.S4). Longer pretreatment of BMDMs with 1μM TTN also did not decrease IL-1β secretion upon Δ*yopM* infection (data not shown). Our combined results from treatment of BMDMs with various phosphatase inhibitors suggest that a PPP other than PP1 and PP2B is required to activate pyrin.

Longer pretreatment with 100 nM OA reduces pyrin dephosphorylation and inflammasome assembly

As mentioned previously, OA penetration inside cells is slow and using a lower concentration of

OA generally requires longer pretreatment time to achieve sufficient inhibitory levels inside the

cell. In contrast CA readily partitions inside cells (41, 48, 49). We found that WT BMDMs

pretreated for 3 hrs and maintained with 100nM OA during infection with Δ*yopM Yptb* showed

reduced pyrin dephosphorylation (Fig. 4A) and a significant decrease in IL-1β release as

compared to the DMSO control (Fig.4B); whereas 15min pretreatment of BMDMs with 100nM OA

did not have such an inhibitory effect on pyrin activation (Fig. S1). Given the selectivity of 100 nM

OA to preferentially inhibit PP2A/PP4 over PP1/PP5 (49), these results suggest that PP2A or PP4

could be involved in positively regulating pyrin. Since most serine/threonine phosphatase

activities inside the cell are carried out PP1 and PP2A (39, 50) our results with TTN and longer

pretreatment with OA in BMDMs allude to a role for PP2A in positive regulation of pyrin activation.

siRNA knockdown implicates PP2A in dephosphorylation of pyrin S205

We next tested the effect of knocking down PP2A in BMDMs on pyrin inflammasome assembly.

PP2A has two catalytic subunit isoforms, alpha (PP2Aca) and beta (PP2Acb), which only differ in

a few amino acids from each other. We first tested two different siRNAs targeting the mRNA of

the alpha catalytic subunit of PP2A (42). The siRNAs (1000pmol) were electroporated into WT

BMDMs which were also LPS-primed. PKN1 siRNA was used as a positive control (17, 33). RT-

qPCR showed that there was significant and selective knock down of PP2Aca mRNA with both

siRNAs as compared to the positive control and PP2Acb analyzed in parallel (Fig. 5A). The siRNA

electroporations were then repeated and LPS-primed BMDMs were left uninfected or infected

with Δ*yopM Yptb.* Immunoblotting showed that there was full depletion of PKN1 with the

corresponding siRNA but only an apparent ~50% knock down of PP2A with the siRNAs #1 and

#2 (Fig.5B). However, since the PP2A antibody used detects both isoforms the remaining signal

detected could have corresponded to PP2Acb. Upon Δ*yopM* infection there was no inhibition of

dephosphorylation of pyrin at S205 with either PP2Aca siRNAs (Fig. 5B). We next tested the

effect of a combined PP2A alpha and beta catalytic subunit knock down. BMDMs were

electroporated with PKN1 or PP2Aca #2 siRNA and PP2Acb #1 siRNA (500pmol each) and LPS

primed as before. RT-qPCR showed that there was a decrease in the mRNA transcripts of both

PP2A subunits (Fig. 5C). Immunoblotting showed a greater depletion of PP2A protein as

compared to knocking down the alpha subunit alone (compare Fig. 5B and D). Upon Δ*yopM*

infection PS205 pyrin was maintained and a trend toward decreased IL-1β secretion was

observed (Fig. 5D and E). Together, these results implicate PP2A alpha and beta subunits as

acting redundantly to dephosphorylate pyrin S205.

DISCUSSION

In this study, we aimed to understand if a PPP activity is important for positive regulation of

inflammasome assembly by oligomeric pyrin. There are seven different PPPs in the mammalian

cell: PP1, PP2A, PP2B (PP3), PP4, PP5, PP6 and PP7. In the human genome there are ~428

Ser/Thr kinases but only ~30 Ser/Thr phosphatases (50). The remarkable diversity of PPP

substrates, substrate recognition and cellular localization is achieved by the association of

catalytic subunits with non-catalytic subunits to form multimeric holoenzymes (39). The active

sites of these phosphatases share 100% similarity, and most phosphatase inhibitors which bind

to the PPP active sites are broadly specific for this reason.

CA is a relatively non-specific and hydrophobic PPP inhibitor. It can easily partition through cell

membranes and potently inhibits most PPPs except for PP2B and PP7 (49). Using 10nM CA we

obtained evidence that a PPP activity is important for pyrin S205 dephosphorylation and

inflammasome assembly in BMDMs in response to RhoA inactivation by TcdB or YopE/T during

$\Delta yopM$ $Yptb$ infection. The same CA treatment also prevented TecA-dependent

dephosphorylation of pyrin S242 in THP-1 cells infected with $B.$ $cenocepacia.$ It has been shown

previously in mouse peritoneal macrophages that 50nM CA prevented activation of NLRP3, AIM2

and NLRC4 inflammasomes, indicating that PPP activity can be important for inflammasome

activation in general, and is perhaps acting at a latter step (51). We observed that 10nM CA

prevented infection-induced pro-IL-1β synthesis in THP-1 cells infected with $B.$ $cenocepacia$

illustrating that results with this inhibitor need to be carefully controlled and interpreted since it

can block multiple steps in an inflammasome pathway. Results from our experiments with

treatment of BMDMs with CA and $\Delta yopK$ $Yptb$ mutant infection showed that there was a non-

significant decrease in IL-1β release which can be attributed to inhibition of NLRP3 but not of the

NLRC4 inflammasome. Luheshi et al used a higher CA concentration of 50nM (51) which could

explain why we didn't observe inhibition of NLRC4 inflammasome in response to $\Delta yopK$ infection

with 10nM CA treatment.

We were also able to show using BN-PAGE that murine pyrin forms oligomers consisting of

dimers and higher order oligomers that are phosphorylated on S205 when pyrin is inactive. Due

to the insolubility of active pyrin in non-ionic detergent we were unable to determine if

dephosphorylation changes the oligomeric status of pyrin. However, PPP inhibition with CA

resulted in the appearance of a very high molecular weight mystery oligomer which indicates that

phosphorylation increases oligomerization of pyrin. Pyrin belongs to the TRIM family of proteins

and Trim5α which is involved in HIV viral capsid recognition is known to form dimers and higher

order lattices consisting of dimers of dimers or trimers of dimers (26). The higher order murine

pyrin oligomers (~480-720kDa) that we observe on BN-PAGE immunoblots could represent

dimers of dimers or trimers of dimers similar to those formed by Trim5α. It has been previously

shown through crosslinking studies in THP-1 cell lysates that human pyrin can form trimers which

is mediated through its coiled coil domains (28). In the same study it was also shown that the B-

box domain of human pyrin interacts with the N-terminal pyrin domain to sequester it and prevent

interaction with ASC (28). However, Weinert et al reported that a truncated version of human pyrin

forms an antiparallel dimer in vitro through interaction of coiled coil domains (27). These

discrepancies in trimer vs dimer formation by human pyrin could be attributed to experimental and

technical differences. It is expected that regulation of human and murine pyrin by oligomerization

would be conserved but due to structural differences, it is possible that murine pyrin forms dimers

and human pyrin forms trimers through their respective coiled coil domains. To rule out technical

differences it would be informative to investigate oligomerization of human pyrin by BN-PAGE

instead of crosslinking. The phosphorylation status of human pyrin oligomers should also be

determined.

CA treatment resulted in an upshift in molecular weights of PS205 and total murine pyrin bands

in all conditions on BN-PAGE immunoblots, as wells as on some SDS-PAGE immunoblots. In

addition, CA treatment of THP-1 cells resulted in hyperphosphorylation of S242 in human pyrin.

This indicates that a PPP acts constitutively on the second linker site in pyrin to dephosphorylate

it and when this PPP is inhibited the second serine becomes hyperphosphorylated. To our

knowledge this is the first evidence that one of the linker sites can be hypophosphorylated yet

pyrin remains inactive. This finding is surprising given the data that phosphorylation of the second

linker site is critical for negative regulation of pyrin, based on alanine substitution of this residue

in ectopically expressed pyrin variants (33, 34). In addition, the S242R substitution causes

PAAND in which the pyrin variant is considered constitutively active (35). Pull down experiments

in this study also showed that 14-3-3 proteins are not bound to the S242R variant and that pyrin

is dephosphorylated (35). Jeru et al. showed previously that a S208A pyrin variant has reduced

14-3-3 binding while the S242A mutant abolishes 14-3-3 binding (52). This suggested that 14-3-

3 binding could follow a gatekeeper hypothesis in which PS242 is the dominant site which

promotes 14-3-3 binding and PS208 is a secondary site (52). These findings may indicate that
hypophosphorylation of the second linker site is sufficient to keep pyrin oligomers inactive. This
could facilitate fast dephosphorylation of the hypophosphorylated second sites, resulting in rapid
activation of pyrin in response to RhoA inactivation by pathogens. A caveat of this concept is our
limitation of not being able to measure the phosphorylation status of S241 in murine pyrin under
the conditions used.

We were also able to use one and 2D BN-PAGE and immunoblotting to detect interaction of
murine pyrin with YopM during WT *Yptb* infection of BMDMs and found that this interaction was
sensitive to PPP inhibition. The fact that YopM targets the oligomers argues that they are poised
to become activated upon dephosphorylation. We hypothesize that the faint YopM signal on BN-
PAGE immunoblots from the *Mefv*[-/-] BMDMs infected with WT *Yptb* could result from self-
oligomerization of the YopM protein. The YopM protein from *Y. pestis* is known to form tetramers
in vitro (36, 53), so it is conceivable that the *Yptb* YopM protein can form higher order oligomers
on its own in *Mefv*[-/-] BMDMs. One explanation for the selective loss of the YopM signal at the
position of the higher order oligomers from the CA-treated and WT *Yptb*-infected samples is that
the affinity of YopM binding to pyrin is reduced by hyperphosphorylation.

OA in comparison to CA has a greater inhibitory constant to bind to the catalytic subunit of PP2A
than of PP1 and is reported to enter cells slowly (54{Swingle, 2007 #2701). Our results indicated
that 15min pretreatment with 1000nM OA is sufficient to inhibit S205 pyrin dephosphorylation and
IL-1β release in BMDMs upon RhoA inactivation. Stutz et al reported that PP2A is important for
NLRP3 activation through dephosphorylation using the same conditions and concentration of OA
(42). We only observed a reduction in S205 dephosphorylation and a significant decrease in IL-
1β secretion in response to Δ*yopM* infection when BMDMs were pretreated with 100nM OA for
3hr but not for 15min, which can be explained by the slow entry of the inhibitor. We hypothesize
from these results that PP2A plays a role in dephosphorylation of S205 and positive regulation of
the pyrin inflammasome in BMDMs.

PP2A, along with PP1, carry out most serine/threonine dephosphorylation events in cells and
partake in many essential activities for the cell (39). PP2A is abundantly present in cells and forms
heterotrimers consisting of a catalytic subunit, a scaffolding A subunit, and a regulatory B subunit
(39). There are two isoforms of the catalytic and scaffolding subunits, and four isoforms of the
regulatory subunits, which account for the diverse activities and substrates regulated by PP2A

(50). The alpha and beta catalytic subunit of PP2A only differ by a few amino acids and PP2Acα

is more abundantly present in cells than PP2Acβ. Our results indicated that siRNAs targeting

PP2Acα did not reduce S205 pyrin dephosphorylation in BMDMs infected with Δ*yopM*. However,

a combined knock down of both catalytic subunits of PP2A showed that S205 pyrin

phosphorylation was maintained and there was a trend toward decreased in IL-1β secretion upon

Δ*yopM* infection. Perhaps the two catalytic subunits of PP2A act redundantly to dephosphorylate

the pyrin linker. This is distinct from the case of NLRP3 where knockdown of the alpha catalytic

subunit of PP2A alone was sufficient to prevent assembly of the inflammasome (42). To confirm

our results, additional siRNAs targeting the alpha and beta catalytic subunit of PP2A should be

tested. In addition, the impact of PP2A knockdowns on PS241 in murine pyrin and PS208/242 in

human pyrin needs to be investigated.

PPPs recognize and bind to regulatory proteins or to substrates via short linear motifs (SLiMs)

(39). For example, the B56 regulatory subunit of PP2A binds through LxxIxE motifs to its

substrates (39). Interestingly, murine pyrin contains two LxxIxE motifs in the coiled-coil domain

and human pyrin has one LxxIxE motif in the linker. Since both murine and human pyrin have a

PP2A-B56 regulatory motif it is possible that PP2A regulates both forms through a conserved

mechanism. This could be tested by inactivation of the SLiM motifs in murine and human pyrin.

We have not ruled out the possible roles of other PPPs in pyrin dephosphorylation and

inflammasome assembly and additional approaches will likely be needed to address this

possibility. Finally, it remains to be determined if the activity of PP2A or other PPP is upregulated

upon RhoA inactivation, or these enzymes simply constitutively counterbalance PKN. In the case

of PP2A activating NLRP3 by dephosphorylation (42), there is evidence that Bruton tyrosine

kinase negatively regulates PP2A by phosphorylation of tyrosine 307 (55). It is theoretically

possible that PKN could negatively regulate PPP activities by phosphorylation and additional

experiments are needed to address this possibility.

MATERIALS AND METHODS

Bacterial strains. *Y. pseudotuberculosis* (*Yptb*) 32777 (serogroup O1) strains used in this study

are as follows: wild-type (56), Δ*yopM* (57), Δ*yopK* (58), *yopE*^{R144A}Δ*yopM* and *yopT*^{C139A}Δ*yopM*

(17). *Yptb* was grown at 28°C on Luria broth (LB) agar plates or with shaking in LB broth,

supplemented with antibiotics when necessary. The *B. cenocepacia* strains used were AU1054

(WT) and Δ*tecA* mutant (59). *B. cenocepacia* was grown at 37°C on LB agar plates or with

shaking in LB broth.

Bone marrow isolation and cell culture conditions. Bone marrow derived macrophages

(BMDMs) were obtained from femurs of 8-week-old WT C57BL/6 mice (Jackson Laboratories)

and/or *Mefv*[-/-] mice (60) as previously described (19). Cells were grown at 37°C in a humidified

incubator in macrophage growth media (MGM) made up of Dulbecco's modified Eagle Medium

(DMEM) plus GlutaMAX (Gibco) containing 10% heat inactivated fetal bovine serum (FBS) (GE

Health Care), 10% L929 cell-conditioned medium, 1 mM sodium pyruvate (Gibco), 10 mM HEPES

(Gibco) for a week. On day 7 differentiated macrophages were divided into six-well plates at a

density of 0.8×10^6 cells/well in a total volume of 3 ml in MGM diluted to contain 10% L929 cell-

conditioned medium (MGM 10/10) and primed for ~18hrs with 100 ng/ml O26:B6 *Escherichia coli*

LPS (Sigma).

BMDM infection, intoxication, and inhibitor treatments. For BMDM infections, *Yptb* strains

were grown overnight in LB at 28°C. The following day, cultures were diluted 1:40 in LB containing

20 mM sodium oxalate and 20 mM MgCl2 and grown at 28°C for 1 hr, then shifted to 37°C for 2

hr. Cultures were centrifuged and the bacterial pellets were resuspended in phosphate-buffered

saline (PBS). Bacterial suspensions were then diluted in media lacking FBS and containing 10%

L929 cell-conditioned medium (MGM 0/10). LPS-primed BMDMs were washed once with PBS

and left uninfected or infected in MGM 0/10 with *Yptb* strains cultured under conditions as

described above at an MOI of 30. Plates were centrifuged for 5 min at 106xg to facilitate contact

between *Yptb* and macrophages. Plates were then incubated at 37°C in 5% CO_2. In experiments

where the glucosyltransferase toxin TcdB from *Clostridium difficile* (List Biological Laboratories,

Inc.) was used as a positive control for pyrin inflammasome activation, BMDMs were intoxicated

with TcdB at a concentration of 0.1 µg/ml for 90min in MGM 0/10 media.

In experiments where phosphatase inhibitors were used and BMDMs were pre-treated for 15min

with either 1.4mM DMSO as vehicle control or calyculin A (10nM diluted from 100µM stock, LC

Laboratories), cyclosporin A (1µM diluted from 1mM stock, Sigma), okadaic acid (100nM or 1µM

diluted from 1mM stock, Tocris), or tautomycetin (1µM or 2µM diluted from 1mM stock, Tocris) in

MGM 0/10. MGM 10/10 was used for BMDMs pre-treated for 3hr with 100nM okadaic acid. For

infection following pre-treatment, MGM 0/10 media containing bacteria and DMSO or the

respective phosphatase inhibitors was added to the BMDMs, followed by centrifugation and

incubation as above. Post-infection, cell supernatants were collected and processed for IL-1β

detection by ELISA and cell lysates were prepared and processed for immunoblot analysis (see

below).

THP-1 cells, infection, and inhibitor treatment. THP-1 cells (ATCC TIB-202) were grown in

filter sterilized RPMI 1640 media (Cat. No 30-2001) containing 10% heat inactivated fetal bovine

serum and 1.7μl 14.3M β-mercaptoethanol (to prevent cells from clumping) in 25cm^2 flasks.

When confluent the THP-1 cells were expanded in 75cm^2 flasks. No cells were used that were

passaged more than 30 times. For infections, THP-1 cells were seeded at a density of 0.8x10^6

cells per well in 1.5ml of media in 6-well plates and pretreated with ? mM DMSO vehicle control

or 10nM calyculin A for 15 min. Overnight (16-hour) cultures of *B. cenocepacia* were subcultured

1:100 in fresh LB on infection day and shaken at 37°C until the cultures reached mid-log phase

(OD$_{600}$ = 0.300). Cultures were then centrifuged, the LB was removed, and the bacterial pellets

were resuspended in PBS and then diluted to an MOI of 20 in warmed RPMI 1640 media without

FBS and containing DMSO or calyculin A. 1.5ml of supplemented media was added to each well

and mixed by pipetting gently up and down. Following incubation for 3 hours at 37°C, the contents

of each well was collected and centrifuged for 10 minutes at 14,000rpm and 4°C. The cell pellet

was collected and lysed for immunoblot analysis as described below.

Protein analysis by SDS-PAGE and immunoblotting. To obtain cell lysates, BMDMs and THP-

1 cells were lysed in mammalian protein extraction reagent (MPER, Thermo Scientific) along with

cOmplete Mini (Roche) protease inhibitor and PhosSTOP (Roche) phosphatase inhibitor. The

lysis buffer for THP-1 cells was supplemented with 10 nM calyculin A. Protein concentration was

normalized after measurement with a bichinonic acid assay (BCA, Thermo Fisher Scientific) and

3-10μg of total protein cell lysates was run on 4 to 12% NuPAGE Bis-Tris SDS-PAGE gels

(Invitrogen by ThermoFisher Scientific) and transferred to polyvinylidene difluoride membranes

(ThermoFisher Scientific) using an iBlot 2 gel transfer device (Life Technologies). Membranes

were blocked in 5% nonfat dairy milk (add company) and incubated with primary antibodies

overnight. Primary antibodies used in this paper were: rabbit-anti-mouse monoclonal total pyrin

antibody (1:1000 dilution, ab195975; abcam), rabbit-anti-mouse pyrin polyclonal (1:1000 dilution,

(60)), rabbit-anti-mouse phospho-serine 205 monoclonal antibody (1:1000 dilution, ab201784;

abcam), rabbit-anti-mouse phospho-serine 241 monoclonal antibody (1:1000 dilution, ab201784;

abcam), rabbit-anti-mouse/human IL-1β (1:1000 dilution, number 12242; Cell Signaling), rabbit-

anti-mouse/human polyclonal β-actin (1:1000 dilution, number 4967; Cell Signaling), mouse-anti-

mouse/human/rat monoclonal PP1a (1:1000 dilution, number MA5-17239, ThermoFisher

Scientific), rabbit-anti-*Y.pestis* polyclonal YopM (1:1000 dilution, provided by Susan Straley),

mouse-anti-mouse monoclonal PKN (1:1000 dilution, number 393344, Santa Cruz

Biotechnology), rabbit-anti-mouse monoclonal PP2A (1:1000 dilution, #2038, Cell Signaling), and

rabbit-anti-mouse GSDMD (1:1000 dilution, ab209845, Abcam). Horseradish peroxidase-

conjugated anti-rabbit or mouse antibody (Jackson Laboratory) were used as secondary

antibodies. Protein bands bound by antibodies were visualized using chemiluminescent detection

reagent (GE Healthcare) on an iBright FL1500 (ThermoFisher Scientific).

Protein analysis by blue native (BN)-PAGE and immunoblotting. BN-PAGE immunoblotting

was performed as described by Kofoed and Vance (45). All reagents for this procedure were

obtained from Invitrogen except as stated. BMDMs were lysed in non-denaturing 1X Native-PAGE

Novex sample buffer with the indicated concentrations of either digitonin (5% stock) or DDM (10%

stock), MPER (Thermo Scientific) or RIPA lysis or extraction buffer (Thermo Scientific), along with

cOmplete Mini (Roche) protease inhibitor and PhosSTOP (Roche) phosphatase inhibitor. Lysates

were centrifuged at 20,817xg for 10min to obtain soluble and insoluble fractions. In some

experiments the insoluble fractions of 1% digitonin lysates were analyzed by SDS-PASGE and

immunoblotting as described above. Protein concentrations of soluble fractions were normalized

after measurement with a bichinonic acid assay (BCA, Thermo Fisher Scientific) and 10µg or

more of total protein cell lysates was mixed with NativePAGE G-250 additive and loaded on 4-

16% NativePAGE Novex 3–12 % Bis-Tris gels. NativeMark unstained protein standard was used

for molecular weight standard visualization. XCell SureLock Mini-Cell was used for running the

gels. Samples were run at room temperature (RT) at constant 150V for 105-120min using pre-

chilled buffers. Dark blue cathode buffer (10ml NativePAGE running buffer, 10ml NativePAGE

cathode additive, 180ml deionized water) was used for the inner chamber 17ntil samples had

run up to 1/3rd of the gel length after which the buffer was replaced with light blue buffer (10ml

NativePAGE running buffer, 1ml NativePAGE cathode additive, 189ml deionized water) for the

remainder of the run. The outer chamber was filled with NativePAGE running buffer. Gels were

soaked in 2X NuPAGE transfer buffer (Invitrogen) for 15min at shaking at room temperature.

Proteins were transferred to polyvinylidene difluoride membranes (ThermoFisher Scientific) using

the template program P3 (20V constant, 7min) and an iBlot 2 gel transfer device (Life

Technologies). Post transfer membranes were incubated in 8% acetic acid for 15min with shaking,

followed by ponceau S staining for 5min. Membranes were washed with distilled water for 5min

to remove excess ponceau S stain then air-dried, reactivated by immersion in methanol, followed

by blocking in 5% non-fat dry milk. Primary and secondary antibodies used are mentioned above.

Protein analysis by two-dimensional blue native (BN)-PAGE and immunoblotting. The

procedures used followed a previously described protocol (45). Samples were prepared as

described above and separated on NativePAGE Novex 3–12 % Bis-Tris gels. Gel slices

containing samples of interest were excised, trimmed to be 3mm shorter in length than the size

of the 2D well in NuPAGE 4–12 % Bis-Tris 2D Gels, then submerged and shaken at RT in 1X

NuPAGE sample buffer containing 0.1M DTT. Gel slices wwere then microwaved for ~20sec to

denature proteins followed by cooling with shaking at 10min at RT. Slices were then fitted into the

2D well and SDS-PAGE immunoblotting was performed as described above.

siRNA knock-down. siRNAs targeting *Pkn1* (s115927), *PP2Aca* (#1: s72067, #2: s72066) and

PP2Acb (#1: s72069) were obtained from Invitrogen. For siRNA gene knock-down experiments,

400pmol of *Pkn1*,1000pmol of *PP2Aca* siRNAs, and for combined PP2Ac knock down, 500pmol

each of PP2Aca #2 siRNA and PP2Acb #1 siRNA were transfected into BMDMs (1×10^6 cells) by

electroporation using the Neon Transfection System (Invitrogen) and electrical parameters of

1400V, 20ms and 2 pulses. BMDMs were then replated in 6-well plates and grown in MGM 10/10

media. 24hr following transfection, media was changed to fresh MGM 10/10 and LPS was added

for overnight incubation. 48hr following transfection BMDMs were infected with Δ*yopM Yptb* as

described above.

Quantitative real-time PCR (RT-qPCR). At 48hr following transfection, uninfected BMDMs were

lysed with trizol reagent (Invitrogen) to isolate RNA according to the manufacturer's instructions.

500-1000ng of RNA was used to make cDNA using SuperScript IV First-Strand Synthesis System

(Invitrogen). Approximately ~5ng of cDNA was quantified using PowerUp SYBR Green Master

Mix (Invitrogen) in the (add machine name). Primer specificity was checked by melt-curve

analysis. Expression of target genes was normalized to the expression of the housekeeping gene

Hprt. For fold change analysis data was transformed using the 'relative standard curve method

and comparative threshold cycle (Ct) method (ΔΔCt)' as described by Applied Biosystems.

Published primer sequences were used for qPCR analysis (42) and the sequences are: Hprt,

forward   5′-TGAAGT   ACTCATTATAGTCAAGGGCA-3′   and   reverse   5′-CTG

GTGAAAAGGACCTCTCG-3′; Ppp2ca, forward 5′-TCTTCCTCTCACTGCCTTGGT-3′ and

reverse 5′-CAG CAAGTCACACATTGGACCC-3′; Ppp2cb, forward 5′-

AAGGCGTTCACCAAGGAGCT-3′ and reverse 5′- ACAGCGGACCTCTTGCACAT-3′.

IL-1β quantification. IL-1β in BMDM supernatants was quantified using a murine ELISA kit

(MLB00C; R&D Systems) by following the manufacturer's instructions.

Statistical analysis. GraphPad Prism was used to perform statistical analyses. IL-1β ELISA data

were analyzed by one-way or two-way analysis of variance (ANOVA; Tukey's multiple-

comparison test).

Acknowledgments. We thank Isha Nasa for help in providing advice on the phosphatase

inhibitors used in Fig. S1. We would also like to thank members of the Bliska lab for providing

helpful suggestions towards this project.

FIGURES

Fig.S1. PPP inhibition with CA and OA prevents pyrin S205 dephosphorylation and reduces IL-
1β release in BMDMs intoxicated with TcdB. LPS-primed BMDMs were pretreated for 15mins
and maintained with either 10nM calyculin A (CA), 100nM okadaic acid (OA), 1000nM OA or
1000nM cyclosporin A (CSA), and left unintoxicated or intoxicated with 0.1µg/mL of TcdB for 90
mins. Unprimed BMDMs or treatment with vehicle DMSO were used as controls. A) Immunoblot
analysis of total and PS205 pyrin in BMDM lysates. Actin was used as a loading control. B)
Mature IL-1β in supernatants as quantified by ELISA. Two-way ANOVA with Bonferroni post hoc
correction was applied to calculate significance and p-values as compared to DMSO+TcdB are
indicated. P-value<0.05 was considered significant; <0.0001 (****); not significant (ns). Each data
group is presented as an average (error bars are standard deviation) of at least 3 independent
experiments.

612
Fig.1. PPP inhibition with CA reduces pyrin dephosphorylation and IL-1β release in BMDMs
infected with Δ*yopM Yptb.* LPS-primed BMDMs were pre-treated for 15mins and maintained with
either DMSO or 10nM CA, and infected with the Δ*yopM* or Δ*yopK* strain at an MOI of 30 for 90
mins. Treated uninfected BMDMs were analyzed in parallel. A) Immunoblot analysis of PS205
and total pyrin in BMDM lysates. Actin was used as a loading control. B) Mature IL-1β in
supernatants as quantified by ELISA. Two-way ANOVA with Bonferroni post hoc correction was
applied to calculate significance and p-values of each treatment as compared to its corresponding
DMSO treated infected strain is indicated. P-value<0.05 was considered significant; <0.001 (***);
not significant (ns). Each data group is presented as an average (error bars are standard
deviation) of at least 3 independent experiments.

624
625

626
Fig.S2. PPP inhibition with CA reduces pyrin dephosphorylation and IL-1β release in BMDMs
infected with Δ*yopM Yptb* strains expressing only active YopE or YopT. LPS-primed BMDMs were
pre-treated for 15min and maintained with either DMSO or 10nM CA, and either infected with the
Δ*yopM Yptb* YopE (*yopT^{C139A}*) or YopT (*yopE^{R144A}*) strains at an MOI of 30 or intoxicated with
0.1μg/mL of TcdB for 90 mins. Unprimed and uninfected BMDMs were analyzed in parallel as
controls. A) Immunoblot analysis of PS205 and total pyrin and pro-IL-1β in BMDM lysates. Actin
was used as a loading control. B) Mature IL-1β in supernatants as quantified by ELISA. Each
data group is presented as an average (error bars are standard deviation) of two technical
replicates.
636

637
638

1 2 3 4 5 6

639
Fig.S3. PPP inhibition with CA results in hyperphosphorylation of pyrin S242. Naïve THP-1 cells
were pre-treated for 15 min and maintained with either DMSO or 10nM CA, and either left
uninfected or infected with *B. cenocepacia* (Bc) WT or Δ*tecA* for 180 min at an MOI of 20.
Immunoblot analysis of total and PS242 pyrin and pro-IL-1β in THP-1 lysates is shown. PS242
pyrin signals are show with long (15m) or short (2m) exposures for comparison. Actin was
immunoblotted as a loading control.
646

647
Fig.S4. BMDM lysates prepared with digitonin are optimal for detection of pyrin oligomers using
BN-PAGE and immunoblotting. LPS-primed WT and *Mefv* BMDMs were lysed in various non-
ionic detergent concentrations as indicated. Lysate samples were separated by BN-PAGE and
immunoblotted to visualize total (A) and PS205 pyrin (B) oligomers. Positions of BN-PAGE
molecular weight standards are shown on the left of each panel, and dimer and higher order
oligomer are indicated on the right of (B).
654

657
Fig.2. Impact of PPP inhibition with CA and *Yptb* infection on pyrin oligomers and phosphorylation
as determined by BN-PAGE and immunoblotting. WT BMDMs left untreated (A,B) or pre-treated
and maintained with DMSO or 10 nM CA (C-E) were left uninfected or infected with WT or Δ*yopM*
Yptb at an MOI of 30 for 90 mins. BMDM lysate samples prepared in 1% digitonin were separated
by BN-PAGE and immunoblotted for total (A,C) or PS205 (B,D) pyrin. BN-PAGE MW standards
are shown on the left and tentative positions of pyrin oligomers are shown on the right. Insoluble
proteins from 1% digitonin lysates of DMSO treated samples were analyzed by SDS-PAGE and
total pyrin immunoblotting (E).
666

668
Fig.3. Detection of YopM interaction with pyrin oligomers by one-dimensional and two-
dimensional BN-PAGE and immunoblotting. Untreated WT BMDMs (A) or WT or *Mefv*⁻/⁻ BMDMs
pre-treated for 15 min and maintained in DMSO or CA (B, C) were either left uninfected or infected
with WT or Δ*yopM Yptb* at an MOI of 30 for 90 mins. Cell lysates were solubilized in 1% digitonin,
separated by one dimensional BN-PAGE and immunoblotted for YopM (A,B). BN-PAGE MW
standards are shown on the left and tentative positions of YopM oligomers are shown on the right.
C) A 1% digitonin sample from WT BMDMS treated with DMSO and infected with WT *Yptb* was
separated in the first dimension by BN-PAGE after which the gel slice was isolated and analyzed
by SDS-PAGE in the second dimension followed by immunoblotting for pyrin and YopM. MW
markers for each dimension are indicated.
679

680
Fig.S4. TTN treatment does not reduce dephosphorylation of pyrin or IL-1β release in BMDMs
infected with *ΔyopM Yptb.* LPS-primed BMDMs were pre-treated for 15 mins and maintained with
either DMSO or 1 or 2μM TTN and infected with *ΔyopM Yptb* at an MOI of 30 for 90 mins. Treated
uninfected BMDMs were analyzed in parallel. A) Immunoblot analysis of PS205 and total pyrin
in BMDM lysates. Actin was used as a loading control. B) Mature IL-1β in supernatants as
quantified by ELISA. Each data group is presented as an average (error bars are standard
deviation) of technical replicates from one experiment.
688

Fig. 4. Longer pretreatment with 100nM OA reduces pyrin dephosphorylation and IL-1β release
in BMDMs infected with ΔyopM Yptb. LPS-primed BMDMs were pre-treated for 3 hrs and
maintained with either DMSO or 100nM OA and infected with the ΔyopM Yptb at an MOI of 30 for
90 mins. Treated uninfected BMDMs were analyzed in parallel. A) Immunoblot analysis of PS205
and total pyrin in BMDM lysates. Actin was used as a loading control. B) Mature IL-1β in
supernatants as quantified by ELISA. Two-way ANOVA with Bonferroni post hoc correction was
applied to calculate significance and p-value of OA treatment as compared to its corresponding
DMSO treated and ΔyopM infected sample is indicated. P-value<0.05 was considered significant;
<0.01 (**). Each data group is presented as an average (error bars are standard deviation) of at
least 3 independent experiments.
700

702
703

704
Fig. 5. siRNA knock down of PP2A in BMDMs reduces pyrin S205 dephosphorylation. WT
BMDMs were electroporated with siRNAs targeting PKN1 (400pmol), PP2Aca (1000pmol) or
PP2Aca+PP2Acb (500pmol each) as indicated. 24hr after electroporation the BMDMs were LPS-
primed. 48hr following electroporation, cells were either left uninfected or infected with Δ*yopM*
Yptb at an MOI of 30 for 90 mins. A and C) RT-qPCR analysis of mRNA transcripts of PP2Aca
and PP2Acb in uninfected BMDMs. Results were normalized to Hprt mRNA levels. A) Each data
group is presented as an average (error bars are standard deviation) of three independent
experiments. One-way ANOVA was applied to calculate significance and p-values as compared
to PKN1 control siRNA are indicated. P-value<0.05 was considered significant; <0.0001 (****). C)
Results from one experiment are shown. B, and D) Immunoblot analysis of PKN1, PP2A, total
and PS205 pyrin, and pro-IL-1β in uninfected and Δ*yopM Yptb*-infected BMDM lysates. Actin was
used as a loading control. E) Mature IL-1β in supernatants as quantified by ELISA. Results
represent averages (error bars are standard deviation) from two independent experiments.
718

REFERENCES

1. Troisfontaines P, Cornelis GR. 2005. Type III secretion: more systems than you think. Physiology (Bethesda) 20:326-39.

2. Raymond B, Young JC, Pallett M, Endres RG, Clements A, Frankel G. 2013. Subversion of trafficking, apoptosis, and innate immunity by type III secretion system effectors. Trends Microbiol 21:430-41.

3. Blander JM, Sander LE. 2012. Beyond pattern recognition: five immune checkpoints for scaling the microbial threat. Nat Rev Immunol 12:215-25.

4. Vance RE, Isberg RR, Portnoy DA. 2009. Patterns of pathogenesis: discrimination of pathogenic and nonpathogenic microbes by the innate immune system. Cell Host Microbe 6:10-21.

5. Sollberger G, Strittmatter GE, Garstkiewicz M, Sand J, Beer HD. 2014. Caspase-1: the inflammasome and beyond. Innate Immun 20:115-25.

6. Shin S, Brodsky IE. 2015. The inflammasome: Learning from bacterial evasion strategies. Semin Immunol doi:10.1016/j.smim.2015.03.006.

7. Kovacs SB, Miao EA. 2017. Gasdermins: Effectors of Pyroptosis. Trends Cell Biol doi:10.1016/j.tcb.2017.05.005.

8. Shi J, Gao W, Shao F. 2017. Pyroptosis: Gasdermin-Mediated Programmed Necrotic Cell Death. Trends Biochem Sci 42:245-254.

9. Shi J, Zhao Y, Wang K, Shi X, Wang Y, Huang H, Zhuang Y, Cai T, Wang F, Shao F. 2015. Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death. Nature 526:660-5.

10. Bergsbaken T, Fink SL, den Hartigh AB, Loomis WP, Cookson BT. 2011. Coordinated host responses during pyroptosis: caspase-1-dependent lysosome exocytosis and inflammatory cytokine maturation. J Immunol 187:2748-54.

11. Brewer SM, Brubaker SW, Monack DM. 2019. Host inflammasome defense mechanisms and bacterial pathogen evasion strategies. Curr Opin Immunol 60:63-70.

12. Cornelis GR. 2006. The type III secretion injectisome. Nat Rev Microbiol 4:811-25.

13. Viboud GI, Bliska JB. 2005. Yersinia outer proteins: role in modulation of host cell signaling responses and pathogenesis. Annu Rev Microbiol 59:69-89.

14. Black DS, Bliska JB. 2000. The RhoGAP activity of the *Yersinia pseudotuberculosis* cytotoxin YopE is required for antiphagocytic function and virulence. Mol Microbiol 37:515-527.

15.   Shao F, Vacratsis PO, Bao Z, Bowers KE, Fierke CA, Dixon JE. 2003. Biochemical
characterization of the *Yersinia* YopT protease: cleavage site and recognition elements
in Rho GTPases. Proc Natl Acad Sci USA 100:904-909.
16.   LaRock CN, Cookson BT. 2012. The Yersinia virulence effector YopM binds caspase-1
to arrest inflammasome assembly and processing. Cell Host Microbe 12:799-805.
17.   Chung LK, Park YH, Zheng Y, Brodsky IE, Hearing P, Kastner DL, Chae JJ, Bliska JB.
2016. The Yersinia Virulence Factor YopM Hijacks Host Kinases to Inhibit Type III
Effector-Triggered Activation of the Pyrin Inflammasome. Cell Host Microbe 20:296-306.
18.   Ratner D, Orning MP, Proulx MK, Wang D, Gavrilin MA, Wewers MD, Alnemri ES,
Johnson PF, Lee B, Mecsas J, Kayagaki N, Goguen JD, Lien E. 2016. The Yersinia
pestis Effector YopM Inhibits Pyrin Inflammasome Activation. PLoS Pathog
12:e1006035.
19.   Brodsky IE, Palm NW, Sadanand S, Ryndak MB, Sutterwala FS, Flavell RA, Bliska JB,
Medzhitov R. 2010. A Yersinia Effector Protein Promotes Virulence by Preventing
Inflammasome Recognition of the Type III Secretion System. Cell Host Microbe 7:376-
387.
20.   Zwack EE, Snyder AG, Wynosky-Dolfi MA, Ruthel G, Philip NH, Marketon MM, Francis
MS, Bliska JB, Brodsky IE. 2015. Inflammasome activation in response to the Yersinia
type III secretion system requires hyperinjection of translocon proteins YopB and YopD.
MBio 6:e02095-14.
21.   Malik HS, Bliska JB. 2020. The pyrin inflammasome and the Yersinia effector interaction.
Immunol Rev 297:96-107.
22.   Medici NP, Rashid M, Bliska JB. 2019. Characterization of pyrin dephosphorylation and
inflammasome activation in macrophages as triggered by the Yersinia effectors YopE
and YopT. Infect Immun doi:10.1128/IAI.00822-18.
23.   Xu H, Yang J, Gao W, Li L, Li P, Zhang L, Gong YN, Peng X, Xi JJ, Chen S, Wang F,
Shao F. 2014. Innate immune sensing of bacterial modifications of Rho GTPases by the
Pyrin inflammasome. Nature 513:237-41.
24.   Aubert DF, Xu H, Yang J, Shi X, Gao W, Li L, Bisaro F, Chen S, Valvano MA, Shao F.
2016. A Burkholderia Type VI Effector Deamidates Rho GTPases to Activate the Pyrin
Inflammasome and Trigger Inflammation. Cell Host Microbe 19:664-74.
25.   Schnappauf O, Chae JJ, Kastner DL, Aksentijevich I. 2019. The Pyrin Inflammasome in
Health and Disease. Front Immunol 10:1745.

26.   Ganser-Pornillos BK, Pornillos O. 2019. Restriction of HIV-1 and other retroviruses by
787   TRIM5. Nat Rev Microbiol 17:546-556.
788   27.   Weinert C, Morger D, Djekic A, Grutter MG, Mittl PR. 2015. Crystal structure of TRIM20
789   C-terminal coiled-coil/B30.2 fragment: implications for the recognition of higher order
790   oligomers. Sci Rep 5:10819.
791   28.   Yu JW, Fernandes-Alnemri T, Datta P, Wu J, Juliana C, Solorzano L, McCormick M,
792   Zhang Z, Alnemri ES. 2007. Pyrin activates the ASC pyroptosome in response to
793   engagement by autoinflammatory PSTPIP1 mutants. Mol Cell 28:214-27.
794   29.   Chae JJ, Centola M, Aksentijevich I, Dutra A, Tran M, Wood G, Nagaraju K, Kingma
795   DW, Liu PP, Kastner DL. 2000. Isolation, genomic organization, and expression analysis
796   of the mouse and rat homologs of MEFV, the gene for familial mediterranean fever.
797   Mamm Genome 11:428-35.
798   30.   Gao W, Yang J, Liu W, Wang Y, Shao F. 2016. Site-specific phosphorylation and
799   microtubule dynamics control Pyrin inflammasome activation. Proc Natl Acad Sci U S A
800   113:E4857-66.
801   31.   Jones JD, Dangl JL. 2006. The plant immune system. nature 444:323-329.
802   32.   Thumkeo D, Watanabe S, Narumiya S. 2013. Physiological roles of Rho and Rho
803   effectors in mammals. Eur J Cell Biol 92:303-15.
804   33.   Park YH, Wood G, Kastner DL, Chae JJ. 2016. Pyrin inflammasome activation and
805   RhoA signaling in the autoinflammatory diseases FMF and HIDS. Nat Immunol 17:914-
806   21.
807   34.   Park YH, Remmers EF, Lee W, Ombrello AK, Chung LK, Shilei Z, Stone DL, Ivanov MI,
808   Loeven NA, Barron KS, Hoffmann P, Nehrebecky M, Akkaya-Ulum YZ, Sag E, Balci-
809   Peynircioglu B, Aksentijevich I, Gul A, Rotimi CN, Chen H, Bliska JB, Ozen S, Kastner
810   DL, Shriner D, Chae JJ. 2020. Ancient familial Mediterranean fever mutations in human
811   pyrin and resistance to Yersinia pestis. Nat Immunol doi:10.1038/s41590-020-0705-6.
812   35.   Masters SL, Lagou V, Jeru I, Baker PJ, Van Eyck L, Parry DA, Lawless D, De Nardo D,
813   Garcia-Perez JE, Dagley LF, Holley CL, Dooley J, Moghaddas F, Pasciuto E, Jeandel
814   PY, Sciot R, Lyras D, Webb AI, Nicholson SE, De Somer L, van Nieuwenhove E, Ruuth-
815   Praz J, Copin B, Cochet E, Medlej-Hashim M, Megarbane A, Schroder K, Savic S, Goris
816   A, Amselem S, Wouters C, Liston A. 2016. Familial autoinflammation with neutrophilic
817   dermatosis reveals a regulatory mechanism of pyrin activation. Sci Transl Med
818   8:332ra45.

819   36.   McCoy MW, Marre ML, Lesser CF, Mecsas J. 2010. The C-terminal tail of Yersinia
820   pseudotuberculosis YopM is critical for interacting with RSK1 and for virulence. Infect
821   Immun 78:2584-98.

822   37.   Jamilloux Y, Magnotti F, Belot A, Henry T. 2018. The pyrin inflammasome: from sensing
823   RhoA GTPases-inhibiting toxins to triggering autoinflammatory syndromes. Pathogens
824   and disease 76:fty020.

825   38.   Magnotti F, Lefeuvre L, Benezech S, Malsot T, Waeckel L, Martin A, Kerever S, Chirita
826   D, Desjonqueres M, Duquesne A, Gerfaud-Valentin M, Laurent A, Seve P, Popoff MR,
827   Walzer T, Belot A, Jamilloux Y, Henry T. 2019. Pyrin dephosphorylation is sufficient to
828   trigger inflammasome activation in familial Mediterranean fever patients. EMBO Mol Med
829   11:e10547.

830   39.   Nasa I, Kettenbach AN. 2018. Coordination of Protein Kinase and Phosphoprotein
831   Phosphatase Activities in Mitosis. Front Cell Dev Biol 6:30.

832   40.   Aktories K. 2011. Bacterial protein toxins that modify host regulatory GTPases. Nat Rev
833   Microbiol 9:487-98.

834   41.   Namboodiripad AN, Jennings ML. 1996. Permeability characteristics of erythrocyte
835   membrane to okadaic acid and calyculin A. American Journal of Physiology-Cell
836   Physiology 270:C449-C456.

837   42.   Stutz A, Kolbe CC, Stahl R, Horvath GL, Franklin BS, van Ray O, Brinkschulte R, Geyer
838   M, Meissner F, Latz E. 2017. NLRP3 inflammasome assembly is regulated by
839   phosphorylation of the pyrin domain. J Exp Med 214:1725-1736.

840   43.   Gavrilin MA, Abdelaziz DH, Mostafa M, Abdulrahman BA, Grandhi J, Akhter A, Abu
841   Khweek A, Aubert DF, Valvano MA, Wewers MD, Amer AO. 2012. Activation of the pyrin
842   inflammasome by intracellular Burkholderia cenocepacia. J Immunol 188:3469-77.

843   44.   Wittig I, Braun HP, Schagger H. 2006. Blue native PAGE. Nat Protoc 1:418-28.

844   45.   Kofoed EM, Vance RE. 2013. Blue native polyacrylamide gel electrophoresis to monitor
845   inflammasome assembly and composition. Methods Mol Biol 1040:169-83.

846   46.   Choy MS, Swingle M, D'Arcy B, Abney K, Rusin SF, Kettenbach AN, Page R, Honkanen
847   RE, Peti W. 2017. PP1: Tautomycetin complex reveals a path toward the development
848   of PP1-specific inhibitors. Journal of the American Chemical Society 139:17703-17706.

849   47.   Mitsuhashi S, Matsuura N, Ubukata M, Oikawa H, Shima H, Kikuchi K. 2001.
850   Tautomycetin is a novel and specific inhibitor of serine/threonine protein phosphatase
851   type 1, PP1. Biochemical and biophysical research communications 287:328-331.

852   48.   Kreienbuhl P, Keller H, Niggli V. 1992. Protein phosphatase inhibitors okadaic acid and
853   calyculin A alter cell shape and F-actin distribution and inhibit stimulus-dependent
854   increases in cytoskeletal actin of human neutrophils.

855   49.   Swingle M, Ni L, Honkanen RE. 2007. Small-molecule inhibitors of Ser/Thr protein
856   phosphatases, p 23-38, Protein Phosphatase Protocols. Springer.

857   50.   Shi Y. 2009. Serine/threonine phosphatases: mechanism through structure. Cell
858   139:468-484.

859   51.   Luheshi NM, Giles JA, Lopez-Castejon G, Brough D. 2012. Sphingosine regulates the
860   NLRP3-inflammasome and IL-1β release from macrophages. European journal of
861   immunology 42:716-725.

862   52.   Jeru I, Papin S, L'Hoste S, Duquesnoy P, Cazeneuve C, Camonis J, Amselem S. 2005.
863   Interaction of pyrin with 14.3.3 in an isoform-specific and phosphorylation-dependent
864   manner regulates its translocation to the nucleus. Arthritis Rheum 52:1848-57.

865   53.   Evdokimov AG, Anderson DE, Routzahn KM, Waugh DS. 2001. Unusual molecular
866   architecture of the Yersinia pestis cytotoxin YopM: a leucine-rich repeat protein with the
867   shortest repeating unit. J Mol Biol 312:807-21.

868   54.   MacKintosh C, Beattie KA, Klumpp S, Cohen P, Codd GA. 1990. Cyanobacterial
869   microcystin-LR is a potent and specific inhibitor of protein phosphatases 1 and 2A from
870   both mammals and higher plants. FEBS letters 264:187-192.

871   55.   Mao L, Kitani A, Hiejima E, Montgomery-Recht K, Zhou W, Fuss I, Wiestner A, Strober
872   W. 2020. Bruton tyrosine kinase deficiency augments NLRP3 inflammasome activation
873   and causes IL-1β–mediated colitis. The Journal of clinical investigation 130:1793-1807.

874   56.   Simonet M, Falkow S. 1992. Invasin expression in *Yersinia pseudotuberculosis*. Infect
875   Immun 60:4414-4417.

876   57.   McPhee JB, Mena P, Bliska JB. 2010. Delineation of regions of the Yersinia YopM
877   protein required for interaction with the RSK1 and PRK2 host kinases and their
878   requirement for interleukin-10 production and virulence. Infect Immun 78:3529-39.

879   58.   Chung LK, Philip NH, Schmidt VA, Koller A, Strowig T, Flavell RA, Brodsky IE, Bliska JB.
880   2014. IQGAP1 is important for activation of caspase-1 in macrophages and is targeted
881   by Yersinia pestis type III effector YopM. MBio 5:e01402-14.

882   59.   Loeven NA, Perault AI, Cotter PA, Hodges CA, Schwartzman JD, Hampton TH, Bliska
883   JB. 2021. The Burkholderia cenocepacia Type VI Secretion System Effector TecA Is a
884   Virulence Factor in Mouse Models of Lung Infection. mBio 12:e0209821.

885   60.   Chae JJ, Komarow HD, Cheng J, Wood G, Raben N, Liu PP, Kastner DL. 2003.

886   Targeted disruption of pyrin, the FMF protein, causes heightened sensitivity to endotoxin

887   and a defect in macrophage apoptosis. Mol Cell 11:591-604.

888