1 **Title: GASOTRANSMITTER MODULATION OF HYPOGLOSSAL MOTONEURON**
2 **ACTIVITY**
3

Abbreviated Title: Heme Oxygenase and Hypoglossal Motoneurons

Authorship: Brigitte Browe[1,2,3], Ying-Jie Peng[1], Jayasri Nanduri[1,3], Nanduri R. Prabhakar[1,2,3],
Alfredo J. Garcia III[1,2,3]

Affiliations:

[1]Institute   for   Integrative   Physiology,   The   University   of   Chicago;
[2]Grossman Institute for Neuroscience, Quantitative Biology & Human Behavior, The University
of Chicago.
[3]Department of Medicine, Section of Emergency Medicine, The University of Chicago.
13

Keywords: hydrogen sulfide, heme oxygenase 2, hypoglossal motor nucleus

Authors Contribution: Experimental design by AJG, NRP, and BB. Experiments and analyses
performed by BB, AJG, SK and JN. BB wrote the first draft. BB, AJG, and NRP revised and
edited the manuscript.

Correspondence: Alfredo Garcia: ajgarcia3@uchicago.edu

Figure Number: 10   **Abstract: 220**

Table Number: 0   **Introduction: 378**

Discussion: 937

Acknowledgements:

Funding Sources:

This work was supported by NIH P01 HL144454 (NRP) and NIH R01NS107421 (AJG).

Abstract

Obstructive sleep apnea (OSA) is characterized by sporadic collapse of the upper airway leading to periodic disruptions in breathing. Upper airway patency governed by genioglossal nerve activity originates from the hypoglossal motor nucleus. Mice with targeted deletion of the gene *Hmox2,* encoding the carbon monoxide (CO) producing enzyme, heme oxygenase-2 (HO-2), exhibit severe OSA, yet the contribution of central HO-2 dysregulation to the phenomenon is unknown. Using the rhythm brainstem slice preparations, we tested the hypothesis that central HO-2 dysregulated hypoglossal motoneuron activity. Disrupting HO-2 activity increased transmission failure as determined by the intermittent failure of the preBötzinger complex (preBötC) rhythm to produce output from the hypoglossal nucleus. Failed transmission was associated with the reduced input-output relationship between the preBötC and the motor nucleus, which was related to smaller inspiratory drive currents and reduced intrinsic excitability among hypoglossal neurons. In addition to HO-2, hypoglossal neurons also expressed the CO-regulated H_2S producing enzyme cystathionine $\square$-lyase (CSE). H_2S abundance was higher in hypoglossal neurons of HO-2 null mice than wild-type controls. Disrupting CSE function normalized transmission in HO-2 null mice and an H_2S donor mimicked the effects of HO-2 dysregulation. These findings demonstrate a hitherto uncharacterized modulation of hypoglossal activity through interaction of HO-2 and CSE-derived H_2S, and supports the perspective centrally derived HO-2 activity is important to regulating upper airway control.

Introduction

Obstructive sleep apnea (OSA) is a prevalent breathing disorder affecting nearly a billion people throughout the world[1,2]. It is characterized by the periodic cessation of breathing during sleep due to sporadic upper airway collapse. When left untreated, OSA predisposes the individual to a variety of diseases including hypertension[3,4], diabetes[5,6], and cognitive decline[7,8]. Multiple factors contribute to the genesis of OSA including compromised pharyngeal anatomy [9,10], inadequate upper airway muscle function[11-13], low arousal threshold[14], and a hypersensitive chemoreflex (i.e., high loop gain [15]).

Peng et al. recently reported mice with deletion of the *Hmox 2* gene, which encodes the enzyme hemeoxygenase 2 (HO-2), exhibit high incidence of OSA[16]. OSA in HO-2 null mice was attributed in part, to the increased loop gain arising from the heightened carotid body chemo reflex[16-19]. While HO-2 produces several bioactive products[20], the enhanced carotid body chemo reflex and subsequent OSA phenotype is attributed to the loss of HO-2 dependent CO production[16]. However, additional factors such as changed activities in the central nervous system may also contribute to the OSA phenotype observed in HO-2 null mice.

Genioglossal nerve activity originating from the hypoglossal nucleus is an important contributor to maintaining upper airway patency. Sporadic loss of neuromuscular control over upper airway muscles is a key contributor to producing obstructive apneas[21,22]. Predisposition to intermittent reductions in airway patency can originate from within the CNS. Tongue muscle activity is regulated by the state-dependent balance between excitation and inhibition received by respiratory hypoglossal motor neurons[23] and direct modulation of intrinsic excitability of hypoglossal neurons[24-26]. It is, however, unknown whether impaired HO-2 signaling in the hypoglossal nucleus influences synaptic and/or intrinsic neuronal properties to alter output from the motor nucleus.

We tested this possibility using a combination of electrophysiological, genetic, and

pharmacological approaches in the rhythmic medullary brainstem slice preparation. We found

that dysregulated HO-2 activity in the hypoglossal nucleus acts through CSE dependent H_2S

signaling to reduce motor neuron excitability. This in turn, diminishes the input-output

relationship between the preBötC and hypoglossal nucleus, and increases the likelihood of

transmission failure between the premotor rhythm and motor nucleus output. These

observations indicate that hypoglossal HO-2 / CO and CSE / H_2S activities interact as important

modulators of hypoglossal output that potentially contributes to changed upper airway tone

when dysregulated.

Methods

Study Approval. In accordance with National Institutes of Health guidelines, all animal

protocols were performed with the approval of the Institute of Animal Care and Use Committee

at The University of Chicago.

Experimental Animals. Experiments were performed using neonatal (P6 to 12) wildtype mice

(C57/BL6; Charles River), HO-2 null mice (from S. H. Snyder; The Johns Hopkins University),

and HO-2:CSE double-null mice. HO-2:CSE double-null mice were created by crossing HO-2

null females with CSE null males (from R. Wang, Department of Biology, Laurentian University,

Sudbury, ON, Canada). Tissues from both sexes were used. No sex-based differences were

observed; therefore, all sexes were pooled for analysis. All litters were housed with their dam in

ALAAC-approved facilities on a 12 hour / 12-hour light-dark cycle.

Pharmacological Agents. Heme oxygenase activity was blocked using bath application of

Chromium (III) Mesoporphyrin IX chloride (ChrMP459, 20µM; Frontiers Sciences, Newark DE).

CORM-3 (20µM; Sigma-Aldrich St. Louis MO), a CO-donor, was bath applied following

ChrMP459 application. NaHS (10µM to 100 µM; Sigma-Aldrich), a H_2S donor, was bath

applied. In all patch clamp experiments, fast synaptic glycinergic and GABAergic inhibition was

blocked by bath application of strychnine (1µM; Sigma-Aldrich), and picrotoxin (50µM; Sigma-

Aldrich), respectively.   Inhibition of CSE production was accomplished by *in vivo* L-

propargylglycine (L-PAG, 30mg/kg (Sigma-Aldrich) administered (*i.p.* injection) 2.5 to 3 hrs prior

to preparation of the rhythmic brainstem slice preparation. Inhibition of potassium channels

SK_{Ca} and ATP-sensitive potassium channel (K_{ATP}) was via bath application of APAMIN (200 µM;

Sigma-Aldrich) and Tolbutamide (100 µM; Sigma-Aldrich) respectively.

Measurement of H_2S Production. Coronal brainstem sections (300µm thick) were cut with a

cryostat at –20°C. The hypoglossal nucleus and control brainstem regions were excised with a

chilled micro-punch needle. Hypoglossal tissue from a single brainstem was not sufficient for

effectively measuring H_2S levels; therefore, we pooled micro punched tissue from two mice for

each sample where H_2S levels measured. H_2S levels were determined as described

previously[27]. Briefly, cell homogenates were prepared in 100 mM potassium phosphate buffer

(pH 7.4). The enzyme reaction was carried out in sealed tubes. The assay mixture in a total

volume of 500µL contained (in final concentration): 100 mM potassium phosphate buffer (pH

7.4), 800µM L-cysteine, 80µM pyridoxal 5′-phosphate with or without L-PAG (20µM) and cell

homogenate (20µg of protein), was incubated at 37°C for 1 h. At the end of the reaction,

alkaline zinc acetate (1% mass / volume; 250µL) and trichloroacetic acid (10% vol/vol) were

sequentially added to trap H_2S and stop the reaction, respectively. The zinc sulfide formed was

reacted with acidic N,N-dimethyl-p-phenylenediamine sulfate (20µM) and ferric chloride (30µM)

and the absorbance was measured at 670 nm using Shimadzu UV-VIS Spectrophotometer. L-

PAG inhibitable H_2S concentration was calculated from a standard curve and values are

expressed as nanomoles of H_2S formed per hour per mg of protein.

Immunohistochemistry. Anaesthetized   mice   (urethane, $1.2g \cdot kg^{-1}$ *i.p.*)   were   perfused

transcardially with heparinized phosphate-buffered saline (PBS) for 20 min followed by 4%

paraformaldehyde in PBS. Brainstems were harvested, post-fixed in 4% paraformaldehyde

overnight, and cryoprotected in 30% sucrose/PBS at 4°C. Frozen tissues were serially

sectioned at a thickness of 20µm (coronal section) and stored at –80°C. Sections were treated

with 20% normal goat serum, 0.1% bovine serum albumin and 0.1% Triton X-100 in PBS for 30

min and incubated with primary antibodies against choline acetyltransferase (ChAT, 1:100;

Millipore; #AB144P), HO-2 (1:200, Novus Biologicals; # NBP1-52849) and CSE (1:250; gift

from Dr. Schenider) followed by Texas Red-conjugated goat anti-mouse IgG or FITC-

conjugated goat anti-rabbit IgG (1:250; Molecular Probes). After rinsing with PBS, sections were

mounted in Vecta shield containing DAPI (Vector Labs) and analyzed using a fluorescent

microscope (Eclipse E600; Nikon).

Brainstem Slice for Electrophysiology. The isolated rhythmic brainstem slice was prepared

as previously described [28]. Briefly, animals were euthanized by decapitation. Brainstems were

rapidly dissected, isolated, and placed into ice cold artificial cerebral spinal fluid (aCSF)

(composition in mM: 118 NaCl, 25 NaHCO$_3$, 1 NaH$_2$PO$_4$, 1 MgCl$_2$, 3 KCl, 30 Glucose, 1.5 CaCl$_2$,

pH=7.4) equilibrated with 95% O$_2$, 5% CO$_2$. The isolated brainstem was glued to an agar block

(dorsal face to agar) with the rostral face up and submerged in aCSF equilibrated with

carbogen. Serial cuts were made through the brainstem until the appearance of anatomical

landmarks such as the narrowing of the fourth ventricle and the hypoglossal axons. The

preBötC and XIIn was retained in a single transverse brainstem slice (thickness: 560 ± 40µm).

The slice was retained and transferred into the recording chamber (~6mL volume) where it was

continuously superfused (12 -15mL/min) with recirculating aCSF. Prior to recording,

extracellular KCl was raised to 8mM and the spontaneous rhythm was allowed to stabilize prior

to the start of every experiment.

Electrophysiology. Extracellular population activity was recorded with glass suction pipettes

filled with aCSF. Pipettes were positioned over the ventral respiratory column containing the

preBötC and over the medial dorsal column containing the hypoglossal nucleus. In some

experiments, a third pipette was position between the preBötC and hypoglossal nucleus just

lateral to the axon tract to record transmission through the premotor field containing

intermediate premotor inspiratory neurons [29,30]. Extracellular population activity was recorded

with glass suction pipettes filled with aCSF [31]. The recorded signal was sampled at 5kHz,

amplified 10,000X, with a lowpass filter of 10 kHz using an A-M instruments (A-M Systems,

Sequim, WA, USA) extracellular amplifier. The signal was then rectified and integrated using

Clampfit electronic filter. Recordings were stored on a computer for *posthoc* analysis.

All recordings intracellular recordings were made using the Multiclamp 700B (Molecular

Devices: https://www.moleculardevices.com/systems/conventional-patch-clamp/multiclamp-

700b-microelectrode-amplifier). Acquisition and post hoc analyses were performed using the

Axon pCLAMP10 software suite (Molecular Devices: https:

www.moleculardevices.com/system/axon-conventional-patch-clamp/pclamp-11-software-suite).

Whole cell patch clamp recordings of hypoglossal motor neurons were obtained using the blind-

patch technique with a sample frequency of 40 kHz. Recordings were made with unpolished

patch electrodes, pulled from borosilicated glass pipettes with a capillary filament [31]. The

electrodes had a resistance of 3.5-8 MΩ when filled with the whole cell patch clamp pipette

solution containing (in mM): 140 K-gluc acid, 1 $CaCl_2$, 10 EGTA, 2 $MgCl_2$, 4 Na_2-ATP, 10

HEPES. Patch clamp experiments were performed with a patch clamp amplifier (Multiclamp

700B, Molecular Devices, Sunnyvale, CA, USA), and the software program pCLAMP 10.0

(Molecular Devices). Neurons located at least two to three cell layers (about 75-250µm) rostral

from the caudal surface of the slice were recorded. The liquid junction potential was calculated

to be -12mV and was subtracted from the membrane potential. The series resistance was

compensated and corrected throughout each experiment. In voltage clamp experiments,

membrane potential was held at -60mV. Current clamp experiments used a holding potential

between 0 and -100pA to establish the baseline resting membrane potential between -55 and -

70mV. In some cases, we determined rheobases using ramp protocol in our current clamp

recordings. This ramp protocol consisted of a hyperpolarizing step (-100pA) succeeded by the

injection of a ramping depolarizing current (122pA/sec; peak current 600pA).

Statistical Analyses. Unless otherwise explicitly stated elsewhere, each n value represents an

individual animal. Transmission was expressed as a percentage of the hypoglossal network

bursts corresponding to the total network bursts from either the preBötC or the premotor field.

Burst were considered corresponding if initial start time of bursts were within 500-750ms of each

other (corresponding time was maximized until only one hypoglossal burst per preBötC was

detected). Mean I/O and transmission values for each slice were calculated using a 120 s

window. In pharmacological experiments, this window was taken at the end of each baseline or

pharmacological agent phase (each phase duration = 600 s). The input-output (I/O) ratio for

each inspiratory event (defined by network a burst in preBötC) was calculated as previously

described in [31]. I/O ratios values for preBötC bursts without a corresponding XIIn burst were

designated as 0. To illustrate the cycle-to-cycle input-output relationships between networks,

heat maps of individual I/O ratio values. As the rhythmic frequency across preparations varied,

the number of events in the 120 s analysis window also varied; therefore, heat maps plotted the

greater of either the total number of cycles or 25 consecutive cycles from in a given analysis

window.

Statistics were performed using Origin 8 Pro (OriginLab, RRID:SCR_014212) or Prism 6

(GraphPad Software; RRID:SCR_015807). In cases where the distribution of data appeared

normal, comparisons between two groups were conducted using either paired or unpaired two-

tailed t-tests with Welch's correction, as appropriate. In cases, where the distribution of

individual data points did not appear normal, the Wilcoxon match-paired signed rank test was

performed.   A one-way ANOVA was performed followed by *posthoc* Dunnett's test comparing

experimental groups to control when a comparison of three or more groups. In plots where the

mean ± S.E.M. are presented, the mean and S.E.M. are overlaid on the individual results from

the corresponding dataset. Differences were defined when the P-value was less than 0.05.

Results

Hypoglossal neurons express hemeoxygenase-2 (HO-2). We first assessed whether

hypoglossal neurons express HO-2. Hypoglossal neurons showed positive

immunohistochemical expression for HO-2 as indicated by co-localization of HO-2 with Choline

acetyl transferase (ChAT), an established marker of these neurons [11] (**Fig.1A**, n=3).

Disrupting HO-2 function impairs hypoglossal inspiratory activity. Extracellular field

recordings in the rhythmic brainstem slice were simultaneously recorded from the preBötC and

the corresponding motor output from the hypoglossal nucleus. Two approaches were employed

to assess the role of HO-2: (1) using Cr(III) Mesoporphyrin IX chloride (ChrMP459, 20µM), an

inhibitor of HO [32]; and (2) using brain slices from HO-2 null mice.

Representative extracellular field recordings from the preBötC and hypoglossal nucleus

prior and during ChrMP459 exposure were shown in **Fig 1B** (n=11). While ChrMP459

suppressed hypoglossal burst amplitude (**Fig 1C**, *left*; n=11; baseline: 99.57 ± 0.60%, ChrM459:

81.80 ± 10.80%, P=0.007), the HO inhibitor had no effect on preBötC burst amplitude (**Fig 1C**,

right, n=11; Baseline: 99.93 ± 0.34%, ChrMP459: 93.23 ± 4.80%, P=0.21)**.** ChrMP459

consistently reduced the cycle-to-cycle input-output relationship between preBötC and the motor

nucleus as revealed by examining the cycle-to-cycle I/O across preparations (**Fig 1D**, baseline:

1.00 ± 0.07 vs. ChrMP459: 0.55 ± 0.11; P=.006). In the extreme, altered input-output

relationships between preBötC and the hypoglossal nucleus may increase the propensity for

transmission failures as determined by the inability of preBötC activity to produce hypoglossal

output at the network level[31]. Indeed, the reduced I/O ratio was associated with an increase in

failed transmissions of the preBötC activity to output from the hypoglossal nucleus (**Fig 1E**,

baseline: 90.85 ± 5.80% vs. ChrMP459: 80.21 ± 7.40%, P=0.039).   Together these findings

suggest that HO inhibition causes a generalized weakening in the activity relationship between

the rhythm generating network and hypoglossal motoneurons.

ChrMP459 is a pan HO inhibitor, which cannot distinguish activities between heme

oxygenase isoforms. To assess the specific contribution of HO-2, we compared rhythmic

activities in brain slices from wild type (n=8) and HO-2 null (n=7) mice (**Fig 2A**). Larger cycle-to-

cycle I/O ratios were observed in wild type slices as compared to HO-2 null slices (**Fig 2B**: wild

type: 1.04 ± 0.04 vs. HO-2 null: 0.72 ± 0.07, P=0.0019)**.** Similarly, transmission of preBötC

activity to the hypoglossal nucleus was greater in wild type than in HO-2 (**Fig 2 C,** wild

type: 97.07 ± 0.02% vs. HO-2 null: 70.15 ± 0.03%, P<0.0001). These findings established that

genetic elimination of HO-2 produces a similar phenomenon to that of pan HO inhibition

indicating that loss of HO-2 activity alone is sufficient for impairing transmission from preBötC to

the hypoglossal nucleus. Given these similarities and the limited availability of HO-2 null mice,

several of the following studies were performed using the ChrMP459 in rhythmic wild type

brainstem slices.

HO inhibitor does not affect premotor neuron activity. Intermediary premotor neurons relay

drive from the preBötC to the hypoglossal nucleus [29,30]. Therefore, it was possible that HO

inhibition impaired transmission of drive from preBötC by perturbing activity from intermediary

premotor neurons. To address this possibility, triple extracellular recordings (n=5) were made

from the preBötC, the field of the ipsilateral premotor neurons, and the hypoglossal nucleus.

Baseline transmission from preBötC to the premotor field and to the hypoglossal nucleus was

reliable and consistent (**Fig. 3A** *middle panel*). However, ChrMP459 disrupted activity in the

hypoglossal nucleus despite unaltered activities in either preBötC or the intermediate premotor

field (**Fig 3A**, *right panel*). Indeed, while neither transmission failures nor the cycle-to-cycle I/O

ratio from preBötC to the premotor field was affected by ChrMP459 (**Fig 3B**: *left,* I/O: baseline:

1.16 ± 0.09 vs ChrMP 1.16 ± 0.14, P=0.816; *right,* Transmission: baseline: 100.0 ± 0.0% vs

ChrMP 86.35 ± 11.81%, P=0.312), the HO inhibitor reduced the transmission of activity and the

cycle-to-cycle I/O ratio between the premotor field and the hypoglossal nucleus (**Fig 3C**: *left,*

I/O: baseline: 1.12±.17 vs ChrMP 0.40±0.09, P=0.04; *right,* Transmission: baseline 89.57±1.3%

vs ChrMP 57.62±13.69%, P=0.04). These results suggested that dysregulated HO-2 modulates

is a postsynaptic phenomenon in the hypoglossal nucleus.

HO inhibition suppresses inspiratory drive currents and reduces excitability in

hypoglossal neurons. To examine the effect of HO inhibition on postsynaptic activity of the

hypoglossal neurons, we performed patch clamp recordings from a total of 27 wild type

hypoglossal neurons exposed to ChrMP459. These hypoglossal neurons were disinhibited from

fast inhibition (50µM picrotoxin and 1µM strychnine) allowing us to focus on inspiratory-related

fast glutamatergic drive. 19 of the 27 hypoglossal neurons received excitatory synaptic drive in-

phase with the preBötC (i.e., inspiratory hypoglossal neurons). Peak inspiratory drive currents

were reduced in ChrMP459 (n=19, **Fig 4A**, baseline: -142.9 ± 22.82 pA vs. ChrMP459: -95.31 ±

21.79 pA, P=0.004). Reduced drive coincided with hypoglossal neurons generating fewer

action potentials per preBötC burst in ChrMP459 (**Fig 4B**, n=17 baseline: 14.68 ±2.23 action

potentials per burst vs. ChrMP459: 6.79± 1.54 action potentials per burst, P<0.0001).

Furthermore, as determined by the injection of a depolarizing ramp current into hypoglossal

neurons, the HO inhibitor increased rheobase among inspiratory hypoglossal neurons (**Fig 4C,**

n=19**,** baseline: 167.5 ± 35.83 pA vs. ChrMP459: 338.0 ± 82.50 pA; P=0.008) yet decreased

rheobase in non-inspiratory hypoglossal neurons (i.e., neurons not receiving drive during

preBötC activity) (n=8, **Supplemental Fig 1**, baseline: 280.5±56.43 pA vs 228.2±47.96pA,

P=0.017).

Elevated H_2S abundance is observed in the hypoglossal nucleus of HO-2 null mice. We

next sought to determine the mechanism(s) by which inhibition of HO-2 affect hypoglossal

neuron activity. Earlier studies[33,34] have reported that HO-2 is a negative regulator of CSE-

dependent H2S production. To test this possibility, we first examined whether the hypoglossal

neurons express CSE. Brainstem sections from the wild type hypoglossal nucleus revealed

CSE expression in ChAT-positive neurons (**Fig 5A,** n=3). We then determined H_2S abundance

in hypoglossal tissue punches from wild type and HO-2 null mice. Relative to the wild type

hypoglossal nucleus (**Fig 5B** *blue*; n=6, 60.58 ± 6.37 $nmol \bullet mg^{-1} \bullet h^{-1}$), H_2S abundance was

greater in the hypoglossal nucleus of HO-2 null mice (**Fig 5B** *red*; n=6,

144.12 ± 8.29 $nmol \bullet mg^{-1} \bullet h^{-1}$), but not different from the inferior olive brainstem region of HO-2

null mice (**Fig 5B** *grey*; n=4, 56.10 ± 2.88 $nmol \bullet mg^{-1} \bullet h^{-1}$). These findings demonstrated that

the hypoglossal nucleus expresses CSE and HO-2 negatively regulates H_2S production in the

hypoglossal nucleus.

H_2S mediates impaired transmission of inspiratory drive caused by disrupted HO-2

function.

If the impaired transmission of inspiratory drive to the hypoglossal nucleus by disrupted HO-2

function involves CSE-derived H_2S then: 1) a H_2S donor should mimic the effects of disrupted

HO-2 activity; 2) CO administration should improve the input-output relationship in respiratory

slices from HO-2 null mice and ChrM459 application; and 3) CSE blockade should restore the

transmission from preBötC to the hypoglossal nucleus. The following experiments tested these

possibilities.

Wild type brainstem slices exhibited a nearly 1:1 ratio of transmission of inspiratory activity from

preBötC to hypoglossal neurons (**Fig 5C**, *left*). Application of NaHS, a H_2S donor reduced

transmission from preBötC to hypoglossal (**Fig. 5C,** *middle, right*) in a dose-dependent manner

(**Fig. 5D**; 0μM NaHS: n=9, 100 ± 0.73%; 10μM NaHS: n=5**,** 90.14 ± 6.08%; 50μM NaHS: n=6,

84.0 ± 3.91%; 100μM NaHS: n=9, 81.2 ± 5.83%), which coincided with a reduction in I/O ratio

by NaHS (**Fig. 5E**; 0 μM NaHS: 1.055 ± 0.028; 10μM NaHS: 0.85 ± 0.09; 50μM NaHS: 0.82 ±

0.08; 100μM NaHS: 0.84 ± 0.07). These findings demonstrated increased H_2S abundance

reduces hypoglossal neuronal activity consistent with findings using either ChrM459 or HO-2

null mice. Thus, the stability of inspiratory transmission from the preBötC to hypoglossal nucleus

appears to be negatively affected either by increasing H_2S abundance or by disrupting HO-2

function.

CO produced by HO-2 is known to inhibit CSE-dependent H_2S production by HO-2 [33,34].

Therefore, we sought to assess how CORM-3 (20µM), a pharmacological CO donor, impacted

activity in ChrMP459-treated wild type rhythmic slices (n= 3, **Fig 6A**) and rhythmic slices from

HO-2 null mice (n=4). Dysregulated HO-2 activity, caused by either pharmacological

(ChrMP459) or genetic (HO-2 null mice) manipulation, is improved by CORM-3 as indicated by

larger I/O ratios (**Fig 6B**, dysregulated HO-2: 0.73 ± 0.04 vs. CORM-3: 1.07 ± 0.75, P=0.004)

and improved transmission (**Fig 6C**, dysregulated HO-2: 79.88 ± 4.69% vs. CORM-3:

93.39 ± 2.93%, P=0.03). As these findings suggested the absence of HO-2 dependent CO

production is a key factor driving transmission failure in the rhythmic slice preparation, we next

determined the involvement of CSE.

Inspiratory activity in the brainstem slice from HO-2:CSE double null mice appeared to be stable

and consistent (**Fig 6D**). Quantification of simultaneous extracellular field recordings of preBötC

activity and hypoglossal nucleus revealed a larger I/O ratio (**Fig 6E,** HO-2:CSE: 1.04 ± 0.02,

n=6; P=0.003) and near absence of transmission failures (**Fig 6F**, HO-2:CSE: 91.6 ± 0.02%;

P=0.0007) when compared to activity in HO-2 null slices.

Given the observations using HO-2:CSE null mice, we next sought to determine whether

acute blockade of CSE could restore transmission relationships between preBötC and XIIn in

the HO-2 null slice. *In vivo* L-PAG treatment improved transmission of preBötC activity to the

hypoglossal nucleus in the rhythm slice (**Fig. 7A,** n=6) as indicated by larger cycle-to-cycle I/O

ratios (**Fig. 7B,** L-PAG = 1.01± 0.03, P=0.008) and greater transmission rates (**Fig. 7C,** L-PAG

96.31 ± 2.62%, P<0.0001) when compared to the respective metrics from untreated HO-2 null

mice. Intermittent transmission failure was also evident in patch clamp recordings from

untreated HO-2 null hypoglossal neurons (**Fig 7D**, *left* shaded cycles) but not in HO-2 null

hypoglossal neurons treated with L-PAG (**Fig. 7D**, *right*). These reduced transmission events

correlated with smaller individual inspiratory drive currents in HO-2 null hypoglossal neurons

when compared to corresponding inspiratory drive currents from L-PAG treated HO-2 mice (**Fig.**

7D-E, HO-2: -36.71 ± 2.14 pA vs. L-PAG -194.3 ± 82.73 pA, P=0.0007).   Together, these

experiments implicated the involvement of CSE-dependent H2S signaling with the effects of

disrupted HO-2 signaling in hypoglossal neurons.

Blockade of small conductance calcium-activated potassium channel (SK$_{Ca}$) activity

restores changes induced by HO-dysregulation in hypoglossal activity. As our

experiments implicated the involvement of H$_2$S signaling, we next sought to determine how H$_2$S

sensitive ion channels may contribute to impairing hypoglossal neuron activity caused by

HO-dysregulation.   H$_2$S has been shown to enhance activity of several different potassium

channels, including SK$_{Ca}$ and ATP-sensitive potassium channel (K$_{ATP}$) activities [35]. As both SK$_{Ca}$

and K$_{ATP}$ are important in the regulation of hypoglossal neuron excitability [36,37], we examined

how blocking these channels affected hypoglossal activity during HO-dysregulation. At the

network level, administration of the selective SK$_{Ca}$ inhibitor, apamin (200µM), increased the

excitability of the hypoglossal neurons in ChrMP459. This enhanced activity exceeded the

original baseline activity (i.e., prior to ChrMP459 administration) causing ectopic bursting in the

hypoglossal nucleus (**Supplemental Figure 2A**) that made analysis of population transmission

and I/O ratios unreliable. Therefore, we proceeded to resolve the effects of apamin on the

influence of ChrMP459 at the level of individual hypoglossal inspiratory motor neurons. While in

some hypoglossal neurons exposed to ChrMP459 apamin substantially increased drive currents

(>100pA; **FIG 8A1**), in others, apamin modestly increased the drive current (<100pA; **FIG 8A2**).

Despite this variability, apamin increased inspiratory drive currents received by ChrMP459

treated hypoglossal neurons (**FIG 8A3,** n= 6, ChrMP459: -85.77± 38.54 pA vs. apamin: -

219.97± 97.76 pA, P=0.031). Apamin also enhanced the number of action potential generated

per preBötC burst during ChrMP459 (**FIG 8B,** n=8**,** ChrMP-459: 12.57 ± 3.7 action potential per

burst vs. apamin 26.05 ± 6.87 action potential per burst, P=0.016). This was consistent with the

ability of apamin to reduce rheobase in ChrMP459 treated inspiratory hypoglossal neurons (**FIG**

8B; n=7, ChrMP459: 532.0 ± 186.5 pA vs. Apamin: 307.09 ± 80.9 pA, P=0.016).

356   To determine how blockade of K_{ATP} impacted hypoglossal activity during ChrMP459, we

used the K_{ATP} channel blocker, tolbutamide (100μM). Tolbutamide did not induce ectopic

bursting in the hypoglossal nucleus during ChrMP459 (**Supplemental Fig 2B**, n=5).

Furthermore, tolbutamide (100μM), did not improve the rate of transmission of preBötC activity

(**Supplemental Fig 2C**, ChrMP459: 69.24±6.0% tolbutamide: 71.23 ± 6.8%, P=0.83) but did

increase the I/O ratio (**Supplemental Fig 2C right**, ChrMP459: 0.68 ± 0.05 tolbutamide:

0.87 ± 0.05, P=0.037). Tolbutamide neither enhanced inspiratory drive currents in ChrMP459

(**Supplemental Fig 2D**, n=4, ChrMP459: -70.51 ± 27.49 pA vs. tolbutamide: -83.06 ± 36.29 pA,

P=0.37) nor increased the number of action potentials per preBötC burst in ChrMP459 (n=4,

ChrMP459: 7.063±2.08 action potential per burst vs. tolbutamide: 9.37±3.18 action potential

per burst, P=0.125). Moreover, tolbutamide did not affect rheobase of inspiratory hypoglossal

neurons treated with ChrMP459 (**Supplemental Fig 2D**, n=4; ChrMP459: 221.01 ± 74.8 pA vs.

tolbutamide: 180.4 ± 68.63 pA, P=0.218). Thus, these results suggested that apamin could

enhance activity of hypoglossal neurons during HO-inhibition; whereas, the efficacy of

tolbutamide to impact activity during HO-inhibition was limited.

Discussion

Our study reveals a previously uncharacterized neuromodulatory interaction between HO-2 and CSE-derived H_2S regulating activity from the hypoglossal nucleus. We documented that both HO-2 and CSE are expressed among ChAT-positive cells of the hypoglossal nucleus. Although we did not resolve expression among hypoglossal motor neurons innervating upper airway muscles, such as the genioglossus, our electrophysiological studies revealed that dysregulated HO-2 activity impacted intrinsic properties of hypoglossal neurons receiving drive from the preBötC. HO-2 dysregulation reduced impaired transmission originating from the rhythm generating preBötC and the motor nucleus, as evidenced by the cycle-to-cycle reductions in the input-output and the intermittent failures for preBötC activity to generate corresponding motor pool output. The increased propensity of transmission failure in response to dysregulated HO-2 activity is reminiscent of the OSA phenotype observed in HO-2 null mice[16] and demonstrates the potential importance of centrally derived interactions between HO-2 and CSE activity in regulating motor neuron output important for upper airway patency.

Both a pharmacological inhibitor of HO and genetic elimination of Hmox-2 impaired transmission of neural drive from preBötC to the hypoglossal nucleus. Mice treated with intermittent hypoxia (IH) patterned after blood O_2 profiles seen during OSA also show failed transmission related to a changed input-output relationship between preBötC and hypoglossal nucleus[31,38]. While IH-induced phenomena are associated with alterations in preBötC activity, the HO inhibitor neither impacted inspiratory rhythmogenesis from the preBötC (**Fig. 1C**) nor perturbed transmission of neuronal drive to intermediate premotor neurons suggesting that changed interneuron neurophysiology was not a major contributing factor to the impaired transmission in the hypoglossal nucleus.

At the level of the hypoglossal nucleus, ChrMP459 had divergent effects on non-inspiratory and inspiratory hypoglossal neurons. HO inhibition increased rheobase among non-inspiratory neurons; whereas, in inspiratory hypoglossal neurons, it decreased the magnitude of drive

currents, increased rheobase, and diminished the number of action potentials generated during

preBötC bursting. Thus, while our studies did not resolve how hypoglossal neurons projecting to

different tongue muscles are impacted by ChrMP459, our findings indicate that HO-

dysregulation can differentially impact hypoglossal neurons that receive drive from the preBötC.

The observed postsynaptic phenomena among hypoglossal motor neurons receiving inspiratory

drive could contribute to the occurrence of obstructive apneas by increasing the probability for

intermittent filtering of upper airway tone from inspiratory drive by hypoglossal motor neurons.

Even if inspiratory activity is successfully produced at in other inspiratory muscles, such as the

diaphragm. Indeed, this phenotype has been documented in HO-2 null mice[16]. However, further

resolution is needed to understand how dysregulation of HO-2 may impact hypoglossal neurons

innervating different muscle groups and to determine extent of the contribution from central HO-

2 dysregulation to loss of upper airway tone and airway collapse *in vivo*.

In HO-2 null mouse, the incidence of OSA is absent with co-inhibition of CSE[17], which is

consistent with reports that CO generated by HO-2 inhibits CSE-dependent H_2S production[33,34].

After documenting CSE expression in hypoglossal neurons and demonstrating an increased

abundance of H_2S in the hypoglossal nucleus of HO-2 null mice, our electrophysiological studies

demonstrated the ability for a CO donor to improved transmission and the input-output

relationship in HO-2 null slices. Furthermore, using a H_2S donor also increased transmission

failures and reduced the I/O ratio similar to perturbing HO-2 activity. Endogenous H_2S activity

could originate from other H_2S producing enzymes, such as cystathionone β-synthase (CBS)

that is expressed primarily in astrocytes throughout the CNS[39]. However, CBS inhibition

appears to have limited impact on inspiratory activity from the hypoglossal nucleus[40]. In

contrast, our genetic and pharmacological manipulations to ablate/diminish CSE activity

improved transmission and the input-output relationship between inspiratory drive and the

hypoglossal nucleus in HO-2 null mice. Together these findings implicated the interaction

between HO-2 / CO and CSE / H_2S within the hypoglossal nucleus to regulate its output.

How might enhanced H_2S signaling reduce drive currents, and excitability of hypoglossal

neurons? While it is possible that H_2S may impact presynaptic release of glutamate and/or

postsynaptic receptor activity of hypoglossal neurons, the ChrMP459 mediated increase in

rheobase among inspiratory hypoglossal neurons implicated involvement of a non-synaptic

conductance(s) downstream of H_2S based signaling caused by perturbations in HO-2 activity.

The H_2S can enhance both K_{ATP} and SK_{Ca} activities[35]. In the hypoglossal neurons, K_{ATP} is

dynamically active causing periodic adjustment of neuronal excitability[36] while SK_{Ca} also

regulates excitability and firing properties of hypoglossal neurons[37]. In ChrMP459, tolbutamide

had a limited effect normalizing transmission as it improved the I/O ratio between preBötC and

the hypoglossal motor nucleus but did not reduce transmission failure. At the neuronal level,

tolbutamide did not increase drive currents nor enhance intrinsic excitability of hypoglossal

neurons in ChrMP459. These results indicated that blockade of K_{ATP} during HO-2 dysregulation

was limited in countering the effects on HO-2 dysregulation in the hypoglossal nucleus. On the

other hand, apamin normalized drive currents and increased excitability of inspiratory

hypoglossal neurons in ChrMP459 demonstrating that blockade of SK_{Ca} sufficiently mitigates

many aspects of HO-2 dysregulation in hypoglossal neurons that receive drive from the

preBötC.

In conclusion, our study provides proof-of-concept for the existence of a central

mechanism by which loss of HO-2 leads to reduced upper airway tone by enhancing H_2S

activity in the hypoglossal nucleus. This mechanism appears to involve antagonistic interactions

between HO-2 and CSE activities to regulate excitability of hypoglossal neurons and is localized

in the neurons themselves. Although OSA in HO-2 null mice has attributed to increased "loop-

gain" arising from the hypersensitive carotid body reflex[16], our findings indicate the potential

involvement of a disrupted interaction between HO-2 / CO and CSE / H_2S in hypoglossal motor

neurons that contribute could to the sporadic loss of upper airway tone observed in OSA.

Figure 1. Disruption of hemeoxygenase-2 (HO-2) in hypoglossal motor neurons impairs inspiratory activity from the hypoglossal nucleus but not preBötC. **A.** HO-2 (red bottom left) is expressed in ChAT$^+$ cells (green bottom right) of the hypoglossal nucleus (XIIn, combined top). **B-E.** Population recordings of rhythmic brain slices were recorded from ipsilateral preBotC and XIIn simultaneously, analyses were performed on 2-minute windows of baseline and 10-minutes into bath application of 20µM ChRMP459 (ChRMP). **B.** Integrated traces of network activity in spontaneously rhythmic brainstem slices (n=11) recorded from XIIn (grey) and preBötC (black) before (top) and after (bottom) bath application, failed transmission events are highlighted (pink boxes); scale bar: 5 sec. **C.** Percentage change of integrated burst amplitude from Baseline to ChRMP in XIIn (top) and preBötC (bottom). Thin pink and blue lines illustrate integrated burst amplitude from individual slices. Symbols connected by a thick line illustrate mean integrated burst amplitude. **D.** Heat maps of cycle-to-cycle I/O ratios analysis window in Baseline and in ChRMP. Grey boxes indicate non-events in recordings from slower rhythms where less than 25 events occurred during the analysis window. **E.** Comparison of inspiratory drive transmission in XIIn between Baseline and ChRMP. Thin grey lines illustrate transmission from individual slices. Symbols connected by a thick line illustrate mean transmission.

Figure 2. Genetic deletion of HO-2 reduces the I/O relationship between preBötC and the hypoglossal nucleus and uncouples of motor output from inspiratory rhythmogenesis. A. Representative integrated traces of network burst activity in preBötC and XIIn in rhythmic slices from wildtype (WT) (left, n=8) and HO-2 null (HO-2$^{-/-}$; right, n=7) mice; scale bar: 2 sec. **B.** Heat maps of cycle-to-cycle I/O ratios in WT (left) and HO-2$^{-/-}$ (right) rhythms during the analysis window. Grey boxes indicate non-events in recordings from slower rhythms where less than 25 events occurred during the analysis window. **C.** Transmission of inspiratory activity from

preBötC to XIIn in slices from WT vs HO-2$^{-/-}$. Symbols illustrate transmission from individual

slices.

Figure 3. While ChrMP459 does not change transmission from the preBötC to the

premotor area, ChrMP459 increases transmission failure from the premotor area to the

hypoglossal nucleus. A. Diagram of medullary brain slice illustrating relative electrode

placement for simultaneous triple extracellular recordings (n=5) from the XIIn (light grey, 1),

premotor field (dark grey, 2), and preBötC (black, 3). Corresponding representative traces of

integrated network activity in Baseline (left) and in 20µM ChrMP (right); scale bar: 1 sec. **B.** The

cycle to cycle I/O ratio (left) and transmission (right) between preBötC and the premotor field. **C.**

The cycle to cycle I/O ratio (left) and transmission (right) between the premotor field and XIIn.

Figure 4. Heme oxygenase inhibition reduces inspiratory drive currents in hypoglossal

neurons. Whole cell patch clamp recordings were made from hypoglossal neurons in rhythmic

brain slices while simultaneously recording ipsilateral preBötC activity in Baseline and in ChrMP.

Neurons were disinhibited from fast synaptic inhibition using 50µM PTX and 1µM Strychnine.

A. (left) Representative voltage clamp recordings from a XIIn neuron ($V_{holding}$= -60mV) aligned

with corresponding integrated network activity from preBötC before (Baseline, top, grey) and

after 20 µM ChrMP (bottom, purple). Scale bar: 1 sec x 10pA. (middle) Magnification of

highlighted (red dotted box) drive currents from Baseline (grey) and ChrMP (purple). Scale bar:

100 msec x 10pA. (right) Comparison of XIIn inspiratory drive current magnitude in Baseline and

ChrMP (n=20). Thin grey lines illustrate individual neuron response. Symbols connected by a

thick line illustrate mean drive current.   **B.** (left) Representative current clamp recordings from a

XIIn neuron spontaneously active with the preBötC network rhythm in Baseline (top, grey) and

20 µM ChrMP (bottom, purple); skipped transmission of action potentials in ChrMP are

highlighted (pink). Scale bar 2sec x 20mV. (middle) Magnification of highlighted neuronal activity

(red box in left). Scale bars: 100msec x 25mV. (right) Number of action potentials generated

per inspiratory burst in Baseline and in ChrMP (n=17). Thin grey lines illustrate individual neuron

response. **C.** (left) Representative trace of current clamp recording in response to ramp current

injection during Baseline (grey trace) and in ChrMP (purple trace); scale bar: 500 msec. (right)

Comparison of rheobase in inspiratory XIIn neurons during Baseline and in ChrMP (n=19). Thin

grey lines illustrate individual neuron response.

Figure 5. CSE-dependent H$_2$S is produced in the hypoglossal nucleus and exogenous

NaHS uncouples hypoglossal nucleus activity from the preBötC. A. CSE (red, bottom left)

is expressed in ChAT$^+$ neurons (green, bottom right) in the XIIn (overlay, top). **B.** CSE-

dependent H$_2$S generation in homogenates from WT and HO-2$^{-/-}$. Homogenates were prepared

from tissue punches from the XIIn (red area in slice diagram) and inferior olive (grey area in

slice diagram) at bregma between -7.20mm and -7.90mm. (WT: XIIn n=6; HO-2$^{-/-}$: XIIn n=6,

inferior olive n=4). Note: Each n value reported in B represents a sample consisting of the

anatomical region pooled from two animals. **C.** Integrated traces from XIIn (top) and preBötC

(bottom) during baseline (black), and in response to the H$_2$S donor, NaHS, at 50 µM (dark grey)

and 100 µM (light grey). After NaHS application, XIIn but not preBötC burst amplitude were

diminished (red dashed box) and in some cases, preBötC drive failed to produce activity in the

XIIn (pink boxes). **D.** Comparison of transmission from preBötC to XIIn after NaHS application at

10µM, 50µM and 100µM. **E.** I/O ratios for each NaHS concentration. (Baseline: n=9; 10µM n=5;

50µM n=6; 100µM n=9).

Figure 6. HO-dependent transmission failures can be recovered with CO-donor CORM-3

and are not present in HO-2:CSE null transmission. A. Representative extracellular

recording from preBotC and XIIn in WT slices during Baseline (left), in ChrMP alone (middle),

and in ChrMP + 20uM CORM-3 (CORM3, right). **B.** Heat maps of cycle-to-cycle I/O ratios

during dysregulated HO-2 (n=7: n=4 HO-2 null and n=3 WT-ChrMP) before and after CORM-3

application. **C.** Transmission from preBötC to XIIn from dysregulated HO-2 slices before and

after bath application of CORM-3. **D.** Representative extracellular recordings from preBötC and

XIIn in slices from HO-2:CSE null (HO-2:CSE$^{-/-}$); scale bar 2 sec. **E.** Heat map of cycle-to-cycle

I/O ratio from preBötC to XIIn in HO-2:CSE$^{-/-}$. The I/O ratio from HO-2:CSE$^{-/-}$ is greater than I/O

ratios from HO-2$^{-/-}$ (P=0.003). **F.** Comparison of transmission from preBötC to XIIn in HO-2$^{-/-}$

and HO-2:CSE$^{-/-}$ (n=6). HO-2$^{-/-}$ data used for comparisons in E and F were originally shown in

Figure 2.

Figure 7. Transmission failures in HO-2 null mice are rescued with CSE inhibitor L-

propargylglycine. A. Representative recordings of integrated activity of the preBötC (bottom)

and XIIn (top) in slices from HO-2 null mice treated with L-PAG (30mg/kg; L-PAG, n=5). Scale

bar 2 sec. **B.** Heat map of cycle-to-cycle I/O ratio in rhythmic slices from HO-2 null mice treated

with L-PAG (n=6). **C.** Comparison of transmission between HO-2$^{-/-}$ (replotted from Fig 2) and L-

PAG slices (n=6). **D.** Representative voltage clamp recordings of inspiratory drive currents

received by hypoglossal neurons from HO-2$^{-/-}$ (n=7) and L-PAG (n=6). Neurons were

disinhibited from fast synaptic inhibition using 50μM PTX and 1μM Strychnine. Scale bar 10

msec x 20pA. Skipped transmission between preBötC (bottom) the XIIn neuron (top) occurs in

untreated   HO-2$^{-/-}$ (highlighted by pink boxes) but not in neurons from L-PAG. **D1.**

Magnified representative drive potentials from HO-2$^{-/-}$ and L-PAG (from highlighted by red-

outlined boxes in **D**). scale bars: 100 msec x 10pA. **E.** Quantification of drive currents received

by XIIn motoneurons from HO-2$^{-/-}$ mice (n=8) produce smaller drive potentials when compared

to L-PAG (n=6).

Figure 8. Apamin improves network input-output relationship between hypoglossal

nucleus and reverses changes to intrinsic and synaptic excitability caused by HO

dysregulation. A1. Representative inspiratory drive current received by hypoglossal neurons in

ChrMP (purple) and in Apamin (200 μM, gold) where apamin increased the drive current >100

pA. Scale bars: 1 s x 50 pA. **A2.** Representative inspiratory drive current hypoglossal neuron in

ChrMP (purple) and in Apamin (gold) where apamin increased the drive current by less than

<100 pA Scale bars: 1 s x 50 pA. **A3.** Comparison of inspiratory drive currents from

hypoglossal neurons exposed to ChrMP (200 μM) then treated with Apamin (n=6). The effect

of ChrMP on baseline disinhibited drive current for each of these neurons were reported in Fig

4A. **B**. (left) Representative trace of bursting of a hypoglossal neuron resulting from preBötC

drive during ChrMP (purple) and in Apamin (gold) (n=8). Scale bars: 20 mV x 500 msec. (right)

Comparison of action potentials generated per preBötC burst during ChrMP and Apamin (n=8).

The effect of ChrMP on baseline action potential generated per preBötC burst for each of these

neurons were reported in Fig 4B. **C.** (left) Representative traces of current clamp recordings in

response to ramp current injection in ChrMP (purple) and in Apamin (gold). Scale bar: 500msec

x 10mV). (right) Rheobase comparison from inspiratory XIIn neurons during ChrMP and Apamin

(n=7). The effect of ChrMP on baseline rheobase for each of these neurons were reported in

Fig 4C.

Supplemental Figure 1: ChrMP459 decreases rheobase of non-inspiratory hypoglossal

neurons. Comparison of rheobase from non-inspiratory hypoglossal neurons in baseline and

ChrMP (n=8). Non-inspiratory hypoglossal neurons were defined as neurons that did not

receive synaptic drive in-phase with a preBötC burst. Neurons were disinhibited from fast

synaptic inhibition using 50μM PTX and 1μM Strychnine.

Supplemental Figure 2: Apamin produce ectopic hypoglossal network activity during

ChrMP459 and tolbutamide has limited effects on network and inspiratory hypoglossal

neurons activity during ChrMP459. A. Representative traces of integrated network activity of

the preBötC (top) and XIIn (bottom) in slices from WT mice   Baseline, ChrMP, and in 200 μM

Apamin. Asterisks (*) indicate detected hypoglossal output in phase with the preBötC. Green

box indicates ectopic network bursting. Scale bar 10 sec. Due to ectopic bursting within and

around rhythmic burst transmissions, analysis of apamin at the network level could not be

accurately detected. **B.** Representative traces of integrated network activity of the preBötC (top)

and XIIn (bottom) in slices from WT mice Baseline, ChrMP, and in tolbutamide (200 μM, Tolb).

Asterisks (*) indicate detected hypoglossal output in phase with the preBötC. Pink box indicates

cycles where preBötC drive failed to produce activity in the XIIn. **C.** Comparisons of

Transmission (left) and I/O ratio (right) in ChrMP459 and in Tolbutamide (n=5). **D.** (top)

Representative traces of inspiratory drive currents in ChrMP (purple) and in Tolb (green). Scale

bar 1 sec x 50 pA. (bottom) Comparison disinhibited inspiratory drive currents in ChrMP and

Tolb (n=4). The effect of ChrMP on baseline drive current for each of these neurons were

reported in Fig 4B. **E.** (left) Representative trace of bursting of a hypoglossal neuron resulting

from preBötC drive during ChrMP (purple) and in Tolb (green) (n=4). Scale bars: 10 mV x

500 msec. (right) Comparison of action potentials generated per preBötC burst during ChrMP

and Tolb (n=4). The effect of ChrMP on baseline action potentials generated per preBötC burst

for each of these neurons were reported in Fig 4C. **F.** (left) Representative traces of current

clamp recordings in response to ramp current injection in ChrMP (purple) and in Tolb (green).

Scale bar: 1 sec x 10mV). (right) Rheobase comparison from inspiratory hypoglossal neurons

during ChrMP and Tolb (n=7). The effect of ChrMP on rheobase for each of these neurons were

reported in Fig 4C.

607   **<u>Citations</u>**

608   1   Lyons, M. M., Bhatt, N. Y., Pack, A. I. & Magalang, U. J. Global burden of sleep-
609   disordered breathing and its implications. *Respirology* **25**, 690-702,
610   doi:10.1111/resp.13838 (2020).

611   2   Malhotra, A. *et al.* Metrics of sleep apnea severity: beyond the apnea-hypopnea index.
612   *Sleep* **44**, doi:10.1093/sleep/zsab030 (2021).

613   3   Mehra, R. Sleep apnea and the heart. *Cleve Clin J Med* **86**, 10-18,
614   doi:10.3949/ccjm.86.s1.03 (2019).

615   4   Yeghiazarians, Y. *et al.* Obstructive Sleep Apnea and Cardiovascular Disease: A
616   Scientific Statement From the American Heart Association. *Circulation* **144**, e56-e67,
617   doi:10.1161/CIR.0000000000000988 (2021).

618   5   Loffler, K. A. *et al.* Continuous Positive Airway Pressure Treatment, Glycemia, and
619   Diabetes Risk in Obstructive Sleep Apnea and Comorbid Cardiovascular Disease.
620   *Diabetes Care* **43**, 1859-1867, doi:10.2337/dc19-2006 (2020).

621   6   Hua, F. New insights into diabetes mellitus and its complications: a narrative review. *Ann
622   Transl Med* **8**, 1689, doi:10.21037/atm-20-7243 (2020).

623   7   Daulatzai, M. A. Cerebral hypoperfusion and glucose hypometabolism: Key
624   pathophysiological modulators promote neurodegeneration, cognitive impairment, and
625   Alzheimer's disease. *J Neurosci Res* **95**, 943-972, doi:10.1002/jnr.23777 (2017).

626   8   Force, U. S. P. S. T. *et al.* Screening for Obstructive Sleep Apnea in Adults: US
627   Preventive Services Task Force Recommendation Statement. *JAMA* **317**, 407-414,
628   doi:10.1001/jama.2016.20325 (2017).

629   9   Castro, D. & Freeman, L. A. in *StatPearls*   (2022).

630   10   Genta, P. R. *et al.* Airflow Shape Is Associated With the Pharyngeal Structure Causing
631   OSA. *Chest* **152**, 537-546, doi:10.1016/j.chest.2017.06.017 (2017).

11 Neelapu, B. C. *et al.* Craniofacial and upper airway morphology in adult obstructive sleep
apnea patients: A systematic review and meta-analysis of cephalometric studies. *Sleep*
Med Rev **31**, 79-90, doi:10.1016/j.smrv.2016.01.007 (2017).

12 Vos, W. G., De Backer, W. A. & Verhulst, S. L. Correlation between the severity of sleep
apnea and upper airway morphology in pediatric and adult patients. *Curr Opin Allergy*
Clin Immunol **10**, 26-33, doi:10.1097/ACI.0b013e328334f659 (2010).

13 Kubin, L. Neural Control of the Upper Airway: Respiratory and State-Dependent
Mechanisms. *Compr Physiol* **6**, 1801-1850, doi:10.1002/cphy.c160002 (2016).

14 Eckert, D. J., White, D. P., Jordan, A. S., Malhotra, A. & Wellman, A. Defining
phenotypic causes of obstructive sleep apnea. Identification of novel therapeutic targets.
Am J Respir Crit Care Med **188**, 996-1004, doi:10.1164/rccm.201303-0448OC (2013).

15 Nemati, S. *et al.* Model-based characterization of ventilatory stability using spontaneous
breathing. *J Appl Physiol (1985)* **111**, 55-67, doi:10.1152/japplphysiol.01358.2010
(2011).

16 Peng, Y. J. *et al.* Complementary roles of gasotransmitters CO and H2S in sleep apnea.
Proc Natl Acad Sci U S A **114**, 1413-1418, doi:10.1073/pnas.1620717114 (2017).

17 Peng, Y. J., Zhang, X., Nanduri, J. & Prabhakar, N. R. Therapeutic Targeting of the
Carotid Body for Treating Sleep Apnea in a Pre-clinical Mouse Model. *Adv Exp Med Biol*
1071, 109-114, doi:10.1007/978-3-319-91137-3_14 (2018).

18 Osman, A. M., Carter, S. G., Carberry, J. C. & Eckert, D. J. Obstructive sleep apnea:
current perspectives. *Nat Sci Sleep* **10**, 21-34, doi:10.2147/NSS.S124657 (2018).

19 Prabhakar, N. R. & Semenza, G. L. Gaseous messengers in oxygen sensing. *J Mol Med*
(Berl) **90**, 265-272, doi:10.1007/s00109-012-0876-1 (2012).

20 Mancuso, C. Heme oxygenase and its products in the nervous system. *Antioxid Redox*
Signal **6**, 878-887, doi:10.1089/ars.2004.6.878 (2004).

21 Schwartz, A. R. *et al.* Obesity and obstructive sleep apnea: pathogenic mechanisms and
therapeutic approaches. *Proc Am Thorac Soc* **5**, 185-192, doi:10.1513/pats.200708-
137MG (2008).

22 Horner, R. L. The neuropharmacology of upper airway motor control in the awake and
asleep states: implications for obstructive sleep apnoea. *Respir Res* **2**, 286-294,
doi:10.1186/rr71 (2001).

23 Horner, R. L. Emerging principles and neural substrates underlying tonic sleep-state-
dependent influences on respiratory motor activity. *Philos Trans R Soc Lond B Biol Sci*
364, 2553-2564, doi:10.1098/rstb.2009.0065 (2009).

24 Horton, G. A. *et al.* Activation of the Hypoglossal to Tongue Musculature Motor Pathway
by Remote Control. *Sci Rep* **7**, 45860, doi:10.1038/srep45860 (2017).

25   Fleury Curado, T. *et al.* Chemogenetic stimulation of the hypoglossal neurons improves
upper airway patency. *Sci Rep* **7**, 44392, doi:10.1038/srep44392 (2017).

26   Fleury Curado, T. A. *et al.* Silencing of Hypoglossal Motoneurons Leads to Sleep
Disordered Breathing in Lean Mice. *Front Neurol* **9**, 962, doi:10.3389/fneur.2018.00962
(2018).

27   Yuan, S. *et al.* Hydrogen sulfide metabolism regulates endothelial solute barrier function.
Redox Biol **9**, 157-166, doi:10.1016/j.redox.2016.08.004 (2016).

28   Garcia, A. J., 3rd, Dashevskiy, T., Khuu, M. A. & Ramirez, J. M. Chronic Intermittent
Hypoxia Differentially Impacts Different States of Inspiratory Activity at the Level of the
preBotzinger Complex. *Front Physiol* **8**, 571, doi:10.3389/fphys.2017.00571 (2017).

29   Koizumi, H. *et al.* Structural-functional properties of identified excitatory and inhibitory
interneurons within pre-Botzinger complex respiratory microcircuits. *J Neurosci* **33**,
2994-3009, doi:10.1523/JNEUROSCI.4427-12.2013 (2013).

30   Revill, A. L. *et al.* Dbx1 precursor cells are a source of inspiratory XII premotoneurons.
Elife **4**, doi:10.7554/eLife.12301 (2015).

31   Garcia, A. J., 3rd *et al.* Chronic Intermittent Hypoxia Alters Local Respiratory Circuit
Function at the Level of the preBotzinger Complex. *Front Neurosci* **10**, 4,
doi:10.3389/fnins.2016.00004 (2016).

32   Ding, Y., Li, Y. L. & Schultz, H. D. Downregulation of carbon monoxide as well as nitric
oxide contributes to peripheral chemoreflex hypersensitivity in heart failure rabbits. *J
Appl Physiol (1985)* **105**, 14-23, doi:10.1152/japplphysiol.01345.2007 (2008).

33   Prabhakar, N. R. Carbon monoxide (CO) and hydrogen sulfide (H(2)S) in hypoxic
sensing by the carotid body. *Respir Physiol Neurobiol* **184**, 165-169,
doi:10.1016/j.resp.2012.05.022 (2012).

34   Morikawa, T. *et al.* Hypoxic regulation of the cerebral microcirculation is mediated by a
carbon monoxide-sensitive hydrogen sulfide pathway. *Proc Natl Acad Sci U S A* **109**,
1293-1298, doi:10.1073/pnas.1119658109 (2012).

35   Mustafa, A. K. *et al.* Hydrogen sulfide as endothelium-derived hyperpolarizing factor
sulfhydrates potassium channels. *Circ Res* **109**, 1259-1268,
doi:10.1161/CIRCRESAHA.111.240242 (2011).

36   Haller, M., Mironov, S. L., Karschin, A. & Richter, D. W. Dynamic activation of K(ATP)
channels in rhythmically active neurons. *J Physiol* **537**, 69-81, doi:10.1111/j.1469-
7793.2001.0069k.x (2001).

37   Lape, R. & Nistri, A. Current and voltage clamp studies of the spike medium
afterhyperpolarization of hypoglossal motoneurons in a rat brain stem slice preparation.
J Neurophysiol **83**, 2987-2995, doi:10.1152/jn.2000.83.5.2987 (2000).

38   Zanella, S. *et al.* When norepinephrine becomes a driver of breathing irregularities: how
intermittent hypoxia fundamentally alters the modulatory response of the respiratory
network. *J Neurosci* **34**, 36-50, doi:10.1523/JNEUROSCI.3644-12.2014 (2014).

39   Enokido, Y. *et al.* Cystathionine beta-synthase, a key enzyme for homocysteine
metabolism, is preferentially expressed in the radial glia/astrocyte lineage of developing
mouse CNS. *FASEB J* **19**, 1854-1856, doi:10.1096/fj.05-3724fje (2005).

40   da Silva, G. S. F. *et al.* Excitatory Modulation of the preBotzinger Complex Inspiratory
Rhythm Generating Network by Endogenous Hydrogen Sulfide. *Front Physiol* **8**, 452,
doi:10.3389/fphys.2017.00452 (2017).

Supplementa