Novel Methods for Multi-ancestry Polygenic Prediction and their Evaluations in 5.1 Million Individuals of Diverse Ancestry

Haoyu Zhang[1,2*], Jianan Zhan[3], Jin Jin[4], Jingning Zhang[4], Wenxuan Lu[5], Ruzhang Zhao[4], Thomas U. Ahearn[1], Zhi Yu[6], Jared O'Connell[3], Yunxuan Jiang[3], Tony Chen[2], Dayne Okuhara[7], 23andMe Research Team, Montserrat Garcia-Closas[1], Xihong Lin[2,6,8], Bertram L. Koelsch[3], Nilanjan Chatterjee[4,9*]

[1] Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD, USA

[2] Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA, USA

[3] 23andMe Inc., Sunnyvale, CA, USA

[4] Department of Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

[5] Department of Applied Mathematics and Statistics, Johns Hopkins University, Baltimore, MD, USA

[6] Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[7] Booz Allen Hamilton, McLean, Virginia, USA

[8] Department of Statistics, Harvard University, Cambridge, MA, USA

[9] Department of Oncology, School of Medicine, Johns Hopkins University, Baltimore, MD, USA

Conflicts of interest: J.Z., Y.J., J.O., and B.L.K. are employed by and hold stock or stock options in 23andMe, Inc.

* Correspondence to: Haoyu Zhang (haoyu.zhang2@nih.gov) and Nilanjan Chatterjee (nilanjan@jhu.edu)

Abstract

Polygenic risk scores are becoming increasingly predictive of complex traits, but their suboptimal performance in non-European ancestry populations raises questions about their clinical applications and impact on health inequities. We develop CT-SLEB, a powerful and scalable method to calculate PRS based on ancestry-specific GWAS summary statistics from multi-ancestry training samples by integrating multiple techniques, including clumping and thresholding, empirical Bayes and super learning. We evaluate the performance of the proposed method and nine alternatives using large-scale simulated GWAS on ~19 million common variants and large datasets from 23andMe Inc., the Global Lipids Genetics Consortium, All of Us and UK Biobank, which include up to 1.18 million individuals from four non-European populations across thirteen complex traits. Results show that the proposed method can substantially improve the performance of PRS in non-European populations relative to simple alternatives and has comparable or superior performance relative to a recent method that requires a higher order of computational time. Furthermore, our simulation studies provide novel insights to sample size requirements and the effect of SNP density on multi-ancestry risk prediction.

Introduction

Genome-wide association studies (GWAS) have identified tens of thousands of single nucleotide polymorphisms (SNPs) associated with complex traits and diseases[1,2]. Polygenetic risk scores (PRSs), which summarize the combined effect of individual SNPs, have the potential to improve risk stratification for various diseases and conditions[3–13]. However, GWAS to date have mostly been conducted in populations predominately comprised of European (EUR) origin individuals[14]. Consequently, the PRSs generated from these studies tend to underperform in non-EUR populations, particularly in African (AFR) ancestry populations[9,15–18]. The lack of representation of non-EUR populations in PRS research has thus raised concerns that the use of current PRSs for clinical applications may exacerbate health inequities[19–22].

In addition to the critical importance of addressing inequalities in representation of non-EUR population in genetic research, there is also an important need to develop statistical methods that leverage genetic data across populations to develop better performing PRS. Most of the PRS methods to date have been developed to analyze data from a single ancestry group[23–32], and subsequently, their performance was primarily evaluated for risk prediction in EUR populations[4–6,8–11]. While the same methods can also be used to build PRS in non-EUR populations, the resulting PRS, irrespective of the methods, tend to have limited performance due to limited sample sizes of the training datasets compared to sample sizes in EUR populations[15,20]. Some studies have conducted meta-analyses of GWAS across diverse populations to develop an underlying multi-ancestry PRS[33–35]. While such an approach may lead to a single PRS that performs more "equally" across diverse groups, it does not account for heterogeneity in linkage disequilibrium (LD) and effect sizes across populations and not designed for deriving the best PRS possible for each of the underlying populations[36,37].

Recent methods have focused on developing more optimal PRS in non-EUR populations by combining available GWAS from the target population of interest with "borrowed" information from larger GWAS in the EUR populations. One such study developed PRS in separate populations and then combined the PRS by optimally

weighting them to maximize performance in the target population[38]. Other studies have

proposed Bayesian methods that leverage multivariate priors for effect-size distribution

to borrow information across populations[36,39–41]. In spite of these developments,

methods for building PRS leveraging multi-ancestry datasets remain limited. Both

theoretical and empirical studies have indicated that the optimal method for building

PRS depends on multiple factors[23,42,43], including sample size, heritability, effect-size

distribution and LD, and thus exploration of alternative methods with complementary

advantages are needed to build optimal PRS is any given setting. Moreover, and

perhaps more importantly, evaluation of the scope of multi-ancestry methods for

building improved PRS remains quite limited to date due to the lack of suitably large

GWAS for various non-EUR populations, especially of African origin, where risk

prediction remains the most challenging.

In this paper, we propose a computationally simple and powerful method for generating

PRSs using GWAS across diverse ancestry population. The method, which we refer to

as CT-SLEB, is a model-free approach and combines the strength of multiple

techniques, including a two-dimensional extension of the popular clumping and

thresholding (CT) method[23,24], a super-learning (SL) model for combining multiple PRS

and an empirical-Bayes (EB) approach to effect-size estimation. We compare the

performance of the proposed method with nine alternative methods based on large-

scale simulated GWAS across five ancestry groups. In addition, we develop and

validate population-specific PRS for thirteen complex traits using GWAS data from

23andMe, Inc., the Global Lipids Genetics Consortium (GLGC), All of Us (AoU) and UK

Biobank (UKBB) across EUR ($N \approx 3.91$ million), AFR (primarily African American (AA),

$N \approx 265K$), Latino ($N \approx 574K$), East Asian (EAS, $N \approx 270K$), and South Asian (SAS,

$N \approx 77K$). Both simulation studies and empirical data analyses indicate that CT-SLEB is

a scalable and powerful method for generating PRS for non-EUR populations. Further,

our simulation studies and evaluation of various methods in large datasets provide

insights into the future yield of multi-ancestry PRSs as GWAS in diverse populations

continues to grow.

Results

Method overview

CT-SLEB is a method designed to generate multi-ancestry PRSs that incorporate

existing large GWAS from EUR populations and smaller GWAS from non-EUR

populations. The method has three key steps (**Figure 1, Supplementary Figure 1**): 1.

Clumping and Thresholding (CT) for selecting SNPs to be included in a PRS for the

target population; 2. Empirical-Bayes (EB) method for estimating the coefficients of the

SNPs; 3. Super-learning (SL) model to combine a series of PRSs generated under

different SNP selection thresholds. The method requires three independent datasets:

(1) GWAS summary statistics from training datasets across EUR and non-EUR

populations; (2) a tuning dataset for the target population to find optimal model

parameters; and (3) a validation dataset for the target population to report the final

prediction performance.

Two-dimensional Clumping and Thresholding

In step one, CT-SLEB uses two-dimensional clumping and thresholding on GWAS

summary-statistics data to incorporate SNPs with either shared effects across the EUR

and the target populations or population-specific effects in the target population (**Figure**

1a). Each SNP is assigned to the following two groups based on the p-value from the

EUR and the target population: 1. SNPs with a p-value smaller in the EUR population;

2. SNPs with a p-value smaller in the target population or those which exist only in the

target population. SNPs in the first group are ranked based on EUR p-value (smallest to

largest) and then clumped using LD estimates from the EUR reference sample. SNPs in

the second group are ranked based on the target population p-value and clumped using

LD estimates from the target population reference samples. Then the clumped SNPs

from the two groups are combined as a candidate set for the next step. In the

thresholding step, p-value thresholds are varied over a two-dimensional set of grid

points. Each dimension corresponds to the threshold used for the p-value obtained from

one of the populations. At any given combination of thresholds, a SNP may be included

in the target population PRS if the corresponding p-value from either the EUR or the

target population achieves the corresponding threshold.

Empirical-Bayes Estimation of Effect Sizes

Since the effect sizes of SNPs are expected to be correlated across populations[44,45], we propose an EB method to efficiently estimate effect sizes for SNPs to be included in the PRSs (**Figure 1b**). Based on the selected SNP set from the CT step, we first estimate an underlying "prior" covariance matrix of effect sizes between the EUR population and the target population. Then, we estimate the effect size for each SNP in the target population based on the corresponding posterior mean, which weighs the effect-size estimate available from each population based on the bias-variance trade-off (**Methods**).

Super Learning

Previous research has shown that combining PRSs under different p-value thresholds can efficiently increase the prediction performance[26]. Thus, as a final step, we propose a super-learning model to predict the outcome using PRSs generated under different tuning parameters as training variables (**Figure 1c**). The super-learning model is a linear combination of different predictors based on multiple supervised learning algorithms[46–49]. The set of prediction algorithms can be self-designed or chosen from classical prediction algorithms. We use the R package SuperLearner version 2.0-26[50] and choose Lasso[51], ridge regression[52], and neural networks[53] as three different candidate models in the implementation. We train the super-learning model on the tuning dataset and evaluate the performance of the final PRS using the independent validation dataset.

Design of Simulation Studies.

We conduct simulation studies to compare ten methods across five broad categories: 1. *single ancestry methods*, which utilize training and tuning data solely from the target population; 2. *EUR PRS*, generated using single ancestry methods to EUR only GWAS data; 3. *weighted PRS,* which applies a single ancestry PRS approach separately to the EUR and target population, and derives an optimal linear combination of the two; 4. *Bayesian method,* which assumes a multi-variate Bayesian framework and uses the

posterior mean of effect sizes to construct the PRS; and 5. our proposed approach CT-SLEB. The single ancestry methods include two representative PRS methods: CT[23,24] and LDpred2[25,32]. The EUR PRSs are generated based on CT and LDpred2. The weighted PRS approaches include three representative PRS methods: a weighted PRS based on CT, a weighted PRS based on LDpred2, and PolyPred-S+[54] that linearly combines PRSs using EUR and target population PRS from SBayesR[27], and EUR PRS from PolyFun-pred[55]. The PolyFun-pred approach integrates functional annotation information to identify causal variant across the genome and thus uses additional information that is not incorporated in the other methods compared. The Bayesian method include: 1. XPASS method[40], which assumes a multivariate normal distribution to model the effect-size and uses the posterior mean of the target population to construct PRS. 2. PRS-CSx[36], which uses a continuous shrinkage Bayesian framework to calculate the posterior mean of effect sizes for the EUR and the non-EUR populations, and subsequently derives an optimal linear combination of all populations using a tuning dataset. All methods are implemented using the target population and EUR population as training data to construct the PRS for the target population. Additionally, CT-SLEB and PRS-CSx are further evaluated using data from all five ancestries. For computational efficiency, most analyses are restricted to ~2.0 million SNPs included in Hapmap3 (HM3)[56], or the Multi-Ethnic Genotyping Arrays (MEGA)[57] chips array, or both. However, the PolyPred-S+ and PRS-CSx methods can currently only be implemented with ~1.3 million HM3 SNPs in the provided software, and thus the application of these methods in our analysis is also restricted to only the HM3 SNPs.

Simulation Study Results

Results from simulation studies (**Figure 2** and **Supplementary Figure 2-6)** show that generally multi-ancestry methods lead to the most accurately PRSs in different settings. When the training data sample size for the target population is small (**Figure 2a, Supplementary Figure 2a, 3-6 a**), PRSs derived from the single ancestry methods perform poorly compared to the EUR-based PRS. On the other hand, when the training sample size for the target population is large (**Figure 2b, Supplementary Figure 2b, 3-6 b-d**), PRSs generated by the single ancestry methods can outperform EUR PRS.

PRS generated from the multi-ancestry methods can lead to substantial improvement in

either setting.

When only EUR and target population data are used to build the PRS, we observe that

both CT-SLEB and PRS-CSx can lead to improvement over other candidate methods in

most settings. When the sample size for target population is large, weighted LDpred2

has comparable performance as CT-SLEB and PRS-CSx. Between CT-SLEB and PRS-

CSx, neither is uniformly superior to the other across all scenarios considered. When

the sample size for the target population is relatively small (N = 15K), PRS-CSx often

outperforms CT-SLEB when the degree of polygenicity is the highest ($p_{causal} = 0.01$) .

On the other hand, in the same sample size setting, CT-SLEB often outperforms PRS-

CSx by a notable margin when the degree of polygenicity is the lowest ($p_{causal} =$

5×10^{-4}). When the sample size for the target population is larger (N=45K-100K), the

difference between the two methods decreases. When all five ancestries data are used

to build the PRS, CT-SLEB and PRS-CSx improves by 6.8% and 23.7% on average,

respectively, compared to only using EUR and the target population data. PRS-CSx

outperforms CT-SLEB in many settings (**Figure 2b**, **Supplementary Figure 2a-b, 3-6**

b-d). Under different simulation settings, the number of SNPs used by CT-SLEB ranged

from 549K to 933K, while PRS-CSx retained all HM3 SNPs (**Supplementary Table 1**).

In a comparison of runtime using data on chromosome 22 to construct PRS for AFR

(**Methods, Supplementary Table 2**), we observed runtime of CT-SLEB is on average

almost 25 times faster than that of PRS-CSx (4.35 vs. 109.11 mins) in two ancestries

analyses, and 91 times faster than that of PRS-CSx in five ancestries setting (4.62 vs.

420.96 mins) using a single core with Intel E5-26840v4 CPU. Sensitivity analyses are

conducted to evaluate the required tuning and validation sample size for CT-SLEB

(**Supplementary Figure 7**). When the tuning and validation sample size is larger, the

prediction performance on average will increase. Meanwhile, the performance of CT-

SLEB is still robust when the tuning and validation sample size is around 2000.

Additional sensitivity analyses are conducted to compare the performance of CT-SLEB

and PRS-CSx when both methods are constrained to HM3 SNPs. Results show that

CT-SLEB maintains advantage over PRS-CSx in low polygenicity setting when training sample size is 15K or 45K (**Supplementary Figure 8a-b**). However, when training sample size is more than 80K, both methods have similar performance in low polygenicity setting (**Supplementary Figure 8c-d**).

Unequal predictive performance of PRS across different populations has been considered as a barrier to ethical implementation of the technology in healthcare. Thus, we examined how large the sample size one may need for training GWAS in various minority populations to bridge the gap in the performance of PRS in comparison to that of the EUR population. Results indicate that when effect sizes for shared causal SNPs are similar across populations (genetic correlation=0.8), the gap can be mostly eliminated for all populations except AFR when the sample size reaches between 45% to 80% of that of the EUR population (**Figure 3**, **Supplementary Figure 8**). For the AFR population, however, the sample size requirement can dramatically vary depending on the underlying genetic architecture of the traits. If the common SNP heritability is assumed to be the same for the AFR population as that of the other populations, then the sample size requirement for the AFR population is dauntingly large because of substantially smaller per-SNP heritability (**Figure 3a-b**, **Supplementary Figure 9a-b**). If we allow the per-SNP heritability to remain the same across populations, but heritability to vary proportionately to the number of common variants, then the sample size requirement for the AFR population is similar to those for the other minority populations (**Figure 3c-d**, **Supplementary Figure 9c)**.

A major advantage of CT-SLEB over PRS-CSx is its computational scalability of the former method to handle much larger number of SNPs. Thus we use CT-SLEB to investigate the effect of SNP density on PRS performance by considering three different SNP sets to be used for PRS building: (1) ~1.3 million SNPs represented in HM3[56] (2) ~2.0 million SNPs that include all HM3 SNPs and additional SNPs represented in the MEGA array (3) All ~19 million common SNPs included in the 1000 Genomes Project (Phase 3)[58] which were used to generate the traits in our simulation studies. We observe that in general performances of PRS in various US minority populations can be

substantially enhanced by inclusion of SNPs in denser panels, and the benefit due to

denser panels is more enhanced when the sample size for the target population is

larger and in settings where the proportions of causal SNPs are smaller (**Figure 4 and**

Supplementary Figure 10).

23andMe data analysis results

We develop and validate population-specific PRS for seven complex traits using GWAS

data from 23andMe, Inc. (**Methods, Supplementary Table 3-4**). We conduct GWAS

using a training dataset for each population adjusting for principal component (PC) 1-5,

sex and age following standard quality control (**Methods**). The Manhattan plots and QQ

plots for GWAS are shown in **Supplementary Figures 11-17**, and no inflation is

observed given the genomic inflation factor (**Supplementary Table 5**). We estimate

heritability for the seven traits in the EUR population using LD-score regression[59]

(**Supplementary Table 6, Methods**).

Results for heart metabolic disease burden and height (**Figure 5, Supplementary**

Table 7) show a similar pattern as our simulation studies. The PRS-CSx using five

ancestries generally lead to the best performing PRS across different populations.

When only EUR and the target population data are used, both CT-SLEB and PRS-CSx

led to good performance across different populations. Compared to the best performing

EUR or single ancestry PRS, the relative gain is often large, especially for the AA

population. The weighted method did not perform well for the AA population, but it

substantially improved performance compared to each component PRS (EUR and

single ancestry) for other populations. PolyPred-S+ has comparable performance as

PRS-CSx and CT-SLEB on EAS and SAS population, but significantly worse on AA

population. We also observe that even with the best performing method and large

sample sizes across all populations, a significant gap remains for the performance of

the PRSs in non-EUR populations compared to those in the EUR population (**Figure 5**).

We also observe similar trends for the analyses of 23andMe data for the five binary

traits: any cardiovascular disease (any CVD), depression, migraine diagnosis, morning

person, and sing back musical note (SBMN) (**Figure 6, Supplementary Table 7**). For most settings, CT-SLEB, PRS-CSx and PolyPred-S+ often produce the best performing PRS and often lead to substantial improvement over best EUR PRS or single ancestry PRS. For CVD, which is the clinically most relevant trait for risk prediction and preventive intervention, we observe that CT-SLEB tends to outperform PRS-CSx and PolyPred-S+ by a notable margin except for the EAS population. We also observe that for the AFR population, which are particularly underrepresented in genetic research, CT-SLEB outperforms PRS-CSx and PolyPred-S+ by a notable margin for several traits (e.g., CVD and morning person). In contrast, PRS-CSx and PolyPred-S+ outperforms CT-SLEB by a significant margin for predicting migraine diagnosis and SBMN in the SAS population. Similar to the continuous traits, we also observe that even with best performing methods and substantially large GWAS in a number of non-EUR populations, major gap often remains for the performance of PRS in these populations compared to those for the EUR population.

GLGC and AoU data analysis results with UKBB as validation dataset

We develop and validate population-specific PRS using GWAS summary data for four traits from GLGC: high-density lipoprotein cholesterol (HDL), low-density lipoprotein cholesterol (LDL), log triglycerides (logTG) and total cholesterol (TC), and two traits from AoU: height and body mass index (BMI) (**Methods, Supplementary Table 4, 8**). The method is evaluated using individual level data from UKBB (**Supplementary Table 9**). The Manhattan plots and QQ plots for GWAS are shown in **Supplementary Figures 18-23**, and no inflation is observed given the genomic inflation factor for most ancestries other than height for Latino population in AoU ($\lambda_{1000} = 1.04$, **Supplementary Table 5**). We estimate heritability for the four traits in the EUR population using LD-score regression[59] (**Supplementary Table 6, Methods**).

In the data analyses using the GLGC data, CT-SLEB, PRS-CSx and PolyPred-S+ outperform the other approaches (**Figure 7**). Among these three methods, CT-SLEB demonstrates superior performance in the AFR population. For example, CT-SLEB leads to an improvement of 140% and 66.4% in adjusted R^2 for log-TG and LDL,

respectively, compared to the PRS-CSx method. This highlights the advantage of model-free approach CT-SLEB in handling biomarker traits, which can be associated with some SNPs with unusually large effects. On the other hand, PRS-CSx clearly outperforms CT-SLEB on the EAS population. For instance, for log-TG and LDL, the ratio of the adjusted R^2 between CT-SLEB and PRS-CSx is only 89.8% and 52.9%, respectively. These results show importance of generating PRS using multiple alternative methods in multi-ancestry settings. Another interesting observation is that, for LDL and TC, the PRS generated by both CT-SLEB and the five ancestries PRS-CSx demonstrated superior performance in the AFR population when compared to the performance of the corresponding EUR-PRSs in the European population (**Figure 7**). Finally, in AoU data analyses, we observe CT-SLEB performs better in predicting BMI, while PRS-CSx has better performance in predicting height, but both show comparable performance to each other (**Figure 8**).

To have a direct comparison between CT-SLEB and PRS-CSx, we report the prediction performance R^2 (continuous traits) or logit-scale variance (converted from Area under the ROC curve (AUC) for binary traits, **Supplementary Note**) between CT-SLEB and PRS-CSx and then average over the thirteen traits within each ancestry in data analyses of 23andMe, GLGC and AoU (**Supplementary Table 10**). When only the EUR and the target population data are used for PRS construction, the averaged performance ratio between CT-SLEB and PRS-CSx is 144%, 88.5%, 103% and 92.1% for AFR (primarily AA), EAS, Latino, SAS, respectively. When data from five ancestries or three ancestries (AoU) is used for PRS construction, the averaged performance ratio is 143%, 91.7%, 94.5%, 89.6% for AFR (primarily AA), EAS, Latino, SAS, respectively.

Discussion

In summary, we have proposed CT-SLEB as a powerful and computationally scalable method to generate optimal PRS across ancestrally distinct groups by utilizing GWAS across diverse populations. We compare the performance of CT-SLEB with those from a variety of both simple and complex methods, in large-scale simulation studies and very large datasets. Results indicate that there is not a uniformly best performing

method across all scenarios, and it is important to generate PRS using alternative

methods across multi-ancestries. Encouragingly, however, the CT-SLEB method led to

marked improvement in performance of PRS compared to alternatives in AFR origin

populations where polygenic prediction has been most challenging. Computationally,

CT-SLEB is an order of magnitude faster than a recently proposed Bayesian method,

PRS-CSx[36], and can more easily handle much larger SNP contents and additional

populations.

A unique contribution of our study is the evaluation of a variety of PRS methodologies in

the unprecedented large and diverse datasets from the 23andMe, Inc., GLGC, AoU and

UKBB GWAS datasets. Our findings provide important insights into the future yield of

emerging large multi-ancestry GWAS. Adult height is often used as a model trait to

explore the genetic architecture of complex traits and the potential for polygenic

prediction. We observe that the standard CT method, when trained in ~2 million EUR

individuals from 23andMe data, leads to a PRS for the underlying population with a

prediction R^2 of approximately 0.276. Application of LD-score regression to the same

23andMe data leads to an estimated GWAS heritability of height of 0.395, indicating

that the PRS has achieved about 69.8% (0.276/0.395) of its maximum potential in the

23andMe EUR population. We observe, however, that even with the best method and

large sample size of the 23andMe GWAS (N_{Latino}~350K and N_{AFR} ~100K), the highest

prediction accuracy of height PRS for non-EUR populations among all the methods is

substantially lower compared to that of the EUR population (Relative R^2~0.48

(0.133/0.276) for AFR, 0.76 (0.210/0.276) for EAS, ~ 0.86 (0.237/0.276) for Latino, and

~ 0.80 (0.222/0.276) for SAS compared to that of EUR. The average relative R^2

(continuous traits) or logit-scale variance (binary traits) for AFR, EAS, Latino, and SAS

across the thirteen evaluated traits is 0.50, 0.76, 0.77, and 0.79, respectively

(**Supplementary Table 11**).

We also observe similar patterns for other traits, including disease outcomes for which

risk prediction is of most interest. For CVD, for example, the CT method, when trained

in a sample of ~700K cases and ~1.3 million controls from the EUR population,

produces a PRS that by itself has a prediction accuracy of the AUC as 0.65. For other

populations, in some of which the sample size is considerably large (N_{case}/$N_{control}$

=32K/66K for AA and N_{case}/$N_{control}$ =84K/270K for Latino populations) but still much

smaller than that of the EUR population, the AUCs for best performing PRSs are close

to 60% or lower. Further, the sample size is not the only driving factor for differential

performances of PRS across populations. For example, the performance of the best

performing CVD-PRS for the Latino and South Asian populations are very similar even

though the sample size for the later population is much smaller. Collectively, these and

additional results from simulation studies, indicate that bridging the gap between PRS

performance across populations will require much more parity in the sample size of the

underlying GWAS.

Both our simulation studies and data analyses indicate that no single PRS method is

expected to be uniformly most powerful in all settings. In the analysis of GLGC data, for

example, we found CT-SLEB led to improvement in PRS performance by a large margin

compared to PRS-CSx for the AFR population, but the converse was true for the EAS

population. In general, the optimal method for generating PRS will depend on the nature

of the underlying multivariate effect-size distribution of the traits across different

populations. While Bayesian methods, in principle, can generate the optimal PRS under

correct specification of underlying effect-size distribution[25,32,36,40], modeling of effect-size

distribution in multi-ancestry settings can be challenging. In the analysis of lipid traits in

GLGC, for example, we found the Bayesian methods could not account well for

existence of large-effect SNPs in the AFR population due to inadequacy of the

underlying model for the effect-size distribution. The CT method and their extensions,

on the other hand, while they do not require strong modeling assumptions about effect-

size distribution, they do not optimally incorporate LD among SNPs. Our analysis

reveals advantages of alternative methods in different settings by contrasting of results

across a variety of complex traits which are known to have distinct architecture in terms

of heritability, polygenicity and number of clusters of distinct effect-sizes[60,61]. We thus

advocate that in future applications, researchers consider generating and evaluating a

variety of PRS obtained from complementary methods. As different PRS may contain

some orthogonal information, at the end, instead of choosing one best PRS, the best strategy could be to combine them using a final super learning step.

Our study has several limitations. While sample sizes for 23andMe datasets are extremely large, the power of genetic risk prediction is likely to have been blunted in this population, compared to other settings, due to the presence of a higher level of environmental heterogeneity. For example, a recent study[62] reported achieving prediction R^2 for height of ~41% for the EUR individuals within the UKBB using a PRS developed on ~1.1 million individuals from the UKBB (N=400K) and 23andMe (N=700K). In comparison, the PRS prediction R^2 for height we could achieve within 23andMe EUR population is only ~30% despite doubling the sample size of the training dataset. We, however, also note that the estimate of heritability in 23andMe ($h_{SNP}^2 = 0.395$) is substantially smaller than those previously reported[63,64] based on the UKBB ($h_{SNP}^2 \sim 0.5 - 0.7$), which is potentially due to the LD differences between 23andMe and 1000 Genomes Project. When we compare the results across the two studies using prediction R^2 relative to the underlying heritability of the respective populations, we do see a significant gain in performance due to the increased sample size of the current study. Thus, we believe that while caution is needed to extrapolate 23andMe study results to other populations, the relative performance of PRSs we observe across different methods and different ancestry groups within this population is likely to be generalizable to other settings.

While we have conducted large-scale simulation across various scenarios, the simulated genotype data from HapGen2 may not fully reflect the levels of differentiation within and across ancestries due to limited haplotype data within 1000 Genomes Project. A major limitation of our proposed method, as well as a variety of other existing methods, is that they are primarily designed to generate PRS across ancestrally distinct populations. However, many populations, such as the AFR and Latino origin in the US, are highly admixed, and for these populations, the development or clinical reporting of PRS can be improved by explicitly taking into account individual-level estimates of admixture proportions[65]. Additionally, our method assumes that individual-level data is

available for model tuning and validation, but this is not a fundamental limitation as summary-statistics-based methods[66,67] could also be used in these steps. Moreover, while we observe a consistent increase of five ancestries CT-SLEB over two ancestries CT-SLEB in simulations, the results of real data analyses don't show the same consistent pattern (**Figure 5-7**)**.** This may be due to the underlying effect-size across different ancestries being much more complex than the simulations, as the simulations only assume that causal SNPs effect-size follow a multi-variate normal distribution.

In conclusion, we have proposed a novel and computationally scalable method for generating powerful PRS using data from GWAS studies in diverse populations. Further, our simulation studies and data analysis across multiple traits involving large 23andMe Inc., GLGC, AoU and UKBB studies provide unique insight into what is likely to emerge from future GWAS in diverse populations for years to come.

Author contribution

H.Z. and N.C. conceived the project. H.Z., J.Z., J.J., J.Z., W.L. and R.Z. carried out all data analyses with supervision from N.C.; J.Z., J.O.C., Y.J. run GWAS for training data from 23andMe Inc. with the supervision from B.L.K.; R.Z. run GWAS for training data from AoU with the supervision from N.C. and H.Z.; H.Z., T.C., and D.O. developed the software and online resources for data sharing; H.Z., J.Z., J.J., J.Z., W.L., R.Z. and N.C. drafted the manuscript, and X.L., M.G.C. and T.U.A. provided comments. All co-authors reviewed and approved the final version of the manuscript. The following members of the 23andMe Research Team contributed to this study: Stella Aslibekyan, Adam Auton, Elizabeth Babalola, Robert K. Bell, Jessica Bielenberg, Katarzyna Bryc, Emily Bullis, Daniella Coker, Gabriel Cuellar Partida, Devika Dhamija, Sayantan Das, Sarah L. Elson, Nicholas Eriksson, Teresa Filshtein, Alison Fitch, Kipper Fletez-Brant, Pierre Fontanillas, Will Freyman, Julie M. Granka, Karl Heilbron, Alejandro Hernandez, Barry Hicks, David A. Hinds, Ethan M. Jewett, Yunxuan Jiang, Katelyn Kukar, Alan Kwong, Keng-Han Lin, Bianca A. Llamas, Maya Lowe, Jey C. McCreight, Matthew H. McIntyre, Steven J. Micheletti, Meghan E. Moreno, Priyanka Nandakumar, Dominique T. Nguyen, Elizabeth S. Noblin, Jared O'Connell, Aaron A. Petrakovitz, G. David Poznik, Alexandra Reynoso, Morgan Schumacher, Anjali J. Shastri, Janie F. Shelton, Jingchunzi Shi, Suyash Shringarpure, Qiaojuan Jane Su, Susana A. Tat, Christophe Toukam Tchakouté, Vinh Tran, Joyce Y. Tung, Xin Wang, Wei Wang, Catherine H. Weldon, Peter Wilton, Corinna D. Wong.

Acknowledgements

We would like to thank the research participants and employees of 23andMe, Inc for making this work possible. We want to thank Liz Noblin, Melissa J. Francis and Emily Voeglein for helping with the research collaboration agreement with Harvard T.H. Chan School of Public Health, Johns Hopkins Bloomberg School of Public Health and 23andMe, Inc. The analysis utilized the high-performance computation Biowulf cluster at National Institutes of Health, USA, Faculty of Arts and Sciences Research Computing Cluster at Harvard University, and the Joint High Performance Computing Exchange at Johns Hopkins Bloomberg School of Public Health. The UK Biobank data was obtained

under the UK Biobank resource application 17712. This work was funded by NIH grants: K99 CA256513-01 (H. Z.), K99 HG012223 (J.J.), NHLBI 5T32HL007604-37 (Z.Y.), R35-CA197449, U19-CA203654, R01-HL163560, U01-HG009088 U01-HG012064 (X. L.), R01 HG010480-01 (N. C. and J. Z.) and U01HG011724 (N. C.). The AoU Research Program is supported by the National Institutes of Health, Office of the Director: Regional Medical Centers: 1 OT2 OD026549; 1 OT2 OD026554; 1 OT2 OD026557; 1 OT2 OD026556; 1 OT2 OD026550; 1 OT2 OD 026552; 1 OT2 OD026553; 1 OT2 OD026548; 1 OT2 OD026551; 1 OT2 OD026555; IAA #: AOD 16037; Federally Qualified Health Centers: HHSN 263201600085U; Data and Research Center: 5 U2C OD023196; Biobank: 1 U24 OD023121; The Participant Center: U24 OD023176; Participant Technology Systems Center: 1 U24 OD023163; Communications and Engagement: 3 OT2 OD023205; 3 OT2 OD023206; and Community Partners: 1 OT2 OD025277; 3 OT2 OD025315; 1 OT2 OD025337; 1 OT2 OD025276. In addition, the AoU Research Program would not be possible without the partnership of its participants.

Code and Data availability

CT-SLEB package and tutorial: https://github.com/andrewhaoyu/CTSLEB

P + T: https://www.cog-genomics.org/plink/1.9/

SCT and LDpred2: https://github.com/privefl/bigsnpr.

XPASS: https://github.com/YangLabHKUST/XPASS

PolyPred-S+: https://github.com/omerwe/polyfun

PRS-CSx: https://github.com/getian107/PRScsx

LDSC: https://github.com/bulik/ldsc

PLINK: https://www.cog-genomics.org/plink/1.9/

Simulated genotype data for 600K subjects from five ancestries:

https://dataverse.harvard.edu/dataset.xhtml?persistentId=doi:10.7910/DVN/COXHAP

GWAS summary level statistics for five ancestries from GLGC:

http://csg.sph.umich.edu/willer/public/glgc-lipids2021/results/ancestry_specific/

The full GWAS summary statistics for the 23andMe discovery data set could be made available through 23andMe to qualified researchers under an agreement with 23andMe that protects the privacy of the 23andMe participants. Please visit

https://research.23andme.com/collaborate/#dataset-access/ for more information and to apply to access the data. Participants provided informed consent and participated in the research online, under a protocol approved by the external AAHRPP-accredited IRB, Ethical & Independent Review Services.

References

1. Buniello, A. *et al.* The NHGRI-EBI GWAS Catalog of published genome-wide association studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res.* **47**, D1005–D1012 (2019).

2. MacArthur, J. *et al.* The new NHGRI-EBI Catalog of published genome-wide association studies (GWAS Catalog). *Nucleic Acids Res.* **45**, D896–D901 (2017).

3. Chatterjee, N., Shi, J. & García-Closas, M. Developing and evaluating polygenic risk prediction models for stratified disease prevention. *Nat. Rev. Genet.* **17**, 392–406 (2016).

4. Khera, A. v. *et al.* Genome-wide polygenic scores for common diseases identify individuals with risk equivalent to monogenic mutations. *Nat. Genet.* **50**, 1219–1224 (2018).

5. Rao, A. S. & Knowles, J. W. Polygenic risk scores in coronary artery disease. *Curr. Opin. Cardiol.* **34**, 435–440 (2019).

6. Mavaddat, N. *et al.* Polygenic Risk Scores for Prediction of Breast Cancer and Breast Cancer Subtypes. *Am. J. Hum. Genet.* **104**, 21–34 (2019).

7. Lambert, S. A., Abraham, G. & Inouye, M. Towards clinical utility of polygenic risk scores. *Hum. Mol. Genet.* **28**, R133–R142 (2019).

8. Jia, G. *et al.* Evaluating the Utility of Polygenic Risk Scores in Identifying High-Risk Individuals for Eight Common Cancers. *JNCI Cancer Spectr.* **4**, (2020).

9. Dikilitas, O. *et al.* Predictive Utility of Polygenic Risk Scores for Coronary Heart Disease in Three Major Racial and Ethnic Groups. *Am. J. Hum. Genet.* **106**, 707–716 (2020).

10. Li, R., Chen, Y., Ritchie, M. D. & Moore, J. H. Electronic health records and polygenic risk scores for predicting disease risk. *Nat. Rev. Genet.* **21**, 493–502 (2020).

11. Zhang, H. *et al.* Genome-wide association study identifies 32 novel breast cancer susceptibility loci from overall and subtype-specific analyses. *Nat. Genet.* **52**, 572–581 (2020).

12. Graff, R. E. *et al.* Cross-cancer evaluation of polygenic risk scores for 16 cancer types in two large cohorts. *Nat. Commun.* **12**, (2021).

13. Wray, N. R. *et al.* From Basic Science to Clinical Application of Polygenic Risk Scores: A Primer. *JAMA Psychiatry* **78**, 101–109 (2021).

14. Fatumo, S. *et al.* A roadmap to increase diversity in genomic studies. *Nat. Med.* **28**, 243–250 (2022).

15. Duncan, L. *et al.* Analysis of polygenic risk score usage and performance in diverse human populations. *Nat. Commun.* **10**, 1–9 (2019).

16. Liu, C. *et al.* Generalizability of Polygenic Risk Scores for Breast Cancer Among Women With European, African, and Latinx Ancestry. *JAMA Netw. Open* **4**, e2119084–e2119084 (2021).

17. Du, Z. *et al.* Evaluating Polygenic Risk Scores for Breast Cancer in Women of African Ancestry. *J. Natl. Cancer. Inst.* **113**, 1168–1176 (2021).

18. Wojcik, G. L. *et al.* Genetic analyses of diverse populations improves discovery for complex traits. *Nature* **570**, 514–518 (2019).

19. Martin, A. R. *et al.* Human Demographic History Impacts Genetic Risk Prediction across Diverse Populations. *Am J Hum Genet* **100**, 635–649 (2017).

20. Martin, A. R. *et al.* Clinical use of current polygenic risk scores may exacerbate health disparities. *Nat Genet* **51**, 584–591 (2019).

21. Wang, Y. *et al.* Theoretical and empirical quantification of the accuracy of polygenic scores in ancestry divergent populations. *Nat Commun* **11**, 1–9 (2020).

22. Kullo, I. J. *et al.* Polygenic scores in biomedical research. *Nat. Rev. Genet.* **23**, 524–532 (2022).

23. Wray, N. R., Goddard, M. E. & Visscher, P. M. Prediction of individual genetic risk to disease from genome-wide association studies. *Genome Res.* **17**, 1520–1528 (2007).

24. Purcell, S. M. *et al.* Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature* **460**, 748–752 (2009).

25. Vilhjálmsson, B. J. *et al.* Modeling Linkage Disequilibrium Increases Accuracy of Polygenic Risk Scores. *Am. J. Hum. Genet.* **97**, 576–592 (2015).

26. Privé, F., Vilhjálmsson, B. J., Aschard, H. & Blum, M. G. B. Making the Most of Clumping and Thresholding for Polygenic Scores. *Am. J. Hum. Genet.* **105**, 1213–1221 (2019).

27. Lloyd-Jones, L. R. *et al.* Improved polygenic prediction by Bayesian multiple regression on summary statistics. *Nat. Commun.* **10**, 1–11 (2019).

28. Newcombe, P. J., Nelson, C. P., Samani, N. J. & Dudbridge, F. A flexible and parallelizable approach to genome-wide polygenic risk scores. *Genet. Epidemiol.* **43**, 730–741 (2019).

29. Ge, T., Chen, C. Y., Ni, Y., Feng, Y. C. A. & Smoller, J. W. Polygenic prediction via Bayesian regression and continuous shrinkage priors. *Nat. Commun.* **10**, 1–10 (2019).

30. Song, S., Jiang, W., Hou, L. & Zhao, H. Leveraging effect size distributions to improve polygenic risk scores derived from summary statistics of genome-wide association studies. *PLoS Comput. Biol.* **16**, e1007565 (2020).

31. Zhou, G. & Zhao, H. A fast and robust Bayesian nonparametric method for prediction of complex traits using summary statistics. *PLoS Genet.* **17**, (2021).

32. Privé, F., Arbel, J. & Vilhjálmsson, B. J. LDpred2: better, faster, stronger. *Bioinformatics* **36**, 5424–5431 (2021).

33. Koyama, S. *et al.* Population-specific and trans-ancestry genome-wide analyses identify distinct and shared genetic risk loci for coronary artery disease. *Nat. Genet.* **52**, 1169–1177 (2020).

34. Sakaue, S. *et al.* Trans-biobank analysis with 676,000 individuals elucidates the association of polygenic risk scores of complex traits with human lifespan. *Nat. Med.* **26**, 542–548 (2020).

35. Agbaedeng, T. A. *et al.* Polygenic risk score and coronary artery disease: A meta-analysis of 979,286 participant data. *Atherosclerosis* **333**, 48–55 (2021).

36. Ruan, Y. *et al.* Improving polygenic prediction in ancestrally diverse populations. *Nat. Genet.* **54**, 573–580 (2022).

37. Tian, P. *et al.* Multiethnic polygenic risk prediction in diverse populations through transfer learning. *Front. Genet.* **13**, 1854 (2022).

38. Márquez-Luna, C. *et al.* Multiethnic polygenic risk scores improve risk prediction in diverse populations. *Genet. Epidemiol.* **41**, 811–823 (2017).

39. Xiao, J. *et al.* XPXP: improving polygenic prediction by cross-population and cross-phenotype analysis. *Bioinformatics* **38**, 1947–1955 (2022).

40. Cai, M. *et al.* A unified framework for cross-population trait prediction by leveraging the genetic correlation of polygenic traits. *Am. J. Hum. Genet.* **108**, 632–655 (2021).

41. Hoggart, C. BridgePRS: A powerful trans-ancestry Polygenic Risk Score method. *Preprint* doi:10.21203/rs.3.rs-1272562/v1.

42. Dudbridge, F. & Wray, N. R. Power and Predictive Accuracy of Polygenic Risk Scores. *PLoS Genet.* **9**, e1003348 (2013).

43. Chatterjee, N. *et al.* Projecting the performance of risk prediction based on polygenic analyses of genome-wide association studies. *Nat. Genet.* **45**, 400–405 (2013).

44. Brown, B. C., Ye, C. J., Price, A. L. & Zaitlen, N. Transethnic Genetic-Correlation Estimates from Summary Statistics. *Am. J. Hum. Genet.* **99**, 76–88 (2016).

45. Shi, H. *et al.* Population-specific causal disease effect sizes in functionally important regions impacted by selection. *Nat. Commun.* **12**, 1–15 (2021).

46. van der Laan, M. J., Polley, E. C. & Hubbard, A. E. Super learner. *Stat. Appl. Genet. Mol. Biol.* **6**, (2007).

47. Polley, E. & van der Laan, M. J. Super Learner In Prediction. *U.C. Berkeley Division of Biostatistics Working Paper Series* (2010).

48. van der Laan, M. J. & Rose, S. *Targeted learning: causal inference for observational and experimental data.* vol. 4 (Springer New York , 2011).

49. Ledell, E., Petersen, M. & van der Laan, M. J. Computationally efficient confidence intervals for cross-validated area under the ROC curve estimates. *Electron. J. Stat.* **9**, 1583–1607 (2015).

50. Polley, E., LeDell, E., Kennedy, C. & van der Laan, M. J. SuperLearner: Super Learner Prediction. *R package version 2.0-26* (2019).

51. Tibshirani, R. Regression Shrinkage and Selection Via the Lasso. *J. R. Stat. Soc. Series B Stat. Methodol.* **58**, 267–288 (1996).

52. Friedman, J., Hastie, T. & Tibshirani, R. Regularization Paths for Generalized Linear Models via Coordinate Descent. *J. Stat. Softw.* **33**, 1 (2010).

53. Ripley, B. D. *Pattern recognition and neural networks*. (Cambridge university press, 2007).

54. Weissbrod, O. *et al.* Leveraging fine-mapping and multipopulation training data to improve cross-population polygenic risk scores. *Nat. Genet.* **54**, 450–458 (2022).

55. Weissbrod, O. *et al.* Functionally informed fine-mapping and polygenic localization of complex trait heritability. *Nat. Genet.* **52**, 1355–1363 (2020).

56. Consortium, T. I. H. 3. Integrating common and rare genetic variation in diverse human populations. *Nature* **467**, 52 (2010).

57. Bien, S. A. *et al.* Strategies for Enriching Variant Coverage in Candidate Disease Loci on a Multiethnic Genotyping Array. *PLoS One* **11**, 167758 (2016).

58. Auton, A. *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74 (2015).

59. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat. Genet.* **47**, 291–295 (2015).

60. Zhang, Y., Qi, G., Park, J. H. & Chatterjee, N. Estimation of complex effect-size distributions using summary-level statistics from genome-wide association studies across 32 complex traits. *Nat. Genet.* **50**, 1318–1326 (2018).

61. Zhang, Y. D. *et al.* Assessment of polygenic architecture and risk prediction based on common variants across fourteen cancers. *Nat. Commun.* **11**, 1–13 (2020).

62. Márquez-Luna, C. *et al.* Incorporating functional priors improves polygenic prediction accuracy in UK Biobank and 23andMe data sets. *Nat. Commun.* **12**, (2021).

63. Ge, T., Chen, C. Y., Neale, B. M., Sabuncu, M. R. & Smoller, J. W. Phenome-wide heritability analysis of the UK Biobank. *PLoS Genet.* **13**, (2017).

64. Yengo, L. *et al.* Meta-analysis of genome-wide association studies for height and body mass index in ∼700000 individuals of European ancestry. *Hum. Mol. Genet.* **27**, 3641–3649 (2018).

65. Ding, Y. *et al.* Polygenic scoring accuracy varies across the genetic ancestry continuum in all human populations. *bioRxiv* (2022) doi:10.1101/2022.09.28.509988.

66. Song, L. *et al.* SummaryAUC: a tool for evaluating the performance of polygenic risk prediction models in validation datasets with only summary level statistics. *Bioinformatics* **35**, 4038–4044 (2019).

67. Zhao, Z. *et al.* PUMAS: fine-tuning polygenic risk scores with GWAS summary statistics. *Genome Biol.* **22**, 1–19 (2021).

68. Pritchard, J. K. & Przeworski, M. Linkage Disequilibrium in Humans: Models and Data. *Am. J. Hum. Genet.* **69**, 1–14 (2001).

69. Su, Z., Marchini, J. & Donnelly, P. HAPGEN2: simulation of multiple disease SNPs. *Bioinformatics* **27**, 2304–2305 (2011).

70. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

71. Foucher, Y. *et al.* RISCA: Causal Inference and Prediction in Cohort-Based Analyses. *R package version 1.01* Preprint at https://cran.r-project.org/package=RISCA (2020).

72. Graham, S. E. *et al.* The power of genetic diversity in genome-wide association studies of lipids. *Nature* **600**, 675–679 (2021).

Online Methods

We assume there are $l = 1, \dots L$ populations with $l = 1$ indexing the EUR population. We

assume that for each population, summary-statistics data from underlying GWAS are

available in the form of $(\hat{\beta}_{kl}, s_{kl}, p_{kl})$ for $k = 1, 2, \dots, K_L$ SNPs, where $\hat{\beta}, s$ and p denote

estimates of effect-size, standard errors and p-values associated with individual SNPs,

respectively. We further assume that additional datasets are available for each target

population of interest, which could be split into tuning and validation sets. Our proposed

CT-SLEB method contains three steps: 1. Two-dimensional clumping and thresholding

(CT); 2. EB procedure; 3. Super-learning algorithm, which will be described in detail in

the following three subsections.

CT. In this step, we extend the traditional CT to a two-dimensional setting so that PRS

for a target population can be built using approximately independent SNPs that show

significant association in at least one of the two populations (majority population and the

target population). The CT method has two components, the Clumping step and the

Thresholding step. In the two-dimensional setting where the lead SNPs might be

informed by GWAS of either the EUR or target population, it is unclear what reference

sample is the most suited for LD clumping. After initial exploration of alternative

approaches through simulation studies, we find the most informative approach is to split

the SNPs into two sets depending on which population they show stronger signals and

then perform LD clumping for each set separately based on the reference sample for

the respective population. For the thresholding step, we select SNPs based on two

distinct thresholds for their respective p-values in the two populations. As the optimal

threshold for p-value selection is known to depend on sample size for underlying

GWAS[24,42,43], and sample sizes for GWAS across EUR and minority populations are

highly differential, we anticipate (and confirm through simulation studies) that a two-

dimensional approach for threshold selection is more optimal than using a single p-

value threshold across both populations. Following, we describe details of the CT step:

1. The clumping r^2-cutoff and base size of the clumping window size w_b vary

across (0.01, 0.05, 0.1, 0.2, 0.5, 0.8) and (50kb, 100kb), respectively. The

clumping window size w_s is defined as w_b/r^2 because LD is inversely proportional to the genetic distance between variants[26,68].

2. Select all SNPs with smaller p-values in EUR ($p_{k1} < p_{k2}$), and then, clump based on p_{k1} using LD estimates from the EUR reference samples with selected r^2 and w_s.

3. Select all variants with smaller p-values in the target population ($p_{k2} < p_{k1}$) and the population-specific SNPs, and then, clump based on p_{k2} using LD estimates from the reference samples of the target population with the same r^2-cutoff and w_b.

4. Combine the post-clumping variants from the second and third steps as the candidate variants set.

5. Define two different p-value cutoffs (p_{t1}, p_{t2}) for the EUR and the target population. A variant is selected if $p_{k1} < p_{t1}$ or $p_{k2} < p_{t2}$. We allow p_{t1} and p_{t2} to vary in the set: $(5 \times 10^{-8}, 5 \times 10^{-7}, 5 \times 10^{-6}, \dots, 5 \times 10^{-1}, 1.0)$. With the cross combination of p_{t1} and p_{t2}, a total of 81 different p-value cutoffs are applied.

6. With the cross combination of p_{t1}, p_{t2}, r^2 and w_b, a total of 972 PRSs (6 r^2 thresholds * 2 w_b windows * 81 value thresholds) are evaluated on the tuning dataset using estimated regression coefficients ($\hat{\beta}_{k2}$) from GWAS of the target population.

EB to calibrate regression coefficients. In the CT step above, we use $\hat{\beta}_{k2}$ from the target population to calculate PRS. However, $\hat{\beta}_{k2}$ can be noisy when the GWAS sample size of the target population is small. Meanwhile, given the high genetic correlation across different ancestries[44,45], effect sizes from other populations can be used to calibrate the regression coefficients for the PRS. Although we only use p-values from GWAS for the EUR and the target population for selecting SNPs in the CT step, the EB step takes advantage of existing GWAS from multiple populations. Suppose $\hat{\boldsymbol{u}}_k = (\hat{u}_{k1}, \dots, \hat{u}_{kL}) = (\hat{\beta}_{k1}\sqrt{2f_{k1}(1 - f_{k1})}, \dots, \hat{\beta}_{kL}\sqrt{2f_{kL}(1 - f_{kL})})$, is the vector of the standardized effect-size for the kth SNP in L different populations, with $\hat{\boldsymbol{s}}_k^* =$

$(s_{k1}^*, \ldots, s_{kL}^*) = \left(s_{k1}\sqrt{2f_{k1}(1-f_{k1})}, \ldots, \hat{s}_{kL}\sqrt{2f_{kL}(1-f_{kL})}\right)$ being the vector of the

corresponding standard errors of $\hat{\boldsymbol{u}}_k$. We assume that $\hat{\boldsymbol{u}}_k|u_k \sim N(\boldsymbol{u}_k, \boldsymbol{\Sigma_k})$, where $\boldsymbol{\Sigma_k} =$

$diag\{(\hat{\boldsymbol{s}}_k^*)^2\}$ given that the GWAS for different populations are independent. Additionally,

we assume that the prior distribution of the mean of $\hat{\boldsymbol{u}}_k$ is $\boldsymbol{u}_k \sim N(\boldsymbol{0}, \boldsymbol{\Sigma_0})$, which assumes

the effect-size follows the strong negative selection model. By integrating the conditional

and prior distribution, we can obtain the marginal distribution of $\hat{\boldsymbol{u}}_k$ as $N(\boldsymbol{0}, \boldsymbol{\Sigma_0} + \boldsymbol{\Sigma_k})$.

Suppose the SNP set selected from the CT step has K^* variants overlapped across all

the populations. We estimate the prior covariance matrix $\boldsymbol{\Sigma_0}$ using the K^* overlapped

variants shared across all populations as:

10
$$\widehat{\boldsymbol{\Sigma_0}} = \frac{1}{K^*-1}\sum_{k=1}^{K^*} \hat{\boldsymbol{u}}_k^T \hat{\boldsymbol{u}}_k - \boldsymbol{\Sigma_k}.$$

11 We note that we ignore any potential correlation across selected SNPs in this step, but

12 the estimate is still expected to be consistent for $\boldsymbol{\Sigma_0}$ which represents marginal variance-

13 covariance matrices for effect sizes associated with an individual SNP across

14 populations. Applying the Bayes formula, the posterior distribution of $\boldsymbol{u}_k$ becomes

15
$$\boldsymbol{u}_k|\hat{\boldsymbol{u}}_k \sim N\left(\widehat{\boldsymbol{\Sigma_0}}(\widehat{\boldsymbol{\Sigma_0}} + \boldsymbol{\Sigma_k})^{-1}\hat{\boldsymbol{u}}_k, \widehat{\boldsymbol{\Sigma_0}}(\widehat{\boldsymbol{\Sigma_0}} + \boldsymbol{\Sigma_k})^{-1}\boldsymbol{\Sigma_k}\right).$$

16 The EB coefficients for the kth SNP are defined as:

17
$$\widehat{\boldsymbol{\beta}}_k^{EB} = \mathbf{F_k}\,\widehat{\boldsymbol{\Sigma_0}}(\widehat{\boldsymbol{\Sigma_0}} + \boldsymbol{\Sigma_k})^{-1}\hat{\boldsymbol{u}}_k,$$

where $\widehat{\boldsymbol{\beta}}_k^{EB}$ is a $L \times 1$ vector for the posterior effect-size of the kth SNP for in each of the

L populations, and $\mathbf{F_k} = \text{diag}\left\{\frac{1}{\sqrt{2f_{kl}(1-f_{kl})}}\right\}_{L \times L}$ is the scaling matrix to scale effect sizes

from the standardized scale back to the original scale. Preliminary simulation studies

indicate that EB step of effect-size calibration leads to distinct improvement in PRS

performance (compared to using effect-size estimates from the target population)

irrespective of all other steps.

To save computational time, we estimate $\widehat{\boldsymbol{\Sigma}}_0$ in the above step only once based on the

SNP set that gives the best PRS in the CT step across all different p-value thresholds,

r^2-cutoff, and window sizes. We then apply the same $\widehat{\boldsymbol{\Sigma}}_0$ to derive the EB-calibrated

effect sizes for SNPs included in all different PRSs corresponding to cross combination

1. of p_{t1}, p_{t2}, r^2-cutoff and w_b. Using the EB-calibrated effect sizes, we can compute 972

2. PRSs for each population (6 r^2 thresholds * 2 w_b windows * 81 value thresholds). In all

3. analyses, we compute the 1944 PRSs using EB-calibrated effect sizes of the target

4. population and EUR. When more than two ancestries are involved, we use data from all

5. populations to derive the EB estimates of effect-sizes for SNPs for each population

6. (**Supplementary Note**). However, to save computational time at the super-learning

7. step, we derive the final PRS for the target population by only incorporating the 1944

8. PRSs derived for the larger EUR population and the target population. All 1944 PRSs

9. are used as input for the super-learning step to predict the outcome for the target

10. population. Because many PRSs are highly correlated with each other, we filter out the

11. highly redundant one with pairwise correlations higher than 0.98. In the simulations, 369

12. of the 1994 PRSs are kept on average after the filtering.

13.

14. **Super learning.** We combine all PRSs generated from the above steps into an input

15. dataset and train them on the tuning dataset to predict the outcome Y. The super-

16. learning algorithm generates an optimally weighted combination from a set of distinct

17. prediction algorithms[46–49] (**Supplementary Note**). The set of prediction algorithms can

18. be self-designed or chosen from classical prediction algorithms e.g., Lasso[51], ridge

19. regression[52], neural networks[53], etc. We use three different prediction algorithms

20. implemented in the SuperLearner package[50] to generate the super learning estimate:

21. Lasso[51], ridge regression[52] and neural networks[53]. For binary traits, since the ridge

22. regression algorithm is currently not supported by the SuperLearner package now, we

23. only use Lasso and neural network in the data analysis. To use AUC as the objective

24. function, we use the flag "method = method.AUC" in the SupearLearner package.

25.

26. **Simulation.** Large-scale multi-ancestry genotype data are generated using HAPGEN2

27. (version 2.1.2)[69] mimicking the LD of EUR, AFR, Americas (AMR), East Asia (EAS) and

28. South Asia (SAS). The 1000 Genomes Project (Phase 3)[58] is used as the reference

29. panel which include 503 EUR, 661 AFR, 347 AMR, 504 EAS and 489 SAS subjects.

30. Biallelic SNPs with MAF more than 0.01 in any of the populations are kept in the

31. reference panel, resulting in ~8.6 million SNPs for EUR, ~14.8 million SNPs for AFR,

~9.8 million SNPs for AMR, ~7.6 million SNPs for EAS, and ~9.0 million SNPs for SAS.

The genotype data are generated with a total of ~19.2 million SNPs. The set of

simulated variants for all five ancestries is same. Because allele frequencies vary

across populations, a common SNP can be population-specific if its MAF is greater than

0.01 in only one population. The proportion of population-specific SNPs range from

2.92% for AMR to 43.84% for AFR, respectively (**Supplementary Figure 24**). If a

particular variant's MAF is less than 0.01 in a given population, we don't include them in

the PRS calculation for that population because the LD estimation for rare variants can

be unstable. We simulate a total of 120,000 independent subjects for each of the

population.

For generating trait values, we select causal SNPs randomly across the whole genome

with the causal SNP proportion being set to 0.01, 0.001, or 5×10^{-4}. We consider two

alternative models for generating heritability distribution within each population: (A)

Constant common-SNP heritability. (B) Constant per-SNP heritability that implies the

total heritability is proportional to the number of common SNPs. We also consider three

different models for negative selection pattern: strong, mild and no negative selection.

We denote by u_{kl} the standardized effect-size for kth causal SNP for the lth population.

Under strong negative selection and constant heritability model, the standardized effect-

sizes are drawn from a multivariate normal distribution of the form:

22
$$u_{kl} \sim N\left(0, \frac{h^2}{C_l}\right), cov\left(u_{kl_1}, u_{kl_2}\right) = \frac{\rho h^2}{\sqrt{C_{l_1} C_{l_2}}},$$

where C_l is the number of causal SNPs with MAF > 0.01 in the lth population, the

heritability h^2 associated with common SNPs for each population is set to 0.4, and the

genetic correlation ρ is set to 0.8. We then generate the phenotype using linear model

of the form $Y_{il} = \sum_{k=1}^{C_l} \frac{G_{ikl}}{\sqrt{2(f_{kl}(1-f_{kl})}} u_{kl} + \epsilon_{il}$ for the ith subject in the lth population,

where f_{kl} is the effect allele frequency for the kth causal SNP in lth population. The error

terms are generated as $\epsilon_{il} \sim N(0, 1 - h^2)$. We also consider mild negative selection ($u_{kl}^2 \propto$

$[f_{kl}(1 - f_{kl})]^{0.75}$) and no negative selection ($u_{kl}^2 \propto [f_{kl}(1 - f_{kl})]$) scenarios (see

Supplemental Notes for details). Finally, we simulate data under an assumption of total

heritability of all ~19 million SNPs being 0.4 across all populations, but the common

SNP heritability varying proportionately to their number within each the populations. The

model assumes per SNP heritability to be the same across all populations and thus

leads to the common SNP heritability value of 0.32, 0.21, 0.16, 0.19 and 0.17 for AFR,

AMR, EAS, EUR and SAS, respectively. The genetic correlation is set to 0.8 or 0.6.

We set the training sample sizes for each target population to 15,000, 45,000, 80,000,

or 100,000. We generate GWAS summary statistics for each population based on the

training samples using PLINK version 1.90 with the command "--linear". We fixed the

sample sizes for the EUR population at 100,000. We further simulate the tuning and

validation dataset of size 10,000 for each target population. The final prediction R^2 is

reported as the average of ten independent simulation replicates for each simulation

setting. For evaluating CT-SLEB that incorporates data across all five ancestries, we

assume the training sample size for the each of the other non-EUR populations to be

the same as that of the target population.

Existing PRS methods. The CT method selects clumped SNPs with different p-value

thresholds and picks a single optimal PRS based on its performance on the tuning

dataset. We implement CT using PLINK version 1.90[70] with the clumping step command

"--clump --clump-r2 0.1 --clump-kb 500". We estimate LD based on 3,000 randomly

selected unrelated subjects from the training dataset for each population. We set the

candidate p-value thresholds to be $(5 \times 10^{-8}, 1 \times 10^{-7}, 5 \times 10^{-7}, 1 \times 10^{-6}, \dots, 5 \times$

$10^{-1}, 1.0)$ and for computing PRS, we use the PLINK command "--score no-sum no-

mean-imputation". The optimal p-value threshold is determined based prediction R^2

(variation explained by the corresponding PRS) on the tuning dataset.

The LDpred2 method infers SNP effect sizes by a shrinkage estimator that combines

GWAS summary statistics with a prior on effect sizes while leveraging LD information

from an external reference panel. LDpred2 is implemented using the R package

"bigsnpr"[32]. The tuning parameters included are: (1) the proportion of causal SNPs, with

candidate values set to a sequence of length 17 that are evenly spaced on a logarithmic scale from 10^{-4} to 1; (2) per-SNP heritability, with candidate values set to 0.7, 1, or 1.4 times the total heritability estimated by LD score regression divided by the number of causal SNPs; (3) "sparse" option, which is set to "yes" or "no" (the "sparse" option sets some weak effects to zero). The method selects tuning parameters based on the performance on the tuning dataset.

The EUR PRS based on CT or LDpred2 are built based on training dataset from the EUR population and estimates tuning parameters based on tuning sample for the EUR population. When the EUR PRSs are evaluated in the target population, we exclude the SNPs that do not exist in the target population.

The weighted-PRS linearly combines the CT or LDpred2 PRS generated from the EUR and from the target population. The weights for EUR PRS and for the target population PRS in the linear combination are estimated using the tuning dataset from the target population through a linear regression. We implement the weighted-PRS using R version 4.0.0.

The PolyPred-S+ method consists of a PolyFun-pred predictor trained on the EUR population and two SBayesR predictors that use the training data from the EUR and the target population respectively. On a tuning dataset from the target population, PolyPred-S+ performs the non-negative least squares regression to compute the mixture weights and linearly combines the predictors. In PolyFun-pred, PolyFun leverages genome-wise functional annotations to compute the prior causal probabilities, which are fed into the fine-mapping method SuSiE for the posterior causal effect estimation. On the other hand, SBayesR estimates the posterior tagging effects by posing a finite normal mixture prior on the effect sizes. For PolyFun-pred, we use the precomputed prior causal probabilities provided by the authors, extract the LD information using EUR population in 1000 Genomes Project (Phase 3), and assume 10 causal SNPs per locus. Using GCTB (2.03 beta version), we train SBayesR with the sparse shrunk LD matrix for HapMap3 variants published by the SBayesR authors. Currently, the shrunk LD matrix

are only available for EUR populations. Therefore, SBayesR predictors for EUR and the target population both used the shrunk LD matrix for EUR. We specify the model parameters and the MCMC settings in the same way as the PolyPred-S+ authors did in their UKBB simulations.

The XPASS method leverages the genetic correlation between the target and the EUR population by assuming a bivariate normal distribution with non-zero covariance for the pairs of effect sizes that correspond to the same SNP in two populations. It can incorporate population-specific covariates as fixed effects to enhance the accuracy of their weight estimation. For each population, we extract the top 20 PCs from the reference genome as the covariates. When estimating the LD matrices, ancestry-specific reference data is inputted in the function. LD matrices are estimated based on the LD blocks for EUR to all datasets, given that XPASS package only offers choices of EUR and EAS.

The PRS-CSx method estimates population-specific SNP effect sizes based on a Bayesian framework using continuous shrinkage priors to jointly model the GWAS summary statistics from multiple populations. Besides the Bayesian modeling step, PRS-CSx further conducts a step similar to weighted-PRS, which is to linearly combine the PRS based on the posterior effect-sizes obtained from the EUR and the target population with the weights in the linear combination being estimated based on the tuning dataset of the target population. We implemented PRS-CSx following the guidance provided in https://github.com/getian107/PRScsx. We set the hyperparameters a and b in the gamma-gamma prior to their default values of 1 and 0.5, respectively. Further, the parameter ϕ is varied over the default set of values 10^{-6}, 10^{-4}, 10^{-2}, and 1. The optimal ϕ is determined based on the performance on the tuning dataset.

Runtimes and memory usage. The computation time and memory usage of CT-SLEB (two ancestries), CT-SLEB (five ancestries), and PRS-CSx are compared based on

their performance on chromosome 22 and assuming AFR is the target population. All

analyses are performed using a single core with Intel E5-26840v4 CPU. The reported

performance is averaged over 100 replicates. The training dataset includes GWAS

summary statistics for AFR (N_{GWAS}=15,000) and for EUR (N_{GWAS}=100,000) population.

The tuning dataset and validation dataset each contains 10,000 subjects. For five

ancestries analyses, the training GWAS sample sizes for AMR, EAS and SAS are all

set to 15,000.

23andMe Data analysis. The individuals included in our analyses are part of the

23andMe participant cohort. All these individuals included have provided informed

consent and answered surveys online according to our human subject protocol

reviewed and approved by Ethical & Independent Review Services, a private

institutional review board (http://www.eandireview.com). Detailed information about

genotyping, quality control, imputation, removing related individuals, and ancestry

determination is provided in **Supplementary Note**. Participants were included in the

analysis based on consent status as checked when data analyses were initiated.

The analyses include five ancestries: AA, EAS, EUR, Latino, and SAS. Meanwhile, the

analyses include two continuous and five binary traits: 1. Heart metabolic disease

burden 2. Height 3. Any CVD 4. Depression 5. Migraine Diagnosis 6. Morning Person 7.

SBMN. We randomly split the data for each population into training, tuning, and

validation datasets with the proportion of 70%, 20%, and 10%, with detailed sample size

in **Supplementary Table 3**. We perform GWAS for the seven traits using the training

dataset for each population, adjusting for PC 1-5, sex, and age using standard quality

control procedures (**Supplementary Note**). SNPs with MAF > 0.01 in at least one of the

five populations are kept in the analyses. We further restrict analyses to SNPs that are

on HM3 + MEGA chips with ~2.0 million SNPs. We use LDSC version 1.01[59] to estimate

the heritability using the GWAS summary statistics of EUR populations for the seven

traits. We estimate the LD score using the 503 unrelated samples of EUR ancestry from

1000 Genomes Project. We restrict heritability analyses to EUR populations since some

non-EUR populations don't have sufficient sample size to get stable estimate from LD-score regression.

We apply seven methods to compare PRS prediction performance, CT, LDPred2, best EUR PRS based on CT and LDpred2, weighted-PRS, PRS-CSx, CT-SLEB using data from EUR and the target population, and CT-SLEB using all five populations. Since individual-level data is not available in the training step, we use the reference data from the 1000 Genomes Project (Phase 3) to estimate the LD for each population. Specifically, we use AFR and AMR from the 1000 Genomes Project as the reference for the AA and Latino population in 23andMe, respectively. All PRS prediction performances are reported based on the independent validation dataset that is independent of the training and tuning datasets. To calculate the adjusted R^2 for continuous traits, we first regress the traits on covariates and then evaluate performance of the PRS to predict residualized trait values. To calculate the adjusted AUC for binary traits, we used the roc.binary function in the R package RISCA version 1.01[71].

GLGC Data analysis. The GWAS summary statistics of four blood lipids traits are obtained from the public available database of GLGC (http://csg.sph.umich.edu/willer/public/glgc-lipids2021/results/ancestry_specific/). UKBB data are removed from the GWAS summary statistics. The details of study design of study, genotyping, quality control and GWAS are described elsewhere[72]. The training data are available for four blood lipid traits: LDL, HDL, logTG, TC from five different ancestries: EUR, AFR (primarily AA), Latino, EAS, and SAS (**Supplementary Table 8**). The tuning + validation data from UKBB dataset are from EUR, AFR, EAS and SAS ancestries (**Supplementary Table 9**). Details of ancestry prediction of UKBB are described in **Supplementary Note**. Latino ancestries in UKBB can't not be well classified using the ancestry prediction algorithm, and the corresponding sample size is too low to provide meaningful comparison of methods. Therefore, Latino population is evaluated using UKBB data. The implementation of the ten different PRSs approaches follow the same steps as the 23andMe data analyses. We use the reference data from

the 1000 Genomes Project (Phase 3) to estimate the LD for each population. The

adjusted R^2 are adjusted by sex, age and genetic PC1-10.

AoU Data analysis. The individuals included in our analyses are part of the AoU

participant cohort. All these individuals' information have been collected according to

AoU Research Program Operational Protocol

(https://allofus.nih.gov/sites/default/files/aou_operational_protocol_v1.7_mar_2018.pdf).

Detailed information about genotyping, ancestry determination, quality control, removing

related individuals are listed in AoU Research Program Genomic Research Data Quality

Report (https://www.researchallofus.org/wp-content/themes/research-hub-wordpress-

theme/media/2022/06/All%20Of%20Us%20Q2%202022%20Release%20Genomic%20

Quality%20Report.pdf).

The analyses include three ancestries: EUR, AFR, and Latino/Admixed American, and

two continuous traits: 1. Height, 2. BMI. We perform GWAS for the two traits using the

unrelated samples for each population, adjusting for PC 1-16, sex, and age, where

some quality control steps are provided in AoU platform.

In AoU platform, the Whole Genome Sequencing (WGS) data and Array data are both

provided, while the WGS data has fewer samples than Array data (98,590 WGS and

165,127 array samples, June 22, 2022, version). The quality control information and

relatedness of samples are only provided within WGS data. For GWAS with respect to

each population, we perform sample level quality control within WGS data. Due to

computation burden, the analyses are conducted using the array SNPs with subjects

that pass the quality control in the WGS data. SNPs with MAF > 0.01 in at least one of

the three populations are kept, while we also restrict analyses to SNPs that are

available on HM3 + MEGA chips. Since the analyses are constrained to Array data, all

analyses are conducted using up to 800K SNPs (**Supplementary Table 4**). We use the

reference data from the 1000 Genomes Project (Phase 3) to estimate the LD for each

population.

The tuning + validation data from UKBB dataset are from EUR and AFR. Latino population is not evaluated on UKBB for the same reason as GLGC analyses. The implementation of the ten different PRSs approaches follow the same steps as the 23andMe data analyses. The adjusted R^2 are adjusted by sex, age and PC1-10.

Figure 1: CT-SLEB Workflow. The method contains three key steps: 1. Two-
dimensional clumping and thresholding method for selecting SNPs (**Figure 1a**); 2.
Empirical-Bayes procedure for incorporating correlation in effect sizes of genetic
variants across populations (**Figure 1b**); 3. Super-learning model for combing the PRSs
derived from the first two steps under different tuning parameters (**Figure 1c**). The
GWAS summary-statistics data are obtained from the training data. The tuning dataset
is used to train the super-learning model. The final prediction performance is evaluated
using an independent validation dataset.

a: Two-dimensional Clumping and Thresholding

b: Empirical-Bayes Estimation of Effect Sizes

c: Super-learning model

Figure 2: Simulation results of various PRSs methods in multi-ancestry settings. Each of the four non-EUR populations has a training sample size of **15,000 (Figure 2a) or 80,000 (Figure 2b)**. For the EUR population, the size of the training sample is set at 100,000. The tuning dataset includes 10,000 samples per population. Prediction R^2 values are reported based on an independent validation dataset with 10,000 subjects per population. Common SNP heritability is assumed to be 0.4 across all populations, and effect-size correlation is assumed to be 0.8 across all pairs of populations. The proportion of causal SNPs varies across 0.01 (top panel), 0.001 (medium panel), 5×10^{-4} (bottom panel), and effect sizes for causal variants are assumed to be related to allele frequency under **<u>a strong negative selection model</u>**. Data are generated based on ~19 million common SNPs across the five populations, but analyses are restricted to ~2.0 million SNPs that are used on Hapmap3 + Multi-Ethnic Genotyping Arrays chip. PolyPred-S+ and PRS-CSx analyses are further restricted to ~1.3 million HM3 SNPs. All approaches are trained using data from the EUR and the target population. CT-SLEB and PRS-CSx are also evaluated using five ancestries data as training data. The red dashed line shows the prediction performance of EUR PRS generated using the single ancestry method (best of CT or LDpred2) in the EUR population.

Figure 3: Comparison of CT-SLEB PRS across different ancestries to single
ancestry EUR PRS in the EUR population. The training sample size for each of the
four non-EUR populations is 15,000, 45000, 80,000, or 100,000. The training sample
size for the EUR population is fixed at 100,000, and PRS performance is evaluated
using single ancestry CT or LDpred2, whichever performs the best in each setting. Two
different genetic architectures are considered: either the common SNP heritability is
fixed (at 0.4) (**Figure 3a and 3b**) or the per-SNP heritability is fixed (**Figure 3c and 3d**)
across the five populations (**Figure 3c and 3d**). The effect-size correlation is assumed
to be 0.8 across all pairs of populations. The effect sizes for causal variants are
assumed to be related to allele frequency under a strong negative selection model. CT-
SLEB uses the summary statistics from all five ancestries.

12
13
14

Figure 4: Prediction performance of CT-SLEB PRS under varying SNP densities.
The analysis of each simulated data based on ~19 million SNPs is restricted to three
different SNP sets: Hapmap3 (~1.3 million SNPs), Hapmap3 + Multi-Ethnic Genotyping
Arrays (~2.0 million SNPs), 1000 Genomes Project (~19 million SNPs). The training
sample size for each of the four non-EUR populations is 15,000 (**Figure 4a**) or 80,000
(**Figure 4b**). The training sample size for the EUR population is fixed at 100,000.
Prediction R^2 values are reported based on independent validation dataset with 10,000
subjects per population. Common SNP heritability is assumed to be 0.4 across all
populations, and effect-size correlation is assumed to be 0.8 across all pairs of
populations. The proportion of causal SNPs proportion vary across 0.01 (top panel),
0.001 (medium panel), 5×10^{-4} (bottom panel), and effect sizes for causal variants are
assumed to be related to allele frequency under a strong negative selection model.

Figure 5: Prediction accuracy of PRSs for heart metabolic disease burden and
height in 23andMe, Inc. datasets. The total sample size for heart metabolic disease
burden and height is, respectively, 2.46 million and 2.93 million for European, 131K and
141K for African American, 375K and 509K for Latino, 110K and 121K for East Asian,
and 29K and 32K for South Asian. The dataset is randomly split into 70%, 20%, 10% for
training, tuning and validation dataset, respectively. The adjusted R^2 values are reported
based on the performance of the PRS in the validation dataset, accounting for PC1-5,
sex and age. The red dashed line represents the prediction performance of EUR PRS
generated using single ancestry method (best of CT or LDpred2) in the EUR population.
Analyses are restricted to ~2.0 million SNPs that are included in Hapmap3, or the Multi-
Ethnic Genotyping Arrays chips array or both. PolyPred-S+ and PRS-CSx analyses are
further restricted to ~1.3 million HM3 SNPs. All approaches are trained using data from
the EUR and the target population. CT-SLEB and PRS-CSx are also evaluated using
training data from five ancestries. From top to bottom panel, two continuous traits are
displayed in the following order: 1. heart metabolic disease burden; 2. height.

Figure 6: Prediction accuracy of five binary traits in 23andMe, Inc datasets. The data are from five populations: European (averaged N ≈ 2.37 million), African American (averaged N ≈ 109K), Latino (averaged N ≈ 401K), East Asian (averaged N≈ 86K), South Asian (averaged N≈ 24K). The datasets are randomly split into 70%, 20%, 10% for training, tuning and validation dataset, respectively. The adjusted AUC values are reported based on the validation dataset accounting for PC1-5, sex and age. The red dashed line represents the prediction performance of EUR PRS generated using single ancestry method (best of CT or LDpred2) in the EUR population. Analyses are restricted to ~2.0 million SNPs that are included in Hapmap3, or the Multi-Ethnic Genotyping Arrays chips array or both. PolyPred-S+ and PRS-CSx analyses are further restricted to ~1.3 million HM3 SNPs as implemented in the provided software. All approaches are trained using data from the EUR and the target population. CT-SLEB and PRS-CSx are also evaluated using training data from five ancestries. From top to bottom panel, five binary traits are displayed in the following order: 1.any cardiovascular disease (any CVD); 2. depression; 3. migraine diagnosis; 4. sing back musical note (SBMN); 5. morning person.

Figure 7: Prediction accuracy of four blood lipids traits from the Global Lipids Genetics Consortium. We used the GWAS summary statistics from five populations as the training data: European (N ≈ 931K), African (primarily African American, N ≈ 93K), Latino (N≈ 50K), East Asian (N ≈ 146K), South Asian (N ≈ 34K). The tuning and validation dataset are from UK Biobank data with three different ancestries African (N = 9,042), East Asian (N = 2,009), South Asian (N = 10,615). The tuning and validation are split with half and half. The adjusted R^2 values are reported based on the performance of the PRS in the validation dataset, while accounting for PC1-10, sex and age. The red dashed line represents the prediction performance of EUR PRS generated using single ancestry method (best of CT or LDpred2) in the EUR population. Analyses are restricted to ~2.0 million SNPs that are included in Hapmap3, or the Multi-Ethnic Genotyping Arrays chips array or both. PolyPred-S+ and PRS-CSx analyses are further restricted to ~1.3 million HM3 SNPs as has been implemented in the provided software. All approaches are trained using data from the EUR and the target population. CT-SLEB and PRS-CSx are also evaluated using training data from five ancestries. From top to bottom panel, four traits are displayed in the following order: 1. high-density lipoprotein cholesterol (HDL), 2. low-density lipoprotein cholesterol (LDL), 3. log triglycerides (logTG) and 4. total cholesterol (TC).

Figure 8: Prediction accuracy of two traits from All of Us dataset. We used the
GWAS summary statistics from three populations as the training data: European (N ≈
48K), African (N ≈ 22K), Latino (averaged N ≈ 15K). The tuning and validation dataset
are from UK Biobank data with African (N = 9,042). The tuning and validation are split
with half and half. The adjusted R^2 values are reported based on the performance of the
PRS in the validation dataset, while accounting for PC1-10, sex and age. The red
dashed line represents the prediction performance of EUR PRS generated using single
ancestry method (best of CT or LDpred2) in the European population. Analyses are
restricted to around 800K SNPs that are genotyped in All of Us dataset for different
ancestries. All approaches are trained using data from the European and African
population. CT-SLEB and PRS-CSx are further evaluated using training data from three
ancestries: African, European and Latino. From top to bottom panel, two traits are
displayed in the following order: 1. body mass index (BMI), and 2. height.

14
15