Distinct bile salt hydrolase substrate preferences dictate *C. difficile* pathogenesis

One sentence summary: Bile salt hydrolase activity inhibits *C. difficile* by shaping the

conventional and non-canonical conjugated bile acid pools

Matthew H. Foley[1,2]†, Morgan E. Walker[3]†, Allison K. Stewart[4], Sarah O'Flaherty[2], Emily C.

Gentry[5,6], Garrison Allen[1], Shakshi Patel[3], Meichen Pan[2], Violet V. Beaty[3], Molly E. Vanhoy[1],

Michael K. Dougherty[7], Sarah K. McGill[7], Ajay Gulati[7,8], Pieter C. Dorrestein[5,6], Erin S. Baker[4],

Matthew R. Redinbo[3,9]*, Rodolphe Barrangou[2]*, Casey M. Theriot[1]*.

1. Department of Pathobiology and Population Health, College of Veterinary Medicine, North

Carolina State University, Raleigh, NC, USA.

2. Department of Food, Bioprocessing and Nutrition Sciences, North Carolina State University,

Raleigh, NC, USA.

3. Department of Chemistry, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA.

4. Department of Chemistry, North Carolina State University, Raleigh, NC, USA

5. Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California San Diego,

La Jolla, San Diego, CA, USA

6. Collaborative Mass Spectrometry Innovation Center, Skaggs School of Pharmacy and

Pharmaceutical Sciences, University of California San Diego, La Jolla, CA, USA

7. Department of Medicine, Division of Gastroenterology and Hepatology, University of North

Carolina at Chapel Hill, Chapel Hill, NC, USA

8. Department of Pathology and Laboratory Medicine, University of North Carolina at Chapel

Hill, Chapel Hill, NC, USA

9. Departments of Biochemistry & Biophysics, and Microbiology & Immunology, and the

Integrated Program in Biological and Genome Sciences, University of North Carolina at Chapel

Hill, Chapel Hill, NC, USA

† Authors contributed equally to this work

* Corresponding Authors Matthew R. Redinbo redinbo@unc.edu; Rodolphe Barrangou,

rbarran@ncsu.edu; Casey M. Theriot, cmtherio@ncsu.edu

Classification

Biological Sciences, Microbiology.

Keywords

bile salt hydrolase, bile acid, live biotherapeutic, *Lactobacillus*, *Clostridioides difficile*

Summary paragraph: Bile acids (BAs) mediate the crosstalk between human and microbial

cells and influence intestinal diseases including *Clostridioides difficile* infection (CDI). While

bile salt hydrolases (BSHs) shape the BA pool by deconjugating conjugated BAs, the basis for

their substrate preferences and impact on *C. difficile* remain elusive. Here, we survey the

diversity of *Lactobacillus* BSHs and unravel the structural basis of their substrate preference. We

show that leveraging BSH activity and specificity is an effective strategy to prevent *C. difficile*

growth in clinically relevant CDI models. A range of non-canonical conjugated BAs is also

identified, comprising unique BSH substrates that also inhibit *C. difficile* spore germination.

These findings establish BSHs as intestinal enzymes essential to BA homeostasis and

colonization resistance against *C. difficile*.

Introduction

Bile acids (BAs) are host-synthesized and microbial-derived metabolites that support intestinal health and homeostasis by providing a scaffold for host-microbiome crosstalk and adaptation ([1, 2]). Host-encoded BA receptors recognize BAs as signaling molecules to regulate key aspects of host immunity ([3]), metabolism ([4]), and circadian rhythms ([5, 6]). BAs also shape the microbiota, with specific taxa being inherently more resistant to bile stress by harboring mechanisms to survive BA exposure ([7, 8]). The influence of BAs on the microbiota composition is reciprocated by the microbial transformation of BAs to unlock the chemical complexity present in the intestinal BA pool ([9]). The resulting mosaicism of the BA pool is a hallmark of host health and is dependent on the collective activity of microbiome-encoded BA altering enzymes ([10]).

One such enzyme, bile salt hydrolase (BSH), comprises a family of microbial enzymes that catalyze a critical step in BA biosynthesis ([11]). Primary conjugated BAs synthesized in the liver, such as the tauro- and glyco- conjugates of cholic acid (TCA, GCA), are deconjugated by BSHs to generate cholic acid (CA) (**Fig. 1A**; BA names and abbreviations are summarized in **Fig. S1**). This vanguard catalytic step acts as a "gatekeeper'' for subsequent BA transformations that turn primary BAs into secondary BAs (i.e. deoxycholic acid -DCA- formed from the 7α-dehydroxylation of CA) ([1, 11]). Dysregulation of BA biosynthesis and BSH activity has been directly implicated in obesity ([12]), cancer ([13]), inflammatory bowel disease ([3]), and colonization resistance against pathogens including *Clostridioides difficile* ([14, 15]). Strategically managing BSH activity to rationally toggle BA metabolism and judiciously alter intestinal pathobiology presents an emerging therapeutic approach. Additionally, the recent discovery of non-canonical conjugated BAs has

Figure 1.

considerably increased the complexity of the BA pool and altered the paradigm of BSH-BA

biology. In particular, BSHs do not appear to process these non-canonical conjugated BAs *in*

vitro, indicating that renewed and expanded investigations of BSH-substrate relationships are

warranted (**Fig. 1A)** (*16-18*).

Here, we examine the phylogenetic, biochemical, structural diversity and

specificity of BSHs in *Lactobacillus*, a prototypical genus for studying BA deconjugation. We

define how the conjugated amino acid moiety drives substrate preferences in distinct BSHs and

how those preferences can be leveraged to prevent *C. difficile* spore germination and

colonization in pre-clinical models. We further reveal a new function for BSHs, the ability to

process an expanded set of substrates including non-canonical conjugated BAs.

Surveying *Lactobacillus* BSH Diversity

To map the range of BSHs harbored by *Lactobacillus*, 3,712 genomes from 274

lactobacilli were assembled to comprehensively define the diversity of the enzyme within this

key intestinal genus (**Table S1**) (*19*). Highly similar BSH sequence (>95% amino acid identity)

were grouped into 84 clusters and a representative sequence from each cluster was used to

construct a phylogenetic tree. Nearly all BSH clusters are identified in vertebrate-adapted species

and ~40% were human-associated, underscoring their role in host-microbiota symbiosis (**Fig.**

1B, Table S1).

To move toward a functional understanding of BSH activity *in vivo*, 17 diverse

BSHs were selected for heterologous expression and purification **(Table S2, Fig. S2)**. BSH

activity was screened on a panel of conjugated primary and secondary BAs to comprehensively

determine BSH substrate preferences **(Fig. 1C)**. BSHs display a well-defined bias for either

glycine or taurine-conjugated BAs. Glycine-preferences dominate the BSHs tested, potentially

reflecting the increased abundance of glycine-conjugated BAs among vertebrates (*20*), whereas

taurine-specificity is restricted to only a few related BSHs. The glycine-preferring clade

containing LaciBSHa, LcriBSHa, and LcriBSHb exhibits the highest activity. Furthermore, the

taurine-preferring LaviBSH, LjohBSHc, and LgasBSHa (corresponding to the clusters

arrafinosis, johnsonii_1, and gasseri_1, respectively) enzymes are all present within a single

major clade while the LsalBSH (cluster salivarius_2) stands alone in a distantly related branch.

This suggests that taurine-preference may have evolved twice within *Lactobacillus*. Overall,

BSH specialization for glycine or taurine illustrates how lactobacilli may have tailored their

metabolism to manage BA exposure.

Structural Basis for BSH Substrate Preference

To dissect the molecular basis of BSH substrate specificity, we determined the

crystal structures of six of the *Lactobacillus* BSHs examined above, four glycine-preferring

(LgasBSHb, LaciBSHa, LingBSH, LreuBSH) and two taurine-preferring BSHs (LgasBSHa,

LjohBSHc; **Fig. 1D, S3A, Table S3**). While most structures were unliganded, LgasBSHa was

resolved in complexes with taurine or taurine plus CDCA. All BSHs form tetramers and each

monomer exhibits the core four-layered αββα fold typical of the Ntn-hydrolase family (*21*), with

RMSD values ranging from 0.5 to 2.0 Å over equivalent Cα positions (**Fig. 2A, S3B**). Within

this standard core structure, we noted that each monomer contains a loop donated from a

neighboring monomer within the tetramer (**Fig. 2A**). Importantly, sequence analysis revealed

that taurine-preferring enzymes contained an S-R-G/S motif in this loop region, while the

glycine-preferring enzymes encoded a G-V/T-G motif (**Fig. 2B, Fig. S4**) (*22*). These particular

amino acids were also placed directly in each enzyme's active site within the functional BSH

tetramers (**Fig. 2C, 2D**). Thus, we hypothesized that these 3-residue motifs play important roles

in BSH substrate preferences.

Figure2.

121   To test this, the 210-GVG-212 region of taurine-preferring LgasBSHa was replaced

with SRS, while the 207-SRS-209 region of glycine-preferring LgasBSHb was replaced with

GVG, and catalytic efficiencies with conjugates of CA were examined. The LgasBSHa-SRS

variant enzyme exhibited significantly reduced catalytic efficiency with TCA, the wild-type

enzyme's preferred substrate, while also reducing its processing of GCA (**Fig. 2E**). Similarly, the

LgasBSHb-GVG variant showed significantly lower catalytic efficiency with its preferred GCA

substrate, and a trend toward increased activity with the non-standard substrate TCA (**Fig. 2F**).

Taken together, these functional data support the structure-guided hypothesis that short sequence

motifs play important roles in defining the marked glycine *vs.* taurine substrate preferences

exhibited by *Lactobacillus* BSHs, reflecting specialization.

Structurally, the presence of R208 in LgasBSHb appears to block the processing of

taurine-conjugated BAs. When the LgasBSHa and LgasBSHb active sites are overlaid, the

distance between LgasBSHb R208 and taurine from the LgasBSHa structure is 2.3 Å, which

would generate a steric clash (**Fig. 2G**). In contrast, V211 in LgasBSHa is 3.6 Å away and would

not occlude the binding of taurine-conjugated BAs (**Fig. 2G**). Furthermore, R208 in LgasBSHb

forms a hydrogen bond with E21 and that interaction creates a more enclosed active site than that

seen in taurine preferring BSHs (Fig. 2G). In support of this conclusion, the active sites of the

four additional BSH structures resolved here (LjohBSHc, LaciBSHa, LingBSH, LreuBSH)

reveal analogous architectures (**Fig. 1D, 2H, I**). Importantly, when a glycine conjugate was

modelled into the active site of the glycine-preferring BSHs, the distance between R208 and

glycine increased to 3.1 Å, indicating that this smaller substrate can be accommodated with this

clade of enzymes (**Fig. 2I**). Together, these data define three-residue motifs as crucial for the

glycine vs. taurine conjugate preferences demonstrated by *Lactobacillus* BSHs.

Bile Acids Inhibit *C. difficile* in a Conjugation-Dependent Manner

C. difficile infection (CDI) is a significant public health problem that can be

difficult to treat with conventional antibiotics (*23*). Even after successful treatment with

vancomycin, there remains a high recurrence rate (~30%) (*24*), making a fecal microbiota

transplant (FMT) the last line of treatment (*25*). Although FMTs are a lifeline for recurrent CDI

(rCDI) patients, the long-term consequences are unknown at this time, making a targeted

therapeutic approach urgently needed. Susceptibility to CDI is predominantly driven by

antibiotic-driven perturbations to the intestinal microbiota, which results in a loss of colonization

resistance (*26, 27*). Strong evidence suggests that microbial BA metabolism is an important

mechanism of colonization resistance against *C. difficile* (*28*), as antibiotic usage causes depleted

BSHs and increases conjugated BAs (*14*).

155   BAs govern fundamental aspects of *C. difficile*'s pathogenesis: TCA acts as a

germinant for *C. difficile* spores (*29*), whereas CDCA can inhibit germination and growth (*30*).

Most studies have focused on the production of deconjugated secondary BAs as inhibitors of *C.*

difficile pathogenesis while few have examined the contributions of BSH activity (*31*). The

therapeutic potential of BSHs has been proposed, but gaps in our knowledge of their pleiotropic

effects on intestinal biology and their exact mechanism of action have hampered their

development (*32, 33*).

162   To determine how BA deconjugation impacts *C. difficile*, we measured the ability

of common BAs to inhibit germination in the presence of TCA (**Fig. 3A-C**). Though all BAs

tested inhibit germination compared to TCA alone, deconjugated BAs are significantly more

inhibitory compared to their taurine-conjugated versions, whereas they are equally or only mildly

more inhibitory relative to their glycine-conjugated variants (**Fig 3B**). We also assayed the

ability of BAs to inhibit *C. difficile* growth, as we hypothesized that the conjugation would

determine BA toxicity (**Fig. 3C**). Taurine-conjugates inhibit growth less while glycine-

conjugates and deconjugated BAs are much more toxic. Moreover, taurine-conjugated BAs are

significantly less disruptive to *C. difficile*'s membrane integrity compared to glycine-conjugated

and deconjugated BAs (**Fig. S5**). Overall, using BSHs to deconjugate taurine-conjugated BAs

may be a potent strategy to restrict *C. difficile* in the intestinal tract.

Figure 3.

175   To determine whether BSHs are sufficient to inhibit *C. difficile* and provide an

alternative approach to using antibiotics or FMT, stool from three rCDI patients was collected

prior to them receiving an FMT (**Fig 3D**). Unlike mice, FMT recipients have a complex mixture

of taurine and glycine-conjugated primary BAs (**Fig S6**). To test whether BSH substrate

preference could impact *C. difficile* growth, pre-FMT stool was used to culture *C. difficile* and

samples were treated with PBS, a cocktail of taurine-preferring BSHs (T-BSH), glycine-

preferring BSHs (G-BSH), or a broadly-acting combination (G&T-BSH) **(Fig. 3E, S7)**. Alone,

the BSH cocktails, glycine, or taurine have no impact on *C. difficile* growth (**Fig. S8**). When

Recipient 1's stool is treated with the T-BSH and G&T-BSH cocktails, *C. difficile* growth is

significantly inhibited at 8 h. Recipient 2's stool doesn't alter *C. difficile* growth when treated

with any BSH cocktail, though the T-BSH and G&T-BSH cocktails slightly suppress growth at

24 h. Recipient 3's stool robustly suppresses *C. difficile* growth at 8 h using all BSH cocktails,

but only the T-BSH and G&T-BSH cocktails maintain that suppression at 24 h.

188   Targeted BA metabolomics was used to assess how the BSH cocktails alter the pre-

FMT BA pool and explain the individual variability between recipients (**Fig 3F, S9**). Recipients

1 and 3 display an increase in CA and CDCA primarily in response to the T-BSH and G&T-BSH

cocktails. Recipient 3's high levels of conjugated BAs prior to BSH treatment explains the BSH

cocktail's robust inhibition (**Fig. S6, 9**), Furthermore, Recipient 2 does not display an increase in

deconjugated BAs from any BSH treatment, likely due to the overall low concentration of

conjugated BAs prior to BSH treatment (**Fig S6**), thereby explaining why no *C. difficile*

inhibition was observed in this sample.

BSH Activity Inhibits *C. difficile* Spore Germination and Growth

197   To interrogate the role of BSHs in colonization resistance against *C. difficile,* we

leveraged an antibiotic treated mouse model of CDI, since mice typically have a relatively rich

amount of taurine conjugated bile acids. *C. difficile* initiates its infectious cycle when spores

arrive and germinate in the small intestine, and then begin to grow and progress to the large

intestine where the majority of toxin production and disease occurs. To examine the contribution

of BSH activity to colonization resistance along the intestinal tract, we collected murine small

intestinal and cecal contents from cefoperazone-treated mice at several times post-antibiotic

treatment and supplemented them with PBS or the G&T-BSH cocktail (**Fig. 4A**). The treated

contents were then inoculated with 10^5 spores or CFUs/mL to measure *ex vivo* spore germination

and growth, respectively. Remarkably, the addition of the BSH cocktail significantly inhibits *C.*

difficile spore germination in the small intestinal contents of all mice and suppresses growth at 8

and 24 h (**Fig. 4B, C, S10**). Inhibition of growth is more subtle in the cecum, likely due to the

lower concentration of conjugated BAs in this region of the intestinal tract (**Fig. 4D, S11**).

Targeted metabolomics was used to corroborate the BSH cocktail's effect on the murine BA

pool. Small intestinal samples from day 0 treated with the BSH cocktail display a significant

increase in deconjugated BAs (CA, CDCA, βMCA, and UDCA) and a decrease in the taurine-

conjugated forms of the same BAs. In the cecum, only CA was significantly deconjugated (**Fig.**

4D, S11). By day 7 post antibiotic, some mice displayed higher amounts of deconjugated BAs,

independent of BSH treatment, coinciding with the return of colonization resistance after

antibiotic treatment.

Figure 4

219   Aside from the increase in deconjugated BAs, the most striking difference in the

BA pool is the unexpected increase in non-canonical conjugated BAs in the BSH-treated small

intestinal samples (**Fig 4D, S11**). An array of intestinal bacteria can catalyze the conjugation of

BAs to a variety of amino acids through an unknown mechanism (**Fig. 1A**) (*16, 17*). Despite

their widespread synthesis, there is no evidence at this time that shows that the non-canonical

conjugated BAs Ile/Leu-CA, Phe-CA, and Tyr-CA can be deconjugated (*16*). Ile-CA, Phe-CA,

and Tyr-CA were among the BAs detected in the small intestinal samples. Given the increased

CA and βMCA in the BSH-treated samples, these BAs are likely re-conjugated by the small

intestinal microbiota to produce the non-canonical BAs observed. While these BAs did not affect

C. difficile growth or induce spore germination (**Fig. S12A, B**), Tyr-CA and Phe-CA are able to

inhibit spore germination, suggesting for the first time a role for the non-canonical conjugated

BAs in colonization resistance against *C. difficile* (**Fig. 4E, S12C**).

BSHs Process Non-Canonical Bile Acids Conjugates

232   To determine whether BSHs are able to process non-canonical conjugated BAs, we

screened a larger range of BSHs and BAs (**Fig. S13**). While activity with these non-canonical

conjugates was low compared to the canonical conjugations, some BSHs, like LingBSH and

LreuBSH, exhibit appreciable activity on non-canonical BAs. To understand the recalcitrance of

non-canonical conjugated BAs, we broadly monitored the deconjugation of 5 BSHs (LingBSH,

LgasBSHa, LgasBSHb, LjohBSHc, and LreuBSH) with assorted preferences (**Fig. S14A**). BSHs

were incubated with non-canonical conjugated BAs pools (AA-CA, AA-CDCA, AA-DCA, and

AA- βMCA) that consist of one BA (CA, CDCA, DCA, or βMCA) conjugated to 22 different

amino acids, excluding taurine and glycine (*18*), and initial BA deconjugation was quantified

using LC-IMS-MS (**Fig. S14B**). LingBSH displays exceptional activity, suggesting that some

BSHs may be uniquely adapted to process this expanded repertoire of BAs (**Fig. 4F, 14B, S15**).

Ala and Ser-conjugated BAs are efficiently processed, and these observations were validated

using purified Ala-CA and Ser-CA demonstrating that their susceptibility to deconjugation is

similar to GCA and TCA (**Fig. S13**). Ala and Ser are the smallest amino acids next to Gly,

suggesting that the steric constraints of the active site could dictate non-canonical conjugate

deconjugation (**Fig. S15, S16**). Notably, amine-containing amino acids such as Lys, Gln, and

Asn are enriched and among the most efficiently processed conjugates (**Fig. S16**).

249   We also examine whether non-canonical conjugated BA processing would occur in

BSH-expressing *L. gasseri* cells. Wild-type, ΔbshA, ΔbshB, and ΔbshAB *L. gasseri* was grown

and reacted with the AA-CDCA pool and supernatants were used to detect deconjugation.

Δ*bshAB* displayed some deconjugation, possibly due to cells sequestering BAs (**Fig. S14C**).

Individual BSH activities and substrate preferences were comparably maintained *in vivo*,

demonstrating that BSH-expressing bacteria are capable of non-canonical BA deconjugation, and

that they do so based on the enzymatic preferences of the BSHs they encode.

Discussion

There is a growing appreciation for the roles that BSH substrate preferences play in

the management of disease (*6, 8*). This work stresses the distinction between the deconjugation

of glycine and taurine-conjugated BAs, the latter of which has been shown to play a role in the

establishment of colonization resistance against *Klebsiella pneumoniae* and the development of

colorectal cancer (*34, 35*). Our analysis of *C. difficile*-BA sensitivities motivated the use of

glycine and taurine-preferring *Lactobacillus* BSHs to optimally inhibit *C. difficile ex vivo* (**Fig.**

3D). Thus, leveraging BA metabolism to restore colonization resistance may be an alternative

therapy that avoids antibiotic damage, the unspecified risks of FMTs, and the variability of live

biotherapeutics.

The extensive variety of recently identified non-canonical conjugated BAs expands

the complexity of the BSH-BA pool and reinforces BSHs as key BA-altering enzymes (*16, 17*).

BSH-generated deconjugated BAs fuel the production of non-canonical conjugated BAs (**Fig.**

4D). While the mechanism and purpose of this re-conjugation is yet to be elucidated, our

observation that non-canonical conjugations can inhibit *C. difficile* spore germination introduces

new mechanistic avenues by which BSH activity can be used to restore colonization resistance

(**Fig. 4E**). We also demonstrate that BSHs can variably act on these BAs to support a model

where BSHs directly shape the non-canonical conjugated BA pool by simultaneously sustaining

and curtailing it (**Fig. 4D, S14**). Given the broad impacts BSH can have on the overall BA pool,

purposefully augmenting the specificity or activity of BSHs could support their potential use as a

precision modality to steer BA metabolism *in vivo*.

278   BSHs perform the important role of BA deconjugation, and perturbations to this

process are implicated in several diseases including CDI. *C. difficile* pathogenesis is

exceptionally responsive to alterations in BSH activity, thereby establishing these enzymes as a

potential targeted approach to restore colonization resistance. The preferences and efficiency by

which BSHs recognize conjugated BAs defines the resulting BA pool, and their ability to

variably process the expansive catalogue of non-canonical conjugated BAs is a newly prescribed

function of BSHs. In the future, BSHs may be leveraged to rationally manipulate the chemical

complexity of the BA pool toward addressing many states of intestinal disease.

References

1.   M. H. Foley, S. O'Flaherty, R. Barrangou, C. M. Theriot, Bile salt hydrolases:

Gatekeepers of bile acid metabolism and host-microbiome crosstalk in the

gastrointestinal tract. *PLoS Pathog* **15**, e1007581 (2019).

2.   C. L. Gentile, T. L. Weir, The gut microbiota at the intersection of diet and human health.

Science **362**, 776-780 (2018).

3.   A. Lavelle, H. Sokol, Gut microbiota-derived metabolites as key actors in inflammatory

bowel disease. *Nat Rev Gastroenterol Hepatol* **17**, 223-237 (2020).

4.   S. N. Chaudhari *et al.*, Bariatric surgery reveals a gut-restricted TGR5 agonist with anti-

diabetic effects. *Nat Chem Biol* **17**, 20-29 (2021).

5.   K. Govindarajan *et al.*, Unconjugated bile acids influence expression of circadian genes:

A potential mechanism for microbe-host crosstalk. *PLoS One* **11**, e0167319 (2016).

6.   L. Yao *et al.*, A selective gut bacterial bile salt hydrolase alters host metabolism. *Elife* **7**,

(2018).

7.   K. B. Islam *et al.*, Bile acid is a host factor that regulates the composition of the cecal

microbiota in rats. *Gastroenterology* **141**, 1773-1781 (2011).

8.   M. H. Foley *et al.*, *Lactobacillus* bile salt hydrolase substrate specificity governs bacterial

fitness and host colonization. *Proc Natl Acad Sci U S A* **118**, (2021).

9.   J. M. Ridlon, S. C. Harris, S. Bhowmik, D. J. Kang, P. B. Hylemon, Consequences of bile

salt biotransformations by intestinal bacteria. *Gut Microbes* **7**, 22-39 (2016).

10.   B. Jia, D. Park, Y. Hahn, C. O. Jeon, Metagenomic analysis of the human microbiome

reveals the association between the abundance of gut bile salt hydrolases and host health.

Gut Microbes **11**, 1300-1313 (2020).

11.　J. M. Ridlon, D. J. Kang, P. B. Hylemon, Bile salt biotransformations by human intestinal
bacteria. *J Lipid Res* **47**, 241-259 (2006).

12.　S. A. Joyce *et al.*, Regulation of host weight gain and lipid metabolism by bacterial bile
acid modification in the gut. *Proc Natl Acad Sci U S A* **111**, 7421-7426 (2014).

13.　C. Ma *et al.*, Gut microbiome-mediated bile acid metabolism regulates liver cancer via
NKT cells. *Science* **360**, (2018).

14.　C. M. Theriot *et al.*, Antibiotic-induced shifts in the mouse gut microbiome and
metabolome increase susceptibility to *Clostridium difficile* infection. *Nat Commun* **5**,
3114 (2014).

15.　B. H. Mullish *et al.*, Microbial bile salt hydrolases mediate the efficacy of faecal
microbiota transplant in the treatment of recurrent *Clostridioides difficile* infection. *Gut*
68, 1791-1800 (2019).

16.　R. A. Quinn *et al.*, Global chemical effects of the microbiome include new bile-acid
conjugations. *Nature* **579**, 123-129 (2020).

17.　L. N. Lucas *et al.*, Dominant bacterial phyla from the human gut show widespread ability
to transform and conjugate bile acids. *mSystems*, e0080521 (2021).

18.　E. Gentry *et al.*, https://www.researchsquare.com/article/rs-820302/v1. *Nature Portfolio*,
(2022).

19.　S. O'Flaherty, A. Briner Crawley, C. M. Theriot, R. Barrangou, The *Lactobacillus* bile
salt hydrolase repertoire reveals niche-specific adaptation. *mSphere* **3**, (2018).

20.　G. A. Nagana Gowda, N. Shanaiah, A. Cooper, M. Maluccio, D. Raftery, Bile acids
conjugation in human bile is not random: new insights from (1)H-NMR spectroscopy at
800 MHz. *Lipids* **44**, 527-535 (2009).

21.   F. Xu *et al.*, The complex structure of bile salt hydrolase from *Lactobacillus salivarius*

333   reveals the structural basis of substrate specificity. *Sci Rep* **9**, 12438 (2019).

334   22.   F. Madeira *et al.*, The EMBL-EBI search and sequence analysis tools APIs in 2019.

335   *Nucleic Acids Res* **47**, W636-W641 (2019).

336   23.   C. Slimings, T. V. Riley, Antibiotics and hospital-acquired *Clostridium difficile* infection:

337   update of systematic review and meta-analysis. *J Antimicrob Chemother* **69**, 881-891

338   (2014).

339   24.   O. A. Cornely, M. A. Miller, T. J. Louie, D. W. Crook, S. L. Gorbach, Treatment of first

340   recurrence of *Clostridium difficile* infection: fidaxomicin versus vancomycin. *Clin Infect*

341   *Dis* **55 Suppl 2**, S154-161 (2012).

342   25.   C. R. Kelly *et al.*, Effect of fecal microbiota transplantation on recurrence in multiply

343   recurrent *Clostridium difficile* infection: A randomized trial. *Ann Intern Med* **165**, 609-

344   616 (2016).

345   26.   R. C. Owens, Jr., C. J. Donskey, R. P. Gaynes, V. G. Loo, C. A. Muto, Antimicrobial-

346   associated risk factors for *Clostridium difficile* infection. *Clin Infect Dis* **46 Suppl 1**, S19-

347   31 (2008).

348   27.   R. A. Britton, V. B. Young, Role of the intestinal microbiota in resistance to colonization

349   by *Clostridium difficile*. *Gastroenterology* **146**, 1547-1553 (2014).

350   28.   A. D. Reed, C. M. Theriot, Contribution of nhibitory metabolites and competition for

351   nutrients to colonization resistance against *Clostridioides difficile* by Commensal

352   Clostridium. *Microorganisms* **9**, (2021).

353   29.   J. A. Sorg, A. L. Sonenshein, Bile salts and glycine as cogerminants for *Clostridium*

354   *difficile* spores. *J Bacteriol* **190**, 2505-2512 (2008).

30.   J. A. Sorg, A. L. Sonenshein, Chenodeoxycholate is an inhibitor of *Clostridium difficile*

spore germination. *J Bacteriol* **191**, 1115-1117 (2009).

31.   R. Thanissery, J. A. Winston, C. M. Theriot, Inhibition of spore germination, growth, and

toxin activity of clinically relevant *C. difficile* strains by gut microbiota derived

secondary bile acids. *Anaerobe* **45**, 86-100 (2017).

32.   M. L. Jones, C. Tomaro-Duchesneau, C. J. Martoni, S. Prakash, Cholesterol lowering

with bile salt hydrolase-active probiotic bacteria, mechanism of action, clinical evidence,

and future direction for heart health applications. *Expert Opin Biol Ther* **13**, 631-642

(2013).

33.   S. A. Joyce, F. Shanahan, C. Hill, C. G. Gahan, Bacterial bile salt hydrolase in host

metabolism: Potential for influencing gastrointestinal microbe-host crosstalk. *Gut*

Microbes **5**, 669-674 (2014).

34.   A. Stacy *et al.*, Infection trains the host for microbiota-enhanced resistance to pathogens.

Cell **184**, 615-627 e617 (2021).

35.   J. M. Ridlon, P. G. Wolf, H. R. Gaskins, Taurocholic acid metabolism by gut microbes

and colon cancer. *Gut Microbes* **7**, 201-215 (2016).

36.   R. M. Duar *et al.*, Lifestyles in transition: evolution and natural history of the genus

Lactobacillus. *FEMS Microbiol Rev* **41**, S27-S48 (2017).

Acknowledgements: This work was performed in part by the Molecular Education, Technology

and Research Innovation Center (METRIC) at NC State University, which is supported by the

State of North Carolina.

Funding: We thank IFF for supporting this study. MHF was supported by the University of

North Carolina Center for Gastrointestinal Biology and Disease postdoctoral fellowship training

grant T32DK07737. MEW was supported through the NIH training grant T32GM008570 and

NSF DGE-1650116. This study was also supported by the NIGMS R35 GM119438, and R01

GM135218 and GM137286; NIEHS P30 ES025128, P42 ES027704, and P42 ES031009; and the

Environmental Protection Agency STAR RD 84003201 award. The FMT study was supported

by the NCTraCS Translational and Clinical Sciences Institute Pilot Grant UNCSUR11609. MD

was supported by NIH training grant T32 DK007634.

Author contributions: MHF, MEW, SOF, ECP, MRR, RB, CMT designed the research. MHF,

MEW, AKS, SOF, ECG, GA, SP, MP, VVB, MEV, and ESB performed, analyzed, and

interpreted experiments. SOF and MP performed bioinformatic analysis of BSHs. MHF and GA

performed all BSH specific activity assays and *C. difficile* experiments. MEW, SP, and VVB

performed crystallographic work. MD, SKM, and AJ collected patient samples. AKS performed

metabolomic analyses. ECG synthesized non-canonical bile acids. MF and MEW wrote the

manuscript which all authors edited.

Competing interests: MRR is a Founder of Symberix, Inc., which is developing microbiome-

targeted therapeutics. MRR is also the recipient of research funding from Merck and Lilly,

although those funds were not used in this project. PCD is a scientific advisor to Cybele and is a

co-founder and scientific founder of Ometa and Enveda, with prior approval by UC-San Diego.

CMT consults for Vedanta Biosciences, Inc. and Summit Therapeutics. CMT is a founder of

CRISPR Biotechnologies. RB is a founder of Ancilia Bioscience and CRISPR Biotechnologies.

MHF, SOF, RB and CMT are inventors on a patent application related to the use of

Lactobacillus BSH enzymes.

Data and materials availability: All data associated with this study are available in the main

text or the Supplementary Materials. All LC-IMS-MS data are publicly available on Panorama

under the Panorama dashboard of the Baker Lab-NCSU within the "0821 BSH Assessment",

"1121 Mouse Germination and Growth Ex-Vivo" and the "1121 Theriot FMT Ex-Vivo Growth"

projects."

List of Supplementary materials:

Materials and Methods

Fig S1 to S17

Table S1 to S3

Figure legends

Figure 1. *Lactobacillus* BSHs have distinct substrate preferences.

(A) Overview of BA metabolism using cholic acid (CA) as an example. BAs are synthesized and

conjugated with a taurine or glycine (T/GCA) by the liver enzyme Baat. Conjugated BAs are

deconjugated by BSHs, allowing transformations such as 7α-dehydroxylation encoded by the *bai*

operon to generate secondary BAs like DCA. Deconjugated BAs can be re-conjugated to

produce non-canonical conjugated BAs like Tyr-CA. (B) The *Lactobacillus* BSH phylogenetic

tree was assembled from 84 BSH clusters. Major clades are distinguished by branch color. Each

cluster name indicates the BSH-encoding species with the inner metadata layers indicating

lifestyle and isolation source (*36*). The outer metadata layer indicates the BSHs cloned for

characterization with diamonds indicating BSH crystal structures presented here. (C) Heatmap of

BSH activity across conjugated BAs. The clade highlighted in blue displays preference for

taurine-conjugated BAs. Values represent mean specific activity (n = 3-4). Phylogenetic

relationships between BSHs are based on amino acid sequences. (D) Superposition of the

taurine- and glycine-preferring BSH enzyme tetramer crystal structures presented here.

Figure 2. BSH loop region dictates enzyme substrate preference.

(A) Superimposed monomers of the six BSH monomer structures presented here highlighting the

substrate preference loop and colored as indicated in panel B. (B) Sequence alignment of the

loop region of *Lactobacillus* BSHs. Resolved structures are colored as indicated, with a constant

coloring scheme throughout the figure. Taurine-preferring enzymes exhibit a G-V/T-G motif,

while glycine-preferring enzymes maintain a S-R-G/S motif. (C) The LgasBSHa tetramer with

the active site shown in the inset, with the catalytic cysteine at position 2, the bound taurine and

CDCA compounds, and the GVG swapped into the active site from the neighboring monomer

highlighted. (D) The LgasBSHb tetramer with the active site shown in the inset, with the

catalytic cysteine at position 2 and the SRS swapped into the active site from the neighboring

monomer highlighted. The taurine and CDCA ligands from the LgasBSHa structure are also

rendered transparently. (E, F) Catalytic efficiency for wild-type and mutant forms of LgasBSHa

and LgasBSHb with GCA and TCA. Bars are the average of n=3 ± standard deviation.

Significant differences were tested by two-way ANOVA with Tukey's multiple comparisons test

(ns not significant, ***$p < 0.001$, ****$p < 0.0001$). (G) Superposition of the active sites of

LgasBSHa and LgasBSHb showing that taurine is accommodated by Val211 of LgasBSHa but

clashes at 2.3 Å with the side chain of Arg208 in LgasBSHb, which is held in place via a salt

bridge with Glu21. (H, I) Superpositions of the active sites of the two taurine-preferring or the

four glycine-preferring BSH structures presented here reveals consistent active site architectures

that accommodate taurine or glycine, respectively.

Figure 3. Deconjugated BAs and BSH activity can inhibit *C. difficile*. (A) Schematic of *in*

vitro BA-dependent inhibition of *C. difficile* germination and growth. (B) Inhibition of spore

germination using various BAs (0.75 mM) with the germinant TCA (2 mM). (C) Inhibition of *C.*

difficile growth as determined by minimum inhibitory concentration (MIC) testing. Statistical

differences in B and C were assessed by a one-tailed Welch's *t* test. (D) Schematic of pre-FMT

collection and *ex vivo* growth. Pre-FMT stool collected from rCDI patient was supplemented

with PBS or a BSH cocktail and subsequently inoculated with *C. difficile*. (E) *C. difficile* growth

was measured at 8h. (F) Targeted metabolomics showing the deconjugated BAs CA and CDCA

from the samples from E. Statistical differences tested by one-way repeated measures ANOVA

with Sidak's correction for multiple comparisons. All bars represent the average of n=3

replicates ± sd. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

Figure 4. BSHs shape the non-canonical conjugated BA pool that governs *C. difficile* spore

germination (A) Schematic of BSH-dependent inhibition of *ex vivo C. difficile* germination and

growth. Intestinal contents harvested from cefoperazone-treated mice (n = 3-4) were

supplemented with a cocktail of BSHs and then inoculated with *C. difficile* spores or cells to

measure germination or growth, respectively. (B) Germination of *C. difficile* spores in murine

small intestinal contents. Contents were treated for 8h with PBS or a BSH cocktail before

inoculation. (C) Growth at 8h of *C. difficile* in murine intestinal contents. Contents were treated

with PBS or a BSH cocktail and inoculated simultaneously. Statistical differences were tested by

ratio paired *t* test. (D) Targeted metabolomics showing the BAs identified in the Day 0 small

intestinal samples from C. (E) Inhibition of *C. difficile* germination using various BAs (0.75

mM) with the germinant TCA (2 mM). Statistical differences were assessed by one-tailed *t* test.

Bars represent mean ± standard deviation. (F) Heatmap of LingBSH across all non-canonical BA

pools. Values represent the mean from n = 3 replicates of % deconjugation after 5 minutes. Bars

represent mean of n=3-4 ± standard deviation. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p <$

0.0001.