The maintenance of self-incompatibility and the genetic architecture of inbreeding

depression

Daniel J. Schoen and Sarah J. Baldwin

Department of Biology, McGill University, Montreal, QC H3A 1B1 Canada

Author for correspondence:

Daniel J. Schoen

E-mail: dan.schoen@mcgill.ca

Word Count:

Summary: 175

Introduction: 567

Methods: 1774

Results: 684

Discussion: 1357

Total: 4557

Tables: 2

Figures: 3

Supplementary Information: Tables S1- S3, Notes (2)

Summary

• Inbreeding depression plays a fundamental role in evolution. To help detect and
characterize viability loci that underlie inbreeding depression, we forced self-pollinated
plants from self-incompatible populations of *Leavenworthia alabamica* to produce
families of progeny that were genotyped at hundreds of mapped single nucleotide
polymorphism (SNP) loci.

• Bayesian analysis of segregation data for each SNP was used to explore support for
different dominance and selection coefficients at linked viability loci in different genomic
regions.

• There was some support for overdominance (or pseudo-overdominance) at a few viability
loci, as well as for recessiveness and underdominance. One recessive viability locus
mapped to the genomic region of the novel self-incompatibility locus in *Leavenworthia*
alabamica, but in general there was no support for strongly recessive viability loci of
major effect.

• The results are consistent with earlier findings showing that inbreeding depression is
recalcitrant to purging in *Leavenworthia alabamica*. The results also help account for the
maintenance of self-incompatibility in this species and are consistent with expectations
from evolutionary genetic theory that recessive, deleterious alleles linked to loci under
balancing selection are sheltered from selection.

Keywords: Viability loci, dominance coefficient, selection coefficient, Bayesian analysis,
sheltered genetic load, mating system evolution

Introduction:

Inbreeding depression is the reduction of fitness in progeny produced by consanguineous

mating versus mating between unrelated parents (Wright, 1977). Inbreeding depression plays a

fundamental role in crop production, conservation of natural populations, and mating system

evolution (Charlesworth & Willis, 2009; Frankham, 2010; Lloyd, 1979; Lande & Schemske,

1985). While recessive deleterious mutations are thought to play a major role in inbreeding

depression, overdominance has also been implicated (Wright, 1977; Charlesworth & Willis,

2009).

Lande & Schemske (1985) considered the evolutionary consequences of inbreeding

depression and the mating system. Their theoretical work predicted when there is selective

purging of recessive lethal and near-lethal alleles, populations of self-fertilizing organisms

should exhibit reduced equilibrium levels of inbreeding depression. Charlesworth &

Charlesworth (1987) extended the study of purging of inbreeding depression to many deleterious

alleles of milder effect with a broader range of dominance and found that fairly high levels of

inbreeding depression can be maintained in populations of self-fertilizing organisms. For species

with self-incompatibility (SI), theory predicts that an additional source of inbreeding depression

is due to the linkage of deleterious recessive alleles to self-recognition loci (Uyenoyama, 1997).

In gametic self-incompatibility systems the self-recognition locus (S-locus) region is maintained

as heterozygous, thereby hindering the selective purging of recessive deleterious mutations, and

leading to "sheltered genetic load" in this region. Likewise, in sporophytic self-incompatibility

systems, depending on the dominance levels of the S-allele, the S-locus region is expected to be

maintained as heterozygous and harbour sheltered S-linked load (Schierup *et al*., 1997; Llaurens

et al., 2009). Theory has shown that sheltered load may affect the trajectory of mating system

evolution (Porcher & Lande, 2005; Gervais *et al*., 2014).

Few studies have provided information on the number and genomic locations of the viability

loci responsible for the expression of inbreeding depression, and there is little information on the

selection and dominance coefficients of the viability effects, factors important to predicting the

evolutionary trajectory of inbreeding depression. For SI species, there is even less information of

this kind (though see Karkkainnen *et al*., 1999).

The genus *Leavenworthia* (Brassicaceae) is a group of self-incompatible and self-compatible

species that has received considerable attention by evolutionary biologists interested in plant

mating system evolution due to both inter- and intraspecific variation in the mating system

(Lloyd, 1965; Busch, 2005; Busch *et al.*, 2011; Herman *et al.*, 2012; Chantha *et al.*, 2013). Self-

incompatible species in this genus exhibit significant levels of inbreeding depression

(Charlesworth *et al.*, 1994; Busch, 2005). Recently, Baldwin & Schoen (2018) conducted

experiments in which plants from self-incompatible populations of *L. alabamica* were subjected

to several generations of selfing. They found that inbreeding depression expressed in these

experiments is resistant to purging.

Marker-based approaches to studying the genetic architecture of inbreeding depression can

provide information on the dominance and selective coefficients at individual viability loci

(Hedrick & Muona, 1990; Fu & Ritland, 1994a,b), which in turn can help shed light on how

inbreeding depression evolves. When available, such information can be coupled with

recombination and physical maps to reveal the approximate genomic location of viability loci

(Mitchell-Olds, 1995; Remington & O'Malley, 2000; Karkkainnen *et al.*, 1999). Here we extract

information on the genetic architecture of inbreeding depression in *Leavenworthia alabamica*

using a genetic marker-based approach. We discuss the relevance of our results to mating system

evolution and to previous findings that have used the marker-based approaches to examine the

genetic architecture of inbreeding depression in plants.

Methods

<u>Plant growth and production of self-fertilized progeny</u>

Two self-incompatible *Leavenworthia alabamica* plants from separate populations in

Alabama were used in this work. The first was from the Hatton population (Latitude, Longitude:

34.509931, −87.442041), while the second was from the M32 population (Latitude, Longitude

34.489356, −87.63559) (Baldwin & Schoen, 2018). These plants were grown in separate 9" pots

in the McGill University Phytotron and forced to self-fertilize by bud or salt pollinations as

described in Chantha *et al.* (2013). Up to 30 pollinations were carried out per plant two times per

week. All seeds were collected at maturity, stratified in a dry environment for at least two

months, and germinated. The progeny from these two separate seed families (Families 1 and 2,

hereafter) were grown to the flowering stage. Sample sizes were 91 and 92 progeny/family,

respectively.

Genotyping-by-sequencing (GBS)

Young leaf tissue samples from the progeny were collected from flowering stage plants

and stored at -80°C. DNA was extracted from the leaf tissue with Qiagen plant mini-DNA kits.

Extractions were repeated if concentrations were lower than 20 ng/ul and were diluted to 30-50

ng/ul if over 50 ng/ul.

The genotyping-by-sequencing (GBS) method was used to produce fragments for

genotyping with restriction-site associated polymorphisms (Elshire *et al*., 2011) using the

protocol of Wallace & Mitchell (2017). The standard protocol was used to generate libraries with

1.44 ng PstI per 100 ng DNA and 15-18 cycles of PCR. Fragments were then sequenced on one

lane of an Illumina High-Seq run (Family 1, N = 91 and Family 2, N = 90). FASTQC was used

to evaluate the basic quality of the sequencing runs. Sequence reads begin with a barcode on the

5' end identifying the plant of origin, then the restriction site, the sequence to be used for

genotyping, followed by another restriction site, and finally the Illumina universal adapter. Reads

were demultiplexed using Process Radtags in STACKS v. 2.59. The *Leavenworthia alabamica*

reference assembly version 2.5 (Haudry *et al.*, 2013, CoGe platform: Genome ID 20247) was

indexed and sorted with Burrows-Wheeler Aligner Tool, BWA-0.7.17 (Li & Durbin, 2010), and

reads were aligned to the reference genome with BWA-0.7.17. SNP locus segregation ratios

derived from the two families were obtained by producing a catalog of segregating loci using the

GSTACKS and POPULATIONS routines in STACKS v. 2.59 (Catchen *et al*., 2013) running in

R (version 4.1.1). The in-house scripts and associated information used for processing the data

are available in the Supplementary Materials section (Table S1).

SNP filtering and treatment of potential genotyping errors

Previous genomic analysis of *Leavenworthia alabamica* indicates that this species

underwent whole genome triplication during its evolutionary history, and chromosome painting

with *Arabidopsis thaliana* BAC probes suggests that that some triplicated genome segments

were retained as duplicates or triplicates (Haudry *et al*., 2013). The reference genome assembly

in this species is only partially complete (N50 = 309 kb); while the haploid chromosome number

of *L. alabamica* is 11, the genome assembly consists of 3,321 scaffolds. It comprises *ca.* 174

million bp in total of the expected 315 million bp genome size, the latter estimated from flow

cytometry (Brian Husband, pers. comm.). Over half of the scaffolds are short, with lengths 5,000

bp or smaller, though larger ones are > 1M bp. While there is sequence divergence of the

duplicated (or triplicated) regions, longer scaffolds in the reference genome likely represent

separate genomic regions (Haudry *et al*., 2013). But many shorter duplicated (or triplicated)

regions of the genome with extended identity have likely been collapsed into single (artifactual)

contigs. GBS products from separate genomic regions that map to such short, artifactual contigs

would be expected to yield heterozygote genotype calls. To minimize these artifacts we: (1)

filtered out heterozygotes when allele balance fell below 0.333 or above 0.667 using the R-

package SNPfiltr v. 1.00 (DeRaad, 2022), (2) restricted our analysis to SNPs that map to

scaffolds > 100 kbp in length (these account for 73% of the *L. alabamica* assembly); (3) filtered

out SNP loci that exhibited only heterozygotes when adjacent SNP loci on the same scaffold

exhibited segregation ratios that were not significantly different from 1:2:1 by Chi-square tests,

on the basis that evidence of strong selection should be shared by adjacent SNP loci; and (4)

restricted our analysis to loci where data were available from > 90% of the progeny of a cross

since genotyping errors are more likely to have occurred when few progeny per family are

genotyped successfully.

<u>Selection and dominance coefficients at linked viability loci</u>

Linkage groups (each containing a minimum of three SNP loci) were constructed

separately for Families 1 and 2 with the software package 'qtl' version 1.50 (Broman *et al*., 2003)

running in R (version 4.1.1). A viability locus located adjacent to a given SNP locus will likely

affect not only the segregation ratio at that SNP but also at other nearby SNPs. The linkage maps

allowed us to examine segregation ratios of each SNP in relation to that of other nearby SNPs. To

arrive at a conservative estimate of the number of viability loci, we limited our analysis to those

SNPs that exhibited departures from the expected (null) 1:2:1 segregation ratio at a false discovery

rate (Benjamini & Hochberg, 1995) of q = 0.05 and 0.10. For linkage groups where departures

from a 1:2:1 ratio were observed, we selected the marker showing the most extreme departure for

the analysis of selection and dominance coefficients at the linked viability locus.

We note that the analysis of marker locus segregation ratios to estimate selection and

dominance at viability loci linked to marker loci assumes departure from the expected 1:2:1

Mendelian segregation ratio is due to viability selection, rather than to meiotic drive and/or gametic

selection. The parent plant that is self-fertilized is assumed to be heterozygous at a selectively

neutral marker locus (*M*) and a linked viability locus (*V*)—e.g., the parent has the genotype

M_1V_1/M_2V_2, and the genotypes V_1/V_1, V_1/V_2, and V_2/V_2 are associated with the fitness values 1-*s*,

1-*hs*, and 1, respectively, where *s* is the selective coefficient or the proportional amount by which

selection at this locus reduces viability in the homozygote V_2/V_2 ($0 \leq s \leq 1$), and *h* is the dominance

coefficient. A value of *h* < 0 implies overdominance, *h* = 0 implies fully recessive, $0 \leq h \leq 0.5$

implies partially recessive, *h* = 0.5 implies additive, $0.5 \leq h \leq 1.0$ implies partial dominance, and

h > 1 implies underdominance. The marker and viability loci are assumed to be linked with

recombination fraction *c*.

A Bayesian statistical genetic approach was developed to explore the posterior support for

different values of *h* and *s* at viability loci based on the observed numbers of marker locus

genotypes. Fu & Ritland (1994a) showed that following self-fertilization, the post-selection

genotype frequencies expected for three marker genotypes (p_{11} for M_1/M_1, p_{12} for M_1/M_2, and p_{22}

for M_2/M_2) are:

$$p_{11} = \frac{c^2\, s(2h-1) - 2\, cs(h-1) + (1-s)}{4 - s(2h+1)}$$

$$p_{12} = \frac{2(-c^2\, s(2h-1) + 2cs\,(2h-1) - hs + 1)}{4 - s(2h+1)} \qquad\qquad (1)$$

$$p_{22} = \frac{c^2\, s(2h-1) - 2\, csh + 1}{4 - s(2h+1)}$$

In these equations p_{11} denotes the frequency of the least common homozygote genotype. Note

that when *s* = 0, the marker genotype frequencies give the null 1:2:1 ratio.

The data are the numbers of each of the three marker genotypes observed among the

surviving progeny of the self-fertilized parent (N_{11} = number of M_1/M_1, number of observed, N_{12}

= number of M_1/M_2 observed, and N_{22} = number of M_2/M_2 observed). The probability

(likelihood) given the data can be written:

$$Pr\,(N_{11}, N_{12}, N_{22} \,|\, s, h, c\,) = \left(\frac{N!}{N_{11}!\, N_{12}!\, N_{22}!}\right)\, p_{11}{}^{N_{11}} p_{12}{}^{N_{12}}\, p_{22}{}^{N_{22}} \qquad\qquad (2)$$

where N is the sum of N_{11}, N_{12}, and N_{22}. The joint density and posterior probability are,

respectively:

$Pr(s, h, c, N_{11}, N_{12}, N_{22}) = Pr(s) \, Pr(h) \, Pr(c) \, Pr(N11, N12, N22 \mid s, h, c)$ (3)

$Pr(s, h, c \mid N_{11} N_{12} N_{22}) = \dfrac{Pr(s) \, Pr(h) \, Pr(c) \, Pr(N11, N12, N22 \mid s,h,c)}{\int_{s,h,c} Pr(s) \, Pr(h) \, Pr(c) Pr(N11, N12, N22 \mid s,h,c) \, ds \, dh \, dc}$ (4)

To explore values of the parameters using a Bayesian approach, we chose uniform priors with 0

$< s < 1$, $-100 < h < 100$, and $0 < c < 0.2$. The broad prior for h allows for the possibility of

extreme overdominance and underdominance. The prior for c (less than the theoretical upper

limit of $c = 0.5$) is justified based on the average distance between markers of < 5 cM in this

study. A program written in the R statistical language (version 4.1.1) to conduct Markov Chain

Monte Carlo method with the Metropolis-Hastings sampler (Weir, 1996) (Supplementary

Materials: Note 1). The program was used to "integrate out" the nuisance parameter c, allowing

the posterior probabilities to be estimated for different values of s and h. For the required burn-in

period, we ran the sampler 5.5 million times before examining the posterior probability

distributions. Tracking the parameter values during the runs showed that the sampler had

stabilized by the end of this burn-in period.

<u>The S-locus region of *Leavenworthia alabamica* and locations of viability loci</u>

221   The *Leavenworthia alabamica* S-locus region has previously been cloned and sequenced

from the same plant material that was used to produce the sequence information in the genome

assembly in the present work (GenBank accession KC981242.1) (Chantha *et al*., 2013). This

allowed us to locate the individual *L. alabamica* genome assembly scaffold containing the S-

locus. The BLAST algorithm (Altschul *et al.,* 1990) implemented in software package Geneious

Prime (version 2.2) was used to conduct the search of *L. alabamica* reference genome version

2.5 using both *L. alabamica* S-locus genes (*Lal2* and *LaSCRL*; Chantha *et al.*, 2013). Selection

and dominance coefficients were estimated for all potential viability loci linked to SNPs in this

region that showed significant departures from 1:2:1 were analysed.

Statistical power test and different modes of gene action

The analysis of the number and mode of action pf the linked viability loci hinges on the

ability to detect departure from the expected 1:2:1 ratio obtained when self-fertilizing a plant

with heterozygous marker loci. Given that the sample size of progeny per family in our work was

limited to a maximum of 91 progeny (the maximum number we could obtain by forced selfing of

each of the SI parent plants used), we were interested in the statistical power of our study to

detect selection and characterize the mode of gene action. We did this by producing expected

segregation ratios for a range of different values of the parameters in equation (1) followed by

standard methods (Cohen, 1988) for examining statistical power, as implemented in the R

program package 'pwr' version 1.3 (Supplementary Materials: Note 2).

Results

SNP segregation ratios, linkage map locations, and significance testing

We found 315 mappable SNP loci in Family 1, and 237 in Family 2. In Family 1 we

found 18 linkage groups with 3 or more SNPs, ranging in length from 2 to 201 cM, for a total

map length of 901 cM. In Family 2 we found 16 linkage groups with 3 or more SNPs, ranging in

length from 3 to 88 cM, for a total map length of 229 cM. SNP grouping by linkage group was

consistent in the two families (see Supplementary Tables S2 and S3 for linkage group, scaffold

positions, and segregation patterns of SNPs). SNP segregation patterns along with and their -

$\log_{10}$ significance levels from the Chi-square test of goodness-of-fit to the null 1:2:1 segregation

ratio varied within each linkage group are shown in Figures 1 and 2.

Selection and dominance of viability loci linked to SNPs

In Family 1, we detected four regions of the genome with putative viability loci (Fig. 1).

In three of these regions (linkage groups 1,6, and 9), Bayesian estimates of h and s support the

existence of viability loci with weakly recessive gene action and large selection coefficients

(Table 1, Fig. 1). In the fourth region (linkage group 6), Bayesian estimates of h and s support

the existence of a viability locus with overdominant gene action and a large selection coefficient,

though it should be noted that our analysis does not permit distinguishing between

overdominance and pseudo-overdominance.

In Family 2, we detected two regions of the genome with putative viability loci (Fig. 2)

and a third that we were unable to map to any linkage group. In linkage group 5, Bayesian

estimates of h and s support the existence of a viability locus with overdominant gene action and

a large selection coefficient, though again it should be noted that our analysis does not permit

distinguishing between overdominance and pseudo-overdominance (Table 1, Fig. 2). In linkage

group 11, a viability locus with underdominant gene action and a large selection coefficient was

inferred (Table 1, Fig. 2). The two genomic regions where these viability loci were detected are

in separate linkage groups from those detected in Family 1 (Table S3). The segregation ratio for

the unmapped viability locus suggests strong overdominance. Though it is also possible that this

is an artifact arising from a locus that maps to duplicated genomic regions as discussed in the

Methods section (Table 1).

<u>The S-locus region of *Leavenworthia alabamica* and locations of viability loci</u>

The tightly linked S-locus genes *Lal2* and *SCRL* along with their intervening non-coding

region are located on scaffold 46, at base pair positions 2,035,545 to 2,044,046. Several genomic

regions on this scaffold in Family 1 showed a strong deviation from the 1:2:1 ratio (Fig. 2 and

Table 2). The segregation data here support recessive gene action and strong selection $h = 0.18$ to

0.45, and $s = 0.80$ to 0.86 (Table 2). In Family 2, however, this region did not harbour any

detectable viability loci.

<u>Statistical power to detect different modes of gene action</u>

Statistical power to detect different modes of gene action in experiments of the type

conducted here depends on the value of the recombination fraction c. For close linkage, e.g., c

near 0 as is likely for most SNPs in this study (Figs. 1 and 2, Table S2), overdominant and

underdominant action are easier to detect than dominant and recessive action (Fig. 3). As the

recombination fraction between marker and inbreeding loci increases, statistical power to detect

underdominant selection increases, whereas statistical power to detect overdominant gene action

decreases (Fig. 3). For dominant and recessive gene actions, statistical power of detection is

reduced with increasing c, but compared with overdominant and underdominant action the

changes are less affected by recombination fraction. Given a sample size of 90 progeny, close

linkage, and overdominant or underdominant genes actions, selection coefficients of *ca*. $s = 0.3$

at viability loci are detectable with statistical power of 80%. For dominant and recessive actions,

viability loci with selection coefficients of *ca*. $s = 0.6$ are detectable with statistical power of

80%.

Discussion

<u>The genetic architecture of inbreeding depression.</u>

We detected a number of significant departures from the 1:2:1 segregation ratio at SNPs

located in different regions of the *Leavenworthia alabamica* genome in the progeny of parental

plants from two SI populations. The segregation data were used to explore support for different

values for selection and dominance coefficients associated with linked viability loci. The

approach we used assumes that meiotic drive, gametic selection, and epistasis do not contribute

to departures from normal segregation. The inability to distinguish the effect of these factors

from viability selection is a limitation of this approach. Nevertheless, there are several reasons to

believe that most departures from the 1:2:1 segregation ratio observed in this study are associated

with inbreeding depression rather than with meiotic drive or gametic selection. First, where

evidence of meiotic drive has been studied in detail in other species (e.g., in *Arabidopsis*

thaliana), the effects were generally found in specific genomic regions, rather than genome wide

as observed in the present study (Salomé *et al.*, 2012; Seymour *et al.*, 2019). Second, gametic

selection favouring a specific allele linked to markers is not expected to produce a heterozygote

excess at linked marker loci, as observed in this study for some of the marker loci in this study.

Several past studies have used marker data to explore the genetic architecture of

inbreeding depression in plants. Hedrick & Muona (1990) used the approach in *Pinus sylvestris*

and detected a single viability locus with a strong contribution to the fitness of progeny. In

another study, using progeny from self-pollinated plants collected from a wild *Mimulus guttatus*

population, Fu & Ritland (1994b) applied a graphical approach that divides the space of possible

segregation ratios into different modes of gene action. They detected 24 viability loci, with the

majority showing underdominance or dominance effects. Mitchell-Olds (1995), studying

inbreeding depression in the progeny of parent of an *Arabidopsis thaliana* F1 hybrid between

different accessions, reported a single viability locus with overdominance that mapped to

chromosome 1. Remington and O'Malley (2000) examined progeny segregation ratios at AFLP

loci in selfed *Pinus taeda* plants and detected both recessive and overdominant action at 19

embryonic viability loci. In the only other study to apply a Bayesian approach to estimating

selection and dominance at loci associated with inbreeding depression, albeit with a different

parameterisation that the one employed here, Kärkkäinen *et al*. (1999) in *Arabidopsis lyrata* ssp.

petraea reported linkages of viability loci to seven separate allozyme loci, mostly with

overdominant effects. Finally, Hedrick *et al.* (2016) examined segregation ratios for nearly

10,000 SNPs in 28 progeny from a self-pollinated plant of *Eucalytus grandis*, and found

widespread heterozygote excess, consistent with overdominance, though their simulation

analysis suggested that pseudo-overdominance is the more likely explanation. Overall, these

empirical studies suggest that modes of gene action for linked viability loci may vary from

species to species.

It is notable that a few of the above-cited studies have reported underdominant viability

loci linked to markers (e.g., especially Fu & Ritland, 1994b, but also Kärkkäinen *et al*. 1999).

Underdominance was also inferred at two marker-linked viability loci in the present

investigation. Underdominance at viability loci cannot contribute to inbreeding depression. We

note from the results of the power analyses we conducted, underdominance is detectable with

smaller sample sizes compared with other modes of action that require larger sample sizes to

detect selection of the same magnitude when $c < 0.3$ (Fig. 3). This could help account for the

unexpected reports of underdominance in studies that have employed few markers (e.g., Fu &

Ritland, 1994b).

<u>Overdominance or pseudo-overdominance of inbreeding depression</u>

For a number of the marker loci showing departures from normal segregation, our

analysis suggests overdominance at linked viability loci contributes to inbreeding depression.

Alternatively, these results could be explained as being due to pseudo-overdominance (Ohta,

1971). Psuedo-overdominance would be detected in a genomic region by the methods used here

if deleterious, recessive viability mutations are linked and in repulsion. In general, compared

with recessive or partly recessive action at viability loci, there is little support from past studies

for true overdominance as an underlying cause of inbreeding depression (Charlesworth and

Willis 2009). Distinguishing between true overdominance and pseudo-overdominance is difficult

but becomes progressively easier with multiple families. True overdominance, if present, should

often be detected at the same viability loci in the progeny of multiple selfed parents, since

selection there should render it frequent. In contrast, viability loci exhibiting pseudo-

overdominance arising from deleterious recessive or partly recessive alleles are less likely to be

shared among parents, as the genomic regions are maintained by the interaction of mutation and

selection. This leaves open the possibility that both true and pseudo-overdominance are present.

<u>Sheltered S-linked load</u>

An inferred viability locus (or possibly, multiple linked loci) with recessive action was

detected only in one genomic region, and one family (Family 1). Interestingly, the region is

within 80 kb from the site of the S-locus. These results can be interpreted as evidence that there

is sheltering from selection of one (or more) recessive, deleterious mutations in the S-locus

region (Table 2).

In Family 2, no such effects were detected in the S-locus region. The finding is not

necessarily at odds with the evidence of sheltered load in the Family 1 data. There are likely

several levels of dominance for S-locus alleles in *L. alabamica* (Busch *et al.*, 2010), and in

sporophytic systems such as this one, the strongest sheltering from selection is expected for those

viability loci linked to S-alleles that exhibit the highest degree of dominance (Llaurens *et al.*,

2009).

<u>Inbreeding depression, purging, and the maintenance of self-incompatibility in *Leavenworthia*</u>

<u>*alabamica*</u>

Busch (2005) found significant levels of inbreeding depression for germination and

survival rate, biomass, flower number, and pollen viability in self-incompatible populations of

Leavenworthia alabamica, and estimated that inbreeding depression for total lifetime fitness

exceeds 0.5. Baldwin & Schoen (2019) studied inbreeding depression for a similar set of fitness

components in *L. alabamica* and again found that inbreeding depression for lifetime fitness

exceeds 0.5. They extended their study to include three consecutive generations of self-

pollination and found no evidence that inbreeding depression was purged by this treatment.

383   Inbreeding depression is thought to be the main barrier to the evolution of self-pollination

384 (Fisher, 1941; Lloyd 1979), and yet with some self-pollination, purging of recessive load is

385 expected. In an earlier set of population surveys Baldwin & Schoen (2017) found that some

386 selfing likely does occur in populations of *Leavenworthia alabamica* that have been formerly

387 characterized at SI. Between 6% and 18% of plants exhibit "leaky self-incompatibilty" in which

388 pollen tubes from self-pollen penetrates the styles. Such plants are presumably capable of setting

389 seed with self-pollen. Baldwin & Schoen (2017) also showed that variation in the degree of this

390 leakiness is heritable, and so at least one requirement for purging inbreeding depression (i.e.,

391 heritable variation for level of self-pollination) is likely present in populations of this species.

392 This raises the question of why SI is maintained. Theory suggests that the spread of mutations

393 that reduce SI occurs mainly when inbreeding depression loci are of major effect and recessive

394 (Porcher & Lande, 2005; Gervais *et al*., 2014). The nearly additive viability loci detected in this

395 study are not expected to contribute significantly to standing inbreeding depression

396 (Charlesworth & Charlesworth, 1987). Moreover, any component of inbreeding depression due

397 to loci with overdominance or pseudo-overdominance (as also detected in this study) should be

398 resistant to purging. Given our findings here, we suggest that inbreeding depression in the self-

399 incompatible populations of this species is likely due to many loci of small effect—or in other

400 words, with effects too mild to be detected in the type of study we reort here. Inbreeding

401 depression of this type would be expected to be resistant to purging (Charlesworth &

402 Charlesworth, 1987) as we observed in our earlier work (Baldwin & Schoen, 2019).

404 **Acknowledgements**

405   We thank Mark Romer and Claire Cooney for assistance with plant growth in the McGill

406 Phytotron. Yong-Bi Fu provided comments on the manuscript. SJB acknowledges scholarship

407 support from the Fonds de Recherche du Québec – Nature et technologies and support from a

408 Discovery Grant from the Natural Sciences and Engineering Research Council of Canada to DJS

410 **Author Contributions**

411 DJS and SJB conceived the study. DJS developed the Bayesian approach for characterizing the

412 genetic architecture of viability loci underlying inbreeding depression, and the power analysis.

SJB and DJS initiated the study. Both authors analyzed and interpreted the data and wrote the

manuscript. The authors are joint first authors.

Data Availability

The *Leavenworthia alabamica* reference assembly version 2.5 is available on the CoGe platform:

https://genomevolution.org/coge/GenomeInfo.pl?gid=20247. FASTQ files of GBS sequence data

for parents and progeny are available on GenBank under SRA accession number PRJNA813545

Supporting Information

423	**File name**	**Description**
424	TableS1.xlsx	Table S1. Scripts and additional information used to process GBS data in
425		STACKS
426	TableS2.xlsx	Table S2. SNPs from Families 1 and 2, their position in linkage groups
427		and scaffolds; and segregation patterns.
428	TableS3.xlsx	Table S3. Cross reference for linkage groups in Families 1 and 2.
429	Note 1	R code for Bayesian analysis
430	Note 2	R code for power analysis in Fig. 3

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search
tool. *Journal of Molecular Biology* **215**: 403-410.

Baldwin SJ, Schoen DJ. 2017. Genetic variation for pseudo-self-compatibility in self-
incompatible populations of *Leavenworthia alabamica* (Brassicaceae). *New Phytologist* **213**:
430-429.

Baldwin SJ, Schoen DJ. 2019. Inbreeding depression is difficult to purge in self-incompatible
populations of *Leavenworthia alabamica*. *New Phytologist* **224**:1330-1338.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful
approach to multiple testing. *Journal of the Royal Statistical Society: Series B*
(Methodological) **57**: 289-300.

Broman KW, Wu H, Sen Ś, Churchill GA. 2003. R/qtl: QTL mapping in experimental crosses.
Bioinformatics **19**: 889-90.

Busch JW. 2005. Inbreeding depression in self-incompatible and self-compatible populations of
Leavenworthia alabamica. *Heredity* **94**: 159-165.

Busch JW, Joly S, Schoen DJ. 2010. Does mate limitation in self-incompatible species promote
the evolution of selfing? The case of *Leavenworthia alabamica. Evolution* 64:1657-1670.

Busch JW, Joly S, Schoen DJ. 2011. Demographic signatures accompanying the evolution of
selfing in *Leavenworthia alabamica*. *Molecular Biology and Evolution* **28**: 1717-1729.

Chantha SC, Herman AC, Platts AE, Vekemans X, Schoen DJ. 2013. Secondary evolution of
a self-incompatibility locus in the Brassicaceae genus *Leavenworthia*. *PLoS Biology* **14**:
e1001560.

Charlesworth D, Lyons EE, Litchfield LB. 1994 Inbreeding depression in two highly
inbreeding populations of *Leavenworthia*. *Proceedings of the Royal Society of London,*
Series B **258**: 209-214.

Charlesworth D, Charlesworth B. 1987. Inbreeding depression and its evolutionary
consequences. *Annual Review of Ecology and Systematics* **18**: 237-268.

Charlesworth D, Willis JH. 2009. The genetics of inbreeding depression. *Nature Reviews*
Genetics **10**: 783-796

Cohen, J. 1988. *Statistical power analysis for the behavioral sciences, 2nd ed*. Hillsdale, NJ
(USA): Lawrence Erlbaum Associates.

DeRaad DA. 2022. SNPfiltR: an R package for interactive and reproducible SNP filtering.
Molecular Ecology Resources (in press).

Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, Mitchell SE. 2011.
A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PloS*
One **6**: e19379.

Frankham R. 2010. Where are we in conservation genetics and where do we need to go?
Conservation Genetics **11**: 661-3.

Fu YB, Ritland K. 1994a. On estimating the linkage of marker genes to viability genes
controlling inbreeding depression. *Theoretical and Applied Genetics* **88**: 925-932.

Fu YB, Ritland K. 1994b. Evidence for the partial dominance of viability genes contributing to
inbreeding depression in *Mimulus guttatus*. *Genetics* **136**: 323-331.

Gervais C, Awad DA, Roze D, Castric V, Billiard S. 2014. Genetic architecture of inbreeding
depression and the maintenance of gametophytic self-incompatibility. *Evolution* **68**: 3317-
3324.

Glémin S. 2003. How are deleterious mutations purged? Drift versus nonrandom mating.
Evolution **57**: 2678-2687.

Haudry A, Platts AE, Vello E, Hoen DR, Leclercq M, Williamson RJ, Forczek E, Joly-
Lopez Z, Steffen JG, Hazzouri KM, Dewar K. 2013. An atlas of over 90,000 conserved
noncoding sequences provides insight into crucifer regulatory regions. *Nature Genetics*
45:891-898.

Hedrick PW, Muona O. 1990. Linkage of viability genes to marker loci in selfing organisms.
Heredity **64**: 67-72.

Hedrick PW, Hellsten U, Grattapaglia D. 2016. Examining the cause of high inbreeding
depression: analysis of whole-genome sequence data in 28 selfed progeny of Eucalyptus
grandis. *New Phytologist* **209**: 600-611.

Herman AC, Busch JW, Schoen DJ. 2012. Phylogeny of Leavenworthia S-alleles suggests
unidirectional mating system evolution and enhanced positive selection following an ancient
population bottleneck. *Evolution* **66**: 1849-1861.

Kärkkäinen K, Kuittinen H, van Treuren R, Vogl C, Oikarinen S, Savolainen O. 1999.

Genetic basis of inbreeding depression in *Arabis petraea*. *Evolution* **53**:1354-1365.

Lande R, Schemske DW. 1985. The evolution of self-fertilization and inbreeding depression in

plants. I. Genetic models. *Evolution* 39: 24-40.

Li H. and Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler

transform. *Bioinformatics* **26**: 589-595.

Llaurens V, Gonthier L, Billiard S. 2009. The sheltered genetic load linked to the S locus in

plants: new insights from theoretical and empirical approaches in sporophytic self-

incompatibility. *Genetics* **183**: 1105–1118.

Lloyd DG. 1965. Evolution of self-compatibility and racial differentiation in Leavenworthia

(Cruciferae). Contributions of the Gray Herbarium **195**:3-134.

Lloyd, DG. 1979. Some reproductive factors affecting the selection of self-fertilization in plants.

The American Naturalist **113:** 67-79.

Mitchell-Olds T. 1995. Interval mapping of viability loci causing heterosis in Arabidopsis.

Genetics **140**: 1105-1109.

Porcher E, Lande R. 2005. The evolution of self-fertilization and inbreeding depression under

pollen discounting and pollen limitation. *Journal of Evolutionary Biology* **18**: 497-508.

Remington DL, O'Malley DM. 2000. Whole-genome characterization of embryonic stage

inbreeding depression in a selfed loblolly pine family. *Genetics* **155**: 337-348.

Schierup MH, Vekemans X, Christiansen FB. 1997. Evolutionary dynamics of sporophytic

self-incompatibility alleles in plants. *Genetics* **147**: 835-846.

Seymour DK, Chae E, Arioz BI, Koenig D, Weigel D. 2019. Transmission ratio distortion is

frequent in *Arabidopsis thaliana* controlled crosses. *Heredity* **122**: 294-304.

Uyenoyama MK. 1997. Genealogical structure among alleles regulating self-incompatibility in

natural populations of flowering plants. *Genetics* **147**: 1389–1400.

Wallace J, Mitchell S. 2017. Genotyping-by-sequencing. *Current Protocols in Plant Biology* **2**:

64-77.

Weir BS. 1996. *Methods for discrete population genetic data. Genetic Data Analysis II.*

Sunderland, MA (USA): Sinauer Associates, Inc.

Wright S. 1984. *Evolution and the genetics of populations, vol. 3: Experimental results and*

evolutionary deductions. Chicago, IL (USA): University of Chicago Press.

Figure Legends

Figure 1. -Log_{10} significance levels from Chi-square goodness-of-fit tests to the 1:2:1

segregation ratio for SNPs in linkage groups of Family 1. Red solid and dashed lines indicate

significance levels corresponding to the false discovery rate of $q = 0.05$ and 0.10, respectively.

Below are the segregation ratios for the SNPs. Blue denotes heterozygote, red and black denote

homozygotes. Above are heat maps of posterior probability densities for dominance (*h*) and

selection (*s)* coefficients of inferred viability loci in linkage groups (positions indicated by

arrows).

Figure 2. -Log_{10} significance levels from Chi-square goodness-of-fit tests to the 1:2:1

segregation ratio for SNPs in linkage groups of Family 2. Red solid and dashed lines indicate

significance levels corresponding to the false discovery rate of $q = 0.05$ and 0.10, respectively.

Below are the segregation ratios for the SNPs. Blue denotes heterozygote, red and black denote

homozygotes. Above are heat maps of posterior probability densities for dominance (*h*) and

selection (*s)* coefficients of inferred viability loci in different linkage groups (positions indicated

by arrows).

Figure 3. Statistical power to reject the 1:2:1 segregation ratio for different modes of action of

viability loci, selection coefficients, and recombination fractions. Shown are four cases:

overdominant with $h = -3.0$ (red), recessive with $h = 0.25$ (black), dominant (blue) with $h = 0.75$,

and underdominant ($h = 1.5$) (orange). From top to bottom graphs, the values of the

recombination fraction are $c = 0$, 0.1, 0.2, and 0.3. The graphs are based on a sample size of 90

progeny.

Fig. 1

Table 1. Marker loci that show the most significant deviations from normal segregation within separate linkage groups. Marker locus (scaffold_base pair position), linkage group (LG), and posterior probabilities of overdominant (OD), recessive (R), dominant (D), and underdominant (UD) gene actions[1]. The values for the selection coefficient (s) and dominance coefficient (h) are the Bayesian estimators (95% credible intervals).

Family	Marker	LG	M_1M_1	M_1M_2	M_2M_2	OD	R	D	UD	s	h
1	61_934679	1	11	43	33	0.11	0.36	0.45	0.07	0.67 (0.34, 1.0)	0.47 (-0.33, 1.14)
1	57_279929	2	11	42	34	0.15	0.38	0.41	0.06	0.68 (0.28, 1.0)	0.42 (-0.45, 1.08)
1	28_130518	6	10	57	22	0.99	0.01	0	0	0.61 (0.23, 0.97)	-1.82 (-3.89, -0.16)
1	46_1956910	9	6	43	36	0	0.51	0.44	0.01	0.86 (0.68, 0.98)	0.45 (0.01, 0.79)
2	27_153823	5	0	69	22	0.99	0.01	0	0	0.91 (0.63, 1.0)	-1.55 (-4.28, -0.23)
2	8_243257	11	18	34	39	0	0.01	0.20	0.79	0.50 (0.20, 0.73)	1.38 (0.74, 2.31)
2	197_101568	-	0	85	0	-	-	-	-	-	-

[1] Cumulative posterior probabilities associated with overdominant (OD), recessive (R), dominant (D), and underdominant (UD) modes of gene actions were determined by summing over the posterior probabilities associated with values of $h < 0$, $0 \leq h < 0.5$, $0.5 \leq h < 1$, and $h \geq 1$, respectively.

Table 2. SNPs on the scaffold containing the *Leavenworthia alabamica* S-locus (*Lal2*)[1] that show significant deviations from 1:2:1 segregation. SNP locus (scaffold_base pair position) and posterior probabilities of overdominant (OD), recessive (R), dominant (D), and underdominant (UD) gene actions[1]. The values for the selection coefficient (*s*) and dominance coefficient (*h*) are the Bayesian estimators (90% credible intervals).

SNP	OD	R	D	UD	*s*	*h*
46_1,497,122	0.26	0.51	0.23	0.00	0.80 (0.52, 0.98)	0.18 (-0.76, 0.71)
46_1,807,312	0.09	0.52	0. 39	0.00	0.82 (0.59 .0.98)	0.39 (-0.18, 0.80)
46_1,807,671	0.10	0.56	0. 33	0.01	0.80 (0.53, 0.95)	0.36 (-0.16,0 .79)
46_1,956,910	0.04	0.51	0.44	0.01	0.86 (0.68, 0.98)	0.45 (0.01, 0.79)
46_1,957,161	0.09	0.52	0.38	0.01	0.85 (0.64, 0.96)	0.38 (-0.18, 0.80)

[1] Base pair position of the start of exon 1 of *LalSCRL* on scaffold 46 is 2,035,545 bp