Discordant calls across genotype discovery approaches elucidate variants with systematic errors

Elizabeth G. Atkinson[1,2,3,*], Mykyta Artomov[1,4,5,6]*, Konrad J. Karczewski[1,2,4], Alexander A. Loboda[1,4,7,8], Heidi

L. Rehm[1,4,9], Daniel G. MacArthur[2,10], Benjamin M. Neale[1,2,4]*, Mark J. Daly[1,2,11]*

[1]Analytic and Translational Genetics Unit, Department of Medicine, Massachusetts General Hospital, Boston,
MA, USA
[2]Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA
[3]Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, USA
[4]Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA
[5]The Steve and Cindy Rasmussen Institute for Genomic Medicine, Nationwide Children's Hospital,
Columbus, OH, USA
[6]Department of Pediatrics, The Ohio State University College of Medicine, Columbus, OH, USA
[7]ITMO University, Saint-Petersburg, Russia
[8]Almazov National Medical Research Center, St. Petersburg, Russia
[9]Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA
[10]Centre for Population Genomics, Garvan Institute of Medical Research and Murdoch Children's Research
Institute, Australia
[11]Institute for Molecular Medicine Finland, University of Helsinki, Helsinki, Finland
*These authors contributed equally to this work.
20

Corresponding author contact information:

Emails: elizabeth.atkinson@bcm.edu; mykyta.artomov@nationwidechildrens.org

Phone: (202) 246-0666

One Baylor Plaza, Taub Research Building Room 621

Department of Molecular and Human Genetics

Baylor College of Medicine

Houston, TX 77030

Running Title: Discordant genotyping across approaches

Keywords: next-generation sequencing; whole exome sequencing; GWAS; variant calling; allele frequencies

Abstract

Large-scale next-generation sequencing datasets have been transformative for informing clinical

variant interpretation and as reference panels for statistical and population genetic efforts. While such

resources are often treated as ground truth, we find that in widely used reference datasets such as the

Genome Aggregation Database (gnomAD), some variants pass gold standard filters yet are systematically

different in their genotype calls across genotype discovery approaches. The inclusion of such discordant

sites in study designs involving multiple genotype discovery strategies could bias results and lead to false-

positive hits in association studies due to technological artifacts rather than a true relationship to the

phenotype. Here, we describe this phenomenon of discordant genotype calls across genotype discovery

approaches, characterize the error mode of wrong calls, provide a blacklist of discordant sites identified in

gnomAD that should be treated with caution in analyses, and present a metric and machine learning

classifier trained on gnomAD data to identify likely discordant variants in other datasets. We find that different

genotype discovery approaches have different sets of variants at which this problem occurs but that there are

characteristic variant features that can be used to predict discordant behavior. Discordant sites are largely

shared across ancestry groups, though different populations are powered for discovery of different variants.

We find that the most common error mode is that of a variant being heterozygous for one approach and

homozygous for the other, with heterozygous in the genomes and homozygous reference in the exomes

making up the majority of miscalls.

Introduction

While next-generation sequencing (NGS) technologies have been transformative for genomics research,

they have an appreciable error rate (Ma et al. 2019) as a cost of their high-throughput capacity. To account

for this, sophisticated pipelines have been developed for the detection and removal of incorrect sequencing

calls (Li et al. 2019; McKenna et al. 2010; Adelson et al. 2019; Highnam et al. 2015; Anderson et al. 2010;

Lam et al. 2019). However, even with gold-standard filtering, spurious genotype calls can infiltrate datasets

and potentially skew results. This is of particular importance with datasets that aggregate calls generated by

multiple genotype discovery approaches, as different strategies have distinct error modes. Identifying

variants that have technical artifacts affecting genotype calls is of major importance, as such loci give

misleading information regarding population allele frequencies, and could be incorrectly identified as being

phenotypically meaningful in gene discovery.

By leveraging the unprecedented size and depth of the gnomAD database (Karczewski et al. 2020; Lek

et al. 2016), we comprehensively characterize trends in genotype calling depending on the sequencing

technology. Specifically, we note that a subset of variants, despite passing standard quality filters

(Karczewski 2017), produce discordant allele frequencies in data generated using different genotype

discovery approaches, stemming from unreliable variant calling. This can not be explained by population

stratification, as this effect is observed even when looking at the same set of individuals. Such unreliably

genotyped variants should therefore be screened out of analyses. Including these variants in gene discovery

efforts, particularly in study designs where case and control data are represented by different combinations

of sequencing platforms or genotype discovery approaches, could result in their appearance as false-positive

associations.

In this article, we comprehensively characterize the observation of discordant genotyping depending on

genotype discovery approach using a large set of diverse individuals from the gnomAD database, including a

subset of participants who underwent both whole-exome and whole-genome sequencing (WES/WGS). We

then validate our findings in two external datasets for which data from multiple genotype discovery

approaches are available: the Thousand Genomes Project and the All of Us Research Project (Auton and

Salcedo 2015; The All of Us Research Program Investigators 2019). Correcting for this technical error,

whether by removing the gnomAD blacklist variants provided here or by identifying user-identified spurious

calls with our freely distributed machine learning predictor, should be incorporated as a step in QC pipelines

to avoid spurious associations, particularly in large-scale studies aggregating data from multiple sources.

Results

Discordance in genotype calls across genotype discovery approaches replicates across ancestries

We sought to compare allele frequencies across variants found within exome and genome sequencing

datasets in gnomAD to test whether there are regions with significant bias associated with genotype

discovery approach. We first focused on the largest population represented in gnomAD 0.2 – the Non-

Finnish Europeans (NFE). Using the full release of gnomAD version 2.1.1, we filtered the data to include only

sites that were present and had a quality determination of PASS in both the genomes and exomes

(Karczewski et al. 2020). To ensure sufficient power, we filtered for sites with allele count (AC) >10 and ran a

Fisher's Exact Test (FET) on the difference in the number of alternate AC to total alleles (allele number, AN)

between these two datasets. A non-negligible fraction of sites were significantly discordant in their calls

(**Figure** 1A). We also tested other less stringent AC thresholds (AC>1 and AC>5) and observed that the

trends of discordance between the genotype discovery approaches were consistent across AC cutoffs

(**Supplemental Figure S**1).

When comparing allele frequencies between sequencing strategies it is critical to control for ancestry, as

populations will have differing frequencies at many loci simply due to demography (Auton and Salcedo 2015;

Gravel et al. 2011; Bergström et al. 2020). To assess if ancestry affected discordance rates, we ran FET

concordance tests across all gnomAD continental groups (**Figure 1B**). The total discordant variant counts

were directly related to the sample size of the population in question and were not enriched for any given

ancestry (**Supplemental Figure S2**). While novel discordant variants were discovered in each population,

most variants observed in other populations were shared with the NFE (86.9%, **Figure 1B, Supplemental**

Table S1). Replication of the same variants across multiple ancestries strengthens the argument of a shared

technical artifact and suggests that ancestral bias between exome and genome datasets is unlikely to be a

confounding factor. Future work may wish to investigate other biological and non-biological factors for an

impact on discordance.

Error mode of discordant calls

Next, we aimed to classify the error mode that results in discordant calls using individuals with both

WES and WGS data. As 946 gnomAD individuals underwent both whole exome and whole genome

sequencing, we were able to examine the rates and error modes of discordant genotype calls without

concern over population structure or differing sample composition; since these are the same individuals, any difference in AF and/or genotype calls can be conclusively determined to be due to technical artifacts. We tallied the number of sites in each pairwise exomes-genomes genotype category (6,333 sites; **Supplemental Figure S**3) to classify miscall error mode. Of the 6 possible error modes (homozygous reference/heterozygous, homozygous reference/homozygous variant, or homozygous variant/heterozygous for both dataset directions), we find that the majority of calls, 57.7%, are a heterozygous genotype call in the genomes but a homozygous reference genotype call in the exomes (**Figure 1D, Supplemental Figure S4**). We also note that different sequencing platforms have different rates of discordant calls, though due to sharing restrictions we can not identify platform names with certainty. Overall, ~16% of the variants that were present and PASS in both exomes and genomes in the overlapping individuals had at least one discordant call.

Next, we examined the discordance of allele frequencies, again with a FET. Usually in cohort-based comparisons, the expected distribution of FET *p value*s is represented by a uniform distribution. Since we are looking at the same individuals, it is expected that allele frequencies should be identical (i.e. p=1). We observe the presence of many variants substantially deviating from expectations, representing loci with significantly different MAF in the exomes vs the genomes (**Figure 1C**).

Figure 1. Discordance in genotype calls across NGS **genotype discovery approaches**. **A**) FET concordance test *p* value for shared, PASS sites in the gnomAD NFE exomes and genomes. Bars are colored by variant type: insertion (ins), deletion (del), or single-nucleotide variant (snv). **B**) 'Bad' sites are replicated across ancestry groups in gnomAD. Sites flagged as discordant in both the NFE and another ancestry group are plotted in grey, those new sites not in the NFE are shown in black. **C**) QQ plot for the FET *p* value of shared variants in a set of 946 individuals for whom both WES and WGS data was available. **D**) Different exome captures' contribution to discordant sites. Bars are colored by the error mode that was observed for the discordant genotype call: heterozygous (het), homozygous reference (HomRef) or homozygous variant (HomVar) in either the exomes (E) or genomes (G).

Identification of discordant sites

We next sought to identify problematic sites failing a FET test of concordant WGS/WES frequency

estimates in the largest subset of gnomAD, the NFE. Based on the distribution of *p* values from this test, we

decided upon a threshold of $p<1\times10^{-5}$ to determine the classification of a variant as 'bad' or 'good' (**Figure 1,**

Supplemental Figure S5). Of the 283,287 PASS/PASS variants tested with MAF>0.01 and AC>10, 51,255

(18.1%) failed the FET test and were deemed 'bad' while 231,631 (81.8%) passed and were deemed 'good'.

Distributions of metadata features for the good vs bad sites do show trends in several features, though no

feature alone perfectly explains the phenomenon (**Supplemental Figure S6**). They also highlight a

difference in discordance patterns of indels vs SNVs (**Supplemental Figure S7**). Specifically, SNVs show a

pattern of higher AF in genomes as compared to exomes, while indels do not have this trend. Indels are also

generally less stable in allele frequency estimates than SNVs. It therefore appears that two distinct technical

error modes might be affecting miscalls in indels vs SNVs, rather than one shared mechanism. As many of

these indels fell in the low complexity regions of the genome, it is likely that a mapping issue is responsible

for their miscalls. A comprehensive description of gnomAD structural variant calling and considerations is

published and can be found in a gnomAD blog post (Collins et al. 2020;

https://gnomad.broadinstitute.org/news/2019-03-structural-variants-in-gnomad/). To correct this, we therefore

recommend excluding the low complexity regions from stringent analyses. In general, when there was

discordance, the genomes were found to have a higher MAF than the exomes (**Supplemental Figure S1B**).

The trend in MAF difference aligns with the most commonly observed error mode in genotyping.

Having confirmed that there was a systematic and significant AF discordance between genotype

discovery approach, we used our FET tests to generate a list of sites harboring this technical artifact that

may be excluded from analyses. Again, these blacklisted sites represent variants that were a PASS in

gnomAD QC in both the exomes and genomes, but are unreliably genotyped depending on the sequencing

technology used. Given a situation where, for example, a case cohort has been exome sequenced while the

control cohort has been genome sequenced, such sites could give false-positive associations due to the

resulting AF differences. We, therefore, recommend for them to be treated with caution or broadly excluded

(in addition to standard cohort QC) unless thorough confirmation of their validity in a particular dataset has

been done.

Our analysis of variant allele frequency discordance reflects technical differences between whole exome

and whole genome next-generation sequencing approaches in recovering coding DNA variation. Similarly,

we performed this concordance analysis in the All of Us Research Program dataset to compare allele

frequencies between whole genome sequencing and microarray genotyping to quantify any similar effect

arising between these genotype discovery approaches in primarily non-coding variation.

We subsampled the All of Us primary release cohort down to the 95,596 samples who have both WGS and

microarray genotyping data available. Call rate, HWE and MAF > 0.05 filters were applied to ensure only good

quality common variants entered the analysis. Out of 102,631 variants (7,944 coding) - 2,344 had Fisher's

exact test $P < 0.05$ (Supplemental File S1). Note that due to identical samples being analyzed, the expected p-

value distribution is centered at 1; **Supplemental Figure S8**. We evaluated the overlap between the variants

flagged in All of Us and gnomAD, finding that only 7 out of them were found in both samples, likely due to the

focus of gnomAD on coding variation (given comparisons included WES) versus on non-coding variation in All

of Us. Out of these 7 variants, rs4951250 was found to be significantly discordant in both datasets ($P<1\times10^{-16}$

genome vs exome; $P=4\times10^{-5}$ genome vs array).*Recovering filtered concordant sites*

In addition to generating this blacklist of bad sites that should be excluded despite being a PASS in

gnomAD QC, we investigated whether additional trustworthy sites could be rescued from the 'Non-PASS' list

based on our AF concordance criteria. Non-PASS variants are those that did not meet all required passing

criteria in the gnomAD QC pipeline (Karczewski et al. 2019). We tested this by conditioning on PASS in one

dataset, Non-PASS in the other, and re-ran the concordance pipeline requiring the following threshold in both

datasets AC>1, DP > 10, and AF>0.01% to add a higher level of stringency for recovering sites. In total,

there were 41,584 sites that met these criteria of which 30,683 were instances where the genomes are a

Non-PASS and the exomes are a PASS. Approximately half of these sites had *p* values greater than 1×10^{-5},

which we consider to be reliable. The exomes represent the vast majority of sequences in gnomAD, which

may make their results more stable. For analyses that require less stringent QC, we provide these sites

which can be optionally retained, given that they pass cohort QC in the individual dataset.

Predicting technical bias for variants.

We used features based on variant annotations generated during variant calling (e.g. variant quality,

mapping quality, etc.) to build a random forest predictor that detects the presence of technical bias for a

particular variant (**Supplemental Table S3**). Model training and validation was performed using several

approaches. First, we used a leave-one-out cross validation procedure using the exome dataset from

gnomAD. In 22 trials (one for each of the 22 autosomes) we set aside one chromosome and used the other

21 for model training. Then, the model was tested on the variants from the chromosome that was not used

for training. Using the FET p value threshold for the discordance analysis we classified the variants into 'bad'

(with $p <$ threshold) and 'good' ($p >$ = threshold) groups. By varying the FET p value threshold discriminating

the groups we performed ROC analysis. Since representation of the classes varies depending on the

selected threshold, we used the class weights for balancing the classes (in this and the following tests the

weight for the 'bad' class was set as the fraction of 'bad' variants in the training data and the weight for the

'good' class was set to 1). The area under the ROC curve (ROC AUC) for such a model was estimated as

0.841 (**Figure 2A**).

Next, we used variant annotations from the gnomAD genomes dataset for training and variant

annotations from the gnomAD exomes as a test sample. These are two separate datasets that are well-

powered to detect biases in allele frequencies and ensure full independence between test and training

samples. In this setting our model again reliably predicted discordant variants, with ROC AUC=0.803

(**Figure 2A**). Feature importance analysis of the model suggests that variant quality, inbreeding coefficient,

and quality by depth are the key parameters discriminating variants with and without evidence for technical

bias. Therefore, current protocols for alignment and variant calling are leaving a notable footprint which can

be used to detect platform biases (**Figure 2B**).

Finally, we used data from the 1000 Genomes Project (1000 Genomes Project Consortium et al. 2012)

as an independent public dataset for testing the predictor. Exome sequences from 1,393 samples

(**Supplemental Table S4**) were used to create the variant call set following GATK best practices. Variant

annotations were used to classify variants using the gnomAD genomes data as a training sample. Due to

sample size, the 1000 Genomes data is significantly less powered to detect technical biases compared to

gnomAD. Therefore, it is harder to confidently identify ground truth for discordant variants. Identical 1000

Genomes samples from exome and genome sequencing were used to detect variants with a signature of

technical bias using the AF concordance FET described here (**Supplemental Table S5**). Due to the power

limitations, instead of performing ROC analysis we compared the FET p values for the variants classified as

having evidence of technical bias ('bad') and those without such an evidence ('good', **Fig 2C**).

Figure 2. Reliable classification of discordant variants based on variant annotations. A) ROC analysis

for two random forest predictor validation approaches using either leave-one-out analysis on the exomes, or

the genomes as a training set to classify exome variants; **B**) Feature importance analysis for the random forest

model; **C**) Comparison of concordance analysis FET *p* values for variants from 1000 Genomes classified using

the random forest predictor trained on the gnomAD genomes dataset. MeanDecreaseGini is a measure of how

each variable contributes to the homogeneity of the nodes and leaves in the resulting random forest. The

This random forest model trained on the gnomAD genome v2.1 dataset was incorporated into a freely

distributed R package called *DNAdiscover* - DNA DISCOrdant Variant IdentifiER

(https://github.com/na89/DNAdiscover). The package uses variant annotations to predict whether a variant is

likely to be 'discordant' or 'concordant' in user input data and performs well with both genome and exome

sequencing data.

Discordant variants are reported in published studies and are predicted to be functionally important

We first observed the phenomenon of variant discordance through the investigation of GWAS variants in

the COVID host genetics initiative and the UK Biobank (COVID-19 Host Genetics Initiative 2021). When

inspecting top associated variants that did not have strong LD friends, we noticed that many had discordant

frequencies between GWAS arrays and gnomAD, and this often corresponded to variants that had

discrepant frequencies between the gnomAD exomes and genomes. We thus suggest that the blacklist

provided herein can be used for quality control of GWAS variants.

To investigate whether discordant variants may be spuriously attributed phenotypic relevance, we

annotated all variants with their predicted functional consequence using the Ensembl Variant Effect Predictor

(VEP) (McLaren et al. 2016). Blacklisted variants appear in all functional consequence categories, including

11536 that are annotated as missense (Supplemental Figure S9, Supplemental Table S5). Such variants are

tempting to prioritize for functional follow-up given their apparent functional importance despite having GWAS

signal likely driven by the observed genotype calling artifact. Technical artifacts are actually expected to be

enriched in functionally important categories given that they are immune from the effects of natural selection,

a phenomenon that has been previously observed for putatively loss-of-function somatic variation (Buckley et

al. 2017).

To see if blacklist sites have been reported in peer-reviewed publications, we intersected genome-wide

significant variants ($p < 5 \times 10^{-8}$) from the GWAS catalog (Welter et al. 2014) with our discordant sites list.

Seventeen bad variants were found in the GWAS catalog, underscoring the importance of controlling for this

artifact, as it may impact downstream interpretation of association findings (Supplemental Table S2).

Variants in this list have been associated with multiple health-related phenotypes including schizophrenia,

telomere length, and blood protein levels. Of these 17, half are multi-allelic and approximately a third are

indels, echoing our earlier results of a higher discordance rate for these variant types than bi-allelic SNVs.

Additionally, more than half of the 17 are present in the first megabase of the chromosome, suggesting that

areas flanking the telomeres should be treated with caution.

Discussion

The need for extremely large sample sizes to obtain sufficient statistical power in genetic studies

requires the creation of datasets that may go beyond the financial capabilities of many individual research

groups. This leads to the creation of meta-datasets that have contributions from many individual studies, thus

creating heterogeneity in the genotype discovery approaches that were used for genotyping. Therefore,

identification of DNA variants that are susceptible to technical bias when genotypes originate from multiple

discovery strategies is vital in order to avoid false-positive associations and analyses of the artificially inflated

allele frequencies. This is of particular concern in instances where cases may originate from one data

generation effort and controls from another.

Here, we identify and describe a technical artifact arising in various genotype discovery approaches that

may affect cohort data variant quality despite the following of gold standard QC procedures. We present our

metric for the identification of discordant sites, provide a blacklist of the discordant variants identified in

gnomAD which should be treated with caution, and release an openly-available software package containing

our random forest predictor that reliably classifies untrustworthy variants in user cohort data. Excluding

variants with signals of discordance across sequencing platforms will result in higher quality results and

reduces the risk of spurious associations in gene discovery. This is particularly important as we observe that technical artifacts are enriched in functionally important annotations.

Additionally, we show that discordance in allele frequencies is also present in the All of Us Research Program dataset, when comparing whole genome sequencing to microarray genotyping for overlapping samples. This finding indicates that variants in both coding and non-coding DNA could have discordant genotype calls. Importantly, in our predictor we use the variant annotations which often are utilized in variant quality score recalibration and filtration pipelines. Our results indicate that stricter filtration thresholds might be helpful for elimination of some discordant variants, however, more cautious consideration of discordance is warranted in heterogeneous datasets.

We note that while we provide a blacklist of variants failing our discordance test in the gnomaD v2 dataset for ready exclusion, the specific sites that are discordant in a given cohort depends on the genotype discovery approach utilized and dataset composition. Therefore, for optimal precision, we recommend identification of discordant sites within user cohorts with the provided classifier rather than a blanket restriction of variants identified in gnomAD. We freely provide an R-package with a predictor trained on gnomAD whole genome sequencing data, *DNAdiscover*, for such use in other cohort data to identify cohort-specific sites with features indicative of unreliable genotype calls.

Based on the examinations presented in this manuscript, we recommend researchers using aggregated cohort data to implement the following conservative QC procedures to ensure the elimination of discordant sites:

- Drop any variant that fails in both the gnomAD exomes and genomes
- Consider dropping any variant that fails in the gnomAD exomes, as these represent the bulk of gnomAD data
- Drop the blacklist variants presented here that are PASS in both the gnomAD exomes and genomes but that are discordant in frequency across genotype discovery approach
- Drop variants that are flagged by our random forest predictor, *DNAdiscover,* in an independent dataset, as each genotype discovery approach has a distinct genotyping error mode
- Remove the low complexity regions

●   Optional: skeptically retain sites that are on the 'recovered' list here

Methods

Characterizing discordance in genotype calls across gnomAD exomes and genomes

All analyses were conducted using the Hail software program on the Google Cloud platform (GCP

2021). Plots were created using Bokeh and ggplot2 (Jolly 2018; Wickham 2011). Concordance metrics for

genotype calls were generated from the overlapping individuals with the command

hail.methods.concordance(). Using the full release of gnomAD version 2.1 (Karczewski et al. 2020), we

filtered to include only sites that were both present and had a quality determination of PASS in the genomes

and exomes. We split multi-allelic variants, retained only sites that are present and PASS in both exomes

and genomes, filtering to only bi-allelic sites with AC > 1 and AF > 0.01% in either dataset for the NFE.

Starting with all sites with at least 1 alternate allele, and subsequently for sites with AC>5 and 10, we

calculated the allele frequency (AF) in the genomes and exomes separately and ran a Fisher's Exact Test

(FET) on the difference in the number of alternate AC to total alleles (allele number, AN) between these two

datasets. Specific filters for various steps are described in their relevant *Results* section.

AllofUs data was subsampled to 95,596 with both whole genome sequencing (WGS) and microarray

genotyping (WGA) available. MAF>0.05, HWE > 0.0001 and MAC>10 filters were applied to keep only

common variants. Multiallelic variants were split. Variant call rate > 0.8 was required in both WGS and WGA

datasets. Variants with call rate difference between datasets greater that 0.05 were also eliminated from

analysis. The R libraries *dplyr, reshape2, pROC, ROCR* and *RandomForest* were used to process variant

annotations, evaluate predictor quality and build a DNAdiscover using R-4.0.3 (Wickham et al. 2015;

Wickham 2012; Robin et al. 2011; Sing et al. 2005; Liaw et al. 2002).

Acknowledgements

We thank the members of the gnomAD and Hail teams for their assistance with this project. This project was supported by the National Institute of Mental Health (K01 MH121659 and T32 MH017119 to E.G.A.) and the National Human Genome Research Institute (U24HG011450). E.G.A was additionally supported by the Caroline Wiess Law Fund for Research in Molecular Medicine and the ARCO Foundation Young Teacher-Investigator Fund at Baylor College of Medicine. A.A.L. was supported by the Ministry of Science and Higher Education of the Russian Federation (Agreement No. 075-15-2022-301, Institutional grant to Almazov National Medical Research Center). M.A. was supported by the Aging Biology Foundation and Nationwide Foundation Pediatric Innovation Fund. The All of Us Research Program is supported by the National Institutes of Health, Office of the Director: Regional Medical Centers: 1 OT2 OD026549; 1 OT2 OD026554; 1 OT2 OD026557; 1 OT2 OD026556; 1 OT2 OD026550; 1 OT2 OD 026552; 1 OT2 OD026553; 1 OT2 OD026548; 1 OT2 OD026551; 1 OT2 OD026555; IAA #: AOD 16037; Federally Qualified Health Centers: HHSN 263201600085U; Data and Research Center: 5 U2C OD023196; Biobank: 1 U24 OD023121; The Participant Center: U24 OD023176; Participant Technology Systems Center: 1 U24 OD023163; Communications and Engagement: 3 OT2 OD023205; 3 OT2 OD023206; and Community Partners: 1 OT2 OD025277; 3 OT2 OD025315; 1 OT2 OD025337; 1 OT2 OD025276. In addition, the All of Us Research Program would not be possible without the partnership of its participants.

Author Contributions

E.G.A. and M.A. designed and implemented pipelines, ran analyses, and drafted the primary manuscript. K.J.K., A.A.L. aided in code implementation and interpretation. H.R., D.M., B.M.N. and M.J.D. supervised and advised on the project. All authors reviewed and approved the final draft.

Data Access

gnomAD summary data is freely available at https://gnomad.broadinstitute.org/. Additional information and a discussion of the best practices for using gnomAD can be found at https://macarthurlab.org/blog/.

DNAdiscover package for prediction of presence of the technical bias in variants coming from NGS and

usage manual are available at https://github.com/na89/DNAdiscover. A blacklist of bad variants failing the

discordance test in gnomAD is provided with this manuscript in the Supplementary Materials.

Competing Interests statement

M.J.D. is a founder of Maze Therapeutics. B.M.N. is a member of the Deep Genomics Scientific Advisory

Board and serves as a consultant for the Camp4 Therapeutics Corporation, Takeda Pharmaceutical and

Biogen. K.J.K is a consultant for Vor Biopharma. The remaining authors declare no competing interests.

References

1000 Genomes Project Consortium, Abecasis GR, Auton A, Brooks LD, DePristo MA, Durbin RM, Handsaker RE, Kang HM, Marth GT, McVean GA. 2012. An integrated map of genetic variation from 1,092 human genomes. *Nature* **491**: 56–65.

Adelson RP, Renton AE, Li W, Barzilai N, Atzmon G, Goate AM, Davies P, Freudenberg-Hua Y. 2019. Empirical design of a variant quality control pipeline for whole genome sequencing data using replicate discordance. *Sci Rep* **9**: 16156.

The All of Us Research Program Investigators. 2019. The "All of Us" Research Program. *New England Journal of Medicine*, *381*(7), 668–676.

Anderson CA, Pettersson FH, Clarke GM, Cardon LR, Morris AP, Zondervan KT. 2010. Data quality control in genetic case-control association studies. *Nat Protoc* **5**: 1564–1573.

Auton A, Salcedo T. 2015. The 1000 Genomes Project. *Assessing Rare Variation in Complex Traits* 71–85. http://dx.doi.org/10.1007/978-1-4939-2824-8_6.

Bergström A, McCarthy SA, Hui R, Almarri MA, Ayub Q, Danecek P, Chen Y, Felkel S, Hallast P, Kamm J, et al. 2020. Insights into human genetic variation and population history from 929 diverse genomes. *Science* **367**. https://science.sciencemag.org/content/367/6484/eaay5012/tab-pdf (Accessed March 19, 2020).

Buckley AR, Standish KA, Bhutani K, Ideker T, Lasken RS, Carter H, Harismendy O, Schork NJ. 2017. Pan-

cancer analysis reveals technical artifacts in TCGA germline variant calls. *BMC Genomics*. Jun 12;18(1):458.

doi: 10.1186/s12864-017-3770-y.

Collins RL, Brand H, Karczewski KJ, Zhao X, Alföldi J, Francioli LC, Khera AV, Lowther C, Gauthier LD,

Wang H, et al. 2020. A structural variation reference for medical and population genetics. *Nature* **581**: 444–

451.

COVID-19 Host Genetics Initiative. 2021. Mapping the human genetic architecture of COVID-19. *Nature*.

http://dx.doi.org/10.1038/s41586-021-03767-x.

Gravel S, Henn BM, Gutenkunst RN, Indap AR, Marth GT, Clark AG, Yu F, Gibbs RA, 1000 Genomes

Project, Bustamante CD. 2011. Demographic history and rare allele sharing among human populations. *Proc*

Natl Acad Sci U S A **108**: 11983–11988.

Highnam G, Wang JJ, Kusler D, Zook J, Vijayan V, Leibovich N, Mittelman D. 2015. An analytical framework

for optimizing variant discovery from personal genomes. *Nat Commun* **6**: 6275.

Jolly K. 2018. *Hands-On Data Visualization with Bokeh: Interactive web plotting for Python using Bokeh*.

Packt Publishing Ltd.

Karczewski K. 2017. The genome Aggregation Database (gnomAD). *MacArthur Lab*.

https://macarthurlab.org/2017/02/27/the-genome-aggregation-database-gnomad/ (Accessed May 13, 2020).

Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, Collins RL, Laricchia KM, Ganna A,

Birnbaum DP, et al. 2020. The mutational constraint spectrum quantified from variation in 141,456 humans.

Nature **581**: 434–443.

Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, Collins RL, Laricchia KM, Ganna A,

Birnbaum DP, et al. 2019. Variation across 141,456 human exomes and genomes reveals the spectrum of

loss-of-function intolerance across human protein-coding genes. *bioRxiv* 531210.

https://www.biorxiv.org/content/10.1101/531210v2 (Accessed May 13, 2020).

Lam M, Awasthi S, Watson HJ, Goldstein J, Panagiotaropoulou G, Trubetskoy V, Karlsson R, Frei O, Fan C-

C, De Witte W, et al. 2019. RICOPILI: Rapid Imputation for COnsortias PIpeLIne. *Bioinformatics*.

http://dx.doi.org/10.1093/bioinformatics/btz633.

Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, O'Donnell-Luria AH, Ware JS, Hill AJ,

Cummings BB, et al. 2016. Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**: 285–

291.

Liaw A, Wiener M, Others. 2002. Classification and regression by randomForest. *R news* **2**: 18–22.

Li J, Jew B, Zhan L, Hwang S, Coppola G, Freimer NB, Sul JH. 2019. ForestQC: Quality control on genetic

variants from next-generation sequencing data using random forest. *PLoS Comput Biol* **15**: e1007556.

Ma X, Shao Y, Tian L, Flasch DA, Mulder HL, Edmonson MN, Liu Y, Chen X, Newman S, Nakitandwe J, et

al. 2019. Analysis of error profiles in deep next-generation sequencing data. *Genome Biol* **20**: 50.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D, Gabriel

S, Daly M, et al. 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation

DNA sequencing data. *Genome Res* **20**: 1297–1303.

McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, Flicek P, Cunningham F. 2016. The

Ensembl Variant Effect Predictor. *Genome Biol* **17**: 122.

Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez J-C, Müller M. 2011. pROC: an open-source

package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics* **12**: 77.

Sing T, Sander O, Beerenwinkel N, Lengauer T. 2005. ROCR: visualizing classifier performance in R.

Bioinformatics **21**: 3940–3941.

Welter D, MacArthur J, Morales J, Burdett T, Hall P, Junkins H, Klemm A, Flicek P, Manolio T, Hindorff L, et

al. 2014. The NHGRI GWAS Catalog, a curated resource of SNP-trait associations. *Nucleic Acids Res* **42**:

D1001–6.

Wickham H. 2011. ggplot2. *Wiley Interdiscip Rev Comput Stat* **3**: 180–185.

Wickham H. 2012. reshape2: Flexibly reshape data: a reboot of the reshape package. *R package version* **1**.

http://cran.ms.unimelb.edu.au/web/packages/reshape2/.

Wickham H, Francois R, Henry L, Müller K, Others. 2015. dplyr: A grammar of data manipulation. *R package*

version 0 4 **3**: 156.

Google Cloud Computing, Hosting Services & APIs. *Google Cloud*. https://cloud.google.com/gcp/ (Accessed

May 13, 2020a).

Hail. https://hail.is/ (Accessed May 13, 2020b).