1

siRNA-mediated gene knockdown via electroporation in hydrozoan jellyfish embryos

[1] Tokiha Masuda-Ozawa, [2] Sosuke Fujita**,** [3] Ryotaro Nakamura, [3] Hiroshi Watanabe, [2] Erina Kuranaga, and [1,4] Yu-ichiro Nakajima*****

1. Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, Sendai, Japan
2. Graduate School of Life Sciences, Tohoku University, Sendai, Japan
3. Evolutionary Neurobiology Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan
4. Graduate School of Pharmaceutical Sciences, The University of Tokyo

*Author for correspondence e-mail: nakaji97@g.ecc.u-tokyo.ac.jp

Tel: +81-3-5841-4863

Running title: **siRNA gene knockdown in jellyfish embryos**

Main text: 4361 words (Abstract: 247 words)

Figures: 6   **Supplementary information** (Figures: 5; Table: 2)

Keywords: siRNA, electroporation, cnidarian, hydrozoan, jellyfish, *Clytia, Cladonema*

Abstract

As the sister group to bilaterians, cnidarians stand in a unique phylogenetic position that provides insights into evolutionary aspects of animal development, physiology, and behavior. While cnidarians are classified into two types, sessile polyps and free-swimming medusae, most studies at the cellular and molecular levels have been conducted on representative polyp-type cnidarians and have established techniques of genetic manipulation. Recently, gene knockdown by delivery of short hairpin RNAs into eggs via electroporation has been introduced to two polyp-type cnidarians, *Nematostella vectensis* and *Hydractinia symbiolongicarpus*, enabling systematic loss-of-function experiments. By contrast, current methods of genetic manipulation for medusa-type cnidarians, or jellyfish, are quite limited, and reliable techniques are required to interrogate function of specific genes. Here, we present a method to knockdown target genes by small interfering RNA (siRNA) delivery into fertilized eggs via electroporation, using the hydrozoan jellyfish, *Clytia hemisphaerica* and *Cladonema paciificum*. We show that siRNAs targeting endogenous *GFP1* and *Wnt3* in *Clytia* efficiently knockdown gene expression and result in known planula phenotypes: loss of green fluorescence and defects in axial patterning, respectively. We also successfully knock down endogenous *Wnt3* in *Cladonema* by siRNAs electroporation, which circumvents the technical difficulty of microinjecting small eggs. Unexpectedly, *Wnt3* knockdown in *Cladonema* does not induce morphological defects in planula, suggesting a distinct mechanism that operates during embryogenesis. Our method for electroporation of siRNAs targeting specific genes is applicable to other animals, including and beyond jellyfish species, and will facilitate the investigation and understanding of myriad aspects of animal development.

Introduction

The phylum Cnidaria, as the sister group to Bilateria, separated from their common ancestor over 500 million years ago. Cnidarians have diversified their morphologies in the forms of corals, sea anemones, hydroids, and jellyfish, all of which are divided into two clades, Anthozoa and Medusozoa; The two differ in that Anthozoa includes only sessile polyp-type animals, while Medusozoa (Hydrozoa, Staurozoa, Scyphozoa, and Cubozoa) includes two forms: polyp and medusa, or jellyfish stage (Fig. 1A)[1,2]. The life cycle of most, though not all, jellyfish includes five forms: gametes, fertilized eggs, planulae, polyps, and medusae[3]. Fertilized eggs undergo embryogenesis to become planula larvae, which metamorphose into sessile polyps. While vegetatively-growing polyps give rise to free-swimming medusae through the process of budding or strobilization, medusae sexually reproduce by releasing gametes. Due to their unique phylogenetic position, studies using cnidarians have provided evolutionary insights into development, regeneration, and behaviors in multicellular animals[1,4,5]. To date, the field's molecular and cellular understanding of cnidarians has been acquired primarily using polyp-type animals such as the anthozoan *Nematostella vectensis* and the hydrozoan *Hydra* and *Hydractinia*. By contrast, jellyfish biology remains largely unestablished.

Among jellyfish species, the hydrozoan *Clytia hemisphaerica* is the best-studied laboratory model[6,7]. The well-established method for maintaining and propagating animals ensures a reliable resource for daily experiments[8]. While the transparent and small-sized medusae allow the whole-mount visualization, relatively large-sized eggs (~180 μm diameter) enable different manipulations with microinjection (Fig. 1B). Indeed, gene silencing by morpholino oligos and mRNA microinjection as well as gene knockout via CRISPR/Cas9 allow for the investigation of the functions of genes of interest[9-13]. Furthermore, the *Clytia* genome assembly is complete, and transcriptome profiles are available for various stages and tissues, even at the single-cell level, which creates the foundation for research at the molecular level[12,14-18]. These resources and techniques, together with recently-established transgenesis[19], accelerate research into different features associated with distinct *Clytia* life stages. Current research topics using *Clytia* range from the establishment of the oral-aboral axis during embryogenesis and

gametogenesis regulated by the light-dark cycle to body regeneration, nervous system
development, and characterizing adult medusa behavior[9,10,12,17,19,20].

By contrast, the hydrozoan *Cladonema pacificum* is an emerging model jellyfish
that has been recently utilized for various studies in development, regeneration, and
physiology[21]. The ease of rearing all stages of *Cladonema* without a filtration system or
large water tank, along with their high spawning rate, enables easy lab maintenance.
The small-sized medusae with branched tentacles allow for investigations of body size
control and tentacle morphogenesis (Fig. 1C)[22-24]. *Cladonema* gametogenesis is
regulated by the light-dark cycle as in *Clytia*, and recent work has identified the
neuropeptides involved in oocyte maturation[25]. Furthermore, *Cladonema* medusae
possess primitive complex eyes (ocelli) in their tentacle bulbs, and studies using the
closely related species *Cladonema radiatum* have identified conserved light-sensitive
opsins and regulators of eye formation (*Pax*, *Six*, and *Eya*)[25-29], providing a model for
the evolutionary developmental biology of photoreceptor organs. Despite these
attractive features of *Cladonema* as a laboratory animal, no genome assembly or
transcriptome is currently available, and genetic manipulation techniques are only just
being developed. One major technical issue in manipulating *Cladonema* is their small
eggs—with a diameter of approximately 60 μm (Fig. 1C), regular microinjection is quite
difficult. Establishing genetic manipulations is required to facilitate the in-depth
investigations needed to further understand the biology of *Cladoenema*, and an
alternative method that eliminates need for microinjection would greatly facilitate that
objective.

RNA interference (RNAi), the phenomenon of double stranded RNA (dsRNA)
mediated silencing of target genes, has been widely exploited in living organisms to
analyze gene function[30], primarily using chemically- or *in vitro*-synthesized double-
stranded small interfering RNAs (siRNAs) or vector-based short hairpin RNAs
(shRNAs)[31]. In cnidarians, siRNA-mediated gene silencing was initially applied to
Nematostella and *Hydra* polyps via soaking or electroporation[32-34]. Additionally, gene
knockdown via shRNA microinjection or electroporation into eggs has recently been
utilized in studies using *Nematostella* and *Hydractinia* to show efficient reduction in
gene expression and associated phenotypes in early developmental stages[35-38]. In
particular, shRNA delivery via electroporation, which does not require the rigors of

microinjection, allows for the experimental gene knockdown of large numbers of

individuals simultaneously[36,38], opening up the possibility of manipulating genes in

different aquatic animals, even those that produce very small eggs like *Cladonema*.

The mechanism of RNAi-mediated gene repression after introducing foreign

dsRNA into cells involves the microRNA (miRNA) pathway, an endogenous gene

repression machinery conserved in both animals and plants[39]. In cnidarians, the

presence of the miRNA pathway has been demonstrated in *Nematostella* and

Hydra[32,40,41]. However, little is known about the endogenous RNAi pathway, especially

the presence and roles of miRNAs and the miRNA-related genes, in jellyfish such as

Clytia and *Cladonema*. Furthermore, in mammalian cultured cells, siRNAs can induce

gene repression independent of endogenous dsRNA processing factors, while shRNAs

cannot[42,43]. On this basis, we elected to use siRNA instead of shRNA for gene

knockdown to avoid the potential pitfall of dsRNA processing in jellyfish.

Here we report a gene knockdown method for jellyfish embryos with siRNA via

electroporation. Using two hydrozoan species, *Clytia hemisphaerica* and *Cladonema*

pacificum, we demonstrate that siRNA delivery into fertilized eggs effectively reduces

the expression of endogenous genes. We also confirm the known loss-of-function

phenotypes in *Clytia* after knocking down *GFP1* or *Wnt3*, which are induced in a dose-

dependent manner with siRNA. We further show that knockdown of *Wnt3* in *Cladonema*

embryos does not cause defects in planula morphology, implicating a potentially distinct

embryogenesis regulating mechanism between these two hydrozoan species. Overall,

our siRNA-mediated knockdown approach allows for the manipulation of a large number

of embryos through electroporation and enables functional analysis of early

development, providing a new experimental platform applicable to different jellyfish

species and other marine invertebrates.

Materials and Methods

Animal culture and spawning induction

Clytia hemisphaerica (Z11, Z4B as females and Z4C2, Z23 as males) and *Cladonema*

pacificum (6W as females and UN2 as males) were used for this research.

Clytia hemisphaerica were cultured using a previously reported method[8] with a

few modifications. Artificial sea water (ASW) was prepared using 220 g SEA LIFE

(Marin Tech) per 5L MilliQ water (Merk Millipore) with antibiotics (40 units/ml of penicillin

and 40 µg/ml of streptomycin). Medusae, embryos, and planula larvae were maintained

at 20ºC. Medusae were fed everyday with Vietnamese brine shrimp (A&A Marine LLC,

Elk Rapids, MI, USA). Spawning timing was controlled by a 13 hours (h) dark /11 h light

cycle, and spawning was induced by light. Male and female medusae were transferred

into V-7 cups (AS ONE) before spawning (60 min for male and 90 min for female after

light stimulation).

Cladonema pacificum were cultured as previously described[23]. *Cladonema*

medusae were maintained at 22ºC in ASW, which was prepared by SEA LIFE (Marin

Tech) dissolved in tap water with chlorine neutralizer (Coroline off, GEX Co. ltd) (24

p.p.t) and antibiotics (40 units/ml penicillin and 40 µg/ml of streptomycin). Spawning

timing was controlled by a 30 min dark/23.5 h light cycle, and spawning of male and

female gametes was induced by dark stimulation. Before dark stimulation, adult females

and males were separately transferred into V7 cups (AS ONE) and 60 mm dishes (BD),

respectively.

Nematostella vectensis were cultured as previously described[44], with a few

modifications. Briefly, adult animals were maintained in brackish water at a salinity one-

third of artificial seawater (35 g/l, pH 7.5-8.0, SEA LIFE, Marin Tech) and fed with

freshly hatched artemia two times per week. Spawning induction was performed at 26°C

under light for at least 11 h.

siRNA and shRNA

siRNA sequences were designed based on CDS sequences and synthesized by

manufacturers (Nippon gene Co.Ltd and Sigma-Aldrich, Merk). shRNAs were

synthesized as described in previous reports[36,38]. Briefly, shRNAs were synthesized by

in vitro transcription (IVT) from double stranded DNA templates using the

AmpliScribeTM T7-flashTM transcription kit (Lucigen, Inc.) and were purified using

Direct-Zol TM RNA Miniprep kits (Zymo Research, R2070). Concentrations of shRNAs

were measured by NanoDrop One (Thermo fisher).

Collection of gametes and electroporation for *Clytia* and *Cladonema* fertilized eggs

For preparation of gametes, sexually-mature *Clytia* medusae were treated following a previously reported method[8], partially modified to fit our experimental equipment. Adult medusae were maintained on a 13 h dark/11 h light cycle, and light stimuli induced gametogenesis in 60 min for sperms and 90 min for eggs (Fig. 2A). For *Cladonema pacificum*, sexually-mature medusae were maintained on a 30 min dark/23.5 h light cycle. Dark stimulation induced gametogenesis in 25 min. Unfertilized eggs were collected and resuspended in 15% Ficoll (Nacalai tesque, Japan) in ASW to prevent the eggs settling from at the bottom of the microtube or sticking to the microtube surface, keeping the egg solution homogeneous (Fig. 2B)[36,38].

Unfertilized *Clytia* and *Cladonema* eggs were collected in dishes and transferred into 1.5 ml tubes coated with Ficoll (Nacalai tesque, Japan) to prevent the eggs from sticking to the tubes. Sperm water was added into eggs, and eggs and sperm were incubated for 5 min at room temperature (20-22ºC). The sperm water was then removed, and fertilized eggs were resuspended in 15% Ficoll in ASW, at which point siRNAs or shRNAs were added into fertilized egg solution. The fertilized eggs and RNA mixture were carefully transferred into a 4 mm cuvette and placed into the shockpod cuvette chamber connected to the Gene Pulser Xcell (Bio-Rad). In most experiments, the Gene Pulser Xcell was used for pulsing with a square wave voltage of 50 V and a single pulse duration of 25 ms, except for optimizing electroporation conditions (Supplementary Figs 1-2). After electroporation, fertilized eggs were transferred to a dish, and incubated for 10 min. The 15% Ficoll solution was removed and replaced with new ASW, and the samples were incubated at 20-22ºC.

Collection of egg sacs, dejellying, and electroporation for *Nematostella* fertilized eggs

The collection of egg sacs, dejellying, and electroporation steps were performed as previously described[36] with several modifications. Briefly, unfertilized egg sacs were

incubated in culture media with sperm at 20-25ºC for 30 min. The fertilized egg sacs

were then dejellied in freshly prepared L-Cysteine solution (0.04 g/ml, Nacalai tesque, in

brackish water; pH adjusted by 5N NaOH at 7.5-8.0) on a shaker for 10 min. The

dejellied eggs were rinsed with brackish water before they were suspended in Ficoll

solution (Sigma-Aldrich, in brackish water) at final concentrations of 15%. The

suspended eggs and siRNA (5 µg/µl stock solution, in MilliQ) were transferred to a 4

mm cuvette (Bio-Rad) at the final volume of 200 µl and were gently mixed. The Bio-rad

Gene Pulser Xcell was used for pulsing with a square wave voltage of 50 V and a pulse

duration of 25 ms. The samples were transferred to a 100 mm dish and maintained in

brackish water for 3 days.

Imaging

GFP and Rhodamine fluorescent imaging and bright field imaging were taken with an

Axio Zoom V16 (ZEISS), and image processing was done with ZEN 3.4 blue edition

(ZEISS) and Fiji/ImageJ software[45]. Intensity of mean values of green fluorescence was

measured using the average brightness (pixel value) of the pixels. To prevent planula

movement, we added 0.5 M sodium azide (final concentration was less than 0.25M).

Morphological quantification of planula larvae

After *Wnt3* knockdown by siRNA electroporation of *Clytia* or *Cladonema*, we

approximated the planula body shape into a two dimensional ellipse, measured both the

long and short axes (Figs. 4C and 6D), and calculated the aspect ratio of those values

using Fiji/ImageJ software (Figs 4D, 6E).

RT-qPCR

For *Clytia* and *Cladonema* samples, total RNA was extracted with RNeasy Mini or Micro

kits (Qiagen). Lysate was treated with DNaseI (Qiagen) for 15 min at room temperature

(RT). cDNA was synthesized with the PrimeScript II 1st strand synthesis kit (Takara bio).

RT-qPCR was performed with CFX connect (Bio-Rad) using iTaqTM universal SYBER

Green Supermix (Bio-Rad) or the QuantStudio 6 Flex Real-Time PCR System (Thermo

Fisher) using TB Green Premix Ex TaqII (Tli RNaseH Plus) (Takara, RR820). Gene

expressions were normalized to the housekeeping gene, *EF1alpha*, and the delta-delta-

ct method was used for quantification (CFX maestro software, Bio-Rad or QuantStudio

6 Flex Real-Time PCR System software, Thermo Fisher).

For *Nematostella* samples, total RNA was extracted using the RNeasy Mini Kit

and RNase-Free DNase Set (Qiagen). cDNA synthesis was conducted using the

SuperScript IV First-Strand Synthesis System (Thermo Fisher). RT-qPCR was

performed with the StepOnePlusTM Real-Time PCR System (Applied Biosystems,

Thermo Fisher) using PowerUP SYBR Green Master Mix (Thermo Fisher). Gene

expressions were normalized to the housekeeping gene, *NvEf1alpha*, and the delta-

delta-ct method was used for quantification (StepOne™ and StepOnePlus™ Software

v2.3).

Sequence alignment

Wnt3 amino acid sequence alignment was performed by ClustalW[46]. GenBank

accession numbers and protein names are ABG49501.1 Wnt3 [*Xenopus tropicalis*],

BAB70502.1 WNT3 [*Homo sapiens*], ABK90822.1 Wnt3 [*Gallus gallus*], ACB15465.1

Wnt3 [*Clytia hemisphaerica*], ABF48092.1 secreted signaling factor Wnt3, partial

[*Nematostella vectensis*], CAK50826.1 Wnt3 protein precursor, partial [*Hydractinia*

echinata], XP_015753662.1 PREDICTED: protein Wnt-3a-like isoform X2 [*Acropora*

digitifera], XP_015753659.1 PREDICTED: protein Wnt-3a-like isoform X1 [*Acropora*

digitifera], XP_032241388.1 protein Wnt-3a [*Nematostella vectensis*]. The alignment

data was visualized and phylogenetic tree was constructed using neighbor joining and

the BLOSUM62 matrix on Jalview software[47].

qPCR primers

Name	Sequence (5'->3')	
CpWnt3 qP FW	CAAATGTGGTCGATCAAACG	This study
CpWnt3 qP RV	TACGCCTTCTGCAACACTTG	This study
CpEF1-a_qPCR_F7	GGTCAATCTCGTTCCCTCCA	This study

CpEF1-a_qPCR_R7	TTTCCACCAGAGGTATCGGC	This study
CheGFP1 qP F	TTGCTGTCCGAATAGTGCAG	Fourrage et al., 2014 Open Biol
CheGFP1 qP RV	GACAACTCCTCCTCCGAGTG	Fourrage et al., 2014 Open Biol
CheWnt3 qP F	ATCATGGCAGGTGGAAACTC	Leclère et al., 2012 Dev Biol
CheWnt3 qP RV	CCCCATTTCCAACCTTCTTC	Leclère et al., 2012 Dev Biol
CheEF1a qP F	TGCTGTTGTCCCAATCTCTG	Leclère et al., 2012 Dev Biol / Fourrage et al., 2014 Open biol
CheEF1a qP RV	AAGACGGAGTGGTTTGGATG	Leclère et al., 2012 Dev Biol / Fourrage et al., 2014 Open biol
NvEf1alpha FW	GGTTGCCTCTTCGCTTACCACT	This strudy
NvEf1alpha RV	CGTTCCTGGCTTTAGGACAC	This study
NvBra1_Fw_1032	CGGGCTCACACTCTCACTTA	This study
NvBra1_Rv1174	CTTGCGGTATGGTGTTCCAG	This study

286 **siRNA design**

Name	Sequence (5'->3')		ORF
siCheGFP1_1 sense	CCAUCAGCUUCGAAAAUGAdTdT	This study	HQ397706.1
antisense	UCAUUUUCGAAGCUGAUGGdTdT	This study	
siCheGFP1_2 sense	UGAUGGCGCUUAUAAAGUUdTdT	This study	HQ397706.1
antisense	AACUUUAUAAGCGCCAUCAdTdT	This study	
siCheGFP1_3 sense	CGGACUCAUUCUUCAAAAAdTdT	This study	HQ397706.1
antisense	UUUUUGAAGAAUGAGUCCGdTdT	This study	
siCheWnt3_1 sense	GCAUGUGACUGUAAAUUUAdTdT	This study	EU374721.1
antisense	UAAAUUUACAGUCACAUGCdTdT	This study	
siCheWnt3_2 sense	ACAGCUUGGUAAAAUGUUAdTdT	This study	EU374721.1
antisense	UAACAUUUUACCAAGCUGUdTdT	This study	
siCpWnt3_1 sense	GUGGAACUGUAGUGUUUCAdTdT	This study	EU374721.1
antisense	UGAAACACUACAGUUCCACdTdT	This study	
siCpWnt3_2 sense	AGCGUGUGCUGAAGGUAAAdTdT	This study	EU374721.1
antisense	UUUACCUUCAGCACACGCUdTdT	This study	
siNvBra1_1 sense	GAAGAGAUCACGAGUCUAAdTdT	This study	AF540387.2
antisense	UUAGACUCGUGAUCUCUUCdTdT	This study	
siNegative_Control-1 for Nv sense	GCAACACGCAGAGTCGTAAdT	Same sequence as Karabulut et al, 2019 Dev Biol	
antisense	TTACGACTCTGCGTGTTGCdT	Same sequence as Karabulut et al, 2019 Dev Biol	

289 **shRNA template**

Name	Sequence (5'->3')	
CheWnt3_SI1_F	TAATACGACTCACTATA GCATGTGACTGTAAATTTA TTCAAGAGA TAAATTTACAGTCACATGC TT	This study

CheWnt3 _SI1_R	AAGCATGTGACTGTAAATTTATCTCTTGAATAAATTTAC AGTCACATGCTATAGTGAGTCGTATTA	This study
CheWnt3 _SI2_F	TAATACGACTCACTATA ACAGCTTGGTAAAATGTTA TTCAAGAGA TAACATTTTACCAAGCTGT TT	This study
CheWnt3 _SI2_R	AAACAGCTTGGTAAAATGTTATCTCTTGAATAACATTTT ACCAAGCTGTTATAGTGAGTCGTATTA	This study

Results

Optimization of electroporation conditions

In both hydrozoan jellyfish *Clytia hemisphaerica* and *Cladonema pacificum*, gametogenesis is regulated by light-dark transitions[6,48]. A previous report showed that *Clytia* medusae release eggs within 110-120 min or sperm within 60-90 min upon light illumination after 3-8 h of darkness[12]. *Cladonema pacificum*, which live along coastal areas in Japan, exhibit two types of gametogenesis depending on habitat: the dark-light transition (light stimulation type) and the light-dark transition (dark stimulation type), respectively. *Cladonema* medusae of the dark stimulation type, which we used in this study, release eggs or sperm within 30 min upon dark stimulus after 20-24 h of constant illumination[25,48]. By utilizing the characteristic gametogenesis of these jellyfish species, we can control egg/sperm release and fertilization at any time of day. Indeed, we routinely obtained *Clytia* sperm within 60 min and *Clytia* eggs within 90 min of light in a 13h dark/11h light cycle (Fig. 2A). Similarly, we collected gametes from *Cladonema* within 25 min of dark stimulation in a 23.5h light/30 min dark cycle (Fig. 2A).

To determine optimal electroporation parameters, we first tested different electroporation conditions and monitored the delivery efficiency of the red fluorescent dye Rhodamine-Dextran into unfertilized *Clytia* and *Cladonema* eggs (Supplementary Figs. 1-2). Given that the average molecular weight of siRNAs (21-mer, ~13,300 Da) are similar to the molecular size of Rhodamine-Dextran (10,000 Da), electroporation trials with Rhodamine-Dextran would visually confirm that small molecules are incorporated into jellyfish eggs, as previously shown in *Hydractinia* eggs[38]. The collected eggs were suspended in 15% Ficoll artificial seawater to prevent precipitation and to make a homogeneous solution (Fig. 2B)[36,38]. We performed electroporation trails using Rhodamine-Dextran using a cuvette with a 4 mm gap and tested several conditions of

voltage (V) and pulse length time (milliseconds, ms) using a conventional
electroporation system (Fig. 2C). As previous work showed that increasing the number
of pulses does not increase the Dextran delivery rate but decreases the survival rate of
embryos[38], we fixed the number of pulses to one and incubated unfertilized eggs with 1
mg/ml Dextrane-Rhodamine solution. *Clytia* eggs have little autofluorescence in the red
channel, and without electroporation (No EP), red fluorescence was rarely detected in
unfertilized eggs (Fig. 2D and Supplementary Fig. 1A). We compared the eight different
electroporation conditions and found that the rhodamine fluorescence was correlatively
detected in *Clytia* eggs depending on the strength of electric voltage, as long as the
voltage was in the range of 50-100 V（Supplementary Figs. 1B-1D）. By contrast, when
the voltage exceeded 200 V, the eggs were ruptured due to physical damage, and
cellular debris were observed（Supplementary Figs, 1E-1G）. We performed similar
electroporation trials on unfertilized *Cladonema* eggs with eight different conditions.
While red autofluorescence was negligible in the *Cladonema* eggs in the absence of
electroporation (Fig. 2D and Supplementary Fig. 2A), Rhodamine fluorescence was
frequently detected with 50 V (Fig. 2D and Supplementally Fig. 2E). In the more intense
conditions beyond 100 V (Supplementally Figs. 2F and 2G), the eggs were severely
damaged. Based on these results, we established the optimal electroporation condition
(50 V, 1 pulse, 25 ms) that allows high frequency of Dextran-positive eggs and low
levels of cell damage for both jellyfish eggs (Fig. 2D). Of note, during the initial trials, we
used unfertilized eggs, but we found that the electroporation process severely affected
the survival rate of embryos. In the case of *Cladonema*, electroporation before
fertilization severely decreased the survival rate of fertilized eggs (7.04% at 1 h post
fertilization) while electroporation after fertilization did not influence the survival rate (No
EP: 59.67%, With EP: 60.15%) (Supplementary Table 1). This dissimilarity is likely due
to the decrease of fertilization rate after electroporation, and both elapsed time after
spawning and physical damage by electroporation can affect fertilization efficiency.
Therefore, we decided to conduct further electroporation experiments with fertilized
eggs.
347
348
Gene knockdown with siRNA in *Nematostella vectensis*

Before attempting gene knockdown via siRNAs electroporation in fertilized jellyfish

eggs, we first used *Nematostella vectensis,* where egg electroporation methodology has

been established[36]. We selected *Nematostella Brachyury* (*NvBra*), a gene expressed in

the blastopore margin that functions in the early embryo[49,50], as a target. After

electroporation of siRNAs targeting *NvBra* in fertilized eggs, we examined *NvBra*

expression in planula larvae at 3 days-post-fertilization (dpf) by RT-qPCR and found

that *NvBra* expression was drastically suppressed compared to no electroporation

controls and negative control siRNAs (Supplementary Fig. 3). We further evaluated

survival rates after electroporation and confirmed that siRNA electroporation does not

severely affect survival rates of 3 dpf planula at any of the tested conditions

(Supplementary Table 2). These results suggest that siRNA delivery into fertilized eggs

via electroporation is an effective gene knockdown method that is comparable to the

effect of shRNA delivery into unfertilized eggs via electroporation[36], supporting our

attempt to perform gene knockdown via siRNA electroporation using fertilized jellyfish

eggs.

Endogenous *GFP1* knockdown with siRNA in *Clytia hemisphaerica*

The hydrozoan jellyfish *Clytia* endogenously expresses four green fluorescence proteins

(*CheGFP1-4*), and the distribution of each GFP expression dynamically changes across

its life cycle. For instance, *Clytia* eggs express maternal *CheGFP2* mRNA, and planula

larvae show typical ring-like *CheGFP1* fluorescence in the lateral ectoderm at 3 dpf[11].

While *CheGFP1* gene expression becomes dominant in the planula stage; *CheGFP2,*

CheGFP3, and *CheGFP4* are expressed in different tissues and organs in the

medusa[51]. Momose *et al*. succeeded in genome editing by injecting CRISPR/Cas9

targeting the endogenous *CheGFP1* into fertilized eggs, and confirmed the knockout

effect by loss of green fluorescence in 3 dpf planula larvae[11]. In addition, *CheGFP1* is

not essential for animal survival, which allows for the monitoring of knockdown effects

after embryogenesis. For these reasons, we chose endogenous *CheGFP1* for the first

target gene of siRNA knockdown.

In order to maximize the effect of gene knockdown, we used a combination of

three different siRNAs against *CheGFP1* in a mixture with a 1:1:1 ratio, and

electroporated 600 ng/µl of siRNA mix (200 ng/µl per each siRNA) into fertilized eggs

(Fig. 3A). To verify the effect of gene knockdown, we measured intensity of green

fluorescence of 2 dpf planula and found that siRNA (600ng/µl siRNA mix) electroporated

embryos showed dramatically reduced fluorescent intensity compared to the no-siRNA

controls (Figs. 3B-D; mean intensity value: 5.7×10^3 for 0 ng/µl control, 1.8×10^3 for 600

ng/µl). To evaluate the knockdown effects at varying doses of siRNA, we next

electroporated different concentrations of the *CheGFP1* siRNA mix (0, 150, 300, and

600 ng/µl) into fertilized eggs. While control 3 dpf planulae with no-electroporation (No

EP) as well as those with no-siRNA showed typical ring-shape GFP fluorescence in the

ectoderm, many planulae with *CheGFP1* siRNA electroporation lost green fluorescence

（Figs. 3E-3H）. To evaluate the GFP knockdown effects, we classified planulae with

typical GFP1 fluorescence as "wild type" and planulae with reduced fluorescence as

"reduced" for further quantification (Figs. 3I and 3J). The percentage of wild type

phenotype dramatically reduced in an siRNA dose-dependent manner (Fig. 3K). To

further quantify the above result at a molecular level, we examined relative *CheGFP1*

mRNA levels by RT-qPCR. Consistent with the image analysis results, we confirmed a

marked reduction of *CheGFP1* gene expression at different concentrations of *CheGFP1*

siRNA (Fig. 3L). These results demonstrate that siRNAs targeting *CheGFP1* repress its

expression upon electroporation and that the knockdown effects are dose-dependent.

Wnt3* knockdown with siRNA in *Clytia hemisphaerica

Since electroporation of siRNAs targeting endogenous *CheGFP1* was shown to be

effective, we next decided to investigate the effects of knocking down other genes that

play important roles in early development. Wnt3 is a secreted signaling protein in the

Wnt/β-catenin pathway that is involved in diverse developmental processes including

cell fate decisions and patterning[52,53]. In *Clytia*, *Wnt3* RNA is maternally localized to the

animal cortex, the future oral side of the egg, and Wnt3 organizes axial patterning as

the main ligand for the Wnt/β-catenin pathway in the early embryonic stage[10]. After

embryogenesis, the *Clytia* planula normally elongates along the oral-aboral axis and

becomes oval in shape （Fig. 4A）. By contrast, after *CheWnt3* morpholino injection, the

morphant loses oral-aboral axis polarity and exhibits a spherical shape[10]. This raises the

possibility that knockdown of *CheWnt3* via siRNA may result in a similar spherical

morphology for *Clytia* larvae as observed in *CheWnt3* morphants. Therefore, to

evaluate siRNA knockdown effects on *Clytia*, we chose *CheWnt3* as our second target.

We used a mixture of two different siRNAs (1:1 ratio) targeting *CheWnt3*, and

carried out electroporation of the siRNA mix into *Clytia* fertilized eggs. After

electroporation of *CheWnt3* siRNAs, 1 dpf planula larvae showed nearly complete

spherical morphology (Fig. 4B), which is reminiscent of *CheWnt3* morphants[10]. To

quantify the effect of siRNA knockdown on *Clytia* larval morphology, we calculated the

aspect ratio in two dimensions, where values greater than 1.0 shows the tendency to be

oval (ellipsoid-like) while a value of 1.0 indicates a perfect circle (sphere) (Fig. 4C). The

control larvae with 0 ng/µl siRNA showed a median aspect ratio of 1.69, indicating a

typical ellipsoid-like morphology. By contrast, larvae that underwent electroporation with

CheWnt3 siRNAs (400 ng/µl) showed a median aspect ratio of 1.16, suggesting a much

greater tendency toward spherical morphology (Fig. 4D). These results quantitively

confirmed the effect of *CheWnt3* knockdown on larval morphology, which phenocopies

CheWnt3 morphants[10]. To analyze the level of gene-specific knockdown at varying

doses of siRNAs, we next electroporated mixtures of 0, 100, 200, and 400 ng/µl of

CheWnt3 siRNAs into *Clytia* fertilized eggs (Figs. 4E-4H). To evaluate *CheWnt3* siRNA

effects on morphology, we classified 1 dpf planulae with elongated ellipsoid-like shapes

as "normal" and planulae with spherical shapes as "spherical" (Figs. 4I and 4J). The

percentage of normal phenotype dramatically reduced after *CheWnt3* siRNA-

electroporation (< 1.0 % in 400 ng/µl), even at the lowest dose (5.2% in 100 ng/µl) (Fig.

4K). To further evaluate the result at a molecular level, we analyzed relative *CheWnt3*

mRNA levels by RT-qPCR and confirmed a significant reduction in *CheWnt3* gene

expression at all concentrations (Fig. 4L). Of note, after electroporation with universal

negative control siRNAs（Nippon gene co.Ltd）, both morphological phenotypes and

relative gene expression were comparable to no-siRNA controls (Supplementary Figs.

4A and 4B), further confirming that siRNA-mediated knockdown by electroporation is

target-specific. These results together indicate that siRNAs targeting endogenous genes

effectively knockdown their expression in *Clytia* embryos.

Identification of *Wnt3* in *Cladonema pacificum*

Identification of *Wnt3* in *Cladonema pacificum*

In contrast to *Clytia*, the sole genetic model jellyfish, previous studies have neither

established genetic manipulation nor performed genome sequencing of *Cladonema*

pacificum, despite its unique biological features. This is partly because the extremely

small size of the *Cladonema* egg makes microinjection difficult. By testing whether

siRNA-mediated knockdown via electroporation can work in *Cladonema*, we can start to

manipulate genes in *Cladonema* while also demonstrating that siRNA electroporation

can be applicable to other jellyfish species. While green fluorescent protein (GFP) is

often an easy early target for gene manipulation in emerging organisms, *Cladonema*

does not exhibit apparent endogenous green fluorescence expression at any stage,

unlike *Clytia*. We instead turned our focus to *Wnt3* as the target for gene knockdown in

Cladonema since Wnt3 plays an important role in early embryonic development in

several cnidarian species[54,55].

To identify the *Wnt3* ortholog in *Cladonema*, we utilized the result of RNA-seq of

the *Cladonema* polyp, stolon, and medusa manubrium (data not shown). We performed

Cladonema Wnt3 CDS annotation using *Clytia* CDS database (MARIMBA, Marine

models database: http://marimba.obs-vlfr.fr) and found one contig annotated with *Wnt3*

(*CpWnt3*). We then performed phylogenic analysis using the neighbor-joining method,

and confirmed *CpWnt3* (TRINITY_DN8032_c0_g1_i1.p1) is grouped with medusozoan

Clytia Wnt3 rather than Anthozoa *Nematostella* and *Acropora digitifera* (Fig. 5A).

Multiple sequence alignment further showed highly conserved amino acids sequences

among different species including bilaterians and cnidarians (Fig. 5B). These findings

raise the possibility that *CpWnt3* has a similar function in controlling the Wnt/β-catenin

pathway in *Cladonema.*

Wnt3* knockdown with siRNA in *Cladonema pacificum

Does *Wnt3* function during *Cladonema* embryogenesis, particularly during axial

patterning? Interestingly, *Cladonema* planula larvae do not show a clear elongated

shape (Fig. 6A), as observed in *Clytia*. It is thus possible that morphogenesis, including

axial patterning and/or the function of the Wnt/β-catenin pathway, differs between these

two jellyfish species.

In order to verify whether siRNA electroporation works in *Cladonema* and *Wnt3*
affects axial patterning, we performed electroporation of siRNAs targeting *CpWnt3*. We
prepared fertilized *Cladonema* eggs as we did in *Clytia*, and used a mixture of two
different siRNAs against *CpWnt3* CDS with a 1:1 ratio (300 ng/µl) and performed
electroporation with the previously established parameters (50V, 1 pulse, 25 ms) in
fertilized *Cladonema* eggs. Strikingly, planula larvae after electroporation of siRNAs for
CpWnt3 did not exhibit morphological differences compared to controls（Figs. 6A-6C;
median aspect ratio: 1.21 for 0 ng/µl control, 1.16 for 300 ng/µl for si*CpWnt3*）. We then
quantified the morphological phenotypes by calculating the aspect ratio of 1 dpf
planulae and confirmed that *CpWnt3* knockdown does not cause a significant
morphological change (Figs. 6D and 6E). One possibility that would explain the above
result is a defect in the siRNA knockdown itself. To test this potential explanation, we
carried out RT-qPCR using mRNA samples from 1 dpf planula after electroporating
different concentrations of *CpWnt3* siRNA mix (0, 150, and 300 ng/µl) into fertilized
eggs. We confirmed a reduction of *Wnt3* gene expression in an siRNA dose-dependent
manner (Fig. 6F), eliminating siRNA knockdown defects as the cause of our initial
result. Taken together this work indicates that repression of *Wnt3* expression does not
have a significant impact on *Cladonema* embryogenesis, at least on planula
morphology. Given that overall oral-aboral polarity is less prominent in *Cladonema*
compared to *Clytia*, the axial patterning mechanism may differ between these two
jellyfish species.
499
500

Discussion

In this study, we have established a method to knock down endogenous genes in two
hydrozoan jellyfish, *Clytia hemisphaerica* and *Cladonema pacificum*, via siRNA
electroporation of fertilized eggs. We showed that knockdown of endogenous *GFP1* in
Clytia causes the loss of GFP fluorescence in the planula stage, as previously achieved
by CRISPR/Cas9-mediated *GFP1* knockout[11]. We also confirmed that knockdown of
Wnt3 in *Clytia* induces spherical morphology of the planula, mirroring the results of
injections of *Wnt3* morpholino antisense oligo[10]. By contrast, although we succeeded in
efficient repression of *Wnt3* gene expression in *Cladonema* after *Wnt3* siRNA

electroporation, the morphological phenotypes did not appear as observed in *Clytia*,

suggesting that the mechanism involved in embryogenesis and planula morphogenesis

may differ across hydrozoan jellyfish species.

Effective gene knockdown via shRNA electroporation has previously been

reported using two cnidarian polyps, *Nematostella vectensis* and *Hydractinia*

symbiolongicarpus[36,38]. We tested whether shRNAs are also effective in the hydrozoan

jellyfish *Clytia*. We designed shRNAs targeting *CheWnt3* using the same 19 bp target

sequence that was used for siRNAs, and performed electroporation on fertilized eggs

with a mix of two different *Wnt3* shRNAs. After electroporation of *Wnt3* shRNAs, relative

Wnt3 mRNA expression in 1 dpf planula was weakly down-regulated (0.83), and was

not significant compared to the striking effects of *Wnt3* siRNA electroporation (0.039) at

the same concentration (400 ng/µl) (Supplementary Fig. 4A). In addition, electroporation

of *Wnt3* siRNAs resulted in severe morphological changes manifested by a spherical

shape, showing almost no "normal" phenotype (0.71%), whereas electroporation of

shRNA had a limited effect (60.4%) (Supplementary Fig. 4B). These results suggest

that the effect of siRNA is much stronger than that of shRNA in *Clytia* when treated with

the same target sequence and concentration.

One possibility that would explain such distinct effects between siRNA and

shRNA electroporation is the existence of differences in the RNAi machinery or

efficiency of RNA processing among cnidarians. During RNAi in mammals, the RNase

III Dicer protein processes shRNA in collaboration with cofactors TRBP (Transactivation

response element RNA-binding protein) and PACT（protein activator of the interferon-

induced protein kinase）to produce a mature form of siRNA[39]. Mature siRNA in the Dicer

and TRBP/PACT complex is associated with Argonaute protein and cleaves

endogenous complementary mRNAs as the RNA-induced silencing complex [30,56].

These small RNA biogenesis factors, Dicers and cofactors (TRBP in bilaterians, HYL1

in plants and cnidarians), are found in *Nematostella vectensis*, *Acropora digitifera*, and

Hydra vulgaris[41]. Although Dicers (TCONS_00004571 and TCONS_00004525) and

Hyl1 (TCONS_00010722) are predicted to exist in *Clytia* from the MARIMBA

transcriptome data, the expression level of Dicers in oocytes and early gastrulation

stages is lower than that in other stages, suggesting the possibility that insufficient

efficiency of shRNA to siRNA processing is responsible for the difference in knockdown

efficiency between siRNA and shRNA electroporation. It will be interesting to elucidate

the detailed molecular mechanism of RNAi in different cnidarian species and address

RNA processing efficiency in distinct developmental stages.

During *Clytia* early embryogenesis, *Wnt3* RNA is localized to the animal cortex of

the egg and contributes to the formation of the oral-aboral axis[10]. Accordingly,

knockdown of *Wnt3* in *Clytia* fertilized eggs induces suppression of axis formation and

planula elongation (Fig. 4). In contrast, no morphological phenotype was observed after

Wnt3 knockdown in *Cladonema*, despite a marked decrease in its gene expression (Fig.

6). This result was in stark contrast to the *Wnt3* loss-of-function phenotypes in *Clytia.*

We thought the divergence in outcome may be due to large differences in *Cladonema*

and *Clytia Wnt3*, but amino acid alignment results demonstrate that *Cladonema Wnt3* is

homologous to *Wnt3* in other cnidarians and is highly homologous to *Clytia Wnt3* (Fig.

5B). Beyond *Clytia*, the Wnt/β-catenin signaling pathway is involved in axis formation in

Nematostella[54,57]. Furthermore, given the similar localization of the Wnt ligand in

Hydractinia[55,58], the Wnt/β-catenin pathway likely functions in early *Cladonema*

development to some extent. One possibility that accounts for the morphological

phenotype of *Wnt3* knockdown in *Cladonema* is that several Wnt ligands act

redundantly. Indeed, we found multiple *Wnt* genes in *Cladonema* from transcriptome

data (Supplementary Fig. 5). Another possibility is that the limited elongation in

Cladonema planulae might hamper phenotypical appearance upon inhibition of Wnt/β-

catenin signaling. Understanding the diversity and differences of gastrulation and

planula morphogenesis in cnidarians is an interesting and unresolved topic for future

studies.

An interesting feature of jellyfish is that their morphology dramatically changes

across their life cycle from planula to polyp and then to medusa. In particular, polyps

and medusae, different adult stages of the same animal, exhibit distinct regenerative

ability, lifespan, and behaviors. Although regeneration mechanisms have been

extensively studied in polyp-only animals such as *Hydra* and *Hydractinia*, classical

studies have used the medusa stage of several jellyfish species and demonstrated their

regenerative potential[21]. Recent work using *Clytia* medusae has further shown a

remarkable remodeling and repatterning mechanism orchestrated by muscle systems

upon organ loss[17]. At the behavior level, the upside-down jellyfish *Cassiopea* medusae

exhibit a sleep-like state[59], which is similarly observed in *Hydra* polyps[60]. More recent

work using transgenic *Clytia* has characterized feeding behaviors in medusae at a

neural-network resolution[19]. To understand how these diverse biological processes in

jellyfish are regulated at the molecular level, it is necessary to analyze the functions of

specific genes at various stages. Although gene knockdown by dsRNA electroporation

into polyps in the scyphozoan jellyfish *Aurelia* had been reported[61], genetic

manipulation in medusae has not yet been achieved. Given that siRNA electroporation

is applicable to *Hydra* polyps[33,62] in addition to cnidarian fertilized eggs, we now have a

better chance to apply this technique to the medusa stage by adjusting established

conditions. Our method of gene knockdown via siRNA electroporation will complement

the method of shRNA electroporation in cnidarian polyps and promise molecular-level

analysis of interesting biological phenomena associated with different life stages in

jellyfish.

References

1 Technau, U. & Steele, R. E. Evolutionary crossroads in developmental biology:

Cnidaria. *Development* **138**, 1447-1458, doi:10.1242/dev.048959 (2011).

2 Zapata, F. *et al.* Phylogenomic Analyses Support Traditional Relationships within

Cnidaria. *PLoS One* **10**, e0139068, doi:10.1371/journal.pone.0139068 (2015).

3 Leclere, L. & Rottinger, E. Diversity of Cnidarian Muscles: Function, Anatomy,

Development and Regeneration. *Front Cell Dev Biol* **4**, 157,

doi:10.3389/fcell.2016.00157 (2016).

4 Holstein, T. W., Hobmayer, E. & Technau, U. Cnidarians: an evolutionarily

conserved model system for regeneration? *Dev Dyn* **226**, 257-267,

doi:10.1002/dvdy.10227 (2003).

5 Bosch, T. C. G. *et al.* Back to the Basics: Cnidarians Start to Fire. *Trends*

Neurosci **40**, 92-105, doi:10.1016/j.tins.2016.11.005 (2017).

6 Houliston, E., Momose, T. & Manuel, M. Clytia hemisphaerica: a jellyfish cousin

joins the laboratory. *Trends Genet* **26**, 159-167, doi:10.1016/j.tig.2010.01.008

(2010).

7   Houliston, E., Leclere, L., Munro, C., Copley, R. R. & Momose, T. Past, present
and future of Clytia hemisphaerica as a laboratory jellyfish. *Curr Top Dev Biol*,
doi:10.1016/bs.ctdb.2021.12.014 (2022).

8   Lechable, M. *et al.* An improved whole life cycle culture protocol for the
hydrozoan genetic model Clytia hemisphaerica. *Biol Open* **9**,
doi:10.1242/bio.051268 (2020).

9   Momose, T. & Houliston, E. Two oppositely localised frizzled RNAs as axis
determinants in a cnidarian embryo. *PLoS Biol* **5**, e70,
doi:10.1371/journal.pbio.0050070 (2007).

10 Momose, T., Derelle, R. & Houliston, E. A maternally localised Wnt ligand
required for axial patterning in the cnidarian Clytia hemisphaerica. *Development*
135, 2105-2113, doi:10.1242/dev.021543 (2008).

11 Momose, T. *et al.* High doses of CRISPR/Cas9 ribonucleoprotein efficiently
induce gene knockout with low mosaicism in the hydrozoan Clytia hemisphaerica
through microhomology-mediated deletion. *Sci Rep* **8**, 11734,
doi:10.1038/s41598-018-30188-0 (2018).

12 Quiroga Artigas, G. *et al.* A gonad-expressed opsin mediates light-induced
spawning in the jellyfish Clytia. *Elife* **7**, doi:10.7554/eLife.29555 (2018).

13 Quiroga Artigas, G. *et al.* A G protein-coupled receptor mediates neuropeptide-
induced oocyte maturation in the jellyfish Clytia. *PLoS Biol* **18**, e3000614,
doi:10.1371/journal.pbio.3000614 (2020).

14 Lapebie, P. *et al.* Differential responses to Wnt and PCP disruption predict
expression and developmental function of conserved and novel genes in a
cnidarian. *PLoS Genet* **10**, e1004590, doi:10.1371/journal.pgen.1004590 (2014).

15 Condamine, T. *et al.* Molecular characterisation of a cellular conveyor belt in
Clytia medusae. *Dev Biol* **456**, 212-225, doi:10.1016/j.ydbio.2019.09.001 (2019).

16 Leclere, L. *et al.* The genome of the jellyfish Clytia hemisphaerica and the
evolution of the cnidarian life-cycle. *Nat Ecol Evol* **3**, 801-810,
doi:10.1038/s41559-019-0833-2 (2019).

17 Sinigaglia, C. *et al.* Pattern regulation in a regenerating jellyfish. *Elife* **9**,
doi:10.7554/eLife.54868 (2020).

18 Chari, T. *et al.* Whole-animal multiplexed single-cell RNA-seq reveals
transcriptional shifts across Clytia medusa cell types. *Sci Adv* **7**, eabh1683,
doi:10.1126/sciadv.abh1683 (2021).

19 Weissbourd, B. *et al.* A genetically tractable jellyfish model for systems and
evolutionary neuroscience. *Cell* **184**, 5854-5868 e5820,
doi:10.1016/j.cell.2021.10.021 (2021).

20 Leclere, L. *et al.* Maternally localized germ plasm mRNAs and germ cell/stem cell
formation in the cnidarian Clytia. *Dev Biol* **364**, 236-248,
doi:10.1016/j.ydbio.2012.01.018 (2012).

21 Fujita, S., Kuranaga, E. & Nakajima, Y. I. Regeneration Potential of Jellyfish:
Cellular Mechanisms and Molecular Insights. *Genes (Basel)* **12**,
doi:10.3390/genes12050758 (2021).

22 Fujiki, A., Hou, S., Nakamoto, A. & Kumano, G. Branching pattern and
morphogenesis of medusa tentacles in the jellyfish Cladonema pacificum
(Hydrozoa, Cnidaria). *Zoological Lett* **5**, 12, doi:10.1186/s40851-019-0124-4
(2019).

23 Fujita, S., Kuranaga, E. & Nakajima, Y. I. Cell proliferation controls body size
growth, tentacle morphogenesis, and regeneration in hydrozoan jellyfish
Cladonema pacificum. *PeerJ* **7**, e7579, doi:10.7717/peerj.7579 (2019).

24 Hou, S., Zhu, J., Shibata, S., Nakamoto, A. & Kumano, G. Repetitive
accumulation of interstitial cells generates the branched structure of Cladonema
medusa tentacles. *Development* **148**, doi:10.1242/dev.199544 (2021).

25 Takeda, N. *et al.* Identification of jellyfish neuropeptides that act directly as
oocyte maturation-inducing hormones. *Development* **145**,
doi:10.1242/dev.156786 (2018).

26 Stierwald, M., Yanze, N., Bamert, R. P., Kammermeier, L. & Schmid, V. The Sine
oculis/Six class family of homeobox genes in jellyfish with and without eyes:
development and eye regeneration. *Dev Biol* **274**, 70-81,
doi:10.1016/j.ydbio.2004.06.018 (2004).

27 Suga, H., Schmid, V. & Gehring, W. J. Evolution and functional diversity of
jellyfish opsins. *Curr Biol* **18**, 51-55, doi:10.1016/j.cub.2007.11.059 (2008).

28   Suga, H. *et al.* Flexibly deployed Pax genes in eye development at the early
668   evolution of animals demonstrated by studies on a hydrozoan jellyfish. *Proc Natl*
669   *Acad Sci U S A* **107**, 14263-14268, doi:10.1073/pnas.1008389107 (2010).
670   29   Graziussi, D. F., Suga, H., Schmid, V. & Gehring, W. J. The "eyes absent" (eya)
671   gene in the eye-bearing hydrozoan jellyfish Cladonema radiatum: conservation of
672   the retinal determination network. *J Exp Zool B Mol Dev Evol* **318**, 257-267,
673   doi:10.1002/jez.b.22442 (2012).
674   30   Meister, G. & Tuschl, T. Mechanisms of gene silencing by double-stranded RNA.
675   *Nature* **431**, 343-349, doi:10.1038/nature02873 (2004).
676   31   Rao, D. D., Vorhies, J. S., Senzer, N. & Nemunaitis, J. siRNA vs. shRNA:
677   similarities and differences. *Adv Drug Deliv Rev* **61**, 746-759,
678   doi:10.1016/j.addr.2009.04.004 (2009).
679   32   Moran, Y. *et al.* Cnidarian microRNAs frequently regulate targets by cleavage.
680   *Genome Res* **24**, 651-663, doi:10.1101/gr.162503.113 (2014).
681   33   Watanabe, H. *et al.* Nodal signalling determines biradial asymmetry in Hydra.
682   *Nature* **515**, 112-115, doi:10.1038/nature13666 (2014).
683   34   Pankow, S. & Bamberger, C. The p53 tumor suppressor-like protein nvp63
684   mediates selective germ cell death in the sea anemone Nematostella vectensis.
685   *PLoS One* **2**, e782, doi:10.1371/journal.pone.0000782 (2007).
686   35   He, S. *et al.* An axial Hox code controls tissue segmentation and body patterning
687   in Nematostella vectensis. *Science* **361**, 1377-1380,
688   doi:10.1126/science.aar8384 (2018).
689   36   Karabulut, A., He, S., Chen, C. Y., McKinney, S. A. & Gibson, M. C.
690   Electroporation of short hairpin RNAs for rapid and efficient gene knockdown in
691   the starlet sea anemone, Nematostella vectensis. *Dev Biol* **448**, 7-15,
692   doi:10.1016/j.ydbio.2019.01.005 (2019).
693   37   DuBuc, T. Q. *et al.* Transcription factor AP2 controls cnidarian germ cell
694   induction. *Science* **367**, 757-762, doi:10.1126/science.aay6782 (2020).
695   38   Quiroga-Artigas, G., Duscher, A., Lundquist, K., Waletich, J. & Schnitzler, C. E.
696   Gene knockdown via electroporation of short hairpin RNAs in embryos of the
697   marine hydroid Hydractinia symbiolongicarpus. *Sci Rep* **10**, 12806,
698   doi:10.1038/s41598-020-69489-8 (2020).

39 Ha, M. & Kim, V. N. Regulation of microRNA biogenesis. *Nat Rev Mol Cell Biol*
15, 509-524, doi:10.1038/nrm3838 (2014).
40 Krishna, S. *et al.* Deep sequencing reveals unique small RNA repertoire that is
regulated during head regeneration in Hydra magnipapillata. *Nucleic Acids Res*
41, 599-616, doi:10.1093/nar/gks1020 (2013).
41 Moran, Y., Praher, D., Fredman, D. & Technau, U. The evolution of microRNA
pathway protein components in Cnidaria. *Mol Biol Evol* **30**, 2541-2552,
doi:10.1093/molbev/mst159 (2013).
42 Murchison, E. P., Partridge, J. F., Tam, O. H., Cheloufi, S. & Hannon, G. J.
Characterization of Dicer-deficient murine embryonic stem cells. *Proc Natl Acad*
Sci U S A **102**, 12135-12140, doi:10.1073/pnas.0505479102 (2005).
43 Kok, K. H., Ng, M. H., Ching, Y. P. & Jin, D. Y. Human TRBP and PACT directly
interact with each other and associate with dicer to facilitate the production of
small interfering RNA. *J Biol Chem* **282**, 17649-17657,
doi:10.1074/jbc.M611768200 (2007).
44 Watanabe, H. *et al.* Sequential actions of beta-catenin and Bmp pattern the oral
nerve net in Nematostella vectensis. *Nat Commun* **5**, 5536,
doi:10.1038/ncomms6536 (2014).
45 Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis.
Nat Methods **9**, 676-682, doi:10.1038/nmeth.2019 (2012).
46 Thompson, J. D., Higgins, D. G. & Gibson, T. J. CLUSTAL W: improving the
sensitivity of progressive multiple sequence alignment through sequence
weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids*
Res **22**, 4673-4680, doi:10.1093/nar/22.22.4673 (1994).
47 Waterhouse, A. M., Procter, J. B., Martin, D. M., Clamp, M. & Barton, G. J.
Jalview Version 2--a multiple sequence alignment editor and analysis workbench.
Bioinformatics **25**, 1189-1191, doi:10.1093/bioinformatics/btp033 (2009).
48 Deguchi, R., Kondoh, E. & Itoh, J. Spatiotemporal characteristics and
mechanisms of intracellular Ca(2+) increases at fertilization in eggs of jellyfish
(Phylum Cnidaria, Class Hydrozoa). *Dev Biol* **279**, 291-307,
doi:10.1016/j.ydbio.2004.11.036 (2005).

49 Scholz, C. B. & Technau, U. The ancestral role of Brachyury: expression of
NemBra1 in the basal cnidarian Nematostella vectensis (Anthozoa). *Dev Genes*
Evol **212**, 563-570, doi:10.1007/s00427-002-0272-x (2003).

50 Servetnick, M. D. *et al.* Cas9-mediated excision of Nematostella brachyury
disrupts endoderm development, pharynx formation and oral-aboral patterning.
Development **144**, 2951-2960, doi:10.1242/dev.145839 (2017).

51 Fourrage, C., Swann, K., Gonzalez Garcia, J. R., Campbell, A. K. & Houliston, E.
An endogenous green fluorescent protein-photoprotein pair in Clytia
hemisphaerica eggs shows co-targeting to mitochondria and efficient
bioluminescence energy transfer. *Open Biol* **4**, 130206, doi:10.1098/rsob.130206
(2014).

52 Guder, C. *et al.* The Wnt code: cnidarians signal the way. *Oncogene* **25**, 7450-
7460, doi:10.1038/sj.onc.1210052 (2006).

53 Lee, P. N., Pang, K., Matus, D. Q. & Martindale, M. Q. A WNT of things to come:
evolution of Wnt signaling and polarity in cnidarians. *Semin Cell Dev Biol* **17**,
157-167, doi:10.1016/j.semcdb.2006.05.002 (2006).

54 Kraus, Y., Aman, A., Technau, U. & Genikhovich, G. Pre-bilaterian origin of the
blastoporal axial organizer. *Nat Commun* **7**, 11694, doi:10.1038/ncomms11694
(2016).

55 Plickert, G., Jacoby, V., Frank, U., Muller, W. A. & Mokady, O. Wnt signaling in
hydroid development: formation of the primary body axis in embryogenesis and
its subsequent patterning. *Dev Biol* **298**, 368-378,
doi:10.1016/j.ydbio.2006.06.043 (2006).

56 Bartel, D. P. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*
116, 281-297, doi:10.1016/s0092-8674(04)00045-5 (2004).

57 DuBuc, T. Q., Stephenson, T. B., Rock, A. Q. & Martindale, M. Q. Hox and Wnt
pattern the primary body axis of an anthozoan cnidarian before gastrulation. *Nat*
Commun **9**, 2007, doi:10.1038/s41467-018-04184-x (2018).

58 Duffy, D. J., Plickert, G., Kuenzel, T., Tilmann, W. & Frank, U. Wnt signaling
promotes oral but suppresses aboral structures in Hydractinia metamorphosis
and regeneration. *Development* **137**, 3057-3066, doi:10.1242/dev.046631
(2010).

59 Nath, R. D. *et al.* The Jellyfish Cassiopea Exhibits a Sleep-like State. *Curr Biol*
27, 2984-2990 e2983, doi:10.1016/j.cub.2017.08.014 (2017).
60 Kanaya, H. J. *et al.* A sleep-like state in Hydra unravels conserved sleep
mechanisms during the evolutionary development of the central nervous system.
Sci Adv **6**, doi:10.1126/sciadv.abb9415 (2020).
61 Fuchs, B. *et al.* Regulation of polyp-to-jellyfish transition in Aurelia aurita. *Curr*
Biol **24**, 263-273, doi:10.1016/j.cub.2013.12.003 (2014).
62 Lommel, M. *et al.* Hydra Mesoglea Proteome Identifies Thrombospondin as a
Conserved Component Active in Head Organizer Restriction. *Sci Rep* **8**, 11753,
doi:10.1038/s41598-018-30035-2 (2018).

772
773

Acknowledgement

We thank R. Deguchi (Miyagi Univ. Education, Japan) for sharing *Cladonema pacificum*
and EMBRC France for sharing *Clytia hemisphaerica*. We thank T. Momose (Sorbonne
University, CNRS, France) for helpful discussion. We thank I. Nagai and A. Sasaki for
technical assistance; and A. Dahal, A. Tanimoto, and J. Higuchi for *Nematostella*
culture. This work was supported by JST grant number JPMJCR1852 to E.K., AMED
under grant number JP21gm6110025 to Y.N., and the JSPS KAKENHI grant numbers
21H05255 to E.K., and JP17H06332 and JP19K22550 to Y.N.

782
783

Author contributions

T.M.-O. conceptualized and designed the project, performed experiments using jellyfish,
analyzed data, prepared figures, and wrote the manuscript. S.F. performed experiments
and analyzed data. R.N. performed experiments using *Nematostella*. H.W.
conceptualized and designed the project. E.K. contributed to reagents. Y.N.
conceptualized and designed the project, prepared figures, and wrote the manuscript.
All authors approved the final manuscript.

791
792

Competing interests

The authors declare no competing interests.

Figure legends

Figure 1. The hydrozoan jellyfish *Clytia hemisphaerica* and *Cladonema pacificum*.

(**A**) The cladogram depicting the phylogenetic position of cnidarian jellyfish. As the sister

group of Bilateria, the phylum Cnidaria is divided into two clades, Anthozoa and

Medusozoa, which consists of four classes: Hydrozoa, Staurozoa, Scyphozoa, and

Cubozoa. Hydorozoa includes polyp type animals without a medusa stage (e.g. *Hydra*

and *Hydractinia*) and jellyfish that have both polyp and medusa stages (e.g. *Clytia* and

Cladonema). (**B** and **C**) Photos of an adult medusa and eggs of *Clytia hemisphaerica*

(**B**) and *Cladonema pacificum* (**C**). Scale bars: 1 mm for medusae; 100 µm for eggs.

Figure 2. The electroporation procedure for jellyfish eggs. (**A**) Schematic of

spawning under the 24 h light/dark cycle. *Clytia hemisphaerica* are maintained in a 13 h

dark/11 h light cycle. After light stimuli, sperm spawning occurs within 60-90 min and

egg spawning occurs within 90-120 min. *Cladonema pacificum* are maintained in a 23.5

h light/0.5 h dark cycle. After dark stimuli, sperm and egg spawning occurs within 25

min. (**B**) Fertilized eggs are resuspended in 15% Ficoll/artificial sea water to prevent

precipitation of eggs during electroporation. (**C**) Electroporation is performed with the

Bio-Rad Gene Pulser Xcell electroporation system and a cuvette with a 4 mm gap. (**D**)

Visualization of the Rhodamine-Dextran delivery into eggs by electroporation.

Rhodamine is shown in red. While little or no fluorescence was observed in eggs

without electroporation (No EP), under the condition of a single 50 V pulse for 25 ms,

rhodamine signals were detected in both *Clytia and Cladonema* eggs without cell

damage. For clarity, egg outlines are indicated with dashed lines for eggs without

electroporation. Scale bars: 200 µm.

Figure 3. **Phenotypes of *GFP1* knockdown with siRNA in *Clytia* fertilized eggs.** (**A**) Schematic of the siRNA electroporation procedure using fertilized eggs. Fertilized eggs in 15% Ficoll/artificial sea water mixed with siRNA are incubated for 5 min, and electroporation is conducted after transferring into a cuvette. (**B** and **C**) Phenotypes of *Clytia* 2-day planula after *CheGFP1* siRNA knockdown. (**D**) Boxplots showing the GFP fluorescence intensity mean values in 2-day planula after electroporation of 0 and 600 ng/µl siRNAs targeting *CheGFP1*. Center lines indicate the medians; x's denote the means; box limits represent the 25th and 75th percentiles; whiskers show the maximum and minimum values. (**E**-**H**) Phenotypes of 3-day planula after siRNA targeting *CheGFP1* electroporation (0, 150, 300, and 600 ng/µl si*CheGFP1*). (**I** and **J**) For quantification of green fluorescence, we classified phenotypes into two categories: planula with ring-like green fluorescence as "wild-type," and planula with reduced green fluorescence as "reduced". (**K**) Quantification of planula larvae positive for *CheGFP1* (wild-type). Bar plots show the percentage of wild-type phenotype after *CheGFP1* knockdown. Number of examined planula: 0 ng/µl, *n*=82; 150 ng/µl, *n*=64; 300 ng/µl, *n*=58; 600 ng/µl, *n*=114. Error bars: maximum and minimum values. (**L**) Quantification of *CheGFP1* mRNA expression levels in 3-day planula by RT-qPCR. *CheEF1a* was used as an internal control. Bar heights represent mean values of at least three independent experiments. *CheGFP1* expression levels are standardized relative to the control (0 ng/µl) condition. Error bars: standard deviation. Scale bars: 200 µm.

Figure 4. Phenotypes of *Wnt3* knockdown with siRNA in *Clytia* fertilized eggs. (**A** and **B**) Typical phenotypes of *Clytia* 1-day planula. While morphologies of control planula (0 ng/µl) have an elongated oval-shape (A), those of *CheWnt3* siRNA (400 ng/µl) are spherical in shape (**B**). (**C**) Schematic of ellipse approximation for planula morphology and calculation of the aspect ratio. The aspect ratio was calculated by dividing the long axis (a) by short axis (b). (**D**) Boxplots showing the aspect ratio of the 1-day planula. Center lines show the medians; x's denote the mean values; box limits indicate the 25th and 75th percentiles; whiskers show maximum and minimum values; inner points and outliers are represented by circles. Number of examined planula: 0 ng/µl, *n*=127; 400 ng/µl, *n*=25. (**E**-**H**) Phenotypes of 1-day planula after siRNA targeting

CheWnt3 electroporation (0, 100, 200, and 400 ng/µl si*CheWnt3*). (**I** and **J**) For

morphology quantification, we classified planula phenotypes into two categories:

elongated oval shape as "normal" and spherical (circle) shape as "spherical". (**K**) Bar

plots show the percentage of normal phenotypes across four different siRNA doses.

Percentages are the mean value of three independent experiments, and error bars

indicate standard deviation. Numbers of examined planula: 0 ng/µl, n=127; 100 ng/µl,

n=75; 200 ng/µl, n=86; 400 ng/µl, n=95. Error bars: maximum and minimum values. (**L**)

Quantification of *CheWnt3* mRNA levels of 1-day planula by RT-qPCR. *CheEF1a* was

used as an internal control. Bar heights represent the mean value of at least three

independent experiments. *CheWnt3* expression levels are standardized relative to the

control (0 ng/µl). Error bars: standard deviation. Scale bars: (A, B, E-H) 400 µm, (I, J)

100 µm.

Figure 5. Identification of *Cladonema Wnt3* by phylogenetic analyses. (**A**) The

results of (**A**) phylogenetic analyses and (**B**) sequence alignment of Wnt3 among

indicated species. Amino acid sequence alignment was performed with ClustalW.

Alignment visualization and phylogenetic tree calculation were done with the Jalview

software. Cnidarian Wnt3 proteins are highlighted in blue and *Cladonema* Wnt3 is

highlighted in red. In sequence alignment, consensus residues are colored. Wnt3

proteins are universally conserved among animals. *He, Hydractinia echinate; Cp,*

Cladonema pacificum; Che, Clytia hemisphaerica; Nv, Nematostella vectensis; Ad,

Acropora digitifera; Hs, Homo sapiens; Gg, Gallus gallus; Xt, Xenopus tropicalis.

GenBank accession numbers are shown in "Materials and methods."

Figure 6. Phenotypes of *Wnt3* knockdown with siRNA and Wnt/β-catenin pathway

inhibitor treatment in *Cladonema* fertilized eggs. (**A**) Typical morphology of

Cladonema egg and planula. The wild-type *Cladonema* planula larvae exhibit a short

oval shape. To deliver siRNAs targeting *CpWnt3,* electroporation was performed in

fertilized eggs (1-cell stage), and phenotypes were confirmed at planula in one day. (**B**

and **C**) Phenotypes of 1-day planula after *CpWnt3* siRNA knockdown. (**D**) Schematic of

ellipse approximation for planula morphology and calculation of the aspect ratio. The

aspect ratio was calculated by dividing the long axis (a) by short axis (b). (**E**) Boxplots

showing the aspect ratio of the 1-day planula after si*CpWnt3* electroporation (0 and 300

ng/µl). Center lines show the medians; x's denote the mean values; box limits indicate

the 25th and 75th percentiles; whiskers show maximum and minimum values; inner

points and outliers are represented by circles. Number of examined planulae: 0 ng/µl,

n=22; 300 ng/µl, *n*=50. (**F**) Quantification of *CpWnt3* mRNA levels in 1-day planula by

RT-qPCR. *CpEF1a* was used as an internal control. Bar heights represent the mean

value of at least three independent experiments. Error bars indicate standard deviation.

CpWnt3 expression levels are standardized relative to the control (0 ng/µl). Scale bars:

(B, C) 200µm.

Supplementary materials

Supplementary Figure 1. Dextran delivery to *Clytia* eggs under different

electroporation conditions. (**A**-**G**) Photos of unfertilized *Clytia* eggs after

electroporation of Dextran-Rhodamine (1 mg/ml) in indicated conditions. Fluorescence

intensity tends to increase in a voltage and time dependent manner. In the severe

conditions above 200 V (**E**-**G**), eggs were severely damaged. Scale bars: 200 µm.

Supplementary Figure 2. Dextran delivery to *Cladonema* eggs under different

electroporation conditions. (**A**-**F**) Photos of unfertilized *Cladonema* eggs after

electroporation of Dextran-Rhodamine (1 mg/ml) in indicated conditions. Fluorescence

was frequently detected beyond 50 V, but in the severe conditions (**E** and **F**), eggs were

severely damaged. Scale bars: 200 µm.

Supplementary Figure 3. Gene knockdown with siRNA via electroporation into

***Nematostella* fertilized eggs.** qPCR quantification of *Nematostella Brachyury* (*NvBra*)

mRNA expression in siRNA-electroporated planulae (3 dpf). *NvEf1alpha* was used as

the internal control. Shown are the results of the mean ± S.D. from a single experiment

in triplicate, representing at least two independent experiments with similar results.

Supplementary Figure 4. Comparison of siRNA and shRNA in *Clytia*. (**A**) *CheWnt3*

relative gene expression using total mRNA samples of 1-day planula after

electroporation. *CheEF1a* was used as an internal control. Bar heights represent the

mean value of at least three independent experiments. Error bars indicate standard

deviation. siNC, siRNA universal negative control. (**B**) Percentage of "normal

phenotype" of 1-day planula after *CheWnt3* knockdown. Number of examined planula:

shRNA 400 ng/μl, n=288; siNC 400 ng/μl, n=45. Data are the average of three

biologically different experiments. Note that the same dataset for 0 and 400 ng/μl from

Fig. 4K and 4L were shown for comparison.

Supplementary Figure 5. Phylogenetic analysis of *Cladonema* Wnt proteins.   To

analyze orthology of *Cladonema* Wnt proteins, phylogenetic analysis was performed by

using amino acids sequences of Wnt proteins from *Cladonema*, *Clytia* and

Nematostella. Maximum likelihood (ML) tree calculation was inferred on ClustalW with

the PhyML bootstrap method. Branch supports were computed out of 100 bootstrapped

trees. Tree was visualized using MEGA-X software (Kumar S, Stecher G, Li M, Knyaz

C, and Tamura K (2018) *Molecular Biology and Evolution* 35:1547-1549). We performed

Cladonema Wnt ligand CDS annotation using *Clytia* CDS database (MARIMBA, Marine

models database: http://marimba.obs-vlfr.fr) and found eight contigs annotated with Wnt

ligands in *Cladonema*. Among the eight Wnt contigs, CpWntX4

(TRINITY_DN27620_c1_g1_i1.p1) is homologous to the *Clytia* Wnt ligand CDS,

CheWntX4 (TCONS_00027333-protein), which has never been registered in GenBank.

We used protein BLAST, but no other homologous cnidarian Wnt was found, so that we

provide the name CpWntX4 for convenience.

Cp, Cladonema pacificum; Che, Clytia hemisphaerica; Nv, Nematostella vectensis.

GenBank accession numbers, transcriptome name or RNAseq contig ID are follows;

CpWnt5a (TRINITY_DN3183_c0_g3_i1.p1), CpWntX1A

(TRINITY_DN4292_c0_g1_i3.p1), CpWnt2 (TRINITY_DN8519_c0_g1_i1.p1), CpWnt9

(TRINITY_DN3403_c0_g1_i3.p1), CpWntX4 (TRINITY_DN27620_c1_g1_i1.p1),

CpWntX2 (TRINITY_DN13473_c0_g1_i3.p1), CpWnt7b

(TRINITY_DN9745_c0_g1_i1.p1), CheWnt2 (TCONS_00012270), CheWntX1A

(ACB15464.1), CheWnt3 (ACB15465.1), CheWntX4 (TCONS_00027333-protein),

CheWntX1B (TCONS_00065525), Wnt5 (ACB15462.1), CheWnt5a

(TCONS_00047033), CheWntX3 (ACB28461.1), CheWntX2 (ACB15461.1), CheWnt7b

(TCONS_00073720), CheWnt9 (TCONS_00056741), NvWnt1 (AAT00640.1), NvWnt3

(ABF48092.1), NvWnt5 (AAW28133.1), NvWnt6 (AAW28134.1)

Supplementary Table 1. **Survival rate of *Cladonema* embryos after**

electroporation. The number of cleavage stage *Cladonema* embryos at 1 h post

fertilization (hpf) was considered embryo survival. Compared to fertilized eggs,

unfertilized eggs were much less tolerant of electroporation.

Supplementary Table 2. **Survival rate of *Nematostella* embryos after**

electroporation. The number of live *Nematostella* larvae was counted at 3 dpf (planula

stage). The survival rate remained around 20-40% regardless of the difference in siRNA

concentration and target.

Figure 1

Masuda-Ozawa et al.

A

Bilateria

Cnidaria

Anthozoa

Medusozoa

Hydrozoa
Polyp: *Hydra, Hydractinia*
Jellyfish: *Clytia, Cladonema*

Staurozoa

Scyphozoa

Cubozoa

B Medusa   Egg

Clytia hemisphaerica

C

Cladonema pacificum

Figure 2
Masuda-Ozawa et al.

A *Clytia hemisphaerica*

Spawn

0   Dark   13   14.5   Light   24 (hour)

Cladonema pacificum

Spawn

0   Light   13   13.5   Light   24 (hour)
      Dark

B

C

D

	No EP	50 V, 1 pulse, 25 ms
Clytia	Rhodamine	
Cladonema		

Figure 3

Masuda-Ozawa et al.

A Fertilized eggs + siRNA 5 min Electroporation

si*CheGFP1* electroporation
2 day planula

B 0 ng/µl

C 600 ng/µl

D GFP intensity mean value (A.U. x10³)

si*CheGFP1* electroporation/3 day planula

E 0 (ng/µl)

F 150

G 300

H 600

I Wild type — No EP

J Reduced — si*GFP1*

K Wild type (%)

L *GFP1* relative gene expression

Figure 4

Masuda-Ozawa et al.

si*CheWnt3* electroporation
1 day planula

A 0 ng/µl

B 400 ng/µl

C

Oral

Ellipse

Planula

Aboral

Aspect ratio

$A = \dfrac{a}{b}$

D

Aspect ratio

0   400 (ng/µl)

1.8

1.2

1.0

si*CheWnt3* electroporation/1 day planula

E 0 (ng/µl)

F 100

G 200

H 400

I No EP

Normal

J si*Wnt3*

Spherical

K

Normal phenotype (%)

0   100   200   400 (ng/µl)

L

Wnt3 relative gene expression

0   100   200   400 (ng/µl)

Figure 6

A Egg — si*CpWnt3* electroporation → 1 day planula

si*CpWnt3* electroporation/1 day planula

B 0 ng/µl

C 300 ng/µl

D

Planula   Ellipse

Aspect ratio

$$A = \frac{a}{b}$$

E

Aspect ratio

0   300   (ng/µl)

1.8

1.2

1.0

F

CpWnt3 relative gene expression

0   150   300 (ng/µl)