1

**siRNA-mediated gene knockdown via electroporation in hydrozoan**

**jellyfish embryos**

[1] Tokiha Masuda-Ozawa, [2] Sosuke Fujita**,** [3] Ryotaro Nakamura, [3] Hiroshi Watanabe,

[2] Erina Kuranaga, and [1,4] Yu-ichiro Nakajima**\***

1. Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, Sendai,

Japan

2. Graduate School of Life Sciences, Tohoku University, Sendai, Japan

3. Evolutionary Neurobiology Unit, Okinawa Institute of Science and Technology

Graduate University, Okinawa, Japan

4. Graduate School of Pharmaceutical Sciences, The University of Tokyo

*Author for correspondence e-mail: nakaji97@g.ecc.u-tokyo.ac.jp

Tel: +81-3-5841-4863

**Running title**: **siRNA gene knockdown in jellyfish embryos**

**Main text**: 6487 words (Abstract: 258 words)

**Figures**: 5 **Supplementary information** (Figures: 7; Table: 5)

**Keywords**: siRNA, electroporation, cnidarian, hydrozoan, jellyfish, *Clytia, Cladonema*

## Abstract

As the sister group to bilaterians, cnidarians stand in a unique phylogenetic position that provides insight into evolutionary aspects of animal development, physiology, and behavior. While cnidarians are classified into two types, sessile polyps and free-swimming medusae, most studies at the cellular and molecular levels have been conducted on representative polyp-type cnidarians and have focused on establishing techniques of genetic manipulation. Recently, gene knockdown by delivery of short hairpin RNAs into eggs via electroporation has been introduced in two polyp-type cnidarians, *Nematostella vectensis* and *Hydractinia symbiolongicarpus*, enabling systematic loss-of-function experiments. By contrast, current methods of genetic manipulation for most medusa-type cnidarians, or jellyfish, are quite limited, except for *Clytia hemisphaerica*, and reliable techniques are required to interrogate function of specific genes in different jellyfish species. Here, we present a method to knock down target genes by delivering small interfering RNA (siRNA) into fertilized eggs via electroporation, using the hydrozoan jellyfish, *Clytia hemisphaerica* and *Cladonema paciificum*. We show that siRNAs targeting endogenous *GFP1* and *Wnt3* in *Clytia* efficiently knock down gene expression and result in known planula phenotypes: loss of green fluorescence and defects in axial patterning, respectively. We also successfully knock down endogenous *Wnt3* in *Cladonema* by siRNA electroporation, which circumvents the technical difficulty of microinjecting small eggs. *Wnt3* knockdown in *Cladonema* causes gene expression changes in axial markers, suggesting a conserved Wnt/β-catenin-mediated pathway that controls axial polarity during embryogenesis. Our gene-targeting siRNA electroporation method is applicable to other animals, including and beyond jellyfish species, and will facilitate the investigation and understanding of myriad aspects of animal development.

## Introduction

The phylum Cnidaria is the sister group to Bilateria, having separated from their common ancestor over 500 million years ago. Cnidarians have diversified their morphologies across an array forms, including corals, sea anemones, hydroids, and jellyfish, all of which are divided into two clades: Anthozoa and Medusozoa. The two differ in that Anthozoa includes only sessile polyp-type animals, while Medusozoa (Hydrozoa, Staurozoa, Scyphozoa, and Cubozoa) contains two forms: polyp and medusa, commonly known as jellyfish (Fig. 1A)[1,2]. The life cycle of most, though not all, jellyfish consists of five forms: gametes, fertilized eggs, planulae, polyps, and medusae[3]. Fertilized eggs undergo embryogenesis to become planula larvae, which metamorphose into sessile polyps. While vegetatively-growing polyps give rise to free-swimming medusae through the process of budding or strobilization, medusae sexually reproduce by releasing gametes. Due to their unique phylogenetic position, studies using cnidarians have provided evolutionary insight into development, regeneration, and behaviors in multicellular animals[1,4,5]. Despite divergent morphologies and lifestyles among cnidarians, to date, the molecular and cellular understanding of cnidarians has been acquired primarily using polyp-type animals such as the anthozoan *Nematostella vectensis* and the hydrozoan *Hydra* and *Hydractinia*. By contrast, jellyfish biology remains largely unestablished.

Among jellyfish species, the hydrozoan *Clytia hemisphaerica* is the best-studied laboratory model[6,7]. The well-established methodology for maintaining and propagating animals ensures a reliable source for daily experiments[8]. Transparent and small-sized medusae make whole-mount visualization achievable, and relatively large-sized eggs (~180 µm diameter) enable different manipulations via microinjection (Fig. 1B). Indeed, gene silencing by morpholino oligos and mRNA microinjection as well as gene knockout via CRISPR/Cas9 allow for the investigation of functions of genes of interest[9-13]. Furthermore, the *Clytia* genome assembly is complete, and transcriptome profiles are available for various stages and tissues, even at the single-cell level, which creates the foundation for research at the molecular level[12,14-18]. These resources and techniques, together with recently-established transgenesis[19], have accelerated research into numerous facets of distinct *Clytia* life stages. While initial studies focused on

embryogenesis, current research topics using *Clytia* include gametogenesis,

regeneration, and behavior exhibited by adult medusa [9,10,12,17,19,20].

Compared to the established model jellyfish *Clytia*, the hydrozoan *Cladonema*

*pacificum* is an emerging model that has recently been utilized for various studies in

development, regeneration, and physiology[21]. The ease of rearing all stages of

*Cladonema* without a filtration system or large water tank, along with their high

spawning rate, enable easy lab maintenance. Small-sized medusae with branched

tentacles allow for investigations of body size control and tentacle morphogenesis (Fig.

1C)[22-24]. *Cladonema* gametogenesis is regulated by a light-dark cycle as in *Clytia*, and

recent work has identified the neuropeptides involved in oocyte maturation[25].

Furthermore, *Cladonema* medusae possess eyes with a complex structure that includes

lenses (ocelli) in their tentacle bulbs, and studies using the closely related species

*Cladonema radiatum* have identified conserved light-sensitive opsins and regulators of

eye formation (*Pax*, *Six*, and *Eya*)[25-29], providing a model for the evolutionary

developmental biology of photoreceptor organs. Despite these attractive features of

*Cladonema* as a laboratory animal, no genome assembly or transcriptome is currently

available, and genetic manipulation techniques are only just being developed. One

major technical issue in manipulating *Cladonema* is their small eggs—with a diameter of

approximately 60 μm (Fig. 1C), regular microinjection is quite difficult. Establishing

genetic manipulations is required to facilitate the in-depth investigations needed to

further understand the biology of *Cladonema*, and an alternative method that eliminates

the need for microinjection would greatly facilitate that objective.

RNA interference (RNAi), the phenomenon of double stranded RNA (dsRNA)

mediated silencing of target genes, has been widely exploited in living organisms to

analyze gene function[30], primarily using chemically- or *in vitro*-synthesized double-

stranded small interfering RNAs (siRNAs) or vector-based short hairpin RNAs

(shRNAs)[31]. In cnidarians, siRNA-mediated gene silencing was initially applied to

*Nematostella* and *Hydra* polyps via soaking or electroporation[32-34]. More recently, gene

knockdown via shRNA microinjection or electroporation into eggs has been utilized in

studies with *Nematostella* and *Hydractinia* to show efficient reduction in gene

expression and associated phenotypes in early developmental stages[35-38]. In particular,

shRNA delivery via electroporation, which does not require the rigors of microinjection,

allows for the experimental gene knockdown of large numbers of individuals

simultaneously[36,38], opening up the possibility of manipulating genes in different aquatic

animals, even those that produce very small eggs like *Cladonema*.

The mechanism of RNAi-mediated gene repression after introducing foreign

dsRNA into cells involves the microRNA (miRNA) pathway, an endogenous gene

repression machinery conserved in both animals and plants[39]. In cnidarians, the

presence of the miRNA pathway has been demonstrated in *Nematostella* and

*Hydra*[32,40,41]. However, little is known about the endogenous RNAi pathway, especially

the presence and roles of miRNAs and the miRNA-related genes, in jellyfish such as

*Clytia* and *Cladonema*. Furthermore, in mammalian cultured cells, siRNAs can induce

gene repression independent of endogenous dsRNA processing factors, while shRNAs

cannot[42,43]. On this basis, we selected siRNA instead of shRNA for gene knockdown to

avoid the potential pitfall of dsRNA processing in jellyfish.

Here we report a gene knockdown method for jellyfish embryos with siRNA via

electroporation. Using two hydrozoan species, *Clytia hemisphaerica* and *Cladonema*

*pacificum*, we demonstrate that siRNA delivery into fertilized eggs effectively reduces

the expression of endogenous genes. We also confirm the known loss-of-function

phenotypes in *Clytia* after knocking down *GFP1* or *Wnt3*, which are induced in a dose-

dependent manner with siRNA. We further show that knockdown of *Wnt3* in *Cladonema*

embryos results in the reduction of gene expression of the oral marker *Brachyury* in

planula, implicating the Wnt/β-catenin pathway in the control of oral-aboral patterning.

Overall, our siRNA-mediated knockdown approach allows for the manipulation of a

large number of embryos through electroporation and enables functional analysis of

early development, providing a new experimental platform applicable to different jellyfish

species and other marine invertebrates.

**Materials and Methods**

**Animal culture and spawning induction**

*Clytia hemisphaerica* (Z11, Z4B as females and Z4C2, Z23 as males) and *Cladonema*

*pacificum* (6W as females and UN2 as males) were used for this research.

*Clytia hemisphaerica* were cultured using a previously reported method[8] with a
few modifications. Artificial sea water (ASW) was prepared using 220 g SEA LIFE
(Marin Tech) per 5L MilliQ water (Merk Millipore) with antibiotics (40 units/ml of penicillin
and 40 µg/ml of streptomycin). Medusae, embryos, and planula larvae were maintained
at 20ºC. Medusae were fed daily with Vietnamese brine shrimp (A&A Marine LLC, Elk
Rapids, MI, USA). Spawning timing was controlled by a 13 hours (h) dark /11 h light
cycle, and spawning was induced by light. Male and female medusae were transferred
into V-7 cups (AS ONE) before spawning (60 min for male and 90 min for female after
light stimulation).

*Cladonema pacificum* were cultured as previously described[23]. *Cladonema*
medusae were maintained at 22ºC in ASW, which was prepared using SEA LIFE (Marin
Tech) dissolved in tap water with chlorine neutralizer (Coroline off, GEX Co. ltd) (24
p.p.t) and antibiotics (40 units/ml penicillin and 40 µg/ml of streptomycin).  Spawning
timing was controlled by a 30 min dark/23.5 h light cycle, and spawning of male and
female gametes was induced by dark stimulation. Before dark stimulation, adult females
and males were separately transferred into V7 cups (AS ONE) and 60 mm dishes (BD),
respectively.

*Nematostella vectensis* were cultured as previously described[44], with a few
modifications. Briefly, adult animals were maintained in brackish water at a salinity one-
third of artificial seawater (35 g/l, pH 7.5-8.0, SEA LIFE, Marin Tech) and fed with
freshly hatched artemia twice per week. Spawning induction was performed at 26°C
under light for at least 11 h.

**183    siRNA and shRNA**
siRNA sequences (19 mer RNA + 2 mer DNA) for *CheGFP1*, *CheWnt3*, and *CpWnt3*
were designed based on their CDS sequences by Nippon Gene Co., Ltd, and siRNA
duplexes were synthesized by manufacturers (Nippon Gene Co., Ltd and Sigma-
Aldrich, Merk). Lyophilized siRNA was resuspended in RNase free water to a final
concentration of 6 µg/µl as stock solution. The siRNA stock solution was diluted with
RNase free water to a total volume of 10 µl and then added to 90 µl of fertilized eggs in
15% Ficoll ASW just prior to electroporation.

shRNAs were synthesized as described in previous reports[36,38]. Briefly, shRNAs

were synthesized by *in vitro* transcription (IVT) from double stranded DNA templates

using the AmpliScribeTM T7-flashTM transcription kit (Lucigen, Inc.) and were purified

using Direct-Zol TM RNA Miniprep kits (Zymo Research, R2070). Concentrations of

shRNA was measured with a NanoDrop One (Thermo fisher).

**Collection of gametes and electroporation for *Clytia* and *Cladonema* fertilized**

**eggs**

For the preparation of gametes, sexually-mature *Clytia* medusae were treated following

a previously reported method[8], partially modified to fit our experimental equipment.

Adult medusae were maintained on a 13 h dark/11 h light cycle, and light stimuli

induced gametogenesis in 60 min for sperms and 90 min for eggs (Fig. 2A). For

*Cladonema pacificum*, sexually-mature medusae were maintained on a 30 min

dark/23.5 h light cycle. Dark stimulation induced gametogenesis in 25 min. Unfertilized

eggs were collected and resuspended in 15% Ficoll (Nacalai tesque, Japan) in ASW to

prevent the eggs settling from at the bottom of the microtube or sticking to the microtube

surface, keeping the egg solution homogeneous (Fig. 2B)[36,38].

Unfertilized *Clytia* and *Cladonema* eggs were collected in dishes and transferred

into 1.5 ml tubes coated with Ficoll (Nacalai tesque, Japan) to prevent the eggs from

sticking to the tubes. Sperm water was added into eggs, and the eggs and sperm were

incubated for 5 min at room temperature (20-22ºC). The sperm water was then

removed, and fertilized eggs were resuspended in 15% Ficoll in ASW, at which point

siRNA or shRNA was added into fertilized egg solution. The fertilized egg and RNA

mixture was carefully transferred into a 4 mm cuvette and placed into the shockpod

cuvette chamber connected to the Gene Pulser Xcell (Bio-Rad). In most experiments,

the Gene Pulser Xcell was used for pulsing with a square wave voltage of 50 V and a

single pulse duration of 25 ms, except when optimizing electroporation conditions

(Supplementary Figs 1-2). After electroporation, fertilized eggs were transferred to a

dish, and incubated for 10 min. The 15% Ficoll solution was removed and replaced with

new ASW, and the samples were incubated at 20-22ºC.

**Collection of egg sacs, dejellying, and electroporation of *Nematostella* fertilized**

**eggs**

The collection of egg sacs, dejellying, and electroporation steps were performed as

previously described[36] with several modifications.  Briefly, unfertilized egg sacs were

incubated in culture media with sperm at 20-25ºC for 30 min. The fertilized egg sacs

were then dejellied in freshly prepared L-Cysteine solution (0.04 g/ml, Nacalai tesque, in

brackish water; pH adjusted by 5N NaOH at 7.5-8.0) on a shaker for 10 min.  The

dejellied eggs were rinsed with brackish water before they were suspended in Ficoll

solution (Sigma-Aldrich, in brackish water) at a final concentration of 15%. The

suspended eggs and siRNA (5 µg/µl stock solution, in MilliQ) were transferred to a 4

mm cuvette (Bio-Rad) with a final volume of 200 µl and were gently mixed. The Bio-Rad

Gene Pulser Xcell was used for pulsing, with a square wave voltage of 50 V and a pulse

duration of 25 ms. The samples were transferred to a 100 mm dish and maintained in

brackish water for 3 days.

**Imaging**

GFP and Rhodamine fluorescent imaging as well as bright field imaging, including *in*

*situ* hybridization imaging, were taken with an Axio Zoom V16 (ZEISS), and image

processing was done with ZEN 3.4 blue edition (ZEISS) and Fiji/ImageJ software[45].

Intensity of mean values of green fluorescence was measured using the average

brightness (pixel value) of the pixels. To prevent planula movement, we added 0.5 M

sodium azide (final concentration was less than 0.25M).

**Morphological quantification of planula larvae**

After *Wnt3* knockdown by siRNA electroporation of *Clytia* or *Cladonema*, we

approximated the planula body shape into a two dimensional ellipse, measured both the

long and short axes (Figs. 4C and 5D), and calculated the aspect ratio of those values

using Fiji/ImageJ software (Figs, 4D, 5E).

**RT-qPCR**

For *Clytia* and *Cladonema* samples, total RNA was extracted with RNeasy Mini or Micro kits (Qiagen). Lysate was treated with DNaseI (Qiagen) for 15 min at room temperature (RT). cDNA was synthesized with the PrimeScript II 1$^{st}$ strand synthesis kit (Takara bio). RT-qPCR was performed with CFX connect (Bio-Rad) using iTaqTM universal SYBER Green Supermix (Bio-Rad) or the QuantStudio 6 Flex Real-Time PCR System (Thermo Fisher) using TB Green Premix Ex TaqII (Tli RNaseH Plus) (Takara, RR820). Gene expression was normalized to the housekeeping gene, *EF1alpha* or *F-actin capping protein subunit beta*, and the delta-delta-ct method was used for quantification (CFX maestro software, Bio-Rad or QuantStudio 6 Flex Real-Time PCR System software, Thermo Fisher).

For *Nematostella* samples, total RNA was extracted using the RNeasy Mini Kit and RNase-Free DNase Set (Qiagen). cDNA synthesis was conducted using the SuperScript IV First-Strand Synthesis System (Thermo Fisher). RT-qPCR was performed with the StepOnePlusTM Real-Time PCR System (Applied Biosystems, Thermo Fisher) using PowerUP SYBR Green Master Mix (Thermo Fisher). Gene expression was normalized to the housekeeping gene, *NvEf1alpha,* and the delta-delta-ct method was used for quantification (StepOne™ and StepOnePlus™ Software v2.3).

*In situ* **hybridization**

Purified total RNA was reverse transcribed into cDNA by PrimeScript$^{TM}$ 2 1st strand cDNA Synthesis Kit (TaKaRa). The target gene fragments (*Brachyury*, *CpBra*) were amplified from a cDNA library. The primer sets used for PCR cloning are as follows: *CpBra*: 5'-GCTCCCATAAGATCCGGTCG -3'(forward) and 5'-TTTGTCGCAGTCGAAGACCA-3'(reverse), or 5'-GAACGGTGATGGACAGGTCA-3'(forward) and 5'-GGATTCCAAGGATTGGGCGT-3'(reverse). PCR products were sub-cloned into the TAK101 vector (TOYOBO). The resulting plasmids were used for RNA probe synthesis with digoxigenin (DIG) labeling mix (Roche) and T7 or T3 RNA polymerase (Roche), according to the insert direction.

Planula larvae were anesthetized in 50% 0.5 M NaN$_3$ in ASW for 5 min and fixed

overnight at 4°C with 4% paraformaldehyde (PFA) in ASW. Fixed samples were washed

three times with PBS containing 0.1% Tween-20 (PBST), followed by pre-hybridization

in hybridization buffer (HB buffer: 5×SSC, 50% formamide, 0.1% Tween-20, 50 µg/ml

tRNA, 50 µg/ml heparin) at 55°C for 2 h. Samples were hybridized with HB Buffer

containing the probes (final probe concentration: 0.5-1 ng/µL in HB Buffer) at 55°C

overnight and washed. The samples were incubated with 0.05% alkaline phosphatase

(AP)-conjugated anti-DIG antibodies (Roche) in 1% blocking reagent (Roche) in Maleic

acid buffer for 4 h at room temperature. Colorimetric reactions were performed by

NBT/BCIP (Roche) in alkaline phosphatase buffer (100 mM NaCl, 100 mM Tris-HCl (pH

9.5), 50 mM MgCl$_2$, 0.1% Tween-20) until the signals were detected.

**Sequence alignment and phylogenetic analysis**

Amino acid sequence alignment of Wnt3 and Wnt family proteins were performed by

ClustalW[46]. GenBank accession numbers and protein names are listed in

Supplementary Table 3 and 4. The alignment data was visualized on Jalview software[47].

Maximum likelihood (ML) tree calculation was inferred on ClustalW with the PhyML

bootstrap method. Branch supports were computed out of 100 bootstrapped trees. The

tree visualization was created using MEGA-X software[48].

**qPCR primers**

| Name | Sequence (5'->3') | |
|---|---|---|
| CpWnt3 qP FW | CAAATGTGGTCGATCAAACG | This study |
| CpWnt3 qP RV | TACGCCTTCTGCAACACTTG | This study |
| CpEF1-a_qPCR_F7 | GGTCAATCTCGTTCCCTCCA | This study |
| CpEF1-a_qPCR_R7 | TTTCCACCAGAGGTATCGGC | This study |
| CpBrachyury F | AAGGCGTATGTTTCCGGTCC | This study |
| CpBrachyury R | CAACGATGGTCTTCGACTGC | This study |
| CpFoxQ2a F | AGTTGGAGAAACAGCGTTCG | This study |
| CpFoxQ2a R | CCGTTCGCAAAGTCGTCAAA | This study |
| CpCapZbeta_qPCR_F | AAAGAAAGCTGGAGACGGTTCA | This study |
| CpCapZbeta_qPCR_R | GTAGTGGGCATTTCTTCCGC | This study |
| CheGFP1 qP F | TTGCTGTCCGAATAGTGCAG | Fourrage et al., 2014 Open Biol |

| Name | Sequence | Reference |
|---|---|---|
| CheGFP1 qP RV | GACAACTCCTCCTCCGAGTG | Fourrage et al., 2014 Open Biol |
| CheGFP1_2 qP F | ACAATCGCGTCACACTTAAAGG | This study |
| CheGFP1_2 qP RV | CGTTGTTTTCTTTGTCCGGC | This study |
| CheWnt3 qP F | ATCATGGCAGGTGGAAACTC | Leclère et al., 2012 Dev Biol |
| CheWnt3 qP RV | CCCCATTTCCAACCTTCTTC | Leclère et al., 2012 Dev Biol |
| CheEF1a qP F | TGCTGTTGTCCCAATCTCTG | Leclère et al., 2012 Dev Biol / Fourrage et al., 2014 Open biol |
| CheEF1a qP RV | AAGACGGAGTGGTTTGGATG | Leclère et al., 2012 Dev Biol / Fourrage et al., 2014 Open biol |
| NvEf1alpha FW | GGTTGCCTCTTCGCTTACCACT | This strudy |
| NvEf1alpha RV | CGTTCCTGGCTTTAGGACAC | This study |
| NvBra1_Fw_1032 | CGGGCTCACACTCTCACTTA | This study |
| NvBra1_Rv1174 | CTTGCGGTATGGTGTTCCAG | This study |

**siRNA design**

| Name | Sequence (5'->3') | | ORF |
|---|---|---|---|
| siCheGFP1_1 sense | CCAUCAGCUUCGAAAAUGAdTdT | This study | HQ397706.1 |
| antisense | UCAUUUUCGAAGCUGAUGGdTdT | This study | |
| siCheGFP1_2 sense | UGAUGGCGCUUAUAAAGUUdTdT | This study | HQ397706.1 |
| antisense | AACUUUAUAAGCGCCAUCAdTdT | This study | |
| siCheGFP1_3 sense | CGGACUCAUUCUUCAAAAAdTdT | This study | HQ397706.1 |
| antisense | UUUUUGAAGAAUGAGUCCGdTdT | This study | |
| siCheWnt3_1 sense | GCAUGUGACUGUAAAUUUAdTdT | This study | EU374721.1 |
| antisense | UAAAUUUACAGUCACAUGCdTdT | This study | |
| siCheWnt3_2 sense | ACAGCUUGGUAAAAUGUUAdTdT | This study | EU374721.1 |
| antisense | UAACAUUUUACCAAGCUGUdTdT | This study | |
| siCpWnt3_1 sense | GUGGAACUGUAGUGUUUCAdTdT | This study | EU374721.1 |
| antisense | UGAAACACUACAGUUCCACdTdT | This study | |
| siCpWnt3_2 sense | AGCGUGUGCUGAAGGUAAAdTdT | This study | EU374721.1 |
| antisense | UUUACCUUCAGCACACGCUdTdT | This study | |
| siNvBra1_1 sense | GAAGAGAUCACGAGUCUAAdTdT | This study | AF540387.2 |
| antisense | UUAGACUCGUGAUCUCUUCdTdT | This study | |
| siNegative_Control-1 for Nv sense | GCAACACGCAGAGTCGTAAdT | Same sequence as Karabulut et al, 2019 Dev Biol | |
| antisense | TTACGACTCTGCGTGTTGCdT | Same sequence as Karabulut et al, 2019 Dev Biol | |

313 **shRNA template**

| Name | Sequence (5'->3') | |
|------|-------------------|---|
| CheWnt3 _SI1_F | TAATACGACTCACTATA GCATGTGACTGTAAATTTA TTCAAGAGA TAAATTTACAGTCACATGC TT | This study |
| CheWnt3 _SI1_R | AAGCATGTGACTGTAAATTTATCTCTTGAATAAATTTAC AGTCACATGCTATAGTGAGTCGTATTA | This study |
| CheWnt3 _SI2_F | TAATACGACTCACTATA ACAGCTTGGTAAAATGTTA TTCAAGAGA TAACATTTTACCAAGCTGT TT | This study |
| CheWnt3 _SI2_R | AAACAGCTTGGTAAAATGTTATCTCTTGAATAACATTTT ACCAAGCTGTTATAGTGAGTCGTATTA | This study |

**Graphs and statistical analysis**

All graphs were prepared in Excel. To assess phenotypes and RT-qPCR statistical

significance, we used the percentage values and delta-delta-Ct values to perform two-

tailed Student's t tests, except in Figure 5F, where a one-tailed Student's t test was

used. For statistics related to fluorescence intensity mean value and aspect ratio,

Kolmogorov–Smirnov normality tests were used. All statistical tests were performed at

https://www.socscistatistics.com.

**Results**

**Optimization of electroporation conditions**

In both hydrozoan jellyfish *Clytia hemisphaerica* and *Cladonema pacificum*,

gametogenesis is regulated by light-dark transitions[6,49]. A previous report showed that

*Clytia* medusae release eggs within 110-120 min and sperm within 60-90 min upon light

illumination after 3-8 h of darkness[12]. *Cladonema pacificum*, which live along coastal

areas in Japan, exhibit two types of gametogenesis depending on habitat: the dark-light

transition (light stimulation type) and the light-dark transition (dark stimulation type),

respectively. Dark stimulation *Cladonema* medusae, which we used in this study,

release eggs or sperm within 30 min of dark stimulus after 20-24 h of constant exposure

to light[25,49]. By utilizing the characteristic gametogenesis of these jellyfish species, we

can control egg/sperm release to induce fertilization at any time of day. Indeed, we

routinely obtained *Clytia* sperm within 60 min and *Clytia* eggs within 90 min of light on a

13h dark/11h light cycle (Fig. 2A). Similarly, we collected gametes from *Cladonema*

within 25 min of dark stimulation on a 23.5h light/30 min dark cycle (Fig. 2A).

To determine optimal electroporation parameters, we first tested different

electroporation conditions and monitored the delivery efficiency of the red fluorescent

dye Rhodamine-Dextran into unfertilized *Clytia* and *Cladonema* eggs (Supplementary

Figs. 1-2). Given that the average molecular weight of siRNA (21-mer, ~13,300 Da) is

similar to the molecular size of Rhodamine-Dextran (10,000 Da), electroporation trials

with Rhodamine-Dextran can visually confirm that small molecules are incorporated into

jellyfish eggs, as previously shown in *Hydractinia* eggs[38]. The collected eggs were

suspended in 15% Ficoll artificial seawater to prevent precipitation and to make a

homogeneous solution (Fig. 2B)[36,38]. We performed electroporation trails for

Rhodamine-Dextran using a cuvette with a 4 mm gap and tested several conditions of

voltage (V) and pulse length time (milliseconds, ms) using a conventional

electroporation system (Fig. 2C). As previous work showed that increasing the number

of pulses does not increase the Dextran delivery rate but decreases the survival rate of

embryos[38], we fixed the number of pulses to one and incubated unfertilized eggs with 1

mg/ml Dextrane-Rhodamine solution. *Clytia* eggs have little autofluorescence in the red

channel, and without electroporation (No EP), red fluorescence was rarely detected in

unfertilized eggs (Fig. 2D and Supplementary Fig. 1A). We compared the eight different

electroporation conditions and found that the rhodamine fluorescence was detected in

*Clytia* eggs and correlated with the strength of electric voltage, as long as the voltage

was in the range of 50-100 V（Supplementary Figs. 1B-1D）. By contrast, when the

voltage exceeded 200 V, the eggs ruptured from physical damage, and cellular debris

was observed（Supplementary Figs, 1E-1G）.

We performed similar electroporation trials on unfertilized *Cladonema* eggs under

eight different conditions. While red autofluorescence was negligible in the *Cladonema*

eggs in the absence of electroporation (Fig. 2D and Supplementary Fig. 2A),

Rhodamine fluorescence was frequently detected with 50 V (Fig. 2D and

Supplementally Fig. 2E). In the more intense conditions beyond 100 V (Supplementally

Figs. 2F and 2G), the eggs were severely damaged. Based on these results, we

established the optimal electroporation conditions (50 V, 1 pulse, 25 ms) that provide

high frequency of Dextran-positive eggs and low levels of cell damage for both jellyfish

eggs (Fig. 2D). Of note, during the initial trials, we used unfertilized eggs, but we found

that the electroporation process severely affected the survival rate of subsequent

embryos. In the case of *Cladonema*, electroporation before fertilization severely

decreased the survival rate of fertilized eggs (7.04% at 1 h post fertilization) while

electroporation after fertilization did not influence the survival rate (No EP: 59.67%, With

EP: 60.15%) (Supplementary Table 1). This dissimilarity is likely due to the decrease of

fertilization rate after electroporation, and both elapsed time after spawning and physical

damage by electroporation can affect fertilization efficiency. Therefore, we decided to

conduct further electroporation experiments with fertilized eggs.

**Gene knockdown with siRNA in *Nematostella vectensis***

Before attempting gene knockdown via siRNAs electroporation in fertilized jellyfish

eggs, we first used *Nematostella vectensis,* where egg electroporation methodology has

been established[36]. We selected *Nematostella Brachyury* (*NvBra*), a gene expressed in

the blastopore margin that functions in the early embryo[50,51], as a target. After

electroporation of siRNA targeting *NvBra* in fertilized eggs, we examined *NvBra*

expression in planula larvae at 3 days-post-fertilization (dpf) by RT-qPCR and found

that *NvBra* expression was drastically suppressed compared to no electroporation

controls and negative control siRNAs (Supplementary Fig. 3). We further evaluated

survival rates after electroporation and confirmed that siRNA electroporation does not

severely affect survival rates of 3 dpf planula in any of the tested conditions

(Supplementary Table 2). These results suggest that siRNA delivery into fertilized eggs

via electroporation is an effective gene knockdown method that is comparable to the

effect of shRNA delivery into unfertilized eggs via electroporation[36], supporting our

attempt to perform gene knockdown via siRNA electroporation using fertilized jellyfish

eggs.

**Endogenous *GFP1* knockdown with siRNA in *Clytia hemisphaerica***

The hydrozoan jellyfish *Clytia* endogenously expresses green fluorescent proteins,

which are encoded by 14 genes grouped into 4 clusters (*CheGFP1-4*)[16]. The expression

pattern for each GFP changes by stage in the life cycle: for instance, *Clytia* eggs

express maternal *CheGFP2* mRNA, and planula larvae show typical ring-like *CheGFP1*

fluorescence in the lateral ectoderm at 3 dpf[11]. While *CheGFP1* gene expression

becomes dominant in the planula stage; *CheGFP2, CheGFP3,* and *CheGFP4* are

expressed in different tissues and organs in the medusa[52].  Momose *et al*. succeeded in

genome editing by injecting CRISPR/Cas9 targeting the endogenous *CheGFP1* into

fertilized eggs, and confirmed the knockout effect by loss of green fluorescence in 3 dpf

planula larvae[11]. In addition, *CheGFP1* is not essential for animal survival, which allows

for the monitoring of knockdown effects after embryogenesis. For these reasons, we

chose endogenous *CheGFP1* for the first target gene of siRNA knockdown.

In order to maximize the effect of gene knockdown, we used a combination of

three different siRNAs against *CheGFP1* (Accession No. HQ397706.1; Supplementary

Fig. 4A) in a mixture with a 1:1:1 ratio, and electroporated 600 ng/µl of siRNA mix (200

ng/µl per each siRNA) into fertilized eggs (Fig. 3A). To verify the effect of gene

knockdown, we measured green fluorescence intensity in 2 dpf planula and found that

siRNA (600ng/µl siRNA mix) electroporated embryos showed dramatically reduced

fluorescence intensity compared to the no-siRNA controls (Figs. 3B-D; mean intensity

value: $5.7 \times 10^3$ for 0 ng/µl control, $1.8 \times 10^3$ for 600 ng/µl). To evaluate the knockdown

effects at varying doses of siRNA, we next electroporated different concentrations of the

*CheGFP1* siRNA mix (0, 150, 300, and 600 ng/µl) into fertilized eggs. While control 3

dpf planulae with no-electroporation (No EP) as well as those with no-siRNA showed

typical ring-shape GFP fluorescence in the ectoderm, many planulae with *CheGFP1*

siRNA electroporation lost green fluorescence（Figs. 3E-3H）. To evaluate the GFP

knockdown effects, we classified planulae with typical GFP1 fluorescence as "wild type"

and planulae with reduced fluorescence as "reduced" for further quantification (Figs. 3I

and 3J). The percentage of wild type phenotype dramatically reduced in an siRNA dose-

dependent manner (Fig. 3K). To further quantify the above result on a molecular level,

we examined relative *CheGFP1* mRNA levels by RT-qPCR using the primer set that

was evaluated in a previously study[52]. Consistent with the image analysis results, we

confirmed a marked reduction of *CheGFP1* gene expression at different concentrations

of *CheGFP1* siRNA (Fig. 3L). These results demonstrate that siRNAs targeting

*CheGFP1* repress its expression upon electroporation and that the knockdown effects

are dose-dependent.

<span style="color:red">Given the presence of multiple *CheGFP1* loci (seven nearly identical *CheGFP1*</span>

<span style="color:red">genes) in the *Clytia* genome[16], it was not clear whether three siRNAs against *GFP1*</span>

<span style="color:red">sufficiently suppress gene expression in different versions of *CheGFP1*, particularly</span>

<span style="color:red">those expressed in the planula stage (Supplementary Fig. 4A）. We thus performed RT-</span>

<span style="color:red">qPCR using the newly designed primer set that amplifies all seven *CheGFP1* genes and</span>

<span style="color:red">found a dramatic reduction of gene expression upon electroporation of *CheGFP1*</span>

<span style="color:red">siRNAs (600 ng/µl)（Supplementary Fig. 4B）. Altogether, these results indicate that our</span>

<span style="color:red">combination of *CheGFP1* siRNAs suppresses gene expression of *CheGFP1* in the</span>

<span style="color:red">*Clytia* planula.</span>

***Wnt3* knockdown with siRNA in *Clytia hemisphaerica***

Since electroporation of siRNAs targeting endogenous *CheGFP1* was shown to be

effective, we next decided to investigate the effects of knocking down other genes that

play important roles in early development. Wnt3 is a secreted signaling protein in the

Wnt/β-catenin pathway that is involved in diverse developmental processes including

cell fate decisions and patterning[53,54]. In *Clytia*, *Wnt3* RNA is maternally localized to the

animal cortex, the future oral side of the <span style="color:red">embryo</span>, and Wnt3 organizes axial patterning

as the main ligand for the Wnt/β-catenin pathway in the early embryonic stage[10]. After

embryogenesis, the *Clytia* planula normally elongates along the oral-aboral axis and

becomes oval in shape（Fig. 4A）. By contrast, after *CheWnt3* morpholino injection, the

morphant loses oral-aboral axis polarity and exhibits a spherical shape[10]. This raises the

possibility that knockdown of *CheWnt3* via siRNA may result in a similar spherical

morphology for *Clytia* larvae as is observed in *CheWnt3* morphants. Therefore, to

evaluate siRNA knockdown effects on *Clytia*, we chose *CheWnt3* as our second target.

We used a mixture of two different siRNAs (1:1 ratio) targeting *CheWnt3*, and

carried out electroporation of the siRNA mix into *Clytia* fertilized eggs. After

electroporation of *CheWnt3* siRNAs, 1 dpf planula larvae showed nearly complete

spherical morphology (Fig. 4B), which is reminiscent of *CheWnt3* morphants[10]. To

quantify the effect of siRNA knockdown on *Clytia* larval morphology, we calculated the

aspect ratio in two dimensions, where values greater than 1.0 shows the tendency to be

oval (ellipsoid-like) while a value of 1.0 indicates a perfect circle (sphere) (Fig. 4C). The

control larvae with 0 ng/µl siRNA showed a median aspect ratio of 1.69, indicating a

typical ellipsoid-like morphology. By contrast, larvae that underwent electroporation with

*CheWnt3* siRNAs (400 ng/µl) showed a median aspect ratio of 1.16, suggesting a much

greater tendency toward spherical morphology (Fig. 4D). These results quantitively

confirmed the effect of *CheWnt*3 knockdown on larval morphology, which phenocopies

*CheWnt3* morphants[10]. To analyze the level of gene-specific knockdown at varying

doses of siRNAs, we next electroporated mixtures of 0, 100, 200, and 400 ng/µl of

*CheWnt3* siRNAs into *Clytia* fertilized eggs (Figs. 4E-4H). To evaluate *CheWnt3* siRNA

effects on morphology, we classified 1 dpf planulae with elongated ellipsoid-like shapes

as "normal" and planulae with spherical shapes as "spherical" (Figs. 4I and 4J). The

percentage of normal phenotype dramatically reduced after *CheWnt3* siRNA-

electroporation (< 1.0 % in 400 ng/µl), even at the lowest dose (5.2% in 100 ng/µl) (Fig.

4K). To further evaluate the result at a molecular level, we analyzed relative *CheWnt3*

mRNA levels by RT-qPCR and confirmed a significant reduction in *CheWnt3* gene

expression at all concentrations (Fig. 4L). Of note, after electroporation with universal

negative control siRNA（Nippon Gene Co., Ltd.）, both morphological phenotypes and

relative gene expression were comparable to no-siRNA controls (Figs. 4K and 4L),

further confirming that siRNA-mediated knockdown by electroporation is target-specific.

These results together indicate that siRNAs targeting endogenous genes effectively

knock down their expression in *Clytia* embryos.

Effective gene knockdown via shRNA electroporation has previously been

reported using two cnidarian polyps, *Nematostella vectensis* and *Hydractinia*

*symbiolongicarpus*[36,38]. To test whether shRNAs are also effective in the hydrozoan

jellyfish *Clytia*, we designed shRNAs targeting *CheWnt3* using the same 19 bp target

sequence that was used for siRNAs, and performed electroporation on fertilized eggs

with a mix of two different *Wnt3* shRNAs. After electroporation of *Wnt3* shRNAs, relative

*Wnt3* mRNA expression in 1 dpf planula was weakly down-regulated (0.83), and was

not significant compared to the striking effects of *Wnt3* siRNA electroporation (0.039) at

the same concentration (400 ng/µl) (Fig. 4L). In addition, electroporation of *Wnt3*

siRNAs resulted in an extensive morphological change toward spherical shaped

planulae, showing almost no "normal" phenotype (0.71%), whereas electroporation of

shRNA had a limited effect (60.4%) (Fig. 4K). These results suggest that the effect of

siRNA is much stronger than that of shRNA in *Clytia* when treated with the same target

sequence and concentration.

**Identification of *Wnt3* in *Cladonema pacificum***

In contrast to *Clytia*, the sole genetic model jellyfish, previous studies have neither

established genetic manipulation nor performed genome sequencing of *Cladonema*

*pacificum*, despite its unique biological features. This is partly because the extremely

small size of the *Cladonema* egg makes microinjection difficult. By testing whether

siRNA-mediated knockdown via electroporation can work in *Cladonema*, we can start to

manipulate genes in *Cladonema* while also demonstrating that siRNA electroporation

can be applicable to other jellyfish species. While green fluorescent protein (GFP) is

often an easy early target for gene manipulation in emerging organisms, *Cladonema*

does not exhibit apparent endogenous green fluorescence expression at any stage,

unlike *Clytia*. We instead turned our focus to *Wnt3* as the target for gene knockdown in

*Cladonema* since Wnt3 plays an important role in early embryonic development in

several cnidarian species[55,56].

To identify the *Wnt3* ortholog in *Cladonema*, we utilized RNA-seq results from the

*Cladonema* polyp, stolon, and medusa manubrium (data not shown). We performed

*Cladonema Wnt3* CDS annotation using the *Clytia* CDS database (MARIMBA, Marine

models database: http://marimba.obs-vlfr.fr) and found one contig annotated with *Wnt3*

(*CpWnt3*) and multiple *Wnt* genes (Supplementary Figs. 5 and 6). We then performed

phylogenic analysis using the neighbor-joining method, and confirmed *CpWnt3*

(LC720435) is grouped with medusozoan *Hydra vulgaris* (*HvWnt3*) and *Clytia Wnt3*

(*CheWnt3*) rather than Anthozoa *Nematostella* and *Acropora digitifera* (Supplementary

Fig. 5A).  Multiple sequence alignment further showed highly conserved amino acids

sequences among different species including bilaterians and cnidarians (Supplementary

Fig. 5B). These findings raise the possibility that *CpWnt3* has a similar function in

controlling the Wnt/β-catenin pathway in *Cladonema.*

***Wnt3* knockdown with siRNA in *Cladonema pacificum***

Does *Wnt3* function during *Cladonema* embryogenesis, particularly during axial patterning? Interestingly, *Cladonema* planula larvae do not show a clear elongated shape (Fig. 5A), as observed in *Clytia*. It is thus possible that morphogenesis, including axial patterning and/or Wnt/β-catenin pathway function, differs between these two jellyfish species. It is also possible that the limited elongation in *Cladonema* planulae might hamper phenotypical appearance upon inhibition of Wnt/β-catenin signaling.

In order to verify that siRNA electroporation works in *Cladonema* and that *Wnt3* affects axial patterning, we performed electroporation of siRNAs targeting *CpWnt3*. We prepared fertilized *Cladonema* eggs as we did in *Clytia*, and used a mixture of two different siRNAs against *CpWnt3* CDS with a 1:1 ratio (300 ng/μl) and performed electroporation with the previously established parameters (50V, 1 pulse, 25 ms) in fertilized *Cladonema* eggs. After electroporation of siRNAs for *CpWnt3*, planula larvae did not exhibit morphological differences compared to controls（Figs. 5A-5C; median aspect ratio: 1.21 for 0 ng/μl control, 1.16 for 300 ng/μl for si*CpWnt3*）. We then quantified the morphological phenotypes by calculating the aspect ratio of 1 dpf planulae and confirmed that *CpWnt3* knockdown does not cause a significant morphological change (Figs. 5D and 5E), which is consistent with the possibility that *Cladonema* planulae simply exhibit limited elongation.

Another possibility that would explain the above result is a defect in the siRNA knockdown itself. To test this potential explanation, we carried out RT-qPCR using mRNA samples from 1 dpf planula after electroporating different concentrations of *CpWnt3* siRNA mix (0, 150, and 300 ng/μl) into fertilized eggs. We confirmed a reduction of *Wnt3* gene expression in an siRNA dose-dependent manner (Fig. 5F), eliminating siRNA knockdown defects as the cause of our initial result.

To further confirm whether the Wnt/β-catenin pathway is affected by *Wnt3* knockdown, we examined the gene expression of axial markers *Brachyury* (*CpBra*) and *FoxQ2a* (*CpFoxQ2a*), whose expression are influenced by the Wnt/β-catenin pathway in *Clytia*[10]. From RT-qPCR, we found that *CpWnt3* knockdown causes a decrease in *CpBra* gene expression and an increase in *CpFoxQ2a* gene expression (Supplementary Fig. 7). We also examined *CpBra* expression by *in situ* hybridization and confirmed the

reduction of *CpBra* expression on the oral side of planula larvae (Figs. 5G-5I), which is

similar to the phenotype exhibited by *CheWnt3* morphants in *Clytia*[10]. Notably, after

*CpWnt3* siRNAs electroporation, the rate of metamorphosis from planula to primary

polyp decreased dramatically (Supplementary Table 5), implying potential

developmental defects upon abrogation of the Wnt/β-catenin pathway. These results

suggest that, although overall oral-aboral polarity is less prominent in *Cladonema*

compared to *Clytia* at the morphological level, the axial patterning mechanism mediated

by Wnt/β-catenin signaling may be conserved between these two jellyfish species.

**Discussion**

In this study, we have established a method to knock down endogenous genes in two

hydrozoan jellyfish, *Clytia hemisphaerica* and *Cladonema pacificum*, via the siRNA

electroporation of fertilized eggs. We showed that knockdown of endogenous *GFP1* in

*Clytia* causes the loss of GFP fluorescence in the planula stage (Fig. 3), as previously

achieved by CRISPR/Cas9-mediated *GFP1* knockout[11]. We also confirmed that

knockdown of *Wnt3* in *Clytia* induces spherical morphology of the planula (Fig. 4),

mirroring the results of injections of *Wnt3* morpholino antisense oligo[10]. We further

succeeded in efficient repression of *Wnt3* gene expression in *Cladonema* after *Wnt3*

siRNA electroporation and found that expression of axial patterning genes is controlled

by the Wnt/β-catenin pathway (Fig. 5), suggesting that the conserved mechanism is

involved in embryogenesis and planula morphogenesis across hydrozoan jellyfish

species.

Our results show that the knockdown efficiency of siRNA is much greater than

that of shRNA for electroporation of *Clytia* fertilized eggs with the same *Wnt3* target

sequence in the same concentration (Fig. 4K-4L). One possibility that would explain

such distinct effects between siRNA and shRNA electroporation is the existence of

differences in the RNAi machinery or RNA processing efficiency among cnidarians.

During RNAi in mammals, the RNase III Dicer protein processes shRNA in collaboration

with cofactors TRBP (Transactivation response element RNA-binding protein) and

PACT（protein activator of the interferon-induced protein kinase）to produce a mature

form of siRNA[39]. Mature siRNA in the Dicer and TRBP/PACT complex is associated

with the Argonaute protein and cleaves endogenous complementary mRNAs as the

RNA-induced silencing complex [30,57]. These small RNA biogenesis factors, Dicers and

cofactors (TRBP in bilaterians, HYL1 in plants and cnidarians), are found in

*Nematostella vectensis, Acropora digitifera,* and *Hydra vulgaris*[41]. Although Dicers

(TCONS_00004571 and TCONS_00004525) and Hyl1 (TCONS_00010722) are

predicted to exist in *Clytia* based on MARIMBA transcriptome data, the expression level

of Dicers in oocytes and early gastrulation stages is lower than that in other stages,

suggesting the possibility that insufficient shRNA to siRNA processing efficiency is

responsible for the difference in knockdown efficiency between siRNA and shRNA

electroporation. It will be interesting to elucidate the detailed molecular mechanism of

RNAi in different cnidarian species and address RNA processing efficiency in distinct

developmental stages. The lower efficiency of knockdown by shRNA compared to

siRNA in *Clytia* could also be simply explained by lower shRNA electroporation

efficiency. This possibility could be tested by inserting shRNA or siRNA into fertilized

eggs by microinjection, instead of electroporation, which would be followed by

assessing gene expression and phenotypes.

During *Clytia* early embryogenesis, *Wnt3* RNA is localized to the animal cortex of

the egg and contributes to the formation of the oral-aboral axis[10]. Accordingly,

knockdown of *Wnt3* in *Clytia* fertilized eggs induces suppression of axis formation and

planula elongation (Fig. 4). In contrast, despite the fact that no morphological phenotype

was observed after *Wnt3* knockdown in *Cladonema*, gene expression of the conserved

axial patterning genes, *Brachyury* and *FoxQ2a,* was disrupted (Figs. 5G-5H and

Supplementary Fig. 7). In addition to *Clytia,* the Wnt/β-catenin pathway is involved in

axis formation during cnidarian embryogenesis including *Nematostella*[44,55,58]*,*

*Acropora*[59]*, and Hydractinia*[56]. Furthermore, given that the Wnt/β-catenin pathway is

also associated with axis formation during metamorphosis in *Hydractinia*[56,60] and

*Clytia*[61], the reduced metamorphosis rate observed in *CpWnt3* knockdown *Cladonema*

may be attributed to disrupted axis formation (Supplementary Table 5). Taken together,

our data support the pivotal role of Wnt/β-catenin signaling in cnidarian development.

Why do *Cladonema* planulae show a less elongated morphology during the

transition from embryos? In the torpedo-shaped *Clytia* planula, the Wnt/Planar Cell

Polarity (PCP) pathway is activated during the gastrulation phase, which causes

elongation of the body axis[62]. In contrast, the PCP pathway is independent of the Wnt/β-catenin pathway in *Nematostella*, which have relatively short planula morphology[63]. It is thus possible that the Wnt/PCP pathway is not associated with cell polarity and elongation in *Cladonema*, but more detailed studies are needed.

An interesting feature of jellyfish is that their morphology dramatically changes across their life cycle from planula to polyp and then to medusa. In particular, polyps and medusae, different adult stages of the same animal, exhibit distinct regenerative ability, lifespan, and behaviors. Although regeneration mechanisms have been extensively studied in polyp-only animals such as *Hydra* and *Hydractinia*, classical studies have used the medusa stage of several jellyfish species and demonstrated their regenerative potential[21]. Recent work using *Clytia* medusae has further shown a remarkable remodeling and repatterning mechanism orchestrated by muscle systems upon organ loss[17]. At the behavior level, the medusae of the upside-down jellyfish *Cassiopea* exhibit a sleep-like state[64], which is similarly observed in *Hydra* polyps[65]. More recent work using transgenic *Clytia* has characterized feeding behaviors in medusae at a neural-network resolution[19]. To understand how these diverse biological processes in jellyfish are regulated at the molecular level, the functions of specific genes at various stages must be analyzed. Although gene knockdown by dsRNA electroporation into polyps in the scyphozoan jellyfish *Aurelia* has been reported[66], genetic manipulation in medusae has not yet been achieved. Given that siRNA electroporation is applicable to *Hydra* polyps[33,67] in addition to cnidarian fertilized eggs, we now have a better chance to apply this technique to the medusa stage, although, because they are susceptible to electric shock, electroporation parameters must be optimized in order to achieve gene knockdown in medusae. Collectively, our gene knockdown via siRNA electroporation method will complement the existing shRNA electroporation approach in cnidarian polyps and will enable molecular-level analysis of the vast biological phenomena exhibited across the different life stages in jellyfish.

## References

1       Technau, U. & Steele, R. E. Evolutionary crossroads in developmental biology: Cnidaria. *Development* **138**, 1447-1458, doi:10.1242/dev.048959 (2011).

2    Zapata, F. *et al.* Phylogenomic Analyses Support Traditional Relationships within
Cnidaria. *PLoS One* **10**, e0139068, doi:10.1371/journal.pone.0139068 (2015).
3    Leclere, L. & Rottinger, E. Diversity of Cnidarian Muscles: Function, Anatomy,
Development and Regeneration. *Front Cell Dev Biol* **4**, 157,
doi:10.3389/fcell.2016.00157 (2016).
4    Holstein, T. W., Hobmayer, E. & Technau, U. Cnidarians: an evolutionarily
conserved model system for regeneration? *Dev Dyn* **226**, 257-267,
doi:10.1002/dvdy.10227 (2003).
5    Bosch, T. C. G. *et al.* Back to the Basics: Cnidarians Start to Fire. *Trends*
*Neurosci* **40**, 92-105, doi:10.1016/j.tins.2016.11.005 (2017).
6    Houliston, E., Momose, T. & Manuel, M. Clytia hemisphaerica: a jellyfish cousin
joins the laboratory. *Trends Genet* **26**, 159-167, doi:10.1016/j.tig.2010.01.008
(2010).
7    Houliston, E., Leclere, L., Munro, C., Copley, R. R. & Momose, T. Past, present
and future of Clytia hemisphaerica as a laboratory jellyfish. *Curr Top Dev Biol*,
doi:10.1016/bs.ctdb.2021.12.014 (2022).
8    Lechable, M. *et al.* An improved whole life cycle culture protocol for the
hydrozoan genetic model Clytia hemisphaerica. *Biol Open* **9**,
doi:10.1242/bio.051268 (2020).
9    Momose, T. & Houliston, E. Two oppositely localised frizzled RNAs as axis
determinants in a cnidarian embryo. *PLoS Biol* **5**, e70,
doi:10.1371/journal.pbio.0050070 (2007).
10    Momose, T., Derelle, R. & Houliston, E. A maternally localised Wnt ligand
required for axial patterning in the cnidarian Clytia hemisphaerica. *Development*
**135**, 2105-2113, doi:10.1242/dev.021543 (2008).
11    Momose, T. *et al.* High doses of CRISPR/Cas9 ribonucleoprotein efficiently
induce gene knockout with low mosaicism in the hydrozoan Clytia hemisphaerica
through microhomology-mediated deletion. *Sci Rep* **8**, 11734,
doi:10.1038/s41598-018-30188-0 (2018).
12    Quiroga Artigas, G. *et al.* A gonad-expressed opsin mediates light-induced
spawning in the jellyfish Clytia. *Elife* **7**, doi:10.7554/eLife.29555 (2018).

13    Quiroga Artigas, G. *et al.* A G protein-coupled receptor mediates neuropeptide-

induced oocyte maturation in the jellyfish Clytia. *PLoS Biol* **18**, e3000614,

doi:10.1371/journal.pbio.3000614 (2020).

14    Lapebie, P. *et al.* Differential responses to Wnt and PCP disruption predict

expression and developmental function of conserved and novel genes in a

cnidarian. *PLoS Genet* **10**, e1004590, doi:10.1371/journal.pgen.1004590 (2014).

15    Condamine, T. *et al.* Molecular characterisation of a cellular conveyor belt in

Clytia medusae. *Dev Biol* **456**, 212-225, doi:10.1016/j.ydbio.2019.09.001 (2019).

16    Leclere, L. *et al.* The genome of the jellyfish Clytia hemisphaerica and the

evolution of the cnidarian life-cycle. *Nat Ecol Evol* **3**, 801-810,

doi:10.1038/s41559-019-0833-2 (2019).

17    Sinigaglia, C. *et al.* Pattern regulation in a regenerating jellyfish. *Elife* **9**,

doi:10.7554/eLife.54868 (2020).

18    Chari, T. *et al.* Whole-animal multiplexed single-cell RNA-seq reveals

transcriptional shifts across Clytia medusa cell types. *Sci Adv* **7**, eabh1683,

doi:10.1126/sciadv.abh1683 (2021).

19    Weissbourd, B. *et al.* A genetically tractable jellyfish model for systems and

evolutionary neuroscience. *Cell* **184**, 5854-5868 e5820,

doi:10.1016/j.cell.2021.10.021 (2021).

20    Leclere, L. *et al.* Maternally localized germ plasm mRNAs and germ cell/stem cell

formation in the cnidarian Clytia. *Dev Biol* **364**, 236-248,

doi:10.1016/j.ydbio.2012.01.018 (2012).

21    Fujita, S., Kuranaga, E. & Nakajima, Y. I. Regeneration Potential of Jellyfish:

Cellular Mechanisms and Molecular Insights. *Genes (Basel)* **12**,

doi:10.3390/genes12050758 (2021).

22    Fujiki, A., Hou, S., Nakamoto, A. & Kumano, G. Branching pattern and

morphogenesis of medusa tentacles in the jellyfish Cladonema pacificum

(Hydrozoa, Cnidaria). *Zoological Lett* **5**, 12, doi:10.1186/s40851-019-0124-4

(2019).

23    Fujita, S., Kuranaga, E. & Nakajima, Y. I. Cell proliferation controls body size

growth, tentacle morphogenesis, and regeneration in hydrozoan jellyfish

Cladonema pacificum. *PeerJ* **7**, e7579, doi:10.7717/peerj.7579 (2019).

24   Hou, S., Zhu, J., Shibata, S., Nakamoto, A. & Kumano, G. Repetitive
accumulation of interstitial cells generates the branched structure of Cladonema
medusa tentacles. *Development* **148**, doi:10.1242/dev.199544 (2021).

25   Takeda, N. *et al.* Identification of jellyfish neuropeptides that act directly as
oocyte maturation-inducing hormones. *Development* **145**,
doi:10.1242/dev.156786 (2018).

26   Stierwald, M., Yanze, N., Bamert, R. P., Kammermeier, L. & Schmid, V. The Sine
oculis/Six class family of homeobox genes in jellyfish with and without eyes:
development and eye regeneration. *Dev Biol* **274**, 70-81,
doi:10.1016/j.ydbio.2004.06.018 (2004).

27   Suga, H., Schmid, V. & Gehring, W. J. Evolution and functional diversity of
jellyfish opsins. *Curr Biol* **18**, 51-55, doi:10.1016/j.cub.2007.11.059 (2008).

28   Suga, H. *et al.* Flexibly deployed Pax genes in eye development at the early
evolution of animals demonstrated by studies on a hydrozoan jellyfish. *Proc Natl*
*Acad Sci U S A* **107**, 14263-14268, doi:10.1073/pnas.1008389107 (2010).

29   Graziussi, D. F., Suga, H., Schmid, V. & Gehring, W. J. The "eyes absent" (eya)
gene in the eye-bearing hydrozoan jellyfish Cladonema radiatum: conservation of
the retinal determination network. *J Exp Zool B Mol Dev Evol* **318**, 257-267,
doi:10.1002/jez.b.22442 (2012).

30   Meister, G. & Tuschl, T. Mechanisms of gene silencing by double-stranded RNA.
*Nature* **431**, 343-349, doi:10.1038/nature02873 (2004).

31   Rao, D. D., Vorhies, J. S., Senzer, N. & Nemunaitis, J. siRNA vs. shRNA:
similarities and differences. *Adv Drug Deliv Rev* **61**, 746-759,
doi:10.1016/j.addr.2009.04.004 (2009).

32   Moran, Y. *et al.* Cnidarian microRNAs frequently regulate targets by cleavage.
*Genome Res* **24**, 651-663, doi:10.1101/gr.162503.113 (2014).

33   Watanabe, H. *et al.* Nodal signalling determines biradial asymmetry in Hydra.
*Nature* **515**, 112-115, doi:10.1038/nature13666 (2014).

34   Pankow, S. & Bamberger, C. The p53 tumor suppressor-like protein nvp63
mediates selective germ cell death in the sea anemone Nematostella vectensis.
*PLoS One* **2**, e782, doi:10.1371/journal.pone.0000782 (2007).

35   He, S. *et al.* An axial Hox code controls tissue segmentation and body patterning
in Nematostella vectensis. *Science* **361**, 1377-1380,
doi:10.1126/science.aar8384 (2018).
36   Karabulut, A., He, S., Chen, C. Y., McKinney, S. A. & Gibson, M. C.
Electroporation of short hairpin RNAs for rapid and efficient gene knockdown in
the starlet sea anemone, Nematostella vectensis. *Dev Biol* **448**, 7-15,
doi:10.1016/j.ydbio.2019.01.005 (2019).
37   DuBuc, T. Q. *et al.* Transcription factor AP2 controls cnidarian germ cell
induction. *Science* **367**, 757-762, doi:10.1126/science.aay6782 (2020).
38   Quiroga-Artigas, G., Duscher, A., Lundquist, K., Waletich, J. & Schnitzler, C. E.
Gene knockdown via electroporation of short hairpin RNAs in embryos of the
marine hydroid Hydractinia symbiolongicarpus. *Sci Rep* **10**, 12806,
doi:10.1038/s41598-020-69489-8 (2020).
39   Ha, M. & Kim, V. N. Regulation of microRNA biogenesis. *Nat Rev Mol Cell Biol*
**15**, 509-524, doi:10.1038/nrm3838 (2014).
40   Krishna, S. *et al.* Deep sequencing reveals unique small RNA repertoire that is
regulated during head regeneration in Hydra magnipapillata. *Nucleic Acids Res*
**41**, 599-616, doi:10.1093/nar/gks1020 (2013).
41   Moran, Y., Praher, D., Fredman, D. & Technau, U. The evolution of microRNA
pathway protein components in Cnidaria. *Mol Biol Evol* **30**, 2541-2552,
doi:10.1093/molbev/mst159 (2013).
42   Murchison, E. P., Partridge, J. F., Tam, O. H., Cheloufi, S. & Hannon, G. J.
Characterization of Dicer-deficient murine embryonic stem cells. *Proc Natl Acad*
*Sci U S A* **102**, 12135-12140, doi:10.1073/pnas.0505479102 (2005).
43   Kok, K. H., Ng, M. H., Ching, Y. P. & Jin, D. Y. Human TRBP and PACT directly
interact with each other and associate with dicer to facilitate the production of
small interfering RNA. *J Biol Chem* **282**, 17649-17657,
doi:10.1074/jbc.M611768200 (2007).
44   Watanabe, H. *et al.* Sequential actions of beta-catenin and Bmp pattern the oral
nerve net in Nematostella vectensis. *Nat Commun* **5**, 5536,
doi:10.1038/ncomms6536 (2014).

45    Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis.
*Nat Methods* **9**, 676-682, doi:10.1038/nmeth.2019 (2012).

46    Thompson, J. D., Higgins, D. G. & Gibson, T. J. CLUSTAL W: improving the
sensitivity of progressive multiple sequence alignment through sequence
weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids*
*Res* **22**, 4673-4680, doi:10.1093/nar/22.22.4673 (1994).

47    Waterhouse, A. M., Procter, J. B., Martin, D. M., Clamp, M. & Barton, G. J.
Jalview Version 2--a multiple sequence alignment editor and analysis workbench.
*Bioinformatics* **25**, 1189-1191, doi:10.1093/bioinformatics/btp033 (2009).

48    Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular
Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol* **35**,
1547-1549, doi:10.1093/molbev/msy096 (2018).

49    Deguchi, R., Kondoh, E. & Itoh, J. Spatiotemporal characteristics and
mechanisms of intracellular Ca(2+) increases at fertilization in eggs of jellyfish
(Phylum Cnidaria, Class Hydrozoa). *Dev Biol* **279**, 291-307,
doi:10.1016/j.ydbio.2004.11.036 (2005).

50    Scholz, C. B. & Technau, U. The ancestral role of Brachyury: expression of
NemBra1 in the basal cnidarian Nematostella vectensis (Anthozoa). *Dev Genes*
*Evol* **212**, 563-570, doi:10.1007/s00427-002-0272-x (2003).

51    Servetnick, M. D. *et al.* Cas9-mediated excision of Nematostella brachyury
disrupts endoderm development, pharynx formation and oral-aboral patterning.
*Development* **144**, 2951-2960, doi:10.1242/dev.145839 (2017).

52    Fourrage, C., Swann, K., Gonzalez Garcia, J. R., Campbell, A. K. & Houliston, E.
An endogenous green fluorescent protein-photoprotein pair in Clytia
hemisphaerica eggs shows co-targeting to mitochondria and efficient
bioluminescence energy transfer. *Open Biol* **4**, 130206, doi:10.1098/rsob.130206
(2014).

53    Guder, C. *et al.* The Wnt code: cnidarians signal the way. *Oncogene* **25**, 7450-
7460, doi:10.1038/sj.onc.1210052 (2006).

54    Lee, P. N., Pang, K., Matus, D. Q. & Martindale, M. Q. A WNT of things to come:
evolution of Wnt signaling and polarity in cnidarians. *Semin Cell Dev Biol* **17**,
157-167, doi:10.1016/j.semcdb.2006.05.002 (2006).

55   Kraus, Y., Aman, A., Technau, U. & Genikhovich, G. Pre-bilaterian origin of the
blastoporal axial organizer. *Nat Commun* **7**, 11694, doi:10.1038/ncomms11694
(2016).

56   Plickert, G., Jacoby, V., Frank, U., Muller, W. A. & Mokady, O. Wnt signaling in
hydroid development: formation of the primary body axis in embryogenesis and
its subsequent patterning. *Dev Biol* **298**, 368-378,
doi:10.1016/j.ydbio.2006.06.043 (2006).

57   Bartel, D. P. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*
**116**, 281-297, doi:10.1016/s0092-8674(04)00045-5 (2004).

58   DuBuc, T. Q., Stephenson, T. B., Rock, A. Q. & Martindale, M. Q. Hox and Wnt
pattern the primary body axis of an anthozoan cnidarian before gastrulation. *Nat*
*Commun* **9**, 2007, doi:10.1038/s41467-018-04184-x (2018).

59   Yasuoka, Y., Shinzato, C. & Satoh, N. The Mesoderm-Forming Gene brachyury
Regulates Ectoderm-Endoderm Demarcation in the Coral Acropora digitifera.
*Curr Biol* **26**, 2885-2892, doi:10.1016/j.cub.2016.08.011 (2016).

60   Duffy, D. J., Plickert, G., Kuenzel, T., Tilmann, W. & Frank, U. Wnt signaling
promotes oral but suppresses aboral structures in Hydractinia metamorphosis
and regeneration. *Development* **137**, 3057-3066, doi:10.1242/dev.046631
(2010).

61   Krasovec, G., Pottin, K., Rosello, M., Queinnec, E. & Chambon, J. P. Apoptosis
and cell proliferation during metamorphosis of the planula larva of Clytia
hemisphaerica (Hydrozoa, Cnidaria). *Dev Dyn* **250**, 1739-1758,
doi:10.1002/dvdy.376 (2021).

62   Momose, T., Kraus, Y. & Houliston, E. A conserved function for Strabismus in
establishing planar cell polarity in the ciliated ectoderm during cnidarian larval
development. *Development* **139**, 4374-4382, doi:10.1242/dev.084251 (2012).

63   Kumburegama, S., Wijesena, N., Xu, R. & Wikramanayake, A. H. Strabismus-
mediated primary archenteron invagination is uncoupled from Wnt/beta-catenin-
dependent endoderm cell fate specification in Nematostella vectensis (Anthozoa,
Cnidaria): Implications for the evolution of gastrulation. *Evodevo* **2**, 2,
doi:10.1186/2041-9139-2-2 (2011).

64   Nath, R. D. *et al.* The Jellyfish Cassiopea Exhibits a Sleep-like State. *Curr Biol*
**27**, 2984-2990 e2983, doi:10.1016/j.cub.2017.08.014 (2017).

65   Kanaya, H. J. *et al.* A sleep-like state in Hydra unravels conserved sleep
mechanisms during the evolutionary development of the central nervous system.
*Sci Adv* **6**, doi:10.1126/sciadv.abb9415 (2020).

66   Fuchs, B. *et al.* Regulation of polyp-to-jellyfish transition in Aurelia aurita. *Curr*
*Biol* **24**, 263-273, doi:10.1016/j.cub.2013.12.003 (2014).

67   Lommel, M. *et al.* Hydra Mesoglea Proteome Identifies Thrombospondin as a
Conserved Component Active in Head Organizer Restriction. *Sci Rep* **8**, 11753,
doi:10.1038/s41598-018-30035-2 (2018).

## Acknowledgement

We thank R. Deguchi (Miyagi Univ. Education, Japan) for sharing *Cladonema pacificum*
and EMBRC France for sharing *Clytia hemisphaerica*. We thank T. Momose (Sorbonne
University, CNRS, France) for helpful discussion. We thank I. Nagai, H. Nakatani, and
A. Sasaki for technical assistance; and A. Dahal, A. Tanimoto, and J. Higuchi for
*Nematostella* culture. This work was supported by JST grant number JPMJCR1852 to
E.K., AMED under grant number JP21gm6110025 to Y.N., and the JSPS KAKENHI
grant numbers JP21H05255 to E.K., and JP17H06332, JP19K22550, JP22H02762 to
Y.N.

## Author contributions

T.M.-O. conceptualized and designed the project, performed experiments using jellyfish,
analyzed data, prepared figures, and wrote the manuscript. S.F. performed experiments
and analyzed data. R.N. performed experiments using *Nematostella*. H.W.
conceptualized and designed the project. E.K. contributed reagents. Y.N.
conceptualized and designed the project, prepared figures, and wrote the manuscript.
All authors approved the final manuscript.

## Data availability

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request and nucleotide sequences are available in the GenBank/EMBL/DDBJ under the accession numbers (*CpWnt1*, LC720432; *CpWnt1b*, LC720433; *CpWnt2*, LC720434; *CpWnt3*, LC720435; *CpWnt5*, LC720436; *CpWnt6*, LC720437; *CpWnt8*, LC720438; *CpWntA*, LC720439; *CpBrachyury*, LC720440; *CpFoxQ2a*, LC720441; *F-actin capping protein subunit beta*, LC720442; *CpEF1alpha*, LC720443).

## Competing interests

The authors declare no competing interests.

## Figure legends

**Figure 1. The hydrozoan jellyfish *Clytia hemisphaerica* and *Cladonema pacificum*.** (**A**) Cladogram depicting the phylogenetic position of cnidarian jellyfish. As the sister group of Bilateria, the phylum Cnidaria is divided into two clades, Anthozoa and Medusozoa, which consists of four classes: Hydrozoa, Staurozoa, Scyphozoa, and Cubozoa. Hydrozoa includes polyp type animals without a medusa stage (e.g. *Hydra* and *Hydractinia*) and jellyfish that have both polyp and medusa stages (e.g. *Clytia* and *Cladonema*). (**B** and **C**) Photos of an adult medusa and the eggs of *Clytia hemisphaerica* (**B**) and *Cladonema pacificum* (**C**). Scale bars: 1 mm for medusae; 100 µm for eggs.

**Figure 2. The electroporation procedure for jellyfish eggs.** (**A**) Schematic of spawning under the 24 h light/dark cycle. *Clytia hemisphaerica* are maintained on a 13 h dark/11 h light cycle. After light stimuli, sperm spawning occurs within 60-90 min and egg spawning occurs within 90-120 min. *Cladonema pacificum* are maintained on a 23.5 h light/0.5 h dark cycle. After dark stimuli, sperm and egg spawning occurs within

25 min. (**B**) Fertilized eggs are resuspended in 15% Ficoll/artificial sea water to prevent

precipitation of eggs during electroporation. (**C**) Electroporation is performed with the

Bio-Rad Gene Pulser Xcell electroporation system and a cuvette with a 4 mm gap. (**D**)

Visualization of the Rhodamine-Dextran delivery into eggs by electroporation.

Rhodamine is shown in red. While little or no fluorescence was observed in eggs

without electroporation (No EP), under the condition of a single 50 V pulse for 25 ms,

rhodamine signals were detected in both *Clytia and Cladonema* eggs without cell

damage. For clarity, egg outlines are indicated with dashed lines for eggs without

electroporation. Scale bars: 200 μm.

**Figure 3**. **Phenotypes of *GFP1* knockdown with siRNA in *Clytia* fertilized eggs.** (**A**)

Schematic of the siRNA electroporation procedure using fertilized eggs. Fertilized eggs

in 15% Ficoll/artificial sea water mixed with siRNA are incubated for 5 min, and

electroporation is conducted after transferring into a cuvette.  (**B** and **C**) Phenotypes of

*Clytia* 2-day planula after *CheGFP1* siRNA knockdown. (**D**) Boxplots showing the GFP

fluorescence intensity mean values in 2-day planula after electroporation of 0 and 600

ng/μl siRNAs targeting *CheGFP1*. Center lines indicate the medians; x's denote the

means; box limits represent the 25th and 75th percentiles; whiskers show the maximum

and minimum values. 0 ng/μl, n=24; 600 ng/μl, n=81. ***p<0.001. (**E**-**H**) Phenotypes of

3-day planula after siRNA targeting *CheGFP1* electroporation (0, 150, 300, and 600

ng/μl si*CheGFP1*). (**I** and **J**) For quantification of green fluorescence, we classified

phenotypes into two categories: planula with ring-like green fluorescence as "wild-type,"

and planula with reduced green fluorescence as "reduced". (**K**) Quantification of planula

larvae positive for *CheGFP1* (wild-type). Bar plots show the percentage of wild-type

phenotype after *CheGFP1* knockdown. Number of examined planula: 0 ng/μl, *n*=82; 150

ng/μl, *n*=64; 300 ng/μl, *n*=58; 600 ng/μl, *n*=114. Error bars: maximum and minimum

values. Experiments were repeated three times. p-values: 150 ng/μl, p=0.064505; 300

ng/μl, p=0.001437; 600 ng/μl, p=0.00066. **p<0.01, ***p<0.001. (**L**) Quantification of

*CheGFP1* mRNA expression levels in 3-day planula by RT-qPCR. *CheEF1alpha* was

used as an internal control. Bar heights represent mean values of at least three

independent experiments. *CheGFP1* expression levels are standardized relative to the

control (0 ng/µl) condition. Error bars: standard deviation. Experiments were performed
in triplicated and repeated at least three times. p-values: 150 ng/µl, p=0.829361; 300
ng/µl, p=0.08588; 600 ng/µl, p=0.037359. *p<0.05. Scale bars: 200 µm.
944
945
**Figure 4. Phenotypes of *Wnt3* knockdown with siRNA in *Clytia* fertilized eggs.** (**A**
and **B**) Typical phenotypes of *Clytia* 1-day planula. While morphologies of control
planula (0 ng/µl) have an elongated oval-shape (A), those of *CheWnt3* siRNA (400
ng/µl) are spherical in shape (**B**). (**C**) Schematic of ellipse approximation for planula
morphology and calculation of the aspect ratio. The aspect ratio was calculated by
dividing the long axis (a) by short axis (b). (**D**) Boxplots showing the aspect ratio of the
1-day planula. Center lines show the medians; x's denote the mean values; box limits
indicate the 25th and 75th percentiles; whiskers show maximum and minimum values;
inner points and outliers are represented by circles. Number of examined planula: 0
ng/µl, *n*=127; 400 ng/µl. *n*=25, ***p<0.001. (**E-H**) Phenotypes of 1-day planula after
siRNA targeting *CheWnt3* electroporation (0, 100, 200, and 400 ng/µl si*CheWnt3*). (**I**
and **J**) For morphology quantification, we classified planula phenotypes into two
categories: elongated oval shape as "normal" and spherical (circle) shape as
"spherical". (**K**) Bar plots show the percentage of normal phenotypes across four
different siRNA doses as well as planulae treated with shRNA and siNC, a siRNA
universal negative control. Percentages are the mean value and error bars indicate
standard deviation. Numbers of examined planula: 0 ng/µl, *n*=127; 100 ng/µl, *n*=75; 200
ng/µl, *n*=86; 400 ng/µl, *n*=95. shRNA 400 ng/µl, *n*=288; siNC 400 ng/µl, *n*=45.
Experiments were repeated three times. Error bars: maximum and minimum values. p-
values: 100 ng/µl, p=0.000599; 200 ng/µl, p=0.000438; 400 ng/µl, p=0.000402; 400
ng/µl of shRNA, p=0.031928; 400 ng/µl of siNC, p=0.283359. *p<0.05, ***p<0.001. (**L**)
Quantification of *CheWnt3* mRNA levels of 1-day planula by RT-qPCR. *CheEF1alpha*
was used as an internal control. Bar heights represent the mean value. *CheWnt3*
expression levels are standardized relative to the control (0 ng/µl). Error bars: standard
deviation. p-values: 100 ng/µl, p= 0.00799; 200 ng/µl, p= 0.000754; 400 ng/µl, p=
0.000733; 400 ng/µl of shRNA, p= 0.095403; 400 ng/µl of siNC, p= 0.900264. **p<0.01,

***p<0.001. n.s., not significant. Experiments were performed in triplicate and repeated

at least three times. Scale bars: (A, B, E-H) 400 μm, (I, J) 100 μm.

**Figure 5. Phenotypes of *Wnt3* knockdown with siRNA in *Cladonema* fertilized**

**eggs.** (**A**) Typical morphology of *Cladonema* egg and planula. The wild-type

*Cladonema* planula larvae exhibit a slight oval shape. To deliver siRNAs targeting

*CpWnt3,* electroporation was performed in fertilized eggs (1-cell stage), and phenotypes

were confirmed at planula in one day. (**B** and **C**) Phenotypes of 1-day planula after

*CpWnt3* siRNA knockdown. (**D**) Schematic of ellipse approximation for planula

morphology and calculation of the aspect ratio. The aspect ratio was calculated by

dividing the long axis (a) by short axis (b). (**E**) Boxplots showing the aspect ratio of the

1-day planula after si*CpWnt3* electroporation (0 and 300 ng/μl). Center lines show the

medians; x's denote the mean values; box limits indicate the 25th and 75th percentiles;

whiskers show maximum and minimum values; inner points and outliers are

represented by circles. Number of examined planulae: 0 ng/μl, *n*=22; 300 ng/μl, *n*=50.

p= 0.658303. n.s., not significant. (**F**) Quantification of *CpWnt3* mRNA levels in 1-day

planula by RT-qPCR. *CpEF1alpha* was used as an internal control. Bar heights

represent the mean value. Error bars indicate standard deviation. *CpWnt3* expression

levels are standardized relative to the control (0 ng/μl). Experiments were performed in

triplicate and repeated three times. p-values: 150 ng/μl, p= 0.376498; 300 ng/μl, p=

0.073956. (**G, H**) Representative *in situ* hybridization (ISH) image of oral *CpBra*

expression in 1-day planula for control (G: 0 ng/μl si*CpWnt3*) and *CpWnt3* siRNA (H:

300 ng/μl si*CpWnt3*). (**I**) Quantification of *CpBra* expression phenotypes based on ISH

images. Stacked bar plots show the percentage of 1-day planulae in each phenotypic

class. *CpBra* expression patterns are categorized into three phenotypes (strong, weak

and none). 0 ng/μl, *n*=74; 300 ng/μl, *n*=58. Scale bars: (B, C) 200 μm, (G, H) 50 μm.

# Figure 3

Masuda-Ozawa et al.

**A**

Fertilized eggs + siRNA | 5 min | Electroporation

si*CheGFP1* electroporation
2 day planula

**B** 0 ng/µl

**C** 600 ng/µl

**D**

GFP intensity mean value (A.U. x10³)

***

0    600  (ng/µl)

si*CheGFP1* electroporation/3 day planula

**E** 0 (ng/µl)

**F** 150

**G** 300

**H** 600

**I** Wild type — No EP

**J** Reduced — si*GFP1*

**K**

Wild type (%)

***
**

0   150   300   600
(ng/µl)

**L**

*GFP1* relative gene expression

*

0   150   300   600
(ng/µl)

**Figure 5**  Masuda-Ozawa et al.