1

siRNA-mediated gene knockdown via electroporation in hydrozoan jellyfish embryos

[1] Tokiha Masuda-Ozawa, [2] Sosuke Fujita**,** [3] Ryotaro Nakamura, [3] Hiroshi Watanabe, [2] Erina Kuranaga, and [1,4] Yu-ichiro Nakajima*****

1. Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, Sendai, Japan
2. Graduate School of Life Sciences, Tohoku University, Sendai, Japan
3. Evolutionary Neurobiology Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan
4. Graduate School of Pharmaceutical Sciences, The University of Tokyo

*Author for correspondence e-mail: nakaji97@g.ecc.u-tokyo.ac.jp
Tel: +81-3-5841-4863

Running title: **siRNA gene knockdown in jellyfish embryos**
Main text: 6487 words (Abstract: 258 words)
Figures: 5   **Supplementary information** (Figures: 7; Table: 5)
Keywords: siRNA, electroporation, cnidarian, hydrozoan, jellyfish, *Clytia, Cladonema*

Abstract

As the sister group to bilaterians, cnidarians stand in a unique phylogenetic position that provides insight into evolutionary aspects of animal development, physiology, and behavior. While cnidarians are classified into two types, sessile polyps and free-swimming medusae, most studies at the cellular and molecular levels have been conducted on representative polyp-type cnidarians and have focused on establishing techniques of genetic manipulation. Recently, gene knockdown by delivery of short hairpin RNAs into eggs via electroporation has been introduced in two polyp-type cnidarians, *Nematostella vectensis* and *Hydractinia symbiolongicarpus*, enabling systematic loss-of-function experiments. By contrast, current methods of genetic manipulation for most medusa-type cnidarians, or jellyfish, are quite limited, except for *Clytia hemisphaerica*, and reliable techniques are required to interrogate function of specific genes in different jellyfish species. Here, we present a method to knock down target genes by delivering small interfering RNA (siRNA) into fertilized eggs via electroporation, using the hydrozoan jellyfish, *Clytia hemisphaerica* and *Cladonema paciificum*. We show that siRNAs targeting endogenous *GFP1* and *Wnt3* in *Clytia* efficiently knock down gene expression and result in known planula phenotypes: loss of green fluorescence and defects in axial patterning, respectively. We also successfully knock down endogenous *Wnt3* in *Cladonema* by siRNA electroporation, which circumvents the technical difficulty of microinjecting small eggs. *Wnt3* knockdown in *Cladonema* causes gene expression changes in axial markers, suggesting a conserved Wnt/β-catenin-mediated pathway that controls axial polarity during embryogenesis. Our gene-targeting siRNA electroporation method is applicable to other animals, including and beyond jellyfish species, and will facilitate the investigation and understanding of myriad aspects of animal development.

Introduction

The phylum Cnidaria is the sister group to Bilateria, having separated from their common ancestor over 500 million years ago. Cnidarians have diversified their morphologies across an array forms, including corals, sea anemones, hydroids, and jellyfish, all of which are divided into two clades: Anthozoa and Medusozoa. The two differ in that Anthozoa includes only sessile polyp-type animals, while Medusozoa (Hydrozoa, Staurozoa, Scyphozoa, and Cubozoa) contains two forms: polyp and medusa, commonly known as jellyfish (Fig. 1A)[1,2]. The life cycle of most, though not all, jellyfish consists of five forms: gametes, fertilized eggs, planulae, polyps, and medusae[3]. Fertilized eggs undergo embryogenesis to become planula larvae, which metamorphose into sessile polyps. While vegetatively-growing polyps give rise to free-swimming medusae through the process of budding or strobilization, medusae sexually reproduce by releasing gametes. Due to their unique phylogenetic position, studies using cnidarians have provided evolutionary insight into development, regeneration, and behaviors in multicellular animals[1,4,5]. Despite divergent morphologies and lifestyles among cnidarians, to date, the molecular and cellular understanding of cnidarians has been acquired primarily using polyp-type animals such as the anthozoan *Nematostella vectensis* and the hydrozoan *Hydra* and *Hydractinia*. By contrast, jellyfish biology remains largely unestablished.

Among jellyfish species, the hydrozoan *Clytia hemisphaerica* is the best-studied laboratory model[6,7]. The well-established methodology for maintaining and propagating animals ensures a reliable source for daily experiments[8]. Transparent and small-sized medusae make whole-mount visualization achievable, and relatively large-sized eggs (~180 μm diameter) enable different manipulations via microinjection (Fig. 1B). Indeed, gene silencing by morpholino oligos and mRNA microinjection as well as gene knockout via CRISPR/Cas9 allow for the investigation of functions of genes of interest[9-13]. Furthermore, the *Clytia* genome assembly is complete, and transcriptome profiles are available for various stages and tissues, even at the single-cell level, which creates the foundation for research at the molecular level[12,14-18]. These resources and techniques, together with recently-established transgenesis[19], have accelerated research into numerous facets of distinct *Clytia* life stages. While initial studies focused on

embryogenesis, current research topics using *Clytia* include gametogenesis,

regeneration, and behavior exhibited by adult medusa [9,10,12,17,19,20].

Compared to the established model jellyfish *Clytia*, the hydrozoan *Cladonema*

pacificum is an emerging model that has recently been utilized for various studies in

development, regeneration, and physiology[21]. The ease of rearing all stages of

Cladonema without a filtration system or large water tank, along with their high

spawning rate, enable easy lab maintenance. Small-sized medusae with branched

tentacles allow for investigations of body size control and tentacle morphogenesis (Fig.

1C)[22-24]. *Cladonema* gametogenesis is regulated by a light-dark cycle as in *Clytia*, and

recent work has identified the neuropeptides involved in oocyte maturation[25].

Furthermore, *Cladonema* medusae possess eyes with a complex structure that includes

lenses (ocelli) in their tentacle bulbs, and studies using the closely related species

Cladonema radiatum have identified conserved light-sensitive opsins and regulators of

eye formation (*Pax*, *Six*, and *Eya*)[25-29], providing a model for the evolutionary

developmental biology of photoreceptor organs. Despite these attractive features of

Cladonema as a laboratory animal, no genome assembly or transcriptome is currently

available, and genetic manipulation techniques are only just being developed. One

major technical issue in manipulating *Cladonema* is their small eggs—with a diameter of

approximately 60 µm (Fig. 1C), regular microinjection is quite difficult. Establishing

genetic manipulations is required to facilitate the in-depth investigations needed to

further understand the biology of *Cladonema*, and an alternative method that eliminates

the need for microinjection would greatly facilitate that objective.

RNA interference (RNAi), the phenomenon of double stranded RNA (dsRNA)

mediated silencing of target genes, has been widely exploited in living organisms to

analyze gene function[30], primarily using chemically- or *in vitro*-synthesized double-

stranded small interfering RNAs (siRNAs) or vector-based short hairpin RNAs

(shRNAs)[31]. In cnidarians, siRNA-mediated gene silencing was initially applied to

Nematostella and *Hydra* polyps via soaking or electroporation[32-34]. More recently, gene

knockdown via shRNA microinjection or electroporation into eggs has been utilized in

studies with *Nematostella* and *Hydractinia* to show efficient reduction in gene

expression and associated phenotypes in early developmental stages[35-38]. In particular,

shRNA delivery via electroporation, which does not require the rigors of microinjection,

allows for the experimental gene knockdown of large numbers of individuals

simultaneously[36,38], opening up the possibility of manipulating genes in different aquatic

animals, even those that produce very small eggs like *Cladonema*.

The mechanism of RNAi-mediated gene repression after introducing foreign

dsRNA into cells involves the microRNA (miRNA) pathway, an endogenous gene

repression machinery conserved in both animals and plants[39]. In cnidarians, the

presence of the miRNA pathway has been demonstrated in *Nematostella* and

Hydra[32,40,41]. However, little is known about the endogenous RNAi pathway, especially

the presence and roles of miRNAs and the miRNA-related genes, in jellyfish such as

Clytia and *Cladonema*. Furthermore, in mammalian cultured cells, siRNAs can induce

gene repression independent of endogenous dsRNA processing factors, while shRNAs

cannot[42,43]. On this basis, we selected siRNA instead of shRNA for gene knockdown to

avoid the potential pitfall of dsRNA processing in jellyfish.

Here we report a gene knockdown method for jellyfish embryos with siRNA via

electroporation. Using two hydrozoan species, *Clytia hemisphaerica* and *Cladonema*

pacificum, we demonstrate that siRNA delivery into fertilized eggs effectively reduces

the expression of endogenous genes. We also confirm the known loss-of-function

phenotypes in *Clytia* after knocking down *GFP1* or *Wnt3*, which are induced in a dose-

dependent manner with siRNA. We further show that knockdown of *Wnt3* in *Cladonema*

embryos results in the reduction of gene expression of the oral marker *Brachyury* in

planula, implicating the Wnt/β-catenin pathway in the control of oral-aboral patterning.

Overall, our siRNA-mediated knockdown approach allows for the manipulation of a

large number of embryos through electroporation and enables functional analysis of

early development, providing a new experimental platform applicable to different jellyfish

species and other marine invertebrates.

Materials and Methods

Animal culture and spawning induction

Clytia hemisphaerica (Z11, Z4B as females and Z4C2, Z23 as males) and *Cladonema*

pacificum (6W as females and UN2 as males) were used for this research.

Clytia hemisphaerica were cultured using a previously reported method[8] with a

few modifications. Artificial sea water (ASW) was prepared using 220 g SEA LIFE

(Marin Tech) per 5L MilliQ water (Merk Millipore) with antibiotics (40 units/ml of penicillin

and 40 µg/ml of streptomycin). Medusae, embryos, and planula larvae were maintained

at 20ºC. Medusae were fed daily with Vietnamese brine shrimp (A&A Marine LLC, Elk

Rapids, MI, USA). Spawning timing was controlled by a 13 hours (h) dark /11 h light

cycle, and spawning was induced by light. Male and female medusae were transferred

into V-7 cups (AS ONE) before spawning (60 min for male and 90 min for female after

light stimulation).

Cladonema pacificum were cultured as previously described[23]. *Cladonema*

medusae were maintained at 22ºC in ASW, which was prepared using SEA LIFE (Marin

Tech) dissolved in tap water with chlorine neutralizer (Coroline off, GEX Co. ltd) (24

p.p.t) and antibiotics (40 units/ml penicillin and 40 µg/ml of streptomycin). Spawning

timing was controlled by a 30 min dark/23.5 h light cycle, and spawning of male and

female gametes was induced by dark stimulation. Before dark stimulation, adult females

and males were separately transferred into V7 cups (AS ONE) and 60 mm dishes (BD),

respectively.

Nematostella vectensis were cultured as previously described[44], with a few

modifications. Briefly, adult animals were maintained in brackish water at a salinity one-

third of artificial seawater (35 g/l, pH 7.5-8.0, SEA LIFE, Marin Tech) and fed with

freshly hatched artemia twice per week. Spawning induction was performed at 26°C

under light for at least 11 h.

siRNA and shRNA

siRNA sequences (19 mer RNA + 2 mer DNA) for *CheGFP1*, *CheWnt3*, and *CpWnt3*

were designed based on their CDS sequences by Nippon Gene Co., Ltd, and siRNA

duplexes were synthesized by manufacturers (Nippon Gene Co., Ltd and Sigma-

Aldrich, Merk). Lyophilized siRNA was resuspended in RNase free water to a final

concentration of 6 µg/µl as stock solution. The siRNA stock solution was diluted with

RNase free water to a total volume of 10 µl and then added to 90 µl of fertilized eggs in

15% Ficoll ASW just prior to electroporation.

shRNAs were synthesized as described in previous reports[36,38]. Briefly, shRNAs

were synthesized by *in vitro* transcription (IVT) from double stranded DNA templates

using the AmpliScribeTM T7-flashTM transcription kit (Lucigen, Inc.) and were purified

using Direct-Zol TM RNA Miniprep kits (Zymo Research, R2070). Concentrations of

shRNA was measured with a NanoDrop One (Thermo fisher).

Collection of gametes and electroporation for *Clytia* and *Cladonema* fertilized

eggs

For the preparation of gametes, sexually-mature *Clytia* medusae were treated following

a previously reported method[8], partially modified to fit our experimental equipment.

Adult male and female medusae were maintained on a 13 h dark/11 h light cycle, and

light stimuli induced gametogenesis in 60 min for sperm and 90 min for eggs (Fig. 2A).

For *Cladonema pacificum*, sexually-mature medusae were maintained on a 30 min

dark/23.5 h light cycle. Dark stimulation induced gametogenesis in 25 min for both

sexes.

Unfertilized *Clytia* and *Cladonema* eggs were collected and resuspended in 15%

Ficoll (Nacalai tesque, Japan) in ASW to prevent the eggs settling from at the bottom of

the microtube or sticking to the microtube surface, keeping the egg solution

homogeneous (Fig. 2B)[36,38]. Sperm water was added into the egg solution, and the

eggs and sperm were incubated for 5 min at room temperature (20-22ºC) until

fertilization occurred. The sperm water was then removed from the egg solution, and

fertilized eggs were resuspended in 15% Ficoll in ASW to adjust the total volume

needed for the number of electroporations. For each electroporation, 200-400 of *Clytia*

fertilized eggs or 200-700 of *Cladonema* fertilized eggs were prepared in 90 µl of 15%

Ficoll in ASW. 10 µl of siRNA/shRNA solution that was adjusted to the proper

concentration in RNase free water was added to the fertilized egg solution.

The total 100 µl of fertilized egg and RNA mixture was carefully transferred into a

4 mm cuvette using a 200 µl pipet tip and placed into the shockpod cuvette chamber

connected to the Gene Pulser Xcell (Bio-Rad). In most experiments, the Gene Pulser

Xcell was used for pulsing with a square wave voltage of 50 V and a single pulse

duration of 25 ms, except when optimizing electroporation conditions (Supplementary

Figs 1-2). After electroporation, fertilized eggs were transferred to a dish, and incubated

for 10 min. The 15% Ficoll solution was removed and replaced with new ASW, and the

samples were incubated at 20-22ºC.

Collection of egg sacs, dejellying, and electroporation of *Nematostella* fertilized

eggs

The collection of egg sacs, dejellying, and electroporation steps were performed as

previously described[36] with several modifications. Briefly, unfertilized egg sacs were

incubated in culture media with sperm at 20-25ºC for 30 min. The fertilized egg sacs

were then dejellied in freshly prepared L-Cysteine solution (0.04 g/ml, Nacalai tesque, in

brackish water; pH adjusted by 5N NaOH at 7.5-8.0) on a shaker for 10 min. The

dejellied eggs were rinsed with brackish water before they were suspended in Ficoll

solution (Sigma-Aldrich, in brackish water) at a final concentration of 15%. The

suspended eggs and siRNA (5 µg/µl stock solution, in MilliQ) were transferred to a 4

mm cuvette (Bio-Rad) with a final volume of 200 µl and were gently mixed. The Bio-Rad

Gene Pulser Xcell was used for pulsing, with a square wave voltage of 50 V and a pulse

duration of 25 ms. The samples were transferred to a 100 mm dish and maintained in

brackish water for 3 days.

Imaging

GFP and Rhodamine fluorescent imaging as well as bright field imaging, including *in*

situ hybridization imaging, were taken with an Axio Zoom V16 (ZEISS), and image

processing was done with ZEN 3.4 blue edition (ZEISS) and Fiji/ImageJ software[45].

Intensity of mean values of green fluorescence was measured using the average

brightness (pixel value) of the pixels. To prevent planula movement, we added 0.5 M

sodium azide (final concentration was less than 0.25M).

Morphological quantification of planula larvae

After *Wnt3* knockdown by siRNA electroporation of *Clytia* or *Cladonema*, we

approximated the planula body shape into a two dimensional ellipse, measured both the

long and short axes (Figs. 4C and 5D), and calculated the aspect ratio of those values

using Fiji/ImageJ software (Figs, 4D, 5E).

RT-qPCR

For *Clytia* and *Cladonema* samples, total RNA was extracted with RNeasy Mini or Micro

kits (Qiagen). Lysate was treated with DNaseI (Qiagen) for 15 min at room temperature

(RT). cDNA was synthesized with the PrimeScript II 1st strand synthesis kit (Takara bio).

RT-qPCR was performed with CFX connect (Bio-Rad) using iTaqTM universal SYBER

Green Supermix (Bio-Rad) or the QuantStudio 6 Flex Real-Time PCR System (Thermo

Fisher) using TB Green Premix Ex TaqII (Tli RNaseH Plus) (Takara, RR820). Gene

expression was normalized to the housekeeping gene, *EF1alpha* or *F-actin capping*

protein subunit beta, and the delta-delta-ct method was used for quantification (CFX

maestro software, Bio-Rad or QuantStudio 6 Flex Real-Time PCR System software,

Thermo Fisher).

For *Nematostella* samples, total RNA was extracted using the RNeasy Mini Kit

and RNase-Free DNase Set (Qiagen). cDNA synthesis was conducted using the

SuperScript IV First-Strand Synthesis System (Thermo Fisher). RT-qPCR was

performed with the StepOnePlusTM Real-Time PCR System (Applied Biosystems,

Thermo Fisher) using PowerUP SYBR Green Master Mix (Thermo Fisher). Gene

expression was normalized to the housekeeping gene, *NvEf1alpha,* and the delta-delta-

ct method was used for quantification (StepOne™ and StepOnePlus™ Software v2.3).

In situ **hybridization**

Purified total RNA was reverse transcribed into cDNA by PrimeScript^{TM} 2 1st strand

cDNA Synthesis Kit (TaKaRa). The target gene fragments (*Brachyury*, *CpBra*) were

amplified from a cDNA library. The primer sets used for PCR cloning are as follows:

CpBra: 5'-GCTCCCATAAGATCCGGTCG -3'(forward) and 5'-

TTTGTCGCAGTCGAAGACCA-3'(reverse), or 5'-GAACGGTGATGGACAGGTCA-3'

(forward) and 5'-GGATTCCAAGGATTGGGCGT-3' (reverse). PCR products were sub-

cloned into the TAK101 vector (TOYOBO). The resulting plasmids were used for RNA

probe synthesis with digoxigenin (DIG) labeling mix (Roche), and T7 or T3 RNA

polymerase (Roche) was used, according to the insert direction. The detail for the probe

synthesis was referred to the published protocol[46].

Planula larvae were anesthetized in 50% 0.5 M NaN_3 in ASW for 5 min and fixed

overnight at 4°C with 4% paraformaldehyde (PFA) in ASW. The following whole mount

in situ hybridization was performed following the protocol in Hou et al.[24] with slight

modifications. Briefly, fixed samples were washed three times with PBS containing 0.1%

Tween-20 (PBST), followed by pre-hybridization in hybridization buffer (HB buffer:

5×SSC, 50% formamide, 0.1% Tween-20, 50 μg/ml tRNA, 50 μg/ml heparin) at 55°C for

2 h. Samples were hybridized with HB Buffer containing the antisense probes (final

probe concentration: 0.5-1 ng/μL in HB Buffer) at 55°C overnight. The samples were

washed each twice with wash buffer 1 (5x SSC, 50% formamide, 0.1% Tween-20),

wash buffer 2 (2x SSC, 50% formamide, 0.1% Tween-20) and 2x SSC. The samples

were then washed with 0.1% PBST and incubated in 1% blocking buffer (1% blocking

reagent [Roche, 11175041910] in Maleic acid) for 1 h. The samples were incubated with

alkaline phosphatase (AP)-conjugated anti-DIG antibodies (1:2000, Roche,

11175041910) in 1% blocking buffer for 4 h at room temperature. Colorimetric reactions

were performed by NBT/BCIP (Roche, 11175041910) in alkaline phosphatase buffer

(100 mM NaCl, 100 mM Tris-HCl (pH 9.5), 50 mM $MgCl_2$, 0.1% Tween-20) until the

signals were detected.

Sequence alignment and phylogenetic analysis

Amino acid sequence alignment of Wnt3 and Wnt family proteins were performed by

ClustalW[47]. GenBank accession numbers and protein names are listed in

Supplementary Table 3 and 4. The alignment data was visualized on Jalview software[48].

Maximum likelihood (ML) tree calculation was inferred on ClustalW with the PhyML

bootstrap method. Branch supports were computed out of 100 bootstrapped trees. The

tree visualization was created using MEGA-X software[49].

qPCR primers

Name	Sequence (5'->3')	
CpWnt3 qP FW	CAAATGTGGTCGATCAAACG	This study
CpWnt3 qP RV	TACGCCTTCTGCAACACTTG	This study
CpEF1-a_qPCR_F7	GGTCAATCTCGTTCCCTCCA	This study
CpEF1-a_qPCR_R7	TTTCCACCAGAGGTATCGGC	This study
CpBrachyury F	AAGGCGTATGTTTCCGGTCC	This study
CpBrachyury R	CAACGATGGTCTTCGACTGC	This study
CpFoxQ2a F	AGTTGGAGAAACAGCGTTCG	This study
CpFoxQ2a R	CCGTTCGCAAAGTCGTCAAA	This study
CpCapZbeta_qPCR_F	AAAGAAAGCTGGAGACGGTTCA	This study
CpCapZbeta_qPCR_R	GTAGTGGGCATTTCTTCCGC	This study
CheGFP1 qP F	TTGCTGTCCGAATAGTGCAG	Fourrage et al., 2014 Open Biol
CheGFP1 qP RV	GACAACTCCTCCTCCGAGTG	Fourrage et al., 2014 Open Biol
CheGFP1_2 qP F	ACAATCGCGTCACACTTAAAGG	This study
CheGFP1_2 qP RV	CGTTGTTTTCTTTGTCCGGC	This study
CheWnt3 qP F	ATCATGGCAGGTGGAAACTC	Leclère et al., 2012 Dev Biol
CheWnt3 qP RV	CCCCATTTCCAACCTTCTTC	Leclère et al., 2012 Dev Biol
CheEF1a qP F	TGCTGTTGTCCCAATCTCTG	Leclère et al., 2012 Dev Biol / Fourrage et al., 2014 Open biol
CheEF1a qP RV	AAGACGGAGTGGTTTGGATG	Leclère et al., 2012 Dev Biol / Fourrage et al., 2014 Open biol
NvEf1alpha FW	GGTTGCCTCTTCGCTTACCACT	This strudy
NvEf1alpha RV	CGTTCCTGGCTTTAGGACAC	This study
NvBra1_Fw_1032	CGGGCTCACACTCTCACTTA	This study
NvBra1_Rv1174	CTTGCGGTATGGTGTTCCAG	This study

siRNA design

Name	Sequence (5'->3')		ORF
siCheGFP1_1 sense	CCAUCAGCUUCGAAAAUGAdTdT	This study	HQ397706.1
antisense	UCAUUUUCGAAGCUGAUGGdTdT	This study	
siCheGFP1_2 sense	UGAUGGCGCUUAUAAAGUUdTdT	This study	HQ397706.1
antisense	AACUUUAUAAGCGCCAUCAdTdT	This study	
siCheGFP1_3 sense	CGGACUCAUUCUUCAAAAAdTdT	This study	HQ397706.1
antisense	UUUUUGAAGAAUGAGUCCGdTdT	This study	
siCheWnt3_1 sense	GCAUGUGACUGUAAAUUUAdTdT	This study	EU374721.1

antisense	UAAAUUUACAGUCACAUGCdTdT	This study	
siCheWnt3_2 sense	ACAGCUUGGUAAAUGUUAdTdT	This study	EU374721.1
antisense	UAACAUUUUACCAAGCUGUdTdT	This study	
siCpWnt3_1 sense	GUGGAACUGUAGUGUUUCAdTdT	This study	EU374721.1
antisense	UGAAACACUACAGUUCCACdTdT	This study	
siCpWnt3_2 sense	AGCGUGUGCUGAAGGUAAAdTdT	This study	EU374721.1
antisense	UUUACCUUCAGCACACGCUdTdT	This study	
siNvBra1_1 sense	GAAGAGAUCACGAGUCUAAdTdT	This study	AF540387.2
antisense	UUAGACUCGUGAUCUCUUCdTdT	This study	
siNegative_Control-1 for Nv sense	GCAACACGCAGAGTCGTAAdT	Same sequence as Karabulut et al, 2019 Dev Biol	
antisense	TTACGACTCTGCGTGTTGCdT	Same sequence as Karabulut et al, 2019 Dev Biol	

324 **shRNA template**

Name	Sequence (5'->3')	
CheWnt3 _SI1_F	TAATACGACTCACTATA GCATGTGACTGTAAATTTA TTCAAGAGA TAAATTTACAGTCACATGC TT	This study
CheWnt3 _SI1_R	AAGCATGTGACTGTAAATTTATCTCTTGAATAAATTTAC AGTCACATGCTATAGTGAGTCGTATTA	This study
CheWnt3 _SI2_F	TAATACGACTCACTATA ACAGCTTGGTAAAATGTTA TTCAAGAGA TAACATTTTACCAAGCTGT TT	This study
CheWnt3 _SI2_R	AAACAGCTTGGTAAAATGTTATCTCTTGAATAACATTTT ACCAAGCTGTTATAGTGAGTCGTATTA	This study

Graphs and statistical analysis

All graphs were prepared in Excel. To assess phenotypes and RT-qPCR statistical

significance, we used the percentage values and delta-delta-Ct values to perform two-

tailed Student's t tests, except in Figure 5F, where a one-tailed Student's t test was

used. For statistics related to fluorescence intensity mean value and aspect ratio,

Kolmogorov–Smirnov normality tests were used. All statistical tests were performed at

https://www.socscistatistics.com.

Results

Optimization of electroporation conditions

In both hydrozoan jellyfish *Clytia hemisphaerica* and *Cladonema pacificum*,
gametogenesis is regulated by light-dark transitions[6,50]. A previous report showed that
Clytia medusae release eggs within 110-120 min and sperm within 60-90 min upon light
illumination after 3-8 h of darkness[12]. *Cladonema pacificum*, which live along coastal
areas in Japan, exhibit two types of gametogenesis depending on habitat: the dark-light
transition (light stimulation type) and the light-dark transition (dark stimulation type),
respectively. Dark stimulation *Cladonema* medusae, which we used in this study,
release eggs or sperm within 30 min of dark stimulus after 20-24 h of constant exposure
to light[25,50]. By utilizing the characteristic gametogenesis of these jellyfish species, we
can control egg/sperm release to induce fertilization at any time of day. Indeed, we
routinely obtained *Clytia* sperm within 60 min and *Clytia* eggs within 90 min of light on a
13h dark/11h light cycle (Fig. 2A). Similarly, we collected gametes from *Cladonema*
within 25 min of dark stimulation on a 23.5h light/30 min dark cycle (Fig. 2A).

To determine optimal electroporation parameters, we first tested different
electroporation conditions and monitored the delivery efficiency of the red fluorescent
dye Rhodamine-Dextran into unfertilized *Clytia* and *Cladonema* eggs (Supplementary
Figs. 1-2). Given that the average molecular weight of siRNA (21-mer, ~13,300 Da) is
similar to the molecular size of Rhodamine-Dextran (10,000 Da), electroporation trials
with Rhodamine-Dextran can visually confirm that small molecules are incorporated into
jellyfish eggs, as previously shown in *Hydractinia* eggs[38]. The collected eggs were
suspended in 15% Ficoll artificial seawater to prevent precipitation and to make a
homogeneous solution (Fig. 2B)[36,38]. We performed electroporation trails for
Rhodamine-Dextran using a cuvette with a 4 mm gap and tested several conditions of
voltage (V) and pulse length time (milliseconds, ms) using a conventional
electroporation system (Fig. 2C). As previous work showed that increasing the number
of pulses does not increase the Dextran delivery rate but decreases the survival rate of
embryos[38], we fixed the number of pulses to one and incubated unfertilized eggs with 1
mg/ml Dextrane-Rhodamine solution. *Clytia* eggs have little autofluorescence in the red
channel, and without electroporation (No EP), red fluorescence was rarely detected in
unfertilized eggs (Fig. 2D and Supplementary Fig. 1A). We compared the eight different
electroporation conditions and found that the rhodamine fluorescence was detected in

Clytia eggs and correlated with the strength of electric voltage, as long as the voltage
was in the range of 50-100 V（Supplementary Figs. 1B-1D）. By contrast, when the
voltage exceeded 200 V, the eggs ruptured from physical damage, and cellular debris
was observed（Supplementary Figs, 1E-1G）.

We performed similar electroporation trials on unfertilized *Cladonema* eggs under
eight different conditions. While red autofluorescence was negligible in the *Cladonema*
eggs in the absence of electroporation (Fig. 2D and Supplementary Fig. 2A),
Rhodamine fluorescence was frequently detected with 50 V (Fig. 2D and
Supplementally Fig. 2E). In the more intense conditions beyond 100 V (Supplementally
Figs. 2F and 2G), the eggs were severely damaged. Based on these results, we
established the optimal electroporation conditions (50 V, 1 pulse, 25 ms) that provide
high frequency of Dextran-positive eggs and low levels of cell damage for both jellyfish
eggs (Fig. 2D). Of note, during the initial trials, we used unfertilized eggs, but we found
that the electroporation process severely affected the survival rate of subsequent
embryos. In the case of *Cladonema*, electroporation before fertilization severely
decreased the survival rate of fertilized eggs (7.04% at 1 h post fertilization) while
electroporation after fertilization did not influence the survival rate (No EP: 59.67%, With
EP: 60.15%) (Supplementary Table 1). This dissimilarity is likely due to the decrease of
fertilization rate after electroporation, and both elapsed time after spawning and physical
damage by electroporation can affect fertilization efficiency. Therefore, we decided to
conduct further electroporation experiments with fertilized eggs.

Gene knockdown with siRNA in *Nematostella vectensis*
Before attempting gene knockdown via siRNAs electroporation in fertilized jellyfish
eggs, we first used *Nematostella vectensis,* where egg electroporation methodology has
been established[36]. We selected *Nematostella Brachyury* (*NvBra*), a gene expressed in
the blastopore margin that functions in the early embryo[51,52], as a target. After
electroporation of siRNA targeting *NvBra* in fertilized eggs, we examined *NvBra*
expression in planula larvae at 3 days-post-fertilization (dpf) by RT-qPCR and found
that *NvBra* expression was drastically suppressed compared to no electroporation
controls and negative control siRNAs (Supplementary Fig. 3). We further evaluated

survival rates after electroporation and confirmed that siRNA electroporation does not

severely affect survival rates of 3 dpf planula in any of the tested conditions

(Supplementary Table 2). These results suggest that siRNA delivery into fertilized eggs

via electroporation is an effective gene knockdown method that is comparable to the

effect of shRNA delivery into unfertilized eggs via electroporation[36], supporting our

attempt to perform gene knockdown via siRNA electroporation using fertilized jellyfish

eggs.

Endogenous *GFP1* knockdown with siRNA in *Clytia hemisphaerica*

The hydrozoan jellyfish *Clytia* endogenously expresses green fluorescent proteins,

which are encoded by 14 genes grouped into 4 clusters (*CheGFP1-4*)[16]. The expression

pattern for each GFP changes by stage in the life cycle: for instance, *Clytia* eggs

express maternal *CheGFP2* mRNA, and planula larvae show typical ring-like *CheGFP1*

fluorescence in the lateral ectoderm at 3 dpf[11]. While *CheGFP1* gene expression

becomes dominant in the planula stage; *CheGFP2, CheGFP3,* and *CheGFP4* are

expressed in different tissues and organs in the medusa[53]. Momose *et al*. succeeded in

genome editing by injecting CRISPR/Cas9 targeting the endogenous *CheGFP1* into

fertilized eggs, and confirmed the knockout effect by loss of green fluorescence in 3 dpf

planula larvae[11]. In addition, *CheGFP1* is not essential for animal survival, which allows

for the monitoring of knockdown effects after embryogenesis. For these reasons, we

chose endogenous *CheGFP1* for the first target gene of siRNA knockdown.

In order to maximize the effect of gene knockdown, we used a combination of

three different siRNAs against *CheGFP1* (Accession No. HQ397706.1; Supplementary

Fig. 4A) in a mixture with a 1:1:1 ratio, and electroporated 600 ng/µl of siRNA mix (200

ng/µl per each siRNA) into fertilized eggs (Fig. 3A). To verify the effect of gene

knockdown, we measured green fluorescence intensity in 2 dpf planula and found that

siRNA (600ng/µl siRNA mix) electroporated embryos showed dramatically reduced

fluorescence intensity compared to the no-siRNA controls (Figs. 3B-D; mean intensity

value: 5.7×10^3 for 0 ng/µl control, 1.8×10^3 for 600 ng/µl). To evaluate the knockdown

effects at varying doses of siRNA, we next electroporated different concentrations of the

CheGFP1 siRNA mix (0, 150, 300, and 600 ng/µl) into fertilized eggs. While control 3

dpf planulae with no-electroporation (No EP) as well as those with no-siRNA showed

typical ring-shape GFP fluorescence in the ectoderm, many planulae with *CheGFP1*

siRNA electroporation lost green fluorescence（Figs. 3E-3H）. To evaluate the GFP

knockdown effects, we classified planulae with typical GFP1 fluorescence as "wild type"

and planulae with reduced fluorescence as "reduced" for further quantification (Figs. 3I

and 3J). The percentage of wild type phenotype dramatically reduced in an siRNA dose-

dependent manner (Fig. 3K). To further quantify the above result on a molecular level,

we examined relative *CheGFP1* mRNA levels by RT-qPCR using the primer set that

was evaluated in a previously study[53]. Consistent with the image analysis results, we

confirmed a marked reduction of *CheGFP1* gene expression at different concentrations

of *CheGFP1* siRNA (Fig. 3L). These results demonstrate that siRNAs targeting

CheGFP1 repress its expression upon electroporation and that the knockdown effects

are dose-dependent.

Given the presence of multiple *CheGFP1* loci (seven nearly identical *CheGFP1*

genes) in the *Clytia* genome[16], it was not clear whether three siRNAs against *GFP1*

sufficiently suppress gene expression in different versions of *CheGFP1*, particularly

those expressed in the planula stage (Supplementary Fig. 4A）. We thus performed RT-

qPCR using the newly designed primer set that amplifies all seven *CheGFP1* genes and

found a dramatic reduction of gene expression upon electroporation of *CheGFP1*

siRNAs (600 ng/μl)（Supplementary Fig. 4B）. Altogether, these results indicate that our

combination of *CheGFP1* siRNAs suppresses gene expression of *CheGFP1* in the

Clytia planula.

Wnt3 knockdown with siRNA in *Clytia hemisphaerica*

Since electroporation of siRNAs targeting endogenous *CheGFP1* was shown to be

effective, we next decided to investigate the effects of knocking down other genes that

play important roles in early development. Wnt3 is a secreted signaling protein in the

Wnt/β-catenin pathway that is involved in diverse developmental processes including

cell fate decisions and patterning[54,55]. In *Clytia*, *Wnt3* RNA is maternally localized to the

animal cortex, the future oral side of the embryo, and Wnt3 organizes axial patterning

as the main ligand for the Wnt/β-catenin pathway in the early embryonic stage[10]. After

embryogenesis, the *Clytia* planula normally elongates along the oral-aboral axis and

becomes oval in shape（Fig. 4A）. By contrast, after *CheWnt3* morpholino injection, the

morphant loses oral-aboral axis polarity and exhibits a spherical shape[10]. This raises the

possibility that knockdown of *CheWnt3* via siRNA may result in a similar spherical

morphology for *Clytia* larvae as is observed in *CheWnt3* morphants. Therefore, to

evaluate siRNA knockdown effects on *Clytia*, we chose *CheWnt3* as our second target.

We used a mixture of two different siRNAs (1:1 ratio) targeting *CheWnt3*, and

carried out electroporation of the siRNA mix into *Clytia* fertilized eggs. After

electroporation of *CheWnt3* siRNAs, 1 dpf planula larvae showed nearly complete

spherical morphology (Fig. 4B), which is reminiscent of *CheWnt3* morphants[10]. To

quantify the effect of siRNA knockdown on *Clytia* larval morphology, we calculated the

aspect ratio in two dimensions, where values greater than 1.0 shows the tendency to be

oval (ellipsoid-like) while a value of 1.0 indicates a perfect circle (sphere) (Fig. 4C). The

control larvae with 0 ng/µl siRNA showed a median aspect ratio of 1.69, indicating a

typical ellipsoid-like morphology. By contrast, larvae that underwent electroporation with

CheWnt3 siRNAs (400 ng/µl) showed a median aspect ratio of 1.16, suggesting a much

greater tendency toward spherical morphology (Fig. 4D). These results quantitively

confirmed the effect of *CheWnt3* knockdown on larval morphology, which phenocopies

CheWnt3 morphants[10]. To analyze the level of gene-specific knockdown at varying

doses of siRNAs, we next electroporated mixtures of 0, 100, 200, and 400 ng/µl of

CheWnt3 siRNAs into *Clytia* fertilized eggs (Figs. 4E-4H). To evaluate *CheWnt3* siRNA

effects on morphology, we classified 1 dpf planulae with elongated ellipsoid-like shapes

as "normal" and planulae with spherical shapes as "spherical" (Figs. 4I and 4J). The

percentage of normal phenotype dramatically reduced after *CheWnt3* siRNA-

electroporation (< 1.0 % in 400 ng/µl), even at the lowest dose (5.2% in 100 ng/µl) (Fig.

4K). To further evaluate the result at a molecular level, we analyzed relative *CheWnt3*

mRNA levels by RT-qPCR and confirmed a significant reduction in *CheWnt3* gene

expression at all concentrations (Fig. 4L). Of note, after electroporation with universal

negative control siRNA（Nippon Gene Co., Ltd.）, both morphological phenotypes and

relative gene expression were comparable to no-siRNA controls (Figs. 4K and 4L),

further confirming that siRNA-mediated knockdown by electroporation is target-specific.

These results together indicate that siRNAs targeting endogenous genes effectively

knock down their expression in *Clytia* embryos.

Effective gene knockdown via shRNA electroporation has previously been

reported using two cnidarian polyps, *Nematostella vectensis* and *Hydractinia*

symbiolongicarpus[36,38]. To test whether shRNAs are also effective in the hydrozoan

jellyfish *Clytia*, we designed shRNAs targeting *CheWnt3* using the same 19 bp target

sequence that was used for siRNAs, and performed electroporation on fertilized eggs

with a mix of two different *Wnt3* shRNAs. After electroporation of *Wnt3* shRNAs, relative

Wnt3 mRNA expression in 1 dpf planula was weakly down-regulated (0.83), and was

not significant compared to the striking effects of *Wnt3* siRNA electroporation (0.039) at

the same concentration (400 ng/µl) (Fig. 4L). In addition, electroporation of *Wnt3*

siRNAs resulted in an extensive morphological change toward spherical shaped

planulae, showing almost no "normal" phenotype (0.71%), whereas electroporation of

shRNA had a limited effect (60.4%) (Fig. 4K). These results suggest that the effect of

siRNA is much stronger than that of shRNA in *Clytia* when treated with the same target

sequence and concentration.

Identification of *Wnt3* in *Cladonema pacificum*

In contrast to *Clytia*, the sole genetic model jellyfish, previous studies have neither

established genetic manipulation nor performed genome sequencing of *Cladonema*

pacificum, despite its unique biological features. This is partly because the extremely

small size of the *Cladonema* egg makes microinjection difficult. By testing whether

siRNA-mediated knockdown via electroporation can work in *Cladonema*, we can start to

manipulate genes in *Cladonema* while also demonstrating that siRNA electroporation

can be applicable to other jellyfish species. While green fluorescent protein (GFP) is

often an easy early target for gene manipulation in emerging organisms, *Cladonema*

does not exhibit apparent endogenous green fluorescence expression at any stage,

unlike *Clytia*. We instead turned our focus to *Wnt3* as the target for gene knockdown in

Cladonema since Wnt3 plays an important role in early embryonic development in

several cnidarian species[56,57].

To identify the *Wnt3* ortholog in *Cladonema*, we utilized RNA-seq results from the

Cladonema polyp, stolon, and medusa manubrium (data not shown). We performed

Cladonema Wnt3 CDS annotation using the *Clytia* CDS database (MARIMBA, Marine

models database: http://marimba.obs-vlfr.fr) and found one contig annotated with *Wnt3*

(*CpWnt3*) and multiple *Wnt* genes (Supplementary Figs. 5 and 6). We then performed

phylogenic analysis using the neighbor-joining method, and confirmed *CpWnt3*

(LC720435) is grouped with medusozoan *Hydra vulgaris* (*HvWnt3*) and *Clytia Wnt3*

(*CheWnt3*) rather than Anthozoa *Nematostella* and *Acropora digitifera* (Supplementary

Fig. 5A). Multiple sequence alignment further showed highly conserved amino acids

sequences among different species including bilaterians and cnidarians (Supplementary

Fig. 5B). These findings raise the possibility that *CpWnt3* has a similar function in

controlling the Wnt/β-catenin pathway in *Cladonema.*

Wnt3 knockdown with siRNA in Cladonema pacificum

Does *Wnt3* function during *Cladonema* embryogenesis, particularly during axial

patterning? Interestingly, *Cladonema* planula larvae do not show a clear elongated

shape (Fig. 5A), as observed in *Clytia*. It is thus possible that morphogenesis, including

axial patterning and/or Wnt/β-catenin pathway function, differs between these two

jellyfish species. It is also possible that the limited elongation in *Cladonema* planulae

might hamper phenotypical appearance upon inhibition of Wnt/β-catenin signaling.

In order to verify that siRNA electroporation works in *Cladonema* and that *Wnt3*

affects axial patterning, we performed electroporation of siRNAs targeting *CpWnt3*. We

prepared fertilized *Cladonema* eggs as we did in *Clytia*, and used a mixture of two

different siRNAs against *CpWnt3* CDS with a 1:1 ratio (300 ng/µl) and performed

electroporation with the previously established parameters (50V, 1 pulse, 25 ms) in

fertilized *Cladonema* eggs. After electroporation of siRNAs for *CpWnt3*, planula larvae

did not exhibit morphological differences compared to controls（Figs. 5A-5C; median

aspect ratio: 1.21 for 0 ng/µl control, 1.16 for 300 ng/µl for si*CpWnt3*）. We then

quantified the morphological phenotypes by calculating the aspect ratio of 1 dpf

planulae and confirmed that *CpWnt3* knockdown does not cause a significant

morphological change (Figs. 5D and 5E), which is consistent with the possibility that

Cladonema planulae simply exhibit limited elongation.

Another possibility that would explain the above result is a defect in the siRNA

knockdown itself. To test this potential explanation, we carried out RT-qPCR using

mRNA samples from 1 dpf planula after electroporating different concentrations of

CpWnt3 siRNA mix (0, 150, and 300 ng/μl) into fertilized eggs. We confirmed a

reduction of *Wnt3* gene expression in an siRNA dose-dependent manner (Fig. 5F),

eliminating siRNA knockdown defects as the cause of our initial result.

To further confirm whether the Wnt/β-catenin pathway is affected by *Wnt3*

knockdown, we examined the gene expression of axial markers *Brachyury* (*CpBra*) and

FoxQ2a (*CpFoxQ2a*), whose expression are influenced by the Wnt/β-catenin pathway in

Clytia[10]. From RT-qPCR, we found that *CpWnt3* knockdown causes a decrease in

CpBra gene expression and an increase in *CpFoxQ2a* gene expression (Supplementary

Fig. 7). We also examined *CpBra* expression by *in situ* hybridization and confirmed the

reduction of *CpBra* expression on the oral side of planula larvae (Figs. 5G-5I), which is

similar to the phenotype exhibited by *CheWnt3* morphants in *Clytia*[10]. Notably, after

CpWnt3 siRNAs electroporation, the rate of metamorphosis from planula to primary

polyp decreased dramatically (Supplementary Table 5), implying potential

developmental defects upon abrogation of the Wnt/β-catenin pathway. These results

suggest that, although overall oral-aboral polarity is less prominent in *Cladonema*

compared to *Clytia* at the morphological level, the axial patterning mechanism mediated

by Wnt/β-catenin signaling may be conserved between these two jellyfish species.

Discussion

In this study, we have established a method to knock down endogenous genes in two

hydrozoan jellyfish, *Clytia hemisphaerica* and *Cladonema pacificum*, via the siRNA

electroporation of fertilized eggs. We showed that knockdown of endogenous *GFP1* in

Clytia causes the loss of GFP fluorescence in the planula stage (Fig. 3), as previously

achieved by CRISPR/Cas9-mediated *GFP1* knockout[11]. We also confirmed that

knockdown of *Wnt3* in *Clytia* induces spherical morphology of the planula (Fig. 4),

mirroring the results of injections of *Wnt3* morpholino antisense oligo[10]. We further

succeeded in efficient repression of *Wnt3* gene expression in *Cladonema* after *Wnt3*

siRNA electroporation and found that expression of axial patterning genes is controlled

by the Wnt/β-catenin pathway (Fig. 5), suggesting that the conserved mechanism is

involved in embryogenesis and planula morphogenesis across hydrozoan jellyfish

species.

Our results show that the knockdown efficiency of siRNA is much greater than

that of shRNA for electroporation of *Clytia* fertilized eggs with the same *Wnt3* target

sequence in the same concentration (Fig. 4K-4L). One possibility that would explain

such distinct effects between siRNA and shRNA electroporation is the existence of

differences in the RNAi machinery or RNA processing efficiency among cnidarians.

During RNAi in mammals, the RNase III Dicer protein processes shRNA in collaboration

with cofactors TRBP (Transactivation response element RNA-binding protein) and

PACT（protein activator of the interferon-induced protein kinase）to produce a mature

form of siRNA[39]. Mature siRNA in the Dicer and TRBP/PACT complex is associated

with the Argonaute protein and cleaves endogenous complementary mRNAs as the

RNA-induced silencing complex [30,58]. These small RNA biogenesis factors, Dicers and

cofactors (TRBP in bilaterians, HYL1 in plants and cnidarians), are found in

Nematostella vectensis, Acropora digitifera, and *Hydra vulgaris*[41]. Although Dicers

(TCONS_00004571 and TCONS_00004525) and Hyl1 (TCONS_00010722) are

predicted to exist in *Clytia* based on MARIMBA transcriptome data, the expression level

of Dicers in oocytes and early gastrulation stages is lower than that in other stages,

suggesting the possibility that insufficient shRNA to siRNA processing efficiency is

responsible for the difference in knockdown efficiency between siRNA and shRNA

electroporation. It will be interesting to elucidate the detailed molecular mechanism of

RNAi in different cnidarian species and address RNA processing efficiency in distinct

developmental stages. The lower efficiency of knockdown by shRNA compared to

siRNA in *Clytia* could also be simply explained by lower shRNA electroporation

efficiency. This possibility could be tested by inserting shRNA or siRNA into fertilized

eggs by microinjection, instead of electroporation, which would be followed by

assessing gene expression and phenotypes.

During *Clytia* early embryogenesis, *Wnt3* RNA is localized to the animal cortex of

the egg and contributes to the formation of the oral-aboral axis[10]. Accordingly,

knockdown of *Wnt3* in *Clytia* fertilized eggs induces suppression of axis formation and

planula elongation (Fig. 4). In contrast, despite the fact that no morphological phenotype

was observed after *Wnt3* knockdown in *Cladonema*, gene expression of the conserved

axial patterning genes, *Brachyury* and *FoxQ2a,* was disrupted (Figs. 5G-5H and

Supplementary Fig. 7). In addition to *Clytia,* the Wnt/β-catenin pathway is involved in

axis formation during cnidarian embryogenesis including *Nematostella*[44,56,59],

Acropora[60]*, and Hydractinia*[57]. Furthermore, given that the Wnt/β-catenin pathway is

also associated with axis formation during metamorphosis in *Hydractinia*[57,61] and

Clytia[62], the reduced metamorphosis rate observed in *CpWnt3* knockdown *Cladonema*

may be attributed to disrupted axis formation (Supplementary Table 5). Taken together,

our data support the pivotal role of Wnt/β-catenin signaling in cnidarian development.

An interesting feature of jellyfish is that their morphology dramatically changes

across their life cycle from planula to polyp and then to medusa. In particular, polyps

and medusae, different adult stages of the same animal, exhibit distinct regenerative

ability, lifespan, and behaviors. Although regeneration mechanisms have been

extensively studied in polyp-only animals such as *Hydra* and *Hydractinia*, classical

studies have used the medusa stage of several jellyfish species and demonstrated their

regenerative potential[21]. Recent work using *Clytia* medusae has further shown a

remarkable remodeling and repatterning mechanism orchestrated by muscle systems

upon organ loss[17]. At the behavior level, the medusae of the upside-down

jellyfish *Cassiopea* exhibit a sleep-like state[63], which is similarly observed in *Hydra*

polyps[64]. More recent work using transgenic *Clytia* has characterized feeding behaviors

in medusae at a neural-network resolution[19]. To understand how these diverse

biological processes in jellyfish are regulated at the molecular level, the functions of

specific genes at various stages must be analyzed. Although gene knockdown by

dsRNA electroporation into polyps in the scyphozoan jellyfish *Aurelia* has been

reported[65], gene knockdown at the medusa stage has not yet been achieved. Given that

siRNA electroporation is applicable to *Hydra* polyps[33,66] in addition to cnidarian fertilized

eggs, we now have a better chance to apply this technique to the medusa stage,

although, because they are susceptible to electric shock, electroporation parameters

must be optimized in order to achieve gene knockdown in medusae. Collectively, our

gene knockdown via siRNA electroporation method will complement the existing shRNA

electroporation approach in cnidarian polyps and will enable molecular-level analysis of

the vast biological phenomena exhibited across the different life stages in jellyfish.

References

1   Technau, U. & Steele, R. E. Evolutionary crossroads in developmental biology: Cnidaria. *Development* **138**, 1447-1458, doi:10.1242/dev.048959 (2011).

2   Zapata, F. *et al.* Phylogenomic Analyses Support Traditional Relationships within Cnidaria. *PLoS One* **10**, e0139068, doi:10.1371/journal.pone.0139068 (2015).

3   Leclere, L. & Rottinger, E. Diversity of Cnidarian Muscles: Function, Anatomy, Development and Regeneration. *Front Cell Dev Biol* **4**, 157, doi:10.3389/fcell.2016.00157 (2016).

4   Holstein, T. W., Hobmayer, E. & Technau, U. Cnidarians: an evolutionarily conserved model system for regeneration? *Dev Dyn* **226**, 257-267, doi:10.1002/dvdy.10227 (2003).

5   Bosch, T. C. G. *et al.* Back to the Basics: Cnidarians Start to Fire. *Trends Neurosci* **40**, 92-105, doi:10.1016/j.tins.2016.11.005 (2017).

6   Houliston, E., Momose, T. & Manuel, M. Clytia hemisphaerica: a jellyfish cousin joins the laboratory. *Trends Genet* **26**, 159-167, doi:10.1016/j.tig.2010.01.008 (2010).

7   Houliston, E., Leclere, L., Munro, C., Copley, R. R. & Momose, T. Past, present and future of Clytia hemisphaerica as a laboratory jellyfish. *Curr Top Dev Biol*, doi:10.1016/bs.ctdb.2021.12.014 (2022).

8   Lechable, M. *et al.* An improved whole life cycle culture protocol for the hydrozoan genetic model Clytia hemisphaerica. *Biol Open* **9**, doi:10.1242/bio.051268 (2020).

9   Momose, T. & Houliston, E. Two oppositely localised frizzled RNAs as axis determinants in a cnidarian embryo. *PLoS Biol* **5**, e70, doi:10.1371/journal.pbio.0050070 (2007).

10   Momose, T., Derelle, R. & Houliston, E. A maternally localised Wnt ligand required for axial patterning in the cnidarian Clytia hemisphaerica. *Development* **135**, 2105-2113, doi:10.1242/dev.021543 (2008).

11   Momose, T. *et al.* High doses of CRISPR/Cas9 ribonucleoprotein efficiently induce gene knockout with low mosaicism in the hydrozoan Clytia hemisphaerica

through microhomology-mediated deletion. *Sci Rep* **8**, 11734,
doi:10.1038/s41598-018-30188-0 (2018).

12   Quiroga Artigas, G. *et al.* A gonad-expressed opsin mediates light-induced
spawning in the jellyfish Clytia. *Elife* **7**, doi:10.7554/eLife.29555 (2018).

13   Quiroga Artigas, G. *et al.* A G protein-coupled receptor mediates neuropeptide-
induced oocyte maturation in the jellyfish Clytia. *PLoS Biol* **18**, e3000614,
doi:10.1371/journal.pbio.3000614 (2020).

14   Lapebie, P. *et al.* Differential responses to Wnt and PCP disruption predict
expression and developmental function of conserved and novel genes in a
cnidarian. *PLoS Genet* **10**, e1004590, doi:10.1371/journal.pgen.1004590 (2014).

15   Condamine, T. *et al.* Molecular characterisation of a cellular conveyor belt in
Clytia medusae. *Dev Biol* **456**, 212-225, doi:10.1016/j.ydbio.2019.09.001 (2019).

16   Leclere, L. *et al.* The genome of the jellyfish Clytia hemisphaerica and the
evolution of the cnidarian life-cycle. *Nat Ecol Evol* **3**, 801-810,
doi:10.1038/s41559-019-0833-2 (2019).

17   Sinigaglia, C. *et al.* Pattern regulation in a regenerating jellyfish. *Elife* **9**,
doi:10.7554/eLife.54868 (2020).

18   Chari, T. *et al.* Whole-animal multiplexed single-cell RNA-seq reveals
transcriptional shifts across Clytia medusa cell types. *Sci Adv* **7**, eabh1683,
doi:10.1126/sciadv.abh1683 (2021).

19   Weissbourd, B. *et al.* A genetically tractable jellyfish model for systems and
evolutionary neuroscience. *Cell* **184**, 5854-5868 e5820,
doi:10.1016/j.cell.2021.10.021 (2021).

20   Leclere, L. *et al.* Maternally localized germ plasm mRNAs and germ cell/stem cell
formation in the cnidarian Clytia. *Dev Biol* **364**, 236-248,
doi:10.1016/j.ydbio.2012.01.018 (2012).

21   Fujita, S., Kuranaga, E. & Nakajima, Y. I. Regeneration Potential of Jellyfish:
Cellular Mechanisms and Molecular Insights. *Genes (Basel)* **12**,
doi:10.3390/genes12050758 (2021).

22   Fujiki, A., Hou, S., Nakamoto, A. & Kumano, G. Branching pattern and
morphogenesis of medusa tentacles in the jellyfish Cladonema pacificum

(Hydrozoa, Cnidaria). *Zoological Lett* **5**, 12, doi:10.1186/s40851-019-0124-4

(2019).

23   Fujita, S., Kuranaga, E. & Nakajima, Y. I. Cell proliferation controls body size

growth, tentacle morphogenesis, and regeneration in hydrozoan jellyfish

Cladonema pacificum. *PeerJ* **7**, e7579, doi:10.7717/peerj.7579 (2019).

24   Hou, S., Zhu, J., Shibata, S., Nakamoto, A. & Kumano, G. Repetitive

accumulation of interstitial cells generates the branched structure of Cladonema

medusa tentacles. *Development* **148**, doi:10.1242/dev.199544 (2021).

25   Takeda, N. *et al.* Identification of jellyfish neuropeptides that act directly as

oocyte maturation-inducing hormones. *Development* **145**,

doi:10.1242/dev.156786 (2018).

26   Stierwald, M., Yanze, N., Bamert, R. P., Kammermeier, L. & Schmid, V. The Sine

oculis/Six class family of homeobox genes in jellyfish with and without eyes:

development and eye regeneration. *Dev Biol* **274**, 70-81,

doi:10.1016/j.ydbio.2004.06.018 (2004).

27   Suga, H., Schmid, V. & Gehring, W. J. Evolution and functional diversity of

jellyfish opsins. *Curr Biol* **18**, 51-55, doi:10.1016/j.cub.2007.11.059 (2008).

28   Suga, H. *et al.* Flexibly deployed Pax genes in eye development at the early

evolution of animals demonstrated by studies on a hydrozoan jellyfish. *Proc Natl*

Acad Sci U S A **107**, 14263-14268, doi:10.1073/pnas.1008389107 (2010).

29   Graziussi, D. F., Suga, H., Schmid, V. & Gehring, W. J. The "eyes absent" (eya)

gene in the eye-bearing hydrozoan jellyfish Cladonema radiatum: conservation of

the retinal determination network. *J Exp Zool B Mol Dev Evol* **318**, 257-267,

doi:10.1002/jez.b.22442 (2012).

30   Meister, G. & Tuschl, T. Mechanisms of gene silencing by double-stranded RNA.

Nature **431**, 343-349, doi:10.1038/nature02873 (2004).

31   Rao, D. D., Vorhies, J. S., Senzer, N. & Nemunaitis, J. siRNA vs. shRNA:

similarities and differences. *Adv Drug Deliv Rev* **61**, 746-759,

doi:10.1016/j.addr.2009.04.004 (2009).

32   Moran, Y. *et al.* Cnidarian microRNAs frequently regulate targets by cleavage.

Genome Res **24**, 651-663, doi:10.1101/gr.162503.113 (2014).

33 Watanabe, H. *et al.* Nodal signalling determines biradial asymmetry in Hydra.
Nature **515**, 112-115, doi:10.1038/nature13666 (2014).

34 Pankow, S. & Bamberger, C. The p53 tumor suppressor-like protein nvp63
mediates selective germ cell death in the sea anemone Nematostella vectensis.
PLoS One **2**, e782, doi:10.1371/journal.pone.0000782 (2007).

35 He, S. *et al.* An axial Hox code controls tissue segmentation and body patterning
in Nematostella vectensis. *Science* **361**, 1377-1380,
doi:10.1126/science.aar8384 (2018).

36 Karabulut, A., He, S., Chen, C. Y., McKinney, S. A. & Gibson, M. C.
Electroporation of short hairpin RNAs for rapid and efficient gene knockdown in
the starlet sea anemone, Nematostella vectensis. *Dev Biol* **448**, 7-15,
doi:10.1016/j.ydbio.2019.01.005 (2019).

37 DuBuc, T. Q. *et al.* Transcription factor AP2 controls cnidarian germ cell
induction. *Science* **367**, 757-762, doi:10.1126/science.aay6782 (2020).

38 Quiroga-Artigas, G., Duscher, A., Lundquist, K., Waletich, J. & Schnitzler, C. E.
Gene knockdown via electroporation of short hairpin RNAs in embryos of the
marine hydroid Hydractinia symbiolongicarpus. *Sci Rep* **10**, 12806,
doi:10.1038/s41598-020-69489-8 (2020).

39 Ha, M. & Kim, V. N. Regulation of microRNA biogenesis. *Nat Rev Mol Cell Biol*
15, 509-524, doi:10.1038/nrm3838 (2014).

40 Krishna, S. *et al.* Deep sequencing reveals unique small RNA repertoire that is
regulated during head regeneration in Hydra magnipapillata. *Nucleic Acids Res*
41, 599-616, doi:10.1093/nar/gks1020 (2013).

41 Moran, Y., Praher, D., Fredman, D. & Technau, U. The evolution of microRNA
pathway protein components in Cnidaria. *Mol Biol Evol* **30**, 2541-2552,
doi:10.1093/molbev/mst159 (2013).

42 Murchison, E. P., Partridge, J. F., Tam, O. H., Cheloufi, S. & Hannon, G. J.
Characterization of Dicer-deficient murine embryonic stem cells. *Proc Natl Acad*
Sci U S A **102**, 12135-12140, doi:10.1073/pnas.0505479102 (2005).

43 Kok, K. H., Ng, M. H., Ching, Y. P. & Jin, D. Y. Human TRBP and PACT directly
interact with each other and associate with dicer to facilitate the production of

small interfering RNA. *J Biol Chem* **282**, 17649-17657,

doi:10.1074/jbc.M611768200 (2007).

44   Watanabe, H. *et al.* Sequential actions of beta-catenin and Bmp pattern the oral

nerve net in Nematostella vectensis. *Nat Commun* **5**, 5536,

doi:10.1038/ncomms6536 (2014).

45   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis.

Nat Methods **9**, 676-682, doi:10.1038/nmeth.2019 (2012).

46   Fujita, S., Kuranaga, E., Miura, M. & Nakajima, Y. I. Fluorescent In Situ

Hybridization and 5-Ethynyl-2'-Deoxyuridine Labeling for Stem-like Cells in the

Hydrozoan Jellyfish Cladonema pacificum. *J Vis Exp*, doi:10.3791/64285 (2022).

47   Thompson, J. D., Higgins, D. G. & Gibson, T. J. CLUSTAL W: improving the

sensitivity of progressive multiple sequence alignment through sequence

weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids*

Res **22**, 4673-4680, doi:10.1093/nar/22.22.4673 (1994).

48   Waterhouse, A. M., Procter, J. B., Martin, D. M., Clamp, M. & Barton, G. J.

Jalview Version 2--a multiple sequence alignment editor and analysis workbench.

Bioinformatics **25**, 1189-1191, doi:10.1093/bioinformatics/btp033 (2009).

49   Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular

Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol* **35**,

1547-1549, doi:10.1093/molbev/msy096 (2018).

50   Deguchi, R., Kondoh, E. & Itoh, J. Spatiotemporal characteristics and

mechanisms of intracellular Ca(2+) increases at fertilization in eggs of jellyfish

(Phylum Cnidaria, Class Hydrozoa). *Dev Biol* **279**, 291-307,

doi:10.1016/j.ydbio.2004.11.036 (2005).

51   Scholz, C. B. & Technau, U. The ancestral role of Brachyury: expression of

NemBra1 in the basal cnidarian Nematostella vectensis (Anthozoa). *Dev Genes*

Evol **212**, 563-570, doi:10.1007/s00427-002-0272-x (2003).

52   Servetnick, M. D. *et al.* Cas9-mediated excision of Nematostella brachyury

disrupts endoderm development, pharynx formation and oral-aboral patterning.

Development **144**, 2951-2960, doi:10.1242/dev.145839 (2017).

53   Fourrage, C., Swann, K., Gonzalez Garcia, J. R., Campbell, A. K. & Houliston, E.

An endogenous green fluorescent protein-photoprotein pair in Clytia

hemisphaerica eggs shows co-targeting to mitochondria and efficient

bioluminescence energy transfer. *Open Biol* **4**, 130206, doi:10.1098/rsob.130206

(2014).

54   Guder, C. *et al.* The Wnt code: cnidarians signal the way. *Oncogene* **25**, 7450-

7460, doi:10.1038/sj.onc.1210052 (2006).

55   Lee, P. N., Pang, K., Matus, D. Q. & Martindale, M. Q. A WNT of things to come:

evolution of Wnt signaling and polarity in cnidarians. *Semin Cell Dev Biol* **17**,

157-167, doi:10.1016/j.semcdb.2006.05.002 (2006).

56   Kraus, Y., Aman, A., Technau, U. & Genikhovich, G. Pre-bilaterian origin of the

blastoporal axial organizer. *Nat Commun* **7**, 11694, doi:10.1038/ncomms11694

(2016).

57   Plickert, G., Jacoby, V., Frank, U., Muller, W. A. & Mokady, O. Wnt signaling in

hydroid development: formation of the primary body axis in embryogenesis and

its subsequent patterning. *Dev Biol* **298**, 368-378,

doi:10.1016/j.ydbio.2006.06.043 (2006).

58   Bartel, D. P. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*

116, 281-297, doi:10.1016/s0092-8674(04)00045-5 (2004).

59   DuBuc, T. Q., Stephenson, T. B., Rock, A. Q. & Martindale, M. Q. Hox and Wnt

pattern the primary body axis of an anthozoan cnidarian before gastrulation. *Nat*

Commun **9**, 2007, doi:10.1038/s41467-018-04184-x (2018).

60   Yasuoka, Y., Shinzato, C. & Satoh, N. The Mesoderm-Forming Gene brachyury

Regulates Ectoderm-Endoderm Demarcation in the Coral Acropora digitifera.

Curr Biol **26**, 2885-2892, doi:10.1016/j.cub.2016.08.011 (2016).

61   Duffy, D. J., Plickert, G., Kuenzel, T., Tilmann, W. & Frank, U. Wnt signaling

promotes oral but suppresses aboral structures in Hydractinia metamorphosis

and regeneration. *Development* **137**, 3057-3066, doi:10.1242/dev.046631

(2010).

62   Krasovec, G., Pottin, K., Rosello, M., Queinnec, E. & Chambon, J. P. Apoptosis

and cell proliferation during metamorphosis of the planula larva of Clytia

hemisphaerica (Hydrozoa, Cnidaria). *Dev Dyn* **250**, 1739-1758,

doi:10.1002/dvdy.376 (2021).

63   Nath, R. D. *et al.* The Jellyfish Cassiopea Exhibits a Sleep-like State. *Curr Biol*
27, 2984-2990 e2983, doi:10.1016/j.cub.2017.08.014 (2017).

64   Kanaya, H. J. *et al.* A sleep-like state in Hydra unravels conserved sleep
mechanisms during the evolutionary development of the central nervous system.
Sci Adv **6**, doi:10.1126/sciadv.abb9415 (2020).

65   Fuchs, B. *et al.* Regulation of polyp-to-jellyfish transition in Aurelia aurita. *Curr*
Biol **24**, 263-273, doi:10.1016/j.cub.2013.12.003 (2014).

66   Lommel, M. *et al.* Hydra Mesoglea Proteome Identifies Thrombospondin as a
Conserved Component Active in Head Organizer Restriction. *Sci Rep* **8**, 11753,
doi:10.1038/s41598-018-30035-2 (2018).

854
855

Acknowledgement

We thank R. Deguchi (Miyagi Univ. Education, Japan) for sharing *Cladonema pacificum*
and EMBRC France for sharing *Clytia hemisphaerica*. We thank T. Momose (Sorbonne
University, CNRS, France) for helpful discussion. We thank I. Nagai, H. Nakatani, and
A. Sasaki for technical assistance; and A. Dahal, A. Tanimoto, and J. Higuchi for
Nematostella culture. This work was supported by JST grant number JPMJCR1852 to
E.K., AMED under grant number JP21gm6110025 to Y.N., and the JSPS KAKENHI
grant numbers JP21H05255 to E.K., and JP17H06332, JP19K22550, JP22H02762 to
Y.N.

865
866

Author contributions

T.M.-O. conceptualized and designed the project, performed experiments using jellyfish,
analyzed data, prepared figures, and wrote the manuscript. S.F. performed experiments
and analyzed data. R.N. performed experiments using *Nematostella*. H.W.
conceptualized and designed the project. E.K. contributed reagents. Y.N.
conceptualized and designed the project, prepared figures, and wrote the manuscript.
All authors approved the final manuscript.

874
875

Data availability

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request and nucleotide sequences are available in the GenBank/EMBL/DDBJ under the accession numbers (*CpWnt1*, LC720432; *CpWnt1b*, LC720433; *CpWnt2*, LC720434; *CpWnt3*, LC720435; *CpWnt5*, LC720436; *CpWnt6*, LC720437; *CpWnt8*, LC720438; *CpWntA*, LC720439; *CpBrachyury*, LC720440; *CpFoxQ2a*, LC720441; *F-actin capping protein subunit beta*, LC720442; *CpEF1alpha*, LC720443).

Competing interests

The authors declare no competing interests.

Figure legends

Figure 1. The hydrozoan jellyfish *Clytia hemisphaerica* and *Cladonema pacificum*. (**A**) Cladogram depicting the phylogenetic position of cnidarian jellyfish. As the sister group of Bilateria, the phylum Cnidaria is divided into two clades, Anthozoa and Medusozoa, which consists of four classes: Hydrozoa, Staurozoa, Scyphozoa, and Cubozoa. Hydrozoa includes polyp type animals without a medusa stage (e.g. *Hydra* and *Hydractinia*) and jellyfish that have both polyp and medusa stages (e.g. *Clytia* and *Cladonema*). (**B** and **C**) Photos of an adult medusa and the eggs of *Clytia hemisphaerica* (**B**) and *Cladonema pacificum* (**C**). Scale bars: 1 mm for medusae; 100 µm for eggs.

Figure 2. The electroporation procedure for jellyfish eggs. (**A**) Schematic of spawning under the 24 h light/dark cycle. *Clytia hemisphaerica* are maintained on a 13 h dark/11 h light cycle. After light stimuli, sperm spawning occurs within 60-90 min and egg spawning occurs within 90-120 min. *Cladonema pacificum* are maintained on a 23.5 h light/0.5 h dark cycle. After dark stimuli, sperm and egg spawning occurs within

25 min. (**B**) Fertilized eggs are resuspended in 15% Ficoll/artificial sea water to prevent

precipitation of eggs during electroporation. (**C**) Electroporation is performed with the

Bio-Rad Gene Pulser Xcell electroporation system and a cuvette with a 4 mm gap. (**D**)

Visualization of the Rhodamine-Dextran delivery into eggs by electroporation.

Rhodamine is shown in red. While little or no fluorescence was observed in eggs

without electroporation (No EP), under the condition of a single 50 V pulse for 25 ms,

rhodamine signals were detected in both *Clytia and Cladonema* eggs without cell

damage. For clarity, egg outlines are indicated with dashed lines for eggs without

electroporation. Scale bars: 200 μm.

Figure 3. **Phenotypes of *GFP1* knockdown with siRNA in *Clytia* fertilized eggs.** (**A**)

Schematic of the siRNA electroporation procedure using fertilized eggs. Fertilized eggs

in 15% Ficoll/artificial sea water mixed with siRNA are incubated for 5 min, and

electroporation is conducted after transferring into a cuvette. (**B** and **C**) Phenotypes of

Clytia 2-day planula after *CheGFP1* siRNA knockdown. (**D**) Boxplots showing the GFP

fluorescence intensity mean values in 2-day planula after electroporation of 0 and 600

ng/μl siRNAs targeting *CheGFP1*. Center lines indicate the medians; x's denote the

means; box limits represent the 25th and 75th percentiles; whiskers show the maximum

and minimum values. 0 ng/μl, n=24; 600 ng/μl, n=81. ***p<0.001. (**E**-**H**) Phenotypes of

3-day planula after siRNA targeting *CheGFP1* electroporation (0, 150, 300, and 600

ng/μl si*CheGFP1*). (**I** and **J**) For quantification of green fluorescence, we classified

phenotypes into two categories: planula with ring-like green fluorescence as "wild-type,"

and planula with reduced green fluorescence as "reduced". (**K**) Quantification of planula

larvae positive for *CheGFP1* (wild-type). Bar plots show the percentage of wild-type

phenotype after *CheGFP1* knockdown. Number of examined planula: 0 ng/μl, *n*=82; 150

ng/μl, *n*=64; 300 ng/μl, *n*=58; 600 ng/μl, *n*=114. Error bars: maximum and minimum

values. Experiments were repeated three times. p-values: 150 ng/μl, p=0.064505; 300

ng/μl, p=0.001437; 600 ng/μl, p=0.00066. **p<0.01, ***p<0.001. (**L**) Quantification of

CheGFP1 mRNA expression levels in 3-day planula by RT-qPCR. *CheEF1alpha* was

used as an internal control. Bar heights represent mean values of at least three

independent experiments. *CheGFP1* expression levels are standardized relative to the

control (0 ng/µl) condition. Error bars: standard deviation. Experiments were performed

in triplicated and repeated at least three times. p-values: 150 ng/µl, p=0.829361; 300

ng/µl, p=0.08588; 600 ng/µl, p=0.037359. *p<0.05. Scale bars: 200 µm.

Figure 4. Phenotypes of *Wnt3* knockdown with siRNA in *Clytia* fertilized eggs. (**A**

and **B**) Typical phenotypes of *Clytia* 1-day planula. While morphologies of control

planula (0 ng/µl) have an elongated oval-shape (A), those of *CheWnt3* siRNA (400

ng/µl) are spherical in shape (**B**). (**C**) Schematic of ellipse approximation for planula

morphology and calculation of the aspect ratio. The aspect ratio was calculated by

dividing the long axis (a) by short axis (b). (**D**) Boxplots showing the aspect ratio of the

1-day planula. Center lines show the medians; x's denote the mean values; box limits

indicate the 25th and 75th percentiles; whiskers show maximum and minimum values;

inner points and outliers are represented by circles. Number of examined planula: 0

ng/µl, *n*=127; 400 ng/µl. *n*=25, ***p<0.001. (**E**-**H**) Phenotypes of 1-day planula after

siRNA targeting *CheWnt3* electroporation (0, 100, 200, and 400 ng/µl si*CheWnt3*). (**I**

and **J**) For morphology quantification, we classified planula phenotypes into two

categories: elongated oval shape as "normal" and spherical (circle) shape as

"spherical". (**K**) Bar plots show the percentage of normal phenotypes across four

different siRNA doses as well as planulae treated with shRNA and siNC, a siRNA

universal <u>n</u>egative <u>c</u>ontrol. Percentages are the mean value and error bars indicate

standard deviation. Numbers of examined planula: 0 ng/µl, *n*=127; 100 ng/µl, *n*=75; 200

ng/µl, *n*=86; 400 ng/µl, *n*=95. shRNA 400 ng/µl, *n*=288; siNC 400 ng/µl, *n*=45.

Experiments were repeated three times. Error bars: maximum and minimum values. p-

values: 100 ng/µl, p=0.000599; 200 ng/µl, p=0.000438; 400 ng/µl, p=0.000402; 400

ng/µl of shRNA, p=0.031928; 400 ng/µl of siNC, p=0.283359. *p<0.05, ***p<0.001. (**L**)

Quantification of *CheWnt3* mRNA levels of 1-day planula by RT-qPCR. *CheEF1alpha*

was used as an internal control. Bar heights represent the mean value. *CheWnt3*

expression levels are standardized relative to the control (0 ng/µl). Error bars: standard

deviation. p-values: 100 ng/µl, p= 0.00799; 200 ng/µl, p= 0.000754; 400 ng/µl, p=

0.000733; 400 ng/µl of shRNA, p= 0.095403; 400 ng/µl of siNC, p= 0.900264. **p<0.01,

***p<0.001. n.s., not significant. Experiments were performed in triplicate and repeated

at least three times. Scale bars: (A, B, E-H) 400 µm, (I, J) 100 µm.

Figure 5. Phenotypes of *Wnt3* knockdown with siRNA in *Cladonema* fertilized

eggs. (**A**) Typical morphology of *Cladonema* egg and planula. The wild-type

Cladonema planula larvae exhibit a slight oval shape. To deliver siRNAs targeting

CpWnt3, electroporation was performed in fertilized eggs (1-cell stage), and phenotypes

were confirmed at planula in one day. (**B** and **C**) Phenotypes of 1-day planula after

CpWnt3 siRNA knockdown. (**D**) Schematic of ellipse approximation for planula

morphology and calculation of the aspect ratio. The aspect ratio was calculated by

dividing the long axis (a) by short axis (b). (**E**) Boxplots showing the aspect ratio of the

1-day planula after si*CpWnt3* electroporation (0 and 300 ng/µl). Center lines show the

medians; x's denote the mean values; box limits indicate the 25th and 75th percentiles;

whiskers show maximum and minimum values; inner points and outliers are

represented by circles. Number of examined planulae: 0 ng/µl, *n*=22; 300 ng/µl, *n*=50.

p= 0.658303. n.s., not significant. (**F**) Quantification of *CpWnt3* mRNA levels in 1-day

planula by RT-qPCR. *CpEF1alpha* was used as an internal control. Bar heights

represent the mean value. Error bars indicate standard deviation. *CpWnt3* expression

levels are standardized relative to the control (0 ng/µl). Experiments were performed in

triplicate and repeated three times. p-values: 150 ng/µl, p= 0.376498; 300 ng/µl, p=

0.073956. (**G, H**) Representative *in situ* hybridization (ISH) image of oral *CpBra*

expression in 1-day planula for control (G: 0 ng/µl si*CpWnt3*) and *CpWnt3* siRNA (H:

300 ng/µl si*CpWnt3*). The oral pole is indicated by an asterisk. (**I**) Quantification of

CpBra expression phenotypes based on ISH images. Stacked bar plots show the

percentage of 1-day planulae in each phenotypic class. *CpBra* expression patterns are

categorized into three phenotypes (strong, weak and none). 0 ng/µl, *n*=74; 300 ng/µl,

n=58. Scale bars: (B, C) 200 µm, (G, H) 50 µm.

A

Bilateria

Cnidaria

Anthozoa

Medusozoa

Hydrozoa
Polyp: *Hydra, Hydractinia*
Jellyfish: *Clytia, Cladonema*

Staurozoa

Scyphozoa

Cubozoa

B Medusa   Egg

Clytia hemisphaerica

C

Cladonema pacificum

Figure 2

Figure 3

Masuda-Ozawa et al.

A Fertilized eggs + siRNA — 5 min — Electroporation

si*CheGFP1* electroporation
2 day planula

B 0 ng/µl

C 600 ng/µl

D GFP intensity mean value (A.U. x10³)

***

0   600 (ng/µl)

si*CheGFP1* electroporation/3 day planula

E 0 (ng/µl)

F 150

G 300

H 600

I Wild type — No EP

J Reduced — si*GFP1*

K Wild type (%)

***

**

0   150   300   600
(ng/µl)

L *GFP1* relative gene expression

*

0   150   300   600
(ng/µl)

A si*CheWnt3* electroporation
1 day planula

A 0 ng/μl B 400 ng/μl

C Oral — Ellipse — Planula — Aboral

Aspect ratio
$$A = \frac{a}{b}$$

D Aspect ratio — *** — 0 — 400 (ng/μl) — 1.8 / 1.2 / 1.0

si*CheWnt3* electroporation/1 day planula

E 0 (ng/μl) F 100 G 200 H 400

I No EP — Normal
J si*Wnt3* — Spherical

K Normal phenotype (%) — *** / *** / *** / * — 0 100 200 400 sh siNC — siRNA (ng/μl)

L *Wnt3* relative gene expression — ** / *** / *** / n.s. — 0 100 200 400 sh siNC — siRNA (ng/μl)

Figure 5

A Egg — si*CpWnt3* electroporation → 1 day planula

si*CpWnt3* electroporation/1 day planula

B 0 ng/µl **C** 300 ng/µl

D Ellipse — Aspect ratio
Planula $A = \dfrac{a}{b}$

E Aspect ratio — 1.8, 1.2, 1.0 — n.s. — 0 / 300 (ng/µl)

F *CpWnt3* relative gene expression — 0 / 150 / 300 (ng/µl)

CpBra ISH

Oral / Aboral

G Control (0 ng/µl) **H** si*CpWnt3* (300 ng/µl)

I *CpBra* ISH phenotype (%)

Control (0 ng/µl): Strong 50.0, Weak 45.9, None 4.1
si*CpWnt3* (300 ng/µl): Strong 20.7, Weak 56.9, None 22.4

Strong / Weak / None

☐ Strong ☐ Weak ☐ None