Title:

Nr2f1-dependent shaping of mitochondrial architecture in newborn mouse hippocampal neurons unravels

novel insights into the neurodevelopmental disorder BBSOAS

Authors:

Bonzano Sara*[1,2], Michelon Filippo[1,2], Crisci Isabella[1,2], Molineris Ivan[1,3], Neri Francesco[1,3], Oliviero Salvatore[1,3],

Beckervordersandforth Ruth[4], Lie Dieter Chichung[4], Peretto Paolo[1,2], Bovetti Serena[1,2], Studer Michèle[5] and De Marchis

Silvia*[1,2]

co-corresponding authors

Affiliations:

[1]Department of Life Sciences and Systems Biology (DBIOS), University of Turin, Via Accademia Albertina 13, Turin, Italy

[2]Neuroscience Institute Cavalieri Ottolenghi (NICO), Regione Gonzole 10, Orbassano (Turin), Italy

[3]IIGM Foundation-Italian Institute for Genomic Medicine, Turin, Italy.

[4]Institut für Biochemie, Friedrich-Alexander Universität Erlangen-Nürnberg (FAU), Fahrstrasse 17, Erlangen, Germany.

[5]Institute de Biologie Valrose (iBV), Université Côte d'Azur (UCA), CNRS 7277, Inserm 1091, Avenue Valrose 28, Nice,

France.

Co-corresponding authors:

Sara Bonzano

sara.bonzano@unito.it

+39 011 670 6632 4685

Silvia De Marchis

silvia.demarchis@unito.it

+39 011 670 6632 4685

Abstract

The nuclear receptor Nr2f1, also known as COUP-TFI, plays a key role during cortical development by acting as a strong transcriptional regulator in embryonic and adult neural cells. In humans, NR2F1 mutations cause the Bosch-Boonstra-Schaaf optic atrophy-intellectual syndrome (BBSOAS), a rare neurodevelopmental disorder characterized by multiple clinical features including optic nerve atrophy, cerebral visual impairment, seizures, intellectual disability, and autistic traits. Moreover, alterations in mitochondrial energy supply have been recently reported in some BBSOAS patients. Although mitochondrial dysfunction is increasingly implicated in neurodevelopmental disorders, whether and how Nr2f1 can regulate mitochondria function in neural cells is still completely unknown. To address these questions, we have combined genome-wide and *in silico* analyses with mouse genetics, neuroanatomical and imaging approaches. First, we found that key factors for mitochondrial function and dynamics represent potential genomic targets under direct Nr2f1 transcriptional control in neurons. To genetically manipulate Nr2f1 function in a restricted neuronal population, we focused on the adult mouse hippocampal dentate gyrus, a site of persistent neurogenesis in rodents and where Nr2f1 is highly expressed. We showed that both loss- and gain-of-function of Nr2f1 within the hippocampal neurogenic niche led to altered mitochondrial architecture in newborn neurons likely reflecting alteration in mitochondrial dynamics. Lastly, we identified the fusion factor Mfn2 as a putative Nr2f1 target and demonstrated its downregulation following both Nr2f1 adult depletion and embryonic haploinsufficiency in mice. Overall, our study provides the first evidence of a significant role of Nr2f1 in shaping mitochondrial architecture in neurons and opens a promising avenue for the identification of new mechanisms implicated in BBSOAS pathogenesis involving mitochondrial dysfunction.

Keywords

COUP-TFI, mitochondria, dentate gyrus, neurogenesis, adult-born neurons, Mfn2

Introduction

The role of the key transcriptional regulator Nr2f1 (Nuclear receptor subfamily 2, group f, member 1), also known as COUP-TFI (Chicken Ovalbumin Upstream Promoter-Transcription Factor 1), has been intensively investigated during brain development, where it exerts pleiotropic functions including the control of neural stem cell competency, cortical area and cell-type specification, and axonal pathfinding (Bertacchi *et al*, 2019b). In humans, *NR2F1* has recently emerged as a disease gene: multiple *NR2F1* pathological variants cause the Bosch-Boonstra-Schaaf optic atrophy-intellectual syndrome (BBSOAS; OMIM: 615722), a rare monogenic autosomal-dominant disorder characterized by multiple clinical features including global developmental delay, mild-to-severe intellectual disability (ID), optic nerve atrophy, seizures and traits characteristic of the autism spectrum disorder (ASD) (Bosch *et al*, 2014; Chen *et al*, 2016; Kaiwar *et al*, 2017). Most variants in the NR2F1 gene of patients described so far are deletions and/or mutations predominantly located in the DNA-binding domain (DBD). Most of the pathogenic variants lead to haploinsufficiency or dominant-negative effects, thus compromising and/or completely abolishing NR2F1 transcriptional regulatory activity (Chen *et al*, 2016; Bosch *et al*, 2016; Kaiwar *et al*, 2017; Rech *et al*, 2020; Billiet *et al*, 2021). Recently, alterations in mitochondrial energy supply due to a defective function of the electron transport chain (ETC), have been reported in muscle biopsies from two BBSOAS patients (Martín-Hernández *et al*, 2018; Hobbs *et al*, 2020), suggesting that altered NR2F1 transcriptional activity might lead to mitochondria-related functional defects. Furthermore, several neurological symptoms - such as optic atrophy, hypotonia, and seizure - described in BBSOAS patients are often associated to mitochondrial dysfunction in the nervous system (Valenti *et al*, 2014; Zsurka & Kunz, 2015; Frye, 2020; Lenaers *et al*, 2021). Nevertheless, whether and how NR2F1 can impact on mitochondrial dynamics and/or function in neural cells is still unknown. Importantly, no data are available on the cellular/molecular mechanisms by which NR2F1 loss-of-function might trigger mitochondrial dysfunction in BBSOAS patients. To address these questions, we decided to focus on a restricted neuronal population by exploiting adult neurogenesis in the mouse hippocampal dentate gyrus (DG), where Nr2f1 is highly expressed (Bonzano *et al*, 2018). Indeed, in the DG subgranular zone (SGZ), adult neural stem/progenitor cells (aNSPCs) ensure the continuous generation of new neurons whose integration into the preexisting circuitries is crucial for proper adaptive behaviors and cognitive functions (such as learning and memory) (Aimone *et al*, 2014). We have previously demonstrated that Nr2f1 drives aNSPCs toward a neuronal lineage inhibiting an astroglial fate (Bonzano *et al*, 2018). Interestingly, DG immature neurons show higher Nr2f1 expression (Bonzano *et al*, 2018; Artegiani *et al*, 2017) and an overall increase in mitochondrial mass compared to aNSPCs (Steib *et al*, 2014; Beckervordersandforth, 2017). Moreover, mitochondrial dysfunction impairs DG neurogenesis (Beckervordersandforth *et al*, 2017; Beckervordersandforth, 2017; Khacho *et al*, 2016; Steib *et al*, 2014). Thus, the adult hippocampal neurogenic process represents an excellent model to dissect the role of Nr2f1 on mitochondria.

In this study, we demonstrate that Nr2f1 shapes mitochondrial architecture in adult-born DG neurons by various approaches, ranging from genome-wide and *in silico* analyses to mouse genetics and state-of-the-art neuroanatomical and imaging techniques. We provide evidence that this process occurs through the regulation of nuclear genes encoding proteins essential for mitochondrial dynamics and function. Among them, we identified the mitochondrial membrane protein mitofusin-2 (Mfn2) that plays a central role in regulating mitochondrial fusion, as a putative target gene of Nr2f1 transcriptional activity. We found that Mfn2 is downregulated in the DG of mice heterozygous for Nr2f1, a validated BBSOAS mouse model (Bertacchi *et al*, 2019a; Chen *et al*, 2020; Jurkute *et al*, 2021), and we confirmed its downregulation

109 in adult-born DG neurons conditionally depleted for Nr2f1, supporting a cell-intrinsic effect. Overall, our data provide new

110 insights into the functional role of Nr2f1 by demonstrating its direct involvement in shaping mitochondria in adult-born DG

111 neurons. This paves the way for further research on the mechanisms and role of mitochondrial dysfunction in the

112 pathogenesis of BBSOAS, opening new possibilities for therapeutic intervention.

Results and Discussion

Nr2f1 nuclear genomic targets are enriched in key mitochondrial factors

An involvement of mitochondrial dysfunction in BBSOAS patients has been recently reported in muscles without any signs

of mitochondrial DNA pathological variants (Martín-Hernández *et al*, 2018; Hobbs *et al*, 2020) and multiple clinical

features associated to BBSOAS are compatible with altered mitochondrial function in the brain (Valenti *et al*, 2014; Zsurka

& Kunz, 2015; Frye, 2020; Lenaers *et al*, 2021). These evidences suggest that the nuclear transcription factor Nr2f1 might

impinge on mitochondria in the neural tissue by directly binding and regulating the expression of nuclear-encoded

mitochondrial genes that code for more than 95% of the mitochondrial proteins (Calvo *et al*, 2016; Misgeld & Schwarz,

2017). We thus carried out a genome-wide analysis of Nr2f1 occupancy by Chromatin immunoprecipitation followed by

deep sequencing (ChIP☐seq) (**Figure 1a**) on the adult neocortex, which is highly enriched in Nr2f1 expressing neurons,

with virtually no Nr2f1 expression in glia cells (**Figure S1a,b**) (Foglio *et al*, 2021). By peak calling analysis, we identified

2119 binding sites for Nr2f1 and revealed that these sites were enriched in both CpG islands and promoter regions (i.e.

☐3kb/+2kb from the Transcription Starting Site - TSS - of annotated genes) (**Figure 1b and Figure S1c**). Almost all Nr2f1-

bound promoters were positive for H3K4me3, a histone mark highly enriched at active promoters (Liang *et al*, 2004), but

not all H3K4me3+ promoters turned to be bound by Nr2f1 (**Figure 1a**), indicating specificity. Notably, predicting TF

binding by the Homer motif discovery tool confirmed that the observed genomic peaks are highly enriched in the putative

Nr2f1 consensus sequence (**Figure 1c and Figure S1d**), indicating direct binding. Together, these data show that in neurons

the transcription factor Nr2f1 binds to promoters predominantly in a chromatin permissive state, in line with a previous

study reporting a high association of Nr2f1/2 with open chromatin characterized by high levels of p300 and H3K27ac in

neural crest cells (Rada-Iglesias *et al*, 2012).

The set of genes identified as putative Nr2f1 genomic targets by ChIP-seq was then analyzed by Gene Ontology (GO) term

annotation through the PANTHER classification system. Concerning the "cellular component" class, the analysis revealed

a significant enrichment in genes involved in the control of mitochondrial structure (e.g. mitochondrial membranes and

matrix organization) (**Figure 1d**). Notably, GO clustering of Nr2f1-bound genes according to the cellular localization of

their products clearly showed a higher fold-enrichment and more significant over-representation for mitochondria (e.g.,

"mitochondrion", fold-enrichment=2.04; FDR=8.20e-21) rather than non-mitochondrial subcellular location (e.g.,

"cytoskeleton", fold-enrichment=1.27, FDR=0.0425; "Golgi apparatus", fold-enrichment=1.41, FDR=0.0061). In line,

analysis of the "biological process" class revealed an enrichment in genes involved in mitochondrial function, including the

organization of mitochondrial membranes as well as mitochondria-related metabolism and cellular respiration (**Figure 1d**).

Thus, Nr2f1 likely acts as a transcriptional regulator of key mitochondrial factors in the adult mouse neural tissue.

Interestingly, a previous study reported altered expression of nuclear genes encoding mitochondrial proteins (e.g.,

SLC25A1, ECHS1, MRPL45, TIMM23) in the *Nr2f1-null* embryonic neocortex (Montemayor *et al*, 2010). Moreover, by

interrogating a recently published RNA-sequencing dataset (Chen *et al*, 2020), we found altered expression of multiple

genes coding for mitochondrial proteins (e.g., *OPA1, GDAP1, ACO2, ATP5G2*) in the hippocampi of adult *Nr2f1-*

heterozygous mice. Taken together, these data suggest that Nr2f1 may exert a direct role on the transcriptional control of

nuclear genes crucial for mitochondria in neurons. Nevertheless, to date there is no clear evidence for Nr2f1 in controlling

mitochondrial organization and/or function *in vivo*.

Nr2f1 loss-of-function decreases the mitochondrial mass of adult-born hippocampal neuron

To study the role of Nr2f1 on mitochondria, we focused on the neurogenic niche of the adult hippocampal neurogenic niche.

Here, generation of new neurons from adult neural stem/progenitor cells (aNSPCs) is accompanied by increasing levels of

Nr2f1 expression (Bonzano *et al*, 2018; Artegiani *et al*, 2017) and by a dramatic reorganization of the mitochondrial

network leading to increased mitochondrial mass in newborn neurons (Steib *et al*, 2014; Beckervordersandforth, 2017;

Beckervordersandforth *et al*, 2017; Khacho *et al*, 2019). We used an *in vivo* conditional loss-of-function (LOF) approach

targeting the aNSPC lineage *via* tamoxifen (TAM) administration in triple transgenic mice carrying the inducible *Cre-*

recombinase (*CreERT2*) under the Glast promoter, both Nr2f1 alleles flanked by loxP sites and the reporter gene encoding

the yellow fluorescent protein (YFP) for lineage tracing (herein named *Nr2f1-icKO*; **Figure 2a**) (Bonzano *et al*, 2018;

Armentano *et al*, 2007). Animals carrying both *Cre* and YFP alleles but wildtype for *Nr2f1* were used as controls (*Ctrl*). To

label mitochondria in adult-born hippocampal neurons, a retroviral vector encoding the mitochondria-targeted fluorescent

protein DsRed (RV-mitoDsRed) was stereotaxically injected into the DG two weeks after induction by TAM. Mice were

then analyzed at 17 days post viral-injection (**Figure 2a'**), allowing recombined and transduced newborn cells to develop

into doublecortin-expressing neurons (i.e., DCX+mitoDsRed+YFP+ cells). Notably, while in *Ctrl* mice YFP+ cells were

virtually all Nr2f1-positive, in *Nr2f1-icKO* mice the population of YFP+ cells - including the specific subset transduced by

the mitoDsRed virus - were substantially devoid of Nr2f1 staining (**Figure 2b,c**), thus validating *Nr2f1* inactivation in the

aNSPC neuronal lineage of *Nr2f1-icKO* mice.

To perform the analyses, we selected DCX+mitoDsRed+YFP+ newborn neurons from *Nr2f1-icKO* and *Ctrl* mice showing

comparable patterns of the dendritic trees throughout the DG layers (i.e., the primary dendrite branching in the outer part of

the granule cell layer and the dendritic arborization spanning the molecular cell layer; **Figure S2a and Figure 3a,b**).

Morphometric analyses on those cells revealed no changes in the size of perikarya (**Figure S2a'**) but a decrease in the total

dendritic length and in the number of dendritic branching points (**Figure S2a'',a'''**), thus suggesting a reduced complexity

of the dendritic tree.

To directly assess the consequences of *Nr2f1* LOF on mitochondria in DG newborn neurons, we analyzed the mitochondrial

occupancy expressed as the percentage of the area occupied by the mitoDsRed+ signal over the area encompassing the

perikaryon and the cone-shaped proximal portion of the primary dendrite (named soma hereafter) (**Figure 2d and Figure**

S2c). MitoDsRed+ mitochondria appear densely packed and tightly organized into complex networks in the soma of control

DCX+mitoDsRed+YFP+ newborn neurons (**Figure 2d,e and Figure S2b-b'',c**). Interestingly, we observed a reduction by

about 20% in the mitochondrial occupancy within the soma in Nr2f1-depleted neurons compared to control ones (**Figure**

2d,e). By exploiting the Mitochondrial Network Analysis (MiNA) toolset (Valente *et al*, 2017), we confirmed a decreased

mitochondrial mass (i.e., mitochondrial footprint) and we further revealed a reduced complexity of the mitochondrial networks in Nr2f1-deficient newborn neurons (**Figure S2d**). In line, we found a reduction by 28% in the area covered by mitoDsRed+ mitochondria within the cylindrical portion of the primary dendrite of Nr2f1-deficient newborn neurons (**Figure 2d,f** and **Figure S2b''',e**).

All together, these findings provide the first evidence of reduced mitochondrial mass and network complexity due to *Nr2f1* LOF in neurons.

Nr2f1 loss-/gain- of function in adult-born DG neurons alters dendritic mitochondrial number and morphology

We next evaluated the mitochondrial content and architecture in the distal dendrites of DCX+mitoDsRed+YFP+ newborn neurons, where high-resolution confocal microscopy allows the analysis of single mitoDsRed+ mitochondria (**Figure 3a-b'**) and thus enables quantification and in-depth morphometric assessment. We found a net reduction in the abundance of mitochondria in Nr2f1-depleted neurons not only when quantified as total number of mitochondria per dendritic arborization, but also when normalized to the total length of the dendritic arbor (**Figure 3c,d**). Even if the number of mitochondria linearly correlates to the dendritic arborization length both in *Ctrl* and in *Nr2f1-icKO* neurons (Spearman correlation: *Ctrl*, r=0.7868, p=0.0021; *Nr2f1-icKO*, r=0.8326, p<0.001), *Nr2f1-icKO* cells always show a lower number of mitochondria compared to control cells of akin dendritic length (**Figure 3e**), thus supporting that alteration in mitochondrial content is not secondary to morphological defects of the dendritic tree. On the other hand, previous studies indicate that mitochondria play a key role in the architecture of the dendritic arbor (Kimura & Murakami, 2014; Fang *et al*, 2016; Steib *et al*, 2014), thus altered dendritic mitochondria in newborn neurons lacking Nr2f1 might influence dendritic length and branching.

Furthermore, we observed a reduction in the maximum length of single dendritic mitochondria in *Nr2f1-icKO* neurons compared to control ones (**Figure 3f-h**). Notably, by classifying mitochondria into four different categories based on their length (adapted from (Khacho *et al*, 2016)), we observed an increase in fragmented mitochondria concomitantly with a reduction in tubular ones in the dendrites of *Nr2f1-icKO* neurons (**Figure 3i**). In line, estimation of the linear mitochondrial occupancy (i.e., the percentage of the dendritic length covered by the mitoDsRed+ signal) unveiled a drastic decrease (about 55%) in Nr2f1-depleted neurons compared to controls (**Figure 3j**). Overall, our morphometric analyses on mitoDsRed+ mitochondria revealed that loss of Nr2f1 function in adult-born hippocampal neurons induces a global impairment in the mitochondrial mass that appears progressively more dramatic from the proximal to the distal cell compartments (**Figure S2f**).

To further confirm the involvement of Nr2f1 function on mitochondrial architecture, we exploited a complementary genetic strategy to increase *Nr2f1* expression (i.e. gain-of-function, GOF) in adult-born hippocampal neurons, by crossing the transgenic *CAGGS-lox-stop-lox-hCOUP-TFI* mouse line (Wu *et al*, 2010) to the *Glast::CreERT2* and *YFP* reporter line (herein *Nr2f1-O/E*; **Figure S3a**) (Bonzano *et al*, 2018). Analysis of the mitochondrial content on the dendritic arborization of DCX+mitoDsRed+YFP+ newborn neurons in *Nr2f1-O/E* mice revealed the occurrence of longer mitochondria without changes in their number, leading ultimately to an overall increase in the linear dendritic mitochondrial occupancy in newborn granule neurons (**Figure S3b-d**).

Data obtained by *Nr2f1* LOF and GOF, together with GO annotations disclosing a stronger enrichment for nuclear-encoded

mitochondrial genes than that for other structural genes (e.g., cytoskeleton-related genes)-as Nr2f1 direct targets (**Figure 1**),

strongly support a primary effect of Nr2f1 manipulation on mitochondria in adult-born DG neurons.

The mitochondrial fusion factor Mfn2 is downregulated in mouse models of BBSOAS

Mitochondria undergo continuous morphological transition by dynamic processes thanks to the coordinated action of

proteins promoting mitochondrial fission (e.g., Drp1, Fis1) and fusion (e.g., Mfn1/2, Opa1), as well as their transport (e.g.,

Miro, Milton) (Course & Wang, 2016; Misgeld & Schwarz, 2017; Giacomello *et al*, 2020). It is well known that altered

expression and function of those factors leads to permanent changes in the overall shape of mitochondria (Chan, 2020) that

could underlie the observed mitochondrial phenotype following *Nr2f1* LOF and GOF. We thus wondered whether altered

Nr2f1 activity might impinge on mitochondrial dynamics. To this purpose, we exploited our ChIP-seq dataset for Nr2f1

(**Figure 1**) and unveiled that three of the key nuclear genes encoding proteins crucial for mitochondrial dynamics, namely

dynamin-2 (Dnm2), mitochondrial dynamics protein of 51kDa (MiD5) and mitofusin-2 (Mfn2), are indeed bound by Nr2f1

on their promoter regions (**Figure 4a**). Based on the fragmented and elongated mitochondrial shape observed in *Nr2f1-icKO*

and *Nr2f1-O/E* neurons respectively, we focused our attention on the fusion factor *Mfn2* as a putative target for Nr2f1

action. Notably, Nr2f1 binds to a promoter region closed to the transcription starting site (TSS) of the *Mfn2* gene, which is

associated to the H3K4me3 in the adult neural tissue (**Figure 4a**). We next crossed our Chip-seq dataset with previously

published transcriptome data on the hippocampi of adult *Nr2f1-heterozygous* mice (*Nr2f1-het)* (Chen *et al*, 2020), a

validated BBSOAS mouse model (Bertacchi *et al*, 2019a; Jurkute *et al*, 2021; Tocco *et al*, 2021). By Gene Set Enrichment

Analysis (GSEA), we found that among the 1053 enriched genes approximately 8% (82/1053) code for mitochondrial

proteins, including the *Mfn2* gene (**Figure 4b**). In addition, we did observe a reduction in Mfn2 protein levels (as the area

occupied by Mfn2+ signal and its intensity) in the DG of *Nr2f1-het* mice by *in situ* densitometric analysis (**Figure 4c,d,e**).

To validate Mfn2 downregulation upon *Nr2f1* LOF we quantified Mfn2 fluorescence labelling within the soma of triple-

labeled Mfn2+DCX+YFP+ newborn neurons one months after TAM administration in *Nr2f1-icKO* mice (**Figure 4f,g**). In

line with the observed reduction in mitochondrial occupancy in *Nr2f1-icKO* newborn neurons (**Figure 2e**), we found a 15%

reduction in the percentage of area occupied by the Mfn2+ signal compared to controls (**Figure 4h,i**). Notably, analysis of

the intensity of Mfn2 staining normalized to the area occupied by Mfn2+ signal (CTMF; about 40% decrease; **Figure 4j,k**),

revealed a drastic drop, indicating that Nr2f1 deletion lowers Mfn2 levels in a cell-autonomous manner.

It is worth noting that the morphological changes we did observe in adult-born neurons lacking Nr2f1, including

mitochondrial fragmentation and decreased mitochondrial mass, as well as a progressively stronger phenotype proceeding

from proximal to distal cell compartments (**Figure S2f**), recapitulate the phenotype of Mfn2 downregulation described in

differentiated excitatory neurons *in vitro* (Fang *et al*, 2016). Overall, it is thus highly conceivable that Nr2f1 might act as

upstream regulator of Mfn2 expression controlling mitochondrial dynamics in adult-born DG neurons.

Conclusion

Mitochondria, which exert essential roles in both bioenergetic and non-energetic biological processes, are increasingly

recognized as major players in neuronal development and function, including neural stem cell activation, dendritogenesis,

synaptic transmission and plasticity (Khacho *et al*, 2016; Kimura & Murakami, 2014; Faits *et al*, 2016; Steib *et al*, 2014;
Khacho *et al*, 2019; Rangaraju *et al*, 2019b, 2019a). According to their crucial functions, it is becoming increasingly clear
that mitochondrial dysfunction dramatically contributes to the pathogenesis of various neurodegenerative and
neurodevelopmental disorders (Jurcau, 2021; Monzio Compagnoni *et al*, 2020; Rojas-Charry *et al*, 2021). Interestingly, a
mitochondrial involvement has been recently reported for the BBSOAS, a human monogenic neurodevelopmental
syndrome caused by NR2F1 haploinsufficiency (Bosch *et al*, 2014; Kaiwar *et al*, 2017; Chen *et al*, 2016) and characterized
by several symptoms including intellectual disability and optic atrophy (Bosch *et al*, 2014; Kaiwar *et al*, 2017; Rech *et al*,
2020). Specifically, a mitochondrial dysfunction was described in the hypotonic skeletal muscles in two BBSOAS patients
(Martín-Hernández *et al*, 2018; Hobbs *et al*, 2020).

Here, by exploiting genetic mouse models of BBSOAS we unveil a novel and important link between Nr2f1 function and
mitochondria in the central nervous system. First, we discovered several mitochondrial factors, including the mitochondrial
fusion factor Mfn2, among the putative direct genomic targets for the transcription factor Nr2f1. Second, thanks to
conditional inactivation of Nr2f1 in adult-born hippocampal neurons and to in-depth morphometric analyses of
mitochondria at a single cell resolution *in vivo*, we clearly demonstrated the occurrence of a mitochondrial phenotype in
neurons lacking *Nr2f1* function, including altered mitochondrial mass and shape. Last, we provided evidence of altered
expression of Mfn2 upon Nr2f1 loss-of-function/haploinsufficiency, in line with the observed mitochondrial phenotype. In
light of our findings, the mitochondrial abnormalities recently described in the optic nerve and retina of *Nr2f1* constitutive
knockout/het mice (Bertacchi *et al*, 2019a) might be a direct consequence of Nr2f1 mutations and contribute to the optic
atrophy in Nr2f1 mouse models. As the homology between human and mouse *Nr2f1* is very high (especially in the DNA
binding domain where there is 100% of amino acid sequence homology) (Qiu *et al*, 1995; Alfano & Studer, 2013), their
functions and targets are likely to be conserved in both species. In this study, we have only scratched the surface of the
mitochondrial dysfunction underlying the complex phenotype of *Nr2f1* haploinsufficiency in the brain and future
investigations are needed to clarify the role of dysfunctional mitochondria in the pathogenesis of BBSAOS. Altogether, our
data point to mitochondrial dysfunction in neural tissue as a key pathological mechanism in BBSOAS, suggesting that the
current estimation of mitochondrial involvement in BBSOAS patients might be underestimated.

Material and Methods

Animals and treatments
In vivo experiments were performed on two to three-month-old C57BL/6J mice (Charles Rivers) of both genders. Triple
transgenic mice (C57BL/6J background) were also used to manipulate Nr2f1 expression through tamoxifen administration.
Glast::CreERT2$^{+/wt}$;R26-loxP-stop-loxP-YFP$^{yfp+/yfp+}$;Nr2f1$^{fl/fl}$ (*Nr2f1-icKO*) were used for *in vivo* loss-of-function
experiments and *Glast::CreERT2$^{+/wt}$;R26-loxP-stop-loxP-YFP$^{yfp+/yfp+}$;CAG-S-loxP-stop-loxP-hCOUP-TFI$^{+/wt}$* (*Nr2f1-O/E*)
were used for *in vivo* Nr2f1 overexpression (see (Bonzano *et al*, 2018)). *Glast::CreERT2$^{+/wt}$;R26-loxP-stop-loxP-*
YFP$^{yfp+/yfp+}$ were used as control mice (*Ctrl*). For activation of the *CreERT2*-recombinase in the Glast+ lineage, animals
were administered with tamoxifen (TAM, SIGMA, T-5648) at the dose of 2,5 mg/mouse/day dissolved into corn-oil

(SIGMA) by means of intraperitoneal injections (i.p.) for 2 consecutive days at the age of two-months (Bonzano *et al*,

2018). Adult (four-month-old) constitutive *Nr2f1* heterozygous mice (i.e. *Nr2f1^(wt/null)*, named *Nr2f1-het*) were generated and

genotyped as previously described (Jurkute *et al*, 2021). Littermates of *Nr2f1-het* mice with wildtype Nr2f1 alleles were

used as controls (i.e., *Nr2f1^(wt/wt)*, WT). Both *Nr2f1-het* and WT littermates were bred in a 129S2/SvPas background. Mice

were housed under standard laboratory conditions (n=2-4 mice/standard cage) with basic recommended environmental

enrichment (paper tubes and litter, igloo) under a 12 h light/dark cycle with access to food and water *ad libitum*. All

procedures were conducted in accordance with the Guide for the Care and Use of Laboratory Animals of the European

Community Council Directives (2010/63/EU and 86/609/EEC) and approved by local bioethics committees, the Italian

Ministry of Health (Authorization number 864/2018-PR, protocol E6B5E.3) and the French Ministry of Education,

Research and Innovation CIEPAL NCE/2019–548 (Nice) under authorization #15 349 and #15 350.

Retroviral production

Replication-deficient recombinant murine Moloney leukemia (MML) retroviruses specifically transduce proliferating cells

and allow the dating of their birth, as well as label precursor cells and their progeny in the adult hippocampal neurogenic

lineage (Zhao *et al*, 2006). CAG-IRES-mitochondrial Discosoma red (mitoDsRed) were generated from the pCAG IRES–

GFP vector (Jagasia *et al*, 2009) by replacing the GFP coding sequence with cDNA for mitochondrially targeted DsRed.

Retroviruses were generated as described previously(Zhao *et al*, 2006; Steib *et al*, 2014). Virus-containing supernatant was

harvested four times every 48 h after transfection and concentrated by two rounds of ultracentrifugation. Viral titers were

determined by transduction of HEK293T cells for 72 h with limiting dilution of MMLV suspension and counting of reporter

expressing cell spots under a fluorescent microscope (Leica Microsystems). Titers applied were ~5×10^8 colony-forming

units (CFU)/ml (Zhao *et al*, 2006).

Surgical procedure for retroviral injection

Two weeks after TAM administration, adult mice were anesthetized in a constant flow of Isofluorane (3%) in oxygen,

positioned in a stereotaxic apparatus (Stoelting) and injected with a pneumatic pressure injection apparatus (Picospritzer II,

General Valve Corporation). The skull was exposed by an incision in the scalp and a small hole (about 1 mm) drilled

through the skull. 0.8 µl of retrovirus-mitoDsRed was injected in the DG using a sharpened glass capillary at the following

stereotaxic coordinates: -2 mm (antero-posterior), 1.5 mm (lateral) to Bregma and -2.0 mm below the surface of the skull.

Mice (n=3/4 genotype/experiments) were analyzed 17 days after retroviral injection.

Tissue preparation and immunofluorescence staining

For immunostaining, adult mice were deeply anesthetized with an intraperitoneal injection of a mixture of tiletamine and

zolazepam (40-80 mg/kg; Zoletil; Virbac) and perfused transcardially with ice-cold 0.9% saline solution followed by ice-

cold 4% paraformaldehyde (PFA) in 0.1 M phosphate buffer (PB), pH 7.4. Brains were removed from the skull, post-fixed

for 4 hours in the same PFA solution, cryoprotected in a 30% sucrose solution (in 0.1 M PB, pH 7.4), embedded with cryo-

embedding matrix (OCT; Killik, BioOptica), frozen at -80°C and finally sectioned using cryostat (Leica Microsystems).

Free-floating coronal serial sections (50µm-thick for the hemisphere ipsilateral to the viral injection; 40µm-thick for the

contralateral hemisphere) were collected in series on multi-well dishes (6 to 8 wells per animal). Sections were stored at –20°C in antifreeze solution (30% ethylene glycol, 30% glycerol, 10% PB: 189 mM NaH$_2$PO$_4$, 192.5 mM NaOH; pH 7.4) until use. Immunofluorescence reactions were performed on free-floating coronal serial sections as detailed below: sections were incubated in blocking solution (0.01M PBS - pH 7.4, 1 or 2% Triton X-100 for 40 and 50 µm-thick slices respectively, and 10% normal serum of the same species of the secondary antibody, i.e., normal donkey serum – NDS) for two hours at room temperature. Afterwards, slices were incubated for 48 hours at 4°C with primary antibodies (see Table 1) diluted in 0.01M PBS (pH 7.4), 1% to 2% Triton X-100 (for 40 and 50 µm-thick slices respectively), and 1% NDS. For immunostainings against Mfn2, sections were subjected to antigen retrieval. Slices were incubated in 10mM Tris, 1mM EDTA, 0.05% Tween 20 for 2 min at 99°C and washed three times with MilliQ water followed by three washing step with 0.01M PBS (pH 7.4) prior incubation with primary antibody. Sections were washed in PBS and incubated for 2 hours or overnight (for 40 and 50 µm-thick slices respectively) at 4°C with secondary antibodies (see Table 2) in 0.01 M PBS (pH 7.4), 0.2% Triton X-100, and NDS (1%). Sections were washed in 0.01M PBS (pH 7.4) and incubated for 20 minutes at room temperature with 4,6-diamidino-2-phenylindole (DAPI, 1 µg/ml) to label nuclei. Sections were washed in 0.01M PBS (pH 7.4) then mounted on gelatine coated slides, air dried, and coverslipped with antifade mounting medium Mowiol (4-88 reagent, Calbiochem 475904).

Table 1. List of the primary antibodies

Antigen name	Host	Dilution	Source	Catalogue number
Primary antibodies				
COUP-TFI/Nr2f1	Rabbit	1:1000	Abcam	ab181137
COUP-TFI/Nr2f1	Rabbit	1:1000	Studer's lab	-
DCX	Goat	1:1500	Santa Cruz Biotechnology	Sc-8066
GFAP	Goat	1:2000	Abcam	ab53554
GFP	Chicken	1:1000	AvesLab	GFP-1020
Mfn2	Rabbit	1:500	Immunological Science	MAB-94608
NeuN	Mouse	1:1000	Chemicon	MAB377

Table 2. List of the secondary antibodies

Antigen name	Host	Dilution	Source	Catalogue number
Secondary antibodies				
AlexaFluor488 Anti-Ck	Donkey	1:400	Jackson ImmunoResearch	703-545-155
AlexaFluor488 Anti-Gt	Donkey	1:400	Jackson ImmunoResearch	705-545-147
AlexaFluor488 Anti-Ms	Donkey	1:400	Jackson ImmunoResearch	715-545-151
AlexaFluor647 Anti-Gt	Donkey	1:600	Jackson ImmunoResearch	705-605-147

AlexaFluor647 Anti-Ms	Donkey	1:600	Jackson ImmunoResearch	715-605-151	360
AlexaFluor647 Anti-Rb	Donkey	1:600	Jackson ImmunoResearch	711-605-152	361
Cy3 Anti-Ms	Donkey	1:800	Jackson ImmunoResearch	715-165-151	362
Cy3 Anti-Gt	Donkey	1:800	Jackson ImmunoResearch	705-165-147	363
Cy3 Anti-Rb	Donkey	1:800	Jackson ImmunoResearch	711-165-152	364

Microscope acquisition and quantifications

Images of multiple immunofluorescence on tissue sections were acquired with a TCS SP5 confocal microscope (Leica Microsystems) or a Nikon microscope coupled with a computer-assisted image analysis system (Neurolucida software, MicroBrightField). Confocal z-stacked images used for morphometric analyses on both cellular and mitochondrial architecture were captured through the thickness of the slice (50 µm) at 0.5-µm optical steps with an objective 63X/1.4 N.A. oil immersion lens with an additional zoom (2x) and a resolution of 8-bit, 1024/1024 pixels and 50Hz scan-speed (1 voxel=133.6*133.6*395.2 nm, xyz). mitoDsRed+YFP+ newborn granule neurons residing in the upper and lower blades of the DG were used for analyses only when satisfying all the following criteria: (i) to be positive for doublecortin (i.e. triple DCX+mitoDsRed+YFP+ newborn cells); (ii) to bear an apical dendrite that arborize into the molecular cell layer (MCL); (iii) to show a high and homogeneous mitoDsRed signal; and (iv) to include the whole dendritic morphology within the 50-µm section thickness. Morphometric analyses of the selected DCX+mitoDsRed+YFP+ newborn neurons was performed by using the Simple Neurite Tracer (SNT of the Neuroanatomy toolkit) plug-in on Fiji (https://fiji.sc/). Perikarya size was assessed by manually drawing the largest cross-sectional area of each cell soma based on the DCX+YFP+ double staining using the polygon selection tab in Fiji. Manual counting of dendritic mitochondria number and evaluation of their length was carried out on 2D images obtained by the Max intensity projection tool applied on the optical slices that include the whole dendritic arborization. The number of dendritic mitochondria was manually analyzed by the Cell Counter and channel tool plug-ins; measurement of the dendritic mitochondrial length was carried out by drawing the length (parallel to the dendrite) of each mitochondrion using the segmented line tool on Fiji.

Analyses of the mitochondrial mass in the soma and the primary dendrite compartments were carried out as described below. First, for each cell compartments a maximum z-projection of all the optical slices including the region of interest (ROI) was applied and the ROI based on the cytoplasmic YFP+ signal on Fiji, drawn and measured as area in μm^2. In particular, the soma area comprises the perikaryon area *plus* the cone-shaped portion of the primary dendrite directly stemming from the soma, whereas the area of the primary dendrite includes the cylindric part of primary apical dendrite to its first ramification (see **Figure S2b**). Afterwards the channel containing the mitochondrial signal (i.e., mitoDsRed) was binarized after thresholding and the resulting binary image was used to calculate the area covered by mitochondrial signal

and expressed as a percentage on the ROI area. In addition, the MiNA toolkit developed by the StuartLab (Valente *et al*,
2017) was used for the semi-automated analyses of the mitochondrial network features in the soma compartment.

To quantify Mfn2 immunostaining within the soma of DCX+YFP+ neurons, confocal image z-stacks were captured
throughout the thickness of the cell bodies of DCX+YFP+ neurons. Acquisitions were done with 0.5-µm optical step size
using an objective 63X/1.4 N.A. oil immersion lens and by adding an additional zoom (3x) with a resolution of 8-bit,
1024/1024 pixels and 50Hz scan-speed (1 voxel=120.3*120.3*395.2 xyz). Importantly, all the parameters (i.e., laser power,
gain and offset) were kept the same among acquisitions. We developed a custom-made workflow allowing an unbiased and
semi-automated quantitative analysis of Mfn2 staining coverage and intensity on triple labeled Mfn2+DCX+YFP+ neurons
on Fiji. First, a max z-projection of the optical slices containing the whole soma area was applied and the soma ROI was
manually drawn based on the cytoplasmic YFP+ signal. The obtained ROI was applied to the channel including the naïve
Mfn2 signal to calculate the Mfn2 staining intensity within the soma shaft compartment and expressed as the Corrected
Total Cell Fluorescence (CTCF) calculated by the formula CTCF = Integrated Density – (ROI X Mean fluorescence of
background readings). The background readings were calculated by averaging three mean fluorescence obtained by three
small ROI (5x5µm) without Mfn2 staining (i.e., blood vessel lumen or cell nuclei close to the cell of interest). To evaluate
the fractional area covered by Mfn2 staining in the soma area, we first applied a "pre-processing" step on the acquired
channel including the Mfn2 signal as follow: (i) unsharp mask (radius=2, mask=0.4); (ii) substract background (rolling=50);
(iii) enhance local contrast (i.e., CLAHE; blocksize=9, histogram=256, maximum=4, no mask; slow); (iv) median filtering
(radius=1). Afterwards, the channel containing the Mfn2 staining was given a threshold (by Moments algorithm) and the
binary image was cleaned outside the ROI to get the area occupied by Mfn2 staining (i.e., Mfn2 ROI) within the selected
cell. Finally, we calculated the area fraction covered by Mfn2 signal on the soma area (i.e., Mfn2 staining occupancy) and
expressed it as a percentage. The Mfn2 ROI was also applied to the naive channel containing the Mfn2 signal to calculate
the Mfn2 staining intensity both as Integrated Density (IntDens) and as corrected total mitochondrial fluorescence (CTMF)
wherein the ROI was the Mfn2 signal or calculated as the CTCF wherein the ROI was the soma area. To calculate the Mfn2
staining coverage on the DG of adult constitutive *Nr2f1-het* mice, we adopted the same strategy described above but on the
complete image.

For all the analyses the number of cells, animals, and mitochondria are described in the figure legends.

Chromatin immunoprecipitation followed by sequencing (ChIP-seq)
Adult neocortical tissue was acutely isolated from two-month-old C57BL/6J mice (n=3 males). Animals were deeply
anesthetized with an intraperitoneal injection of a mixture of tiletamine and zolazepam (40 mg/kg; Zoletil; Virbac) and
xylazine (5 mg/kg; Rompun; Bayer) and the brain removed after rapid decapitation. Neocortices from both hemispheres
were microdissected and pooled together, frozen and stored at -80°C until further analysis. Around 10 mg of neural tissue
were chopped with a scalpel, harvested in 5 ml of 0.01 M PBS (pH 7.4) and crosslinked by adding formaldehyde to a final
concentration of 1.5%, incubated for 30 min at RT on a rotator, and quenched with 0.125 M glycine for 5 min at RT. After
crosslinking, the tissue was washed twice in cold 0.01 M PBS (pH 7.4) and centrifuged at 1.000xg for 5 min at 4°C. Tissue
pellet was resuspended in 0.25ml of sodium dodecyl sulfate (SDS) lysis buffer (50 mM Tris pH 8.0, 1% SDS, 10 mM
EDTA, anti-proteases) and incubated on a rotator for 30 min at 4°C. Afterwards, sample was sonicated for 18 cycles on high

power setting (30s ON, 30s OFF) using the Bioruptor Next Gen (Diagenode) and then centrifuged at 20.000xg for 10 min at
4°C.

The isolated chromatin was diluted 10-fold with ChIP dilution buffer (16.7 mM Tris-HCl pH 8.0, 0.01% SDS, 1.1 % Triton
X-100, 1.2 mM EDTA, 167 mM NaCl) (1/10 was kept as input) and incubated with 2 μg of rabbit anti-COUP-TFI/Nr2f1
antibody from Studer's lab (Tripodi *et al*, 2004) overnight at 4°C on a rotator. Protein G-conjugated magnetic beads (Dynal,
Thermo Fisher Scientific) were saturated with PBS/1% BSA overnight at 4°C. Next day, samples were incubated with
saturated beads for two hours at 4°C on a rotator, and subsequently washed with 1 ml of cold Low salt buffer (20 mM Tris-
HCl pH 8.0, 0.1 % SDS, 1% Triton X-100, 2 mM EDTA, 150 mM NaCl), 1 ml of cold High salt buffer (20 mM Tris-HCl
pH 8.0, 0.1 % SDS, 1% Triton X-100, 2 mM EDTA, 500 mM NaCl), 1 ml of cold LiCl buffer (10 mM Tris-HCl pH 8.0, 1%
DOC, 250 mM LiCl, 1 mM EDTA, 1% NP-40), and twice with 1 ml of cold TE buffer (10 mM Tris-HCl pH 8.0, 1 mM
EDTA). The immunoprecipitated chromatin was eluted with 200 μl of Elution buffer (10 mM Tris-HCl pH 8.0, 1 mM
EDTA, 1% SDS, 150 mM NaCl, 5mM DTT) for 30 min at RT on a rotator, and decrosslinked at 65°C overnight. The
decrosslinked DNA was purified using QIAquick PCR Purification Kit (Qiagen) according to the manufacture's instruction.
For genome-wide analysis of binding, sequencing libraries were constructed using the NEBNext ChIP-seq Library Prep
Reagent Set for Illumina and a NextSeq 500 Illumina sequencer (New England BioLabs Inc.).

Bioinformatic analyses

The reads from sequencing were mapped to the mouse genome (mm9 assembly) using Bowtie version 0.12.7, reporting
only unique hits with up to two mismatches. The redundant reads were collapsed, and peak calling was performed using
MACS version 1.4.1 (Zhang *et al*, 2008) with normalization for IgG ChIP at a fixed p-value $=$ 1E-8. Homer version 4.11
was used to perform de-novo motif discovery and peak annotation. A ChIP-seq peak was considered associated with a gene
if it showed an overlap of -3kb/+2kb region around a transcription start site (TSS) of the gene. Gene Ontology (GO)
enrichment analysis was performed using PANTHER Overrepresentation Test version 14 (Mi *et al*, 2019).

We performed a pre-rank Gene Set Enrichment Analysis (GSEA) using as "rank value" the DeSeq2 statistic of genes
expression in the hippocampi of adult *Nr2f1-het vs.* WT mice obtained by bulk RNA-seq by Chen and colleagues (Chen *et*
al, 2020), and as "pathway" the genes bound on the promoter by Nr2f1.

Statistics

Data are derived from at least three different animals/group. A two-tailed unpaired t-test was performed to compare the
differences between two groups, moreover, an F-test of equality of variances was conducted to compare variances, and
Welch's correction was applied in case of unequal variance distribution. Data distribution was tested to be normal with
either the Shapiro-Wilk test (alpha=0.05; if $3<n<50$) or D'Agostino-Pearson (if $n>50$). When comparing two populations of
data, two-tailed unpaired Student's t test was used to calculate statistical significance in cases of Gaussian distribution;
otherwise, the non-parametric Mann-Whitney test was used. For Sholl analysis, two-way repeated measures ANOVA with
Sidak's post-hoc test was used. The confidence interval was expressed with 95% confidence. For the analysis of normalized
data (i.e., fold change) a Kruskal-Wallis test followed by Dunn's post-hoc test was used. Kolmogorov-Smirnov test was
carried out to compare cumulative distributions. The statistical significance was defined as follows: *p<0.05, **p<0.01, and
***p<0.001. All statistical analyses were performed using Graphpad Prism 8 software (version 8.0.2; GraphPad Software,

San Diego California USA, www.graphpad.com). Fisher's exact test, with the Benjamini–Hochberg False Discovery Rate

(FDR) correction for multiple testing, has been added as the default algorithm for the over-representation test for GO

analyses (Mi *et al*, 2019).

Acknowledgements

478 This work received the financial support of the Jerome Lejeune Foundation (Cycle 2020a – Project #1977) and was

479 supported by the ex60% (University of Turin) to S. De Marchis and the Foundation Cassa Risparmio Torino (CRT –

480 2019/2254) to S. Bovetti. Postdoctoral fellowships to S. Bonzano were funded by: Compagnia San Paolo (Turin) "Bando

481 per l'internazionalizzazione della ricerca"; Fondazione Umberto Veronesi (Milan); the Accademia Nazionale dei Lincei

482 "Centro Linceo Interdisciplinare Beniamino Segre" (Rome).

Author Information

References

Aimone JB, Li Y, Lee SW, Clemenson GD, Deng W & Gage FH (2014) Regulation and function of adult neurogenesis:

from genes to cognition. *Physiol Rev* 94: 991–1026

Alfano C & Studer M (2013) Neocortical arealization: Evolution, mechanisms, and open questions. *Dev Neurobiol* 73: 411–

447

Armentano M, Chou SJ, Srubek Tomassy G, Leingärtner A, O'Leary DDM & Studer M (2007) COUP-TFI regulates the

balance of cortical patterning between frontal/motor and sensory areas. *Nat Neurosci* 10: 1277–1286

Artegiani B, Lyubimova A, Muraro M, van Es JH, van Oudenaarden A & Clevers H (2017) A Single-Cell RNA Sequencing

Study Reveals Cellular and Molecular Dynamics of the Hippocampal Neurogenic Niche. *Cell Rep* 21: 3271–3284

Beckervordersandforth R (2017) Mitochondrial Metabolism-Mediated Regulation of Adult Neurogenesis. *Brain Plast* 3:

73–87

Beckervordersandforth R, Ebert B, Schäffner I, Moss J, Fiebig C, Shin J, Moore DL, Ghosh L, Trinchero MF, Stockburger

C, *et al* (2017) Role of Mitochondrial Metabolism in the Control of Early Lineage Progression and Aging Phenotypes

in Adult Hippocampal Neurogenesis. *Neuron* 93: 560-573.e6

Bertacchi M, Gruart A, Kaimakis P, Allet C, Serra L, Giacobini P, Delgado☐García JM, Bovolenta P & Studer M (2019a)

Mouse Nr2f1 haploinsufficiency unveils new pathological mechanisms of a human optic atrophy syndrome . *EMBO*

Mol Med 11: 1–18

Bertacchi M, Parisot J & Studer M (2019b) The pleiotropic transcriptional regulator COUP-TFI plays multiple roles in

neural development and disease. *Brain Res* 1705: 75–94

Billiet B, Amati☐Bonneau P, Desquiret☐Dumas V, Guehlouz K, Milea D, Gohier P, Lenaers G, Mirebeau☐Prunier D, den

Dunnen JT, Reynier P, *et al* (2021) NR2F1 database: 112 variants and 84 patients support refining the clinical

synopsis of Bosch–Boonstra–Schaaf optic atrophy syndrome . *Hum Mutat*

Bonzano S, Crisci I, Podlesny-Drabiniok A, Rolando C, Krezel W, Studer M & De Marchis S (2018) Neuron-Astroglia Cell

Fate Decision in the Adult Mouse Hippocampal Neurogenic Niche Is Cell-Intrinsically Controlled by COUP-TFI In

Vivo. *Cell Rep* 24: 329–341

Bosch DGM, Boonstra FN, Gonzaga-Jauregui C, Xu M, De Ligt J, Jhangiani S, Wiszniewski W, Muzny DM, Yntema HG,

Pfundt R, *et al* (2014) NR2F1 mutations cause optic atrophy with intellectual disability. *Am J Hum Genet* 94: 303–

309

Bosch DGM, Boonstra FN, De Leeuw N, Pfundt R, Nillesen WM, De Ligt J, Gilissen C, Jhangiani S, Lupski JR, Cremers

FPM, *et al* (2016) Novel genetic causes for cerebral visual impairment. *Eur J Hum Genet* 24: 660–665

Calvo SE, Clauser KR & Mootha VK (2016) MitoCarta2.0: An updated inventory of mammalian mitochondrial proteins.

Nucleic Acids Res 44: D1251–D1257

Chan DC (2020) Mitochondrial Dynamics and Its Involvement in Disease. *Annu Rev Pathol Mech Dis* 15: 235–259

Chen CA, Bosch DGM, Cho MT, Rosenfeld JA, Shinawi M, Lewis RA, Mann J, Jayakar P, Payne K, Walsh L, *et al* (2016)

The expanding clinical phenotype of Bosch-Boonstra-Schaaf optic atrophy syndrome: 20 new cases and possible

genotype-phenotype correlations. *Genet Med* 18: 1143–1150

Chen CA, Wang W, Pedersen SE, Raman A, Seymour ML, Ruiz FR, Xia A, Van Der Heijden ME, Wang L, Yin J, *et al*

(2020) Nr2f1 heterozygous knockout mice recapitulate neurological phenotypes of Bosch-Boonstra-Schaaf optic

atrophy syndrome and show impaired hippocampal synaptic plasticity. *Hum Mol Genet* 29: 705–715

Course MM & Wang X (2016) Transporting mitochondria in neurons. *F1000Research* 5: 1735

Faits MC, Zhang C, Soto F & Kerschensteiner D (2016) Dendritic mitochondria reach stable positions during circuit

development. *Elife* 5: 1–17

Fang D, Yan S, Yu Q, Chen D & Yan SS (2016) Mfn2 is required for mitochondrial development and synapse formation in

human induced pluripotent stem cells/hiPSC derived cortical neurons. *Sci Rep* 6: 1–13

Foglio B, Rossini L, Garbelli R, Regondi MC, Mercurio S, Bertacchi M, Avagliano L, Bulfamante G, Coras R, Maiorana A,

et al (2021) Dynamic expression of NR2F1 and SOX2 in developing and adult human cortex: comparison with

cortical malformations. *Brain Struct Funct* 226: 1303–1322

Frye RE (2020) Mitochondrial Dysfunction in Autism Spectrum Disorder: Unique Abnormalities and Targeted Treatments.

Semin Pediatr Neurol 35: 100829

Giacomello M, Pyakurel A, Glytsou C & Scorrano L (2020) The cell biology of mitochondrial membrane dynamics. *Nat*

Rev Mol Cell Biol 21: 204–224

Hobbs MM, Wolters WC, Rayapati AO & Kahn DA (2020) Bosch-Boonstra-Schaaf optic atrophy syndrome presenting as

new-onset psychosis in a 32-year-old man: A case report and literature review. *J Psychiatr Pract* 26: 58–62

Jagasia R, Steib K, Englberger E, Herold S, Faus-Kessler T, Saxe M, Gage FH, Song H & Lie DC (2009) GABA-cAMP

response element-binding protein signaling regulates maturation and survival of newly generated neurons in the adult

hippocampus. *J Neurosci* 29: 7966–7977

Jurcau A (2021) Insights into the pathogenesis of neurodegenerative diseases: Focus on mitochondrial dysfunction and

oxidative stress. *Int J Mol Sci* 22

Jurkute N, Bertacchi M, Arno G, Tocco C, Kim US, Kruszewski AM, Avery RA, Bedoukian EC, Han J, Ahn SJ, *et al*

(2021) Pathogenic NR2F1 variants cause a developmental ocular phenotype recapitulated in a mutant mouse model.
Brain Commun 3: 1–21

Kaiwar C, Zimmermann MT, Ferber MJ, Niu Z, Urrutia RA, Klee EW & Babovic-Vuksanovic D (2017) Novel NR2F1
variants likely disrupt DNA binding: molecular modeling in two cases, review of published cases, genotype-
phenotype correlation, and phenotypic expansion of the Bosch-Boonstra-Schaaf optic atrophy syndrome. *Cold Spring*
Harb Mol case Stud 3

Khacho M, Clark A, Svoboda DS, Azzi J, MacLaurin JG, Meghaizel C, Sesaki H, Lagace DC, Germain M, Harper ME, *et*
al (2016) Mitochondrial Dynamics Impacts Stem Cell Identity and Fate Decisions by Regulating a Nuclear
Transcriptional Program. *Cell Stem Cell* 19: 232–247

Khacho M, Harris R & Slack RS (2019) Mitochondria as central regulators of neural stem cell fate and cognitive function.
Nat Rev Neurosci 20: 34–48

Kimura T & Murakami F (2014) Evidence that dendritic mitochondria negatively regulate dendritic branching in pyramidal
neurons in the neocortex. *J Neurosci* 34: 6938–6951

Lenaers G, Neutzner A, Le Dantec Y, Jüschke C, Xiao T, Decembrini S, Swirski S, Kieninger S, Agca C, Kim US, *et al*
(2021) Dominant optic atrophy: Culprit mitochondria in the optic nerve. *Prog Retin Eye Res* 83: 100935

Liang G, Lin JCY, Wei V, Yoo C, Cheng JC, Nguyen CT, Weisenberger DJ, Egger G, Takai D, Gonzales FA, *et al* (2004)
Distinct localization of histone H3 acetylation and H3-K4 methylation to the transcription start sites in the human
genome. *Proc Natl Acad Sci U S A* 101: 7357–7362

Martín-Hernández E, Rodríguez-García ME, Chen CA, Cotrina-Vinagre FJ, Carnicero-Rodríguez P, Bellusci M, Schaaf CP
& Martínez-Azorín F (2018) Mitochondrial involvement in a Bosch-Boonstra-Schaaf optic atrophy syndrome patient
with a novel de novo NR2F1 gene mutation. *J Hum Genet* 63: 525–528

Mi H, Muruganujan A, Huang X, Ebert D, Mills C, Guo X & Thomas PD (2019) Protocol Update for large-scale genome
and gene function analysis with the PANTHER classification system (v.14.0). *Nat Protoc* 14: 703–721

Misgeld T & Schwarz TL (2017) Mitostasis in Neurons: Maintaining Mitochondria in an Extended Cellular Architecture.
Neuron 96: 651–666

Montemayor C, Montemayor OA, Ridgeway A, Lin F, Wheeler DA, Pletcher SD & Pereira FA (2010) Genome-wide
analysis of binding sites and direct target genes of the orphan nuclear receptor NR2F1/COUP-TFI. *PLoS One* 5

Monzio Compagnoni G, Di Fonzo A, Corti S, Comi GP, Bresolin N & Masliah E (2020) The Role of Mitochondria in
Neurodegenerative Diseases: the Lesson from Alzheimer's Disease and Parkinson's Disease. *Mol Neurobiol* 57:
2959–2980

Qiu Y, Krishnan V, Zeng Z, Gilbert DJ, Copeland NG, Gibson L, Yang-Feng T, Jenkins NA, Tsai MJ & Tsai SY (1995)
Isolation, characterization, and chromosomal localization of mouse and human COUP-TF I and II Genes. *Genomics*
29: 240–246 doi:10.1006/geno.1995.1237 [PREPRINT]

Rada-Iglesias A, Bajpai R, Prescott S, Brugmann SA, Swigut T & Wysocka J (2012) Epigenomic annotation of enhancers
predicts transcriptional regulators of human neural crest. *Cell Stem Cell* 11: 633–648

Rangaraju V, Lauterbach M & Schuman EM (2019a) Spatially Stable Mitochondrial Compartments Fuel Local Translation
during Plasticity. *Cell* 176: 73-84.e15

Rangaraju V, Lewis TL, Hirabayashi Y, Bergami M, Motori E, Cartoni R, Kwon SK & Courchet J (2019b) Pleiotropic

mitochondria: The influence of mitochondria on neuronal development and disease. *J Neurosci* 39: 8200–8208

Rech ME, McCarthy JM, Chen CA, Edmond JC, Shah VS, Bosch DGM, Berry GT, Williams L, Madan-Khetarpal S,

Niyazov D, *et al* (2020) Phenotypic expansion of Bosch–Boonstra–Schaaf optic atrophy syndrome and further

evidence for genotype–phenotype correlations. *Am J Med Genet Part A* 182: 1426–1437

Rojas-Charry L, Nardi L, Methner A & Schmeisser MJ (2021) Abnormalities of synaptic mitochondria in autism spectrum

disorder and related neurodevelopmental disorders. *J Mol Med* 99: 161–178

Steib K, Schäffner I, Jagasia R, Ebert B & Chichung Lie D (2014) Mitochondria modify exercise-induced development of

stem cell-derived neurons in the adult brain. *J Neurosci* 34: 6624–6633

Tocco C, Bertacchi M & Studer M (2021) Structural and Functional Aspects of the Neurodevelopmental Gene NR2F1:

From Animal Models to Human Pathology. *Front Mol Neurosci* 14

Tripodi M, Filosa A, Armentano M & Studer M (2004) The COUP-TF nuclear receptors regulate cell migration in the

mammalian basal forebrain. *Development* 131: 6119–6129

Valente AJ, Maddalena LA, Robb EL, Moradi F & Stuart JA (2017) A simple ImageJ macro tool for analyzing

mitochondrial network morphology in mammalian cell culture. *Acta Histochem* 119: 315–326

Valenti D, de Bari L, De Filippis B, Henrion-Caude A & Vacca RA (2014) Mitochondrial dysfunction as a central actor in

intellectual disability-related diseases: An overview of Down syndrome, autism, Fragile X and Rett syndrome.

Neurosci Biobehav Rev 46: 202–217

Wu SP, Lee DK, DeMayo FJ, Tsai SY & Tsai MJ (2010) Generation of ES cells for conditional expression of nuclear

receptors and coregulators in vivo. *Mol Endocrinol* 24: 1297–1304

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nussbaum C, Myers RM, Brown M, Li W, *et al*

(2008) Model-based analysis of ChIP-Seq (MACS). *Genome Biol* 9

Zhao C, Teng EM, Summers RG, Ming GL & Gage FH (2006) Distinct morphological stages of dentate granule neuron

maturation in the adult mouse hippocampus. *J Neurosci* 26: 3–11

Zsurka G & Kunz WS (2015) Mitochondrial dysfunction and seizures: The neuronal energy crisis. *Lancet Neurol* 14: 956–

966

Figures Legends

Figure 1. ChIP-seq analysis on adult mouse neocortex revealed direct binding of Nr2f1 on mitochondrial genes. a.

Example of a genomic peak on a promoter showing Nr2f1 binding associated to the H3K4me3 (gene on the left) compared

with a promoter region showing H3K4me3 without Nr2f1 binding (gene on the right). **b.** Pie chart illustrating the relative

distribution (expressed as percentage) of the 2119 Nr2f1-bound sequences. **c.** De novo motif discovery of TF motif in Nr2f1

binding sites (forward) – see Figure S1e for matches to known motifs. **d.** Bar graph illustrating gene ontology (GO) term

analysis wherein the 2119 genes bound by Nr2f1 were ranked for cellular components (top) and biological processes

(bottom). The full lists of enriched categories are listed in Supplementary Table S1.

Figure 2. Nr2f1 LOF leads to decreased mitochondrial mass in the soma and proximal dendritic compartment of adult-born hippocampal neurons *in vivo*. a. Overview of the Cre-mediated gene rearrangements of Rosa26 (R26) and Nr2f1 loci in *Glast::CreERT2;R26-floxed STOP-Nr2f1fl/fl* mice. **a'.** Experimental design. **b.** Representative confocal images of adult-born neurons transduced with the RV-mitoDsRed (red) and immunostained for YFP (green) and Nr2f1 (magenta) in *Ctrl* and *Nr2f1-icKO* mice. Nuclei are counterstained with DAPI (white). **c.** Histogram reporting data validating Nr2f1 loss in the Glast-lineage. While in control DG, the vast majority of YFP+ and mitoDsRed+YFP+ cells showed nuclei that are positive for Nr2f1 (Nr2f1+), virtually all YFP+ and mitoDsRed+YFP+ cells in mutant DG are negative for Nr2f1. Nr2f1+ on total YFP+ cells: n=229/230 cells out of 3 *Ctrl* animals; N=5/340 cells out of 3 *Nr2f1-icKO* animals; Nr2f1+ on mitoDsRed+YFP+ cells: n=16/16 cells out of 3 *Ctrl* animals; n=0/20 cells out of 3 *Nr2f1-icKO* animals. **d.** Representative confocal images (max projection) showing the mitoDsRed+ mitochondria (red) within the soma area (solid lines) and primary dendrite (dotted lines) in control and *Nr2f1-icKO* mitoDsRed+DCX+YFP+ newborn neurons. **e,f.** Graph reporting the mitochondrial occupancy expressed as the percentage of the soma area (e) or of the primary dendrite area (f) covered by the mitoDsRed signal. Mann-Whitney test, p=0.0066 (e); p=0.0001 (f). For the analyses reported in e and f, n=13 (*Ctrl*) and 16 (*Nr2f1-icKO*) and n=14 (*Ctrl*) and 18 (*Nr2f1-icKO*) cells out of 3 animals per genotypes and at least 3 cells/animals were used. Data are shown as mean±SD. Each dot represents a cell. Scale bars: b, 20 µm; b (insets), 10 µm; d, 10 µm. *p<0.05, **p<0.01, ***p<0.001.

Figure 3. Alteration in dendritic mitochondrial number, architecture and mass in newborn neurons lacking Nr2f1.
a,b. Representative confocal images of two recombined and mitoDsRed transduced (red) adult-born neurons immunolabelled with doublecortin (DCX, white) in the DG of both *Ctrl* (a) and *Nr2f1-icKO* (b). **a',b'.** Representative drawing reporting the position of single mitochondrion (red dots) on the dendritic arborization of the two cells shown in a-b boxes. **c.** Histogram reporting the total number of mitochondria present in each dendritic arborization of the analyzed mitoDsRed+DCX+YFP+ cells. Mann-Whitney, p=0.0058. **d.** Histogram showing the linear density of dendritic mitochondria obtained by normalizing the total number of dendritic mitochondria to the total dendritic arborization length. Unpaired Student's t test, p=0.0001. **e.** Graph showing the relationship between the dendritic arborization length (x axis) and the total number of dendritic mitochondria (y axis) per each mitoDsRed+DCX+YFP+ cell analyzed; linear regression with Pearson's correlation was performed with R^2, p-values and best fitted line equation displayed. Note the shift of the regression fitted line of *Nr2f1-icKO* cells without changes in its steepness compared to the *Ctrl* fitted line. **F.** Representative confocal images showing high magnification of the mitochondria in comparable level of dendrites in *Ctrl* and *Nr2f1-icKO* mitoDsRed+DCX+YFP+ adult-born neurons. **G.** Quantification of dendritic mitochondrial length in control and Nr2f1-depleted adult-born hippocampal neurons. Each data point represents an individual mitochondrial size; n=459 mitochondria out of 13 control cells, n=279 mitochondria out of 14 *Nr2f1-icKO* cells; Mann Whitney test, p<0.0001. **h.** Cumulative frequency distribution for the length of dendritic mitochondria in control vs. Nr2f1-ablated mitoDsRed+DCX+YFP+ adult-born neurons. Kolmogorov-Smirnov test, p<0.0001. **i.** Quantification of the distribution of dendritic mitochondrial morphologies into the four different subclasses based on their length (i.e., fragmented, <1 µm; elongated, from 1 to 2 µm; medium/large, from 2 to 3 µm; tubular, > 3 µm). Mann Whitney test: fragmented, p=0.001; elongated, p=0.6760; medium/large, p= 0.0917; tubular, p<0.0001. Each dot represents a cell. **j.** Graph reporting the mitochondrial occupancy expressed as the percentage of dendritic length covered by the mitoDsRed staining. Mann Whitney, p<0.0001. For all the

659 analyses, n=13 (*Ctrl*) and n=14 (*Nr2f1-icKO*) cells out of 3 animals per genotype and at least 3 cells/animal were used for

660 each analysis. Data are shown as: mean±SD (c,d,i,j); best regression fitted line in black, the band represent the 95%

661 confidence interval and each dot represents a cell (e); box and whiskers plot with median (middle line), upper/lower

662 quartiles, and error bars ranging from min to max values (g). Scale bars: a,b,a',b', 20 µm; f,i, 5 µm. *p<0.05, **p<0.01,

663 ***p<0.001.

665 **Figure 4. The fusion factor Mfn2 as a direct target for Nr2f1 transcriptional activity. a.** Schematic illustration showing

666 the main factors involved in the mitochondrial dynamics (fission in orange; fusion in light blue) in mammalian cells: among

667 them, those underlined in yellow show specific Nr2f1 binding on their promoter regions as revealed by peak calling analysis

668 on ChIP-seq data. Sequence tag accumulation of Nr2f1 ChIP-seq of around the TSS regions of Dmn2, Smcr7l and Mfn2 are

669 shown. **b.** Gene Set Enrichment Analysis (GSEA) revealed Mfn2 among the genes encoding mitochondrial proteins bound

670 by Nr2f1 and deregulated in constitutive heterozygous (*Nr2f1-het)* hippocampi. **c.** Representative confocal images showing

671 Mfn2 staining within the DG of adult wildtype (*WT*) and *Nr2f1-het* mice. Insets show higher magnifications, the binarized

672 images used to calculate the area occupied by Mfn2 staining and heat-map images of Mfn2+ signal in WT and Nr2f1-het

673 DG. **d,e.** Graphs reporting the area occupied by Mfn2+ signal (d) and the intensity of Mfn2 protein staining (e) in *WT* and

674 *Nr2f1-het* DG. Mann Whitney test: p=0.0095 (d) and unpaired Student's t-test p<0.0001 (e). N=4 *WT* and 6 *Nr2f1-het* mice.

675 **f.** Experimental design. **g.** Representative confocal images of the DG immunostained for YFP (green), Mfn2 (magenta),

676 DCX (white) and counterstained with DAPI (blue) in *Ctrl* (top) and *Nr2f1-icKO* (bottom) mice one months after tamoxifen

677 administration. **h.** Representative binarized images showing the area occupied by Mfn2 staining within the soma area in one

678 control (top) and one Nr2f1-depleted (bottom) recombined newborn neurons (i.e. DCX+YFP+). **i.** Graph reporting the

679 mitochondrial Mfn2 occupancy expressed as the percentage of the soma area covered by the Mfn2+ signal; Mann-Whitney

680 test, p <0.0001; n=45 cells out of 3 Ctrl mice and n=45 cells out of 3 *Nr2f1-icKO* mice. **j.** Representative heat-map images

681 of Mfn2+ signal in control (top) and Nr2f1-deficient (bottom) recombined newborn neurons. **k.** Graph reporting the

682 mitochondrial Mfn2 staining intensity normalized on Mfn2 area residing in the cell compartment comprising the soma area

683 (expressed as CTMF, cell threshold mitochondrial fluorescence); Unpaired t-test, p<0.0001; n=45 cells out of 3 Ctrl mice

684 and n=45cells out of 3 *Nr2f1-icKO* mice. Data are shown as box and whiskers plot with median (middle line), upper and

685 lower quartile, and the error bars ranging from min to max values. Each dot represents an animal (d,e) or a cell (i,k). c, 50

686 µm (low magnification) and 10 µm (higher magnification). Scale bars: g,h,j, 10 µm; ***p<0.0001.

Figure 1

Figure 2

Figure 3

Figure 4

Supplemental Figures with Legends: S1, S2, S3

Figure S1 - Related to Figure 1. a,b. Representative confocal images showing Nr2f1 immunofluorescence in the neuronal population within the adult mouse neocortex., The vast majority of Nr2f1+ cells belongs to the neuronal population, as revealed by the high colocalization of Nr2f1 immunofluorescence with the neuronal marker NeuN (a) and no co-expression within the recombined astrocytic population labelled by YFP in *Glast::CreERT2;Rosa26-lox-stop-lox-YFP* mice following a chase of two weeks after tamoxifen administration (b). **c.** Graph reporting the logarithmic ratio between the observed genomic Nr2f1-bound peaks versus the expected ones. Numbers in each line represent the quantity of annotated

peaks/annotation category. **d.** Raw data showing the *de novo* motif revealed by ChIP-seq for Nr2f1 in the adult brain and one of the best matches with known ChIP-seq database. Scale bars: a,b, 50 µm (low magnification); b, 10 µm (higher magnification with resliced).

Figure S2 - Related to Figure 2 and 3. a. Representative reconstruction of triple labelled mitoDsRed+DCX+YFP+ DG newborn neurons at 17dpi and used for the analyses. **a'.** Histogram reporting no difference in the perikarya of control vs. Nr2f1-deficient mitoDsRed+DCX+YFP+ newborn neurons. Unpaired Student's t-test. **a'',a'''.** Histograms showing the total dendritic length (a''; unpaired Student's t-test p=0.0296) and the number of dendritic branches (a'''; Mann-Whitney t-test p=0.0153) in Nr2f1 deficient mitoDsRed+DCX+YFP+ newborn neurons and control neurons. For all the analyses in a'-a''', n=13 and n=14 cells out of 3 animals per genotypes and at least 3 cells/animals were used for each analysis. **b.** Drawing showing the ROIs used for the quantifications of mitochondrial occupancy and quantified in b', b'' and b'''. **b'-b'''.** Quantification of the morphometric parameters used to calculate the mitochondrial occupancy. Student's unpaired t test, exact p-values in the graphs. n=13 *Ctrl* and n=14 *Nr2f1-icKO* cells out of 3 animals per genotype and at least 4 cells/animal. Data are shown as mean±SD. **c.** Representative binarized images showing the mitoDsRed+ mitochondria (black) within the soma (c) in control and Nr2f1-deficient mitoDsRed+DCX+YFP+ newborn neurons. **d.** Graphs showing the results obtained by applying the MiNA toolkit on the mitoDsRed signal within the soma compartment of control and *Nr2f1-icKO* newborn mitoDsRed+DCX+YFP+ neurons. From the left to the right: unpaired Student's t-test with Welch's correction, p=0.0061; unpaired Student's t-test, p=0.0465; Mann-Whitney test, p=0.0248; Mann-Whitney test, p=0.0341. Data are shown as box and whiskers plot with median (middle line), upper and lower quartile, and the error bars ranging from min to max values. A coherent reduction in the parameter of mitochondrial footprint (i.e. the total area consumed by mitochondrial signal after being separated from the background) with a coincident decrease in the length of branches (i.e. mean length of all the lines used to represent the mitochondrial structures) and in the summed branch length (i.e. the mean of the sum of the lengths of branches for each independent mitochondrial structure divided by the number of independent mitochondrial structures) as well as in the number of network branches (i.e. the mean number of attached lines used to represent each mitochondrial structure) was found in *Nr2f1-icKO* neurons compared to controls. **e.** Representative binarized images showing the mitoDsRed+ mitochondria (black) within the primary dendrite in control and Nr2f1-deficient mitoDsRed+DCX+YFP+ newborn neurons. **f.** Histogram showing the reduction (expressed as fold change) in the mitochondrial mass within the different subcellular compartments in Nr2f1-depleted mitoDdRed+DCX+YFP+ cells compared to the one measured in the compartments of the control cells. Statistics (asterisks) were obtained by Mann Whitney test of fold change values obtained from Nr2f1-icKO cells and normalized to control values (Nr2f1-icKO vs. Ctrl): soma, p=0.0047; primary dendrite, p=0.0002; dendritic arborization, p<0.0001. Each dot represents a cell. Scale bars: a, 30 µm; c,e, 10 µm. *p<0.05, **p<0.01, ***p<0.001.

Figure S3. Nr2f1 overexpression leads to increased mitochondrial content in adult-born DG neurons. a. Overview of the Cre-mediated gene rearrangements of Rosa26 loci and the transgene *CAG-loxP-stop-LoxP-hNR2F1*. Experimental design is the same used in Figure 2A'. **b.** Histogram showing the linear density of dendritic mitochondria obtained by normalizing the total number of dendritic mitochondria to the total dendritic arborization length in both Ctrl and Nr2f1-O/E newborn neurons. Unpaired Student's t test, p=0.4434. **d.** Cumulative frequency distribution for the length of dendritic mitochondria in control vs. Nr2f1 overexpressing mitoDsRed+DCX+YFP+ adult-born neurons. Kolmogoror-Smirnov test, p<0.0002. **d.** Graph reporting the mitochondrial occupancy expressed as the percentage of dendritic length covered by the mitoDsRed signal. Mann Whitney test, p=0.0301. Data are shown as mean±SD. Each dot represents a cell. Data are shown as mean±SD. *p<0.05.