1 **Domestication in dry-cured meat *Penicillium* fungi: convergent specific phenotypes and**

2 **horizontal gene transfers without strong genetic subdivision**

Running title: Footprints of domestication in dry-cured meat *Penicillium* fungi

Ying-Chu Lo†[1], Jade Bruxaux†[1,2], Ricardo C Rodríguez de la Vega[1], Samuel O'Donnell[1],

Alodie Snirc[1], Monika Coton[3], Mélanie Le Piver[4], Stéphanie Le Prieur[1], Daniel Roueyre[4],

Joëlle Dupont[5], Jos Houbraken[6], Robert Debuchy[7], Jeanne Ropars[1], Tatiana Giraud*[1] and

Antoine Branca*[8]

[1]IDEEV - Laboratoire Ecologie, Systématique et Evolution, CNRS, AgroParisTech,

Université Paris Saclay, 12 route 128, 91198 Gif sur Yvette, France

[2]Department of Ecology and Environmental Science, UPSC, Umeå University, 90187 Umeå,

Sweden

[3]Univ Brest, Laboratoire Universitaire de Biodiversité et Ecologie Microbienne, 29280

Plouzané, France

[4]Laboratoire Interprofessionnel de Production - SAS L.I.P, 34 Rue de Salers, 15000 Aurillac,

France

[5]Origine, Structure, Evolution de la Biodiversité, UMR 7205 CNRS-MNHN, Muséum

National d'Histoire Naturelle, CP39, 57 rue Cuvier, 75231 Paris Cedex 05, France

[6]Westerdijk Fungal Biodiversity Institute, Uppsalalaan 8, 3584 CT Utrecht, the Netherlands

[7]Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC),

91198, Gif-sur-Yvette, France.

[8]IDEEV - Laboratoire Evolution, Génomes Comportement, Ecologie, CNRS Université Paris

Saclay UMR 9191, IRD UMR 247, 12 route 128, 91198 Gif sur Yvette, France

†These authors contributed equally

*These authors jointly supervised the study

Corresponding author: Jade Bruxaux bruxaux.jade@gmail.com

Abstract:

Some fungi have been domesticated for food production, with genetic differentiation between

populations from food and wild environments, and food populations often acquiring

beneficial traits through horizontal gene transfers (HGTs). Studying their adaptation to

human-made substrates are of fundamental and applied importance, for understanding

adaptation processes and for further strain improvement. We studied here the population

structures and phenotypes of two distantly related *Penicillium* species used for dry-cured meat

production, *P. nalgiovense*, the most common species in the dry-cured meat food industry,

and *P. salamii*, used locally by farms. Both species displayed low genetic diversity, lacking

differentiation between strains isolated from dry-cured meat and those from other

environments. Nevertheless, the strains collected from dry-cured meat within each species

displayed slower proteolysis and lipolysis than their wild conspecifics, and those of *P.*

nalgiovense were whiter. Phenotypically, the non-dry-cured meat strains were more similar to

their sister species than to their conspecific dry-cured meat strains, indicating an evolution of

specific phenotypes in dry-cured meat strains. A comparison of available *Penicillium*

genomes from various environments revealed HGTs, particularly between *P. nalgiovense* and

P. salamii (representing almost 1.5 Mb of cumulative length). HGTs additionally involved *P.*

biforme, also found in dry-cured meat products. We further detected positive selection based

on amino-acid changes. Our findings suggest that selection by humans has shaped the *P.*

salamii and *P. nalgiovense* populations used for dry-cured meat production, which constitutes

domestication. Several genetic and phenotypic changes were similar in *P. salamii*, *P.*

nalgiovense, and *P. biforme*, indicating convergent adaptation to the same human-made

environment. Our findings have implications for fundamental knowledge on adaptation and

for the food industry: the discovery of different phenotypes and of two mating types paves the

way for strain improvement by conventional breeding, to elucidate the genomic bases of

beneficial phenotypes and to generate diversity.

Keywords: fungi, parallel adaptation, convergence, sausage, domestication

Introduction

The way in which populations adapt to their environment is a fundamental issue in biology.

Domestication is widely used as a model for understanding adaptation as it corresponds to a

recent adaptation process caused by strong, human-driven selection (Darwin, 1859, 1868).

Humans have grown and bred various organisms, choosing individuals with the most

interesting features for their particular uses or for consumption. For instance, many

morphological traits differ between domesticated maize and its wild ancestor, including floral

morphology, seed size, dispersal ability, stem number and length (Hufford et al., 2012; Stitzer

& Ross-Ibarra, 2018). Charles Darwin used domesticated pigeons as a model for investigating

adaptation and natural selection processes (Darwin, 1859, 1868). The parallel adaptation of

distantly related species to the same environment provides an excellent study model for

investigating whether evolution is repeatable — i.e., whether independent adaptation events to

the same ecological niche involve similar phenotypic and genomic changes — a situation

known as convergence (Chan et al., 2010; Colosimo et al., 2004; Duboué et al., 2011). In

addition to adaptive changes, domesticated populations may suffer from bottlenecks, in which

a drastic loss of genetic diversity occurs due to the strong selection exerted by humans. Such

bottlenecks may jeopardize the ability of the domesticated populations to adapt (Maruyama &

Fuerst, 1985; Nei et al., 1975). Studying domesticated organisms also has applied importance,

for understanding the genomic bases of desired traits, for further variety generation and for

maintaining genetic diversity that could allow future improvement.

Fungi are good models for studies of adaptation using domestication. Their experimental

tractability makes it possible to perform fitness tests in controlled environments, and their

small and compact genomes facilitate genomic studies to elucidate the genetic basis of

adaptation (Gibbons et al., 2012; Gladieux et al., 2013; Legras et al., 2018; Ropars et al., 2015;

Watarai et al., 2019; Wolfe & Dutton, 2015). Many fungi have been domesticated for food

production (Dupont et al., 2017; Steensels et al., 2019), the chief examples being

Saccharomyces, Aspergillus and *Penicillium. Saccharomyces cerevisiae* is the most studied

domesticated fungal species, with different populations used for winemaking, baking and

brewing (Almeida et al., 2014; Gallone et al., 2016; C. Gonçalves et al., 2018; Legras et al.,

2018; Peter et al., 2018). Fungi are also used for cheese ripening. For example, *P. roqueforti*

is used for blue-veined cheeses and *P. camemberti* is used for soft cheeses, such as Brie and

Camembert. Both fungi have recently emerged as useful models for adaptation and

domestication studies (Bodinaku et al., 2019; Cheeseman et al., 2014; Dumas et al., 2020;

Ekseth et al., 2014; Ropars et al., 2015, 2020; Williams et al., 1986). The *Saccharomyces*

cerevisiae, P. camemberti and *P. roqueforti* populations used in the food industry have

specific traits that are beneficial for food production and enable them to thrive in their human-

made environments to a much greater extent than other populations (Dumas et al., 2020;

Marsit & Dequin, 2015; Ropars et al., 2020), providing evidence of genuine domestication.

The genomic basis of several of these adaptive traits has been determined in *Saccharomyces*

and *Penicillium* food populations, with the identification of gene duplications, interspecific

hybridizations and horizontal gene transfers (Almeida et al., 2014; Cheeseman et al., 2014; C.

Gonçalves et al., 2018; Guillamón & Barrio, 2017; Marsit et al., 2015, 2017; Ropars et al.,

2015). Several adaptive horizontal gene transfers have occurred between distantly related

cheese *Penicillium* fungi, leading to rapid convergent adaptation through the very same genes

and genomic mechanisms (Cheeseman et al., 2014; Ropars et al., 2015). As expected from a

domestication process in selfers or asexuals, the fungal populations used for food production

have lost much of their genetic variability, as demonstrated for *S. cerevisiae* from wine and

beer and *Penicillium* fungi from cheese (Dumas et al., 2020; Legras et al., 2018; Ropars et al.,

2020). Most fungi can be cultured asexually, leading to a particularly large loss of genetic

variability in domesticated fungi (Dumas et al., 2020; Ropars et al., 2020).

Like cheese fungi, *Penicillium nalgiovense* is an excellent model for studying domestication

and adaptation. This species has been used to inoculate the surface of most dry-cured meat

products for decades, to enhance the ripening process (M. Coton et al., 2021; Garcia et al.,

2001; Ludemann et al., 2004; Sunesen & Stahnke, 2003). Another recently described fungus,

Penicillium salamii, also occurs in dry-cured meats (Perrone et al., 2015) and may be included

in starter cultures or form part of the natural mycobiota at local production sites (M. Coton et

al., 2021; Magistà et al., 2016). Several traits have probably been subject to similar selection

processes in both these fungi, either deliberately through human intervention or unwittingly,

to enable these fungi to thrive in dry-cured meats and produce high quality products, with an

appealing appearance and aroma, and a better shelf life and product safety. The fungi colonize

the surface, thereby helping to preserve the dry-cured meat and to protect it against excessive

desiccation, irradiation by light, oxygen and undesirable microorganisms, such as

contaminating toxigenic molds (Chávez et al., 2011; Nielsen et al., 1998). These fungal

species also help to improve the aroma of the dry-cured meat, through the metabolites they

produce with their lipase, protease, transaminase and dehydrogenase activities (Leroy et al.,

2006; Sunesen & Stahnke, 2003). However, if proteolytic and lipolytic activities are too high,

this can result in an undesirable texture, bitterness and a rancid taste (Toldrá, 1998). Another

important trait is the color of *Penicillium* fungi, which is also under selection, to meet human

preferences (Filtenborg et al., 1996) or because the protection that pigments provide in nature,

e.g., against desiccation, radiation and various stresses, may not be needed in food products

anymore (Lin & Xu, 2020). Salt is added to dry-cured meats during production as a taste

enhancer; it also helps to exclude undesirable microorganisms by reducing water activity, one

of the key mechanisms of preservation in these products. The *Penicillium* fungi used to

inoculate dry-cured meats may, therefore, have evolved a high degree of salt tolerance.

Penicillium nalgiovense and *P. salamii* have also been isolated from environments other than

dry-cured meats, such as the soil, plants, sewage and dog bones (Visagie et al., 2014).

However, it remains unknown whether the dry-cured meat *Penicillium* fungi display

particular phenotypes and/or form populations that are genetically isolated from those found

in other environments. *Penicillium nalgiovense* and *P. salamii* lineages have diverged at least

30 million years ago (Houbraken et al., 2020; Perrone et al., 2015; Steenwyk et al., 2019),

which corresponds, approximately, to the divergence between humans and Cercopithecoids

(macaques, langurs…) (Pozzi et al., 2014; Reis et al., 2018); a convergent adaptation to dry-

cured meat, if observed, would therefore constitute an interesting case of parallel adaptation.

The aim of this study was to determine whether, within these two *Penicillium* species, the

strains used in dry-cured meat production had features typical of domesticated organisms, by

investigating whether dry-cured meat strains 1) were genetically different from the strains

isolated from other environments, 2) displayed a genome-wide reduction of genetic diversity

potentially indicative of strong population reduction and/or clonality, 3) had specific traits that

would make them better suited to human needs or preferences than other strains, 4) displayed

footprints of genomic adaptation, and 5) displayed convergent phenotypic and/or genomic

changes. We addressed these questions by isolating strains from dry-cured meat, sequencing

the genomes of multiple *P. nalgiovense* and *P. salamii* strains collected from dry-cured meat

and other environments, and searching for genetic differentiation, recombination footprints,

horizontal gene transfers, specific genome arrangements and traces of positive selection in

terms of high rates of non-synonymous changes. Phenotypic differences between *Penicillium*

fungi from dry-cured meat environments and other species/populations were also investigated

by comparing their growth rates at different salt concentrations, color, lipid and protein

degradation rates and their ability to produce harmful toxins. For interpretation of the

phenotypic differences between strains from dry-cured meat and other environments in terms

of domestication, we further compared the phenotypes of the non-dry-cured meat strains to

those of their respective sister species (*P. chrysogenum* for *P. nalgiovense* and *P. olsonii* for *P.*

salamii). Under the hypothesis that dry-cured meat populations are domesticated, we would

expect traits essential for dry-cured meat products to have evolved specifically in dry-cured

meat strains, and to be absent in strains from other environments and in the corresponding

sister species. In addition, because we found horizontally transferred genomic regions shared

between *P. nalgiovense, P. salamii* and *P. biforme* and because *P. biforme* is also used to

inoculate dry-cured meat, albeit rarely, we also compared phenotypes between strains from

dry-cured meat and other environments for *P. biforme* and its sister species, *P. palitans*. We

cannot entirely rule out that feral strains could be present among fungi isolated from non-dry-

cured meat environments (i.e., escaped from dry-cured meat). In that case, however, they

could only reduce the differences between dry-cured meat and non-dry-cured meat groups.

Hence, any differences found between dry-cured meat and non-dry-cured meat groups are

conservative regarding the occurrence of feral strains. Furthermore, we tried to identify feral

strains by looking for strains isolated from non-dry-cured meat environments but resembling

phenotypically and/or genetically dry-cured meat strains.

Material and Methods

Strain collection and DNA extraction

We used 94 strains for experiments and genome analyses including 22 isolated from dry-

cured meat samples in this study (see below; Table S1), 33 from the CBS collection housed at

the Westerdijk Fungal Biodiversity Institute (Utrecht, the Netherlands), 33 from the public

collection of the national museum of natural history in Paris (LCP) and six from previous

studies. Strain isolation, DNA extraction and species identification protocols are described in

the Supplementary methods

Genome sequencing and genome assembly

For short-read sequencing, DNA libraries and sequencing were performed by the INRAE

GenoToul platform (Toulouse, France). Paired-end libraries of 2x150 bp fragments were

prepared with TruSeq Nano DNA Library Prep Kits (Illumina), and sequencing was

performed on a HiSeq3000 Illumina sequencer, with a mean final coverage of 38x and a

median of 10x (genomes of 27 to 35 Mb). We sequenced the genomes of 55 strains collected

from either dry-cured meat or other environments (Table S1).

We used the P6/C4 Pacific Biosciences SMRT technology to obtain a high-quality reference

genome sequence of the LCP06525 *P. salamii* strain and the ESE00252 *P. nalgiovense* strain,

both from dry-cured meat. Nuclear genomic DNA was extracted with the

cetyltrimethylammonium bromide (CTAB) extraction procedure described previously

(Cheeseman et al., 2014). Sequencing was performed by the UCSD IGM Genomics Facility

La Jolla, CA, USA. Long- and short-read assemblies are described in the Supplementary

methods. All the genomes were deposited on ENA under the study accession number

PRJEB44534 (Table S1).

Population genetic analyses

We mapped cleaned sequencing reads to the reference genomes LCP06525 (*P. salamii*) and

LCP06099 (FM193; *P. nalgiovense*; accession numbers HG815136–HG815288 and

HG815290–HG816004; Ropars et al., 2015) using Stampy with default parameters (v1.0.32;

Lunter & Goodson, 2011). SNPs were called for each individual using the genome analysis

toolkit GATK (v4.1.0.0; McKenna et al., 2010) with the *RealignerTargetCreator*,

IndelRealigner and *UnifiedGenotyper* functions using default parameters except that we used

a ploidy of one. For each species, we merged the variant call formats (VCF) files using the

CombineVariants function in GATK and tagged the sites as follows: quality (with low qual-

tag), quality by depth (QD < 2.0), mapping quality (MQ < 40), Fisher strand bias (FS > 60; a

statistic to detect strand bias in variant calling in reverse and forward strands), mapping

quality rank sum (MQRankSum < 12.5; to test read quality differences between reference and

alternative alleles), read position rank (ReadPosRank < 8.0; based on the distance from the

end of reads compared with alternative allele) and missing data (allowing maximum 10%

missing data). The filter-tags were prepared using vcftools with PASS/filtered tag and GATK

with the *VariantFiltration* function. Indels and tagged SNPs were entirely removed from the

original VCF files using vcftools (v0.1.16; Danecek et al., 2011). SNPs were used to analyze

population structure and linkage disequilibrium (LD). To produce an alignment for a

phylogenetic analysis, we had to include the *P. nalgiovense* PacBio sequence (ESE00252) and

the respective outgroups which were available as fasta files and therefore absent in the vcf file.

We thus used NUCmer v3.1 from the MUMmer3 package (Kurtz et al., 2004) to align those

sequences against the two reference genomes (LCP06099 for *P. nalgiovense* and LCP06525

for *P. salamii*) and called SNPs. We then merged those SNPs with the previous vcf files in R

v3.6.2 (R Core Team, 2019). We built population trees using RAxML (v8.2.7; Stamatakis,

2014), run with the general time-reversible model of nucleotide substitution with the gamma

model of rate heterogeneity (GTRGAMMA) and 100 bootstraps.

In order to analyze population subdivision, we filtered our SNP dataset with bcftools (v1.7; Li,

2011) to keep only distant SNPs to avoid too strong LD (r^2 lower than 0.2 in a window of

1kbp), with a minor allele present in at least two individuals per species (minor allele

frequency of 0.05). We then ran STRUCTURE (v2.3.4; Pritchard et al., 2000) with the

admixture model during 20,000 iterations and removed the first 10,000 iterations as burn-in,

for a number of populations (K) varying from 2 to 10, with 10 independent runs for each K.

We combined the results with the R package pophelper v2.3.0 (Francis, 2017). We used the

full fasta alignment built to estimate the population tree to run a principal component analysis

following the code used in Konishi et al. (2019) and ran it in R v3.6.2 (R Core Team, 2019).

We performed neighbor-net analyses with the *phangorn* R package (Schliep, 2011). The

substitution model used for building the distance matrix was JC69 (Jukes & Cantor, 1969; Li,

2011). The two most divergent *P. nalgiovense* strains, CBS297.97 and DTO 204-F1, were

removed from the neighbor-net analyses to better see the remaining relationships.

We searched for the mating-type locus in the genomes and identified the alleles as MAT1-1 or

MAT1-2 with tblastx v2.6.0 and the *P. chrysogenum* strain ATCC02889 MAT1-1

(AM904544.1) and strain NRRL1249B21 MAT1-2 (AM904545.1) sequences as references.

The presence of the two different mating types in a population could potentially allow sexual

reproduction and recombination (Böhm et al., 2015; Ropars et al., 2014), while the presence

of a single mating type suggests strict asexual reproduction. Fungi able to undergo sexual

reproduction between strains carrying the same mating-type idiomorph in their haploid

genome are indeed extremely rare and this phenomenon is not known in any *Penicillium*

fungus (Wilson et al., 2021).

Detection of horizontal gene transfers between *P. salamii* and *P. nalgiovense*

We aimed at detecting very recent horizontal gene transfers (HGTs) that could be involved in

parallel adaptation to the dry-cured meat environment, having therefore occurred in both *P.*

salamii and *P. nalgiovense* during domestication and thus human times. Genomic regions

resulting from such recent HGTs were expected to be nearly identical between the two distant

species *P. salamii* and *P. nalgiovense* in contrast to the genomic background, and lacking in

many non-dry-cured meat strains and/or species (e.g. Cheeseman et al., 2014). By the very

nature of such recent HGTs, the resulting genomic regions (HTRs for horizontally transferred

regions) are lacking in many strains and are almost identical in the strains in which they are

present, which prevents using phylogenetic incongruence between different regions in the

genomes to detect or validate HGTs, but provides stronger evidence. Other HTRs, present

only in one of these species, may be involved in adaptation to dry-cured meat. However, we

focused on the shared ones, because we were particularly interested in parallel adaptation.

In order to identify HTRs, we first searched for thresholds above which we could consider

that regions were more similar across a longer length than expected given the species

phylogenetic distance. Given that the phylogenetic distance between *P. olsonii* and *P.*

chrysogenum or *P. rubens* (two species formerly considered as one) is the same as the

distance between *P. nalgiovense* and *P. salamii*, sequences that would be more similar and

longer than the ones typically detected between *P. olsonii* and *P. chrysogenum* or *P. rubens*

would not be expected given the phylogenetic distance separating *P. nalgiovense* from *P.*

salamii, and would thus more likely correspond to recent gene flow, i.e. HGT or introgression.

More details are given on the detection of horizontal gene transfers in Supplementary

Methods, as well as on transposable element detection and filtering, and on gene function

enrichment.

For plotting similarity levels along genomes, we merged several "reference" HTRs based on

overlapping sequences, and we kept only those larger than 35 kbp, collectively adding to 1.1

Mbp. Global alignments between these long HTRs and each of the dry-cured meat *Penicillum*

genome assemblies were obtained with minimap2 (Li, 2018) using the -cs asm20 and --

cs=long options. Plotting was performed based on the Pairwise mApping Format (*.paf) files

with plot_coverage function of pafr (https://dwinter.github.io/pafr/index.html) for contig

tracks and Rideogram (Hao et al., 2020) for synteny links in R v3.6.2 (R Core Team, 2019).

The percentage of identity was calculated as the number of matching bases divided by the

length of the query.

Presence and similarity of HTRs in other species

We searched for the presence of the candidate HTRs in other *Penicillium* (n=47), *Aspergillus*

(n=4) and *Monascus* (n=1) species by globally aligning the consensus sequences of the eight

largest HTRs with genome assemblies (see Table S2 in Supplementary data) using minimap2

(-cs asm20 and --cs=long options). We kept matches that either: 1) covered the entire HTR; 2)

covered an entire contig on the assembly with at least 1 kbp of matched bases, or 3) that were

larger than 10 kbp and found at the edges of larger contigs, allowing to a maximum of two

such matches per HTR. We calculated the fraction of each HTR present in the genomes as the

sum of non-ambiguous matches length divided by the total length of the focal HTR, and the

percentage of identity as the number of matched bases to the HTR consensus sequences

divided by the sum of non-ambiguous matches length.

Detection of selection: SnIPRE analyses and Tajima's D

We calculated Tajima's D (Tajima, 1989) using the *strataG* R package (v2.0.2; Archer et al.,

2017) and tested genome-wide neutral evolution or recent change in demography by

comparing the observed values to the 95% confidence interval around zero. A negative

Tajima's D indicates recent selective sweeps, weak negative selection or recent population

expansion while a positive Tajima's D indicates balancing selection, population structure or

recent population contraction. We obtained nucleotide diversity indexes (π, the average

number of pairwise nucleotide differences per site (Tajima, 1989); and θ_W, the observed

nucleotide diversity calculated from the number of segregating sites (Watterson, 1975)) using

the strataG (v2.0.2; Archer et al., 2017) and pegas (v0.14; Paradis, 2010) packages and

scaling to the length of the reference genome. We used vcftools to measure LD, in windows

of 15,000 bp and with a minor allele frequency cutoff of 0.2 (Danecek et al., 2011).

We also aimed at detecting individual genes evolving under positive selection in terms of high

rates of non-synonymous substitutions. Because the two species studied here are used in

industrial contexts (Garcia et al., 2001; Ludemann et al., 2004; Magistà et al., 2016; Sunesen

& Stahnke, 2003), and therefore potentially cultured in high quantities after an initial

bottleneck, we cannot assume a constant effective population size through time. Even more

importantly, these fungi are cultivated asexually to be sold as starters in the industry, therefore

constituting clonal lineages (see results). Most of the classic frameworks for detecting genes

under positive selection are therefore unsuitable. We decided to use SnIPRE (Eilertson et al.,

2012), a Bayesian generalization of the McDonald and Kreitman analyses (McDonald &

Kreitman, 1991), which does not make any assumption on demography or reproduction mode.

This method detects genes in which amino-acid changes are more or less frequent than

expected under neutrality (inferring positive or purifying selection, respectively), by

identifying genes in which non-synonymous divergence to an outgroup is higher or lower than

expected based on observed synonymous and nonsynonymous SNPs in the in-group, thus

accounting for gene-specific mutation rates. We ran the tests for *P. nalgiovense* using *P.*

chrysogenum as an outgroup and for *P. salamii* using *P. olsonii* as an outgroup. For this goal,

we applied the method described in supplementary for the phylogeny reconstruction to obtain

single-copy genes present in all the *P. nalgiovense* strains and five *P. chrysogenum* on one

dataset, and in all the *P. salamii* strains and five *P. olsonii* on a second dataset, allowing one

missing sequence per orthologous gene alignment. From these alignments, we recovered the

number of synonymous and non-synonymous sites, and among each category the number of

fixed and polymorphic sites in each of *P. salamii* and *P. nalgiovense* gene with the MKtest

program of libsequence (Thornton, 2003). We estimated the mean number of synonymous

and polymorphic sites by performing a weighted average on the polydNdS output. The

MKtest and polydNdS programs belong to the package "analysis" of libsequence (Thornton,

2003). We collected the results of these analyses and used them as input for the SnIPRE

analysis. We ran it in its Bayesian implementation following the script providing by the

authors for 25,000 iterations and removed a burn-in of 10,000 iterations, with a thinning value

of four. To test whether the proportion of genes under positive, negative, or neutral selection

was different between one specific COG (cluster of orthologous groups) functional category

and the rest of the genome, we used Fisher's exact tests as described in the supplementary for

the HTRs. As strains from dry-cured meat did not cluster together neither in the phylogenetic

trees nor in the population genomic analyses (see results), we did not compare these strains

with others within each species.

Phenotypic tests

We compared various phenotypic traits between species and strains from different

environments: growth rate at different salt concentrations, spore production, lipolysis, and

proteolysis abilities. We first studied *P. nalgiovense* and *P. salamii*, the most common species

inoculated for dry-cured meat production, with their respective sister species *P. chrysogenum*

and *P. olsonii*. However, we found shared horizontally transferred regions between these dry-

cured meat species and the *P. biforme* complex, including the two clonal lineages *P.*

camemberti var. *camemberti* and *P. camemberti* var. *caseifulvum* (Ropars et al., 2020), and

these can also be, even if more rarely, inoculated in dry-cured meat. We therefore also

analyzed phenotypes in *P. biforme*, that display genetic and phenotypic variability, in contrast

to the cheese clonal *P. camemberti* varieties (Ropars et al., 2020), and compared them to

phenotypes of their sister species *P. palitans*. For all the experiments below, we used all

available *P. nalgiovense*, *P. salamii* and *P. biforme* strains that we could find in public

collections, from non-food environments, and similar numbers of strains, chosen at random,

from dry-cured meat and from their closely related outgroup species. The Table S1 gives the

number and identity of strains used for each species, type of environment and experiment.

The Supplementary methods provide details on phenotypic measures as well as on mycotoxin

or penicillin production.

Statistical analyses of the phenotypic measures

We analyzed several phenotypic traits to test our hypothesis that dry-cured meat fungi are

domesticated: growth and spore production at different salt concentrations using malt extract

agar medium, colony color on malt extract agar medium, and lipolytic and proteolytic

activities. If dry-cured meat fungi have been domesticated, we expected that: 1) there would

be significant differences between dry-cured meat and non-dry-cured meat strains within each

species, and 2) there would be no significant differences between the non-dry-cured meat

strains and the sister species not used for dry-cured meat production; this would indeed

suggest that only the fungi used for dry-cured meat production have evolved specific

phenotypes related to adaptations to the dry-cured meat substrate. Because we found that *P.*

nalgiovense and *P. salamii* shared horizontally transferred regions with the *P. biforme*

complex, which can also be inoculated in dry-cured meat, we included *P. biforme* in

phenotypic assays. We therefore tested first whether there were significant phenotypic

differences between dry-cured meat and non-dry-cured meat strains within each of the species

used for dry-cured meat maturation, i.e., *P. nalgiovense, P. salamii* and *P. biforme.* Then, we

tested whether there were significant differences between the non-dry-cured meat strains

(excluding dry-cured meat strains from this analysis) and the sister species populations, not

used for dry-cured meat production (i.e., *P. olsonii* for *P. salamii, P. chrysogenum* for *P.*

nalgiovense and *P. palitans* for *P. biforme*). To be able to compare sister species, we included

in the tests a "clade" effect, clades corresponding to sister species pairs, to control for the

phylogenetic relatedness in the analysis.

For all analyses, fixed effects were added sequentially to test for their significance through

likelihood ratio tests until reaching the full model with all possible fixed effects. We thereafter

only introduce the full model. All post-hoc tests were conducted using the emmeans function

of package emmeans (Russell et al., 2018) with Tukey correction for pairwise mean

comparison. Post-hoc tests were run when we found a significant effect either between sister

species or between populations from dry-cured meat and non-dry-cured meat.

For analyzing growth at different salt concentrations, we fitted two linear mixed models on

the log-transformed colony area, keeping the day of measurement as a random variable and

the salt concentration as fixed effects in each of them. In the first one, testing the differences

between strains from dry-cured meat and other environments, we further included as fixed

effects the species and its interaction with salt concentration, as well as the isolation

environment (dry-cured meat or not) nested within species and its interaction with salt

concentration. In the second one, where we tested the differences between non-dry-cured

meat strains and the sister species, we included as fixed effects the clade and its interaction

with salt concentration, as well as the species nested within the clade and its interactions with

salt concentration.

For quantifying spore number after growth at various salt concentrations, as too few spores

were recovered at 18% salt concentration, we only analyzed growth data for 0%, 2% and 10%

salt concentration. As spores were only counted once (after the 17-day growth experiment),

we did not include any random effect in the model. We therefore fitted two linear models

using as a dependent variable the log-transformed number of spores divided by the colony

area to normalize the effect of growth on sporulation.

For lipolysis and proteolysis analyses, we fitted two linear mixed models on the distance to

the lysis zone using the day of measurement and the batch number as random effects. To test

for differences between dry-cured meat and other environment strains, we used as fixed

effects species and isolation environment (dry-cured meat or not) nested within species. To

test for differences between non-dry-cured meat populations and the sister species, we

included as fixed effects the clade and the species nested within the clade.

For colony whiteness, we fitted two linear mixed models using the same fixed effects as for

the two models used for proteolysis and lipolysis tests. As three color measurements per strain

were done, we added the measured circle location and strain ID as random effects. For colony

color, because red and green values were strongly correlated (r=0.998), we only used red and

blue components in RGB space to fit generalized linear mixed models with a binomial

response. We considered as random effects the measured circle location and strain ID. Each

model used the same fixed effects as those for proteolysis and lipolysis tests.

All linear mixed models were fitted using the lmer() function of *lme4* R package (Bates et al.,

2015; R Core Team, 2019) and glmer() for the generalized linear mixed models. Simple linear

models were fitted with lm() function (R Core Team, 2019). Likelihood ratio tests between

models were performed using anova() function in R (R Core Team, 2019). For all tests, we

first checked that assumptions were met, i.e., homoscedasticity, independence of residuals,

and normal and identical distribution of residuals.

Results

Genetic diversity and population structure in *P. salamii* and *P. nalgiovense*

We collected 140 dry-cured meat casings, from which we isolated 134 monospore-derived

strains. The most frequent *Penicillium* species isolated from dry-cured meat was *P.

nalgiovense* (*n*=70) followed by *P. salamii* (*n*=16). We also found *P. biforme* (*n*=5), *P.

chrysogenum* (*n*=5), *P. solitum* (*n*=5) and *P. camemberti* (*n*=5) and, sporadically, other

Penicillium species and other genera (Table 1). In six cases (5% of the 106 samples with these

six most abundant *Penicillium* species), we isolated two different species from the same

sample, but never *P. nalgiovense* and *P. salamii* together (Fig. S1).

We obtained high-quality genome assemblies for *P. salamii* LCP06525 (37.00 Mbp) and *P.

nalgiovense* ESE00252 (34.56 Mbp) with long-read (PacBio) sequencing technology and we

sequenced the genomes of 55 strains with Illumina technology, including some of the strains

we isolated as well as other available strains (Table S1). We identified 1,395,641 SNPs by

mapping reads from the 21 Illumina-sequenced *P. salamii* genomes onto the *P. salamii*

reference genome. A population tree based on 2,977,098 SNPs after the addition of the

outgroup (Fig. 1A) revealed genetic differentiation by sampling environment in *P. salamii*.

Two clades contained mostly dry-cured meat strains, whereas the other two clades contained

mostly non-dry-cured meat strains. The frequencies of the two isolation environments differed

significantly between the three clades with more than one strain (Chi²= 7.6723; df = 2; *p*-

value = 0.02). No association with geographic origin was observed for *P. salamii* strains; the

least diverse clade included only strains from Europe, but was also a clade containing mostly

dry-cured meat strains (Fig. 1A).

For the 28 *P. nalgiovense* genomes obtained by Illumina sequencing, reads were mapped onto

the *P. nalgiovense* reference genome (LCP06099=FM193), leading to the identification of

only 380,884 SNPs (381,996 when the other available PacBio genome ESE00252 was added),

indicating a very low level of polymorphism. The *P. nalgiovense* population tree, based on a

2,367,990 SNPs with the outgroup (Fig. 1B), had very short branches for most strains,

suggesting that these strains represent a clonal lineage. No genetic differentiation was

observed, with no effect of sampling environment or geographic location in particular. Only

two strains displayed some genetic differences from the main lineage.

We analyzed 1,766 distant SNPs for *P. nalgiovense* and 7,685 for *P. salamii*, to investigate

the population structures of these two species. For *P. salamii*, the STRUCTURE analysis

supported four well-delimited genetic clusters, consistent with the population tree (Fig. 1A,

Fig. S2A). For *P. nalgiovense,* the STRUCTURE analysis was also consistent with the

population tree, with two well-delimited genetic clusters (Fig. 1B, Fig. S2B); the other genetic

clusters corresponded only to admixed strains, and were not, therefore, considered to

correspond to genuine clusters. The PCA (Fig. 1) recovered patterns similar to those revealed

by the population trees and STRUCTURE analyses: four clusters for *P. salamii*, with a

tendency towards separation by geographic origin, and weaker structure for *P. nalgiovense*,

with the most differentiated strain being from North America.

We identified six different genomic architecture types among *P. salamii* strains (Fig. 1A, Fig.

S3). The genomes of *P. nalgiovense* strains were highly collinear, except for two strains

displaying large rearrangements relative to the other strains (Fig. 1B). These collinearity

groups in both *P. salamii* and *P. nalgiovense* corresponded to the genetic clusters detected

above (Fig. 1). In *P. salamii*, genomic rearrangements differentiated the four strains obtained

outside of Europe (USA and Middle East).

Nucleotide and genetic diversities were much higher in *P. salamii* (π = 0.0111 bp^{-1}, SE = 1 x

10^{-5}; θ_W = 0.0105 bp^{-1}, SE = 7 x 10^{-4}) than in *P. nalgiovense* (π = 0.0009 bp^{-1}, SE = 1 x 10^{-6};

θ_W = 0.0030 bp^{-1}, SE = 2 x 10^{-4}), which had diversity levels similar to those in *P. roqueforti* (π

= 0.0011 bp^{-1}; θ_W = 0.0007 bp^{-1}; Dumas et al., 2020) and *P. biforme* (π = 0.0011 bp^{-1}; θ_W =

0.0011 bp^{-1}; Ropars et al., 2020). We detected both mating types (MAT1-1 and MAT1-2) in

P. salamii, in balanced proportions and with a distribution across strains corresponding to the

genetic structure and genomic architecture types (Fig. 1A). Each of the two MAT groups

displayed relatively low nucleotide and genetic diversities (MAT1-1 group: π = 0.0007 bp^{-1},

SE = 3 x 10^{-6}; θ_W = 0.0006 bp^{-1}, SE = 1 x 10^{-4}; MAT1-2 group: π = 2 x 10^{-5} bp^{-1}, SE = 4 x 10^{-7};

θ_W = 3 x 10^{-5} bp^{-1}, SE = 3 x 10^{-6}). In *P. nalgiovense*, only the two most divergent strains

carried MAT1-1, all the other strains being identified as MAT1-2, supporting the hypothesis

of the use of a single clonal lineage for dry-cured meat production (Fig. 1B). The MAT1-2

group displayed very low diversity (π = 4 x 10^{-5} bp^{-1}, SE = 5 x 10^{-7}; θ_W = 8 x 10^{-5} bp^{-1}, SE = 5

x 10^{-6}).

Overall, *P. nalgiovense* displayed high levels of genome-wide LD, with no detectable decay

over physical distance (Fig. S4). This finding is consistent with a complete absence of

recombination due to exclusively asexual multiplication in this lineage. In *P. salamii,* we

observed a sharp decay in LD within 200 bp, but LD level remained high. Analysis of LD

within each of the two *P. salamii* populations with sufficient numbers of strains and both

mating types revealed the same LD pattern as in *P. nalgiovense* (Fig. S4). This LD pattern in

P. salamii is consistent with the asexual multiplication of two lineages of opposite mating

types, with recombination events occurring in the ancestral population, before the asexual

multiplication of these two lineages. The obtained neighbor-net networks further support the

population structure identified and recent clonality in *P. salamii* and *P. nalgiovense* (Fig. 1).

Detection of horizontal gene transfers between *P. nalgiovense* and *P. salamii*

We aimed to detect very recent horizontally transferred regions (HTRs), present in both *P.*

nalgiovense and *P. salamii, i.e.,* resulting from events (HGTs) during domestication by

humans that could potentially be involved in parallel adaptation to the dry-cured meat

environment. The genomic regions resulting from recent horizontal gene transfers are

expected to be almost identical between the distantly related *P. salamii* and *P. nalgiovense*

species, would not necessarily be present in all strains and would be absent from many non-

dry-cured meat species (e.g. Cheeseman et al., 2014).

We searched for recent HTRs between *P. nalgiovense* and *P. salamii*, by identifying

thresholds above which regions could be considered to be more similar across a longer length

than expected from the phylogenetic distance between the species. For comparisons of

genomic sequences of more than 1,000 bp between the non-dry-cured meat sister species and

their respective sister species (i.e., comparisons of *P. chrysogenum* and *P. rubens*, two species

formerly considered as a single species, with *P. olsonii*), 98.21% was the highest level of

similarity (Fig. S5A). We therefore set the identity threshold for identifying potential HTRs at

98%, because lower levels of similarity might be expected to occur by chance for such long

sequences between sister species. The longest of the nearly identical sequences was 11,414 bp

long (Fig. S5B). However, as the second longest sequence was only 6,972 bp long, we

decided to set the threshold length for identifying potential HTRs to 7,000 bp, because shorter

fragments with such high levels of similarity might occur by chance between the sister species

found in dry-cured meat products.

Our analysis of genomic regions of more than 7,000 bp in length with more than 98%

similarity between pairs of *P. nalgiovense* and *P. salamii* strains identified 26 putative HTRs.

These putative HTRs were between 7,224 bp and 496,686 bp long (Table S4). The cumulative

total length of the putative HTRs was 1,447,064 bp, including 145,640 bp assigned to

transposable elements (TEs) of known families and 558,247 bp assigned to TEs of unknown

families (10% and 38.6%, respectively; Table S4). We identified 34 (65%) of the 52 (2x26)

consensus HTR extremities (the very last bases) as belonging to TEs, and all but three of the

extremities had a detected TE within 1,000 bp, suggesting a possible role for TEs in the

horizontal transfer of some of these regions. After the removal of all TEs of known and

unknown families identified in the *P. nalgiovense* and *P. salamii* reference genomes ("repeat-

masked" analyses), three putative HTRs had lengths below 500 bp and were therefore

discarded for subsequent analyses (Table S4).

Plotting the identified putative HTRs on the contigs of the *P. salamii* and *P. nalgiovense*

reference genomes showed that some groups of HTRs clustered very close to each other (Fig.

2), suggesting that they may represent a single horizontal transfer event in these strains (in

particular HTRs 4, 5, 6, 7 and 8). The other HTRs were located in very small contigs and may

therefore also actually belong to the same regions but being not assembled due to repetitive

elements. We found three occurrences of the DUF3435 domain, characteristic of the Starship

elements involved in HGTs in fungi (Gluck-Thaler et al., 2022), in the two large clusters of

HTRs. The HTRs were collinear between the reference genomes of *P. salamii* and *P.*

nalgiovense despite widespread genomic rearrangements elsewhere (Fig. 2), which brings

further support to the inference of recent horizontal transfers. We present hereafter results

separating the 23 initially identified HTRs as no criterion appeared optimal to decide which

should be merged. There has nevertheless likely been just one or a few independent horizontal

transfer events with cumulative length of recent HTRs reaching almost 1.5 Mb.

We identified 283 coding genes in the putative HTRs remaining after repeat masking, 199 of

which yielded hits in one of the egg-NOG databases, mostly with the fungal egg-NOG

database (87.4%, 173 genes), suggesting that most of the putative HTRs identified originated

from fungi. The other 25 genes had best hits in the eukaryote egg-NOG database (12.1%) or

the principal egg-NOG database (0.5%). They could, therefore, also be of fungal origin, but

better annotated in more phylogenetically distant groups (e.g. mammals), resulting in a best

match at a higher taxonomical level.

All but the two most divergent *P. nalgiovense* strains contained the same putative HTRs,

whereas there was more variability in terms of presence/absence of HTRs among *P. salamii*

strains, with some strains even carrying none of the putative HTRs (Fig. 3; Figs. S6-8). The

number and length of the putative HTRs largely accounted for the differences in genome size

between *P. salamii* strains (adjusted R^2: 0.67; Fig. S9). The fraction of each putative HTR

present in genomes differed between strains (Fig. 3; Figs. S6-8) but the present sequences

were always very similar to that of the reference putative HTR (99-100% identity; Fig. 3; Figs.

S6-8). Among the eight largest HTRs, only HTR5 had around a third of its sequence with

identity below 98% among *P. nalgiovense* strains (Fig. 3). There were two different versions

of the sequence in *P. nalgiovense* (with 89% identity between them), where the strains were in

this region either 100% identical to one or the other. They may therefore correspond to recent

HGTs of the same region from two different donor species or populations. The finding of near

100% identity for most HTRs between strains and species suggests HGTs during

domestication by humans and preclude any dating attempt.

We constructed a phylogenetic species tree of 68 genomes from other *Penicillium, Aspergillus*

or *Monascus* species with 1102 single-copy orthologous genes. The resulting species tree (Fig.

4) was consistent with previous studies (Houbraken et al., 2014; Perrone et al., 2015;

Steenwyk et al., 2019) and was well supported. The putative HTRs were absent from most of

the other species analyzed (Fig. 4, Figs. S10-S12). Only 25 species carried parts of at least

one putative HTR, and 19 species carried parts of more than one putative HTR. The species

carrying the largest number of the 23 putative HTRs belonged to genus *Penicillium* (Fig. 4,

Figs. S10-S12). Two *Penicillium* species had large fragments of seven putative HTRs: *P.*

biforme and both varieties of *P. camemberti* (Fig. 4, Figs. S10-S12). Interestingly, these two

species belong to a species complex found in dry-cured meat and sometimes even used for

inoculation for dry-cured meat maturation, but mostly used for cheese production. The two

varieties of *P. camemberti,* var. *camemberti* and var. *caseifulvum,* are two closely related

clonal lineages displaying very little variability, having diverged recently from *P. biforme,*

which harbors higher levels of genetic and phenotypic variability (Ropars et al., 2020). Most

of the other *Penicillium* species harbored only small parts of the putative HTRs. HTR 6 and

HTR 15 were present in more species than the other putative HTRs and had similar

distributions across species (Fig. 4, Figs. S10-S12). The similarity between the putative HTRs

present in the different species and the reference putative HTR was very high, despite the

phylogenetic distances between the species harboring them (typically more than 95%, Fig. 4,

Figs. S10-S12), providing further support for the view that they result from horizontal gene

transfers.

We, thus, identified genomic regions that were almost identical between distantly related

species, absent in more closely related species, rich in TEs (accounting for a mean of 44% of

the regions, ranging from 9 to 91%), including some firmly linked to known TE families, and

located in different regions of the genomes among species, or even the population. These

findings provide compelling evidence that these regions resulted from horizontal gene

transfers. Given the very nature of the putative HTRs, and their complete absence in many

strains and species, it is not possible to use genealogies to test for horizontal gene transfer, but

the evidence provided above is even stronger. These very high identity levels also prevent

using gene genealogies to identify the donor species for HTRs. The genetic diversity within

species was not useful either to identify the donor species as we had only one or two genomes

for each species, except *P. nalgiovense* and *P. salamii* in which they were nearly identical.

Given the distance separating species in our tree, the *P. nalgiovense* – *P. salamii* pair

appeared to have a particularly high shared HTR content, in terms of total length. Only the *P.*

nalgiovense – *P. biforme* pair had a longer total length of HTRs (Fig. S13), which,

interestingly, also corresponds to a pair of species used in dry-cured meat production. More

generally, all the species that were phylogenetically distantly related but had a large number

of putative HTRs in common were known to be present and sometimes used in either cheese

or dry-cured meat production (Fig. S13).

An analysis of functional categories in clusters of orthologous groups present in the HTRs

identified eight categories present in most strains and four categories that were less present

(Table S5). None of the *P. nalgiovense* or *P. salamii* strains displayed an enrichment in

certain functions in the HTRs relative to the rest of the genome. The seven HTRs present in *P.*

biforme and the two varieties of *P. camemberti* carried 42 and 47 genes, but only seven to 20

of them had predicted functions. These genes were involved in the cytoskeleton, transport and

metabolism of carbohydrates, inorganic ions, lipids or secondary metabolites, post-

translational modifications, or intracellular trafficking (Table S5). The two regions present in

the largest number of species (HTR6 and HTR15) carried 62 and 11 genes, respectively, 45

and 10 of which, respectively, were annotated, and only 24 and seven, respectively, with

predicted functions, belonging to 11 different COGs. The carbohydrate transport and

metabolism category was the only one containing five genes, three being the maximum for the

other categories. Therefore, some gene functions in the HTRs may be beneficial for dry-cured

meat production as they are involved in the metabolism of meat nutrients, but the lack of

significant enrichment renders conclusions challenging.

Detection of positive selection

For *P. salamii*, we found no evidence of genome-wide positive or negative selection or a

recent change in demography (Tajima's $D = 0.148$, not significantly different from zero, 95%

confidence interval [-1.657; 2.153]; *p*-value: 0.432). By contrast, for *P. nalgiovense*, we found

evidence of widespread selective sweeps, background selection or recent population

expansion (Tajima's $D = -2.834$, 95% confidence interval [-4.641; -0.814]; *p*-value: 6e-08).

We looked for footprints of selection outside of HTRs with a Bayesian generalization of the

McDonald and Kreitman test (McDonald & Kreitman, 1991), which identifies genes in which

amino-acid changes are more or less frequent than expected under neutrality (inferring

positive or purifying selection, respectively), by detecting higher or lower non-synonymous

divergence from an outgroup than expected based on observed synonymous and

nonsynonymous SNPs in the in-group. We analyzed 7,155 genes present in all, or all but one

of the strains of *P. nalgiovense*, and 7,010 genes present in all, or all but one of the *P. salamii*

strains. We found that 1,572 of these genes in *P. nalgiovense* and 466 in *P. salamii* were

evolving under purifying selection, and that 71 of these genes were common to the two

species (Table S6). We identified 751 genes in *P. salamii* and 30 in *P. nalgiovense* evolving

under positive selection, with only three of these genes common to the two species (one with

no predicted function, one predicted to encode a phosphotransferase and the third predicted to

be involved in meiosis; Table S6). The genes under positive or purifying selection displayed

no specific clustering within genomes. We detected a significant excess of positive selection

in *P. nalgiovense* for genes involved in secondary metabolite biosynthesis, transport, and

catabolism, and in *P. salamii* for genes involved in transcription, posttranslational

modification, protein turnover, chaperones, signal transduction and defense mechanisms

(Table S6). We detected a significant excess of functions under purifying selection for

functions involved in the transport and metabolism of carbohydrates and amino acids in *P.*

nalgiovense and for functions involved in energy production and conversion, and in secondary

metabolite biosynthesis, transport, and catabolism in *P. salamii* (Table S6).

Phenotypic experiments to test the hypothesis of domestication

We analyzed several phenotypic traits, to test the hypothesis that dry-cured meat fungi are

domesticated: growth and spore production on malt extract agar medium at different salt

concentrations, colony color on malt extract agar medium, and lipolytic and proteolytic

activities. If dry-cured meat fungi have been domesticated, then we would expect to see: 1)

significant differences between dry-cured meat and non-dry-cured meat strains within each

species, and 2) no significant differences between the non-dry-cured meat strains and the

sister species not used for dry-cured meat production. These findings would indicate that only

the fungi used for dry-cured meat production had evolved specific phenotypes related to

adaptations to the dry-cured meat substrate. We found that *P. nalgiovense* and *P. salamii*

contained horizontally transferred regions common to the *P. biforme* complex, which can also

be used to inoculate dry-cured meat. We therefore also included *P. biforme* in phenotypic

assays. We investigated the possibility that there were phenotypic differences between dry-

cured meat and non-dry-cured meat strains within each of the species used for dry-cured meat

maturation: *P. nalgiovense, P. salamii* and *P. biforme.* We also investigated whether there

were differences between the non-dry-cured meat strains (the dry-cured meat strains were

excluded from this analysis) and sister species not used for dry-cured meat production (i.e., *P.*

olsonii for *P. salamii, P. chrysogenum* for *P. nalgiovense* and *P. palitans* for *P. biforme*). For

the comparison of sister species, we included a "clade" effect in this test, with clades

corresponding to sister-species pairs, to control for phylogenetic relatedness in the analysis.

Salt tolerance: effect of salt concentration on growth and spore number

In comparisons of growth between non-dry-cured meat populations and their sister species in

the three clades under various salt concentrations, we found significant effects of salt

concentration, clade and an interaction between clade and salt concentration, but no effect of

species identity within a clade (Table 2, Fig. 5). Thus, the non-dry-cured meat strains and

their sister species had similar responses to salt. All pairs of species grew faster in the

presence of 2% salt than without salt, but growth rates rapidly decreased with increasing salt

concentration beyond 2% (Table 2, Fig. 5).

In the analysis including dry-cured meat strains, we found no difference in growth between

populations from dry-cured meat and populations from other environments (Table 2, Fig. 5).

We found a significant interaction between isolation environment and salt concentration, a

significant interaction between species and salt concentration and separate independent effects

for salt concentration and species. This suggests that salt tolerance levels may differ between

dry-cured meat and non-dry-cured meat strains (Table 2, Fig. 5). However, no post-hoc test of

the impact of isolation environments was significant, in any species. The strongest difference

between different isolation environments was for *P. salamii*, in which dry-cured meat strains

grew better than non-dry-cured-meat strains in the absence of salt, although this difference

was not significant in a post-hoc test (df =327; *t*=-3.285; *p*-value =0.1533).

We also measured spore production after 17 days of growth on malt extract agar medium

containing various concentrations of salt. Spore numbers were very low on medium

containing 18% salt. We therefore removed this category from the analysis. An analysis

including only non-dry-cured meat strains and their sister species revealed a significant and

strong effect of salt, with higher concentrations impairing spore production (Table 2, Fig. 5);

we also found a weaker, but nevertheless significant within-clade species effect (Table 2, Fig.

5). In the analysis limited to dry-cured meat species, we found no difference between

populations from dry-cured meat and those from other environments (Table 2, Fig. 5). We

found a significant interaction between species identity and salt concentration (Table 2, Fig.

5). Significantly higher levels of sporulation were observed for *P. nalgiovense* than for *P.*

salamii, but only in the total absence of salt (df=113, *t*=3.416, *p*-value=0.0240).

Proteolysis and lipolysis activity

We investigated whether the dry-cured meat strains had evolved specific proteolysis and/or

lipolysis activities in *P. nalgiovense, P. salamii* and *P. biforme.* A comparison of species

without dry-cured meat strains revealed a significant effect of clade on both proteolysis and

lipolysis (Table 2, Fig. 5). We detected no within-clade effect of species: the non-dry-cured

meat populations and their sister species had similar levels of activity (Table 2, Fig. 5).

In the analysis including both dry-cured meat and non-dry-cured meat strains, the within-

species effect of the isolation environment was significant for both proteolysis and lipolysis,

and there was also a significant species effect (Table 2, Fig. 5). Post-hoc tests showed that, in

P. biforme and *P. salamii,* proteolysis was significantly slower in strains from dry-cured meat

than in strains from other environments (Table 2). We observed a similar tendency in *P.*

nalgiovense, but it was not significant. Lipolysis rates were also lower in strains from dry-

cured meat than in those from other environments in *P. nalgiovense* and in *P. salamii*, but no

such effect was detected in *P. biforme* (Table 2).

The differences between dry-cured meat and non-dry-cured meat strains, together with the

lack of difference between the non-dry-cured meat populations and their sister species suggest

that dry-cured meat strains have evolved specific catabolic abilities, constituting

domestication (Table 2, Fig. 5).

Brightness and color

We tested whether colony color was closer to white in dry-cured meat populations, as this

may have been a criterion of selection by humans or pigments may not be useful anymore in

the human-made environment, where UV are less present (Lin & Xu, 2020). We found no

significant species or clade effect in the analysis including only non-dry-cured meat strains

and their sister species (Table 2, Fig. 6). In the analysis including all strains of the dry-cured

meat species, we found a significant effect of species and of isolation environment within

species on the distance from whiteness in the RGB (red, green and blue) space (Table 2, Fig.

6). Post-hoc tests showed that only *P. nalgiovense* colonies from dry-cured meat were

significantly closer to white than colonies from other environments (Table 2). Visual

inspection, indeed, suggested that the colonies on malt extract agar medium were typically

white with a smooth surface in *P. nalgiovense* strains from dry-cured meat, whereas the

colonies were more greenish, with a smooth or cotton-like surface in *P. salamii* (Fig. 6).

These results indicate that colony color in the dry-cured meat population in *P. nalgiovense* has

evolved toward greater whiteness.

We then investigated whether the dry-cured meat strains had evolved specific colony colors in

terms of their red, green, and blue (RGB) components. In analyses limited to non-dry-cured

meat strains and their sister species, the colors of fungal colonies were significantly affected

by clade identity but not by the species within a clade (Table 2, Fig. 6). In analyses limited to

dry-cured meat species, both isolation environment and species had significant effects on

colony color (Table 2, Fig. 6). Post-hoc tests showed that *P. biforme* dry-cured meat strains

were redder and/or less blue (Table 2), whereas the differences were not significant in the

other two species. Thus, *P. biforme* dry-cured meat strains have evolved to have less blue

colonies.

Mycotoxins and penicillin production

We detected none of the mycotoxins or penicillin assessed (andrastin A, ermefortins A & B,

(iso)-fumigaclavin A, meleagrin, mycophenolic acid, ochratoxin A, penicillic acid, penicillin

G, penitrem A, roquefortine C and sterigmatocystin) in any of the *P. salamii* and *P.*

nalgiovense dry-cured meat strains tested (Table S7). None of these extrolites were detected

in *P. biforme* in a previous study (Ropars et al., 2020). All the sequenced strains from both *P.*

salamii and *P. nalgiovense* harbored genes for the entire pathway for penicillin production in

their genomes (*PCBAB, PCBC, penDE* and *phl* genes, and the *RFX* transcription factor gene).

None of the extrolites were detected in *P. olsonii*. We observed two distinct extrolite

production profiles in *P. chrysogenum*, the sister species of *P. nalgiovense*. Five strains

produced low levels of roquefortine C and no other extrolite (Table S7). The other nine *P.*

chrysogenum strains produced high levels of andrastin A, meleagrin, roquefortine C and

penicillin G.

Possible feral strains

We tried to check whether some strains could be feral, i.e. escaped from dry-cured meat and

isolated from other environment, with the rationale that they would be genetically and/or

phenotypically close to dry-cured meat strains but isolated in another substrate. Among the

six *P. nalgiovense* potential "feral" strains (i.e. closely related to dry-cured meat strains:

CBS112438, DTO259.D5, LCP6099, LCP03915, LCP03991, LCP05232) four were tested for

all the phenotypic traits (CBS112438, DTO259.D5, LCP03915, LCP05232) and one for the

color (LCP03991). Four of them (CBS112438, LCP03915, LCP03991, LCP05232) exhibited

phenotypes close to those of dry-cured meat strains (lower proteolysis and lipolysis, whiter

color), and it was therefore not possible to rule out a feral origin. In contrast, the strain

DTO259.D5 displayed phenotypes really similar to other non-dry-cured meat strains (high

proteolysis and lipolysis, darker color), suggesting that it was a genuine wild strain rather than

a recent escape from an industrial environment or that *P. nalgiovense* is able to revert back

quickly to wild phenotypes. Two *P. salamii* strains were also genetically related to dry-cured

meat strains despite having been collected in other environments (DTO309.F9 and

CBS135407). Both were phenotyped and did not differ from the non-dry-cured strains, ruling

out a potential "feral" origin.

Discussion

The use of dry-curing as a method for conserving a perishable raw meat material can be traced back to Greek and Roman times (Frost, 1999). However, it remains unclear when the inoculation of dry-cured meat casings with *Penicillium* fungi began. *Penicillium nalgiovense* is the commonest species in the dry-cured meat food industry (Andersen & Frisvad, 1994; Larsen & Breinholt, 1999; Sunesen & Stahnke, 2003). *Penicillium salamii* was recently identified as an alternative ripening agent and has only been used or recovered from local dry-cured meat production sites, often producing naturally fermented meat products (M. Coton et al., 2021; Sinch & Dincho, 1994; Sunesen & Stahnke, 2003). This difference in usage was reflected in our samples from dry-cured meats, for which *P. nalgiovense* was the most frequently isolated species, followed by *P. salamii.* These two most abundant species have however not been found to co-occur in our samples despite six cases of species co-isolation, suggesting that they may be used as alternatives for dry-cured meat production. The wider use of *P. nalgiovense* may explain the difference in genetic diversity and the differentiation between the two species: the very low level of genetic diversity in *P. nalgiovense* may result from the selection of a single clonal lineage widely used in commercial dry-cured meat production, resulting in a strong bottleneck. Alternatively, the low level of genetic diversity may be explained by a possible relatively recent introduction of this species into Europe. Indeed, we had only four strains from outside Europe (USA and the Middle East), two of which were genetically different from the main lineage. *Penicillium nalgiovense* strains from non-food environments displayed no specific clustering, and most originated from human-related environments, such as the car upholstery, cheese, and dog bone. *Penicillium salamii* had a higher level of genetic diversity, consistent with the lack of wide-scale commercial use

for this species (Magistà et al., 2016; Perrone et al., 2015). Genetic differentiation was

observed in *P. salamii*, with clusters including different proportions of strains from dry-cured

meats or other environments (e.g., sewage plant, tea leaves or soil), indicating that migration

between dry-cured meat and other environments is still occurring.

We found no evidence of recombination in *P. nalgiovense*, with little LD decay, and only one

mating type among the dry-cured meat strains, consistent with the use of asexual cultures of

commercial starters. In the cheese fungi *P. camemberti* and *P. roqueforti*, cheese populations

also have low levels of genetic diversity and clonal population structures (Dumas et al., 2020;

Ropars et al., 2020). Non-cheese *P. roqueforti* populations have much higher levels of genetic

diversity, with a larger number of recombination footprints, and higher levels of sexual

fertility (Ropars et al., 2014, 2016), contrasting with our findings for *P. nalgiovense*. However,

the alternative mating type was detected in the non-dry-cured meat strains. In *P. salamii*, we

identified both mating types among the dry-cured meat strains, suggesting that sexual

reproduction may occur in this environment. However, the LD patterns and the presence of

only one mating type in each *P. salamii* population suggests an absence of recent sexual

reproduction. The detection of both mating types in the two dry-cured meat species

nevertheless opens up possibilities for future strain breeding.

Despite the lack of strong genetic differentiation between dry-cured meat strains and other

strains, we found domestication footprints when comparing phenotypes relevant to dry-cured

meat production. Indeed, dry-cured meat strains of both *P. salamii* and *P. nalgiovense*

displayed slower proteolysis and lipolysis than strains from other environments. *Penicillium*

biforme presented the same pattern for proteolysis. Excessive protein degradation is

detrimental for dry-cured meat products, potentially rendering the product less firm and

imparting bitter and metallic tastes (Toldrá, 1998), whereas excessive lipolysis would dry the

product (Demyer et al., 1974) and have a negative impact on flavor. We found no major

difference in lipolysis or proteolysis between the non-dry-cured meat strains of *P. salamii*, *P.*

nalgiovense and *P. biforme* and their respective sister species, indicating that the dry-cured

meat strains had evolved specific and convergent phenotypes, slower lipolysis and proteolysis,

probably under selection by humans.

Fungal colonization of the surface of the meat casing has a direct impact on the observed

color of the product, and this fungal coating has probably been subject to selection by humans

for dry-cured meat fungal strains. Indeed, the visual appearance of the product is crucial for

dry-cured meat-eaters, and a white color typically makes such products more attractive.

Penicillium nalgiovense strains sampled from dry-cured meat were whiter than strains from

other environments, consistent with the selection of dry-cured meat starters on the basis of

color (Leistner, 1990). Dry-cured meat strains of *P. biforme* were less blue than non-dry-

cured meat strains of this fungus. *Penicillium biforme* is widely used for cheese-making and is

whiter than the closely related species *P. fuscoglaucum* on cheese media (Ropars et al., 2020).

Whiter *P. biforme* strains may have been selected because it generates a more appealing

aspect for both cheese and dry-cured meat production, especially as the strains isolated from

these two types of product belong to the same population (Ropars et al., 2020). The clonal

cheese fungus *P. camemberti* is known from historical records to have been selected as a

white mutant to make Brie cheeses that looked less "moldy" (Ropars et al., 2020). White

color in dry-cured meat fungi could alternatively be the result of relaxed selection in human-

made environments because pigments would not be useful anymore to protect against UV or

other stresses (Lin & Xu, 2020). By contrast, we observed no difference in color between *P.*

salamii strains from different environments, probably due to weaker or more recent selection

by humans, possibly even with colonization of the product principally from the environment

rather than through active inoculation (Perrone et al., 2015), or due to contrasting selection

pressures. Demand for local and traditional products is increasing, including that for dry-

cured meats (Iaccarino et al., 2006), and this may have fostered an interest in less

homogeneous and less white fungi on sausage casings. In the analysis limited to non-dry-

cured meat strains, the lack of major color differences between the species used for dry-cured

meat and their sister species indicated that the changes occurred specifically in dry-cured meat

strains, further supporting the occurrence of domestication.

Phenotypic tests thus suggest that there may have been selection in *Penicillium* fungi for

certain strains relevant to the production of dry-cured meat, with convergent evolution toward

similar phenotypes, especially for lipolysis and proteolysis. Some of the strains isolated from

non-dry-cured meat environments may be feral, but this would only make differences between

strains from dry-cured meat and other environments more difficult to detect, this could not

have generated or inflated the differences we found between strains from dry-cured meat and

other environments, so our findings are robust. By contrast, we detected no specific traits

relating to the response to salt concentration in dry-cured meat strains. *Penicillium* fungi,

particularly those present in fermented food, are salt-tolerant and, therefore, able to grow in

the salt-rich environments found in dry-cured meats and cheeses (Yadav et al., 2018); there

may have been no further advantage of higher levels of salt tolerance or evolutionary

constraints may have prevented the development of this tolerance.

None of the targeted extrolites, including mycotoxins and penicillin, were detected in *P.*

nalgiovense or *P. salamii* dry-cured meat strains (M. Coton et al., 2021). However, the

production of penicillin has been reported in some *P. nalgiovense* strains (Andersen &

Frisvad, 1994) and the closely related species *P. chrysogenum* is known to produce mycotoxins (M. Coton et al., 2019; Frisvad et al., 2004). The lack of antibiotic or mycotoxin production in dry-cured meat strains may be due to relaxed selection in a human-made environment containing fewer competitors or to active selection for safe products. In *Penicillium* cheese fungi, some of the mycotoxin biosynthesis pathway genes harbor deletions, impairing these functions specifically in cheese populations and indicating selection for safe food production or relaxed selection (E. Coton et al., 2020; Gillot et al., 2017; Ropars et al., 2020). The absence of mycotoxin production in both *P. salamii* and its sister species *P. olsonii* suggests that this feature did not result from selection by humans in this clade. However, the culture medium is known to affect toxin production in *Penicillium* fungi (Frisvad et al., 2004).

We investigated genomic footprints potentially associated with adaptation to dry-cured meat production. Given the clonal structure of the dry-cured meat strains, especially in *P. nalgiovense*, it was not possible to perform genome-wide scans for the detection of selection based on either local decreases in genetic diversity or local genomic differentiation between populations. Indeed, under asexual reproduction, the selection of a beneficial variant results in a hitchhiking of the whole genome without the generation of typical local selective sweeps or islands of genomic differentiation. We therefore looked for footprints of horizontal gene transfers, which have been shown to be an important and frequent mechanism of rapid adaptation under selection by humans in fungi (Cheeseman et al., 2014; Marsit et al., 2015; Novo et al., 2013; Ropars et al., 2015). By comparing available *Penicillium* genomes from various environments, we found evidence of horizontal gene transfers, and, specifically, of horizontal transfers between *P. nalgiovense* and *P. salamii* despite the relatively large genetic distance between these species. The cumulative length of the HTRs shared between *P.*

nalgiovense and *P. salamii* reached nearly 1.5 Mb. We could not generate genealogical trees

of these regions to support the inference of horizontal transfer because most species

completely lack these regions, but these absences, together with the very high levels of

identity between these regions and their different locations within the genome, provide even

stronger support for the inference of horizontal transfer. Indeed, introgression by interspecific

hybridization would lead to the high-similarity regions being located at the same loci across

species and being present in all species. In addition, given the very low collinearity (Fig. 2)

and an estimate of 30 My divergence between *P. nalgiovense* and *P. salamii* (Houbraken et

al., 2020; Perrone et al., 2015; Steenwyk et al., 2019), it is very unlikely that sexual

reproduction could be successful. The inference of horizontal transfer is further supported by

the presence of the DUF3435 domain in two large clusters of HTRs, as these domains are

characteristic of the Starship mobile elements (Gluck-Thaler et al., 2022). By contrast to

findings for cheese fungi, the HTRs were not specific to the dry-cured meat strains

(Cheeseman et al., 2014; Ropars et al., 2015), which may be due to some non-dry-cured meat

strains being feral strains, to gene flow between dry-cured meat and non-dry-cured meat

populations, or to a horizontal transfer older than the divergence between these strains. In

such cases, this could mean that humans chose fungi with desirable traits for dry-cured meat

production, acquired by horizontal transfers before human selection. However, HTRs were

nearly 100% identical between species and strains, indicating horizontal transfers in human

times. Several HTRs also appeared to be present in other species, particularly in *P. biforme*

but also in *P. camemberti*, which are also used commercially in the food industry, mostly for

the inoculation of cheeses, but also in dry sausages (this study; Cheeseman et al., 2014;

Ropars et al., 2015, 2020). The abundance of shared HTRs was higher in dry-cured meat

species than in available *Penicillium* genome pairs, suggesting their possible selection in this

rich, human-made environment, as previously reported for cheese *Penicillium* fungi

(Cheeseman et al., 2014; Ropars et al., 2015) and *Saccharomyces* yeasts used in wine-making

(Marsit et al., 2015). We were unable to identify any particular functions overrepresented in

these regions relative to the rest of the genome, possibly due to the low proportion of genes

with predicted functions or to the diversity of functions under selection. Other HTRs may be

present in only *P. salamii* or only *P. nalgiovense* and involved in adaptation to the dry-cured

meat environment but our approach cannot detect such HTRs. We were indeed mostly

interested in parallel adaptation to dry-cured meat.

Despite the low level of genetic differentiation between strains from dry-cured meat and other

environments, we were able to detect phenotypic differences, even in *P. nalgiovense,* in which

overall genetic diversity was low. However, a few SNPs may have a strong phenotypic impact,

and the insertion of transposable elements may also modify gene regulation. In order to detect

genes under selection for adaptation to the dry-cured meat environment, we also ran tests to

detect non-synonymous substitutions that were more or less frequent relative to synonymous

substitutions than would be expected under neutrality. *Penicillium nalgiovense* had few genes

under positive selection and many under purifying selection, consistent with a high frequency

of asexual reproduction in this species and, thus, probable genome-wide hitchhiking effects

and background selection. The number of genes under positive selection was 25 times greater

in *P. salamii* than in *P. nalgiovense,* and the genes concerned were involved in different

functions. This finding may reflect the lower adaptation ability of this species, with low levels

of genetic polymorphism. A few categories of genes seemed to be overrepresented among the

genes under positive or negative selection, including some that could be linked to the dry-

cured meat environment (carbohydrate metabolism and transport, defense mechanisms and

secondary metabolite production). It is therefore likely that, in addition to horizontal gene

transfers, positive selection on genes already present in the genome has contributed to the

adaptation to dry-cured meat in *P. salamii* and *P. nalgiovense*. Other types of adaptive

changes may also have occurred that could not be studied here because of the low genetic

diversity, such as selective sweeps in regulatory regions.

Overall, our findings suggest that selection by humans has induced convergent phenotypic

changes in *P. biforme*, *P. salamii* and *P. nalgiovense*, and that *Penicillium* dry-cured meat

fungi have acquired beneficial traits for dry sausage production. These fungi have, therefore,

been domesticated. The low level of genetic differentiation between dry-cured meat and non-

dry-cured meat strains is likely due to the domestication process being very recent. Currently,

P. nalgiovense strains are cultivated and sold to be inoculated in dry-cured meat (e.g.

https://www.lip-sas.fr/index.php/nos-produits/salaisons/nos-propositions), so that the control-

over-reproduction component of domestication is clearly present. The convergence in the

evolution of dry-cured meat populations was most striking for proteolysis, with lower levels

of proteolysis in dry-cured meat populations for *P. nalgiovense*, *P. salamii* and *P. biforme*.

Our findings, thus, add to the growing evidence of domestication in food fungi, such as in

beer and wine *Saccharomyces* yeast, koji *Aspergillus oryzae* (Gibbons et al., 2012), and

cheese *Penicillium* fungi (Dumas et al., 2020; Gallone et al., 2016; M. Gonçalves et al., 2016;

Ropars et al., 2020; Sicard & Legras, 2011). Genomic analyses in *Saccharomyces cerevisiae*

have revealed a differentiation of populations between wild and food environments, and even

specific populations corresponding to different uses of yeast, for the fermentation of bread,

beer, wine or cheese (Gallone et al., 2016; M. Gonçalves et al., 2016). The different clades

have contrasting fermentation traits, such as the capacity to use malt, aroma production and

ethanol production. In the blue cheese fungus *P. roqueforti*, population differentiation was

also observed between cheese and other environments, with a more rapid growth of cheese

strains on cheese medium, higher levels of lipolytic activity, more efficient cheese cavity

colonization, more diverse and positive aromas, and a higher salt tolerance (Caron et al., 2020;

Dumas et al., 2020; Ropars et al., 2015). In both *S. cerevisiae* and *Penicillium* cheese-making

fungi, HGT events have been implicated in adaptation (Cheeseman et al., 2014; Ropars et al.,

2015).

In conclusion, we found clear convergent adaptation in two distantly related dry-cured

meat species of fungi, in terms of proteolysis and lipolysis, and in terms of shared horizontal

gene transfers. Such studies on parallel adaptation to the same ecological niche are important

for understanding the process of evolution and whether it is repeatable. However, only a few

studies on a handful of models have investigated parallel adaptation. Studies on natural

populations have shown that adaptation to similar ecological niches has led to the convergent

evolution of phenotypes, such as size, color and form, in *Anolis* lizards, three-spined

sticklebacks and cichlid fishes in African rift lakes, for example (Colosimo et al., 2004;

Duboué et al., 2011; Elmer et al., 2014; Mahler et al., 2013; Salzburger, 2009). Dry-cured

meat species of *Penicillium* therefore appear to be good models for investigating general

questions about adaptation. We also found a substantial loss of diversity in dry-cured meat

populations, as previously reported for cheese fungi (Dumas et al., 2020; Ropars et al., 2020).

This lack of diversity may jeopardize future strain improvement and lead to degeneration. Our

findings therefore also have implications for the food industry. The discovery of different

phenotypes and the presence of two mating types in both species paves the way for strain

improvement by conventional breeding, which could also allow studying the genomic bases

of beneficial phenotypes for dry-cured meat making and generating further genetic and

phenotypic diversity.

1019

1020

1021　**Data accessibility**

1022　All sequencing data have been deposited to the European Nucleotide Archive under the

1023 accession PRJEB44534. The HTR sequences and their annotations are available under the

1024 publicly available gitlab project https://gitlab.com/abranca/consensus_htr.

1025

1026

Acknowledgments

This work was supported by the ERC starting grant GenomeFun 309403 and Louis D

foundation award to TG, a CNRS PEPS adaptation grant to AB and the FUNGADAPT

ANR☐19☐CE20☐0002 ANR grant to TG, AB and MC. YCL acknowledges a PhD grant

from the France–Taiwan joint program Campus France. We thank Alice Feurtey, Fanny

Hartmann and Fantin Carpentier for help with genome analyses. We thank MNHN for

granting access to strains with the help of Manuela Lopez-Villavicencio and Sandrine Lacoste.

We thank all the people who sent dry-cured meat casings from around the world. We thank

the GenoToul platform for sequencing. We are also grateful to the GenoToul bioinformatics

platform Toulouse Occitanie (Bioinfo Genotoul, doi: 10.15454/1.5572369328961167E12) for

providing some computing resources. PacBio sequencing was conducted at the IGM

Genomics Center, University of California, San Diego, La Jolla, CA. We thank Federico

Laich and Giancarlo Perronne for providing *P. salamii* strains. Some sequence data (Table S2)

were produced by the US Department of Energy Joint Genome Institute

http://www.jgi.doe.gov/ in collaboration with the user community.

1042

References:

Almeida, P., Gonçalves, C., Teixeira, S., Libkind, D., Bontrager, M., Masneuf-Pomarède, I.,

Albertin, W., Durrens, P., Sherman, D. J., Marullo, P., Todd Hittinger, C., Gonçalves,

P., & Sampaio, J. P. (2014). A Gondwanan imprint on global diversity and

domestication of wine and cider yeast *Saccharomyces uvarum*. *Nature*

Communications, 5, 4044. http://dx.doi.org/10.1038/ncomms5044

Andersen, S. J., & Frisvad, J. C. (1994). Penicillin production by *Penicillium nalgiovense*.

Letters in Applied Microbiology, 19(6), 486–488. https://doi.org/10.1111/j.1472-

765X.1994.tb00988.x

Archer, F. I., Adams, P. E., & Schneiders, B. B. (2017). stratag: An r package for

manipulating, summarizing and analysing population genetic data. *Molecular Ecology*

Resources, 17(1), 5–11. https://doi.org/10.1111/1755-0998.12559

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. *Journal of Statistical Software*, 67, 1–48.

https://doi.org/10.18637/jss.v067.i01

Bodinaku, I., Shaffer, J., Connors, A. B., Steenwyk, J. L., Kastman, E., Rokas, A., Robbat, A.,

& Wolfe, B. (2019). Rapid phenotypic and metabolomic domestication of wild

Penicillium molds on cheese. *BioRxiv*, 647172. https://doi.org/10.1101/647172

Böhm, J., Dahlmann, T. A., Gümüşer, H., & Kück, U. (2015). A MAT1-2 wild-type strain

from *Penicillium chrysogenum*: functional mating-type locus characterization, genome

sequencing and mating with an industrial penicillin-producing strain. *Molecular*

Microbiology, 95(5), 859–874. https://doi.org/10.1111/mmi.12909

Caron, T., Le Piver, M., Péron, A.-C., Lieben, P., Lavigne, R., Brunel, S., Roueyre, D., Place,

M., Bonnarme, P., Giraud, T., Branca, A., Landaud, S., & Chassard, C. (2020). Strong

effect of *Penicillium roqueforti* populations on volatile and metabolic compounds

responsible for aromas, flavour and texture in blue cheeses. *BioRxiv*,

2020.03.02.974352. https://doi.org/10.1101/2020.03.02.974352

Chan, Y. F., Marks, M. E., Jones, F. C., Villarreal, G., Shapiro, M. D., Brady, S. D.,

Southwick, A. M., Absher, D. M., Grimwood, J., Schmutz, J., Myers, R. M., Petrov,

D., Jónsson, B., Schluter, D., Bell, M. A., & Kingsley, D. M. (2010). Adaptive

Evolution of Pelvic Reduction in Sticklebacks by Recurrent Deletion of a Pitx1

Enhancer. *Science*, 327(5963), 302. https://doi.org/10.1126/science.1182213

Chávez, R., Fierro, F., García-Rico, R. O., & Laich, F. (2011). Mold-fermented foods:

Penicillium spp. as ripening agents in the elaboration of cheese and meat products.

https://doi.org/10.2174/97816080522331101010073

Cheeseman, K., Ropars, J., Renault, P., Dupont, J., Gouzy, J., Branca, A., Abraham, A.-L.,

Ceppi, M., Conseiller, E., Debuchy, R., Malagnac, F., Goarin, A., Silar, P., Lacoste, S.,

Sallet, E., Bensimon, A., Giraud, T., & Brygoo, Y. (2014). Multiple recent horizontal

transfers of a large genomic region in cheese making fungi. *Nature Communications*,

5, 2876. http://dx.doi.org/10.1038/ncomms3876

Colosimo, P. F., Peichel, C. L., Nereng, K., Blackman, B. K., Shapiro, M. D., Schluter, D., &

Kingsley, D. M. (2004). The Genetic Architecture of Parallel Armor Plate Reduction

in Threespine Sticklebacks. *PLOS Biology*, 2(5), e109.

https://doi.org/10.1371/journal.pbio.0020109

Coton, E., Coton, M., Hymery, N., Mounier, J., & Jany, J.-L. (2020). *Penicillium roqueforti*:

an overview of its genetics, physiology, metabolism and biotechnological applications.

Fungal Biology Reviews, 34(2), 59–73. https://doi.org/10.1016/j.fbr.2020.03.001

Coton, M., Auffret, A., Poirier, E., Debaets, S., Coton, E., & Dantigny, P. (2019). Production

and migration of ochratoxin A and citrinin in Comté cheese by an isolate of

Penicillium verrucosum selected among *Penicillium* spp. mycotoxin producers in YES

medium. *Food Microbiology*, 82, 551–559. https://doi.org/10.1016/j.fm.2019.03.026

Coton, M., Deniel, F., Mounier, J., Joubrel, R., Robieu, E., Pawtowski, A., Jeuge, S.,

Taminiau, B., Daube, G., Coton, E., & Frémaux, B. (2021). Microbial Ecology of

French Dry Fermented Sausages and Mycotoxin Risk Evaluation During Storage.

Frontiers in Microbiology, 12, 737140–737140. PubMed.

https://doi.org/10.3389/fmicb.2021.737140

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker,

R. E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R., & 1000

Genomes Project Analysis Group. (2011). The variant call format and VCFtools.

Bioinformatics, 27(15), 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

Darwin, C. (1859). The origin of species. London, UK: John Murray.

https://search.library.wisc.edu/catalog/9934839413602122

Darwin, C. (1868). The Variation of Animals and Plants Under Domestication. London, UK:

John Murray.

Demyer, D., Hoozee, J., & Mesdom, H. (1974). Specificity of lipolysis during dry sausage

ripening. *Journal of Food Science*, 39(2), 293–296. https://doi.org/10.1111/j.1365-

2621.1974.tb02878.x

Duboué, E. R., Keene, A. C., & Borowsky, R. L. (2011). Evolutionary Convergence on Sleep

Loss in Cavefish Populations. *Current Biology*, 21(8), 671–676.

https://doi.org/10.1016/j.cub.2011.03.020

Dumas, E., Feurtey, A., Rodríguez de la Vega, R. C., Le Prieur, S., Snirc, A., Coton, M.,

Thierry, A., Coton, E., Le Piver, M., Roueyre, D., Ropars, J., Branca, A., & Giraud, T.

(2020). Independent domestication events in the blue-cheese fungus *Penicillium*

roqueforti. *Molecular Ecology*, 29(14), 2639–2660. https://doi.org/10.1111/mec.15359

Dupont, J., Dequin, S., Giraud, T., Le Tacon, F., Marsit, S., Ropars, J., Richard, F., & Selosse,

M.-A. (2017). Fungi as a Source of Food. *Microbiology Spectrum*, 5(3).

http://www.asmscience.org/content/journal/microbiolspec/10.1128/microbiolspec.FU

NK-0030-2016

Eilertson, K. E., Booth, J. G., & Bustamante, C. D. (2012). SnIPRE: Selection Inference

Using a Poisson Random Effects Model. *PLOS Computational Biology*, 8(12),

e1002806. https://doi.org/10.1371/journal.pcbi.1002806

Ekseth, O. K., Kuiper, M., & Mironov, V. (2014). orthAgogue: an agile tool for the rapid

prediction of orthology relations. *Bioinformatics*, 30(5), 734–736.

https://doi.org/10.1093/bioinformatics/btt582

Elmer, K. R., Fan, S., Kusche, H., Luise Spreitzer, M., Kautt, A. F., Franchini, P., & Meyer,

A. (2014). Parallel evolution of Nicaraguan crater lake cichlid fishes via non-parallel

routes. *Nature Communications*, 5, 5168. https://doi.org/10.1038/ncomms6168

Filtenborg, O., Frisvad, J. C., & Thrane, U. (1996). Moulds in food spoilage. *Specific*

Spoilage Organisms, 33(1), 85–102. https://doi.org/10.1016/0168-1605(96)01153-1

Francis, R. M. (2017). pophelper: an R package and web app to analyse and

visualize population structure. *Molecular Ecology Resources*, 17(1), 27–32.

https://doi.org/10.1111/1755-0998.12509

Frisvad, J., Smedsgaard, J., Larsen, T. O., & Samson, R. (2004). Mycotoxins, drugs and other

extrolites produced by species in *Penicillium subgenus Penicillium*. *Stud. Mycol*, 49,

201–241. https://www.semanticscholar.org/paper/Mycotoxins%2C-drugs-and-other-

extrolites-produced-by-Frisvad-

Smedsgaard/bcf2c69758b38d2914f5fae4d954a81d5b9a37b4

Frost, F. J. (1999). Sausage and Meat Preservation in Antiquity. *Greek Roman and Byzantine*

Studies, 40, 241–252.

Gallone, B., Steensels, J., Prahl, T., Soriaga, L., Saels, V., Herrera-Malaver, B., Merlevede, A.,

Roncoroni, M., Voordeckers, K., Miraglia, L., Teiling, C., Steffy, B., Taylor, M.,

Schwartz, A., Richardson, T., White, C., Baele, G., Maere, S., & Verstrepen, K. J.

1145 (2016). Domestication and Divergence of *Saccharomyces cerevisiae* Beer Yeasts. *Cell*,

1146 166(6), 1397-1410.e16. https://doi.org/10.1016/j.cell.2016.08.020

1147 Garcia, M. L., Casas, C., Toledo, V. M., & Selgas, M. D. (2001). Effect of selected mould

1148 strains on the sensory properties of dry fermented sausages. *European Food Research*

1149 *and Technology*, 212(3), 287–291. https://doi.org/10.1007/s002170000258

1150 Gibbons, J. G., Salichos, L., Slot, J. C., Rinker, D. C., McGary, K. L., King, J. G., Klich, M.

1151 A., Tabb, D. L., McDonald, W. H., & Rokas, A. (2012). The evolutionary imprint of

1152 domestication on genome variation and function of the filamentous fungus *Aspergillus*

1153 *oryzae*. *Current Biology: CB*, 22(15), 1403–1409.

1154 https://doi.org/10.1016/j.cub.2012.05.033

1155 Gillot, G., Jany, J.-L., Poirier, E., Maillard, M.-B., Debaets, S., Thierry, A., Coton, E., &

1156 Coton, M. (2017). Functional diversity within the *Penicillium roqueforti* species.

1157 *International Journal of Food Microbiology*, 241, 141–150.

1158 https://doi.org/10.1016/j.ijfoodmicro.2016.10.001

1159 Gladieux, P., Ropars, J., Badouin, H., Branca, A., Aguileta, G., M., V. D., C., R. de la V. R.,

1160 Sara, B., & Tatiana, G. (2013). Fungal evolutionary genomics provides insight into the

1161 mechanisms of adaptive divergence in eukaryotes. *Molecular Ecology*, 23(4), 753–773.

1162 https://doi.org/10.1111/mec.12631

1163 Gluck-Thaler, E., Ralston, T., Konkel, Z., Ocampos, C. G., Ganeshan, V. D., Dorrance, A. E.,

1164 Niblack, T. L., Wood, C. W., Slot, J. C., Lopez-Nicora, H. D., & Vogan, A. A. (2022).

1165 Giant Starship Elements Mobilize Accessory Genes in Fungal Genomes. *Molecular*

1166 *Biology and Evolution*, 39(5), msac109. https://doi.org/10.1093/molbev/msac109

1167 Gonçalves, C., Wisecaver, J. H., Kominek, J., Oom, M. S., Leandro, M. J., Shen, X.-X.,

1168 Opulente, D. A., Zhou, X., Peris, D., Kurtzman, C. P., Hittinger, C. T., Rokas, A., &

1169   Gonçalves, P. (2018). Evidence for loss and reacquisition of alcoholic fermentation in

1170    a fructophilic yeast lineage. *ELife*, 7, e33034. https://doi.org/10.7554/eLife.33034

1171 Gonçalves, M., Pontes, A., Almeida, P., Barbosa, R., Serra, M., Libkind, D., Hutzler, M.,

1172    Gonçalves, P., & Sampaio, J. P. (2016). Distinct Domestication Trajectories in Top-

1173    Fermenting Beer Yeasts and Wine Yeasts. *Current Biology*, 26(20), 2750–2761.

1174    https://doi.org/10.1016/j.cub.2016.08.040

1175 Guillamón, J. M., & Barrio, E. (2017). Genetic Polymorphism in Wine Yeasts: Mechanisms

1176    and Methods for Its Detection. *Frontiers in Microbiology*, 8, 806.

1177    https://doi.org/10.3389/fmicb.2017.00806

1178 Hao, Z., Lv, D., Ge, Y., Shi, J., Weijers, D., Yu, G., & Chen, J. (2020). RIdeogram: drawing

1179    SVG graphics to visualize and map genome-wide data on the idiograms. *PeerJ.*

1180    *Computer Science*, 6, e251. https://doi.org/10.7717/peerj-cs.251

1181 Houbraken, J., de Vries, R. P., & Samson, R. A. (2014). Chapter Four - Modern Taxonomy of

1182    Biotechnologically Important *Aspergillus* and *Penicillium* Species. In S. S. & G. G.M.

1183    (Eds.), Advances in Applied Microbiology (Vol. 86, pp. 199–249). Academic Press.

1184    https://doi.org/10.1016/B978-0-12-800262-9.00004-4

1185 Houbraken, J., Kocsubé, S., Visagie, C. M., Yilmaz, N., Wang, X.-C., Meijer, M., Kraak, B.,

1186    Hubka, V., Bensch, K., Samson, R. A., & Frisvad, J. C. (2020). Classification of

1187    *Aspergillus*, *Penicillium*, *Talaromyces* and related genera (*Eurotiales*): An overview of

1188    families, genera, subgenera, sections, series and species. *Genera of Hyphomycetes: A*

1189    *Tribute to Keith A. Seifert*, 95, 5–169. https://doi.org/10.1016/j.simyco.2020.05.002

1190 Hufford, M. B., Xu, X., van Heerwaarden, J., Pyhäjärvi, T., Chia, J.-M., Cartwright, R. A.,

1191    Elishire, R. J., Glaubitz, J. C., Guill, K. E., Kaeppler, S. M., Lai, J., Morrell, P. L.,

1192    Shannon, L. M., Song, C., Springer, N. M., Swanson-Wagner, R. A., Tiffin, P., Wang,

1193  J., Zhang, G., … Ross-Ibarra, J. (2012). Comparative population genomics of maize

1194  domestication and improvement. *Nature Genetics*, 44(7), 808–811. PASCAL.

1195 Iaccarino, T., Monaco, R. di, Mincione, A., Cavella, S., & Masi, P. (2006). Influence of

1196  information on origin and technology on the consumer response: the case of

1197  soppressata salami. *Food Quality and Preference*, 17(1/2), 76–84.

1198  https://eurekamag.com/research/004/442/004442522.php

1199 Jukes, T., & Cantor, C. (1969). Evolution of protein molecules. *Mammalian Protein*

1200  *Metabolism*, 21–132.

1201 Konishi, T., Matsukuma, S., Fuji, H., Nakamura, D., Satou, N., & Okano, K. (2019). Principal

1202  Component Analysis applied directly to Sequence Matrix. *Scientific Reports*, 9(1),

1203  19297. https://doi.org/10.1038/s41598-019-55253-0

1204 Kurtz, S., Phillippy, A., Delcher, A. L., Smoot, M., Shumway, M., Antonescu, C., & Salzberg,

1205  S. L. (2004). Versatile and open software for comparing large genomes. *Genome*

1206  *Biology*, 5(2), R12–R12. PubMed. https://doi.org/10.1186/gb-2004-5-2-r12

1207 Larsen, T. O., & Breinholt, J. (1999). Dichlorodiaportin, Diaportinol, and Diaportinic Acid:

1208  Three Novel Isocoumarins from *Penicillium nalgiovense*. *Journal of Natural Products*,

1209  62(8), 1182–1184. https://doi.org/10.1021/np990066b

1210 Legras, J.-L., Galeote, V., Bigey, F., Camarasa, C., Marsit, S., Nidelet, T., Sanchez, I.,

1211  Couloux, A., Guy, J., Franco-Duarte, R., Marcet-Houben, M., Gabaldon, T., Schuller,

1212  D., Sampaio, J. P., & Dequin, S. (2018). Adaptation of *S. cerevisiae* to Fermented

1213  Food Environments Reveals Remarkable Genome Plasticity and the Footprints of

1214  Domestication. *Molecular Biology and Evolution*, 35(7), 1712–1727. PubMed.

1215  https://doi.org/10.1093/molbev/msy066

1216 Leistner, L. (1990). Mould‐fermented foods: Recent developments. *Food Biotechnology*,

1217  4(1), 433–441. https://doi.org/10.1080/08905439009549755

Leroy, F., Verluyten, J., & De Vuyst, L. (2006). Functional meat starter cultures for improved

sausage fermentation. *International Journal of Food Microbiology*, 106(3), 270–285.

https://doi.org/10.1016/j.ijfoodmicro.2005.06.027

Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association

mapping and population genetical parameter estimation from sequencing data.

Bioinformatics, 27(21), 2987–2993. https://doi.org/10.1093/bioinformatics/btr509

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*,

34(18), 3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Lin, L., & Xu, J. (2020). Fungal Pigments and Their Roles Associated with Human Health.

Journal of Fungi (Basel, Switzerland), 6(4). https://doi.org/10.3390/jof6040280

Ludemann, V., Pose, G., Pollio, M. L., & Segura, J. (2004). Determination of growth

characteristics and lipolytic and proteolytic activities of *Penicillium* strains isolated

from Argentinean salami. *International Journal of Food Microbiology*, 96(1), 13–18.

https://doi.org/10.1016/j.ijfoodmicro.2004.03.003

Lunter, G., & Goodson, M. (2011). Stampy: A statistical algorithm for sensitive and fast

mapping of Illumina sequence reads. *Genome Research*, 21(6), 936–939. PMC.

https://doi.org/10.1101/gr.111120.110

Magistà, D., Ferrara, M., Del Nobile, M. A., Gammariello, D., Conte, A., & Perrone, G.

(2016). *Penicillium salamii* strain ITEM 15302: A new promising fungal starter for

salami production. *International Journal of Food Microbiology*, 231, 33–41.

https://doi.org/10.1016/j.ijfoodmicro.2016.04.029

Mahler, D. L., Ingram, T., Revell, L. J., & Losos, J. B. (2013). Exceptional Convergence on

the Macroevolutionary Landscape in Island Lizard Radiations. *Science*, 341(6143),

292. https://doi.org/10.1126/science.1232392

Marsit, S., & Dequin, S. (2015). Diversity and adaptive evolution of *Saccharomyces* wine

yeast: a review. *FEMS Yeast Research*, 15(7), fov067. PMC.

https://doi.org/10.1093/femsyr/fov067

Marsit, S., Leducq, J.-B., Durand, É., Marchant, A., Filteau, M., & Landry, C. R. (2017).

Evolutionary biology through the lens of budding yeast comparative genomics. *Nature*

Reviews Genetics, 18, 581. http://dx.doi.org/10.1038/nrg.2017.49

Marsit, S., Mena, A., Bigey, F., Sauvage, F.-X., Couloux, A., Guy, J., Legras, J.-L., Barrio, E.,

Dequin, S., & Galeote, V. (2015). Evolutionary Advantage Conferred by an

Eukaryote-to-Eukaryote Gene Transfer Event in Wine Yeasts. *Molecular Biology and*

Evolution, 32(7), 1695–1707. https://doi.org/10.1093/molbev/msv057

Maruyama, T., & Fuerst, P. A. (1985). Population bottlenecks and nonequilibrium models in

population genetics. II. Number of alleles in a small population that was formed by

recent bottleneck. *Genetics*, 111(3), 675.

http://www.genetics.org/content/111/3/675.abstract

McDonald, J. H., & Kreitman, M. (1991). Adaptive protein evolution at the Adh locus in

Drosophila. *Nature*, 351(6328), 652–654. https://doi.org/10.1038/351652a0

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella,

K., Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The Genome

Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA

sequencing data. *Genome Research*, 20(9), 1297–1303. PMC.

https://doi.org/10.1101/gr.107524.110

Nei, M., Maruyama, T., & Chakraborty, R. (1975). The bottleneck effect and genetic

variability in populations. *Evolution*, 29(1), 1–10. https://doi.org/10.1111/j.1558-

5646.1975.tb00807.x

Nielsen, M. S., Frisvad, J. C., & Nielsen, P. V. (1998). Protection by fungal starters against

growth and secondary metabolite production of fungal spoilers of cheese.

International Journal of Food Microbiology, 42(1), 91–99.

https://doi.org/10.1016/S0168-1605(98)00070-1

Novo, M., Mangado, A., Quirós, M., Morales, P., Salvadó, Z., & Gonzalez, R. (2013).

Genome-Wide Study of the Adaptation of *Saccharomyces cerevisiae* to the Early

Stages of Wine Fermentation. *PLOS ONE*, 8(9), e74086.

https://doi.org/10.1371/journal.pone.0074086

Paradis, E. (2010). pegas: an R package for population genetics with an integrated–modular

approach. *Bioinformatics*, 26(3), 419–420.

https://doi.org/10.1093/bioinformatics/btp696

Perrone, G., Samson, R. A., Frisvad, J. C., Susca, A., Gunde-Cimerman, N., Epifani, F., &

Houbraken, J. (2015). *Penicillium salamii*, a new species occurring during seasoning

of dry-cured meat. *International Journal of Food Microbiology*, 193, 91–98.

https://doi.org/10.1016/j.ijfoodmicro.2014.10.023

Peter, J., De Chiara, M., Friedrich, A., Yue, J.-X., Pflieger, D., Bergström, A., Sigwalt, A.,

Barre, B., Freel, K., Llored, A., Cruaud, C., Labadie, K., Aury, J.-M., Istace, B.,

Lebrigand, K., Barbry, P., Engelen, S., Lemainque, A., Wincker, P., … Schacherer, J.

(2018). Genome evolution across 1,011 *Saccharomyces cerevisiae* isolates. *Nature*,

556(7701), 339–344. https://doi.org/10.1038/s41586-018-0030-5

Pozzi, L., Hodgson, J. A., Burrell, A. S., Sterner, K. N., Raaum, R. L., & Disotell, T. R.

(2014). Primate phylogenetic relationships and divergence dates inferred from

complete mitochondrial genomes. *Molecular Phylogenetics and Evolution*, 75, 165–

183. https://doi.org/10.1016/j.ympev.2014.02.023

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using

multilocus genotype data. *Genetics*, 155(2), 945–959. PubMed.

https://pubmed.ncbi.nlm.nih.gov/10835412

R Core Team. (2019). R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria.

Reis, M. D., Gunnell, G. F., Barba-Montoya, J., Wilkins, A., Yang, Z., & Yoder, A. D. (2018).

Using Phylogenomic Data to Explore the Effects of Relaxed Clocks and Calibration

Strategies on Divergence Time Estimation: Primates as a Test Case. *Systematic*

Biology, 67(4), 594–615. https://doi.org/10.1093/sysbio/syy001

Ropars, J., Didiot, E., Rodríguez de la Vega, R. C., Bennetot, B., Coton, M., Poirier, E.,

Coton, E., Snirc, A., Le Prieur, S., & Giraud, T. (2020). Domestication of the

emblematic white cheese-making fungus *Penicillium camemberti* and its

diversification into two varieties. *Current Biology*, 30(22), 4441–4453.

https://doi.org/10.1016/j.cub.2020.08.082

Ropars, J., López-Villavicencio, M., Dupont, J., Snirc, A., Gillot, G., Coton, M., Jany, J.-L.,

Coton, E., & Giraud, T. (2014). Induction of sexual reproduction and genetic diversity

in the cheese fungus *Penicillium roqueforti*. *Evolutionary Applications*, 7(4), 433–441.

PMC. https://doi.org/10.1111/eva.12140

Ropars, J., Rodríguez de la Vega, R. C., López-Villavicencio, M., Gouzy, J., Dupont, J.,

Swennen, D., Dumas, E., Giraud, T., & Branca, A. (2016). Diversity and Mechanisms

of Genomic Adaptation in Penicillium. In Aspergillus and Penicillium in the post-

genomic era. (R. P. de Vries, G. I. Benoit, & R. Andersen M., Eds.). Caister Academic

Press.

Ropars, J., Rodríguez de la Vega, R. C., López-Villavicencio, M., Gouzy, J., Sallet, E.,

Dumas, É., Lacoste, S., Debuchy, R., Dupont, J., Branca, A., & Giraud, T. (2015).

Adaptive Horizontal Gene Transfers between Multiple Cheese-Associated Fungi.

Current Biology, 25(19), 2562–2569. https://doi.org/10.1016/j.cub.2015.08.025

Russell, V. L., Paul, B., Maxime, H., Jonathon, L., Hannes, R., & Henrik, S. (2018). emmeans:

Estimated Marginal Means, aka Least-Squares Means version 1.7.0 from CRAN

(Version 1) [R]. https://rdrr.io/cran/emmeans/

Salzburger, W. (2009). The interaction of sexually and naturally selected traits in the adaptive

radiations of cichlid fishes. *Molecular Ecology*, 18(2), 169–185.

https://doi.org/10.1111/j.1365-294X.2008.03981.x

Schliep, K. P. (2011). phangorn: phylogenetic analysis in R. *Bioinformatics*, 27(4), 592–593.

https://doi.org/10.1093/bioinformatics/btq706

Sicard, D., & Legras, J.-L. (2011). Bread, beer and wine: Yeast domestication in the

Saccharomyces sensu stricto complex. *Comptes Rendus Biologies*, 334(3), 229–236.

https://doi.org/10.1016/j.crvi.2010.12.016

Sinch, B. J., & Dincho, D. (1994). Molds as Protective Cultures for Raw Dry Sausages.

Journal of Food Protection, 57(10), 928–930. https://doi.org/10.4315/0362-028X-

57.10.928

Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis

of large phylogenies. *Bioinformatics*, 30(9), 1312–1313. PMC.

https://doi.org/10.1093/bioinformatics/btu033

Steensels, J., Gallone, B., Voordeckers, K., & Verstrepen, K. J. (2019). Domestication of

Industrial Microbes. *Current Biology : CB*, 29(10), R381–R393.

https://doi.org/10.1016/j.cub.2019.04.025

Steenwyk, J. L., Shen, X.-X., Lind, A. L., Goldman, G. H., & Rokas, A. (2019). A Robust

Phylogenomic Time Tree for Biotechnologically and Medically Important Fungi in the

Genera *Aspergillus* and *Penicillium*. *MBio*, 10(4), e00925-19.

https://doi.org/10.1128/mBio.00925-19

Stitzer, M. C., & Ross-Ibarra, J. (2018). Maize domestication and gene interaction. *New Phytologist*, 220(2), 395–408.

Sunesen, L. O., & Stahnke, L. H. (2003). Mould starter cultures for dry sausages—selection, application and effects. *Meat Science*, 65(3), 935–948. https://doi.org/10.1016/S0309-1740(02)00281-4

Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*, 123(3), 585–595.

Thornton, K. (2003). libsequence: a C++ class library for evolutionary genetic analysis. *Bioinformatics*, 19(17), 2325–2327. https://doi.org/10.1093/bioinformatics/btg316

Toldrá, F. (1998). Proteolysis and lipolysis in flavour development of dry-cured meat products. *Meat Science*, 49, S101–S110. https://doi.org/10.1016/S0309-1740(98)90041-9

Visagie, C. M., Houbraken, J., Frisvad, J. C., Hong, S.-B., Klaassen, C. H. W., Perrone, G., Seifert, K. A., Varga, J., Yaguchi, T., & Samson, R. A. (2014). Identification and nomenclature of the genus *Penicillium*. *Studies in Mycology*, 78, 343–371. https://doi.org/10.1016/j.simyco.2014.09.001

Watarai, N., Yamamoto, N., Sawada, K., & Yamada, T. (2019). Evolution of *Aspergillus oryzae* before and after domestication inferred by large-scale comparative genomic analysis. *DNA Research*, 26(6), 465–472. https://doi.org/10.1093/dnares/dsz024

Watterson, G. A. (1975). On the number of segregating sites in genetical models without recombination. *Theoretical Population Biology*, 7(2), 256–276. https://doi.org/10.1016/0040-5809(75)90020-9

Williams, A. P., Pitt, J. I., & Hocking, A. D. (1986). The Closely Related Species of

Subgenus *Penicillium* — A Phylogenic Exploration. In R. A. Samson & J. I. Pitt

(Eds.), Advances in Penicillium and Aspergillus Systematics (pp. 121–128). Springer

US. https://doi.org/10.1007/978-1-4757-1856-0_12

Wilson, A. M., Gabriel, R., Singer, S. W., Schuerg, T., Wilken, P. M., van der Nest, M. A.,

Wingfield, M. J., & Wingfield, B. D. (2021). Doing it alone: Unisexual reproduction

in filamentous ascomycete fungi. *Fungal Biology Reviews*, 35, 1–13.

https://doi.org/10.1016/j.fbr.2020.12.003

Wolfe, B. E., & Dutton, R. J. (2015). Fermented Foods as Experimentally Tractable Microbial

Ecosystems. *Cell*, 161(1), 49–55. https://doi.org/10.1016/j.cell.2015.02.034

Yadav, A. N., Verma, Dr. P., Kumar, V., Sangwan, P., Mishra, S., Panjiar, N., Gupta, V. K.,

& Saxena, A. (2018). Biodiversity of the Genus *Penicillium* in Different Habitats. In

New and Future Developments in Microbial Biotechnology and Bioengineering (pp.

3–18). Elsevier. https://doi.org/10.1016/B978-0-444-63501-3.00001-6

Zhang, C., Rabiee, M., Sayyari, E., & Mirarab, S. (2018). ASTRAL-III: polynomial time

species tree reconstruction from partially resolved gene trees. *BMC Bioinformatics*,

19(6), 153. https://doi.org/10.1186/s12859-018-2129-y

1380

1381

1382

1383

1384 **Table 1. Species identified among the 134 monospore-derived strains isolated from dry-**

1385 **cured meat casings.**

Species	Number of identified monospores
Total *Penicillium*	124
P. nalgiovense	70
P. salamii	16
P. biforme	5
P. camemberti	5
P. chrysogenum	5
P. solitum	5
P. polonicum	4
P. brevicompactum	2
P. chermesinum	2
P. crustosum	2
P. nordicum	2
P. expansum	1
P. glabrum	1
P. italicum	1
P. olsonii	1
P. roqueforti	1
P. tularense	1
Total other genera	10
Aspergillus **sp.**	2
Fusarium proliferatum	2
Scopulariopsis **sp.**	2
Alternaria **sp.**	1
Cladosporium salinae	1
Cladosporium spinulosum	1
Geotrichum candidum	1

1386

Table 2. Statistics on the phenotypes measured in this study. Model A corresponds to the

model testing whether non-dry-cured meat population in species used for dry-cured meat

harbored different phenotypes than their sister species and Model B aimed at testing whether

non-dry-cured samples showed significant differences relative to dry-cured meat samples. A

clade corresponds to the species used in dry-cured-meat production and their sister species.

Environment effect corresponds to either dry-cured meat or non-dry-cured meat. Some

variables being non-independent (a species belong to a clade, and an environment is

characteristic of a species), they are included as nested parameters and shown with squared

brackets. Interactions are represented by colons. Cells with red bold font represent the main

focus of the test, that is whether there is a difference between dry-cured meat and non-dry-

cured meat populations. Cells with yellow background highlight significant differences with

p-value < 0.05. Stars after the p-values correspond to a value between 0.05 and 0.01, between

0.01 and 0.001, and below 0.001 for one, two and three stars respectively.

	Dependent variable ->	Growth on Salt			Sporulation on Salt				Proteolysis			Lipolysis			Color (Red/Blue)			Whiteness		
	Explanatory variables	Chi Df	Chi Sq^5	p-value	Df	SumOfSq	F-value^5	p-value	Chi Df	ChiSq^5	p-value	Chi Df	ChiSq^5	p-value	Chi Df	ChiSq^5	p-value	Chi Df	ChiSq^5	p-value
Model A	**Clade**	2	7.084	0.0290 *	2	0.321	0.1974	0.8211	2	33.618	<0.001 ***	2	35.7256	<0.001 ***	2	6.0635	0.0482 *	2	1.3504	0.5091
	Species[Clade]	3	7.0541	0.0702	3	7.772	3.1869	0.0266 *	3	4.6392	0.2002	3	1.4852	0.6857	3	3.9425	0.2677	3	3.1148	0.3743
	Salt Concentration	3	546.1269	<0.001 ***	2	18.291	11.2509	<0.001 ***												
	Clade:Salt Concentration	6	46.0191	<0.001 ***	4	4.170	1.2823	0.2814												
	Species[Clade]:Salt Concentration	9	11.5286	0.2412	6	0.785	0.1609	0.9864												
Model B	**Species**	2	15.6135	<0.001 ***	2	11.823	9.5065	<0.001 ***	2	14.164	<0.001 ***	2	9.1309	0.0104 *	2	28.708	<0.001 ***	2	13.526	0.0012 **
	Environment[Species]	3	0.6689	0.8805	3	1.663	0.8912	0.4481	3	63.867	<0.001 ***[1]	3	53.2360	<0.001 ***[2]	3	21.779	<0.001 ***[3]	3	14.073	0.0028 **[4]
	Salt Concentration	3	547.9128	<0.001 ***	2	10.188	8.1918	<0.001 ***												
	Species:Salt Concentration	6	38.4008	<0.001 ***	4	7.652	3.0761	0.0191 *												
	Environment[Species]:Salt Concentration	9	17.2739	0.0446 *	6	4.250	1.1391	0.3444												

1400

[1] Post-hoc tests for *P. biforme*: df=165, t=4.397, p-value=0.0003; *P. nalgiovense*: df=165, t=2.670, p-value=0.0869 and *P. salamii*: df=167, t=7.606, p-value<0.0001.

[2] Post-hoc tests for *P. biforme*: df=154.17, t=-1.113, p-value=0.8752; *P. nalgiovense*: df=165.34, t=5.283, p-value<0.0001 and *P. salamii*: df=61.68, t=5.154, p-value<0.0001.

[3] Post-hoc tests for *P. biforme*: z=4.586, p-value=0.0001; *P. nalgiovense*: z=0.794, p-value=0.9687 and *P. salamii*: z=0.034, p-value=1.

[4] Post-hoc tests for *P. biforme*: df=51, t=-0.294, p-value=0.997; *P. nalgiovense*: df=51, t=-3.779, p-value=0.0052 and *P. salamii*: df=51, t=-0.136, p-value=1.

[5] We present F values for linear models when the response variable followed a normal distribution and Chi-square values otherwise.

 Figure legends

Figure 1. Rooted maximum likelihood population trees, principal component analysis

(PCA), STRUCTURE and splitstree results based on genome-wide single nucleotide

polymorphisms (SNPs) of *Penicillium salamii* **(A) and** *P. nalgiovense* **(B), with**

information on genomic architecture types and mating-types. Bootstraps are indicated for

the well-supported nodes (i.e. bootstraps higher than 70) and the scale indicates branch length

(substitution rate per polymorphic site). The strain geographical origins are shown with colors

(blue: Europe, purple: America, yellow: Australia, light green: Africa, green: Middle East).

The color circles indicate environment of isolation, with dry-cured meat in red and other

environments in blue. Architecture types correspond to groups of genomes with high

collinearity. For the STRUCTURE plot, each bar represents an individual, with its estimated

proportions of genetic information assigned to different population clusters, represented by

colors, for K = 4 for *P. salamii* and K = 2 for *P. nalgiovense*. See Fig. S2 for barplots at other

K values, giving similar patterns. In the PCA, the shape indicates the environment of isolation,

and the color follows the STRUCTURE results. For each PCA axis, the percentage of

variance explained is indicated between brackets. For the splitstrees, the bubbles colors follow

the STRUCTURE results. For *P. nalgiovense*, the two most divergent strains CBS297.97 and

DTO204.F1 were removed from the left splitstree. The *P. nalgiovense* tree has been rooted

with *P. rubens* instead of *P. chrysogenum* because of the availability of a better genome

assembly at the time of the analysis and of the very high genetic similarity between *P. rubens*

and *P. chrysogenum*.

1429

Figure 2. Synteny and similarity between the *Penicillium nalgiovense* **and** *P. salamii*

reference genomes and location of the horizontally transferred regions (HTRs). Contigs

based on long read assemblies for the genomes of *P. nalgiovense* ESE00252 are shown at the

top and *P. salamii* LCP06525 at the bottom. Only contigs larger than 300 kbp or matching a

HTR are shown. The links indicate matching blocks larger than 1 kbp, colored according to

averaged nucleotide identity (%) between *P. salamii* genomes and *P. nalgiovense* ESE00252.

Only links present in at least half of the pairwise comparisons are shown, or those matching

HTRs and being at least half covered by non-overlapping contigs (to avoid taking repeated

elements as evidence of presence). The HTRs are identified on the contigs by colors as

indicated on the figure key, on which HTRs are magnified 2X for clarity.

1440

Figure 3. Presence and identity level of the eight largest putative horizontally

transferred regions (HTRs) in *Penicillium nalgiovense* (top) and *P. salamii* (bottom)

genomes. The color darkness on the contigs indicates the average nucleotide identity (%) of

HTRs present in each strain compared to the reference genomes. White indicates complete

lack of the corresponding region or ambiguous mapping (mostly due to repetitive elements).

The length of each rectangle is proportional to the size indicated by the scale. The shades

under the names of the strains on the left are colored according to the genetic clusters as in

Figure 1. The strain geographical origins are shown on the right by the color of the squares

(blue: Europe, purple: America, yellow: Australia, light green: Africa, green: Middle East).

The color circles indicate environment of isolation, with dry-cured meat in red and other

environments in blue. See figures S6 to S8 for the information on presence/absence of all the

HTRs in all the strains.

1453

Figure 4. Presence and similarity of the eight largest HTR (horizontally transferred

regions) across 54 *Penicillium*, *Aspergillus* or *Monascus* genomes. In each genome, the

fraction of each HTR covered by non-ambiguous matches was obtained by dividing the total

length of non-overlapping matches over the total length of the corresponding HTR. The

percentages of identity (% identity) correspond to the number of identical bases to the

consensus HTR sequences over the total matched length. The phylogenetic tree on the left

was built from 1102 single-copy genes with Astral v5.6.3. (Zhang et al., 2018) and rooted on

the *Aspergillus* clade. The distance in the internal branches is expressed in coalescent units.

Stars on the right end of each branch correspond to a full support of the relationship between

the four leaves surrounding this branch.

1464

Figure 5. Proteolysis (A) and lipolysis (B) dynamics, and growth (C) and spore

production (D) in response to different salt concentrations in *Penicillium biforme*, *P.*

nalgiovense* and *P. salamii* and their respective sister species, *P. palitans*, *P. chrysogenum

and *P. olsonii*. (A) and (B): The x-axis represents the number of days since the start of the

experiment, the y-axis represents the front of lysis for proteolysis or lipolysis in mm in

measure tubes. The colored area around each logistic line represents the standard error, and

colors correspond to the environment of isolation (red=dry-cured meat, blue and

turquoise=other environments excluding dry-cured meat). Stars indicate significant post-hoc

tests (p-value < 0.05), squares marginally significant ones (0.05 < p-value < 0.1). (C) Growth

rates represented as colony area in cm^2 at days 8 (left panel) and 17 (right panel) post-

inoculation at different salt concentrations (X axis). (D) Spore production is estimated at day

17 for the same species and the same salt concentrations. In (C) and (D), circles indicate

species used for dry-cured meat production, i.e., *P. biforme*, *P. nalgiovense* and *P. salamii*,

and squares indicate their respective sister species not used for producing dry-cured meat (*P.*

palitans, *P. chrysogenum* and *P. olsonii*, respectively). The strains were collected either from

dry-cured meat (red) or from other environments excluding dry-cured meat (blue). The error

bars correspond to the standard error. We did not detect any significant difference between

dry-cured meat and non-dry-cured meat samples in each species with post-hoc tests.

1483

Figure 6. Degree of white, color composition and pictures on malt extract agar medium

of *Penicillium nalgiovense* and *P. chrysogenum* (A)*, P. salamii* and *P. olsonii* (B)*, P.*

***biforme* and *P. palitans* (C), separated by their environment of isolation.** The left panel

plots show the degree of white on a scale from 50 to 550 (represented as dark grey to light

grey). The whitest point for *P. salamii* is circled in black for clarity. The middle panel shows

the distribution among the red, blue and green values in the color of each *Penicillium* strain.

The colors and shapes of the points correspond to the environment of isolation (red

circle=dry-cured meat, blue circle=non-dry-cured meat, blue triangle=non-dry-cured meat

sister species). Stars indicate significant post-hoc tests (p-value < 0.05). The right panel shows

pictures of *P. nalgiovense, P. salamii, P. biforme* and their respective sister species on malt

extract agar medium: the first row corresponds to strains collected on dry-cured meat, the

second (and the sister species) to strains isolated from other environments.

1496