Title: Two modes of mid-frontal theta suggest a role in conflict and error processing

Authors: Vignesh Muralidharan, Adam Aron, Micheal X Cohen* and Robert Schmidt*

(* - shared senior authors)

Author affiliations: [1]Department of Psychology, University of California San Diego, 9500 Gilman Drive, La Jolla, California 92093, USA, [2]Department of Psychology, The University of Sheffield, UK, [3]Radboud University Medical Centre, Nijmegen, Netherlands, and Donders Centre for Neuroscience

Corresponding author: Vignesh Muralidharan, vimuralidharan@ucsd.edu

Number of pages: 40

Number of figures: Main Text – 6; Supplementary - 1

Number of words: 5357 (excluding Methods and Figure Captions)

Conflicts of interest: The authors declare no competing interests.

Acknowledgements: This work was supported by the National Institutes of Health (DA026452 and NS106822). MXC is funded by an ERC-StG 638589 and a Hypatia fellowship from the Radboud University Medical Center. RS was supported by the EU Horizon 2020 programme through the FET Flagship Human Brain Project (HBP-SGA3, 945539).

ABSTRACT

Mid-frontal theta is well-known to be involved in conflict and error-processing. In paradigms known to evoke response conflict, the temporal sequence of events leading up to conflict or error resolution and theta's involvement in such scenarios is not well understood. Using single-trial analyses of electrophysiological data from participants performing the Flanker (N = 28) and Simon paradigm (N = 18), we probed the relationship between transient theta activity and the underlying metrics of response conflict. We specifically investigated "partial errors", in which a small burst of muscle activity in the incorrect response effector occurred, quickly followed by a correction. We found evidence for two modes of theta activity, one of which was strongly related to the partial errors. The temporal distribution of this mode of theta was different in the partial error trials compared to the correct trials with no partial error. Furthermore, we saw that this mode of theta occurred later in trials in which a full error was committed. Our findings point to a possible role of theta in error-correction. This adds to the wider theoretical understanding of mid-frontal theta and the possibility of multiple functional roles of theta in different contexts.

INTRODUCTION

Response conflict monitoring is a key process that helps us to select appropriate actions among several competing ones. The neural mechanisms associated with conflict and error monitoring have been linked to increases in theta-band activity (4-8Hz) in mid-frontal brain regions (Cavanagh & Frank, 2014; Cohen & Cavanagh, 2011; Cohen & Donner, 2013; Cohen & van Gaal, 2014; Nigbur, Cohen, Ridderinkhof, & Stürmer, 2012) and coherence in the theta-band between mid-frontal cortex and basal ganglia, specifically implicating the hyperdirect pathway (Zavala et al., 2016). Theories on mid-frontal control processes have related theta activity increases to the need for control, which arises in situations requiring the resolution of a conflict at the level of stimulus, response, or even rewards (Cavanagh & Frank, 2014). Mechanistically, this has been attributed to theta's ability to modulate reaction times by allowing time for evidence accumulation by increasing decision thresholds (Cavanagh et al., 2011). However, while theta increases are seen in trials in which conflict was resolved successfully, they also occur in trials after an error was committed, suggesting multiple roles of theta with potentially different neural sources (Swick & Turken, 2002). Therefore, it is unclear how the temporal dynamics of theta within a trial relate to an impending error and to the processes that allow the correct response to get elicited.

In response-conflict paradigms, a subset of correct trials contain "potential errors," meaning that the participants started to initiate an incorrect response (measured through electromyogram or force grips), but quickly corrected the erroneous action to make a correct response (Cohen & van Gaal, 2014; Rochet, Spieser, Casini, Hasbroucq, & Burle, 2014; Vidal,

Burle, & Hasbroucq, 2019). Partial error trials are useful because they provide temporally precise single-trial insights into the amount of response conflict experienced by the research participants.

Recent studies on action and motor control have revealed that average increases in oscillatory power in a particular frequency band can be decomposed into transient increases or burst-like activity in single trials (Feingold, Gibson, DePasquale, & Graybiel, 2015; Little, Bonaiuto, Barnes, & Bestmann, 2019; Torrecillos et al., 2018). For instance, beta activity at single trials, also referred to as beta bursts, in motor and frontal cortical areas are more tightly linked to behavior compared to trial-average increases in oscillations (Little et al., 2019; Muralidharan, Aron, & Schmidt, 2021; Muralidharan & Aron, 2021; Soh, Hynd, Rangel, & Wessel, 2021). Here we extend this approach to theta oscillations and provide evidence that transient, mid-frontal theta events relate to cognitive and motor processes of conflict induced error and its correction.

In this study we analyzed EEG data from two well-known paradigms known to elicit response conflict: the Flanker task and the Simon task. We examined conflict-related mid-frontal theta by decomposing average theta into single-trial theta "events." We then investigated the relationship between these theta events and muscle-level metrics of conflict, i.e. partial errors at a single trial level. We found that theta activity could be dissociated into several modes based on the presence, or absence, of a partial error. While one mode of theta was linked to the stimulus conflict, in line with a role of theta in conflict resolution (Cohen & Cavanagh, 2011; Cohen & Donner, 2013; Cohen & van Gaal, 2014; Cohen & Ridderinkhof, 2013), a second mode of theta was strongly linked to partial errors, pointing to a possible role of theta also in error-correction.

METHODS

Participants

There were in total forty-six participants in the study, with 28 performing the Flanker task and 18 performing the Simon task. Subjects were recruited from the University of Amsterdam community, and participated in these studies in exchange for course credit or 14 Euros. Each study was approved by the local ethics committee and subjects signed an informed consent document. Subjects had normal or corrected-to-normal vision and were self-reported free of neuro-logical disorders and history of physical head trauma. Four participants from the Flanker task and three participants from the Simon task were excluded due to extremely noisy muscle activity which led to sub-optimal detection of trials with partial errors and their onset timings.

Tasks and Data

We analyzed behavior and neural data from two different tasks which are known to induce response conflict: the Flanker task and the Simon task. Non-overlapping results from these data have been published previously (Cohen & Ridderinkhof, 2013; Cohen, 2015). The Flanker task was a modified version of the standard task, where participants responded to a central target which was either the letter T or I with their right or left thumb (Appelbaum, Smith, Boehler, Chen, & Woldorff, 2011). They were surrounded by distractor stimuli, two on either side of the central target such that there were three stimulus conditions. First was that the distractors were the same as the central target, or bilaterally congruent. Secondly, the distractor could be either left or right incongruent or partially incongruent, i.e., flanked by distractors different from the central target on either the right or left side respectively. Finally, the flanked distractors were all

different from the central target or bilaterally incongruent. Conflict in this task is created due to the presence of the different surrounding stimuli. The stimulus was presented for 150ms, and the inter-stimulus interval was 1300-1700ms.

The Simon task was a simple visual association with a colored circle mapping onto either a right or a left response (Cohen & Ridderinkhof, 2013). The colored circle stimulus was presented on either the right or left side of a fixation cross, thus leading to visuo-spatial conflict. Conflict and stimulus repetition effects were balanced by using stimuli of four different colors. In this task, conflict occurs when the correct response is opposite to the side the stimulus is presented in, for instance a blue-colored circle presented on the left side of the screen which requires a right-hand response, also called the Incongruent trials. If present on the same side, they are the Congruent trials. Stimuli were presented on the screen for 200ms with 1000ms to respond and a varying inter-trial interval of 700-1200ms.

Electroencephalography and Electromyography

EEG/EMG acquisition and analysis procedures were the same across both studies. EEG data were acquired at 512 Hz from 64 channels placed according to the international 10–20 system, and from both earlobes, using Biosemi equipment. Electromyographic (EMG) recordings were taken from the flexor pollicis brevis muscle of each thumb using a pair of surface electrodes, placed on a subject-by-subject basis approximately 5 mm apart on the thenar eminence.

Data Analysis

All analyses were performed using EEGLAB13 and MATLAB R2016b.

Behavior and Partial Errors

The reaction time was estimated for the button press responses; hereafter referred as response time (RT). A trial was classified as a correct or an error trial based on the response outcome, which we used to compute the accuracy for a given stimulus condition. We then identified the partial error trials using an algorithmic approach in line with a previous study (Cohen & van Gaal, 2014). We took the derivative of the EMG traces from both hands, Z-transformed across the entire trial period and then rectified. This controlled for several things, including differences in impedance and signal strength across subjects and hands. A trial was then classified as a partial error if the differentiated and z-normalized EMG trace of the incorrect hand in that trial exceeded two standard deviations within the time of the stimulus onset and the response time. Additionally, the peak of the partial error had to be more than twice the maximum of the baseline activity seen 300ms prior to the stimulus onset. Using this approach, we could reduce the chance for detecting partial errors which may have occurred due to noisy EMG. We will refer to these trials as partial error trials. We then estimated the EMG onsets of the partial errors and correct response, by first starting at the time of peak of both the EMG traces and back-tracking till the EMG trace for the corresponding hand fell below 20% of the peak for five consecutive milliseconds. In cases where the peak did not fall below 20% till the onset of the stimulus, we used 30% as the threshold to determine EMG onsets. Given that the number of trials with partial errors within a participant is low (~20%), for comparison we picked the correct trials which were

matched for both the response times and stimulus type (for instance the bilaterally incongruent, partially incongruent and bilaterally congruent in the Flanker scenario). To do so for each detected partial error trial, we picked a correct trial which had the closest response time to it and had the same stimulus type. We will refer to these trials as the full-correct matched trials.

EEG

Offline, continuous EEG data were high-pass filtered at 0.5 Hz and epoched from −1.5 to +2 s surrounding stimulus onset of each trial. All trials were visually inspected and those containing facial EMG or other artifacts not related to blinks were manually removed. Independent components analysis was computed using eeglab software (Delorme and Makeig, 2004), and components containing blink/oculo-motor artifacts or other artifacts that could be clearly distinguished from brain-driven EEG signals were subtracted from the data.

Spatial Filtering using GED: Mid-frontal source

In both task datasets, we performed a guided source separation analysis, i.e., generalized eigen decomposition, GED (Blankertz, Tomioka, Lemm, Kawanabe, & Muller, 2008; Cohen, 2017, 2022; Parra & Sajda, 2003), to obtain a mid-frontal spatial filter for extracting the conflict-related theta activity. GED allows for a more guided approach to spatial filtering by allowing us to estimate a spatial filter based on previous hypotheses regarding the spatial topography, frequency band and time-period from the data. Here, we were interested in mid-frontal theta activity (4-8Hz) around the time where conflict related processing would occur

(i.e., in relation to the stimulus onset, 200ms to 100ms after the mean RTs in the high conflict condition). Thus, by pooling all correct trials and filtering the data in theta frequency (4-8Hz), we applied GED with respect to the broadband data, i.e., GED was performed between the theta filtered covariance matrix and the broadband covariance matrix (using the *eig* function in MATLAB). This provided us with spatial filter weights, aka GED components, which maximized the signal to noise ratio in theta frequency band. These weights were then forward-projected onto the theta-filtered covariance matrix to obtain spatial activation maps to visualize each component's scalp distribution. We then selected the filter having a mid-frontal spatial distribution. In scenarios where more than one mid-frontal filter was obtained, we picked the one which explained the maximal variance in the data, as estimated using the generalized eigenvalues. Finally, the epoched EEG data were projected onto the selected mid-frontal spatial filter for time-frequency as well as theta event analyses.

Time-Frequency analyses

Time-frequency analysis was done in EEGLAB13 using the *newtimef* function. For the event-related spectral perturbations (ERSPs), the time-frequency decomposition was done using Morlet wavelets with frequencies ranging from 4-30Hz. We started with 3 cycles at the lowest frequency, linearly increasing as the frequencies increased such that it was 11.25 cycles at the highest frequency. The ERSPs were baselined to a period prior to the stimulus (-500 to -100ms in both task datasets). From the ERSPs of the condition potentially capable of evoking maximal response, i.e. the bilaterally incongruent trials in the Flanker and Incongruent trials in the Simon paradigm, we estimated at an individual level the peak-theta frequency within the stimulus to the

corresponding condition's mean RT. Our rationale for this approach was to select a participant-specific theta which would best reflect the underlying conflict-related processing. We used the peak-theta frequency value to extract single-trial theta events for each participant.

Extracting Single-trial Theta Events

From the GED component time series, we extracted single-trial theta events by first applying a temporal filter. The filter was a frequency-domain gaussian with a peak at the individual's peak-theta frequency and a full-width half maximum of 5 Hz. We found that this was a good tradeoff between obtaining frequency specificity and temporal precision. The complex analytic time-series thus obtained was subjected to a Hilbert transform to compute its amplitude/envelope. From the theta amplitudes in a baseline period (-500ms to -100ms prior to the stimulus onset), we derived threshold values to ascertain which periods of time could be classified as a theta event. Firstly, we used an event definition threshold (median + 1.75SD). Any theta amplitude above this threshold was considered a theta event. Next, once all the theta events in a trial were evaluated, we lowered the threshold value in order to estimate both the timings and durations of the events identified using the higher threshold, i.e., an event length threshold (median +1.0SD). This method helps prevent brief increases in amplitude to be classified as events. Moreover, it has shown to prevent underestimation of the duration of such transient single-trial events (Little et al., 2019). From the theta events we first estimated the theater event probability. Theta event % was estimated by binarizing the time periods in a trial where the theta amplitude was above the event length threshold and then averaging across trials within each condition.

Within a trial, three theta timings were estimated: the theta onset which is the time at which theta exceeds the event length threshold, the theta peak which is time corresponding to the peak of the theta event, and theta offset which is the time when the theta event falls below the threshold (also see Fig. 2b). For correct trials, theta events (i.e. the peak) that occurred before the response times were considered. If a trial had more than one theta event with peak times t_1 and t_2, it was split into two trials with events at time t_1 and t_2 respectively. These timings were then related to the timings of the conflict metrics, i.e., the partial error onset and the correct response onset within a participant using Pearson's correlation (r).

Gaussian Mixture Modeling of Theta Events

The theta timings (specifically the onset) were analyzed in relation to the partial error onset times using gaussian mixture modeling. We used this approach to dissociate the contribution of different theta modes, given that we observed bimodality in the shape of the difference distributions between theta onset and partial error times. The expectation maximization algorithm was used to fit models with K gaussians. In our case K was set to either 1, 2 or 3. The covariance type was selected as full, with each component assuming any possible shape. We also set a 1% regularization to the data, with a maximum iteration of 1000. From the model fits, we obtained the Akaike Information Criterion (AIC) value and estimated the probability of each model (K = 1, 2 and 3) as following $P_K = exp(\frac{AIC_{min} - AIC_K}{2})$. We then tested which of the models best predicted the underlying distribution of the theta timings. We also evaluated the mean and variance of the fitted gaussians models.

Fixed Effects Analysis: Joint Probability Distributions

The temporal structure of the theta activity was ascertained by computing the joint probability distribution of the theta onset times and the correct response EMG onset times. We pooled trials from both tasks to perform fixed-effects analysis. The probability value for a particular correct response and theta onset, say for instance at 300ms and 100ms respectively, was computed by first taking all the correct response onsets within 300 ± 25ms, and then looking at the number of trials that had a theta onset within 100 ± 25ms. This was then done for a range of correct response onsets (200 to 800ms) and theta event onset (-100 to 800ms). These probability maps were then thresholded for statistical significance by performing a permutation test against a null distribution which considered that the theta onsets were uniformly distributed across time (i.e. -100ms prior to stimulus to response time + 400ms) and corrected for multiple comparisons using false discovery rates (Benjamini & Hochberg, 1995). The same map for the trials in which a full error had occurred was computed differently. Since in these trials there are no correct response times, a pseudo "correct" RT was estimated by adding the individual's mean correction time to the error response, i.e. the time from the partial error to the correct response onset. For the same trials, we considered those theta events which occurred between the stimulus and error response + participant-specific max correction time (i.e., average of the correction times greater than the 75th percentile). This was done to make sure we consider the events which occur even after the full error EMG onset, under the assumption that there could be temporal changes between trials where the error was partial compared to when it became a full error. This also

ensures that the time window is long enough to detect any theta that might have occurred later, but within the potential time that would have been available to correct the error.

Statistical analyses

Data were analyzed using several different approaches, as appropriate for the specific comparison (detailed in the Results section). Some comparisons were performed using t-tests, either one sample or paired sample tests. The effect sizes for these tests were estimated using the Bayes factor in favor of the alternate hypothesis (small: BF_{10} = 1-3; medium: BF_{10} = 3-10; large: BF_{10} > 10). For distributions violating normality, we performed non-parametric Wilcoxon tests. For data with multiple levels, we used repeated measures (rm) ANOVA followed by post hoc t-test comparisons between specific conditions, corrected for multiple comparisons using Bonferroni correction. We reported the effect sizes for the ANOVA as partial eta square (η_p^2), interpreted as a small (0.01-0.06), medium (0.06-0.14) and large effect (>0.14). Data sphericity was controlled using the Greenhouse-Geisser correction. For the time-frequency analyses, the ERSPs, we used a non-parametric bootstrap method to compute regions of significance in the time-frequency space. We then used the false discovery rate to correct for multiple comparisons in these cases (Benjamini & Hochberg, 1995). Finally for the group-level comparisons of correlations we first used the Fisher-transform to convert the correlations to z values and then used rmANOVA and/or t-tests to compare between specific conditions. All data are presented as mean ± s.e.m.

RESULTS

Behavior

We analyzed behavior of subjects in the Flanker and Simon task by measuring RTs and accuracies in different trial types depending on the stimulus congruence. In a trial of the Flanker task the distractor stimuli could be different from the target stimulus (i.e. high conflict, bilaterally incongruent), the distractor stimuli on only one side could be different from the target stimulus (i.e. medium conflict, partially incongruent), or the distractor stimuli on both sides could be identical to the target stimulus (i.e. low conflict, bilaterally congruent). In line with previous observations, behavior from the Flanker task exhibited the expected patterns of slower RTs and decreased accuracy for incongruent compared to congruent trials (Appelbaum et al., 2011). The slowest RTs were observed in the bilaterally incongruent trials, followed by the partially incongruent and bilaterally congruent trials ($F_{1.5,39.2} = 219.5$, $p < 0.001$, $\eta_p^2 = 0.9$, Fig. 1a). The accuracy in each condition also scaled with the stimulus condition, with the lowest accuracy in the bilaterally incongruent trials ($F_{1.2,31.3} = 50.9$, $p < 0.001$, $\eta_p^2 = 0.7$).

In the Simon task, the stimulus could appear on the side of the fixation cross that is opposite to the correct response (high conflict, incongruent trial) or on the same side as the correct response (low conflict, congruent trial). Furthermore, behavior depends on the previous trial condition, i.e., the well-known sequence or the Gratton effect (Egner, 2007; Gratton, Coles, Sirevaag, Eriksen, & Donchin, 1988). Accordingly, conflict is increased when the congruence switches between two successive trials, which lead to slower RTs for both congruent-incongruent and incongruent-congruent trial sequences (Fig. 1b). In contrast, congruent-congruent and incongruent-incongruent trial sequences had faster RTs, reflected by a significant interaction

$(F_{1,17} = 48.5, p < 0.001, \eta_p^2 = 0.7)$. The same interaction effect was observed for the accuracies as well, with lower accuracy in the incongruent trials compared to the congruent, especially when the previous trial was congruent and vice versa $(F_{1,17} = 26.9, p < 0.001, \eta_p^2 = 0.6)$. Therefore, behavior in both tasks was affected by conflict as expected (Appelbaum et al., 2011; Cohen & Ridderinkhof, 2013).

Figure 1. Behavior and Partial Error Metrics. a) Response times and accuracies in the Flanker task are shown for the bilaterally incongruent (BI), partially incongruent (PI) and bilaterally congruent (BC) trial types. **b)** Response times and accuracies in the Simon task for the congruent and incongruent trials. Lower-case letters indicate whether the previous trial was congruent (c) or incongruent (i); upper case letters indicate the corresponding for the present trial. **c)** The left plot shows the percentage of trials with partial errors in each participant, with the red dotted line showing the average across all participants. The right plot shows the percentage of partial errors based on the trial type (dots indicate individual subjects, bars show average). **d)** Analysis corresponding to **(c)** is shown for the Simon task, with the right plot showing the proportion of partial errors as a function of previous trial conditions. **e)** Left plot shows for the Flanker task, the percentage of partial error trials as a function of response times. The middle plot

shows the response times of all trials with no partial errors (i.e. full-correct trials, FC), of partial error trials (PE) and of trials matched for response time and stimulus type (full-correct matched trials, FC_{Match}). The left plot average correction time (dots indicate individual subjects, bars show average), which is time from the partial error EMG onset to the correct response EMG onset. **f)** The same analysis as **(e)** is shown for the Simon task.

EEG analyses focused on partial error trials, defined as correct trials in which the incorrect (prepotent) response was partially activated (observed as a small burst in the EMG of the incorrect response hand), but did not develop into a full response and was followed by the correct response. In both the datasets, we identified partial error trials (see *Methods* for more details) and confirmed that their occurrence was as expected (Cohen & van Gaal, 2014). On average there were ~20% of trials with partial errors in both tasks (Fig. 1c and 1d). The percentage of partial errors scaled with the level of conflict, with more partial errors seen in the bilaterally incongruent trials in the Flanker task ($F_{1.1,26.1} = 94.1$, $p < 0.001$, $\eta_p^2 = 0.8$). In the Simon task, there was again a modulation based on previous trial condition, with more partial errors seen in incongruent-congruent and congruent-incongruent trial sequences compared to congruent-congruent and incongruent-incongruent sequences, leading to a significant interaction ($F_{1,14} = 18.4$, $p < 0.001$, $\eta_p^2 = 0.6$). The percentage of partial errors also increased with slower RTs (Fig. 1e and 1f). As a result, all full-correct trials, i.e. trials that were correct and did not have a partial error based on our algorithm, were faster on average compared to the partial error trials (Fig. 1e and 1f). To control for the fact that there were more full-correct trials than partial error trials and that they had faster RTs, we selected a subset of trials from the full-correct trials to match for both RTs and stimulus type, referred to as full-correct matched trials (see *Methods* and Fig. 1e and f). These trials were then used for comparison with the partial error trials. Finally, we also estimated the correction time, which is the time taken from the onset of the

partial error EMG to the onset of the correct response EMG. These typically range from 100-200ms (mean ± SEM: Flanker: 142 ± 6ms; Simon: 178 ± 6ms). Next, we sought to extract mid-frontal theta activity to understand the trial-by-trial dynamics and relationship to the mechanisms resolving response conflict.

Theta event probability increases in Partial Error trials

To identify and characterize single-trial mid-frontal theta events, we crafted a spatiotemporal filter that maximized midfrontal theta-band for each participant. The EEG data were projected onto the selected filter and then temporally filtered at that individual's peak theta frequency. Next, using the median theta activity in a baseline period prior to the stimulus, a threshold (event definition threshold, median + 1.75SD) on the theta baseline amplitude was defined. Using this threshold, peaks of theta activity that exceeded this threshold were identified. From the identified peaks, using a lower threshold (or event length threshold, median + 1SD), a theta "event" was defined as any time period where the theta amplitude was above this threshold (Fig. 2a, see *Methods* for more details). Within a participant, there was both an increase in the number of theta events and variability in the timing of these events in and around the time of response (Fig. 2a, bottom left). Furthermore, the duration of these theta events irrespective of the stimulus condition or time window were on average around 200-250ms (Flanker: 229 ± 6ms, Simon: 259 ± 4ms) pointing towards their transient nature in comparison to the entire response which was around 500ms.

Figure 2. Theta Event Extraction. a) Shows the analysis steps for extracting theta events within a participant. Top left shows a mid-frontal spatial filter which was identified in an exemplar participant using GED in the Flanker task. The data were then projected on to the selected mid-frontal spatial filter. Top right shows the participant-specific peak theta estimated by looking at the normalized power spectrum of those trials capable of inducing response conflict the most, i.e., the bilaterally incongruent trials in the Flanker task and the incongruent trials in the Simon task (top right). All trials were filtered at this peak theta frequency using a frequency domain gaussian. Bottom left shows theta amplitude distribution from a baseline period (-500 to -100ms in relation to the stimulus) used to evaluate the event thresholds. To identify a theta event, median + 1.75SD was used as the threshold (red dotted line), and once an event was detected, its onset and offsets were determined by a lower threshold median + 1.0SD (black dotted line). Bottom right shows the raster plot of the timings of the theta peak for all trials, with the dotted blue line representing the mean response time across all trials. **b)** Example trials within a participant showing the theta event, its timing (onset, peak and offset) in relation to the timing of the conflict related muscle activity, the partial error onset, the correct response onset and the response time.

The classic event-related spectral perturbation analysis (ERSPs) for all conditions of the

mid-frontal projected data revealed the typical increase in theta power for both tasks (Fig. 3a).

Furthermore, a comparison of the theta power increase in the full-correct matched trials and

partial error trials confirmed higher theta power in partial error trials in both tasks, which is an

observation from earlier studies (Cohen & van Gaal, 2014). At a single-trial level we found that

theta oscillations manifested as transient events and could be characterized by an onset, a peak and an offset (partial error trials shown in Fig. 2b).

We then examined how the number of theta events is modulated over time and found that the increase in theta power (Fig. 3a) could be attributed to a significant increase in the number of theta events in the partial error trials compared to both full-correct and full-correct matched trials in both task paradigms (Fig. 3b). This was also visible in single participants suggesting that some theta activity could have been evoked by the partial errors. Furthermore, this highlights the utility of decomposing theta into single-trial events in providing a better characterization of neural activity occurring during both trial scenarios in comparison to just oscillatory power. Next, we investigated how the timing of these single-trial theta events relate to the conflict-related EMG timings.

Figure 3. Theta Event % in Partial Error and Full-correct trials. a) Group-level ERSPs (p <0.05 FDR corrected maps) for both tasks, showing all trials and the difference between the partial error and full-correct matched trials. The arrow represents the region of theta power difference seen in both tasks for the PE-FC$_{Match}$ ERSPs. **b)** The left plot shows theta event % seen in the partial error trials compared to all full-correct and the matched trials in the

Flanker task. The middle and right plots are single subject examples of theta events sorted by response time. **c)** Same as (**b**) but for the Simon task.

Timing of the theta events relate to response correction

For trials in which theta events occurred between the stimulus onset and the response time, we estimated the time when theta events occurred in relation to the EMG signals of partial errors and correct responses. Firstly, we established the general timeline of theta events and EMG responses aligning theta and EMG signals to the time of the stimulus. On average, in both tasks, theta event onsets preceded partial errors, followed by the peak of the theta event. After the peak of the theta event, the EMG signal of the correct response occurred, which was then followed by the offset of the theta event (Fig. 4a and 4b).

Interestingly, in relation to the stimulus, the theta event timings (onset, peak and offset) occurred later for the partial error trials compared to the full-correct matched trials. This suggests that differences in the timing of the theta events could relate to the partial error occurrences.

Secondly, we computed the same timings, but now aligned to the correct response EMG onsets (Fig. 4c and 4d). The overall timing relationships remained the same, i.e., theta onsets on an average occurred before the partial errors, followed by theta peak and offset. However, now the difference in the timings between the partial error and full-correct matched trials was evident only for the theta offsets, suggesting that theta events occurring in partial error trials are longer in duration. Indeed, that was the case as average theta durations were longer for the partial error trials compared to the full-correct matched trials (Flanker: PE = 276 ± 8ms, FC_{Match} = 261 ± 8ms,

W = 229, p = 0.023, BF_{10} = 2.1; Simon: PE = 315 ± 9ms, FC_{Match} = 286 ± 5ms, $t_{1,14}$ = 3.5, p = 0.003, BF_{10} = 14.2, Fig. 4c and 4d insets).

Figure 4. Timing relationship of theta events to conflict-related EMG timings. a) Average timings of the theta event onset, partial error onset, theta peak, correct response onset and the theta offset in relation to the stimulus for the partial error and the full-correct matched trials in the Flanker task. **b)** Same as **(a)** but for the Simon task. **c)** The theta and EMG timings in relation to the correct response onset. Inset shows the average theta height and duration across participants, which are both higher for the partial error trials. **d)** Same as **(c)** but for the Simon task. **e)** The relationship (z-transformed correlation) between theta timings and the EMG timings in the partial error trials in the

Flanker task. The relationship is stronger for the correct response onset for all theta timings. **f)** Same as **(e)** but for the Simon task.

To further understand the temporal relationship between theta events and the EMG responses, we performed trial-by-trial correlations at the participant level in each task. We correlated each of the theta time points (onset, peak time and offset) with the EMG time points in both the partial error and full-correct trials. The correlation coefficients were analyzed using a 3x2 repeated measures ANOVA with two factors: theta time points (onset, peak and offset) and EMG time points (onset of partial error EMG and onset of correct EMG in the partial error trials). We saw a main effect of both the theta time points (Flanker: $F_{1.5,35.5} = 11.3$, p < 0.001, η_p^2 = 0.3; Simon: $F_{1.2,16.8} = 4.6$, p = 0.040, $\eta_p^2 = 0.25$) and EMG time points (Flanker: $F_{1,23} = 60.5$, p < 0.001, $\eta_p^2 = 0.7$; Simon: $F_{1,14} = 18.8$, p < 0.001, $\eta_p^2 = 0.6$) in both task datasets.

Investigating each condition, we observed that all theta time points had a stronger relationship with the correct response EMG time points compared to the partial error time points (Fig. 4e and 4f). Specifically, the correlation with the correct response was strongest for the theta peak time. This highlights two patterns in the data: one is that theta activity seems to be linked to the emergence of the correct response, which fits well with its existing role of theta holding responses from completion till enough evidence has been accumulated for resolving the conflict (Cavanagh & Frank, 2014; Cavanagh et al., 2011); secondly, the partial error onsets were seen between the theta onset and peak time. Given this close proximity of the partial error to the onset of theta, there is a possibility that theta might be involved in error-related processing. Thus, we looked into the timing relationship between theta and the error more specifically.

Event onset times reveal two modes of mid-frontal theta activity

Although the theta event onset on average occurred prior to the partial error onset (Fig. 4a and 4b), within participants there was considerable variability in the timing of theta activity. Furthermore, we noticed that in individual participants the distribution of theta onset times relative to the partial error times was typically bimodal (Fig. 5a and b insets). Using gaussian mixture modeling with K = 1, 2 and 3 number of gaussians, we probed which model best captured individual distributions (see *Methods* for details on the modeling). We found that the model with two gaussian components fitted the distributions better than one (Fig. 5a and b; Flanker: W = 30.0, p < 0.001, BF_{10} = 82.9; Simon: W = 14.0, p = 0.007, BF_{10} = 16.6). However, this increase in the model probability was not just due to increase in the parameters, as K = 3 did not improve the fit in comparison to K = 2 (Flanker: W = 174.0, p = 0.509, BF_{10} = 0.2; Simon: W = 81.0, p = 0.252, BF_{10} = 0.4).

For each participant we then fitted the parameters using the K = 2 model, yielding a mean μ and a variance σ for each gaussian. The bimodal distribution of theta supported by the best fit of the K=2 model, suggested two modes of theta activity. The first mode occurred prior to the partial error onset (**G1** mean ± SEM, Flanker: -333 ± 62ms and Simon: -185 ± 26ms), while the second mode occurred around the partial error onset (**G2** mean ± SEM, Flanker: 2 ± 10ms and Simon: 11 ± 9ms). In addition, G2 had a smaller variability than G1 in both tasks (Fig. 5a and 5b bottom right), in line with interpretations tying the G2 theta events to the possible error that could have been committed. As the exact timing of the G2 theta events relative to the partial

errors possibly constrains potential functions related to error monitoring or correcting, we next examined the different theta modes in more detail.

In each participant we grouped theta events into G1 and G2 events by comparing the probability density functions of the two fitted gaussians at a given theta onset time. The theta event was then assigned to either G1 or G2 based on which had the higher relative likelihood. We then correlated the theta onset times belonging to G1 and G2 to the partial error EMG timings. We found that the partial error onset time points correlated stronger with the G2 theta onset time points compared to G1 (Fig. 4c and 4d; Flanker: $t_{1,12} = 5.3$, $p < 0.001$, $BF_{10} > 100$; Simon: $t_{1,12} = 3.8$, $p = 0.003$, $BF_{10} = 17.6$). Furthermore, the correlation of G2 with partial errors was stronger than the correlation of G2 with correct response onsets (Fig. 4c and 4d; Flanker: $t_{1,12} = 3.9$, $p = 0.002$, $BF_{10} = 20.6$; Simon: $t_{1,12} = 4.8$, $p < 0.001$, $BF_{10} = 78.8$). This supports that theta activity can be linked to partial errors.

Figure 5. Different modes of theta activity. a) Top left plot shows the model probability for the gaussian mixture modeling (GMM) with K = 1, 2 and 3 gaussian components in the Flanker task. Top right shows a single subject exemplar for the distribution of theta event onsets in relation to the partial error onsets and the gaussians fitted using K = 2 model. Bottom left and right plots show the fitted parameters mu and sigma for both the gaussians respectively. The parameters suggest that there are two modes of theta, with one peaking prior to the partial error (G1) and the other peaking around it (G2). **b)** Same as **(a)** but for the Simon task. **c)** and **d)** Correlational analysis between the theta onsets belonging to either G1 or G2 and the EMG timings (partial error – PEMG and correct response – CEMG) for the Flanker and Simon task respectively. Analysis shows that the relationship is strongest for the theta onset times belonging to G2 and the partial error onset, suggesting that one mode of theta is strongly linked to the partial error.

To further confirm whether these findings are specific to the partial error trials, we investigated the full-correct matched trials as well. We assigned a surrogate "partial error onset" time to each full-correct trial, which was obtained from the partial error trial that was the closest

match to the full-correct trial in terms of the response time. We then looked at the same

relationship between the surrogate "partial error onset" and the G2 theta from the matched

full-correct trials. We obtained G1 and G2 distribution for the matched trials using the same

approach as the partial error trial, but using the surrogate "partial error onsets". We again found

that the correlation of partial error EMG onset was stronger for the G2 theta from partial error

trials compared to the G2 theta in full-correct matched trials (Flanker: $t_{1,12} = 2.3$, $p = 0.038$, BF_{10}

$= 2.0$; Simon: $t_{1,12} = 2.3$, $p = 0.042$, $BF_{10} = 1.8$). Due to the mixture model fitting, not all

participants had a sufficient number of trials for this analysis in either G1 or G2. Therefore, the

results are based on participants who had more than 15 data points (i.e., theta onsets) in both G1

and G2, which led to having 13/24 and 13/15 participants in the Flanker and Simon task,

respectively. To see whether the results depended on the specific inclusion criteria, we repeated

the analysis and included participants with more than 3 data points, leading to 17/24 and 14/15

participants from the Flanker and the Simon task, respectively. The results were the same for the

Flanker task. For the Simon task, all the results were the same except for the relationship

between G2 theta onsets and partial error onsets in partial error trials and full-correct matched

trials which followed the same trend but was not significant (see Supplementary Information,

Fig. S1). In addition, for these participants (N = 17 and N = 14) we examined the duration and

amplitude of theta events in partial error trials (data not shown). We found that the theta event

durations from G2 were smaller compared to those from G1 (Flanker: $324 \pm 14ms$ vs $258 \pm$

$8ms$; $t_{1,16} = 5.3$, $p < 0.001$, $BF_{10} > 100$; Simon: $352 \pm 10ms$ vs $296 \pm 11ms$; $t_{1,13} = 5.0$, $p < 0.001$,

$BF_{10} > 100$). The theta amplitude, however, was stronger for G2 theta events compared to G1

(Flanker: 0.032 ± 0.002 vs 0.034 ± 0.002 ; W = 19.0, p = 0.005, BF$_{10}$ = 50.2; Simon: 0.039 ± 0.001 vs 0.042 ± 0.001 ; W = 9.0, p = 0.004, BF$_{10}$ = 25.5).

These results support the idea that there are several modes of theta activity, likely driven by different events. The first mode (captured by G1) is possibly related to the level of detected conflict, while the second mode (captured by G2) may be more directly linked to the processing of the partial error.

Temporal structure of theta activity suggests possible role in error-correction

The stronger relationship between one of the theta modes (i.e. from G2) and the onset of partial errors suggested that there could be a different underlying theta timing distribution in the partial error trials compared to the full-correct trials. To investigate this, we performed a fixed effects analysis where we separately pooled all the partial error and full-correct matched trials from all subjects across both tasks (given that all of the theta event results were the same across both the tasks). We then looked at the joint probability distribution of the theta onset times in these trials as a function of the correct response EMG onset. The goal was to obtain the underlying temporal structure of theta across time that was yielding the correlations analyzed above (Fig. 5c and 5d).

We found that the theta activity from G1 in both partial error and full-correct matched trials was concentrated prior to the partial error onsets and surrogate partial error onsets, respectively (cyan dots in Fig. 6a). In contrast, the theta events from G2 were distributed around the (surrogate) partial error time onsets, following them in orientation (Fig. 6c). For partial error trials this was evident across the whole temporal range of correct response onsets. However, for

the full-correct matched trials, in particular for slower correct response onsets, the theta events

seemed to occur after the surrogate partial error time and were rather oriented along the correct

RTs. This is in line with the corresponding weaker correlations found above (Fig. 5c and 5d). The

tight concentration of G2 theta events around the partial errors in partial error trials points to a

potential role of theta in the correction of the error that has already started to activate the

muscles.

Figure 6. Temporal structure of theta onsets. a) The joint probability distribution of theta onsets from the G1 component of the GMM fit and the correct response onsets (dotted magenta line indicates where theta onset equals correct response onset) for the partial error, full-correct matched and full error trials. **b)** Same as (**a**) but for the theta onsets belonging to the G2 component. In both (**a**) and (**b**), the cyan line indicates the average partial error onset times for the corresponding correct response onsets. For the full-correct matched trials, these are the partial error onsets from the matched partial error trial. On the other hand for the full error trials, the cyan line indicates the full error EMG onset times and the magenta line corresponds to the potential "correct" response obtained by adding the participant-specific mean correction time.The contoured regions represent fdr corrected significance maps, showing increase in probability in comparison to a null distribution assuming that theta onsets were uniformly distributed. **c)** Single subject exemplar of the theta onset distributions for the full error trials with G1 and G2 components superimposed. **d**) The gaussian mixture model parameters (mu and sigma) for the full error and partial error trials for both tasks, specifically for the theta onsets from G2.

To examine a potential role of theta in the correction of errors, we also investigated trials in which the participants made a response error, i.e. the full error trials. In these trials, we analyzed theta events that happened in between the cue onset and the error response plus the participant-specific max correction time. The rationale was that if theta events reflect a process that prevents errors, theta would have been evoked within this time period (see Methods for more details). Since there were only a small percentage of error trials per participant, we decided to combine the data from both tasks. We again performed a gaussian mixture modeling analysis on theta onsets in the error trials. We kept the model parameters the same as for the partial error trials, i.e. K=2. Similar to the partial error trials, full error trials also had a mode prior to the full error onset (G1) and another mode in and around the full error onset (G2, single subject example shown in Fig. 6c). In the temporal structure of the full error theta onsets from G2, the probability distribution aligned well with the full error onsets, but appeared to be shifted in time compared to how theta aligned in partial error trials (Fig. 6b). This shift was evident in the parameters of the gaussian fits with the average theta onsets from G2 being significantly later than the full error onsets (Fig. 6d, 37 ± 10ms, $t_{1,38} = 3.7$, $p < 0.001$, $BF_{10} = 43.2$), and more importantly their timing was significantly later than the G2 theta onsets in the partial error trials (37 ± 10ms vs 5 ± 7ms,

$t_{1,38} = 2.9$, p = 0.007, $BF_{10} = 5.6$). Moreover, the variability of the G2 distributions from the full error and partial error trials did not change, suggesting that they could be the same process which just got engaged a bit too late in the full error trials (W = 293.0, p = 0.180, $BF_{10} = 0.5$). This supports the idea that the theta related to the partial error might be involved in error-correction.

DISCUSSION

In this study we investigated the relationship between single-trial theta events and metrics of response conflict behavior. We analyzed neural data from the well-known Flanker and Simon tasks known to induce response conflict. Average increases in theta commonly seen during these tasks appeared as short transient increases in activity, akin to bursts of oscillations typically seen in other domains of motor control, for instance beta bursts and motor inhibition (Errington, Woodman, & Schall, 2020; Jana, Hannah, Muralidharan, & Aron, 2020; Muralidharan et al., 2021; Wessel, 2020). This transient activity, which we termed "theta events," was variable in both its timing and occurrence based on the trial type with different degrees of conflict. Specifically, by investigating those trials in which an error was partially committed, as identified from the muscle activity of the incorrect response, we showed that these theta events were increased compared to matched trials without partial errors, indicating that this theta could be evoked by potential, upcoming errors. The timing of these theta events showed that they started on average prior to the partial error, and overall related to the time of the onset of the correct response muscle activity. However, on closer inspection in single trials, our analyses revealed two modes of theta, one of which was strongly related to the partial error onsets. This implies

that there are different forms of theta activity linked to the underlying processing of conflicts and potential errors.

Theta timing points to a role in error-correction

Most studies on theta and conflict-resolution focus on the relationship between theta power and RTs. However, in our study decomposing theta activity into single-trial events revealed finer temporal relationships, specifically to conflict- and error-related processing. From the identified two modes of theta, one mode (theta events from G1) was weakly modulated by the error and could have been evoked by the conflict detected at the level of the stimulus given its presence prior to the partial errors. Once conflict was detected, the rapid engagement of theta could be sufficient in most cases to prevent errors in the full-correct trials. On the other hand, in the partial error trials, this mode of theta could still have been evoked as a result of stimulus processing but perhaps not enough to prevent the error from eliciting a muscle response. For clarity we refer to this theta here as "conflict-related theta". The other mode (theta events from G2) seemed to strongly track the timing of partial errors, suggesting that this theta could also have been elicited in reaction to a potential, upcoming error. Possible functions of this mode of theta include the detection, the monitoring, and even the correction of the error, so we refer to it as "error-related theta". This error-related theta would then reflect the cognitive processes aiming to prevent an error from releasing completely.

Evidence that theta activity represents neural processing involved in error-correction has been previously shown, with greater theta power correlating to shorter correction times in partial error trials (Cohen & van Gaal, 2014). From our study, two pieces of evidence add to the

proposed error-correcting role of theta. Firstly, the stronger relationship between theta activity

from this mode and partial error onsets revealed a different temporal structure of theta activity

than in full-correct trials. Secondly, the timing of the error-related theta occurred a bit later in the

full error trials compared to the partial error trials. Theta activity over different time-scales has

been shown to interact with each other to help in online behavioral adaptation (Cohen, 2016). It

could be that the error-related theta events we observed in the data, and their short-timescale,

could be involved in the rapid correction of the partial error, and thus if they occur a bit later in

time, this leads to a full error.

Theta modes might have different underlying sources and relate to different cognitive

strategies

Results of our study suggest that the modes of theta we observe seem to represent processes

involved in conflict and error-processing. Previous studies have shown that both conflict and

error-related processing could be functionally dissociated both in terms of the areas they

originate from and the frequency bands in which they operate (Cohen & van Gaal, 2014; Nee,

Kastner, & Brown, 2011; Ullsperger & von Cramon, 2001). In our case, these modes of theta

also differ in terms of their strength and duration, with the error-related theta events being

stronger and more punctate, compared to the conflict-related theta. The dissociation with respect

to their functional role and properties could reflect that these two modes of theta are generated by

different underlying brain networks. This is supported by studies that have demonstrated the

presence of multiple sources of mid-frontal theta (Töllner et al., 2017; Zuure, Hinkley, Tiesinga,

Nagarajan, & Cohen, 2020), and related them to both conflict and other conflict-independent processes (Töllner et al., 2017).

If the two theta modes indeed reflect two different functions in the processing of conflicts and errors, it seems plausible that there is also some individual variation in how these two systems are engaged when dealing with conflict. For example, we noticed that for a minority of participants a single gaussian provided a better fit than two gaussians in the gaussian mixture modeling of the theta onsets in the partial error trials (Fig. 5a and 5b). Thus, it is possible that this was due to exerting a particular control strategy, such as initiating theta activity proactively to prevent errors from occuring, or relying more on the error-correcting mode of theta instead. Although mid-frontal theta is considered to be a general mechanism activated whenever there is a need for control (Cavanagh & Frank, 2014; Cavanagh & Shackman, 2015; Cohen & Donner, 2013), theta has been also implicated in different control strategies (Cooper et al., 2015; Eisma, Rawls, Long, Mach, & Lamm, 2021; Messel, Raud, Hoff, Stubberud, & Huster, 2021), such as proactive or reactive (Eisma et al., 2021). While the present data did not allow us to test these ideas, it points to potentially interesting future studies examining how different degrees and strategies of cognitive control affect theta signatures. For example, one prediction would be that a shift in the cognitive strategy is accompanied by a shift between the two modes of theta.

Models of theta function and conclusions

Our findings have implications for rise-to-threshold models incorporating theta activity, specifically the interpretation that theta activity affects decision/response thresholds or "hold your horses" forms of control (Cavanagh & Frank, 2014; Cavanagh et al., 2011; Zavala et al.,

2013). For example, if the function of both modes of theta involves increasing decision thresholds, important questions arise on how the error-correction would manifest mechanistically, and what the mechanistic differences are between the two modes. One possibility is that they differ in terms of their selectivity, so that e.g. the error-related theta raises thresholds selectively for the incorrect response. To the contrary, there is evidence that cortico-spinal excitability is decreased during conflict resolution (Duque, Olivier, & Rushworth, 2013; Wessel, Waller, & Greenlee, 2019), which is non-selective to the action effector, suggesting a global mechanism at play to resolve competition between responses. However, these studies have not looked at partial error trials or the possibility of theta having different modes. Therefore, it is possible that the conflict-related theta might reflect a "caution" signal, globally holding responses from execution, and the error-related theta represents processes more specific to the incorrect response. Future studies aiming to resolve this would further help in understanding theta's role in error-processing better.

Alternatively, there could be other mechanisms implementing error-correction apart from rise-to-threshold model interpretations. For instance, theta's interaction with downstream subcortical regions, specifically the basal ganglia, could be relevant for the different theta modes (Zavala et al., 2013, 2016, 2017, 2018) . Our results point to the importance of the timing of the error-related theta, as its onset was overall delayed in the full error trials compared to the partial error trials. This could mean that how cortical theta, specifically the error-related theta, interacts with subcortical regions (e.g. the subthalamic nucleus) might drive successful error correction. Finally, it is also possible that the error-related theta represents processes involved in monitoring the error and using that information to modify behavior on successive trials.

To conclude, we investigated the single-trial relationship between transient theta events and muscle-level metrics of response conflict. We showed that there is evidence for two modes of theta, one of which is strongly linked to the potential upcoming error. Future studies might further address the origin of the theta events and their interaction with neural circuits involved in error processing.

REFERENCES

Appelbaum, L. G., Smith, D. V., Boehler, C. N., Chen, W. D., & Woldorff, M. G. (2011). Rapid modulation of sensory processing induced by stimulus conflict. *Journal of Cognitive Neuroscience*, *23*(9), 2620–2628.

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57*(1), 289–300.

Blankertz, B., Tomioka, R., Lemm, S., Kawanabe, M., & Muller, K. (2008). Optimizing Spatial filters for Robust EEG Single-Trial Analysis. *IEEE signal processing magazine*, *25*(1), 41–56.

Cavanagh, J. F., & Frank, M. J. (2014). Frontal theta as a mechanism for cognitive control. *Trends in Cognitive Sciences*, *18*(8), 414–421.

Cavanagh, J. F., & Shackman, A. J. (2015). Frontal midline theta reflects anxiety and cognitive control: meta-analytic evidence. *Journal of physiology, Paris*, *109*(1–3), 3–15.

Cavanagh, J. F., Wiecki, T. V., Cohen, M. X., Figueroa, C. M., Samanta, J., Sherman, S. J., & Frank, M. J. (2011). Subthalamic nucleus stimulation reverses mediofrontal influence over

decision threshold. *Nature Neuroscience*, *14*(11), 1462–1467.

Cohen, M. X., & Cavanagh, J. F. (2011). Single-trial regression elucidates the role of prefrontal theta oscillations in response conflict. *Frontiers in Psychology*, *2*, 30.

Cohen, M. X., & Donner, T. H. (2013). Midfrontal conflict-related theta-band power reflects neural oscillations that predict behavior. *Journal of Neurophysiology*, *110*(12), 2752–2763.

Cohen, M. X., & van Gaal, S. (2014). Subthreshold muscle twitches dissociate oscillatory neural signatures of conflicts from errors. *Neuroimage*, *86*, 503–513.

Cohen, M. X., & Ridderinkhof, K. R. (2013). EEG source reconstruction reveals frontal-parietal dynamics of spatial conflict processing. *Plos One*, *8*(2), e57293.

Cohen, M. X. (2015). Comparison of different spatial transformations applied to EEG data: A case study of error processing. *International Journal of Psychophysiology*, *97*(3), 245–257.

Cohen, M. X. (2016). Midfrontal theta tracks action monitoring over multiple interactive time scales. *Neuroimage*, *141*, 262–272.

Cohen, M. X. (2017). Comparison of linear spatial filters for identifying oscillatory activity in multichannel data. *Journal of Neuroscience Methods*, *278*, 1–12.

Cohen, M. X. (2022). A tutorial on generalized eigendecomposition for denoising, contrast enhancement, and dimension reduction in multichannel electrophysiology. *Neuroimage*, *247*, 118809.

Cooper, P. S., Wong, A. S. W., Fulham, W. R., Thienel, R., Mansfield, E., Michie, P. T., & Karayanidis, F. (2015). Theta frontoparietal connectivity associated with proactive and reactive cognitive control processes. *Neuroimage*, *108*, 354–363.

Duque, J., Olivier, E., & Rushworth, M. (2013). Top-down inhibitory control exerted by the

medial frontal cortex during action selection under conflict. *Journal of Cognitive Neuroscience*, *25*(10), 1634–1648.

Egner, T. (2007). Congruency sequence effects and cognitive control. *Cognitive, Affective & Behavioral Neuroscience*, *7*(4), 380–390.

Eisma, J., Rawls, E., Long, S., Mach, R., & Lamm, C. (2021). Frontal midline theta differentiates separate cognitive control strategies while still generalizing the need for cognitive control. *Scientific Reports*, *11*(1), 14641.

Errington, S. P., Woodman, G. F., & Schall, J. D. (2020). Dissociation of Medial Frontal β-Bursts and Executive Control. *The Journal of Neuroscience*, *40*(48), 9272–9282.

Feingold, J., Gibson, D. J., DePasquale, B., & Graybiel, A. M. (2015). Bursts of beta oscillation differentiate postperformance activity in the striatum and motor cortex of monkeys performing movement tasks. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(44), 13687–13692.

Gratton, G., Coles, M. G., Sirevaag, E. J., Eriksen, C. W., & Donchin, E. (1988). Pre- and poststimulus activation of response channels: a psychophysiological analysis. *Journal of Experimental Psychology. Human Perception and Performance*, *14*(3), 331–344.

Jana, S., Hannah, R., Muralidharan, V., & Aron, A. R. (2020). Temporal cascade of frontal, motor and muscle processes underlying human action-stopping. *eLife*, *9*.

Little, S., Bonaiuto, J., Barnes, G., & Bestmann, S. (2019). Human motor cortical beta bursts relate to movement planning and response errors. *PLoS Biology*, *17*(10), e3000479.

Messel, M. S., Raud, L., Hoff, P. K., Stubberud, J., & Huster, R. J. (2021). Frontal-midline theta reflects different mechanisms associated with proactive and reactive control of inhibition.

Neuroimage, *241*, 118400.

Muralidharan, V., Aron, A. R., & Schmidt, R. (2021). Transient beta modulates decision

thresholds during human action-stopping. *BioRxiv*.

Muralidharan, V., & Aron, A. R. (2021). Behavioral induction of a high beta state in

sensorimotor cortex leads to movement slowing. *Journal of Cognitive Neuroscience*, *33*(7),

1311–1328.

Nee, D. E., Kastner, S., & Brown, J. W. (2011). Functional heterogeneity of conflict, error,

task-switching, and unexpectedness effects within medial prefrontal cortex. *Neuroimage*,

54(1), 528–540.

Nigbur, R., Cohen, M. X., Ridderinkhof, K. R., & Stürmer, B. (2012). Theta dynamics reveal

domain-specific control over stimulus and response conflict. *Journal of Cognitive

Neuroscience*, *24*(5), 1264–1274.

Parra, L., & Sajda, P. (2003). Blind source separation via generalized eigenvalue decomposition.

The Journal of Machine Learning Research.

Rochet, N., Spieser, L., Casini, L., Hasbroucq, T., & Burle, B. (2014). Detecting and correcting

partial errors: Evidence for efficient control without conscious access. *Cognitive, Affective &

Behavioral Neuroscience*, *14*(3), 970–982.

Soh, C., Hynd, M., Rangel, B. O., & Wessel, J. R. (2021). Adjustments to Proactive Motor

Inhibition without Effector-Specific Foreknowledge Are Reflected in a Bilateral

Upregulation of Sensorimotor β-Burst Rates. *Journal of Cognitive Neuroscience*, *33*(5),

784–798.

Swick, D., & Turken, A. U. (2002). Dissociation between conflict detection and error monitoring

in the human anterior cingulate cortex. *Proceedings of the National Academy of Sciences of the United States of America*, *99*(25), 16354–16359.

Töllner, T., Wang, Y., Makeig, S., Müller, H. J., Jung, T.-P., & Gramann, K. (2017). Two Independent Frontal Midline Theta Oscillations during Conflict Detection and Adaptation in a Simon-Type Manual Reaching Task. *The Journal of Neuroscience*, *37*(9), 2504–2515.

Torrecillos, F., Tinkhauser, G., Fischer, P., Green, A. L., Aziz, T. Z., Foltynie, T., Limousin, P., et al. (2018). Modulation of beta bursts in the subthalamic nucleus predicts motor performance. *The Journal of Neuroscience*, *38*(41), 8905–8917.

Ullsperger, M., & von Cramon, D. Y. (2001). Subprocesses of performance monitoring: a dissociation of error processing and response competition revealed by event-related fMRI and ERPs. *Neuroimage*, *14*(6), 1387–1401.

Vidal, F., Burle, B., & Hasbroucq, T. (2019). Errors and action monitoring: errare humanum est sed corrigere possibile. *Frontiers in Human Neuroscience*, *13*, 453.

Wessel, J. R., Waller, D. A., & Greenlee, J. D. (2019). Non-selective inhibition of inappropriate motor-tendencies during response-conflict by a fronto-subthalamic mechanism. *eLife*, *8*.

Wessel, J. R. (2020). β-Bursts Reveal the Trial-to-Trial Dynamics of Movement Initiation and Cancellation. *The Journal of Neuroscience*, *40*(2), 411–423.

Zavala, B., Brittain, J.-S., Jenkinson, N., Ashkan, K., Foltynie, T., Limousin, P., Zrinzo, L., et al. (2013). Subthalamic nucleus local field potential activity during the Eriksen flanker task reveals a novel role for theta phase during conflict monitoring. *The Journal of Neuroscience*, *33*(37), 14758–14766.

Zavala, B., Damera, S., Dong, J. W., Lungu, C., Brown, P., & Zaghloul, K. A. (2017). Human

subthalamic nucleus theta and beta oscillations entrain neuronal firing during sensorimotor conflict. *Cerebral Cortex*, *27*(1), 496–508.

Zavala, B., Jang, A., Trotta, M., Lungu, C. I., Brown, P., & Zaghloul, K. A. (2018). Cognitive control involves theta power within trials and beta power across trials in the prefrontal-subthalamic network. *Brain: A Journal of Neurology*, *141*(12), 3361–3376.

Zavala, B., Tan, H., Ashkan, K., Foltynie, T., Limousin, P., Zrinzo, L., Zaghloul, K., et al. (2016). Human subthalamic nucleus-medial frontal cortex theta phase coherence is involved in conflict and error related cortical monitoring. *Neuroimage*, *137*, 178–187.

Zuure, M. B., Hinkley, L. B., Tiesinga, P. H. E., Nagarajan, S. S., & Cohen, M. X. (2020). Multiple midfrontal thetas revealed by source separation of simultaneous MEG and EEG. *The Journal of Neuroscience*, *40*(40), 7702–7713.

SUPPLEMENTARY INFORMATION

Relationship between partial error onsets and theta onsets: Dependence on selection parameters

We tested whether the relationship seen between partial error/correct response onset and theta onset times from G1 and G2 (as seen in Fig. 5c and 5d) depended on the specific participant inclusion criteria. It was the case that in certain participants there were not sufficient trials in either G1 or G2 as a result of the fitting procedure and thus we analyzed those subjects who had more than 15 data points (i.e. theta onsets) in both G1 and G2 for the partial error and the full-correct matched trials. We repeated the same analysis with subjects who had more than 3 data points for both G1 and G2. This resulted in including 17/24 and 14/15 in both Flanker and Simon datasets. We found that the results did not change for the Flanker dataset (Fig. S1). For the Simon dataset, the comparison of correlations between partial error onsets vs G2 theta onsets in the partial error trials with surrogate partial error onsets vs G2 theta onsets in the full-correct matched trials did not reach statistical significance, while the other results remained the same (see Fig. S1). Overall, this supports our results that one of the modes of theta (i.e. from G2) is related to the error.

Figure S1. **Dependence of correlations on inclusion parameters**. **c)** and **d)** Correlational analysis between the theta onsets belonging to either G1 or G2 and the EMG timings (partial error – PEMG and correct response – CEMG) for the Flanker and Simon task respectively using different participant inclusion criterion. Analysis reveals similar results as in the main manuscript, Fig. 5c and 5d.