Main Manuscript

Human neutrophils direct epithelial cell extrusion to enhance intestinal epithelial host

defense during *Salmonella* infection

Authors/affiliations

Anna-Lisa E. Lawrence[a], Ryan P. Berger[a], David R. Hill[b], Sha Huang[c], Veda K.

Yadagiri[b], Brooke Bons[b], Courtney Fields[b], Gautam J. Sule[b], Jason S. Knight[b],

Christiane E. Wobus[a], Jason R. Spence[c], Vincent B. Young[b], Mary X. O'Riordan[a1,3], and

Basel H. Abuaita[a1,2]

[a] Department of Microbiology and Immunology, University of Michigan Medical School,

Ann Arbor, MI, USA

[b] Department of Internal Medicine, University of Michigan Medical School, Ann Arbor,

MI, USA

[c] Department of Cell and Developmental Biology, University of Michigan Medical

School, Ann Arbor, MI, USA

Author list footnotes

[1] M.X.O and B.H.A contributed equally to the work

[2] B.H.A Present address: Department of Pathobiological Sciences, Louisiana State

University School of Veterinary Medicine, Baton Rouge, LA, USA

[3] Corresponding authors Email: **babuaita@lsu.edu**, **oriordan@umich.edu**

Author contributions

A-L.E.L and B.H.A performed the experiments and analyzed the data. A-L.E.L, M.X.O

and B.H.A designed the experiments and wrote the manuscript. R.P.B and D.R.H

assisted in RNA-seq analysis. S.H, V.K.Y, B.B and C.F assisted in HIO differentiation

and culturing. G.J.S, J.R.S, V.B.Y, C.E.W and J.S.K assisted in experimental

preparation and data interpretation.

Declaration of interests

The authors declare no competing interests.

Classification

Biological Sciences: Immunology and Inflammation

Keywords

Intestinal host defense, Cell extrusion, inflammasome

Abstract

Infection of the human gut by *Salmonella enterica* Typhimurium (STM) results in a localized inflammatory disease that is not mimicked in murine infections. To determine mechanisms by which neutrophils, as early responders to bacterial challenge, direct inflammatory programming of human intestinal epithelium, we established a multi-component human intestinal organoid (HIO) model of STM infection. HIOs were micro-injected with STM and then seeded with primary human polymorphonuclear leukocytes (PMN-HIOs), specifically neutrophils and analyzed for bacterial growth and host cell survival. Surprisingly, PMNs did not affect luminal colonization of *Salmonella,* but their presence reduced intraepithelial bacterial burden. Adding PMNs to infected HIOs resulted in substantial accumulation of shed intestinal epithelial cells that could be blocked by Caspase-1 or Caspase-3 inhibition. Cleaved Caspase-3 was present in epithelial cells, but expression of the inflammasome adaptor, ASC, was only detected in PMNs. Caspase inhibition also increased bacterial burden in the epithelium of the PMN-HIO, suggesting PMNs enhance activation of cell death pathways in human intestinal epithelial cells as a protective response to infection. These data support a critical function for neutrophils beyond their antimicrobial role whereby they amplify cell death and extrusion of epithelial cells from the *Salmonella*-infected intestinal monolayer.

Significance statement

Neutrophils are early responders to *Salmonella* intestinal infection, but how they influence infection progression and outcome is unknown. Here we use a co-culture model of human intestinal organoids and human primary neutrophils to study the

contribution of human neutrophils to *Salmonella* infection of the intestinal epithelium.

We found that neutrophils markedly enhanced epithelial defenses, including enhancing

cell extrusion to reduce intraepithelial burden of *Salmonella* and association with the

epithelium, rather than directly killing *Salmonella* in the HIO lumen. These findings

reveal a novel role for neutrophils in the gut beyond killing invading pathogens and

illuminate how neutrophils can reprogram cells in the gut environment to enhance

antimicrobial defenses.

Main text

Introduction

Salmonella enterica is one of the most common causes of foodborne disease,

responsible for an estimated 1.35 million infections in the United States each year (1).

S. enterica serovar Typhimurium (STM), one of the most prevalent *S. enterica* serovars,

infects via the fecal-oral route and stimulates robust inflammation in the host intestinal

milieu, leading to gastroenteritis and diarrheal disease (2). *Salmonella* pathogenesis is

commonly studied *in vivo* using mouse infection models and now more recently *in vitro*

using human derived organoid and enteroid models (3–7). Through these studies a

critical role for epithelial cell death and extrusion of *S. enterica* infected cells has been

shown to regulate infection outcome by reducing epithelial bacterial burden and

restricting the infection to the intestine (5, 8–11).

Epithelium intrinsic induction of cell death and extrusion pathways are essential in

maintenance of normal intestinal homeostasis during health and infection, however it is

also known that innate immune cells play a dominant role in determining infection

outcome by bacterial infections, including *Salmonella* (12, 13). One of the earliest

responders and the most abundant cell types found in *Salmonella*-infected individuals

are polymorphonuclear leukocytes (PMNs), specifically neutrophils (14, 15). PMNs

defend against bacterial infections through multiple mechanisms: antimicrobial effectors

like degradative proteases and ion chelators, production of reactive oxygen species and

formation of sticky antimicrobial neutrophil extracellular traps (NETs) (16). And while

PMNs are very effective at killing extracellular bacteria (17), the role of PMNs in a more

complex infection where a large proportion of bacteria reside within host cells, like the

intestinal epithelium, is unknown.

More recently, it is becoming appreciated that PMNs serve other functions beyond

killing extracellular pathogens, e.g., changing the microenvironment via molecular

oxygen depletion, regulating nutrient availability, and through production of inflammatory

mediators (18, 19). Notably, Gopinath et al. found that neutrophilia induced a super-

shedder phenotype in a mouse infection model of *Salmonella* (20), but how the

interaction between epithelial cells and PMNs affects the outcome of bacterial infections

is still poorly understood. To address this gap in knowledge, we generated a co-culture

model of primary human PMNs, specifically neutrophils, with human intestinal organoids

(HIOs) termed PMN-HIOs to study the contribution of PMNs during infection with

Salmonella enterica serovar Typhimurium (STM).

Using this PMN-HIO model, we evaluated how PMNs modulate intestinal epithelial host

defenses during infection, compared to infected HIOs alone. We show here that the

presence of PMNs elevates the overall inflammatory tone of the epithelium and

markedly promotes cell death and extrusion of epithelial cells, thereby reducing

Salmonella intraepithelial burden.

Results

Human PMNs transmigrate into the HIO lumen during infection and reduce

***Salmonella* intraepithelial burden.**

PMNs are known to transmigrate across intestinal epithelial layers during early stages

of inflammation (21, 22), therefore we asked whether PMNs would transmigrate into the

HIO lumen during infection. 10^5 *Salmonella enterica* Typhimurium (STM) were

microinjected into the HIO lumen and cultured with PMNs isolated from healthy human

volunteers for 8h (PMN-HIOs). To quantify PMN recruitment to infected HIOs, PMNs

were pre-labeled with Carboxyfluorescein succinimidyl ester (CFSE) prior to co-culture

with HIOs. PMN-HIOs were collected at 8h post-infection (hpi), washed to remove

unassociated neutrophils, dissociated into a single cell suspension and the percentage

of CFSE-positive cells was enumerated by flow cytometry. There was a significant

increase in the number of PMNs associated with infected HIOs compared to PBS

controls, with approximately 5% of total cells present in PMN-HIOs staining positive for

CFSE (Fig. 1A). Immunofluorescent staining for neutrophil-specific Myeloperoxidase

(MPO) was performed on paraffin sections to further monitor localization of PMNs within

PMN-HIOs. In contrast to PBS-injected controls, MPO-positive cells were observed in

the lumen of STM-infected HIOs, confirming that PMNs transmigrate into the HIO lumen

during infection (Fig. 1B). Since PMNs are potent killers of bacterial pathogens, we

tested whether PMNs controlled *Salmonella* colonization within the HIO. Although

PMNs killed STM in pure PMN cultures, with ~30% of STM killed by 4hpi (SI Appendix,

Fig. S1), PMNs did not alter the total levels of STM in the HIOs (Fig. 1C). This was not

due to lack of PMN activation in the HIOs as we detected formation of NETs in the

lumen of STM-infected PMN-HIOs (SI Appendix, Fig. S2). In addition, culture

supernatants were analyzed for production of antimicrobial effectors via ELISA (SI

Appendix, Fig. S3). Some antimicrobial effectors such as Elafin (PI3), a small cationic

peptide secreted at mucosal surfaces (23), and Calprotectin (S100A8 and S100A9)

were produced at higher levels in PMN-HIOs, compared to HIOs alone, confirming that

there was no defect in the antimicrobial response to STM and in contrast suggests that

PMNs augment epithelial host defenses. However, since *Salmonella* has evolved

mechanisms to overcome Calprotectin-mediated immunity and thrive under these

conditions; upregulation of these specific antimicrobial effectors is likely insufficient to

reduce *Salmonella* colonization in the PMN-HIOs (24, 25).

During intestinal infection, *Salmonella* reside in both the lumen of the intestine and

within epithelial cells, and it has been suggested that the intracellular pool of bacteria

are important for reseeding the gut lumen to prolong infection and promote fecal

shedding (8, 26). To assess what impact PMNs have on the intracellular bacterial

burden, paraffin sections of *Salmonella*-infected HIOs and PMN-HIOs were stained to

detect both epithelial cells and *Salmonella* and the epithelial bacterial burden was

quantified by fluorescence microscopy (Fig. 1D, 1E). This analysis revealed that there

were significantly fewer intracellular bacteria in the epithelial lining of PMN-HIOs

compared to HIOs alone indicating that PMNs aid in reducing epithelial cell bacterial

burden. Interestingly, we also observed a reduction in epithelial-surface associated

bacteria suggesting that PMNs also reduce STM attachment to further protect the

epithelial lining. Together, these results show that although PMNs transmigrate into the

HIO lumen during *Salmonella* infection, they do not directly kill *Salmonella*, but instead

enhance epithelial defenses to reduce the bacterial burden within the epithelial layer.

PMNs enhance shedding of epithelial cells during *Salmonella* infection.

In addition to measuring a significant reduction in intraepithelial bacterial burden and

association of STM with epithelial cells, we observed robust accumulation of DAPI-

positive cells in the lumen of STM-infected PMN-HIOs that were negative for the PMN

marker MPO (Fig. 1B, F). Shedding of *Salmonella*-infected cells from the gut via

programmed cell death pathways is an important defense mechanism used to protect

the host from invasive *Salmonellosis* and helps reduce intestinal bacterial burden to

resolve the infection and so we hypothesized that in a human infection model PMNs

may enhance this process (5, 8–11). To determine whether these luminal cells were

dead epithelial cells shed from the HIO epithelial lining, we performed Terminal

deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) on HIOs and PMN-HIOs

microinjected with either STM or PBS control. The presence of PMNs induced robust

accumulation of TUNEL-positive cells in the lumen of infected HIOs (Fig. 2A, 2B). While

we also detected a substantial number of TUNEL-positive cells in the mesenchyme this

phenotype was present in all conditions so likely was not caused by either *Salmonella*

or PMNs. Accumulation of luminal TUNEL-positive cells was selectively induced by

PMNs during infection, as neither infected HIOs or uninfected PMN-HIOs showed this

phenotype. To confirm that these cells were epithelial cells, we stained for the epithelial

marker E-cadherin, and found that the vast majority of TUNEL-positive cells in PMN-

HIOs were epithelial cells (Fig. 2C, 2D). Previously we reported that STM infection in

HIOs induces significant induction of TUNEL-positive cells that are retained in the

epithelial lining (4), consistent with these findings, our results suggest that PMNs are

enhancing shedding of these TUNEL-positive cells from the monolayer as we observe

very few TUNEL-positive cells in the epithelial lining of infected PMN-HIOs (Fig. 2D).

TUNEL staining is classically associated with apoptotic pathways, and since activated

PMNs have been shown to induce apoptotic processes in lung epithelium (27), we

hypothesized that PMNs induce epithelial cell apoptosis to reduce bacteria associated

with the epithelial lining. To test our hypothesis, we stained STM-infected PMN-HIO

sections for cleaved Caspase-3 as a marker of apoptosis. We found that many, but not

all luminal epithelial cells were positive for cleaved Caspase-3 (Fig. 2E) suggesting that

multiple forms of cell death were occurring in the PMN-HIO likely including Caspase-4/5

mediated shedding of *Salmonella*-infected epithelial cells as has previously been

established in a human enteroid infection model (5, 6) and more recently shown in a

Caco-2 model of *Salmonella* infection (28). Interestingly, PMN-induced cell shedding

was not restricted to infected cells, as we observed both infected and uninfected cells in

the PMN-HIO lumen (SI Appendix, Fig. S4). Together these results suggest that PMNs

promote programmed cell death pathways and epithelial cell shedding during

Salmonella infection.

Inflammasome activation and IL-1 production is mediated by PMNs during

infection.

PMNs are known to affect epithelial cell function through multiple mechanisms including

through NET formation. Inflammasome activation is known to occur in PMNs during

bacterial infections (29) and it is known that inflammasomes also play a key role in

regulating intestinal inflammation (30). While activation of noncanonical inflammasomes

requiring Caspase-4/5 in human epithelial cells is a hallmark of *Salmonella* infection,

activation of Caspase-1 dependent inflammasomes in innate immune cells is also

critical for effective responses against invading pathogens (31). To assess how PMNs

shape inflammasome activation in the PMN-HIO model, we first examined gene level

expression data from RNA-seq data of HIOs and PMN-HIOs microinjected with PBS or

STM (SI Appendix, Table S1). We found that PMNs significantly contributed to

upregulation of several genes in inflammasome/cell death signaling during STM

infection (Fig. 3A). While there was weak upregulation of IL-1β and IL-1α in STM-

infected HIOs alone, we did not observe significant changes in expression of other

mediators or machinery required for NLRP3 inflammasome assembly such as CASP1,

NLRP3, or PYCARD (encoding ASC which was not differentially expressed under any

condition) (Fig. 3A). However, consistent with previous reports studying cell death

responses to STM-infection in human epithelial models, we did measure a significant

increase in expression of Caspase-4 and Caspase-5. In contrast, when PMNs were

added to infected HIOs, we observed stronger upregulation of *IL-1* genes and effectors

involved in inflammasome activation including the upregulation of *NLRP3* and *Caspase-*

1 (*CASP1*). To further characterize this phenotype, we collected culture supernatants

from HIOs and PMN-HIO and quantified levels of IL-1 family cytokines during infection

(Fig. 3B). IL-1β or IL-1α was undetectable in infected HIOs; release of these cytokines

required the presence of PMNs as IL-1β or IL-1α levels significantly increased in STM-

infected PMN-HIOs. These results suggest that PMNs significantly contribute to

production of IL-1 cytokines in this infection model. This is consistent with previous

reports that the human epithelium is not a dominant source of IL-1 cytokines (32) and

that Caspase-4/5 activation is not classically associated with significant IL-1β

processing (33). Interestingly, we also observed production of IL-1RA, the antagonist of

the IL-1 receptor which is usually co-expressed with IL-1α/β (34), in infected PMN-HIOs

revealing an additional role for PMNs in inducing signaling processes that tune the

magnitude of immune activation. In contrast, IL-33, another important IL-1 family

alarmin in mucosal immunity (35), was produced in all conditions independent of the

presence of PMNs. IL-33 specifically and in contrast to other IL-1 family cytokines, is

released constitutively by epithelial cells where it is then processed extracellularly by

serine proteases including elastase, which is released by PMNs (36). This processed

form is thought to enhance inflammatory signaling. Notably, we observed significantly

lower levels of IL-33 in STM-infected PMN-HIOs, which may be caused by PMN

processing. To further define which cells within the PMN-HIOs contribute to

inflammasome activation and IL-1 processing, paraffin sections of STM-infected PMN-

HIOs were stained for ASC, an adaptor protein required for inflammasome assembly

(Fig. 3C, 3D) (37). ASC-positive signal was not observed in HIO epithelial cells,

consistent with a recent report that *Salmonella*-induced epithelial cell death occurs

independently of ASC (28), but instead was associated with cells positive for vimentin, a

protein expressed by PMNs and mesenchymal cells within the PMN-HIOs. ASC and

Vimentin double-positive cells were primarily located within the lumen of PMN-HIOs,

suggesting that these cells are PMNs. Closer examination of nuclear morphology of the

ASC-positive cells by DAPI staining revealed multi-lobed nuclei, further supporting that

inflammasome activation and IL-1 processing occur in PMNs. Together, these findings

are consistent with a model where PMNs are the primary site of Caspase-1 dependent

inflammasome activation and the production of IL-1 family cytokines during infection in

the PMN-HIO model.

Caspase-1 and Caspase-3 inhibition reduces shedding of epithelial cells and

increases the association of *Salmonella* with the epithelium.

Epithelial cell death and shedding serve to reduce bacterial burden in the intestinal

epithelium (5, 6, 8–11). To define the functional consequences of PMN-induced

epithelial cell death on host defense and determine which caspases were involved, we

treated PMN-HIOs with various caspase inhibitors. Accumulation of TUNEL-positive

epithelial cells was monitored in infected PMN-HIOs in the presence or absence of

Caspase-1 (z-YVAD-FMK) or Caspase-3 (z-DEVD-FMK) inhibitors. We performed

TUNEL staining on paraffin sections from infected PMN-HIOs at 8hpi (Fig. 4A, 4B). Both

Caspase-1 and Caspase-3 inhibition significantly reduced accumulation of TUNEL-

positive cells in the lumen of infected PMN-HIOs, indicating that PMN-dependent

Caspase-1 and -3 activation is required for efficient shedding of epithelial cells. To test

how caspase inhibition and therefore reduced shedding affected STM infection, PMN-

HIOs were stained with an anti-*Salmonella* antibody to characterize the localization of

bacteria in the HIO by quantifying the percentage of infected cells, number of bacteria

per cell, and epithelium associated bacteria (Fig. 4C-E, SI Appendix, Fig. S5).

Consistent with our hypothesis that PMNs enhance shedding of epithelial cells through

caspase activation, there were greater numbers of bacteria per cell in Caspase-1

inhibitor-treated PMN-HIOs, but surprisingly not in Caspase-3 inhibitor-treated PMN-

HIOs (Fig. 4D). Caspase-4/5 activation in human epithelial cells is known to be

important in shedding of *Salmonella*-infected cells and suggests that PMNs may

enhance epithelial Caspase-4/5 signaling via Caspase-1 activity to enhance cell

shedding instead of utilizing Caspase-3 for this process. We also observed a trending

increase in the percentage of infected cells with Caspase-1 inhibition (SI Appendix, Fig.

S5). In contrast, when PMN-HIOs were treated with the Caspase-3 inhibitor, there was

a significant increase in epithelium associated bacteria (Fig. 4D). There is some

evidence that activated PMNs induce apoptosis of intestinal epithelial cells (27) and so

NET formation during *Salmonella* infection in PMN-HIOs may result in enhanced

epithelial cell shedding independent of PMN Caspase-1 to increase the rate of cell

turnover and therefore reduce the association of bacteria with the apical surface of the

epithelium which would protect the epithelium from future bacterial invasion. These data

suggest that both Caspase-1 and Caspase-3 inhibition reduce accumulation of dead

cells in the PMN-HIO lumen, but Caspase-1 activity is important for directly regulating

epithelial bacterial burden while Caspase-3 reduces association of bacteria with the

epithelium to protect the epithelium from the next round of infection. Taken together, our

data support a model where Caspase-1 dependent inflammasome activation in PMNs

enhances epithelial cell shedding via two independent pathways to control *Salmonella*

infection.

Discussion

Neutrophils (PMNs) dominate the early response to *Salmonella* infection in the gut (15,

38–40), but their functions in regulating intestinal epithelial cell host defense and

infection outcome are not well understood. Here we used a co-culture model of human

intestinal organoids (HIOs) with primary human PMNs, termed PMN-HIOs, to elucidate

these roles. We found that while there was no difference in luminal colonization of

Salmonella in the HIOs, PMNs did reduce intraepithelial bacterial burden in the

epithelium and reduced the association of *Salmonella* with the apical surface of the

epithelial monolayer. PMNs were associated with elevated epithelial cell death and

promoted extrusion of these cells into the lumen of infected PMN-HIOs. We found that

Caspase-1 activation was required for epithelial shedding, noting that the ASC

inflammasome adaptor was only present in PMNs, and that inhibition of Caspase-1

activity increased bacterial burden in HIO epithelial cells. Independently, we uncovered

an important role for Caspase-3 activation in epithelial cells in reducing association of

Salmonella with the apical surface, which was enhanced by the presence of neutrophils.

Thus, we propose a model where PMNs enhance shedding of epithelial cells from the

intestinal barrier via two distinct mechanisms to reduce intracellular bacterial loads and

adherence of *Salmonella* to the epithelium, potentially tilting infection outcome favorably

toward the host.

Our observation that PMNs did not affect total bacterial colonization in the HIO lumen

was rather unexpected, both because antimicrobial effectors were produced in the

PMN-HIO cultures and our findings, and that of others, indicate that PMNs kill

Salmonella in the absence of HIOs (17). We also observed transmigration and NET

formation by PMNs in the HIO lumen, so the lack of *Salmonella* killing in the HIO lumen

could not be explained by lack of PMNs at the site of infection. These observations

suggest the possibility that *Salmonella* may employ specific mechanisms to overcome

PMN effector functions in the HIO luminal environment. Alternatively, while PMNs can

be potent killers of invading bacteria, mechanisms used by PMNs to kill pathogens are

not selective and therefore PMN activation near the epithelial barrier may be tightly

regulated to avoid tissue damage (41). Consistent with this idea, we observed robust

production of Elafin (Fig. S3), which is annotated as an antimicrobial peptide in

Reactome (42). Neutrophil elastase strongly contributes to the antimicrobial function of

PMNs for both intracellular killing via phagocytosis and decorating NETs (17, 43, 44).

Elafin can also inhibit neutrophil elastase to reduce tissue damage caused by neutrophil

overactivation (45, 46). Thus, expression of host Elafin in the presence of PMNs may

trade off the ability to kill luminal *Salmonella* with protection of epithelial barrier integrity.

Addition of PMNs to the infected HIOs resulted in decreased intracellular bacterial

burden, which was unexpected since intracellular bacteria are often considered

protected from neutrophil killing. We therefore hypothesized that this result was due to

extrusion of infected cells from the epithelial monolayer, a process mediated by caspase

activation. Previous reports have highlighted the importance of epithelial cell extrusion

in preventing dissemination of *Salmonella* beyond the intestine and that there are

epithelial cell-intrinsic mechanisms to rid the epithelial lining of *Salmonella* (5, 6, 9–11,

47). Of note, the role of the inflammatory caspases in these processes likely differs

between human and mouse, with caspase-1 playing a prominent role in mouse

epithelium, while Caspase-4 plays a more prominent role in human epithelium in the

absence of neutrophils. Although we could detect low levels of shed epithelial cells in

STM-infected HIOs alone consistent with prior studies, this phenotype was markedly

enhanced in the presence of PMNs, and inhibition with a selective Caspase-1 peptide

inhibitor decreased TUNEL+ cells, as well as number of bacteria per epithelial cell.

These data suggest a previously unappreciated role for PMNs in enhancing cell death in

Salmonella-infected epithelial cells and implicate Caspase-1 in that process.

In STM-infected HIOs, we observed close association of clusters of bacteria with the

epithelial surface, which would presumably advantage bacterial pathogens by spatial

proximity to the monolayer. Strikingly, the introduction of PMNs into the infected HIO

led to dispersal of the bacterial clusters into the HIO lumen, concomitant with a

substantial increase in TUNEL+ cells. There are numerous mechanisms by which

PMNs can drive epithelial cell death including oxidant production, which can activate

apoptotic pathways in the epithelium (48, 49). NET formation, which we observed in our

infected PMN-HIOs, also can induce epithelial cell death (27). This process was

reported to be largely dependent on extracellular histones released during NET

formation. We therefore consider that *Salmonella*-induced NET formation in PMN-HIOs

may contribute to overall shedding of epithelial cells and the dispersal of epithelial-

associated bacteria. These possibilities are consistent with our findings that both

infected and uninfected epithelial cells are shed from the intestinal monolayer in the

infected PMN-HIOs, but not the infected HIOs alone. By increasing epithelial cell

shedding, this neutrophil-dependent process contributes to reducing bacterial burden

within and associated with the epithelium.

Our findings pointed to Caspase-1 activity as a driver of accumulation of shed epithelial

cells in the HIO lumen, but markers of inflammasome activation were only observed

when PMNs were present. Although the importance of Caspase-1 in PMN activation

and antimicrobial functions are not fully understood, some evidence implicates

Caspase-1 in NET formation (29, 50). Our prior studies demonstrated that

inflammasome activation in PMNs occurs during bacterial infection (29) and another

study showed that inflammasome-mediated processing of Gasdermin D was required

for NET formation in mice (50). While there may be additional signaling roles of

Caspase-1 in PMN activation, we propose a model whereby PMNs transmigrate into the

inflamed intestine, undergo Caspase-1 mediated NET formation to trigger epithelial cell

death and shedding of infected cells to protect the epithelium from ongoing infection.

We reason that in the environment of the infected gut, in contrast to infected tissue, the

ratio of commensal and pathogenic bacteria to neutrophils may preclude substantive

bacterial killing through direct anti-microbial mechanisms.   Therefore, signaling

mechanisms whereby neutrophils are able to direct protective epithelial responses may

more be more advantageous to the host.

Materials and Methods

Contact for Reagent and Resource Sharing

Reagents and resources can be obtained by directing requests to the corresponding

authors, Basel Abuaita (babuaita@lsu.edu) and Mary O'Riordan (oriordan@umich.edu).

Human Intestinal Organoids (HIOs)

HIOs were generated by the *In Vivo* Animal and Human Studies Core at the University

of Michigan Center for Gastrointestinal Research as previously described (51). Prior to

experiments, HIOs were removed from the Matrigel, washed with DMEM:F12 media,

and re-plated with 5 HIOs/well in 50µl of Matrigel (Corning) in ENR media ((DMEM:F12,

1X B27 supplement, 2mM L-glutamine, 100ng/ml EGF, 100ng/ml Noggin, 500ng/ml

Rspondin1, and 15mM HEPES). Media was exchanged every 2-3 days for 7 days.

Human Polymorphonuclear Leukocytes (PMNs)

PMNs were isolated from blood of healthy human volunteers as previously described

(29). The purity of PMNs was assessed by flow cytometry using APC anti-CD16 and

FITC anti-CD15 antibodies (Miltenyi Biotec); markers characteristic of human

neutrophils. PMNs were labeled with cell trace CFSE dye (Thermo Fisher). PMNs were

incubated at room temperature for 20 minutes in PBS containing 5 µM CFSE. Cells

were washed twice with PBS to remove excess dye and collected by centrifugation.

CFSE-labeled PMNs were then co-cultured with STM-infected HIOs or PBS control to

monitor the association of PMNs with intestinal epithelial cells. PMN-HIOs were washed

twice with PBS to remove unassociated PMNs, mechanically dissociated into single-cell

suspension using a 70 µm cell strainer and analyzed on FACSCanto flow cytometer.

Percent of CFSE-positive cells were determined using FlowJo software.

Bacterial Growth and HIO Microinjection

Salmonella enterica serovar Typhimurium SL1344 (STM) was used throughout the

manuscript. Bacteria were stored at -80°C in Luria-Bertani (LB, Fisher) medium

containing 20% glycerol and cultured on LB agar plates. Individual colonies were grown

overnight at 37ºC under static conditions in LB liquid broth. Bacteria were pelleted,

washed and re-suspended in PBS. Bacterial inoculum was estimated based on OD_{600}

and verified by plating serial dilutions on agar plates to determine colony forming units

(CFU). The lumen of individual HIOs were microinjected with glass caliber needles with

1µl of PBS or STM (10^5 CFU/HIO) as previously described (3, 4, 52). HIOs were then

washed with PBS and incubated for 2h at 37°C in ENR media. HIOs were treated with

100µg/ml gentamicin for 15 min to kill any bacteria outside the HIOs, then incubated in

fresh medium +/- PMNs (5 X 10^5 PMNs/5HIOs/well in a 24-well plate). Where indicated,

PMNs-HIOs were treated with the following inhibitors after microinjection: 4 µM of

Caspase-1 inhibitor, Z-YVAD-FMK or 4 µM Caspase-3 inhibitor, Z-DEVD-FMK.

Bacterial Burden and Cytokine Analyses

Bacterial burden was assessed per HIO. Individual HIOs were removed from Matrigel,

washed with PBS and homogenized in PBS. Total CFU/HIO were enumerated by serial

dilution and plating on LB agar. For cytokine analysis, media from each well containing

5 HIOs/well were collected at 8hpi. Cytokines, chemokines and antimicrobial proteins

were quantified by ELISA at the University of Michigan Cancer Center Immunology

Core.

Immunofluorescence Staining and Microscopy

HIOs were fixed with 10% neutral formalin for 2 days and embedded in paraffin.

Histology sections (5µm) were collected by the University of Michigan Cancer Center

Histology Core. Sections were deparaffinized and antigen retrieval was performed in

sodium citrate buffer (10mM sodium citrate, 0.05% Tween 20, pH 6.0). Sections were

permeabilized with PBS+ 0.2% Triton X-100 for 30 min, then incubated in blocking

buffer (PBS, 5% BSA, and 10% normal goat serum) for 1h. Primary antibodies; anti-E-

Cadherin (BD Biosciences, clone 36), anti-MPO (Agilent, clone A0398), anti-Vimentin

(DSHB, Cat# AMF-17b), anti-ASC (Cell Signaling, Cat#13833) and anti-cleaved

Caspase-3 (Cell Signaling, Cat# 9661) were added to the histology sections in blocking

buffer overnight at 4°C. Goat anti-mouse and anti-rabbit secondary antibodies

conjugated to Alexa-488, Alexa-594 or Alexa-647 were used according to

manufacturer's instructions (Thermo Fisher) for 1h RT in blocking buffer. DAPI (Thermo

Fisher) was used to stain DNA. Bacteria were stained using anti-*Salmonella*

Typhimurium FITC-conjugated antibody (Santa Cruz, Cat# sc-52223). Sections were

mounted using coverslips (#1.5) and Prolong Diamond or Prolong Glass Antifade

Mountant (Thermo Fisher). Images were taken on Olympus BX60 upright compound

microscope, Nikon A1 confocal microscope or Nikon X1 Yokogawa spinning disc

confocal microscope and processed using ImageJ and quantitation was performed in

ImageJ or CellProfiler.

TUNEL Assay

Apoptosis was analyzed by fluorescence microscopy using *In Situ Cell Death Detection*

Kit (Roche) or CF594 TUNEL Assay Apoptosis Detection Kit (Biotium) according to the

manufacturers' protocols. Histology sections were permeabilized using Proteinase K

(20µg/ml) or 0.2% Triton X-100 in PBS and blocked using PBS+ 5% BSA. Sections

461 were stained with primary antibodies overnight at 4°C in blocking buffer and then were

462 incubated in the terminal deoxynucleotidyl transferase end labeling (TUNEL) buffer for

463 1h at 37°C. Slides were washed with PBS and incubated with fluorescent conjugated

464 secondary antibodies. Sections were then counterstained with DAPI to label DNA. To

465 quantify the TUNEL signal in the HIOs, the percent of the HIO lumen filled with TUNEL+

466 cells was quantified using ImageJ software.

468 **RNA Sequencing and Analysis**

469 Total RNA was isolated from 5 HIOs per group with a total of 4 replicates per condition

470 using the mirVana miRNA Isolation Kit (Thermo Fisher). The quality of RNA was

471 confirmed, ensuring the RNA integrity number (RIN)> 8.5, using the Agilent TapeStation

472 system. cDNA libraries were prepared by the University of Michigan DNA Sequencing

473 Core using the TruSeq Stranded mRNA Kit (Illumina) according to the manufacturer's

474 protocol. Libraries were sequenced on Illumina HiSeq 2500 platforms (single-end, 50 bp

475 read length). All samples were sequenced at a depth of 10.5 million reads per sample or

476 greater. Sequencing generated FASTQ files of transcript reads that were pseudoaligned

477 to the human genome (GRCh38.p12) using kallisto software (53). Transcripts were

478 converted to estimated gene counts using the tximport package (54) with gene

479 annotation from Ensembl (55).

481 **Gene Expression and Pathway Enrichment Analysis**

Differential expression analysis was performed using the DESeq2 package (56) with *P*

values calculated by the Wald test and adjusted *P* values calculated using the

Benjamani & Hochberg method (57).

Quantification and Statistical Methods

RNA-seq data analysis was done using RStudio version 1.1.453. Plots were generated

using ggplot2 (58) with data manipulation done using dplyr (59). Other data were

analyzed using Graphpad Prism 9. Statistical differences were determined using

statistical tests indicated in the fig. legends. The mean of at least 2 independent

experiments were presented with error bars showing standard deviation (SD). *P* values

of less than 0.05 were considered significant and designated by: *$*P < 0.05$, **$**P < 0.01$,

$***P < 0.001$ and *$**** P < 0.0001$.

Ethics Statement

Blood samples were obtained from healthy adult donors according to the protocol

approved by the University of Michigan Institutional Review Board (HUM00044257).

Written consent was obtained from all donors.

Acknowledgements

This work was supported by NIH awards U19AI116482-01 (V.B.Y, J.R.S, C.E.W, and

M.X.O), R21AI13540 (M.X.O) and funds from the University of Michigan Medical School

(J.S.K). A-L.E.L was supported by NIH T32 AI007528. We thank the Host Microbiome

Initiative, Microscopy and Image Analysis Laboratory (MIL), the Comprehensive Cancer

Center Immunology (supported by the NCI and NIH award P30CA046592) and
Histology Cores and the DNA Sequencing Core at University of Michigan Medical
School. We gratefully acknowledge the O'Riordan lab members for helpful discussions,
and Dr. Roberto Cieza for assistance with data management.

Figure legends

Fig. 1. PMNs migrate into the lumen during infection and reduce both the number of infected epithelial cells and the association of bacteria with the epithelial surface

(A) Transmigration of PMNs into the lumen of HIOs was quantified using flow cytometry. HIOs were microinjected with STM or PBS and co-cultured with CFSE-labeled PMNs for 8h. PMNs-HIOs were washed to remove any unassociated PMNs, dissociated into a single cell suspension and subjected to flow cytometry. Percentage of PMNs relative to total cells acquired per PMN-HIO was determined by FlowJo software. (B) Immunofluorescent staining of HIOs microinjected with PBS or STM and co-cultured with PMNs. E-cadherin (green) marks the epithelial lining, MPO (red) is specific to PMNs and DNA is stained with DAPI. (C) Total bacterial burden per HIO or PMN-HIO was enumerated at 8hpi. (D) Quantitation of percent infected cells/HIO or PMN-HIO based on 3 fields per view per HIO. (E) Quantitation of epithelium associated bacteria. Number of bacteria within 50μm of the apical epithelial surface based on E-cadherin staining were counted and normalized per 100μm distance. (F) Representative immunofluorescent staining of HIOs and PMN-HIOs infected with STM. *Salmonella* is stained in green, E-cadherin to mark epithelial cells is shown in red, and DNA stained with DAPI in blue. Graphs show the mean and SD of n≥ 10 HIOs represented by dots from at least two independent experiments. Outliers were removed using the ROUT method with Q=0.1%. Unless otherwise stated, significance was determined by Mann-Whitney test with *p<0.05, **p<0.01.

Fig. 2. PMNs induce epithelial cell death and shedding during *Salmonella* infection

(A) Immunofluorescent images of TUNEL staining of histology sections of HIOs and PMN-HIOs injected with PBS or STM at 8hpi. (B) Quantitation of TUNEL positive cells in the lumen of HIOs and PMN-HIOs from (A). Graphs show the mean and SD of HIOs from 2 independent experiments with n>12 HIOs per group. (C) Quantitation of percent of TUNEL-positive epithelial cells in HIO lumen. The percentage of TUNEL-positive cells that stained positive for E-cadherin in the HIO lumen were assessed. (D) Representative confocal microscopy images of histology sections from STM-injected HIOs or PMN-HIOs at 8h. Sections were co-stained with TUNEL (green), epithelial cell marker E-cadherin (red), and DNA marker DAPI (blue). (E) Confocal microscopy images of histology sections of HIOs and PMN-HIOs that were stained for E-cadherin (red), cleaved Caspase-3 (green), and DNA (blue). Arrows point to cleaved Caspase-3 positive epithelial cells whereas arrowheads point to cleaved Caspase-3 negative luminal epithelial cells.

Outliers were removed using the ROUT method with Q=0.1%. Significance was determined via one-way ANOVA with post-Tukey's test for multiple comparisons where ****$p<0.0001$.

Fig. 3. Inflammasome activation and IL-1 production is mediated by PMNs during infection

(A) Gene expression data presented as $\log_2$(fold change) relative to PBS-injected HIOs

for genes involved in inflammasome/cell death signaling. All genes are significantly

changed from PBS-injected HIOs in at least one condition with p-adjusted value <0.05.

(B) Cytokine levels in culture media of HIOs and PMN-HIOs were quantified using

ELISA. Graphs indicate the mean of n=4 biological replicates +/- SD from media

sampled at 8hpi with 5 HIOs or PMN-HIOs per well. (C) Immunofluorescent staining of

histology sections of PMN-HIOs. Sections were stained for ASC expression (green),

Vimentin (red) to mark PMNs and mesenchymal cells, and DNA (blue) was labeled with

DAPI. (D) Zoom of (C) showing luminal ASC-positive cells (green) with multilobed PMN

nuclei.

Statistical significance was determined by 2-way ANOVA where *p<0.05, ***p<0.001,

****p<0.0001.

Fig. 4. Caspase-1 and Caspase-3 inhibition reduces shedding of infected

epithelial cells in the lumen of PMN-HIOs and differentially affect bacterial burden

and bacterial association with the epithelium

(A) Representative fluorescence microscopy images of TUNEL staining of HIO and

PMN-HIO histology sections. HIOs were microinjected with STM and either cultured

alone or co-cultured with PMNs in the presence of inhibitors for Caspase-1 (z-YVAD),

Caspase-3 (z-DEVD), or DMSO control. (B) Quantitation of the percent of lumen filled

with TUNEL-positive cells of STM-infected HIOs or PMN-HIOs with indicated

treatments. (C) Quantitation of the number of bacteria per cell based on 3 fields per

view per HIO. (D) Quantitation of epithelium associated bacteria. Number of bacteria

within 50µm of the apical epithelial surface were counted and normalized per 100µm

distance. (E) Fluorescent microscopy images of STM-infected PMN-HIO histology

sections. Samples were stained for *Salmonella* (green), E-cadherin (red), and DAPI

(blue).

Unless otherwise stated, graphs show the mean +/-SD of n≥ 10 HIOs represented by

dots from at least two independent experiments. Outliers were removed using the

ROUT method with Q=0.1%. Significance was determined by one-way ANOVA with

post-Tukey's test for multiple comparisons where $*p<0.05$, $**p<0.01$.

REFERENCES

1. , Salmonella Homepage (2021) (September 10, 2021).

2. A. C. Baird-Parker, Foodborne salmonellosis. *Lancet* **336**, 1231–1235 (1990).

3. A.-L. E. Lawrence, *et al.*, Salmonella enterica Serovar Typhimurium SPI-1 and SPI-2 Shape the Global Transcriptional Landscape in a Human Intestinal Organoid Model System. *MBio* **12** (2021).

4. B. H. Abuaita, *et al.*, Comparative transcriptional profiling of the early host response to infection by typhoidal and non-typhoidal Salmonella serovars in human intestinal organoids. *PLoS Pathog.* **17**, e1009987 (2021).

5. M. K. Holly, *et al.*, Salmonella enterica Infection of Murine and Human Enteroid-Derived Monolayers Elicits Differential Activation of Epithelium-Intrinsic Inflammasomes. *Infect. Immun.* **88** (2020).

6. L. A. Knodler, *et al.*, Noncanonical inflammasome activation of caspase-4/caspase-11 mediates epithelial defenses against enteric bacterial pathogens. *Cell Host Microbe* **16**, 249–256 (2014).

7. J. L. Forbester, *et al.*, Interaction of salmonella enterica serovar Typhimurium with intestinal organoids derived from human induced pluripotent stem cells. *Infect. Immun.* **83**, 2926–2934 (2015).

8. L. A. Knodler, *et al.*, Dissemination of invasive Salmonella via bacterial-induced extrusion of mucosal epithelia. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 17733–17738 (2010).

9. M. E. Sellin, *et al.*, Epithelium-intrinsic NAIP/NLRC4 inflammasome drives infected enterocyte expulsion to restrict Salmonella replication in the intestinal mucosa. *Cell Host Microbe* **16**, 237–248 (2014).

10. I. Rauch, *et al.*, NAIP-NLRC4 Inflammasomes Coordinate Intestinal Epithelial Cell Expulsion with Eicosanoid and IL-18 Release via Activation of Caspase-1 and -8. *Immunity* **46**, 649–659 (2017).

11. S. M. Crowley, *et al.*, Intestinal restriction of Salmonella Typhimurium requires caspase-1 and caspase-11 epithelial intrinsic inflammasomes. *PLoS Pathog.* **16**, e1008498 (2020).

12. M. A. Kinnebrew, E. G. Pamer, Innate immune signaling in defense against intestinal microbes. *Immunol. Rev.* **245**, 113–131 (2012).

13. O. H. Pham, S. J. McSorley, Protective host immune responses to Salmonella infection. *Future Microbiol.* **10**, 101–110 (2015).

14. A. Rydström, M. J. Wick, Monocyte recruitment, activation, and function in the gut-
associated lymphoid tissue during oral Salmonella infection. *J. Immunol.* **178**,
5789–5801 (2007).

15. C. Tükel, *et al.*, Neutrophil influx during non-typhoidal salmonellosis: who is in the
driver's seat? *FEMS Immunol. Med. Microbiol.* **46**, 320–329 (2006).

16. B. Amulic, C. Cazalet, G. L. Hayes, K. D. Metzler, A. Zychlinsky, Neutrophil
function: from mechanisms to disease. *Annu. Rev. Immunol.* **30**, 459–489 (2012).

17. V. Brinkmann, *et al.*, Neutrophil extracellular traps kill bacteria. *Science* **303**, 1532–
1535 (2004).

18. E. L. Campbell, *et al.*, Transmigrating neutrophils shape the mucosal
microenvironment through localized oxygen depletion to influence resolution of
inflammation. *Immunity* **40**, 66–77 (2014).

19. B. M. Fournier, C. A. Parkos, The role of neutrophils during intestinal inflammation.
Mucosal Immunol. **5**, 354–366 (2012).

20. S. Gopinath, A. Hotson, J. Johns, G. Nolan, D. Monack, The systemic immune state
of super-shedder mice is characterized by a unique neutrophil-dependent blunting
of TH1 responses. *PLoS Pathog.* **9**, e1003408 (2013).

21. A. C. Chin, C. A. Parkos, Pathobiology of neutrophil transepithelial migration:
implications in mediating epithelial injury. *Annu. Rev. Pathol.* **2**, 111–143 (2007).

22. K. L. Mumy, B. A. McCormick, The role of neutrophils in the event of intestinal
inflammation. *Curr. Opin. Pharmacol.* **9**, 697–701 (2009).

23. A. J. Simpson, A. I. Maxwell, J. R. Govan, C. Haslett, J. M. Sallenave, Elafin
(elastase-specific inhibitor) has anti-microbial activity against gram-positive and
gram-negative respiratory pathogens. *FEBS Lett.* **452**, 309–313 (1999).

24. V. E. Diaz-Ochoa, *et al.*, Salmonella Mitigates Oxidative Stress and Thrives in the
Inflamed Gut by Evading Calprotectin-Mediated Manganese Sequestration. *Cell
Host Microbe* **19**, 814–825 (2016).

25. J. Z. Liu, *et al.*, Zinc Sequestration by the Neutrophil Protein Calprotectin Enhances
Salmonella Growth in the Inflamed Gut. *Cell Host Microbe* **11**, 227–239 (2012).

26. A. Chong, *et al.*, Cytosolic replication in epithelial cells fuels intestinal expansion
and chronic fecal shedding of Salmonella Typhimurium. *Cell Host Microbe* **29**,
1177-1185.e6 (2021).

27. M. Saffarzadeh, *et al.*, Neutrophil extracellular traps directly induce epithelial and
endothelial cell death: a predominant role of histones. *PLoS One* **7**, e32366 (2012).

28. N. Naseer, *et al.*, Salmonella Typhimurium induces NAIP/NLRC4- and NLRP3/ASC-
independent, caspase-1/4-dependent inflammasome activation in human intestinal
epithelial cells. *bioRxiv*, 2021.12.09.472040 (2021).

29. B. H. Abuaita, *et al.*, The IRE1α Stress Signaling Axis Is a Key Regulator of
Neutrophil Antimicrobial Effector Function. *J. Immunol.* (2021)
https:/doi.org/10.4049/jimmunol.2001321.

30. S. A. Hirota, *et al.*, NLRP3 inflammasome plays a key role in the regulation of
intestinal homeostasis. *Inflamm. Bowel Dis.* **17**, 1359–1372 (2011).

31. L. Vande Walle, M. Lamkanfi, Inflammasomes: caspase-1-activating platforms with
critical roles in host defense. *Front. Microbiol.* **2**, 3 (2011).

32. S. M. Man, Inflammasomes in the gastrointestinal tract: infection, cancer and gut
microbiota homeostasis. *Nat. Rev. Gastroenterol. Hepatol.* **15**, 721–737 (2018).

33. P. J. Baker, *et al.*, NLRP3 inflammasome activation downstream of cytoplasmic
LPS recognition by both caspase-4 and caspase-5. *Eur. J. Immunol.* **45**, 2918–2926
(2015).

34. C. P. McEntee, C. M. Finlay, E. C. Lavelle, Divergent Roles for the IL-1 Family in
Gastrointestinal Homeostasis and Inflammation. *Front. Immunol.* **10**, 1266 (2019).

35. Z. Hodzic, E. M. Schill, A. M. Bolock, M. Good, IL-33 and the intestine: The good,
the bad, and the inflammatory. *Cytokine* **100**, 1–10 (2017).

36. E. Lefrançais, *et al.*, IL-33 is processed into mature bioactive forms by neutrophil
elastase and cathepsin G. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 1673–1678 (2012).

37. F. Martinon, K. Burns, J. Tschopp, The inflammasome: a molecular platform
triggering activation of inflammatory caspases and processing of proIL-beta. *Mol.*
Cell **10**, 417–426 (2002).

38. J. R. Kurtz, J. A. Goggins, J. B. McLachlan, Salmonella infection: Interplay between
the bacteria and host immune system. *Immunol. Lett.* **190**, 42–50 (2017).

39. C. A. Lee, *et al.*, A secreted Salmonella protein induces a proinflammatory
response in epithelial cells, which promotes neutrophil migration. *Proc. Natl. Acad.*
Sci. U. S. A. **97**, 12283–12288 (2000).

40. C. Cheminay, D. Chakravortty, M. Hensel, Role of neutrophils in murine
salmonellosis. *Infect. Immun.* **72**, 468–477 (2004).

41. C. A. Parkos, Neutrophil-Epithelial Interactions: A Double-Edged Sword. *Am. J.*
Pathol. **186**, 1404–1416 (2016).

42. Q. L. Ying, S. R. Simon, Kinetics of the inhibition of human leukocyte elastase by
elafin, a 6-kilodalton elastase-specific inhibitor from human skin. *Biochemistry* **32**,
1866–1874 (1993).

43. C. T. N. Pham, Neutrophil serine proteases: specific regulators of inflammation.
Nat. Rev. Immunol. **6**, 541–550 (2006).

44. E. P. Reeves, *et al.*, Killing activity of neutrophils is mediated through activation of
proteases by K+ flux. *Nature* **416**, 291–297 (2002).

45. J.-P. Motta, *et al.*, Food-grade bacteria expressing elafin protect against
inflammation and restore colon homeostasis. *Sci. Transl. Med.* **4**, 158ra144 (2012).

46. J.-P. Motta, *et al.*, Modifying the protease, antiprotease pattern by elafin
overexpression protects mice from colitis. *Gastroenterology* **140**, 1272–1282
(2011).

47. A. Hausmann, *et al.*, Intestinal epithelial NAIP/NLRC4 restricts systemic
dissemination of the adapted pathogen Salmonella Typhimurium due to site-specific
bacterial PAMP expression. *Mucosal Immunol.* **13**, 530–544 (2020).

48. S. H. Jia, *et al.*, Activated neutrophils induce epithelial cell apoptosis through
oxidant-dependent tyrosine dephosphorylation of caspase-8. *Am. J. Pathol.* **184**,
1030–1040 (2014).

49. S. A. Gudipaty, J. Rosenblatt, Epithelial cell extrusion: Pathways and pathologies.
Semin. Cell Dev. Biol. **67**, 132–140 (2017).

50. K. W. Chen, *et al.*, Noncanonical inflammasome signaling elicits gasdermin D-
dependent neutrophil extracellular traps. *Sci. Immunol.* **3**, eaar6676 (2018).

51. J. R. Spence, *et al.*, Directed differentiation of human pluripotent stem cells into
intestinal tissue in vitro. *Nature* **470**, 105–109 (2011).

52. D. R. Hill, S. Huang, Y.-H. Tsai, J. R. Spence, V. B. Young, Real-time Measurement
of Epithelial Barrier Permeability in Human Intestinal Organoids. *J. Vis. Exp.* (2017)
https:/doi.org/10.3791/56960.

53. N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq
quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

54. C. Soneson, M. I. Love, M. D. Robinson, Differential analyses for RNA-seq:
transcript-level estimates improve gene-level inferences. *F1000Res.* **4**, 1521
(2015).

55. J. Rainer, EnsDb.Hsapiens.v75: Ensembl based annotation package. R package
version 2.99.0 (2017).

56. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

57. Y. Benjamini, Y. Hochberg, Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society:*
Series B (Methodological) **57**, 289–300 (1995).

58. H. Wickham, *ggplot2: Elegant Graphics for Data Analysis* (Springer, 2016).

59. H. Wickham, R. Francois, L. Henry, K. Müller, Others, dplyr: A grammar of data
manipulation. *R package version 0. 4* **3** (2015).

STM-infected HIO　　　　　　　　　　　**STM-infected PMN-HIO**

Figure 1

A PMN transmigration

PMN transmigration (% of total cells)

PBS   STM

B

PMN-HIO

PBS   STM

Lumen   Lumen

50 μm   E-cadherin MPO DAPI   8hpi

C Total HIO colonization

CFU/HIO

HIO   PMN-HIO

D Percent infected cells

%infected cells/HIO section

HIO   PMN-HIO

E Epithelium associated bacteria

Ave# STM within 50μm of apical surface

HIO   PMN-HIO

F

HIO

Lumen

Epithelium associated bacteria

Epithelium

Salmonella   E-cadherin   Merge

PMN-HIO

Lumen

Epithelium associated bacteria

Epithelium

Salmonella   E-cadherin   Merge

50μm

Figure 2

A

PBS STM

HIO

PMN-HIO

500 µm

TUNEL DAPI

B

%TUNEL+ in HIO lumen

%TUNEL⁺ area

**** ****

PBS STM PBS STM
HIO   PMN-HIO

C

TUNEL⁺ E-cadherin cells in HIO lumen

%TUNEL⁺ E-cadherin⁺ cells/HIO

PMN-HIO

D

STM-infected HIO STM-infected PMN-HIO

DAPI

TUNEL

E-cadherin

Merge

200µm

E

STM-infected PMN-HIO

50µm

Cleaved Caspase-3 E-cadherin DAPI

Figure 3

A Inflammasome/Cell Death

IL1B
IL1RN
IL1A
NLRP3
NFKB2
CASP5
CASP10
IL33
IL1R1
NFKB1
CASP1
CASP4
TXN
CASP3
PANX1
RELA
SUGT1
TXNIP
NLRP1

log₂(fold change)

STM / HIO
PBS STM / PMN-HIO

B

IL1β — pg/ml — PBS STM | PBS STM — HIOs | PMN-HIOs — **** ****

IL1α — pg/ml — PBS STM | PBS STM — HIOs | PMN-HIOs — **** ****

IL1RA (IL1RN) — pg/ml — PBS STM | PBS STM — HIOs | PMN-HIOs — **** ****

IL33 — pg/ml — PBS STM | PBS STM — HIOs | PMN-HIOs — *** *

C STM-infected PMN-HIO

50μm

ASC Vimentin DAPI

D STM-infected PMN-HIO

5μm

ASC Vimentin DAPI

Figure 4

B %TUNEL⁺ in HIO lumen

C Number of bacteria per cell

D Epithelium associated bacteria

Scale bars: 250μm, 50μm

TUNEL DAPI

Salmonella E-cadherin DAPI