Titles: Mark loss can strongly bias demographic rates in multi-state models: a case study with simulated and empirical datasets

Authors

Frédéric Touzalin[1,2], Eric J. Petit[3], Emmanuelle Cam[4], Claire Stagier[1], Emma C. Teeling[1], Sébastien J. Puechmaille[5,6,7]

[1]School of Biology and Environmental Science, Science Centre West, University College Dublin, Dublin, Ireland.

[2]Bretagne Vivante-SEPNB, Brest, France.

[3]DECOD (Ecosystem Dynamics and Sustainability), INRAE, Institut Agro, Ifremer, Rennes, France.

[4]Université de Bretagne occidentale, Brest, LEMAR, CNRS, IRD, Ifremer, F-29280 Plouzane, France.

[5]Zoological Institute and Museum, University of Greifswald, Greifswald, Germany.

[6]ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France.

[7]Institut Universitaire de France, Paris, France

Correspondence

Frédéric Touzalin

Email: fredtouzalin@gmail.com

Abstract

1. The development of methods for individual identification in wild species and the refinement of

Capture-Mark-Recapture (CMR) models over the past few decades have greatly improved the

assessment of population demographic rates to address ecological and conservation questions. In

particular, multi-state models, which offer flexibility in analysing complex study systems, have

gained popularity within the ecological community. In this study, we focus on the issue of mark loss

and the associated recycling of remarked individuals, which requires further exploration given the

increasing use of these models.

2. To fill this knowledge gap, we employed a wide range of simulation scenarios that reflect

commonly encountered real case studies, drawing inspiration from the survival rates of 700

vertebrate species. Using a multi-state, Arnason-Schwartz (AS) modelling framework, we estimated

the effects of mark loss and recycled individuals on parameter estimates. We assessed parameter

bias by simulating a metapopulation system with varying capture and survival rates. Additionally,

we demonstrated how mark loss can be easily estimated and accounted for using a 10-year

empirical CMR dataset of bats. The bats were individually identified using Passive Integrated

Transponder (PIT) tag technology as potentially lost marks and multi-locus genotypes as 'permanent

marks'.

3. Our simulation results revealed that the occurrence of bias and the affected parameters were

highly dependent on the study system, making it difficult to establish general rules to predict bias *a*

priori. The model structure and the interdependency among parameters pose challenges in

predicting the impact of bias on estimates.

4. Our findings underscore the importance of assessing the effect of mark loss when using AS

models. Ignoring such violations of model assumptions can have significant implications for

ecological inferences and conservation policies. In general, the use of permanent marks, such as

genotypes, should always be preferred when modelling population dynamics. If that is not feasible,

an alternative is to combine two independent types of temporary marks, such as PIT tags and bands.

5. Analysis of our empirical dataset on *Myotis myotis* bats revealed that tag loss is higher in

juveniles than in adults during the first year after tagging. The use of surgical glue to close the

injection hole reduces tag loss rate from 28% to 19% in juveniles, while it has no effect on the tag

loss rate in adults (~10%). The main bias observed in our metapopulation system appears in the

survival rate, with up to a 20% underestimation if tag loss is not accounted for.

KEYWORDS

Arnason-Schwarz model, Bayesian, bats, capture-mark-recapture, mark retention, *Myotis myotis*,

multi-state, surgical glue.

1   Introduction

Capture-mark-recapture (CMR) methods have become a standard approach for estimating

demographic rates of wild species (Johnson *et al.*, 1986; Williams *et al.*, 2002). Accurately

quantifying population dynamic parameters is crucial for assessing population status, understanding

dynamics, and making effective management and conservation decisions. However, CMR models

rely on certain assumptions of homogeneity (Johnson *et al.*, 1986; Williams *et al.*, 2002) that can

introduce biases if violated. One common violation of CMR assumptions, first identified four

decades ago (Nelson *et al.*, 1980), is the loss of marks (see Supporting Information 1, Table S2).

Mark loss has two implications: (1) when marks are shed, it leads to non-identifiability of

individuals (detection heterogeneity), causing them to be considered dead or out of the study area,

even when they are alive and present; (2) if these individuals are recaptured, they will go

unrecognized and be counted as newly recruited individuals, referred to as "recycled" individuals

(Malcolm-White *et al.*, 2020). In open population models, estimates of abundance in the Jolly-Seber

(JS) model (Jolly, 1965; Seber, 1965) or survival in the Cormac-Jolly-Seber (CJS) model (Cormack,

1964; Jolly, 1965; Seber, 1965) can be affected by mark loss (Arnason & Mills, 1981). Several

statistical techniques have been developed to mitigate the confounding effect of mark loss in these

models (Arnason & Mills, 1981; Cowen & Schwarz, 2006; Robson & Regier, 1966; Seber &

Felton, 1981). However, the impact of mark loss on state transition in multi-state models, which

have undergone significant development (Lebreton *et al.*, 2009), has not been extensively studied.

In multi-state models, if survival is state dependent, it is the product of true survival and mark

retention rate for individuals in a specific state (Lebreton *et al.*, 1992; Lebreton *et al.*, 2009). If the

retention rate drops below one without being considered in the model, the estimation of survival in

a particular state is underestimated and becomes confounded with the probability of presence of the

mark. This is especially true if true survival is high, but it is unclear how state transitions are

affected.

Let's consider an example encounter history: "1011", where "1" indicates the individual was

captured and "0" indicates otherwise. If we denote ϕ_t as the survival rate between occasions t and

t+1, and p_t as the capture probability at occasion t (with $q_t = 1 - p_t$), this encounter history occurs

with probability $\phi_1 q_2 \phi_2 p_3 \phi_3 p_4$. We can break down this probability product as follows: the individual

survives between t_1 and t_2 (ϕ_1), was not captured in t_2 (q_2), survives between t_2 and t_3 (ϕ_2), was

captured in t_3 (p_3), and finally survives between t_3 and t_4 (ϕ_3) and was captured in t_4 (p_3). Now, let's

consider that the individual can transition between two states, and its history becomes "1022", with

the individual in state "1" at t_1 and in state "2" at t_3 and t_4. At t_2, when the individual was not

detected, two possibilities arise: either it stayed in state 1 or made a transition to state 2. To account

for these two possible histories, we introduce $\psi^{i,j}$ as the transition probability from state *i* to state *j*

(where *i* and *j* are in {1,2}), conditional on survival. The new encounter history is now the sum of

two components o account for the two possible his histories "1122" or "1222":

$\phi_1^{(1)}\psi^{(1,1)}q_2^{(1)}\phi_2^{(2)}\psi^{(1,2)}p_3^{(2)}\phi_3^{(2)}\psi^{(2,2)}p_4^{(2)} + \phi_1^{(1)}\psi^{(1,2)}q_2^{(2)}\phi_2^{(2)}\psi^{(2,2)}p_3^{(2)}\phi_3^{(2)}\psi^{(2,2)}p_4^{(2)}$, with the superscript

denoting state-specific parameters in parentheses. However, if this individual loses its mark after its

first capture (indicated when recaptured at t_3 and not recognized), its encounter history becomes two

different histories from two different individuals: one becomes "1000", and the second becomes

"0022". In this case, survival and mark loss patterns differ. Not only is survival underestimated (at

least for the "first" history), but the transition probabilities are also underestimated, since there is no

longer a change of state (the second history starts directly at state 2).

Most studies using CMR techniques are affected by mark loss, which can vary based on several

factors such as species (see Supporting Information 1, Table S2), mark type (Smout *et al.*, 2011a),

sex (Conn *et al.*, 2004), mass (Schwarz *et al.*, 2012), size (Acolas *et al.*, 2007), mark location

(Kaemingk *et al.*, 2011), or physiological stage (Besnard *et al.*, 2007). This mark loss has been

previously shown to introduce negative bias in survival estimates and detection (Nichols *et al.*,

1992; Nichols & Hines, 1993). To address situations where an individual's "state" (e.g., location,

behaviour, physiology, reproductive or social status) may affect its survival or detection probability,

and where the individual can change "state" during its lifespan, multi-state models have been

developed (reviewed in Lebreton *et al.*, 2009). These models have gained popularity due to their

flexibility in studying a wide range of ecological systems and biological questions. Biologists can

easily use user-friendly software like Mark (White & Burnham, 1999), WinBUGS (Lunn *et al.*,

2000), JAGS (Plummer, 2003), E-SURGE (Choquet, *et al.*, 2009b), MultiBUGS (Goudie *et al.*,

2020), NIMBLE (de Valpine *et al.*, 2017), and STAN (STAN Development Team, 2022) to

implement these models. They are used to examine ecological and evolutionary hypotheses related

to variations in life history traits (state transitions) over an individual's lifespan (Nichols & Kendall,

1995; see also Cam 2009 for a detailed discussion), density dependence effects (Schofield &

Barker, 2008), co-evolution (Benkman *et al.*, 2005), dispersal probability among subpopulations or

habitats (Hestbeck *et al.*, 1991; Spendelow *et al.*, 1995), and disease prevalence in wild populations

(Jennelle *et al.*, 2007). However, there is limited literature on the impact of mark loss on the

behaviour of multi-state models, and further exploration is needed in this area (Seber & Schofield,

2019).

To bridge this gap, we used simulation-based Arnason-Schwartz (AS) model approaches (Arnason,

1972, 1973; Schwarz *et al.*, 1993) to investigate the impact of mark loss on estimates of model

parameters within a Bayesian framework. With the increasing utilization of multi-state models, our

aim is to evaluate the potential bias in the marginal posterior distributions of demographic

parameter estimates using a metapopulation context, based on biologically realistic scenarios, and

provide comprehensive guidelines for both fieldwork and data analyses. The AS model shares

assumptions with the CJS model, specifically regarding mark loss, but additionally assumes that

states are recorded without error. Similarly to the CJS model, we anticipated that the AS model

might underestimate survival and transition probabilities in the event of tag loss and recycling

(Nichols & Hines, 1993). Since transition probabilities are conditional on survival and detection of

state in our AS model, we expected estimation errors to have varying effects on model parameters

depending on the state transition rate.

To illustrate our approach with an empirical example, we used our decade-long mark recapture

dataset of PIT-tagged and genotyped greater mouse-eared bat (*Myotis myotis*), a taxonomic group

particularly vulnerable to PIT-tag loss (Freeland & Fry, 1995). We employed genotypes as

individual permanent marks to estimate the bias between models that account for mark loss and

recycling versus those that do not, and finally, we suggest recommendations for future studies.

2   Material and method

To quantify the potential bias induced by mark loss on parameter estimates in the AS framework,

we defined various scenarios based on representative situations derived from a compilation data

obtained from the Demographic Species Knowledge Index (Conde *et al.*, 2019) and literature

sources for fish and bat species (Fig. 1, Supporting Information 1, Table S1). These data were

limited to published CMR studies or data from controlled conditions, specifically marked

individuals with known outcomes, such as those observed in zoos. Among the six vertebrate classes

(Actinopterygii, Chondrichthyes, Amphibia, Aves, Reptilia, and Mammalia) encompassing 700

species, the survival rates exhibit a wide range of values (Fig. 1). The relationship between adult

and juvenile survival, available for 143 species (Supporting Information 1, Fig. S1), reveals that low

adult survival is associated with low juvenile survival, while high adult survival can be

accompanied by a diverse range of juvenile survival rates. To limit the number of scenarios to

explore, we selected values towards the extremes. Consequently, we considered two hypothetical

populations: one consisting of long-lived species with high survival rates in both age classes (e.g.,

large mammals), and the other consisting of short-lived species with low survival rates in both age

classes (e.g., amphibians). For each population, we explored cases with high and low detection

probabilities, and for each case, we tested three different mark loss rates, obtained from relevant

literature sources (Supporting Information 1, Table S2). Initially, we present the procedure for

generating our simulated scenarios, followed by the description of the two different models used to

analyse these data: one model that does not account for mark loss, and another model that

incorporates mark loss by utilizing a second permanent mark, allowing for comparison of

estimations. Lastly, we outline the metrics used to evaluate the potential bias in parameter estimates

when mark loss is not accounted for in the AS framework.

167 **2.1 Data generation**

For each scenario, we conducted a simulation using data from a ten-year study period with multiple

capture occasions. During these capture occasions, individuals could be in one of five possible

states: "A", "B", "C", "D", or death itself. State "D" was considered an absorbing state, meaning that

once an individual reached this state, it could not change to any other state. This can be used to

model permanent emigration, for example. In the initial occasion, the individuals in states "A", "B",

and "C" consisted of 40 juveniles (with a 1:1 sex ratio) and 60 adults (80% females, 20% males).

No individuals were initially in state "D". In each subsequent occasion, we marked 40 juveniles and

5 adults in each state ("A", "B", "C"), except for state "D", where recapture was not possible and

only observation was feasible. This conceptual system can be seen as having three breeding sites

("A", "B", "C") where capture and resighting occurred annually, and a surrounding area ("D")

where only resighting was possible. For males only, we allowed a permanent transition to the "D"

state. It is worth noting that sexual asymmetry in state change is common in various taxa, where one

sex is more likely to disperse through permanent emigration (absorbing state). For each simulated

scenario, all juveniles captured at a specific occasion were considered adults in the following

occasion. All scenarios followed the same steps for data generation, as shown in Figure 2. We

initiated the simulation by generating the survival of individual i at occasion t using a Bernoulli

distribution:

$$a_{i,t+1} \sim \begin{cases} Bernoulli(a_{i,t}\phi_{i,t}), \text{if } a_{i,t} = 1, \\ 0, \text{if } a_{i,t} = 0 \end{cases} \tag{1}$$

185   where $a_{i,t} = 1$ if individual i is alive at time t, and 0 if not. $\Phi_{i,t}$ represents the specific survival

186   probability for each individual at a given state, time, and age. To introduce stochastic variations in

187   annual survival for the two age classes, we drew survival probabilities from a logistic distribution.

188   If an individual survived, we simulated the state transition using a Categorical distribution:

$$s_{i,t} \sim Categorical(\psi_{i,t}), \text{ if } A_{i,t} = 1, \tag{2}$$

where $\psi_{i,t}$ represents a state-, sex-, and age-specific transition probability. The study examined the

transitions of females between states "A", "B", and "C" at a constant rate, regardless of their age.

However, females were not allowed to transition to state "D". Juvenile males, on the other hand,

could only transition to state "D", with the proportion depending on their initial state. Adult males,

however, did not change their state throughout the study. The mark loss/retention process was

simulated using a Bernoulli distribution:

$$m_{i,t} \sim \begin{cases} Bernoulli(pr_{i,t}), \text{ if } ts_{i,t-1} = 1 \text{ and not captured at } t, \\ 1, \text{ if captured at } t \text{ (newly marked or recycled)}, \\ 0, \text{ if } ts_{i,t-1} = 0 \text{ and not captured at } t \end{cases} \tag{3}$$

195   with $m_{i,t}$ indicating the presence of a mark on individual i at occasion t and $pr_{i,t}$ representing the

196   probability of retention (see Supporting Information 2, Fig. S2), which is the complement of the

197   mark loss probability pl ($pl_{i,t} = 1-pr_{i,t}$). The detection process was simulated using a Categorical

198   distribution:

$$z_{i,t} \sim \begin{cases} Categorical(pd_{i,t}), & \text{if } A_{i,t} = 1, \\ \text{not detected,} & \text{if } A_{i,t} = 0 \end{cases} \tag{4}$$

with $pd_{i,t}$ representing the state-specific detection probability (Supporting Information 2, section

1.2). We examined two common methods for tracking individuals in our study: physical capture and

resighting. Physical capture involves either recycling the individuals by re-marking them or

checking their existing marks. Resighting, on the other hand, is a passive detection method that only

includes individuals who have retained their marks. This approach is based on the observation that,

in many studies, the probability of resighting is usually higher than the probability of recapture,

making them susceptible to different estimation biases. We categorized the probability of detection

into seven categories, with each category conditioned on the retention of the mark.

$$pd_{i,t} = \begin{cases} pc_{i,t}(1 - ts_{i,t}p.r_{i,t}), & \text{if mark retained and caught in state "A","B" or "C",} \\ (1 - pc_{i,t})ts_{i,t}p.r_{i,t}, & \text{if mark retained and resighted in state "A","B" or "C",} \\ pc_{i,t}ts_{i,t}p.r_{i,t}, & \text{if mark retained and both caught and resighted in state "A","B" or "C",} \\ pb, & \text{if mark retained and resighted in state "D",} \\ 1 - pb, & \text{if mark retained and not resighted in state "D",} \\ (1 - pc_{i,t})(1 - ts_{i,t}p.r_{i,t}), & \text{if undetected in state "A","B" or "C",} \\ 0, & \text{if mark lost} \end{cases} \tag{5}$$

where $pc_{i,t}$ was a state specific capture probability, $p.r_{i,t}$ a state specific resighting probability and pb

the detection probability in state "D".

Since we can expect higher rates of recycling when recapture rates are high, and little recycling

otherwise, we considered four scenarios (Table 1): (1) long-lived species with a high detection rate;

(2) long-lived species with a low detection rate; (3) short-lived species with a low detection rate; (4)

short-lived species with a high detection rate. The values of the simulated parameters can be found

in Table 2 and Figure 3. Mark loss often depends on the time since marking, and in many species, it

occurs soon after marking (Fabrizio *et al.,* 1999; Fokidis *et al.,* 2006; Kremers, 1988; Nichols &

Hines, 1993). In our scenarios, we included this process and determined a range of mark loss rates

from the literature (see Supporting Information 1, Table S2). We simulated three mark loss

probabilities: low (*pl* = 0.05), medium (*pl* = 0.25), and high (*pl* = 0.4) during the first year after

marking, and a constant rate of 0.05 thereafter. This generated a variety of cases of mark loss and

recycled individuals (see Supporting Information 2, Fig. S2-3). This process enabled us to generate

datasets that took mark loss into account in the presence of a second permanent mark. In order to

generate datasets where mark loss was not considered (no second permanent mark), we created

recycled individuals with life histories corresponding to the portion of their lives after mark loss,

and replaced the original life history from mark loss with zeros. For example, a life history "1111"

of an individual that lost its mark between occasions 2 and 3, and was re-marked at occasion 3,

would become two new histories: (1) "1100", representing the first part of the life before mark loss,

and (2) "0011", representing a second individual that was not recognized in the absence of a

permanent mark, but considered newly recruited. Using this data generation process, we simulated 2

x 50 datasets for each of the 12 combinations of parameters (50 with and 50 without recycling),

resulting in a total of 1200 simulated datasets (see Supporting Information 2, Fig. S1). The

computational codes for a fully reproducible example dataset are provided in the Supporting

Information 2.

233   2.2   Statistical models

As described, we simulated two different datasets for each combination, in which the assumption of

permanent marking is violated. In the first dataset, individuals can still be identified even after the

loss of their mark, thanks to a second permanent mark. In the second dataset, recycling occurred

because there was no second permanent mark present (Supporting Information 2, Fig. S3). We

developed two AS models to analyse these data. ModelA, used for the first dataset, included the

estimation of mark loss. ModelW, used for the second dataset, ignored mark loss (Supporting

Information 2, sections 1.7 and 2.2).

To estimate the state, time, and age-dependent survival Φ for both models, we used a Bernoulli

distribution (see Eq. 1 above).

$$p(a_{i,t+1}|a_{i,t}) \sim Bernoulli(a_{i,t}\phi_{i,t}),$$
$$logit(\phi_{i,t}) = \alpha_{state_{i,t+1}} + \beta_{t+1} + \delta_{age_{i,t+1}} + \gamma_{state_{i,t+1},t+1,age_{i,t+1}}$$

(6)

where $a_{i,t}$ is the life status of individual i at time t and coefficient α is a state effect, β a time effect, δ

an age effect and γ a simultaneous effect of state, age and time. The transition and detection

processes were estimated using the same distributions as described for data generation (see Eq. 2, 4,

5 above). Only in Model A, the fate of marks (whether they were retained or lost) for each

individual was tracked. Various combinations of single, double, and permanent marks have been

used to estimate mark loss (Laake *et al.*, 2014). In Model A, we used a single mark loss approach,

with the second mark being permanent. This allowed us to determine whether the non-permanent

mark was lost or retained at each capture occasion, and resighting was conditional on mark

retention (see Eq. 5). The mark retention process was directly incorporated into the model using a

state space approach. While it is possible to model the presence-absence of marks as states with a

transition matrix (McMahon & White, 2009), or as a hidden Markov process for unobserved

individuals where mark retention status is unknown (Laake *et al.*, 2014), we chose to model the

mark retention process using a Bernoulli distribution with p($m_{i,t}$|$m_{i,t-1}$) ~ Bernoulli($pr_{i,t}$), if the

individual was marked or retained its mark at *t-1* ($m_{i,t-1}$ = 1). We estimated three retention

probabilities (*pr*) as a function of age and time since marking:

$$pr = \begin{cases} pr_{juv.}, & \text{if marked when juvenile,} \\ pr_{adlt.,1}, & \text{at the first occasion after marking, if marked as an adult,} \\ pr_{adlt.,2}, & \text{for subsequent occasions (retained at least 2 occasions), if marked as an adult.} \end{cases}$$

(7)

258 When an individual was lost and subsequently recaptured, a new mark was applied in modelA

259 without altering its identity. In contrast, modelW does not incorporate mark retention, so if an

260 individual lost its mark and was recaptured, it was regarded as newly recruited. The priors utilized

261 for each parameter can be found in Supporting Information 2, section 1.6.

2.3 Application on a bats dataset

The simulations we conducted served two purposes: first, to test general hypotheses regarding the impact of mark loss on parameter estimates in multi-state models, and second, to validate a parametrization approach that can accurately estimate these parameters without bias. Based on this, we are able to accurately estimate the probability of mark loss in our own dataset and determine the importance of using a permanent mark in the long term (Juillet *et al.*, 2011). Our empirical dataset consists of a 10-year study on the greater-mouse eared bat (*Myotis myotis*) in Brittany, France (2010-2019). We marked a total of 2,561 individuals in 5 maternity roosts: La Roche Bernard (47°31'N, 2°18'W), Férel (47°28'N, 2°20'W), Noyal-Muzillac (47°35'N, 2°27'W), Béganne (47°35'N, 2°14'W), and Limerzel (47°38'N, 2°21'W). The bats were individually tagged using Passive Integrated Transponders (PIT) tags, specifically the ID-100C (Troven®), which had a unique 10-digit code. These small tags (2.12x11mm, 0.1gr) allowed for identification using passive readers. All individuals captured in roosts without PIT-tags were consistently tagged, including both those who lost their tags and those who were never tagged before. Additionally, a second permanent marking method involved genotyping all newly tagged individuals. Genotypes were determined from DNA extracted from wing biopsies of all tagged individuals and all untagged males captured during swarming surveys (n=4,258 samples; see Supporting Information 3, Fig. S1). A total of 16 microsatellite markers optimized for *Myotis myotis* were used to establish individual genotypes (Foley *et al.*, 2020). To minimize genotyping errors, we analysed two different samples per individual, whenever available. All samples were genotyped and scored twice by two different individuals (Frantz *et al.*, 2003; Puechmaille & Petit, 2007). We also conducted genetic profile comparisons to identify errors. The error rate of genotypes was hypothesized to be sufficiently low to be negligible, and this source of uncertainty was not included in the models (but see Winiarski & McGarigal, 2016). In addition, we checked for lost tags on the floor of the maternity roosts during each winter. This allowed us to partially identify individuals that lost their tags (61.5% compared to genotype), with the remaining losses occurring outside of the roosts. Most tag losses occurred

during the first year, as supported by the absence of records from passive reading detectors

(Supporting Information 3, Fig. S2). Out of a total of 2,561 individuals, 252 (~10%) were identified

as having lost their tag at least once. Of these individuals, 94 were recaptured and retagged once,

while three were retagged twice. For 13 individuals, retagging occurred during the last capture

occasion, resulting in the recycling of 81 individuals out of the 94. To analyse these data, we

employed a multisite model that accounted for movement between maternity roosts as transitions

(similar to the AS model used for simulated data, Supporting Information 3). We divided the

population into two age classes for survival analysis: juveniles (in their first year of life) and adults

(one year and older). To examine departures from the AS model assumptions, we conducted

goodness-of-fit tests using R2ucare (Gimenez *et al.*, 2018b), an R package based on U-CARE

(Choquet *et al.*, 2009a). Minor overdispersion was observed in tests for transience ($\hat{c}_{3G.SR}$=1.82) and

memory ($\hat{c}_{WBWA}$ = 1.96). Insufficient data in the individual contingency tables prevented the

performance of other goodness-of-fit tests. Hence, we retained our two-age class structure for

survival modelling. To account for unmeasured individual survival heterogeneity, we included a

normally distributed random effect in the estimation of survival probability. This was necessary as

other covariates were unable to capture the relevant variation. Individual heterogeneity plays a

crucial role in population dynamics and evolution and is prevalent in wild populations (Gimenez *et

al.*, 2018a).

Emigration from the five subpopulations under study was assessed using capture and resighting data

collected at swarming and wintering sites between capture occasions. We defined eight different

detection states, which enabled us to separately estimate capture and resighting probabilities (see

Supporting Information 3, Table S1). Based on empirical data suggesting potential movement of

individuals between all subpopulations and outside areas, we did not restrict transitions between

subpopulations, except for the movement of juveniles from outside the maternity roosts, which was

not possible due to the absence of tagging outside the five roosts. Therefore, the probability for this

type of movement was set to 0 (see Supporting Information 3, Fig. S3). This study also examined

the impact of using surgical adhesive (Vetbond®) after PIT-tag injection to reduce tag loss

compared to self-healing techniques (Lebl & Ruf, 2010; van Harten *et al.*, 2020). In this model, tag

retention probabilities were modelled similar to Eq. 7, taking into account time since marking

(categorized as first year or subsequent years), individual age class (juvenile or adult), and the use

of surgical adhesive (yes or no). Similar to the simulated datasets, two models were run on two

datasets: one for estimating tag loss using genotyping (as a second permanent mark), and another

model that ignored this information on a transformed dataset that included recycled individuals

(following the same process as the simulated data).

322 2.4   Estimation procedures and assessments

Despite the possibility of using a frequentist approach for analysing simulated data (Lebreton *et al.*,

2009), we opted for a Bayesian approach to ensure consistency with the analysis of empirical data,

where these methods offer greater flexibility in accounting for individual heterogeneity in survival

(Gimenez *et al.*, 2018a). The simulated and empirical data were analysed using JAGS (Kruschke,

2014; Plummer, 2003) through the jagsUI package (Kellner, 2016) in R 3.6.0 (R Core Team, 2019).

We employed four Monte Carlo Markov chains (MCMC) with 150,000 iterations each and

discarded the initial 50,000 iterations (burn-in) when sampling from posterior distributions. We

retained every twentieth iteration, resulting in 20,000 samples from the posterior distribution for

each parameter. Chain convergence was assessed using the Gelman-Rubin statistic, denoted R-hat

(Brooks & Gelman, 1998). Among the 1,200 simulations, some parameters exhibited R-hat values >

1.05, indicating convergence failure. Less than 0.4% of the estimated parameters for the model

accounting for mark loss did not converge (Supporting Information 2, Table S1), particularly the

coefficient γ (combined effect of state, time, and age on survival probability) and $\gamma.c$ (combined

effect of state and time on detection probability). The mean R-hat values for these parameters were

less than 1.2 (Supporting Information 2, Fig. S4). For models that did not account for mark loss,

1.3% of the estimated parameters did not converge (Supporting Information 2, Table S2), especially

in scenarios with low detection or low detection and survival rates (scenario 2 and 3, respectively),

and mark loss probabilities set to 0.25 or 0.4. In most cases, the parameters that failed to converge

exhibited a mean R-hat value less than 1.2, with only a few exceeding 1.5 (Supporting Information

2, Fig. S5). Convergence failures did not pertain to the probability of mark loss in any of the

simulations. To avoid excessively long computing times, we did not increase the number of

iterations to achieve complete convergence of the MCMC chains for these parameters in the

affected simulations. Our results are based on 50 simulated datasets per scenario, and we assume

that the lack of convergence for these few parameters has no substantial influence on our results. To

assess bias in parameter estimates when mark loss or recycling is not accounted for, we calculated

the Earth Mover Distance (EMD) using the EMD-L1 algorithm (Ling & Okada, 2007), which

quantifies the difference between two distributions. This metric measures the minimum cost of

transforming one distribution into another point by point. For each scenario, we also estimated a

Region Of Practical Equivalence (ROPE, Kruschke, 2018) to assess the degree of difference

between the distributions represented by the EMD metric. To define the ROPE for each scenario,

we randomly constructed 1,000 pairs of models from the 50 simulations and calculated the

associated 1,000 EMDs from the posterior distributions of the estimated parameters (for more

details, see Supporting Information 2, Fig. S76). The resulting distributions of EMDs represented

the expected variations in inferences obtained from simulations initiated with the same parameter

values. The ROPE was then defined as the range between 0 and the upper value of the 80% highest

posterior density interval (hdi) from the distribution of these EMDs. Finally, the proportion of

EMDs for each simulated case that fell outside the ROPE was computed, providing a direct

indication of bias; the higher this proportion, the greater the bias. Comparisons of EMDs between

the models that accounted for tag loss and recycling, and those that did not, in relation to their

respective ROPEs, depict cases in which not accounting for tag loss leads to estimates that

substantially differ from estimates obtained when considering tag loss. We also evaluated parameter

bias as the difference between the median of the posterior distribution and the simulated true value

(median – truth). In the analysis of empirical data, we used the median of the posterior distribution

of parameters from the model accounting for a secondary permanent mark as the true value. Finally,

we calculated the precision of the parameter estimates in the simulated data using mean squared

errors (MSE = bias^2 + variance).

369 3   Results

370 3.1   Simulation results

The estimated numbers of lost marks and recycled individuals increased in scenarios with higher

survival, detection, and mark loss rates (Supporting Information 2, Fig. S3). As expected, when the

recapture rate was high (i.e. in scenarios 1 and 4), the proportion of recycled individuals relative to

the number of lost marks was higher for an equivalent rate of mark loss. Regardless of the

scenarios, there was no estimation bias on demographic parameters when the mark loss rate was set

to 0.05 (Supporting Information 2, Fig. S28-S31). However, the number of parameters with biased

estimates increased with the rate of mark loss, but the magnitude of bias varied across the simulated

scenarios. Specifically, in scenario 1 (Fig. 4 & 5), there were substantial underestimations in some

adult and juvenile survival rates. The probability of remaining in the same state was also

underestimated among juveniles, leading to an overestimation of their probability of transitioning to

another state (Fig. 4 & 6). The resighting probability was underestimated in all states except B (Fig.

4 & 6). The biases in state transitions were particularly high for transitions to absorbing states, such

as juveniles transitioning to state "D" (Supporting Information 2, Fig. S57-S71). In simulations with

the highest mark loss rate (0.4), scenarios 1 and 4 (which had high recapture probabilities) showed

underestimates of adult survival only in the early years of the study (Supporting Information 2, Fig.

S21-S22 and S27). The bias in juvenile survival was less pronounced, although moderate

underestimations occurred for high mark loss rates, especially for state A and B in scenarios 1 and 4

(Supporting Information 2, Fig. S21-S22 and S27). Lack of precision in the estimate of juvenile

survival was also observed when the model did not account for tag loss and the mark loss rate was

high. The resighting probability showed substantial bias, with underestimates mainly in states A and

C (Fig. 4 and 7.a), as well as a lack of precision for all scenarios and mark loss rates (Supporting

Information 2, Fig. S28-S49). However, recapture, the second component of detection, showed little

bias except during the second capture occasion for states A and C in scenario 1 when the mark loss

rate was 0.25, and scenarios 1 and 4 when the mark loss rate was 0.4. There was also a decrease in

the precision of the recapture parameter at high mark loss rates. When mark loss and recycling were

ignored, a large percentage of the transition probability estimates were biased (Fig. 6b-e), with an

overall underestimation of the probability to remain in the same state and, as a result, an

overestimation of the probability of changing state. This decrease in precision mainly occurred at

high mark loss rates (Supporting Information 2, Fig. S50-S71). Overestimations of transition

probabilities were observed in juvenile males, except for transitions from state C to D, where the

transition rate was set to 0.9. For females, when mark loss was set at a high level, the same pattern

was observed in states A and B, from which state transitions were set at a low level. On the contrary,

when the transition from state C to B was set at a high level (0.4), there was an underestimation of

the transition probability and an overestimation of the probability to remain in state C, particularly

in scenarios with low detection rates (scenario 2 and 3, Supporting Information 2, Fig. S67-S70).

3.2   Bat metapopulation

Most of the estimated parameters, such as survival, capture, resighting, and state transition

probabilities, exhibited both negative and positive biases without a clear pattern (Fig. 7, Supporting

Information 3, Fig. S4-S11). The largest biases were observed in survival estimates, with a median

underestimation of survival of over 0.26 in juveniles (Fig. 7a) and 0.21 in adults (Fig. 7b).

Emigration, defined as movement outside the studied maternity sites ("Out"), was overestimated by

an average of 0.05 throughout the study (Fig. 7b). The probability of recapture varied from

overestimation to underestimation by up to 0.1, depending on the occasion and roost (Fig. 7c). The

estimated bias in the other parameters was minimal (Fig. 7d and 7e). Tag loss probability was

significantly higher in juveniles, but the use of surgical glue substantially reduced this rate (Fig. 8).

Specifically, the tag loss probability decreased by one third from 0.28 (90%hdi [0.23,0.33]) to 0.19

(90%hdi [0.16,0.22]) with the use of surgical glue. In adults, the use of surgical glue did not affect

tag loss rate, with a 69% overlap in the probability distributions. The tag loss rate in adults was

approximately 0.1, which is half the rate observed in juveniles when surgical glue was used. When

considering the period following one year post-tagging, the probability of tag loss was higher when

surgical glue was used (median 0.03, 90%hdi [0.02, 0.04]) compared to when it was not used

(median 0.02, 90%hdi [0.01, 0.02]). However, this difference may be an artefact due to a lack of

search for lost tags on the ground of the maternity roost in the first years of the study

(Supplementary Information 3, part 3 and Fig. S12). Additional parameter estimates can be found in

Supporting Information 3, part 2.6.

4 Discussion

The consideration of mark loss as a violation of CMR model assumptions has been the topic of

numerous studies and model developments. Most of the existing research has focused on estimating

survival, birth, or population size within the framework of Jolly-Seber models (Arnason & Mills,

1981; Malcolm-White *et al.*, 2020; Schwarz *et al.*, 2012; Smout *et al.*, 2011a), recovery models

(Kremers, 1988; Robson & Regier, 1966), CJS models (Laake *et al.*, 2014; Nelson *et al.*, 1980), and

integrated population models (Riecke *et al.*, 2019). However, there have been few proposed

developments to account for mark loss specifically in AS models, such as those suggested by

Besnard *et al.* (2007), Conn *et al.* (2004), and Johnson *et al.* (2016). These approaches have

typically involved implicitly modelling mark status or using an adjustment factor, as seen in the

work of Nishizawa *et al.* (2018). Additionally, we did not find any exploratory studies that have

examined the impact of mark loss on parameter estimates. Therefore, in this study, we aim to fill

this gap by exploring the effects of mark loss and recycled individuals on parameter estimates

through simulations. We also examine the state of the mark (retained or lost) by modelling it as an

independent Bernoulli process. By doing so, we can gain insights into how mark loss can affect the

state transition of individuals when capture and survival probabilities vary over time, survival rates

vary across age classes, and mark loss processes depend on the time since marking. Our results

demonstrate that the violation of CMR model assumptions due to mark loss can significantly impact

not only estimates of survival, but also capture, resighting, and state transition probabilities.

Specifically, we found that survival is generally underestimated in cases where capture and

detection rates are high. This underestimation becomes even more pronounced when survival rates

are also high, thus moderating previous studies that suggested biases mainly occurred in species

with high survival rates, catchability, and mark loss (Diefenbach & Alt, 1998). Our simulation

results confirmed that the inaccuracy of model estimates is positively associated with the rate of

mark loss. However, this inaccuracy can still occur even when the tag loss rate is low (e.g. 5%), and

it can be independent of survival and capture rates. In datasets with a limited number of recycled

individuals, which corresponds to low survival and capture rates, transition and resighting

probabilities can be severely biased if mark loss is high. This implies that caution should be

exercised when interpreting results from studies with low survival and capture rates if mark loss is

suspected but not accounted for. Specifically, the probability of individuals remaining in the same

state is underestimated when the transition from this state is low, but overestimated when the

transition probabilities are high. The severity of these biases can also vary over time, with biases in

survival and recapture decreasing over time as observed in our simulated datasets. This trend is

partly influenced by the chosen mark loss pattern, highlighting the fact that even in studies

conducted over short periods, parameters can still be substantially biased. It should be noted that if

mark loss increases with time since marking (which was not investigated in this study), biases

would be expected to increase accordingly.

The combination of simulation and empirical studies highlights how the complexity and

interdependence of parameters can either exacerbate or mitigate estimation biases when mark loss

modelling is absent. While the simulations revealed some general trends in the biases' direction, the

real-world example demonstrated the unexpected nature of bias patterns. It should also be noted that

our simulation did not encompass the full range of parameter combinations that can be encountered

in nature, despite aiming to cover demographic variations typically observed in vertebrates. This

suggests that most study systems and monitoring methods have their own unique characteristics,

and it is misleading to predict biases without simulating them. The uncertainty propagation in

parameter estimates caused by mark loss remains challenging to predict and grows with system

complexity. Hence, prior to planning a CMR study, we advocate for researchers and managers to

conduct simulations to evaluate the conditions (i.e., parameter combinations) under which their

study would yield reliable estimates of the parameters of interest (e.g., demographic, state

transition). Preliminary studies with multiple marks could also be considered whenever possible

(Smout *et al.*, 2011a). This approach would allow for the optimization of CMR study design before

its implementation, thereby minimizing biases from the outset.

Previous studies have developed AS models to estimate movement between sites, recruitment,

dispersal, and temporary or permanent emigration (Lebreton *et al.*, 2003, 2009; Schaub *et al.*,

2004). Our simulation results indicate that state transition probabilities are sensitive to mark loss,

even at low rates. For example, the probabilities of staying in the same state (philopatry, if

transitions involve movement) or changing state (e.g., emigration) exhibited both underestimations

and overestimations. These parameters are often essential in addressing ecological and demographic

questions and used for management and conservation purposes (Cam *et al.*, 2004; Horton *et al.*,

2011). Although mark loss is frequently reported for various tag types and taxa, it is only marginally

considered in studies focusing on estimating population dynamics parameters (Nelson *et al.*, 1980;

Ostrand *et al.*, 2012; Smout *et al.*, 2011b; Tavecchia *et al.*, 2012). Most model developments to

account for mark loss have focused on the JS model for abundance estimation, as mark loss and

recycling in this model can introduce significant biases (Malcolm-White *et al.*, 2020). Despite the

growing use of AS models in ecology, demography, management, and conservation (Huntsman *et*

al., 2020; Melnychuk *et al.*, 2017), the issue of mark loss remains largely overlooked. Based on our

study, we recommend the use of permanent or double temporary marks, ideally independent in

terms of loss, or considering dependence in loss (Laake *et al.*, 2014; McMahon & White, 2009;

Schwarz *et al.*, 2012). This is because any analysis of CMR data is potentially affected by this

violation of model assumptions (Riecke *et al.*, 2019).

Despite the fact that PIT tags are suitable for an increasing number of studies and allow data

collection without physically re-capturing individuals, our case study emphasizes the importance of

a second marking method to avoid potential bias in demographic rate estimations. Tag loss has long

been known to occur in small mammal species, especially those that fly or glide (Freeland & Fry,

1995). In our study, we have confirmed that the use of surgical adhesive can reduce PIT tag

shedding in the short term (Lebl & Ruf, 2010; van Harten *et al.*, 2020).

However, our findings show that surgical adhesive alone is not enough to eliminate tag loss, and the

use of additional data (e.g., evidence of tag loss) or a permanent mark (e.g., genotype) is necessary

for the entire or a part of the studied population (Laake *et al.*, 2014). Similar situations, where

permanent marks should be considered, arise when marks deteriorate and become unreadable over

time, such as with neck collars or ear tags (Conn *et al.*, 2004; Diefenbach & Alt, 1998). In such

cases, the accuracy of model parameter estimates is expected to decrease throughout the study

duration, further supporting the use of permanent marks in capture-mark-recapture (CMR) studies.

Mark loss is typically not taken into account from ecological and management perspectives, except

when researchers aim to understand factors influencing mark failures or improve their marking

methods. Our results emphasize the need to assess the impact of mark loss whenever mark failure is

suspected to avoid drawing misleading conclusions about the dynamics of the studied species.

Based on our experience and the literature, PIT tags are prone to being shed regardless of the taxa

being studied, often in the short term but sometimes even in the long term. A recent study on

Gould's wattled bats (*Chalinolobus gouldii*) over a relatively short period of 13-14 months reported

low tag shedding (2.7%) and generalized this finding to all insectivorous bats (van Harten *et al.*,

2020). However, our study shows that this generalization is partly incorrect and suggests that it is

challenging to generalize such conclusions, as the pattern of mark loss is highly species-dependent,

among other factors. Therefore, mark loss should be carefully considered in all CMR analyses and

potentially in other studies using similar datasets. Explicit modelling of mark loss should be

implemented when necessary to achieve more accurate estimations of population dynamics.

Acknowledgements

This project was funded by an Irish Research Council Postdoctoral Fellowship Grant

GOIPD/2018/256 awarded to F.T., a European Research Council Research Grant ERC-2012-

StG311000 and an Irish Research Council Laureate Award awarded to E.C.T. The French field

study was supported by the European Regional Development Fund EU000141 and a Contrat Nature

grant awarded to Bretagne Vivante (BV). S.J.P. was supported by a Junior chair from the Institut

Universitaire de France. We would like to thank the volunteers of BV for assistance in the

fieldwork. Sampling was carried out in accordance with the French ethical and sampling guidelines

issued in 3 successive permits delivered by the Préfet du Morbihan (Brittany) awarded to E.J.P., F.T.

and S.J.P. for the period 2010–2020.

Author contribution

F.T. designed the project with the other co-authors. E.C.T., F.T. acquired the funding. E.C.T., E.J.P.,

S.J.P. and E.C. supervised the project. E.C.T., F.T., E.J.P., and S.J.P supervised the fieldwork.

E.C.T., E.J.P., F.T. and S.J.P. collected samples. S.J.P. & F.T. supervised the microsatelite

genotyping. C.S. generated the genotypes. F.T. and E.C. developed the R scripts for simulating and

analysing data with help of E.J.P. and S.J.P. F.T. led the writing of the manuscript and all authors

contributed to manuscript revisions and gave final approval for publication.

542 **Data accessibility**

543 R scripts for simulating the data, and analysing the data with JAGS, are accessible on Zenodo

544 (https://zenodo.org/record/6453953).

546 **References**

Acolas, M. L., Roussel, J. M., Lebel, J. M., & Bagliniere, J. L. (2007). Laboratory experiment on survival, growth and tag retention following PIT injection into the body cavity of juvenile brown trout (*Salmo trutta*). *Fisheries Research*, *86*(2–3), 280–284. https://doi.org/10.1016/j.fishres.2007.05.011

Arnason, A. N. (1972). Parameter estimates from mark-recapture experiments on two populations subject to migration and death. *Population Ecology*, *13*(2), 97–113. https://doi.org/10.1007/BF02521971

Arnason, A. N. (1973). The estimation of population size, migration rates and survival in a stratified population. *Researches on Population Ecology*, *15*(2), 1–8. https://doi.org/10.1007/BF02510705

Arnason, A. N., & Mills, K. H. (1981). Bias and Loss of Precision Due to Tag Loss in Jolly–Seber Estimates for Mark–Recapture Experiments. *Canadian Journal of Fisheries and Aquatic Sciences*, *38*(9), 1077–1095. https://doi.org/10.1139/f81-148

Benkman, C. W., Colquitt, J. S., Gould, W. R., Fetz, T., Keenan, P. C., & Santisteban, L. (2005). Can Selection by an Ectoparasite Drive a Population of Red Crossbills from Its Adaptive Peak? *Evolution*, *59*(9), 2025–2032. https://doi.org/10.1111/j.0014-3820.2005.tb01071.x

Besnard, A., Piry, S., Berthier, K., Lebreton, J.-D., & Streiff, R. (2007). Modeling survival and mark loss in molting animals: Recapture, dead recoveries, and exuvia recoveries. *Ecology*, *88*(2), 289–295. https://doi.org/10.1890/05-0811

Brooks, S., & Gelman, A. (1998). General methods for monitoring convergence of iterative simulations. *Journal of Computational and Graphical Statistics*, *7*(4), 434–455. https://doi.org/10.2307/1390675

Cam, E. (2009). Contribution of Capture-Mark-Recapture Modeling to Studies of Evolution by Natural Selection. In D. L. Thomson, E. G. Cooch, & M. J. Conroy (Eds.), *Modeling*

Demographic Processes In Marked Populations (pp. 83–129). *Springer US*.

https://doi.org/10.1007/978-0-387-78151-8_5

Cam, E., Monnat, J.-Y. M., & Royle, A. (2004). Dispersal and individual quality in a long lived

species. *Oikos, 106*(2), 386–398. https://doi.org/10.1111/j.0030-1299.2003.13097.x

Choquet, R., Lebreton, J. D., Gimenez, O., Reboulet, A. M., & Pradel, R. (2009a). U-CARE:

Utilities for performing goodness of fit tests and manipulating CApture-REcapture data.

Ecography, 32(6), 1071–1074.

Choquet, R., Rouan, L., & Pradel, R. (2009b). Program E-Surge: A Software Application for Fitting

Multievent Models. In D. L. Thomson, E. G. Cooch, & M. J. Conroy (Eds.), *Modeling*

Demographic Processes In Marked Populations (Vol. 3, pp. 845–865). Springer US.

https://doi.org/10.1007/978-0-387-78151-8_39

Conde, D. A., Staerk, J., Colchero, F., da Silva, R., Schöley, J., Baden, H. M., Jouvet, L., Fa, J. E.,

Syed, H., Jongejans, E., Meiri, S., Gaillard, J.-M., Chamberlain, S., Wilcken, J., Jones, O.

R., Dahlgren, J. P., Steiner, U. K., Bland, L. M., Gomez-Mestre, I., … Vaupel, J. W. (2019).

Data gaps and opportunities for comparative and conservation biology. *Proceedings of the*

National Academy of Sciences, 116(19), 9658–9664.

https://doi.org/10.1073/pnas.1816367116

Conn, P. B., Kendall, W. L., & Samuel, M. D. (2004). A General Model for the Analysis of Mark-

Resight, Mark-Recapture, and Band-Recovery Data under Tag Loss. *Biometrics, 60*(4), 900–

909. https://doi.org/10.1111/j.0006-341X.2004.00245.x

Cormack, R. (1964). Estimates of Survival from Sighting of Marked Animals. *Biometrika, 51*(3–4),

429–438. https://doi.org/10.2307/2334149

Cowen, L., & Schwarz, C. J. (2006). The Jolly–Seber Model with Tag Loss. *Biometrics, 62*(3), 699–

705. https://doi.org/10.1111/j.1541-0420.2006.00523.x

de Valpine, P., Turek, D., Paciorek, C. J., Anderson-Bergman, C., Lang, D. T., & Bodik, R. (2017).

Programming With Models: Writing Statistical Algorithms for General Model Structures

With NIMBLE. *Journal of Computational and Graphical Statistics, 26*(2), 403–413.

https://doi.org/10.1080/10618600.2016.1172487

Diefenbach, D. R., & Alt, G. L. (1998). Modeling and Evaluation of Ear Tag Loss in Black Bears.

The Journal of Wildlife Management, 62(4), 1292–1300. https://doi.org/10.2307/3801993

Fabrizio, M. C., Nichols, J. D., Hines, J., Swanson, B. L., & Schram, S. T. (1999). Modeling data from double-tagging experiments to estimate heterogeneous rates of tag shedding in lake trout (*Salvelinus namaycush*). *Canadian Journal of Fisheries and Aquatic Sciences*, *56*(8), 1409–1419. https://doi.org/10.1139/F99-069

Fokidis, H. B., Robertson, C., & Risch, T. S. (2006). Keeping tabs: Are redundant marking systems needed for rodents? *Wildlife Society Bulletin*, *34*(3), 764–771. https://doi.org/10.2193/0091-7648(2006)34[764:KTARMS]2.0.CO;2

Foley, N. M., Petit, E. J., Brazier, T., Finarelli, J. A., Hughes, G. M., Touzalin, F., Puechmaille, S. J., & Teeling, E. C. (2020). Drivers of longitudinal telomere dynamics in a long-lived bat species, *Myotis myotis*. *Molecular Ecology*, *29*(16), 2963–2977. https://doi.org/10.1111/mec.15395

Frantz, A. C., Pope, L. C., Carpenter, P. J., Roper, T. J., Wilson, G. J., Delahay, R. J., & Burke, T. (2003). Reliable microsatellite genotyping of the Eurasian badger (*Meles meles*) using faecal DNA. *Molecular Ecology*, *12*(6), 1649–1661. https://doi.org/10.1046/j.1365-294x.2003.01848.x

Freeland, W. J., & Fry, K. (1995). Suitability of passive integrated transponder tags for marking live animals for trade. *Wildlife Research*, *22*(6), 767–773. https://doi.org/10.1071/WR9950767

Gimenez, O., Cam, E., & Gaillard, J.-M. (2018a). Individual heterogeneity and capture–recapture models: What, why and how? *Oikos*, *127*(5), 664–686. https://doi.org/10.1111/oik.04532

Gimenez, O., Lebreton, J.-D., Choquet, R., & Pradel, R. (2018b). R2ucare: An r package to perform goodness-of-fit tests for capture–recapture models. *Methods in Ecology and Evolution*, *9*(7), 1749–1754. https://doi.org/10.1111/2041-210X.13014

Goudie, R. J. B., Turner, R. M., De Angelis, D., & Thomas, A. (2020). MultiBUGS: A Parallel Implementation of the BUGS Modelling Framework for Faster Bayesian Inference. *Journal of Statistical Software*, *95*, 7. https://doi.org/10.18637/jss.v095.i07

Hestbeck, J. B., Nichols, J. D., & Malecki, R. A. (1991). Estimates of Movement and Site Fidelity Using Mark-Resight Data of Wintering Canada Geese. *Ecology*, *72*(2), 523–533. https://doi.org/10.2307/2937193

Horton, G. E., Letcher, B. H., & Kendall, W. L. (2011). A Multistate Capture–Recapture Modeling Strategy to Separate True Survival from Permanent Emigration for a Passive Integrated

Transponder Tagged Population of Stream Fish. *Transactions of the American Fisheries Society*, *140*(2), 320–333. https://doi.org/10.1080/00028487.2011.567861

Huntsman, B. M., Venarsky, M. P., Abadi, F., Huryn, A. D., Kuhajda, B. R., Cox, C. L., & Benstead, J. P. (2020). Evolutionary history and sex are significant drivers of crayfish demography in resource-limited cave ecosystems. *Evolutionary Ecology*, *34*(2), 235–255. https://doi.org/10.1007/s10682-019-10029-w

Jennelle, C. S., Cooch, E. G., Conroy, M. J., & Senar, J. C. (2007). State-specific detection probabilities and disease prevalence. *Ecological Applications*, *17*(1), 154–167. https://doi.org/10.1890/1051-0761(2007)017[0154:SDPADP]2.0.CO;2

Johnson, D. H., Burnham, K. P., & Nichols, J. D. (1986). The role of heterogeneity in animal population dynamics. *Proceedings of the International Biometrics Conference, 13,* 5.3.1-5.3.15.

Johnson, D. S., Laake, J. L., Melin, S. R., & DeLong, R. L. (2016). Multivariate State Hidden Markov Models for Mark-Recapture Data. *Statistical Science*, *31*(2), 233–244. https://doi.org/10.1214/15-STS542

Jolly, G. (1965). Explicit Estimates from Capture-Recapture Data with Both Death and Immigration-Stochastic Model. *Biometrika*, *52*, 225–247. https://doi.org/10.2307/2333826

Juillet, C., Choquet, R., Gauthier, G., & Pradel, R. (2011). A Capture–Recapture Model with Double-Marking, Live and Dead Encounters, and Heterogeneity of Reporting Due to Auxiliary Mark Loss. *Journal of Agricultural, Biological, and Environmental Statistics*, *16*(1), 88–104. https://doi.org/10.1007/s13253-010-0035-5

Kaemingk, M. A., Weber, M. J., McKenna, P. R., & Brown, M. L. (2011). Effect of Passive Integrated Transponder Tag Implantation Site on Tag Retention, Growth, and Survival of Two Sizes of Juvenile Bluegills and Yellow Perch. *North American Journal of Fisheries Management*, *31*(4), 726–732. https://doi.org/10.1080/02755947.2011.611863

Kellner, K. (2016). *JagsUI: A Wrapper Around "rjags" to Streamline "JAGS" Analyses*. https://CRAN.R-project.org/package=jagsUI

Kremers, W. K. (1988). Estimation of Survival Rates from a Mark-Recapture Study with Tag Loss. *Biometrics*, *44*(1), 117–130. JSTOR. https://doi.org/10.2307/2531900

Kruschke, J. (2014). Doing Bayesian Data Analysis: A Tutorial with R, JAGS, and Stan (2 edition). *Academic Press*.

Kruschke, J. K. (2018). Rejecting or Accepting Parameter Values in Bayesian Estimation. *Advances in Methods and Practices in Psychological Science, 1*(2), 270–280. https://doi.org/10.1177/2515245918771304

Laake, J. L., Johnson, D. S., Diefenbach, D. R., & Ternent, M. A. (2014). Hidden Markov Model for Dependent Mark Loss and Survival Estimation. *Journal of Agricultural Biological and Environmental Statistics, 19*(4), 524–540. https://doi.org/10.1007/s13253-014-0190-1

Lebl, K., & Ruf, T. (2010). An easy way to reduce PIT-tag loss in rodents. *Ecological Research, 25*(1), 251–253. https://doi.org/10.1007/s11284-009-0629-y

Lebreton, J. D., Burnham, K. P., Clobert, J., & Anderson, D. R. (1992). Modeling Survival and Testing Biological Hypotheses Using Marked Animals—A Unified Approach with Case-Studies. *Ecological Monographs, 62*(1), 67–118.

Lebreton, J.-D., Hines, J. E., Pradel, R., Nichols, J. D., & Spendelow, J. A. (2003). Estimation by capture-recapture of recruitment and dispersal over several sites. *Oikos, 101*(2), 253–264. https://doi.org/10.1034/j.1600-0706.2003.11848.x

Lebreton, J.-D., Nichols, J. D., Barker, R. J., Pradel, R., & Spendelow, J. A. (2009). Chapter 3 Modeling Individual Animal Histories with Multistate Capture–Recapture Models. In Hal Caswell (Ed.), *Advances in Ecological Research: Vol. Volume 41* (pp. 87–173). Academic Press. http://www.sciencedirect.com/science/article/pii/S0065250409004036

Ling, H., & Okada, K. (2007). An Efficient Earth Mover's Distance Algorithm for Robust Histogram Comparison. *IEEE Transactions on Pattern Analysis and Machine Intelligence, 29*(5), 840–853. https://doi.org/10.1109/TPAMI.2007.1058

Lunn, D. J., Thomas, A., Best, N., & Spiegelhalter, D. (2000). WinBUGS - A Bayesian modelling framework: Concepts, structure, and extensibility. *Statistics and Computing, 10*(4), 325–337. https://doi.org/10.1023/A:1008929526011

Malcolm-White, E., McMahon, C. R., & Cowen, L. L. E. (2020). Complete tag loss in capture–recapture studies affects abundance estimates: An elephant seal case study. *Ecology and Evolution, 10*(5), 2377–2384. https://doi.org/10.1002/ece3.6052

McMahon, C. R., & White, G. C. (2009). Tag loss probabilities are not independent: Assessing and quantifying the assumption of independent tag transition probabilities from direct observations. *Journal of Experimental Marine Biology and Ecology*, *372*(1–2), 36–42. https://doi.org/10.1016/j.jembe.2009.02.006

Melnychuk, M. C., Dunton, K. J., Jordaan, A., McKown, K. A., & Frisk, M. G. (2017). Informing conservation strategies for the endangered Atlantic sturgeon using acoustic telemetry and multi-state mark-recapture models. *Journal of Applied Ecology*, *54*(3), 914–925. https://doi.org/10.1111/1365-2664.12799

Nelson, L. J., Anderson, D. R., & Burnham, K. P. (1980). The effect of band loss on estimates of annual survival. In *Journal of Field Ornithology* (Vol. 51, Issue 1, p. 9).

Nichols, J. D., Bart, J., Limpert, R. J., Sladen, W. J. L., & Hines, J. E. (1992). Annual Survival Rates of Adult and Immature Eastern Population Tundra Swans. *The Journal of Wildlife Management*, *56*(3), 485–494. https://doi.org/10.2307/3808863

Nichols, J. D., & Hines, J. E. (1993). Survival rate estimation in the presence of tag loss using joint analysis of capture-recapture and resighting data. In *Marked Individuals in the Study of Bird Population* (Birkhauser Verlag, pp. 229–243). https://pubs.er.usgs.gov/publication/5210628

Nichols, J. D., & Kendall, W. L. (1995). The use of multi-state capture-recapture models to address questions in evolutionary ecology. *Journal of Applied Statistics*, *22*(5–6), 835–846. https://doi.org/10.1080/02664769524658

Nishizawa, H., Joseph, J., Chew, V. Y.-C., Liew, H.-C., & Chan, E.-H. (2018). Assessing tag loss and survival probabilities in green turtles (*Chelonia mydas*) nesting in Malaysia. *Journal of the Marine Biological Association of the United Kingdom*, *98*(4), 961–972. https://doi.org/10.1017/S0025315417000224

Ostrand, K. G., Zydlewski, G. B., Gale, W. L., & Zydlewski, J. D. (2012). Long Term Retention, Survival, Growth, and Physiological Indicators of Juvenile Salmonids Marked with Passive Integrated Transponder Tags. In J. R. McKenzie, B. Parsons, A. C. Seitz, R. K. Kopf, M. G. Mesa, & Q. Phelps (Eds.), *Advances in Fish Tagging and Marking Technology* (Vol. 76, pp. 135–145). Amer Fisheries Soc.

Plummer, M. (2003). JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling. *3rd International Workshop on Distributed Statistical Computing (DSC 2003)*, 1–10.

Puechmaille, S. J., & Petit, E. J. (2007). Empirical evaluation of non-invasive capture–mark–recapture estimation of population size based on a single sampling session. *Journal of Applied Ecology*, *44*(4), 843–852. https://doi.org/10.1111/j.1365-2664.2007.01321.x

R Core Team. (2019). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. https://www.R-project.org/

Riecke, T. V., Williams, P. J., Behnke, T. L., Gibson, D., Leach, A. G., Sedinger, B. S., Street, P. A., & Sedinger, J. S. (2019). Integrated population models: Model assumptions and inference. *Methods in Ecology and Evolution*, *10*(7), 1072–1082. https://doi.org/10.1111/2041-210X.13195

Robson, D. S., & Regier, H. A. (1966). Estimates of Tag Loss from Recoveries of Fish Tagged and Permanently Marked. *Transactions of the American Fisheries Society*, *95*(1), 56–59. https://doi.org/10.1577/1548-8659(1966)95[56:EOTLFR]2.0.CO;2

Schaub, M., Gimenez, O., Schmidt, B. R., & Pradel, R. (2004). Estimating survival and temporary emigration in the multistate capture-recapture framework. *Ecology*, *85*(8), 2107–2113.

Schofield, M. R., & Barker, R. J. (2008). A Unified Capture-Recapture Framework. *Journal of Agricultural Biological and Environmental Statistics*, *13*(4), 458–477. https://doi.org/10.1198/108571108X383465

Schwarz, C. J., Schweigert, J. F., & Arnason, A. N. (1993). Estimating Migration Rates Using Tag-Recovery Data. *Biometrics*, *49*(1), 177–193. https://doi.org/10.2307/2532612

Schwarz, L. K., Hindell, M. A., McMahon, C. R., & Costa, D. P. (2012). The implications of assuming independent tag loss in southern elephant seals. *Ecosphere*, *3*(9), art81. https://doi.org/10.1890/ES12-00132.1

Seber, G. (1965). A Note on Multiple-Recapture Census. *Biometrika*, *52*, 249–260.

Seber, G. A. F., & Felton, R. (1981). Tag loss and the Petersen mark-recapture experiment. *Biometrika*, *68*(1), 211–219. https://doi.org/10.1093/biomet/68.1.211

Seber, G. A. F., & Schofield, M. R. (2019). Tagging Methods and Tag Loss. In G. A. F. Seber & M. R. Schofield (Eds.), Capture-Recapture: Parameter Estimation for Open Animal Populations

(pp. 13–37). *Springer International Publishing*. https://doi.org/10.1007/978-3-030-18187-1_2

Smout, S., King, R., & Pomeroy, P. (2011a). Estimating demographic parameters for capture-recapture data in the presence of multiple mark types. *Environmental and Ecological Statistics*, *18*(2), 331–347. https://doi.org/10.1007/s10651-010-0135-y

Smout, S., King, R., & Pomeroy, P. (2011b). Integrating heterogeneity of detection and mark loss to estimate survival and transience in UK grey seal colonies. *Journal of Applied Ecology*, *48*(2), 364–372. https://doi.org/10.1111/j.1365-2664.2010.01913.x

Spendelow, J. A., Nichols, J. D., Nisbet, I. C. T., Hays, H., Cormons, G. D., Burger, J., Safina, C., Hines, J. E., & Gochfeld, M. (1995). Estimating Annual Survival and Movement Rates of Adults within a Metapopulation of Roseate Terns. *Ecology*, *76*(8), 2415–2428. https://doi.org/10.2307/2265817

STAN Development Team. (2022). Stan Modeling Language Users Guide and Reference Manual, 2.31. http://mc-stan.org/

Tavecchia, G., Adrover, J., Navarro, A. M., & Pradel, R. (2012). Modelling mortality causes in longitudinal data in the presence of tag loss: Application to raptor poisoning and electrocution. *Journal of Applied Ecology*, *49*(1), 297–305. https://doi.org/10.1111/j.1365-2664.2011.02074.x

van Harten, E., Reardon, T., Holz, P. H., Lawrence, R., Prowse, T. A. A., & Lumsden, L. F. (2020). Recovery of southern bent-winged bats (*Miniopterus orianae bassanii*) after PIT-tagging and the use of surgical adhesive. *Australian Mammalogy*, *42*(2), 216–219. https://doi.org/10.1071/AM19024

White, G., & Burnham, K. (1999). Program MARK: survival estimation from populations of marked animals. *Bird Study*, *46*(S), 120–139.

Williams, B. K., Nichols, J. D., & Conroy, M. J. (2002). Analysis and Management of Animal Populations (1 edition). *Academic Press*.

Winiarski, K. J., & McGarigal, K. (2016). Effects of Photo and Genotype-Based Misidentification Error on Estimates of Survival, Detection and State Transition using Multistate Survival Models. *PLOS ONE*, *11*(1), e0145640. https://doi.org/10.1371/journal.pone.0145640

Figure 1: Density of probability of survival across age class and taxa for 700 species. Sample size are indicated by labels, with colour corresponding to the state of individual sampled.

Scenario	Survival	Detection
Scenario 1	Long-lived (high survival)	High detection rate (capture & resighting)
Scenario 2	Long-lived (high survival)	Low detection rate (capture & resighting)
Scenario 3	Short-lived (low survival)	Low detection rate (capture & resighting)
Scenario 4	Short-lived (low survival)	High detection rate (capture & resighting)

Table 1: Summary of the characteristics of each simulated scenario.

Parameter	Definition	Value	
		Long-lived species	Short-lived species
$\phi_{ad.}$	Adult survival in state "A", "B", "C"	logit(N(2.5,0.3)) [0.92]	logit(N(2.5,0.3)-0.3) [0.62]
$\phi_{ad.}$	Adult survival in state "D"	logit(N(1.5,0.3)) [0.81]	logit(N(1.5,0.3)-0.3) [0.51]
$\phi_{juv.}$	Juvenile survival in state "A", "B", "C"	logit(N(0.2,0.3)) [0.55]	logit(N(0.2,0.3)-0.2) [0.35]
$\phi_{juv.}$	Juvenile survival in state "D"	NA	NA
		High	Low
pc_A	Capture probability in state "A"	U(0.6,0.7) [0.65]	U(0.6,0.7)-0.5 [0.15]
pc_B	Capture probability in state "B"	U(0.7,0.8) [0.75]	U(0.7,0.8)-0.5 [0.25]
pc_C	Capture probability in state "C"	U(0.65,0.75) [0.7]	U(0.65,0.75)-0.5 [0.2]
pc_D	Capture probability in state "D"	0	0
$p.r_A$	Resighting probability in state "A"	0.85	0.35
$p.r_B$	Resighting probability in state "B"	0.95	0.45
$p.r_C$	Resighting probability in state "C"	0.9	0.4
$p.r_D$	Resighting probability in state "D"	0.7	0.2

Table 2: Parameter values used to simulate the 4 scenarios. For random values generated, the corresponding distribution is indicated with N (a, b) the normal distribution with mean a and variance b, and U(a,b) the uniform distribution with lower bound a and upper bound b. The square brackets show mean values on the probability scale. To simulate survival for short-lived species, we used the same distribution on as long-lived species but subtract generated values by 0.3 for adults and 0.2 for juveniles. In the same way, we obtained the low values of probability of capture and resighting by subtracting 0.5 from the high values. The probability of capture in state "D" is set to 0, as no capture is possible when individuals are in this state. For the transition values between states see Fig. 3.

Figure 2: *The possible fates of an individual between occasion t and t+1. We first consider the state transition process: if the individual dies between occasions, it can only remain dead and can no longer be detected, both with probability 1 (absorbing state). The individual can, however, survive between occasions with probability φ (depending on time and state at t) and can then change state with probability ψ (see Fig. 1). Second, the mark retention process: if it survives between occasions, then the individual can retain its mark with probability pr (depending on age and time since marking) or lose its mark with probability pl = 1-pr. Thirdly, the detection process: if this individual has lost his mark between occasions, he may possibly be recaptured with a probability pc (depending on time and state at t+1) and if this happens, he is marked again before being released. However, if the mark is retained, other events may occur: the individual may also be captured with probability pc, but it may also be resighted with probability p.r (depending on the state at t+1), or both with probability pc*p.r, or finally not be detected at all with probability 1-pd = (1-pc)*(1-p.r).*

Figure 3: Schematic description of parameter values used to simulate data under scenario 2 (long-lived species with low detection). Central graph: solid arrows correspond to possible transitions of females and dashed arrows to those of males. The size of the arrows is proportional to the probability of transition indicated next to them, all were kept constant over time. Peripheral graph: simulated survival (Surv.) and detection (Det.) probabilities were displayed for states "A", "B" and "C". The light blue lines correspond to adult survival, the orange lines to juvenile survival and the green lines to the probability of capture, which are derived from Normal and Uniform distribution and therefore fluctuate over the years (see Table 1). The grey lines correspond to the probability of resighting, they differed between state but were set constant in time.

Figure 4: Comparison of bias for estimates of juvenile (a) and adult (c) survival, transition (b and d), capture (e) and resighting (f) probabilities between model accounting for mark loss (AML) or not (WML). All violin plots show the distribution of bias over 50 simulations from scenario 1 (long-lived species and high detection probabilities), with a simulated probability of mark loss of 0.4. The median of each simulated distribution is shown with a horizontal line. The numbers 1 to 9 are the recapture opportunities, the letters from A to D represent the different states, AdtF the adult females, AdtM the adult males, JuvF the juvenile females and JuvM the juvenile males.

Figure 5: Tile-plots of the proportion of the distribution of the Earth Mover Distance (across 50 simulated datasets) out of the Region of Practical Equivalence (ROPE), between the model accounting for tag loss and recycling (ModelA) and the model ignoring them (ModelW). The ROPE corresponds to the interval including 80% hdi of the posterior density distribution of the "true value" of a parameter which was estimated with ModelA. Each tile represents annual (right axis) juvenile survival (a), adult survival (b) and capture probability (c) for each scenario (y axis) and tag loss probabilities (x axis). The scenarios indicated at the bottom are: (1) long-lived species and high detection rate; (2) long-lived species and low detection rate; (3) short-lived species and low detection rate; (4) short-lived species and high detection rate. At the top of each panel, A, B, C and D correspond to the states.

Figure 6: Tile-plots of the proportion of the distribution of the Earth Mover Distance (across 50 simulated datasets) out of the Region of Practical Equivalence (ROPE), between model accounting for tag loss and recycling (ModelA) and the model ignoring them (ModelW) for each simulated cases. The ROPE corresponds to the interval including 80% hdi of the posterior density distribution of the "true value" of a parameter which was estimated with ModelA. Each tile represents resighting probability (a) and transition probabilities between subpopulations (direction, "from-to", are indicated above each tile-plot, e.g. "A-B" correspond to state transition from A to B) of juvenile female (b), adult female (c), juvenile male (d) and adult male (e) for each scenario and tag loss probabilities. The scenario are indicated at the bottom: (1) long-lived species and high detection rate; (2) long-lived species and low detection rate; (3) short-lived species and low detection rate; (4) short-lived species and high detection rate. At the top of each panel, A, B, C and D correspond to the states.

Figure 7: Differences in the medians of the posterior distributions of juvenile (a) and adult (b) survival, capture (c) resighting (d) and transition (e) probabilities between the model accounting for tag-loss and the model ignoring it, both estimated from the empirical data. Colonies are abbreviated: Beg = Beganne; Fer = Férel; LIM = Limerzel; LRB = La Roche Bernard; NM = Noyal-Muzillac. Movements between sites are indicated on x axis with direction "from-to". Movements (e) are specified by age (Adt. = Adulte, Juv. = juvenile) and sex classes (M = male, F = Female), indicated on the right side of the plot (e).

Figure 8: Posterior distribution of the tag loss probabilities according to age classes and time after marking in the Myotis myotis *dataset. Left panel corresponds to tag shedding rate during the year following the tag injection and the right panel for the following years (constant in time). In blue, distribution if surgical adhesive was used after tag injection and in red, without surgical adhesive.*