Topographic axonal projection at single-cell precision supports local retinotopy in the mouse superior colliculus

Authors: Dmitry Molotkov[1], Leiron Ferrarese[1], Tom Boissonnet[1,2,3], Hiroki Asari*[1]

Affiliations:

1. Epigenetics and Neurobiology Unit, EMBL Rome, European Molecular Biology Laboratory, Monterotondo, 00015, Italy

2. Collaboration for joint PhD degree between EMBL and Université Grenoble Alpes, Grenoble Institut des Neurosciences, La Tronche, 38700, France

3. Current address: Center for Advanced Imaging, Heinrich-Heine-Universität Düsseldorf, Düsseldorf, 40225, Germany

*Corresponding author: Hiroki Asari, E-mail: asari@embl.it

Acknowledgments: This work was supported by research grants from EMBL (H.A.). The EMBL Genetic and Viral Engineering Facility is acknowledged for virus production; EMBL IT Support for provision of computer and data storage servers; and the LAR facility for taking care of animals. We thank Lin Tian (University of California Davis) for providing axon-targeted GCaMP construct; Thomas Euler and Luke Rogerson (University of Tübingen) for sharing Python scripts to generate random water-wave stimuli, originally from Andreas Tolias lab (Baylor College of Medicine); Simone Calabrese, Ilaria Sauve, Grace Cunliffe, and Matteo Tripodi for supporting experiments; and all the Asari lab members as well as Santiago Rompani for many useful discussions.

Author contributions: D.M. and H.A. designed the study; D.M. and L.F performed experiments; D.M, T.B., L.F. and H.A. analyzed the results; and D.M., L.F. and H.A. wrote the manuscript.

Competing interests: The authors declare no competing financial interests.

Abbreviated title: retinotopic RGC axon tiling in SC.

Key words: retinal ganglion cells; retinotopy; axonal imaging; superior colliculus; mouse.

Acronyms: 2D, two-dimensional; AAV, adeno-associated virus; CNMF, constrained non-negative matrix factorization; RF, receptive field; RGC, retinal ganglion cells; SC, superior colliculus; SD, standard deviation.

1 **Abstract**

Retinotopy, like all long-range projections, can arise from the axons themselves or their targets,

but the underlying connectivity pattern remains unknown at the fine scale. To address this

question, we functionally mapped the spatial organization of the input axons and target

neurons in the mouse retinocollicular pathway at single-cell resolution using *in vivo* two-photon

calcium imaging. We found a near-perfect retinotopic tiling of retinal ganglion cell axon

terminals, with an average error below 30 µm or 2 degrees of visual angle. The precision of

retinotopy was relatively lower for local neurons in the superior colliculus. Subsequent data-

driven modelling ascribed it to a low input convergence, on average 5.5 retinal ganglion cell

inputs to a postsynaptic cell in the superior colliculus. These results indicate that retinotopy

arises largely from topographically precise input from presynaptic cells, rather than elaborating

local circuitry to reconstruct the topography by postsynaptic cells.

13 **Introduction**

Topographic organization is central to brain function (Jbabdi et al., 2013; Patel et al., 2014).

Connections between brain regions are often spatially arranged to have one-to-one mappings,

and sensory processing relies on the transmission of topographically preserved information

from receptor organs. In the visual system, for example, topographic visual representations

are first formed in the retina and conveyed to the brain by retinal ganglion cells (RGCs) whose

axons are bundled into an optic nerve (Masland, 2012). In general, neighboring RGCs project

to neighboring regions in their targets (Frisén et al., 1998; McLaughlin et al., 2003; Cang and

Feldheim, 2013; Arroyo and Feller, 2016). This forms a retinotopic map in the primary

retinorecipient areas, such as the lateral geniculate nucleus (Malpeli and Baker, 1975) and

the superior colliculus (SC; Dräger and Hubel, 1976; Cang et al., 2018), and retinotopy is

likewise transferred throughout the entire visual pathways (Wandell et al., 2007; Cang and

Feldheim, 2013).

How precisely is topographic information transmitted between brain regions? Only

global-level relationship is known even for the best characterized cases, such as the retinal

projection to SC (Frisén et al., 1998; McLaughlin et al., 2003; Cang and Feldheim, 2013;

Arroyo and Feller, 2016). During development, RGC axons reach their target locations based

on molecular guidance cues and activity-dependent refinement. Long-range projections of

dense axonal fibers, however, preclude a precise anatomical characterization of the

connectivity patterns at single-cell resolution (Hong et al., 2011). In the optic nerve, RGC

axons are mixed and lose retinotopic organizations (Horton et al., 1979; Colello and Guillery,

1998). A fundamental question is then if RGC axons can nevertheless find precise targets

even after they get "lost" in the long-range projection (Figure 1, Model 1), or if target neurons

need to reconstruct the topography by elaborating local circuitry (Figure 1, Model 2). A recent

study using a high-density electrode has implicated retinotopic RGC projections to the mouse

SC along the probe shank (Sibille et al., 2022), favoring the former scenario. However, the

precision of retinotopy at the level of individual axons and its relationship to that of target

neurons remain elusive in any retinorecipient area.

Figure 1: Possible retinocollicular projection models for topographic information transmission. Retinotopy in SC (color-coded) can arise from either precisely retinotopic projection of RGC axons, innervating exact target locations in SC (Model 1); or roughly retinotopic projection of RGC axons while SC neurons identify appropriate partners to recover the retinotopy (Model 2). In Model 1, retinotopy can be more precise for RGC axons than for SC somata if the synaptic connectivity is not topographically well organized. In Model 2, in contrast, SC somata should show more precise retinotopy than RGC axons. Note a loss of retinotopic organization in the optic nerve (Horton et al., 1979; Colello and Guillery, 1998).

To address this question, we performed functional mapping of RGC axon terminals

and local neurons in SC at single-cell resolution in awake head-fixed mice using two-photon

calcium imaging. Subsequent analyses on their spatial organizations allowed us to quantify

and directly compare the precision of retinotopy between the pre- and post-synaptic sides.

Moreover, these experimental data served as a basis of a computational modelling analysis

for inferring key features on the topographic information transmission in the retinocollicular

pathway. Specifically, by comparing the observed and simulated retinotopy patterns, here we

addressed 1) to what extent RGC axon terminals are deviated from their retinotopically optimal

target locations in SC; and 2) on average how many RGCs connect to individual SC neurons.

Results

For functional mapping of the mouse retinocollicular projections, we expressed axon-targeted

calcium indicators (GCaMP6s; Broussard et al., 2018) in RGCs via intravitreal injection of

recombinant adeno-associated viruses (AAVs) harboring the pan-neuronal human synapsin

(hSyn) promoter, and monitored the visual responses of the labelled RGC axons in SC using

in vivo two-photon microscopy (Figure 2A,B and Supplemental Movie 1). To segment axonal

patches of individual RGCs and isolate their activity, we used constrained non-negative matrix

factorization (CNMF) that allows us to extract morphological and temporal features from noisy

time-lapse calcium activities based on their spatiotemporal correlations (Giovannucci et al.,

2019). For a field of view of ~0.3 mm2 (0.57-by-0.57 mm), we detected 26 ± 9 axonal patches

displaying independent activity patterns (median ± median absolute deviation here and

thereafter unless otherwise noted; N=37 recording sites in total from 15 animals; e.g., Figure

2C,D). The size of the individual axonal patches (135 ± 25 µm; N=969; Supplemental Figure

1A) is consistent with the past anatomical measurement (Hong et al., 2011), with a substantial

overlap between them over the SC surface (49 ± 11 %). This supports a successful signal

extraction and a good coverage of RGC axonal labelling. These data allowed us to faithfully

reconstruct the local two-dimensional (2D) map of the individual RGC axon terminals in SC

(e.g., Figure 2D).

**Figure 2: *In vivo* two-photon calcium imaging of retinal ganglion cell axon terminals in
the mouse superior colliculus.** *A*: Schematic diagram of the experimental set up for RGC
axonal imaging in SC. *B*: Representative image of cranial window. Medial-posterior part of
SC was clearly visible through a cylindrical silicone plug attached to a glass coverslip. *C*:
Average intensity projection of representative axonal imaging data, overlaid with detected
RGC axonal patches (N=21; color-coded). See also Supplemental Movie 1. *D*: Footprint of
three representative RGC axonal patches (#18, 2, and 20 in distinct color; from left to right,
respectively) overlaid with the profile of the rest patches (in grey). *E*: Corresponding RF of the
three representative RGC axonal patches (from D), estimated by reverse-correlation analysis.

Supplemental Figure 1: Probability distributions of axonic and receptive field statistics.
A: Size of the identified RGC axon terminals, d = 2(A/π)0.5, where A is the axonic field area from the CNMF analysis (135 ± 25 µm, median ± median absolute deviation; N=969). *B*: RF size of RGC axons (green; 4.8 ± 1.2 degrees; N=719) and SC somata (purple; 5.1 ± 0.9 degrees; N=1191), measured as the mean of long- and short-diameters (equivalently, the sum of long- and short-radii, a and b, respectively) of the 2D Gaussian profile at 1 σ fitted to the RF (e.g., Figure 2E). *C*: Distance between neighboring RGC axon centers (green; 100 ± 30 µm; N=761 pairs) or SC somata (purple; 56 ± 28 µm; N=2292 pairs). Neighboring pairs are identified by the Delaunay triangulation analysis (e.g., Figure 3C and Supplemental Figure 2F). *D*: Distance between neighboring RF centers of RGC axons (green; 7.2 ± 2.7 degrees; N=776 pairs) or SC somata (purple; 4.1 ± 1.9 degrees; N=2287 pairs). Neighboring pairs are identified by the Delaunay triangulation analysis (e.g., Figure 3D and Supplemental Figure 2G).

Supplemental Movie 1: *In vivo* two-photon calcium imaging of retinal ganglion cell axon terminals in the mouse superior colliculus. A snippet of motion-corrected movie (resampled at 30 Hz; original sampling rate, 15.4 Hz) showing the activity of RGC axons in the mouse SC in response to the random checkerboard stimuli (see Methods for details). See Figure 2 for the segmentation and Figure 3 for the tiling pattern analysis.

Presence of a well-defined receptive field (RF) is a characteristic feature of all RGC

types (Masland, 2012; Baden et al., 2016). To map the RF of the identified RGC axons, we

computed the response-weighted average of the presented random checkerboard stimuli

(frame rate, 4 Hz; rectangular fields, 3.7° in width and 2.9° in height; e.g., Figure 2E) and fitted

a 2D Gaussian at the peak latency to characterize the spatial RF profile. Most identified axonal

patches had RFs within the stimulation screen (±22° in elevation and ±36.5° in azimuth from

the mouse eye; Boissonnet et al., 2022). In accordance with *ex vivo* retinal physiology (Baden

et al., 2016), the average RF size of the RGC axons was 4.8 ± 1.2 degrees (N=719;

Supplemental Figure 1B), estimated as the mean of the long- and short-axis diameters of the

2D Gaussian profile at 1 standard deviation (SD). There was a weak but statistically significant

correlation between the RF size and the RGC axonal patch size (Pearson's r = 0.19, p = 5e-

7). The RFs locally tiled the visual field with 10 ± 5 % overlap at 1 SD Gaussian profiles, where

the center of every RF occupied a unique location in the visual field. This ensures that these

RFs belong to different RGCs because the RF center location of RGC axons should

correspond well to the location of their somata in the retina. Hence, the RF tiling faithfully

represents retinotopy.

How well does the tiling pattern of RGC axons in SC agree with that of their RFs? As

expected from the global retinotopy in SC (Dräger and Hubel, 1976; Cang et al., 2018; Sibille

et al., 2022), relative positions of the RGC axonal patches (e.g., Figure 3A) agreed well with

those of the corresponding RFs (e.g., Figure 3B) regardless of their cell types. For

quantification, we first computed the Delaunay triangulation using the geometric centers of the

individual axonal patches or the RF center locations as landmark points in each space (e.g.,

Figure 3C,D, respectively). This triangulation features the adjacency relationship regardless

of their absolute positions, where all the adjacent pairs of the landmark points in a given space

are connected as a dual graph of the Voronoi tessellation that separates the space into

territories close to each landmark point. The distances between the centers of neighboring

RGC axons and those between their RFs were identified to be 100 ± 30 μm (N=761 pairs;

Supplemental Figure 1C) and 7.2 ± 2.7 degrees (N=776 pairs; Supplemental Figure 1D),

respectively. As a measure of the agreement between the two tiling patterns, we then

calculated the fraction of the common edges between the two Delaunay triangulations (blue

edges; 88.2% for the example in Figure 3C,D). Throughout our datasets, we found a near-

perfect match between the tiling patterns of RGC axons in SC and their RFs (84 ± 5%; N=36

recordings from 12 animals; Figure 4A) regardless of the recording depth within the superficial

SC layer (120-220 µm deep from the surface; Figure 4B). This observation is consistent with

the precise axonal projection model (Model 1 in Figure 1) whereby RGC axon terminals

retinotopically tile the SC surface at single-cell precision.

Figure 3: Retinal ganglion cell axons retinotopically tile the mouse superior colliculus
at single-cell precision. *A,B*: RGC axon patches in SC from a representative recording
session (A; N=21, color-coded; from Figure 2C) and their corresponding RFs (B; 1 SD
Gaussian profile). *C,D*: Delaunay triangulation of the RGC axonal patch locations (C; from A)
and the RF centers (D; from B). The two triangulation patterns are nearly identical (88.2%;
blue, common edges in both patterns; red, unique edges only in either pattern). *E*:
Comparison between the observed tiling pattern of RGC axons (filled circles; from C) and the
retinotopically ideal pattern (open circle) obtained by applying an optimal Affine transformation
to the corresponding RF locations (in D) that minimizes the discrepancy between the two
patterns (Δ = 26 ± 8 µm). *F*: Corresponding comparison between the observed (filled squares;
from D) and ideal (Affine-transformed pattern in C) RF tiling patterns of RGC axons (Δ = 1.7
± 0.3 degrees).

Supplemental Figure 2: Tiling pattern analysis of local neurons in the mouse superior
colliculus. Figure panels are shown in the same format as in Figures 2 and 3. *A*: Schematic
diagram of the experimental set up for SC somatic imaging. *B*:Average intensity projection of
representative imaging data, overlaid with detected SC somata (N=88; color-coded). *C*:
Representative RF of a local SC neuron (#136) from reverse-correlation analysis. *D,E*: Tiling
pattern of SC somata (D) from a representative recording session (B) and their corresponding
RF tiling pattern (E; 1 SD Gaussian profiles with the same color-code as in B). *F,G*: Delaunay
triangulation of the SC somatic locations (F; from D) and the RF centers (G; from E), showing
a good agreement between them (74.1%; blue, common edges in both patterns; red, unique
edges only in either pattern). *H,I*: Comparison between the observed and retinotopically ideal
tiling patterns of SC somata (H; Δ = 28 ± 10 μm) or their RFs (I; Δ = 1.9 ± 0.8 degrees).

Figure 4: Tiling patterns of retinal ganglion cell axons are retinotopically more precise
than those of local neurons in the mouse superior colliculus. *A*: RGC axons (84 ± 5%;
N=36 recordings from 12 animals; e.g., Figure 3) showed a significantly higher precision of
retinotopy than SC somata (77 ± 5%; N=20 recordings from 20 animals; e.g., Supplemental
Figure 2); p = 0.001, rank sum test. The error bars show the median and interquartile range;
and the dotted lines represent the chance level at p = 0.05 (11 - 28%; bootstrap with 10,000
repetitions). *B*: The precision of the retinotopic tiling of RGC axons was not dependent on the
projection depth from the SC surface (N=4 animals; 83 ± 5, 83 ± 6, and 82 ± 5% at 120, 170,
and 220 µm deep, respectively; mean ± SD; p = 0.9, one-way analysis-of-variance). *C*:
Deviation between the observed and retinotopically ideal locations of the RGC axons (27 ± 4
µm; e.g., Figure 3E) or SC somata (23 ± 5 µm; e.g., Supplemental Figure 2H); p = 0.27, rank
sum test. *D*: Deviation between the observed and retinotopically ideal RF locations of the
RGC axons (1.9 ± 0.3 degrees; e.g., Figure 3F) or SC somata (1.8 ± 0.3 degrees; e.g.,
Supplemental Figure 2I); p = 0.66, rank sum test.

182   To further quantify the precision of the RGC axonal projection, we compared the

observed position with the ideal one that forms perfect retinotopy (e.g., Figure 3E). Specifically,

we used a linear method to estimate this retinotopically optimal tiling pattern of RGC axons:

i.e., an Affine transformation that best mapped the observed RF tiling pattern onto the

corresponding observed axonal tiling pattern (see Methods for details). We found that the

average discrepancy between the observed and retinotopically optimal RGC axonal locations

in SC was 27 ± 4 µm (Figure 4C). This is much shorter than the distance between neighboring

RGC axon centers (100 ± 30 µm, N=761 pairs; Supplemental Figure 1C) or the axonal patch

size (135 ± 25 µm, N=969; Supplemental Figure 1A), and thus will not have a measurable

impact on the retinotopy. Likewise, the extent to which the observed RF tiling pattern deviated

from the linear optimal one (1.9 ± 0.3 degrees; Figure 4D; see Figure 3F for example) was

much smaller than the RF size of RGC axons (4.8 ± 1.2 degrees, N=719; Supplemental Figure

1B) or the spacing between the RFs (7.2 ± 2.7 degrees, N=776 pairs; Supplemental Figure

1D). These data support that RGCs can precisely innervate their axons to their target locations

and faithfully transmit the information about retinotopy despite a loss of topographic

organization along the optic nerve (Horton et al., 1979; Colello and Guillery, 1998).

Thus far we have focused on the local retinotopy at the presynaptic input level, and

demonstrated a precise topographic organization of the RGC axons in the mouse SC (Figures

2-4). What about the postsynaptic side? Taking a similar approach, we next examined the

retinotopy of SC somata at single-cell resolution (Supplemental Figure 2). Specifically, using

in vivo two-photon calcium imaging, we mapped the RFs of local neurons in the superficial SC

layer (58 ± 27 cells/recording from 20 animals; RF size, 5.1 ± 0.9 degrees; RF overlap, 37 ±

11 %; N=1191 cells in total; Supplemental Figure 1B), and performed the same tiling pattern

analysis using the Delaunay triangulation (neighboring somata distance, 56 ± 28 µm, N=2292

pairs, Supplemental Figure 1C; neighboring RF distance, 4.1 ± 1.9 degrees, N=2287 pairs,

Supplemental Figure 1D). As expected, we found that the tiling patterns of SC somata and

their corresponding RFs agreed well in general (77 ± 5%, Figure 4A). However, the agreement

was significantly lower for SC somata than for RGC axons (p = 0.001, rank sum test; Figure

4A), indicating that local cellular-level retinotopy is less precise for the postsynaptic neurons

than for the input axons. Moreover, the average discrepancies between the observed and

retinotopically optimal locations of SC somata (23 ± 5 μm; Figure 4C) or their RFs (1.8 ± 0.3

degrees; Figure 4D) were comparable to those for RGC axons (p = 0.27 and 0.66,

respectively; rank sum test), suggesting that the connectivity between RGC axons and SC

neurons is not necessarily made to improve the precision of local retinotopy. Thus, our data

disagree with the selective connectivity model whereby SC neurons selectively integrate

inputs from appropriate presynaptic partners to reconstruct the topography at single-cell

resolution (Model 2 in Figure 1). Instead, we suggest that retinotopy in SC arises primarily

from precise RGC axonal projections (Model 1 in Figure 1), without much need to elaborate

the postsynaptic connectivity.

221
Figure 5: Data-driven model prediction on the precision of retinal ganglion cell axonal
projection to the mouse superior colliculus. *A*: Schematic of a retinocollicular projection
model. We assumed that 1) a jitter of RGC axonal projection follows a Gaussian distribution
$N[0, \sigma^2]$; and 2) the tiling pattern of RGC RF centers corresponds to that of the cell locations
in the retina. See Methods for details. *B,C*: Representative tiling patterns of simulated RGC
RF centers (B; on a 10%-jittered hexagonal lattice) and the corresponding axon centers at
different jitter levels (C; σ = 10, 27 and 40 μm from left to right panels, respectively). Common
(blue) and unique (red) triangulation edges are also shown in each panel of C, when compared
to the RF tiling pattern in B. The hexagonal lattice spacing was set to be 7.2 degrees and 100
μm for RGC RFs and axons, respectively, from the experimental data (Supplemental Figure
1). *D*: Correspondence of the triangulation edges between simulated RGC RF and axon tiling
patterns at different projection jitter levels (median with 95% confidence interval; 1,000
repetitions). The intersection with the experimentally identified value (horizontal dotted line,
84% from Figure 4A) gives a model prediction on the precision of RGC axonal projection
(vertical yellow line; σ = 27 ± 4 μm, with 95% confidence interval).

237   Our tiling pattern analysis showed a near-perfect retinotopy already at the level of the

axonal inputs to the mouse SC and no further improvement in the precision of retinotopy for

local SC neurons (Figures 3 and 4). What are the conditions to achieve such topographic

organizations in the retinocollicular pathway? To address this question, we next performed a

computational modelling analysis (see Methods for details). Specifically, by comparing the

observed and simulated tiling patterns on the pre- and post-synaptic sides, here we quantified

the following two parameters: 1) the precision of RGC axonal projection to a target location in

SC (Figure 5); and 2) the number of connecting RGCs to individual SC neurons (Figure 6).

245   We first modelled the tiling patterns of RGC axons at different jitter levels to identify

how small the projection error needs to be to recapitulate the observed precision of retinotopy

(Figure 5A). The tiling pattern of RGC somata – or equivalently, that of RGC RFs – was

simulated as a 2D hexagonal lattice with a small additive Gaussian noise, where the standard

deviation of the jitter followed 10% of the lattice spacing to replicate the dense packing of the

cell bodies in the retina (e.g., Figure 5B). The tiling pattern of RGC axons in SC was then

simulated by introducing additional Gaussian noise to the simulated RGC RF tiling pattern,

where the standard deviation σ of this additional noise determines the jitter level of the axonal

projection (e.g., Figure 5C). Here we set the axonal lattice spacing to be 100 µm based on our

experimental data (Supplemental Figure 1C), and ran the tiling pattern analysis as we did on

our experimental data to quantify the precision of retinotopy in the model. As expected, the

larger the jitter was, the less precise the retinotopy was (Figure 5D). This allowed us to

determine the jitter size that agreed with the observed precision level of retinotopy (84%;

Figure 4A): i.e., σ = 27 ± 4 µm (with 95% confidence interval). This is consistent with the

average discrepancy between the observed and retinotopically optimal tiling patterns (Δ = 27

µm; Figure 4C), hence validating our modelling framework and further supporting our estimate

on the precision of the RGC axonal projection.

Figure 6: Data-driven model prediction on the number of connecting retinal ganglion
cells to individual neurons in the superior colliculus. *A*: Schematic of a retinocollicular
mapping model. We assumed that 1) each SC neuron integrates inputs from λ nearest
neighbor RGC axons, where λ follows a Poisson distribution; and 2) the connectivity weights
depend on the amount of overlap between SC dendritic field (radius, 200 μm; Gale and Murphy,
2014) and RGC axonic field (135 μm; Supplemental Figure 1A). RGC axonal tiling was
simulated with σ = 27 μm (Figure 5). See Methods for details. *B,C*: Representative tiling
patterns of simulated SC soma centers (B; on a 10%-jittered hexagonal lattice with 56 μm
spacing; Supplemental Figure 1C) and the corresponding RF centers at different integration
levels (G; λ = 3, 5.5, and 8, from left to right panels, respectively), overlaid with common (blue)
and unique (red) triangulation edges. *D*: Correspondence of the triangulation edges between
simulated SC soma and RF tiling patterns at different input convergence levels (median with
95% confidence interval; 1,000 repetitions). The intersection with the experimentally identified
value (horizontal dotted line, 77% from Figure 4A) gives a predicted number of connecting
RGCs to individual SC neurons (vertical yellow line; λ = 5.5 ± 1.0, with 95% confidence
interval).

We next modelled the retinotopy of SC somata on top of the optimal RGC projection

model described above (jitter size, σ = 27 μm), using the average number of RGC inputs to

SC neurons, λ, as a key model parameter (Figure 6A). The simulated tiling pattern of SC

somata (e.g., Figure 6B) was generated in a similar way to that of RGC somata, but with a

lattice spacing of 56 μm based on our experimental data (Supplemental Figure 1C). For each

simulated SC cell, the RF center location was then determined as a weighted average of the

RF centers of neighboring RGC axons (Figure 6C), where we assumed that the number of

connecting RGCs followed a Poisson distribution (mean, λ), and that the connectivity strength

was proportional to the amount of overlap between the SC cell's dendritic field (radius, 200

μm; Gale and Murphy, 2014) and the RGC's axonic field (135 μm; Supplemental Figure 1D).

Here we introduced a rather simple connectivity rule as implicated by our experimental data

solely from the retinotopy viewpoint (Figures 3 and 4), while details on the cell-type specific

connectivity are beyond the scope of our modelling framework. The tiling pattern analysis on

the simulated SC cells then showed that the larger the number of connecting RGCs was, the

more precise the retinotopy was (Figure 6D). This suggests that the observed relatively less

precise retinotopy for SC somata (77% as opposed to 84% for RGC axons; Figure 4A) results

from a low input convergence. Indeed, by comparing the model outcome with the experimental

data, here we derived that on average SC neurons receive inputs from λ = 5.5 ± 1.0 RGCs

(with 95% confidence interval). This is consistent with the observation in the previous

electrophysiological studies (Chandrasekaran et al., 2007; Sibille et al., 2022).

Discussion

Using *in vivo* two-photon axonal imaging, we conducted functional mapping of the mouse

retinocollicular projection, and demonstrated a precise retinotopic tiling of RGC axon terminals

in SC at single-cell resolution (Figures 2 and 3). Here we calculated the projection error size

in two different ways: 1) based on the deviation from a linearly-estimated retinotopically-ideal

target location (Figures 3 and 4), and 2) by data-driven computational modelling (Figure 5).

Both methods consistently found that the projection jitter was below 30 μm, much smaller than

the observed RGC axonic field size (135 μm; Supplemental Figure 1). These axons can thus

be innervated to their exact target locations to faithfully transmit topographic information from

the retina, even after a loss of topographic organization through the long-range projection via

the optic nerve (Horton et al., 1979; Colello and Guillery, 1998).

In contrast, we found that the local retinotopy of SC somata was no better than that of

RGC axons (Figure 4 and Supplemental Figure 2). The connectivity between RGCs and SC

cells is thus not necessarily made to retain or improve the topography. Instead, assuming no

selectivity in the connectivity patterns, our modelling analysis indicates that a reduced

precision of local retinotopy on the postsynaptic side can be a direct consequence of a low

input convergence level (Figure 6). Based on our experimental data, we derived from our

model that on average SC neurons receive inputs from ~5.5 RGCs (Figure 6). This is

consistent with the past electrophysiological measurements (Chandrasekaran et al., 2007;

Sibille et al., 2022), justifying our model framework and conclusion.

Taken together, we suggest that retinotopy in the mouse SC arises largely from

topographically precise projection of RGC axons, rather than local circuit computation by SC

neurons. Nevertheless, the precision of axonal projection was not perfect. Postsynaptic circuit

mechanisms should then be indispensable as well in retaining topography, especially for

higher-order processing because otherwise retinotopy will no longer be recognizable after a

cascade of signal transmission along the visual hierarchy (e.g., below chance level after six

~80% precision transmissions). It is a future challenge to investigate the cellular-level

topographic organization in other brain areas, including the retinotopy in the downstream

visual pathways (Wandell et al., 2007), and clarify the contribution of pre- and post-synaptic

circuit mechanisms in each area.

Having a precise retinotopy at single-cell resolution facilitates spatial information

processing not only at a global level, but also at local circuit levels. For example, looming

detection has been suggested to arise de novo in the superficial SC layer (Lee et al., 2020).

In principle, this can be achieved even in the absence of retinotopy by elaborating the wiring

among local neurons. It is, however, much more efficient to exploit precise topographic

information conveyed from the retina because the connectivity length and its complexity can

be minimized to locally process spatial information at any point in the visual field. This will also

help align different topographic maps in the same brain area to function coherently, such as

the retinotopy, orientation, and ocular dominance maps in SC (Feinberg and Meister, 2015;

de Malmazet et al., 2018). The observed precise spatial organization we demonstrate here

suggests that the wiring efficiency indeed matters for local circuit computation.

How can then such a precise retinotopic projection be formed? Retinotopic map

formation in SC occurs during the first postnatal week in mice, involving both genetic and

activity-dependent factors (Cang and Feldheim, 2013; Arroyo and Feller, 2016). These factors

also play a key role in the development of a fine-scale organization in other sensory systems,

such as the tonotopic map in the cochlear nucleus (Krasewicz and Yu, 2023), and the

chemotopic map in the olfactory bulb where olfactory sensory neurons expressing the same

olfactory receptor type project exclusively to the same single glomerulus (Imai et al., 2010).

While overall sensory map formation is genetically predetermined by molecular cues (e.g.,

ephrin-Eph signaling; Frisén et al., 1998; Krasewicz and Yu, 2023), a precise topography is

established only after refinement that involves spontaneous activity, such as retinal waves

during development (Arroyo and Feller, 2016), and eventually experience-driven alignment

(McLaughlin et al., 2003). In particular, here we suggest that this refinement process of the

retinocollicular projection should be extremely precise, to the extent that retinotopy arises at a

single-cell resolution regardless of the RGC types in adult animals (Figures 3 and 4). It is then

possible that neuronal circuits are in general wired more precisely than previously thought to

exploit topographic information for their function, including long-range projections to other

retinal targets (Liang, et al., 2018) as well as those in other systems, such as the callosal

projections and entorhinal-hippocampal networks (Jbabdi et al., 2013; Patel et al., 2014).

358 Methods

359   No statistical method was used to predetermine the sample size. All experiments involving

360   animals were performed under the license 233/2017-PR from the Italian Ministry of Health.

361   The data analyses were done in Python and Matlab (Mathworks). The statistical significance

362   level was set to be 0.05. All summary statistics were described as median ± median absolute

363   deviation unless otherwise noted.

Animals

Female C57BL/6J mice (15 for axonal imaging; 20 for somatic imaging) were used at around 6-10 weeks of age at the time of the first surgery. Mice were kept on a 12-h light / 12-h dark cycle and given water and food *ad libitum*. After surgery, the animals were kept in groups operated on the same day. Mice were between 12-24 weeks of age at the time of imaging experiments.

Intravitreal viral injections

Intravitreal injection of recombinant adeno-associated virus (AAV) 2, pseudotyped with a hybrid of AAV1 and AAV2 capsids, was used to deliver hSyn-axon-GCaMP6s expression cassette to the mouse retinal ganglion cells (RGCs; Broussard et al., 2018). For the viral injection, mice were anaesthetized (induction, 4% isoflurane in oxygen; maintenance, 1.8-2.0%) and kept on a heated plate (Supertech Physiological Temperature Controller) to avoid hypothermia. Both eyes were protected by saline drop or viscous eye ointment (VitA-POS, Ursapharm). The scleral surface on the left eye was exposed and a small piercing was made with a sterile 28-30G needle in between the sclera and the cornea. An injection pipette (~50 µm tip diameter with 30-40° bevel) prefilled with a virus solution (~1.5×10^{14} vg/mL in phosphate-buffered saline with 0.001% Pluronic F68 and 0.001% FastGreen) was then inserted into the vitreous chamber approximately 1 mm deep. The injection pipette was made from a borosilicate glass capillary (1B120F-3, WPI) with a pipette puller (DMZ, Zeitz) and a microgrinder (EG-45, Narishige). After a good sealing of the pipette was formed, 1.2 µL of the virus solution was injected at a rate of 10 nL/s using a microinjection pump (either Neurostar NanoW or WPI NanoLiter 2010) with mineral oil (Sigma, M5904) filled in the displacement space by a stainless steel plunger. The pipette was slowly withdrawn at least 5 minutes after the completion of the injection, and the treated eye was covered with the eye ointment. The animal was then allowed to recover from anesthesia in a warmed-up chamber and brought back to its home cage.

Intracranial viral injections

Pseudotyped AAV, composed of AAV2 rep and AAV9 cap genes or a hybrid of AAV1 and AAV2 cap genes, was locally injected to the mouse superior colliculus (SC) for the expression of genetically-encoded calcium indicators (jGCaMP7f, jGCaMP8m or jRGECO1a) under pan-neuronal human synapsin (hSyn) promoter. The intracranial viral injection was made at the same time as the cranial implantation as described below. After making a craniotomy over the right SC, an injection pipette (~30 µm tip diameter; WPI 1B120F-3 borosilicate glass capillary pulled with Zeitz DMZ puller) prefilled with a virus solution (~5×10^{12} to ~4×10^{14} vg/mL in phosphate-buffered saline) was inserted across the dura at coordinates from Bregma around -4 mm AP, 0.5-0.7 mm ML, and then slowly advanced until ~1.25 mm deep. The virus solution (0.4-0.6 µL) was injected at a rate of 2 nL/s with a microinjection pump (either Neurostar NanoW or WPI NanoLiter 2010). The pipette was slowly withdrawn at least 10 minutes after the completion of the injection, followed by the cranial window implantation procedure.

Cranial implantations

We adapted methods described in Feinberg and Meister (2015) for the cranial window implantation over the mouse SC. A cranial window assembly was made in advance, where the surface of a circular glass coverslip (5 mm diameter, 0.13-0.15 mm thickness; Assistent Karl Hecht) was activated by a laboratory corona treater (BD-20ACV Electro-Technic Products) and fused to a cylindrical silicone plug (1.5 mm diameter, 0.75 - 1.00 mm height; Kwik-Sil, WPI) by baking it for 24 hours at 70-80°C.

For the implantation, animals were anaesthetized (induction, 4% isoflurane in oxygen; maintenance, 1.5-2.0%) and placed inside a stereotaxic apparatus (Stoelting 51625). Throughout the surgery, temperature was maintained at 37°C using a heated plate (Supertech Physiological Temperature Controller) to avoid hypothermia, and the eyes were protected with eye ointment (VitA-POS, Ursapharm). After disinfecting and removing the scalp (Betadine 10%, Meda Pharma), the skull surface was scratched and cleaned to ensure good cement

adhesion. A craniotomy of a size about 3.0 mm (anterior-posterior; AP) by 2.5 mm (medial-

lateral; ML) was made over the right SC using a high-speed surgical drill (OmniDrill35, WPI)

with a 0.4 mm ball-tip carbide bur (Meisinger). To prevent bleeding, the craniotomy was treated

by hemostatic sponges (Cutanplast, Mascia Brunelli) soaked with sterile cortex buffer (NaCl

125 mM, KCl 5 mM, Glucose 10 mM, HEPES 10 mM, $CaCl_2$ 2 mM, $MgSO_4$ 2 mM, pH 7.4).

For SC somata imaging, viral injections were made as described above. The implant was then

placed in a way to push the transversal sinus and posterior cortex ~0.5 mm forward and

position the silicone plug over the medial-caudal region of the right SC. Tissue adhesive

(Vetbond, 3M) was used to fix and seal the implant. A custom-made titanium headplate (0.8

mm thick) was then cemented to the skull using acrylic cement powder (Paladur, Kulzer) pre-

mixed with cyanoacrylate adhesive (Loctite 401, Henkel).

After the surgery, the animal was recovered from anesthesia in a warmed-up chamber

and returned to its home cage. For postoperative care, animals were given intraperitoneally 5

mg/kg Rimadyl (Zoetis) and 5 mg/kg Baytril (Bayer) daily for 3-5 days. We waited for another

10-15 days until the cranial window completely recovered before starting *in vivo* two-photon

imaging sessions (e.g., Figure 2B).

Visual stimulation

Visual stimuli were presented to the subject animals as described previously (Boissonnet et

al., 2022). In short, a custom gamma-corrected digital light processing device was used to

project images (1280-by-720 pixels; frame rate, 60 Hz) to a spherical screen (radius, 20 cm)

placed ~20 cm to the contralateral side of an animal's eye, stimulating the visual field ±22° in

elevation and ±36.5° in azimuth (Figure 2A and Supplemental Figure 2A). We presented 1)

random water-wave stimuli (2-10 min) for generating binary masks for signal source extraction

in calcium image analysis (see below); and 2) randomly flickering black-and-white

checkerboard stimuli (10 min) for receptive field mapping, with rectangular fields 3.7° in width

and 2.9° in height, each modulated independently by white noise at 4 Hz.

In vivo two-photon imaging

Prior to *in vivo* imaging sessions, animals were habituated to stay head-fixed on a custom-made treadmill disc (8-10 habituation sessions in total over a week, each for 2 hours). For the imaging session, animals were kept on the treadmill with their head fixed for no longer than 2 hours (2-5 sessions/animal). Two-photon calcium imaging was done on a galvo-resonant multiphoton microscope (Scientifica HyperScope with SciScan image acquisition software) equipped with a mode-locked tunable laser (InSight DS+, Spectra-Physics) and a plan fluorite objective (CFI75 LWD 16X W, Nikon). In each imaging session, we performed single-plane time-lapse recordings (field of view, approximately 0.65-by-0.65 mm) at a depth of 120-220 µm from the SC surface. The fluorescent signal (excitation wavelength, 920 nm for axon-GCaMP6, GCaMP7f, and GCaMP8m; 1040 nm for jRGECO1a; average laser power under the objective, 40-80 mW) was bandpass-filtered (BP 527/70 or BP 650/100 after beam-splitter FF580-FDi01, Semrock) and detected with a non-descanned gallium arsenide phosphide photomultiplier tube (Hamamatsu GaAsP PMT). Each frame was acquired with 1024-by-1024 pixels (16-bit depth) at 15.4 Hz for RGC axonal imaging, and 512-by-512 pixels (16-bit depth) at 30.9 Hz for SC somata imaging.

Calcium image analysis

For preprocessing of RGC axon data, the original 1024-by-1024 pixel images were first downsampled to 512-by-512 pixels (2-by-2 pixels averaging) to reduce noise. To correct motion artefacts, we performed two iterations of Fourier-based rigid image registration in ImageJ, followed by cropping the image border by 16 pixels; and then ten iterations of non-rigid motion correction (NoRMCorre) in CaImAn (Giovannucci et al., 2019), followed by a 12-pixel border crop. The resulting images (456-by-456 pixels; e.g., Supplemental Movie 1) represent a field of view of around 0.57-by-0.57 mm (1.3 µm/pixel).

From the preprocessed images, we identified the axonal patches of individual RGCs and extracted their signals in CaImAn (e.g., Figure 2C,D). Specifically, using a part of the

recordings (3,000-5,000 frames representing the random water-wave stimulus presentation

period), we first ran two iterations of constrained non-negative matrix factorization (CNMF) in

CaImAn, where we set the number of expected components (*params.K*) to be 60 as an

initialization parameter. From the identified components, we then manually selected those with

a uniformly-filled oval-like shape that had a size of around 50-150 μm as biologically relevant

ones (Hong et al. 2011), and converted them into binary spatial masks to run two iterations of

masked CNMF for processing the entire time-lapse recordings. The resulting set of spatial

components (*estimates.A*) and deconvolved neural activities (*estimates.S*) was used for the

subsequent analyses.

477   The area of the individual RGC axonal patches P_i was estimated from the identified

478   spatial components in CaImAn (1.3 μm/pixel), from which the radius was estimated as

479    $(P_i/\pi)^{0.5}$ under the assumption of a circular patch shape (Supplemental Figure 1A). The

480   fraction of the overlap between identified axonal patches was calculated as the ratio of the

481   areas between the intersection of any two patches $\cup_{i \neq j}(P_i \cap P_j)$ and the union of all patches

482    $\cup_i P_i$.

483   For SC soma data, we first ran a sequence of the rigid and non-rigid motion corrections

in CaImAn, followed by image cropping from 512-by-512 pixels into 480-by-480 pixels in

ImageJ (1.3 μm/pixel). Using a part of the recordings (3,000 frames from the random water-

wave stimulus presentation period), we then ran two iterations of CNMF in CaImAn, where the

images were divided into 6-by-6 (36 in total) patches and the expected number (*params.K*)

and size (*params.gSig*) of neurons were set to be 5 per patch and 5-by-5 pixels in half size,

respectively, as initialization parameters. From the identified putative cells, we manually

selected those with a uniformly-filled round shape of around 10-20 μm in size as biologically

relevant ones, and converted them into binary spatial masks to run two iterations of masked

CNMF in CaImAn on the entire time-lapse recordings. The resulting set of spatial components

(*estimates.A*) and deconvolved neural activities (*estimates.S*) was used for the subsequent

analyses.

Receptive field analysis

The receptive fields (RFs) of the identified RGC axon patches or SC somata were estimated by reverse-correlation methods using the random checkerboard stimuli (Chichilnisky, 2001). Specifically, we calculated the response-weighted average of the stimulus waveform (0.5 s window; 1/60 s bin width), and characterized its spatial profile by the two-dimensional (2D) Gaussian curve fit at the peak latency (e.g., Figure 2E and Supplemental Figure 2B). The RF center was assigned to the center of that 2D Gaussian profile, and the RF size was estimated as twice the mean standard deviation (SD) of the long and short axes (Supplemental Figure 1B). The fraction of the overlap between the RFs (1 SD Gaussian profiles) was computed similarly as for the axonal patches. The Pearson correlation coefficient between the RF size and the RGC axonal patch size was calculated with the 95% interval of the data to eliminate the outliers. Those cells that had the RF center on the border or outside the stimulus screen were eliminated from the tiling pattern analysis described below. Those recordings that had less than 10 cells with RF centers on the stimulus screen were also excluded from the tiling pattern analysis.

Tiling data analysis

To compare the tiling patterns between RGC axon patches / SC somata and their RFs (e.g., Figure 3 and Supplemental Figure 2), we first computed the Delaunay triangulation of their centroid locations using the Euclidean distance in each space (e.g., Figure 3C,D and Supplemental Figure 2C,D). As a measure of similarity between the two tiling patterns, we then calculated the number of common edges, divided by the mean of the total number of edges in each triangulation. The chance level was calculated by a bootstrap method (10,000 repetitions; Figure 4A).

We used the least squares method to identify an optimal Affine transformation for mapping a given RF tiling pattern onto the corresponding tiling pattern of RGC axons (e.g., Figure 3E), or vice versa (e.g., Figure 3F). The Euclidean distance of the cell or RF locations

521 between the observed and affine-transformed tiling patterns was then used as a measure of

522 the precision of local retinotopy (Figure 4C,D).

523 Modelling of retinocollicular mapping

We modelled the retinocollicular mapping in four steps (Figures 5 and 6).

1. The tiling pattern of RGC somata was simulated as a 2D hexagonal lattice with a

Gaussian jitter $N[0, \sigma^2]$. The standard deviation σ was set to be 10% of the lattice

spacing L to recapitulate the dense packing of the cell bodies in the retina. We

assumed that the tiling pattern of RGC RFs was equivalent to the corresponding

somatic tiling pattern (e.g., Figure 5B; L = 7.2 degrees from Supplemental Figure 1D).

2. The tiling pattern of RGC axons in SC (i.e., retinocollicular projection) was then

simulated by introducing additional Gaussian noise $N[0, \sigma^2]$ to the RGC RF tiling

pattern from step 1 (e.g., Figure 5C) but with L = 100 μm (Supplemental Figure 1C).

When σ = 0 μm, the tiling pattern of RGC axons is identical to the somatic tiling pattern,

showing perfect retinotopy (i.e., 100% tiling pattern match).

3. The tiling pattern of SC somata was simulated as a 2D hexagonal lattice (L = 56 μm;

Supplemental Figure 1C) with a Gaussian jitter (σ = 0.1L) as in step 1 (e.g., Figure 6B).

4. The RF of each SC neuron was calculated by integrating inputs from λ nearest

neighbor RGC axons, where λ follows a Poisson distribution. Specifically, assuming

that the connectivity strength depends on the amount of overlap between SC dendritic

field (radius, 200 μm; Gale and Murphy, 2014) and RGC axonic field (135 μm;

Supplemental Figure 1A), we defined the SC RF center location as the weighted

average of the RF centers of the connecting RGCs (e.g., Figure 6C).

To identify the precision of RGC axonal projection to SC (Figure 5), we ran the steps 1 and 2

at different jitter levels σ (from 0 to 50 μm in steps of 1 μm; 1,000 repetitions each) and

calculated the similarity between the simulated tiling patterns of RGC axons and their RFs

using the triangulation method as described above (Figure 5D). We then determined the jitter

level σ where the simulated tiling pattern similarity agreed with the experimental data (84%;

Figure 4A).

To estimate the average number of connecting RGCs to individual SC neurons (Figure

6), we ran the steps 1-4 at different mean values λ (from 1 to 10 RGCs in steps of 0.25; 1,000

repetitions; σ = 27 µm for step 2 from Figure 5D) and calculated the similarity between the

simulated SC somatic and RF tiling patterns using the triangulation method as described

above (Figure 6D). We then determined the input convergence level λ where the simulated

tiling pattern similarity agreed with the experimental data (77%; Figure 4A).

References

Arroyo, D. A., Feller, M. B. (2016). Spatiotemporal Features of Retinal Waves Instruct the
Wiring of the Visual Circuitry. Front Neural Circuits, 10, 54.

Baden, T., Berens, P., Franke, K., Rosón, M. R., Bethge, M., Euler, T. (2016). The
functional diversity of retinal ganglion cells in the mouse. Nature, 529, 345-350.

Boissonnet, T., Tripodi, M., Asari, H. (2022). Awake responses suggest inefficient dense
coding in the mouse retina. bioRxiv. 10.1101/2022.02.15.480512

Broussard, G. J., Liang, Y., Fridman, M., Unger, E. K., Meng, G., Xiao, X., Ji, N., Petreanu,
L., Tian, L. (2018). In vivo measurement of afferent activity with axon-specific calcium
imaging. Nat Neurosci, 21, 1272-1280.

Cang, J., Feldheim, D. A. (2013). Developmental mechanisms of topographic map
formation and alignment. Annu Rev Neurosci, 36, 51-77.

Cang, J., Savier, E., Barchini, J., Liu, X. (2018). Visual Function, Organization, and
Development of the Mouse Superior Colliculus. Annu Rev Vis Sci, 4, 239-262.

Chandrasekaran, A. R., Shah, R. D., Crair, M. C. (2007). Developmental homeostasis of
mouse retinocollicular synapses. J Neurosci, 27, 1746–1755.

Chichilnisky, E. J. (2001). A simple white noise analysis of neuronal light responses.
Network, 12, 199-213.

Colello, S. J., Guillery, R. W. (1998). The changing pattern of fibre bundles that pass
through the optic chiasm of mice. Eur J Neurosci, 10, 3653-3663.

de Malmazet, D., Kühn, N. K., Farrow, K. (2018). Retinotopic Separation of Nasal and
Temporal Motion Selectivity in the Mouse Superior Colliculus. Curr Biol, 28, 2961-2969.e4.

Dräger, D. C., Hubel, D. H. (1976). Topography of visual and somatosensory projections
to mouse superior colliculus. J Neurophysiol, 39, 91-101.

Fame, R. M., MacDonald, J. L., & Macklis, J. D. (2011). Development, Specification, and
Diversity of Callosal Projection Neurons. Trends Neurosci, 34, 41-50.

Feinberg, E. H., Meister, M. (2015). Orientation columns in the mouse superior colliculus.
Nature, 519, 229-232.

Frisén, J., Yates, P. A., McLaughlin, T., Friedman, G. C., O'Leary, D. D., Barbacid, M.
(1998). Ephrin-A5 (AL-1/RAGS) is essential for proper retinal axon guidance and
topographic mapping in the mammalian visual system. Neuron, 20, 235-243.

Giovannucci, A., Friedrich, J., Gunn, P., Kalfon, J., Brown, B. L., Koay, S. A., Taxidis, J.,
Najafi, F., Gauthier, J. L., Zhou, P., Khakh, B. S., Tank, D. W., Chklovskii, D. B.,
Pnevmatikakis, E. A. (2019). CaImAn an open source tool for scalable calcium imaging
data analysis. eLife, 8, e38173.

Hong, Y. K., Kim, I.-J., Sanes, J. R. (2011). Stereotyped axonal arbors of retinal ganglion
cell subsets in the mouse superior colliculus. J Comp Neurol, 519, 1691-1711.

Horton, J. C., Greenwood, M. M., Hubel, D. H. (1979). Non-retinotopic arrangement of
fibres in cat optic nerve. Nature, 282, 720-722.

Imai, T., Sakano, H., Vosshall, L. B. (2010). Topographic mapping – the olfactory system.
Cold Spring Harb Perspect Biol, 2, a001776.

Jbabdi, S., Sotiropoulos, S. N., Behrens, T. E. (2013). The topographic connectome. Curr
Opin Neurobiol, 23, 207-215.

Krasewicz, K., Yu, W.-M. (2023). Eph and ephrin signaling in the development of the
central auditory system. Dev Dyn, 252, 10-26.

Lee, K. H., Tran, A., Turan, Z., Meister, M. (2020). The sifting of visual information in the
superior colliculus. eLife, 9, e50678.

Liang, L., Fratzl, A., Goldey, G., Ramesh, R. N., Sugden, A. U., Morgan, J. L., Chen, C., Andermann, M. L. (2018). A Fine-Scale Functional Logic to Convergence from Retina to Thalamus. Cell, 173, 1343-1355.e24.

Malpeli, J. G., Baker, F. H. (1975). The representation of the visual field in the lateral geniculate nucleus of Macaca mulatta. J Comp Neurol, 161, 569-594.

Masland, R. (2012). The Neuronal Organization of the Retina. Neuron, 76, 266-280.

McLaughlin, T., Torborg, C. L., Feller, M. B., O'Leary, D. D. (2003). Retinotopic map refinement requires spontaneous retinal waves during a brief critical period of development. Neuron, 40, 1147-1160.

Patel, G. H., Michael, D. K., Snyder, L. H. (2014) Topographic organization in the brain: Searching for general principles. Trends Cogn Sci, 18, 351-363.

Sibille, J., Gehr, C., Benichov, J. I., Balasubramanian, H., Teh, K. L., Lupashina, T., Vallentin, D., Kremkow, J. (2021). Strong and specific connections between retinal axon mosaics and midbrain neurons revealed by large scale paired recordings. bioRxiv. 10.1101/2021.09.09.459396

Wandell, B. A., Dumoulin, S. O., Brewer, A. A. (2007). Visual field maps in human cortex. Neuron, 56, 366-383.