Facultative parthenogenesis: a transient state in transitions between sex and obligate asexuality in stick insects?

Chloé Larose[1]*, Guillaume Lavanchy[1]*, Susana Freitas[1], Darren J. Parker[1,2], Tanja Schwander[1]

[1]Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne.
[2]School of Natural Sciences, Brambell Building, Bangor University, Bangor, LL57 2UW, United Kingdom.
*Joint first authors. Co-first authors can prioritise their names when adding this paper to their résumés.

Corresponding authors: guillaume.lavanchy@unil.ch; tanja.schwander@unil.ch

Abstract

Transitions from obligate sex to obligate parthenogenesis have occurred repeatedly across the tree of life. Whether these transitions occur abruptly or via a transient phase of facultative parthenogenesis is rarely known. We discovered and characterised facultatively parthenogenetic populations of the North American stick insect *Timema douglasi*, a species in which only obligately parthenogenetic populations were known so far. These populations comprised three genetic lineages. Females from all lineages were capable of parthenogenesis (with variable efficiency) but their propensity to reproduce sexually after mating varied extensively. In all three lineages, parthenogenesis resulted in the complete loss of heterozygosity in a single generation. Obligately parthenogenetic *Timema* have also lost all heterozygosity, suggesting that parthenogenesis evolved via a gradual capacity increase while conserving the proximate mechanism. We speculate that facultative parthenogenesis may often be transient and be replaced by obligate strategies (either sex or parthenogenesis) because of a trade-off between the efficiency of the two reproductive modes. Such a trade-off could help explain why facultative parthenogenesis is rare among animals, despite its potential to combine the known benefits of sex and parthenogenesis.

Introduction

The way organisms reproduce varies extensively across the tree of life. While most animals engage in some form of sex to produce offspring (Bell, 1982; The Tree of Sex Consortium, 2014), many species take alternative routes to reproduction. The most widespread alternative is parthenogenesis, whereby females produce offspring without genetic contribution from males (Normark, 2003; Suomalainen et al., 1987).

Most species appear to be either obligately sexual or obligately parthenogenetic in natural populations, even when there is some capacity for the alternative reproductive mode. For example, virgin females in many sexual species can on occasion reproduce through spontaneous parthenogenesis ("tychoparthenogenesis"), typically with very limited success (Bell, 1982; Fields et al., 2015; Liegeois et al., 2021). On the other end of the spectrum, recent studies have revealed rare or cryptic sex in several species previously thought to be obligately parthenogenetic (Boyer et al., 2021; Freitas et al., 2023; Kuhn et al., 2021; Laine et al., 2022). However, even though such species are still capable of sex, this mode of reproduction is used very rarely. Thus, species cluster at the extremes of the continuum and are either largely sexual or largely asexual (Bell, 1982). This is surprising, given that facultative parthenogenesis, whereby the capacity for sex or parthenogenesis of females are approximately equal, has the potential to combine the long-term advantages of recombination and segregation that come along with sex, together with the short-term advantages of asexuality (D'Souza & Michiels, 2010; Otto, 2009).

The relative rarity of facultative parthenogenesis also raises the question of whether obligate parthenogenesis generally evolves abruptly from sexual ancestors. This is the case for example in many or most obligately asexual lineages of hybrid origin (Cuellar, 1974; Drosopoulos, 1978; Janko et al., 2018; Schultz, 1973). Alternatively, the evolution of obligate parthenogenesis without hybridization may involve facultative parthenogenesis as an intermediate step, which would then be transient. We here evaluate these alternatives using the stick insect genus *Timema*.

Timema stick insects are native to the western USA and Mexico and have been studied extensively for their repeated transitions from sex to obligate parthenogenesis. Five of the 21 described species reproduce via female-producing parthenogenesis, while the others are sexual (Sandoval et al., 1998; Vickery & Sandoval, 1999, 2001). In parthenogenetic species, fertilization of oocytes does not occur even when females are mated with males of related sexual species in the laboratory (Schwander et al., 2013). However, cryptic gene flow has been documented in some species, presumably mediated by rare males (Freitas et al. 2023). Conversely, females of the sexual species are largely incapable of parthenogenesis, although spontaneous parthenogenesis in 1-2% unfertilized eggs laid by virgin sexual

females is well documented (Schwander et al., 2010; Schwander & Crespi, 2009). When mated, sexual

females fertilize all their eggs, and population sex ratios are close to 50:50 (Arbuthnott et al., 2015;

Schwander et al., 2010). Thus the *Timema* populations described so far fit well with the general pattern

in animals whereby reproductive strategies are largely obligate.

However, we recently discovered populations with a large proportion of males in the geographic range

of the species *T. douglasi,* a species that was believed to be exclusively parthenogenetic. The male-

containing populations were geographically close to female-only populations, with an apparently abrupt

transition from female-only populations to populations with relatively even sex ratios. Such variable

population sex ratios could, for example, result from mixes between different proportions of sexual and

parthenogenetic females, or from facultative parthenogenesis with sex and parthenogenesis used at

different frequencies. Alternatively, if the newly discovered populations were sexual, variable sex ratios

could also stem from sex ratio distortion. For example, a driving X chromosome, which would be present

in more than 50% of transmitted sperm cells, could also generate female-biased sex ratios (Helleu et al.,

2015; Jaenike, 2001).

In order to investigate these different hypotheses, we sampled populations along two separate transects

and quantified the local sex ratios. We then characterized the reproductive modes of females from

populations with different sex ratios by measuring the capacity for parthenogenesis of virgin females

and by distinguishing parthenogenetic and sexual offspring in controlled crosses via genotyping based

on Restriction-Associated DNA Sequencing (RADseq). We also screened the sex ratio of sexual

offspring for deviations from the expected 50:50 ratio to test whether a distorter could cause the observed

biases in sex ratio. Finally, we also used the RADseq genotypes to assess whether females with different

reproductive modes belonged to different genetic lineages, and to study the phylogenetic relatedness

between these lineages and previously described lineages of parthenogenetic *T. douglasi* and close

sexual relative *T. poppense*.

Methods

Population sampling and reproductive mode characterisation

96   We characterized the population sex ratios at 29 sampling locations along two transects located in

97   northern California, respectively referred to as "Manchester" and "Orr" transect (Fig. 1). For both

98   transects, we chose the sampling locations according to the host plant distribution along the main road,

99   with isolated patches of redwood (*Sequoia sempervirens*) or douglas fir (*Pseudotsuga menziesii*)

100   considered as distinct locations. We sampled each location during 2 hours with approximately constant

sampling intensity and collected up to 270 individuals (all juvenile) per location. We sexed the individuals based on external morphology. In total, we collected 1195 individuals for the Manchester transect (864 females and 331 males in 12 different populations, hereafter called Manchester 1 [westernmost] to Manchester 12 [easternmost]; see Figure 1) and 1074 individuals for the Orr transect (1029 females and 45 males in 12 populations, hereafter called Orr 1 [easternmost] to Orr 17 [westernmost]; no *Timema* were found at 5 locations).

We aimed to characterise the reproductive mode of approximately 10 females from each female-only population and of approximately 20 females from each population with both sexes (when available). However, the final numbers varied considerably because of mortality, and because females sometimes laid too few eggs prior to mating to determine hatching success (see below). Individuals of each population were separated by sex and maintained in cubic mesh cages (30 x 30 x 30 cm) on douglas fir, in a climatic chamber at 23°C, 55% humidity and 12:12h day:night cycle. Since all individuals were collected as juveniles and only adults mate, this allowed us to obtain virgin females for all populations. We isolated 352 females in Petri dishes (14.5 cm diameter) to obtain unfertilized eggs, with fresh food (a small branch of douglas fir), soil (for females to ingest in order to coat their eggs), and a moistened piece of cotton wool (to prevent desiccation) added every two days. Once a female had laid approximately 20 unfertilized eggs, we allowed her to mate with one to three males from sampling locations with high male frequencies (Manchester 1 - 4, 6 - 7, Orr 5). For each female, the eggs laid before and after mating were kept separately until hatching, which occurs after approximately 5-6 months of diapause. The eggs began to hatch on October 27th of the same year. The eggs were then checked every other day and the number of hatchings recorded. Hatchlings were stored in 100% EtOH until further use. No eggs hatched after December 27th.

In addition, we re-analysed previously published data on the capacity for parthenogenesis of seven sexual species (*T. petita, T. podura, T. cristinae, T. chumash, T. knulli, T californicum* and *T. poppense)* from (Schwander et al., 2010; Schwander & Crespi, 2009) and characterized it for the five asexual species (*T. douglasi, T. shepardi, T. monikensis, T. genevievae* and *T. tahoe*). These data represent a total of 450 virgin females (range: 3 – 162 per species) collected in 2007 and 2008. We reared them under the conditions described above, except that they were maintained at room temperature and fed with different host plants according to the natural host plant use of each species (see Schwander et al., 2010 for details). We collected all eggs and assessed the hatching success of all females which had laid at least five eggs (424 females, 14021 eggs).

Genotyping

To test whether the individuals collected along the Manchester and Orr transects belonged to different
genetic lineages, we genotyped 32 females and their mates (1 - 3 per female, 42 males in total). We also
assessed whether eggs produced after mating were fertilized or not, and quantified heterozygosity in
parthenogenetically produced eggs. To do this, we genotyped 3 hatchlings from eggs laid before mating
(when available) for each of the 32 genotyped females (77 hatchlings from 27 females in total; for the 5
remaining females, none of the eggs laid prior to mating hatched), and 7 hatchlings from eggs laid after
mating (when available) for 24 of the 32 females. For the eight remaining families, we genotyped all
available hatchlings from eggs laid after mating (17 - 51), initially for a different project. In total, we
genotyped 365 post-mating hatchlings. Details on the number of hatchlings and males per family are
provided in Table S1.

We extracted the DNA of a leg (adults) or the whole body (hatchlings). We flash-froze the tissue in
liquid nitrogen and ground it via sonication with ceramic beads. We then performed the extraction using
the Qiagen DNA Blood and Tissue kit on a BioSprint 96 workstation, following manufacturer protocols.

We built double-digested RADseq libraries following the protocol by (Brelsford et al., 2016)Brelsford
et al. (2016)Brelsford et al. (2016)Brelsford et al. (2016). In short, we digested genomic DNA with
EcoRI and MseI and ligated Illumina TruSeq adapters to the cut site. The EcoRI adapter included an
individual 8-base barcode. We amplified the ligated fragments in 20 PCR cycles. We then multiplexed
the libraries and size-selected fragments of 300 - 500 bp on an Agarose gel. The resulting libraries were
then sequenced on 10 Illumina HiSeq 2500 lanes (100 - 125 bp, single end) at the Lausanne Genomics
Technology Facility.

We built loci using Stacks version 2.3e (Catchen et al., 2013). We first demultiplexed the raw reads with
the process_radtags module with parameters -c -q -r and --filter-illumina and trimmed all the reads to
92 bases. We then mapped the reads to the reference genome of *T. douglasi* (Bioproject accession
number: PRJEB31411; Jaron et al., 2022) using the mem algorithm of BWA version 0.7.17 (Li &
Durbin, 2010). We built loci using the gstacks module, enabling the --phasing-dont-prune-hets option.
Finally, we output a vcf file with a single SNP per locus using the --write-single-snp option in
populations. We then filtered genotypes in Vcftools version 0.1.15 (Danecek et al., 2011) as follows:
we retained genotype calls if they had a minimum coverage depth of 11, and discarded loci that had a
minor allele count lower than 3 or a genotype call in fewer than 75% of all individuals. We further
discarded five individuals which had more than 60% missing genotypes. Following this filtering
approach, we obtained 8764 SNPs in 517 individuals for further analyses. Because the adults had on
average more missing data than the hatchlings, we generated a second dataset comprising only the 82
wild-collected adults, filtering with the same criteria as above. This second dataset contained 4943 SNPs
and was used to infer the number of genetic lineages present. All downstream analyses were conducted

in R 4.0.3 (R Core Team, 2021). We read the vcf file using the R package vcfR v1.8.0 (Knaus &
Grünwald, 2017).

We amplified and sequenced an 814 bp fragment of the COI mitochondrial gene for all adults in our
dataset, plus reference *T. douglasi* and *T. poppense* from other populations (Table S2) using a nested
PCR. We first amplified DNA with primers 3'-TCCAATGCACTAATCTGCCATATTA-5' and 3'-
GGAACNGGATGAACAGTTTACCCNCC-5'. We then amplified a subset of this fragment using the
former and 3'-CAACATTTATTTTGATTTTTGG-5'. All primer sequences were obtained from Simon
et al., (1994). All reactions were conducted in 20 μL of Promega Buffer 1X containing 0.3 μM of each
primer, 0.2 mM of each dNTP and 0.02 U/μL G2 Taq polymerase (Promega). All cycles consisted of
5mn at 94°C followed by 20 (first PCR) and 35 (nested PCR) cycles of 30 s at 95°C, 1 mn at 50°C, 1
mn at 72°C, and a final extension at 72°C for 10 mn. The PCR products were sequenced both ways at
Microsynth (Switzerland), and we generated the consensus from the forward and reverse sequences
using BioEdit.

Population structure and phylogeny
186
To identify the number of genetic lineages present across the two transects, we conducted a clustering
analysis on the adults using faststructure v1.0 (Raj et al., 2014), with K ranging from 1 to 10. We then
used the chooseK.py script to select the value of K that maximises marginal likelihood.

We also wanted to assess whether individuals collected along the two transects belonged to known
lineages of the parthenogenetic species *T. douglasi* or the sexual species *T. poppense*. Although we did
not expect populations of the sexual species *T. poppense* in the surveyed geographic range (Law &
Crespi 2002), we could not exclude the presence of geographically isolated *T. poppense* populations.
This is because *T. douglasi* is very closely related *T. poppense* and females of these two species cannot
be distinguished morphologically (Vickery & Sandoval, 1999). We therefore used reference individuals
for species assignments. These reference individuals, five *T. douglasi* and 26 *T. poppense* individuals
from different populations (Table S2) were identified in previous studies (Jaron et al., 2022; Schwander
et al., 2011), with DNA extractions still available in the lab. We used these 31 reference individuals to
study their phylogenetic relationship with the genetic lineages identified in our newly sampled
populations for mitochondrial and nuclear (RADseq) markers. We generated RADseq and COI data for
these individuals with the same protocols as described above. For the RADseq data, we retained
genotype calls with a minimum depth of 8 instead of 11, as allelic dropout would cause less bias for
phylogenetic reconstruction than for inference of reproductive modes (largely based on heterozygosity
estimations, see below). We discarded loci that had a minor allele count lower than three and were
genotyped in less than 80% of all individuals. This dataset contained 16391 sites. The COI dataset
contained 159 polymorphic sites. Note that some individuals had multiple COI copies (as indicated by

double peaks in the sequencing trace), as expected for the presence of nuclear copies of mitochondrial genes or heteroplasmy. In these cases, we called the allele corresponding to the largest peak, and used an ambiguous genotype following the IUPAC code when both peaks were equally large. We added the COI sequence from one individual of *T. cristinae* as an outgroup (EU251511.1; (Schwander et al., 2011). We then aligned the sequences using the MUSCLE algorithm (Edgar, 2004). We reconstructed separate maximum likelihood phylogenies for the nuclear and mitochondrial data, with iqtree2 (Minh et al., 2020), using ModelFinder Plus to select the best substitution model in each case. We used the cophylo function of the R package phytools (Revell, 2012) to plot them as a cophylogeny.

<u>Reproductive mode analyses</u>

For individuals of the Manchester and Orr transects, we calculated heterozygosity as the proportion of the SNPs that were heterozygous. Note that because we only retained polymorphic positions in our dataset, this measure overestimates genomic heterozygosity and is hereafter referred to as "relative heterozygosity". We observed that relative heterozygosity increased linearly with the proportion of missing data (Figure S1). This increased heterozygosity is likely due to merged paralogs, which are enriched in individuals with more missing data as paralogues have higher coverage than single-copy loci and are therefore preferentially retained (Supplementary Material). We therefore corrected relative heterozygosity to remove the effect of missing data by using the residuals of the linear regression and adding the intercept value (Figure S1).

To assess whether hatchlings from eggs laid after mating were produced via sex or parthenogenesis, we used two complementary measures. For the first measure, we looked for genotypes in hatchlings that cannot be produced via parthenogenesis. At loci where the mother is homozygous, offspring can only be heterozygous if they were produced from a fertilized egg (in the absence of genotyping errors; see also Brown et al., 2021). For each offspring, we therefore computed the proportion of heterozygous loci that had a homozygous genotype in the mother. Loci at which either the mother or the offspring were not genotyped were excluded.

For the second measure to assess whether hatchlings were produced via sex or parthenogenesis, we compared relative heterozygosity between mothers and their offspring. We expected heterozygosity levels of hatchlings produced via sex to be similar to that of the mother, or higher in hatchlings sired by a male originating from a different lineage than the mother. On the other hand, we expected heterozygosity levels of parthenogenetically produced hatchlings to be lower than that of their mother. A decrease is expected because facultative parthenogenesis generally occurs via automixis (Suomalainen et al., 1987). Under automixis, meiotic divisions occur and diploidy is restored secondarily via fusion or duplication of meiotic products. Similar to selfing, automixis can result in a loss of heterozygosity between mothers and offspring (Pearcy et al., 2006; Suomalainen et al., 1987).

However, heterozygosity losses can only be observed at sites where the mother was heterozygous, and
where heterozygosity losses are not mechanistically constrained or removed by selection (Jaron et al.,
2022). The extent of the heterozygosity loss then depends on the type of automixis and the amount of
recombination (Pearcy et al 2006, Suomalainen et al 1987). In the most extreme case (gamete
duplication) a haploid gamete duplicates to produce a fully homozygous zygote. Other forms of
automixis (e.g., terminal or central fusion) result in a partial loss of heterozygosity. In these cases,
offspring heterozygosity depends on the heterozygosity of the mother, and on the position and frequency
of recombination events. By combining the two heterozygosity-based measures, we were able to
unambiguously assign each of the 365 genotyped hatchlings from eggs produced after mating to either
sexual or parthenogenetic offspring (see Results).

We then asked whether females from different genetic lineages used different reproductive strategies.
To answer this question, we tested whether there were significant differences between lineages in (i)
hatching success of eggs laid before mating, (ii) hatching success of eggs laid after mating, and (iii) the
proportion of post-mating eggs that were fertilized (which measures the propensity to reproduce via sex)
using binomial Generalized Linear Models (GLMs), tested their significance using the Anova function
of the R package car v3.0-10 (Fox & Weisberg, 2019) and performed multiple comparisons using the
glht function implemented in the R package multcomp v1.4-16 (Hothorn et al., 2008). We also looked
for differences in hatching success of eggs before vs after mating within each lineage using binomial
GLMMs implemented in the R package lme4 (Bates et al., 2015), using the mother identity as random
factor.

In order to test whether variable population sex ratios along the Orr and Manchester transects were
generated by sex ratio distortion, we quantified the sex ratio in sexually produced offspring. Because
female and male hatchlings cannot be distinguished morphologically, we sexed offspring genetically by
combining heterozygosity at X-linked loci and the read depth ratio between the X-linked and autosomal
loci. Because *Timema* males have only one copy of the X chromosome and no Y (XX:XO sex
determination; Schwander & Crespi, 2009), there should be no heterozygosity for X-linked loci.
Furthermore, X-linked loci should have half the coverage compared to autosomal loci in males, but
similar coverage in females. To sex sexually produced offspring via these approaches, we first had to
identify X-linked scaffolds in the *T. douglasi* reference genome (Jaron et al., 2022).

To identify X-linked scaffolds we used a combination of available genomic data from *T. douglasi*
females (Bioproject accession number: PRJNA670663) in addition to whole-genome sequencing of two
male *T. douglasi*. DNA extractions were done on adult male carcasses using the Qiagen Mag Attract
HMW DNA kit following the manufacturer instructions. Sequencing libraries were generated for each
male using a TruSeq DNA nano prep kit (550bp insert size). Libraries were then sequenced (150b,

paired-end) using an Illumina HiSeq 4000 at the Lausanne Genomic Technologies Facility. Reads were

trimmed before mapping using Trimmomatic v0.36 (Bolger et al., 2014) to remove adapter and low-

quality sequences (options: ILLUMINACLIP:3:25:6 LEADING:9 TRAILING:9

SLIDINGWINDOW:4:15 MINLEN:90). Reads from each individual were then mapped to the reference

genome using BWA-MEM v0.7.15 (Li & Durbin, 2010). Multi-mapping and poor quality alignments

were filtered (removing reads with XA:Z or SA:Z tags or a mapq < 30). PCR duplicates were removed

with Picard (v. 2.9.0) (http://broadinstitute.github.io/picard/). Coverage was then estimated for all

scaffolds at least 1000 bp in length using BEDTools v2.26.0 (Quinlan & Hall, 2010). To compare

coverage between males and females, coverage was first summed up for all male and all female libraries

per scaffold. Male and female coverage was then normalised by modal coverage to adjust for differences

in overall coverage. X-linked scaffolds were then identified using the log2 ratio of male to female

coverage. Scaffolds were classified as X-linked if the $\log_2$ ratio of male to female coverage was within

0.1 of the value of the X-linked peak (Figure S3), and as autosomal otherwise.

By distinguishing between autosomal and X-linked loci, we then sexed sexually produced offspring

from our controlled crosses using the RADseq data. We first standardised read depth per individual,

dividing the depth value at each locus by the average depth for that individual. We then computed the

X to autosomes depth ratio by dividing average standardised depth of loci on the X chromosome by that

of autosomal loci (see supplementary material). Using this depth ratio in combination with

heterozygosity on the the X chromosome, we were able to sex 176 out of the 211 sexually produced

offspring (see Figure S5; the remaining 35 offspring had insufficient sequencing coverage for these

analyses). We tested for deviations from a 50:50 sex ratio among sexually produced offspring overall

using a chi-squared test. We further investigated whether there were deviations within each family. For

this, we looked whether the 95% binomial proportion confidence interval of the observed sex ratio

overlapped with the expected 50:50 sex ratio. In addition, we conducted Fisher's exact tests for each

family.

Results

We found that adults from geographically close populations differed strongly in their sex ratio, with

significant sex ratio variation among populations overall (Figures 1A, 1C; Manchester: Likelihood ratio

$\chi^2 = 456.49$, df = 11, $p < 2$ x 10^{-16}; Orr: Likelihood ratio $\chi^2 = 106.46$, df = 11, $p < 2$ x 10^{-16}). The Orr

transect consisted of a mosaic of female-biased and female-only populations. Five populations (Orr 1,

5, 8, 9 and 17) had between 77.8% and 91.7% of females, while the others had only females (Figure

1A). By contrast, we found an abrupt transition from roughly even to strongly female-biased or female-

only sex ratios along the Manchester transect. Sex ratios in populations 1 to 7 ranged from 43.2% to
54.7% of females, while populations 8 to 12 consisted of at least 93.5% females (Figure 1C).

We then investigated whether the field-collected individuals belonged to different genetic lineages using
a clustering analysis. The best supported number of lineages among all genotyped field-collected
individuals was three. We found a gradient of genotype incidences along the transects. The first lineage
(blue in Figure 1B) was mostly found along the Orr transect, with only one individual in Manchester 10,
and is hereafter referred to as the "Orr lineage". The second lineage (yellow in Figure 1B) was found
mostly in the eastern section of Manchester and is hereafter referred to as the "eastern Manchester
lineage". Finally, the third lineage (red in Figure 1B) was mainly found in the western section of
Manchester and is hereafter referred to as the "western Manchester lineage". We found little evidence
for admixture between lineages in 14 out of the 15 populations tested, with only one female from Orr
11 (17-2162) appearing to be a hybrid, perhaps triploid, between the three lineages (Figure S2). In one
population however, Manchester 2, all 5 genotyped individuals appeared to be admixed, with
approximately equal contributions of the eastern and western Manchester lineages. These results were
robust with varying numbers of clusters (k in faststructure), indicating that these were most likely
hybrids and not a fourth lineage (Figure S2).

In our nuclear phylogeny, all individuals collected along the two transects formed a monophyletic clade
with the parthenogenetic *T. douglasi* reference individuals, indicating that all three lineages belong to
T. douglasi. All *T. poppense* reference individuals formed a second, monophyletic clade (Figure 1E).
The mitochondrial phylogeny also showed two monophyletic groups. One contained mostly *T.*
poppense, with five *T. douglasi* individuals (including four from our transects). The second contained
mostly *T. douglasi*, with three *T. poppense* individuals (Figure 1F). Individuals belonging to the eastern
Manchester lineage formed two distinct mitochondrial subclades, which were not segregated
geographically.

A: Orr transect

17
N = 49

10
N = 96

9
N = 18

39°13'N
123°25'W

11
N = 10

8
N = 70

7
N = 26

4
N = 204

2
N = 3

1 km

6
N = 165

5
N = 123

3
N = 202

1
N = 108

= ♀   = ♂

B

Population   Individual ID   Assignment probability
0.0   0.5   1.0

1   17–1913
     17–2076
     17–2077
3   17–2087
4   17–2114
5   17–1743
8   17–1925
     17–1927
     17–2135
     17–2138
11   17–2161
     17–2162

C: Manchester transect

12
N = 270

11
N = 38

5
N = 67

4
N = 170

6
N = 49

8
N = 29

10
N = 62

1
N = 148

7
N = 16

9
N = 139

2
N = 112

3
N = 95

1 km

38°58'N
123°28'W

D

1   17–1605
     17–1607
     17–1613
     17–1618
     17–1623
     17–1624
     17–1625
     17–1628
     17–1629
     17–1634
     17–1636
     17–1639
     17–1642
     17–1770
     17–2195

2   17–1654
     17–1663
     17–1664
     17–1665
     17–1669

3   17–1672
     17–1675
     17–1679
     17–1701
     17–1703
     17–1704
     17–1706

4   17–1716
     17–1719
     17–1736
     17–2005
     17–2009
     17–2011
     17–2013

5   17–1790
     17–1793
     17–1796
     17–1798
     17–1800
     17–1801
     17–1806
     17–2201

6   17–1758
     17–1760
     17–1765
     17–1931
     17–2037

7   17–2041
     17–2042
     17–2165
     17–2191
     17–2193
     17–2194
     17–2199

10   17–1810
     17–1811

12   17–1857
     17–1935
     17–1937

E: Nuclear phylogeny

F: Mitochondrial phylogeny

0.1

0.05

● = Reference *T. poppense* (sexual)   ● = Reference *T. douglasi* (parthenogenetic)

Figure 1 Sex ratios in the sampled populations, genetic structure and species identification of the collected individuals. **A** and **C**: Schematic maps of the Orr and Manchester transects, with host plant patches represented in green along the main road. Pie charts represent the sex ratio of individuals collected in the field. Crosses indicate surveyed host plant patches with no *Timema* present (locations 12-16 along the Orr transect). **B** and **D**: Assignment probabilities to each of the 3 genetic lineages for each genotyped individual, split by transect. **E**: ML nuclear phylogeny (16391 SNPs) of the genotyped field-collected individuals plus 26 *T. poppense* (sexual) and five *T. douglasi* (parthenogenetic) reference individuals to infer species assignments for individuals from the transects. **F**: ML mitochondrial phylogeny of the same individuals based on 159 polymorphic sites of the COI mitochondrial gene. One individual of *T. cristinae* was used to root the tree (not displayed). Tip colour refers to genetic lineage (individuals from Orr and Manchester transects) or reference individuals used for species assignment. Individuals denoted in bold with a star had double peaks in their COI sequence and the allele corresponding to the highest peak was retained. Timema photos taken by © Bart Zijlstra - www.bartzijlstra.com

We isolated 352 females to characterise their reproductive mode and assess the hatching success of the eggs they laid as virgins. 265 of these females laid at least five eggs as virgins. At least one unfertilized egg hatched from the clutches produced by 234 (88.3%) of them, indicating that they were able to reproduce via parthenogenesis. In addition, females capable of parthenogenesis were found in all populations tested. The hatching success of unfertilized eggs varied between 3.3 and 100% (Figure 2A).

The hatching success of unfertilized eggs from the Orr lineage was within the range of obligate asexual *Timema* species, while for the two Manchester lineages, it was intermediate between obligate sexuals and asexuals (Figure 2B).

Figure 2 Parthenogenesis capacity in field-collected females from different populations along the Manchester and Orr transects (**A**) and females from different obligate sexual and obligate asexual species (**B**), measured by the hatching success of unfertilized eggs. Dot size in **A** is proportional to the number of unfertilized eggs laid by each female. In **B**, N refers to the number of females for which hatching success was measured.

In order to gain insights into the mechanisms of automictic parthenogenesis in females from the Orr and Manchester transects, we compared relative heterozygosity in mothers and their offspring produced from eggs laid prior to mating. Heterozygosity losses can only be observed at sites where the mother was heterozygous, and where heterozygosity losses are not mechanistically constrained or removed by selection (Jaron et al., 2022). However, if heterozygosity loss does occur, the extent of it can inform on the type of automixis (Pearcy et al 2006). We found extensive loss of heterozygosity or extremely low

relative heterozygosity in all offspring from eggs laid prior to mating (Figure 3), as would be expected from gamete duplication or terminal fusion without recombination. With one exception, relative heterozygosity in these offspring amounted to 2 - 3% and was not correlated with their mothers' heterozygosity (as would be expected i.e., under central or terminal fusion automixis with recombination). In addition, most, if not all of the heterozygosity observed in hatchlings is caused by false positives (see Supplementary Material), meaning that these hatchlings are most likely completely homozygous. Only one parthenogenetic hatchling from female 17-1793 retained some heterozygosity (Figure 3), which could suggest that it was produced by a different parthenogenesis mechanism than the other individuals.

Figure 3 Relative heterozygosity of mothers (circles coloured as in Figure 1) and their offspring produced from unfertilized eggs laid before mating (black crosses, two to three per mother), and of two reference parthenogenetic individuals (Tdi_04 and Tdi_05) from Jaron et al. (2022).Note the much higher heterozygosity in one hatchling from family 17-1793, indicating that it could have been produced via a different mechanism of parthenogenesis.

We were able to unambiguously infer whether each offspring from eggs laid after mating was produced via sex or parthenogenesis by using a combination of relative heterozygosity and proportion of genotype transitions from homozygous in the mother to heterozygous in the offspring (Figure 4; see also Supplementary Material). These inferences revealed that 211 (58.3%) of offspring were produced by sex, and 151 (41.7%) were produced via parthenogenesis.

We then looked at differences in the parthenogenetic capacity or the propensity to use sex following

mating between females from the Orr, western and eastern Manchester lineages. All 32 females for

which we were able to infer the lineage via genotyping were capable of parthenogenesis, but with

varying efficiency (Figure 5A). Note that one of these females (17-1937 from the eastern Manchester

lineage) did not produce any offspring from the 13 eggs she laid before mating, but she was capable of

parthenogenesis since all three offspring produced from the 14 eggs laid after mating were produced via

parthenogenesis. Hatching success of eggs laid before mating was on average higher for females of the

Orr lineage (67%) than of the two other lineages (western Manchester: 42%, $z = 2.51$, $p = 0.032$; eastern

Manchester: 33%, $z = 3.53$, $p = 0.001$; Figure 5A). There was no significant difference between the

lineages in hatching success of eggs laid after mating ($\chi^2 = 0.003$, $df = 2$, $p = 0.99$; Figure 5B). Mating

influenced the egg hatching success in all three lineages. In the eastern and western Manchester lineages,

eggs laid after mating had a significantly higher hatching success than those laid by virgin females

(western Manchester: 55% vs 42%, $z = 2.0$, $p = 0.04$; eastern Manchester: 55% vs 33%, $z = 4.7$, $p =$

$2.7*10^{-6}$). On the other hand, in the Orr lineage, eggs laid after mating hatched at a lower rate than those

laid before mating (56% vs 69%, $z = -4.7$, $p = 2.7*10^{-6}$). We also found that the proportion of eggs laid

after mating that were actually fertilized was highly variable among females. Twelve females fertilized

between 80% and 100% of their eggs laid after mating, while eleven fertilized none of them. Only five

fertilized an intermediate proportion (14% to 50%). There were significant differences in the proportion

of fertilized eggs between lineages. Females from the Orr lineage fertilized on average fewer eggs (10%)

than those from the eastern and western Manchester lineages (79%, $t = 4.0$, $p = 0.0002$ and 95%, $t = 3.5$, $p = 0.001$ respectively). All females from the western Manchester lineage fertilized more than 80% of their post-mating eggs, while seven out of 10 females from the Orr lineage fertilized none of them. Finally, the eastern Manchester lineage consisted mostly of females with either a very high or very low fertilisation frequency (Figure 5C). The variation in fertilization frequency was not explained by the two mitochondrial subclades. The putatively triploid hybrid female (17-2162) was also capable of facultative parthenogenesis: a large portion of eggs laid prior to mating hatched (24 out of 27), and out of the 17 genotyped hatchlings from the eggs she laid after mating, seven were produced via sex and 10 via parthenogenesis.

Figure 5: Reproductive mode differences between females of the three genetic lineages (distinguished by colours as in Figure 1). **A**: Hatching success of eggs laid prior to mating, which measures the capacity to do parthenogenesis. Dot size proportional to the number of eggs laid before mating. **B**: Hatching success of eggs laid after mating. Dot size proportional to the number of eggs laid after mating **C**: The proportion of sexually produced hatchlings that originated from eggs laid after mating, which measures the propensity to reproduce via sex. Dot size proportional to the number of hatchlings genotyped to assess whether they were produced via sex or parthenogenesis. In all panels: the black squares and bars represent the weighted average per lineage +/- standard error. Different letters indicate significantly different groups ($p < 0.05$ in multiple comparison tests after binomial GLMs).

Finally, we tested whether sex ratio distortion could be contributing to female-biased sex ratios by looking at the sex ratio of sexually produced offspring. A driving X chromosome should lead to an excess of female offspring. Across all families, there were 81 female and 95 male offspring, providing

445 no evidence for biased sex ratios ($\chi^2 = 1.1$, $p = 0.29$). The same was observed within families (all $p >$

446 0.5; Figure S6).

Discussion

How transitions between fundamentally different reproductive modes – such as sex and obligate

parthenogenesis – can occur, is unclear. Transitions might be abrupt, with a direct switch from sex to

obligate parthenogenesis within one generation. Alternatively, transitions might be gradual and involve

a period where individuals are capable of facultative parthenogenesis (Schwander & Crespi, 2009

Neiman et al., 2014). In this study, we discovered high frequencies of males in populations of *Timema*

douglasi, a species previously believed to reproduce solely via obligate parthenogenesis. These

populations were characterized by variable sex ratios, which were caused by different frequencies of

sexual versus parthenogenetic reproduction among facultative parthenogens, and not by sex ratio

distortion. These findings raise the possibility that obligate parthenogenesis, which evolved repeatedly

in the genus *Timema*, may generally derive from facultative parthenogenesis, with facultative

parthenogenesis being a transient state. Two genetically differentiated lineages of *T. douglasi*

reproduced through true facultative parthenogenesis, with good hatching success of unfertilized eggs

but a high proportion of sexually produced offspring when mated. They were found geographically close

to a third lineage of the same species (Orr), in which females had a high capacity for parthenogenesis

and mostly did not fertilize their eggs when mated. Furthermore, eggs produced after mating by females

of this lineage suffered from a lower hatching success, suggesting that fertilisation of eggs that would

normally develop into parthenogenetic offspring can sometimes result in developmental failure. These

results may point to a tradeoff between sexual and parthenogenetic pathways as was previously

suggested to be the case in sexual *Timema* populations: the ability for spontaneous parthenogenesis

likely reduces hatching success of fertilized eggs (Schwander et al., 2010). However, lower hatching

success of post-mating eggs could also be due to females being older. This effect would be compensated

by the positive effect of fertilization on egg hatching success in both Manchester lineages.

Females of the two facultatively parthenogenetic lineages from the Manchester transect had a lower

capacity for parthenogenesis than those of the largely obligate one and featured different propensities to

reproduce sexually following mating. Females of the first lineage (western Manchester) generally

fertilised all their eggs when mated, which resulted in higher hatching success than under

parthenogenesis. Females of the second lineage (eastern Manchester) were more variable.

Approximately half (8 out of 15) of them reproduced the same way as the western Manchester lineage:

using mainly sex when mated but being able to reproduce via parthenogenesis. Five females reproduced

only via parthenogenesis, similarly to females of the Orr lineage. The two remaining females seemed to

use both sex and parthenogenesis at approximately equal frequencies (facultative parthenogenesis), but
genotyping larger clutches would be needed to assess whether this was indeed the case.

The mechanisms that maintain reproductive polymorphism (facultative and largely obligate
parthenogenesis) within populations, and those that cause the sharp transition from equal to strongly
female-biased sex ratios over a few hundred meters along the Manchester transect remain an open
question. Since more than half of the females from the eastern Manchester lineage were able to
reproduce via sex and fertilized most or all of their eggs following mating, a male immigrating from a
nearby sexual population would be expected to have extremely high reproductive success. This should
then lead to an increase in male frequency in the next generation. One possible explanation for the
maintenance of different sex ratios between neighbouring populations is that dispersal is very limited in
these wingless insects (Sandoval, 2000), and that it is just a matter of time until the female-biased
Manchester populations shift towards more equal sex ratios. However, variable selection pressures along
the transect could also play a role, favouring respectively sexual or parthenogenetic reproduction (Burke
& Bonduriansky, 2018). This could help explain why males, even though present in small numbers in
the strongly female-biased populations (such as Manchester 12 with <1% males), have not triggered sex
ratio shifts. Identifying such selection pressures would constitute a major leap towards understanding
the costs and benefits of different reproductive modes in natural populations.

Our comparison of heterozygous genotypes in mothers and their parthenogenetically produced offspring
also provides insights into the mechanisms of parthenogenesis and the evolution of obligate
parthenogenesis in *Timema*. Relative heterozygosity in all but two parthenogenetic offspring (out of 228
produced before or after mating) was around 2.5%, regardless of their mothers' heterozygosity. Given
the 2.5% heterozygosity is most likely generated by structural variation between individuals and
divergence between paralogs (Supplementary Material), the main mechanism of parthenogenesis in
Timema thus appears to generate genome-wide homozygosity in a single generation. This would be
expected under specific forms of automixis, notably under gamete duplication or fusion of non-
recombined sister chromatids (i.e., terminal fusion without recombination). The two remaining out of
228 parthenogenetic offspring (one originating from an unfertilized egg laid prior to mating, the other
from an unfertilized egg laid after mating; see Figure 4), produced by different females, had somewhat
higher heterozygosity. This suggests that some females are able to reproduce via two different
parthenogenesis mechanisms in addition to sex, as was described for spontaneous parthenogenesis in
sexual *Daphnia* (Svendsen et al., 2015). Automixis with "complete" heterozygosity loss in a single
generation is also the proposed mechanism of parthenogenesis in obligate asexual *Timema* species
(Jaron et al., 2022) and of spontaneous parthenogenesis in sexual *Timema* species (Schwander & Crespi,
2009). In *Timema*, obligate parthenogenesis thus likely evolved via a gradual increase of the capacity
for parthenogenesis, with a conserved proximate mechanism.

Our finding of facultative parthenogenetic lineages of *T. douglasi* questions the directionality of

transitions between reproductive modes in this genus and beyond. So far, sex with low levels of

tychoparthenogenesis was believed to be the ancestral state in *Timema*, followed by repeated

transitions to obligate parthenogenesis (Schwander & Crespi, 2009). However, our results open the

possibility that facultative parthenogenesis be an ancestral state in this genus, and that the different

lineages have transitioned towards more obligate strategies. This would require that obligate

parthenogenesis has evolved several times in the *T. poppense/douglasi* species complex. Phylogenetic

analyses of the *Timema* genus revealed that *T. douglasi* consisted of multiple lineages that

independently derived from the sexual species *T. poppense* (Schwander et al., 2011). Repeated

transitions towards parthenogenesis are not surprising if the ancestor of these species was already

capable of reproducing via facultative or spontaneous parthenogenesis, and if selection has the

opportunity to increase parthenogenetic capacity, for example due to mate limitation (Schwander et

al., 2010).

Alternatively, the facultative parthenogenetic *T. douglasi* lineages could represent a re-evolution of sex.

Sex could in theory reappear in all-female lineages via accidental male production. Rare male production

in species with XX/X0 sex determination such as *Timema* can occur via aneuploidy at the X

chromosome, because the resulting X0 individuals would develop into males (Pijnacker & Ferwerda,

1980). However, such a re-expression of the sexual pathway would require the molecular bases

underlying the sexual functions of both sexes to be maintained. Re-expression of sex is therefore

possible at least shortly after the transition to parthenogenesis, which may also have been the case in the

stick insects *Clitarchus hookeri* (Morgan-Richards et al., 2019) and *Bacillus rossius* (de Vichet, 1944).

Occasional male production and re-expression of the sexual pathway in otherwise obligate parthenogens

paves the way for cryptic sex, which could contribute to explaining their long-term persistence (Freitas

et al., 2023).

Finally, gene flow between sexual and parthenogenetic lineages could contribute to frequent transitions

between reproductive modes. One such example is contagious parthenogenesis, whereby rare males

from otherwise parthenogenetic lineages transmit the capacity for parthenogenesis to a related sexual

lineage via hybridization. This has been documented in some aphids (Delmotte et al., 2001), wasps

(Sandrock & Vorburger, 2011) and *Daphnia* (Innes & Hebert, 1988). Conversely, gene flow from a

sexual to mostly but not completely parthenogenetic species could in theory result in an increase of the

frequency of sex in the latter. Occasional gene flow between *T. poppense* and *T. douglasi* is suggested

by the mismatches we observed between nuclear and mitochondrial genomes in about 10% of our

individuals. This could be caused by rare hybridization events between the two species in both directions

and contribute to the maintenance of different reproductive modes within populations. Formal tests of

gene flow between *T. poppense* and *T. douglasi* would shed light onto this question.

More generally, the faculty for occasional and facultative sex in *T. douglasi*, a species previously

believed to be obligately parthenogenetic, adds to recent re-evaluation of the obligate status of some

parthenogenetic species (Boyer et al., 2021; Kuhn et al., 2021; Laine et al., 2022). Many such "ancient

asexuals" could in fact have retained the capacity for occasional, cryptic sex (Schurko et al., 2009). They

would thus represent extremes of a continuum from mostly sexual to mostly parthenogenetic

reproduction. On the other end of this continuum, many taxa that were thought to reproduce solely via

sex were recently found to be capable of parthenogenesis, though often at anecdotal frequency and with

limited success (e.g. Booth et al., 2012; Fields et al., 2015; Groot et al., 2003; Ryder et al., 2022, but see

Kratochvíl et al., 2020).

Whether a transient facultative state is a frequent feature in transitions between extremes of the

reproductive continuum still remains to be investigated. Still, the factors that would favour obligate over

more facultative strategies remain elusive. Facultative parthenogenesis is believed to be an efficient

reproductive mode, combining the "best of both worlds" – the long-term advantages of recombination

and segregation along with the short-term advantages of asexuality. However, the loss of sex in

facultative parthenogens could in theory be driven by sexual conflict (if mating always reduces female

fitness; (Burke & Bonduriansky, 2017). Alternatively, the lower parthenogenesis capacity observed in

the facultatively asexual lineages than in both the more obligate lineage (Orr) and the obligate asexual

species, as well as interspecific comparisons in facultatively parthenogenetic mayflies (Liegeois et al.,

2021) suggest that the efficiency of sex and parthenogenesis are traded-off against each other, consistent

with the "jack of all trades, master of none" hypothesis. In this case, whether obligate strategies are

likely to replace facultative ones will depend on local ecological conditions favouring sex or

parthenogenesis, and on the fluctuations of such conditions.

In conclusion, we discovered the first case of facultative parthenogenesis in *Timema* stick insects, a

genus where obligate parthenogenesis has evolved repeatedly from a sexual ancestor, with an apparently

conserved proximate mechanism of parthenogenesis. We found three genetic lineages that differed both

in their capacity to reproduce via parthenogenesis and in their propensity to reproduce sexually. It is

thus far unclear whether these populations reverted to sex starting from a largely asexual ancestor, or

they are sexual relics in the largely asexual *T. douglasi*, where all other populations have transitioned

to obligate parthenogenesis already. The latter would suggest that transitions to asexuality were

numerous during the evolution of this species, and that asexual populations almost always prevailed on

sexuals. In that case, we could be witnessing an ongoing transition to asexuality via facultative

parthenogenesis. Why facultative parthenogenesis would be a transient stage in *Timema* and which

factors would favour a transition to obligate parthenogenesis would thus be exciting foci for future
research.

Acknowledgements

592 We thank Ian S. Ford and Armand Yazdani for their great help in the field, Karine Légeret for her
593 assistance in the daily monitoring of egg laying and hatching in the lab, and Zoé Dumas and Marjorie
594 Labédan for wet lab help. Comments by three anonymous reviewers improved the manuscript. Most
595 analyses were run using the computing infrastructure at the University of Lausanne maintained by the
596 former Vital-IT platform of the SIB Swiss Institute of Bioinformatics (SIB) and the DCSR.
597

Funding

599 We would like to acknowledge funding from the European Research Council (Consolidator Grant No
600 Sex No Conflict), Swiss FNS grant 31003A_182495, and the University of Lausanne.
601

Conflict of interest disclosure

603 The authors declare they have no conflict of interest relating to the content of this article.
604

Data and code availability

606 All demultiplexed RAD-seq reads have been deposited in NCBI's sequence read archive under
607 BioProject ID PRJNA804475 and whole genome sequence reads of *T. douglasi* males under
608 BioProject ID PRJNA808673. COI sequences have been deposited on GenBank under accessions
609 OP379833 – OP379903. The code used to run the analyses presented in this article is available on
610 GitHub under https://github.com/glavanc1/Timema_facultative_partheno.

References

Arbuthnott, D., Crespi, B. J., & Schwander, T. (2015). Female Stick Insects Mate Multiply to Find
Compatible Mates. *The American Naturalist*, *186*(4), 519–530. https://doi.org/10.1086/682675

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

Bell, G. (1982). *The Masterpiece of Nature: The Evolution and Genetics of Sexuality*. University of
California Press.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics*, *30*(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Booth, W., Smith, C. F., Eskridge, P. H., Hoss, S. K., Mendelson, J. R., & Schuett, G. W. (2012).
Facultative parthenogenesis discovered in wild vertebrates. *Biology Letters*, *8*(6), 983–985.
https://doi.org/10.1098/rsbl.2012.0666

Boyer, L., Jabbour-Zahab, R., Mosna, M., Haag, C. R., & Lenormand, T. (2021). Not so clonal
asexuals: Unraveling the secret sex life of *Artemia parthenogenetica*. *Evolution Letters*, *5*(2),
164–174. https://doi.org/10.1002/evl3.216

Brelsford, A., Dufresnes, C., & Perrin, N. (2016). High-density sex-specific linkage maps of a
European tree frog (Hyla arborea) identify the sex chromosome without information on offspring
sex. *Heredity*, *116*(2), 177–181. https://doi.org/10.1038/hdy.2015.83

Brown, T. A., Tsurusaki, N., & Burns, M. (2021). Genomic Determination of Reproductive Mode in
Facultatively Parthenogenetic Opiliones. *Journal of Heredity*, *112*(1), 34–44.
https://doi.org/10.1093/jhered/esaa045

Burke, N. W., & Bonduriansky, R. (2017). Sexual Conflict, Facultative Asexuality, and the True
Paradox of Sex. *Trends in Ecology & Evolution*, *32*(9), 646–652.
https://doi.org/10.1016/j.tree.2017.06.002

Burke, N. W., & Bonduriansky, R. (2018). The geography of sex: sexual conflict, environmental
gradients and local loss of sex in facultatively parthenogenetic animals. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *373*(1757), 20170422.
https://doi.org/10.1098/rstb.2017.0422

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an analysis
tool set for population genomics. *Molecular Ecology*, *22*(11), 3124–3140.
https://doi.org/10.1111/mec.12354

Cuellar, O. (1974). On the origin of parthenogenesis in vertebrates: the cytogenetic factors. *The*
American Naturalist, *108*(963), 625–648.

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R. E.,
Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., & Durbin, R. (2011). The variant call format
and VCFtools. *Bioinformatics*, *27*(15), 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

de Vichet, G. (1944). Découverte dans le sud de la France d'une station de males de Bacillus rossii
(Fabr.) (Orthoptère, Phasmidae). In *Publications de la Société Linnéenne de Lyon* (Vol. 13, Issue
6, pp. 92–94).

Delmotte, F., Leterme, N., Bonhomme, J., Rispe, C., & Simon, J.-C. (2001). Multiple routes to
asexuality in an aphid species. *Proceedings of the Royal Society of London. Series B: Biological*
Sciences, *268*(1483), 2291–2299. https://doi.org/10.1098/rspb.2001.1778

Drosopoulos, S. (1978). Laboratory Synthesis of a Pseudogamous Triploid "Species" of the Genus
Muellerianella (Homoptera, Delphacidae). *Evolution*, *32*(4), 916.
https://doi.org/10.2307/2407505

D'Souza, T. G., & Michiels, N. K. (2010). The Costs and Benefits of Occasional Sex: Theoretical
Predictions and a Case Study. *Journal of Heredity*, *101*(Supplement 1), S34–S41.
https://doi.org/10.1093/jhered/esq005

Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Research, *32*(5), 1792–1797. https://doi.org/10.1093/nar/gkh340

Fields, A. T., Feldheim, K. A., Poulakis, G. R., & Chapman, D. D. (2015). Facultative parthenogenesis
in a critically endangered wild vertebrate. *Current Biology*, *25*(11), R446–R447.
https://doi.org/10.1016/j.cub.2015.04.018

Fox, J., & Weisberg, S. (2019). *An R Companion to Applied Regression* (Third edition). Sage.

Freitas, S., Parker, D. J., Labédan, M., Dumas, Z., & Schwander, T. (2023). Evidence for cryptic gene
flow in parthenogenetic stick insects of the genus Timema. *BioRxiv*,
https://www.biorxiv.org/content/10.1101/2023.01.21.525009v2.

Groot, T. V. M., Bruins, E., & Breeuwer, J. A. J. (2003). Molecular genetic evidence for
parthenogenesis in the Burmese python, Python molurus bivittatus. *Heredity*, *90*(2), 130–135.
https://doi.org/10.1038/sj.hdy.6800210

Helleu, Q., Gérard, P. R., & Montchamp-Moreau, C. (2015). Sex Chromosome Drive. *Cold Spring
Harbor Perspectives in Biology*, *7*(2), a017616. https://doi.org/10.1101/cshperspect.a017616

Hothorn, T., Bretz, F., & Westfall, P. (2008). Simultaneous Inference in General Parametric Models.
Biometrical Journal, *50*(3), 346–363. https://doi.org/10.1002/bimj.200810425

Innes, D. J., & Hebert, P. D. N. (1988). The origin and genetic basis of parthenogenesis in Daphnia
pulex. *Evolution*, *42*(5), 1024–1035. https://doi.org/10.1111/j.1558-5646.1988.tb02521.x

Jaegle, B., Soto-Jiménez, L. M., Burns, R., Rabanal, F. A., & Nordbog, M. (2021). Extensive gene
duplication in Arabidopsis revealed by pseudo-heterozygosity. *BioRxiv*.

Jaenike, J. (2001). Sex Chromosome Meiotic Drive. *Annual Review of Ecology and Systematics*,
32(1), 25–49. https://doi.org/10.1146/annurev.ecolsys.32.081501.113958

Janko, K., Pačes, J., Wilkinson-Herbots, H., Costa, R. J., Roslein, J., Drozd, P., Iakovenko, N., Rídl,
J., Hroudová, M., Kočí, J., Reifová, R., Šlechtová, V., & Choleva, L. (2018). Hybrid asexuality
as a primary postzygotic barrier between nascent species: On the interconnection between
asexuality, hybridization and speciation. *Molecular Ecology*, *27*(1), 248–263.
https://doi.org/10.1111/mec.14377

Jaron, K. S., Parker, D. J., Anselmetti, Y., Tran Van, P., Bast, J., Dumas, Z., Figuet, E., François, C.
M., Hayward, K., Rossier, V., Simion, P., Robinson-Rechavi, M., Galtier, N., & Schwander, T.
(2022). Convergent consequences of parthenogenesis on stick insect genomes. *Science Advances*,
8(8). https://doi.org/10.1126/sciadv.abg3842

Knaus, B. J., & Grünwald, N. J. (2017). vcfr: a package to manipulate and visualize variant call format
data in R. *Molecular Ecology Resources*, *17*(1), 44–53. https://doi.org/10.1111/1755-0998.12549

Kratochvíl, L., Vukić, J., Červenka, J., Kubička, L., Johnson Pokorná, M., Kukačková, D., Rovatsos,
M., & Piálek, L. (2020). Mixed-sex offspring produced via cryptic parthenogenesis in a lizard.
Molecular Ecology, *29*(21). https://doi.org/10.1111/mec.15617

Kuhn, A., Aron, S., & Hardy, O. J. (2021). Detection of Cryptic Sex in Automictic Populations:
Theoretical Expectations and a Case Study in Cataglyphis Desert Ants. *Frontiers in Ecology and
Evolution*, *9*. https://doi.org/10.3389/fevo.2021.741959

Laine, V. N., Sackton, T. B., & Meselson, M. (2022). Genomic signature of sexual reproduction in the
bdelloid rotifer *Macrotrachella quadricornifera*. *Genetics*, *220*(2).
https://doi.org/10.1093/genetics/iyab221

Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows–Wheeler transform.
Bioinformatics, *26*(5), 589–595. https://doi.org/10.1093/bioinformatics/btp698

Liegeois, M., Sartori, M., & Schwander, T. (2021). Extremely Widespread Parthenogenesis and a
Trade-Off Between Alternative Forms of Reproduction in Mayflies (Ephemeroptera). *Journal of*
Heredity, *112*(1), 45–57. https://doi.org/10.1093/jhered/esaa027

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von Haeseler, A., &
Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference
in the Genomic Era. *Molecular Biology and Evolution*, *37*(5), 1530–1534.
https://doi.org/10.1093/molbev/msaa015

Morgan-Richards, M., Langton-Myers, S. S., & Trewick, S. A. (2019). Loss and gain of sexual
reproduction in the same stick insect. *Molecular Ecology*, *28*(17), 3929–3941.
https://doi.org/10.1111/mec.15203

Neiman, M., Sharbel, T. F., & Schwander, T. (2014). Genetic causes of transitions from sexual
reproduction to asexuality in plants and animals. *Journal of Evolutionary Biology*, *27*(7).
https://doi.org/10.1111/jeb.12357

Normark, B. B. (2003). The evolution of alternative genetic systems in insects. *Annual Review of*
Entomology, *48*(1), 397–423. https://doi.org/10.1146/annurev.ento.48.091801.112703

Otto, S. P. (2009). The Evolutionary Enigma of Sex. *The American Naturalist*, *174*(S1), S1–S14.
https://doi.org/10.1086/599084

Pearcy, M., Hardy, O., & Aron, S. (2006). Thelytokous parthenogenesis and its consequences on
inbreeding in an ant. *Heredity*, *96*(5), 377–382. https://doi.org/10.1038/sj.hdy.6800813

Pijnacker, L. P., & Ferwerda, M. A. (1980). Sex Chromosomes and Origin of Males and Sex Mosaics
of the Parthenogenetic Stick Insect Carausius Morosus Br. *Chromosoma*, *79*(1), 105–114.

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics*, *26*(6), 841–842. https://doi.org/10.1093/bioinformatics/btq033

R Core Team. (2021). *R: A language and environment for statistical computing*. R Foundation for
Statistical Computing.

Raj, A., Stephens, M., & Pritchard, J. K. (2014). fastSTRUCTURE: Variational Inference of
Population Structure in Large SNP Data Sets. *Genetics*, *197*(2), 573–589.
https://doi.org/10.1534/genetics.114.164350

Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology (and other things).
Methods in Ecology and Evolution, *3*(2), 217–223. https://doi.org/10.1111/j.2041-
210X.2011.00169.x

Ryder, O. A., Thomas, S., Judson, J. M., Romanov, M. N., Dandekar, S., Papp, J. C., Sidak-Loftis, L.
C., Walker, K., Stalis, I. H., Mace, M., Steiner, C. C., & Chemnick, L. G. (2022). Facultative
Parthenogenesis in California Condors. *Journal of Heredity*, *113*(2), 217–217.
https://doi.org/10.1093/jhered/esab074

Sandoval, C. (2000). Persistence of a Walking-Stick Population (Phasmatoptera: Timematodea) after a
Wildfire. *The Southwestern Naturalist*, *45*(2), 123. https://doi.org/10.2307/3672452

Sandoval, C., Carmean, D. A., & Crespi, B. J. (1998). Molecular phylogenetics of sexual and
parthenogenetic Timema walking–sticks. *Proceedings of the Royal Society of London. Series B:*
Biological Sciences, *265*(1396), 589–595. https://doi.org/10.1098/rspb.1998.0335

Sandrock, C., & Vorburger, C. (2011). Single-Locus Recessive Inheritance of Asexual Reproduction
in a Parasitoid Wasp. *Current Biology*, *21*(5), 433–437.
https://doi.org/10.1016/j.cub.2011.01.070

Schultz, R. J. (1973). Unisexual Fish: Laboratory Synthesis of a "Species." *Science*, *179*(4069), 180–
181. https://doi.org/10.1126/science.179.4069.180

Schurko, A. M., Neiman, M., & Logsdon, J. M. (2009). Signs of sex: what we know and how we
know it. *Trends in Ecology & Evolution*, *24*(4), 208–217.
https://doi.org/10.1016/j.tree.2008.11.010

Schwander, T., & Crespi, B. J. (2009). Multiple direct transitions from sexual reproduction to
apomictic parthenogenesis in *Timema* stick insects. *Evolution*, *63*(1), 84–103.
https://doi.org/10.1111/j.1558-5646.2008.00524.x

Schwander, T., Crespi, B. J., Gries, R., & Gries, G. (2013). Neutral and selection-driven decay of
sexual traits in asexual stick insects. *Proceedings of the Royal Society B: Biological Sciences*,
280(1764), 20130823. https://doi.org/10.1098/rspb.2013.0823

Schwander, T., Henry, L., & Crespi, B. J. (2011). Molecular Evidence for Ancient Asexuality in
Timema Stick Insects. *Current Biology*, *21*(13), 1129–1134.
https://doi.org/10.1016/j.cub.2011.05.026

Schwander, T., Vuilleumier, S., Dubman, J., & Crespi, B. J. (2010). Positive feedback in the transition
from sexual reproduction to parthenogenesis. *Proceedings of the Royal Society B: Biological
Sciences*, *277*(1686), 1435–1442. https://doi.org/10.1098/rspb.2009.2113

Simon, C., Frati, F., Beckenbach, A., Crespi, B., Liu, H., & Flook, P. (1994). Evolution, Weighting,
and Phylogenetic Utility of Mitochondrial Gene Sequences and a Compilation of Conserved
Polymerase Chain Reaction Primers. *Annals of the Entomological Society of America*, *87*(6),
651–701. https://doi.org/10.1093/aesa/87.6.651

Suomalainen, E., Saura, A., & Lokki, J. (1987). *Cytology and Evolution in Parthenogenesis*. CRC
Press. https://doi.org/10.2307/2409571

Svendsen, N., Reisser, C. M. O., Dukić, M., Thuillier, V., Ségard, A., Liautard-Haag, C., Fasel, D.,
Hürlimann, E., Lenormand, T., Galimov, Y., & Haag, C. R. (2015). Uncovering Cryptic
Asexuality in *Daphnia magna* by RAD Sequencing. *Genetics*, *201*(3), 1143–1155.
https://doi.org/10.1534/genetics.115.179879

The Tree of Sex Consortium. (2014). Tree of Sex: A database of sexual systems. *Scientific Data*, *1*(1),
140015. https://doi.org/10.1038/sdata.2014.15

Vickery, V. R., & Sandoval, C. O. (1999). Two new species of Timema (Phasmatoptera:
Timematodea: Timematidae), one parthenogenetic, in California. *Journa of Orthoptera
Research*, *8*(Nov), 45–47.

Vickery, V. R., & Sandoval, C. P. (2001). Descriptions of three new species of Timema
(Phasmatoptera: Timematodea: Timematidae) and notes on three other species. *J. Orthoptera
Research*.

Supplementary material

Figure S1 Relative heterozygosity (based solely on polymorphic loci) in hatchlings from eggs laid before mating. Black dots: raw heterozygosity value, measured as the proportion of heterozygous SNPs for each individual, as a function of the proportion of missing data. Blue dots: corrected heterozygosity value for the same individuals, obtained by summing the residuals of a linear model predicting heterozygosity with the proportion of missing data (dashed line) and the intercept of that model (dotted line). This correction removes the variance in heterozygosity explained by missing data. The model is built on hatchlings from unfertilized eggs laid prior to mating only because of their extremely low heterozygosity. The effect of missing data is masked in normally heterozygous individuals because the effect size of the missing data on heterozygosity is smaller than the natural variation in heterozygosity levels (Figures 3 and 4). The model was then used to correct heterozygosity for all individuals.

Figure S2 Relative heterozygosity and genetic structure of all field-collected individuals. Top row: corrected relative heterozygosity. Following rows: fastSTRUCTURE results with different levels of K. For K = 3, colours as in main figures. Individuals ordered by location, as in Figures 1B and 1D. Note the high heterozygosity of individual 17-2162, which has a genetic contribution from all three inferred genetic lineages (k=3). This may indicate that this individual is a triploid hybrid between these lineages, but additional data would be needed to confirm (see Figure S5). Note also that all genotyped individuals (all males) from the Manchester 2 population are hybrids between the eastern and western Manchester lineages, with approximately equal contributions from the two. They are likely the result of past hybridization rather than F$_1$ hybrids, as their heterozygosity levels are similar to non-hybrid individuals. Surprisingly they have maintained a high frequency of alleles from the eastern Manchester lineage while neighbouring populations are fixed for alleles from the western Manchester lineage. This further supports the very limited dispersal in these wingless insects.

814
815 **Figure S3** Frequency distribution of $\log_2$ ratio of male to female coverage for all contigs >= 1000 bp in
816 length. Autosomal contigs should have equal coverage in males and females ($\log_2$ ratio of male to
817 female coverage = 0). X-linked contigs should have half the coverage in males than in females ($\log_2$
818 ratio of male to female ratio coverage = -1). X-linked scaffolds were classed as contigs with a $\log_2$
819 ratio of male to female coverage within 0.1 of the X linked peak (black dotted line).
820
821

Assessing the amount of false-positive heterozygosity

We found that all but one offspring produced from unfertilized eggs laid prior to mating had similar and very low relative heterozygosity (~ 2-3%), similar to the heterozygosity measured in obligately parthenogenetic reference individuals, and independently of the heterozygosity of their mother (Figure 3). Previous work revealed that the extremely low heterozygosity measured in obligately parthenogenetic individuals is largely due to divergence between paralogs and structural variation between individuals (Jaron et al., 2022), i.e., represents mostly false-positive heterozygosity. False-positive heterozygosity calls generated by structural variation among individuals is increasingly recognized as a general problem in sequencing data sets, even in model organisms with very well resolved genomes such as *Arabidopsis* (Jaegle et al., 2021). For all these reasons we suspected that the 2-3% relative heterozygosity measured in parthenogenetically produced offspring was also largely generated by repeated regions and structural variation, and not due to heterozygosity from allele divergence.

We investigated whether this was indeed the case by looking at heterozygosity on the X chromosome. Because the X chromosome is monosomic in males (*Timema* have XX:XO sex determination), all heterozygosity measured on the X in males can be classified as false positives. Thus, by comparing heterozygosity on the X chromosome of males with the heterozygosity of the X chromosome of parthenogenetic (diploid) offspring, we can assess whether the heterozygosity in parthenogenetic offspring is also generated by false positives. This would be the case if heterozygosity on the X was similar in the two groups.

We found no difference in heterozygosity levels on the X chromosome of males and parthenogenetic offspring (t-test: $t = 0.296$, $df = 39.741$, $p = 0.77$; Figure S4A). This means that parthenogenetic hatchlings are fully homozygous on their X chromosome. Interestingly, heterozygosity in parthenogenetic offspring was higher on the X chromosome than genome-wide, indicating that repeated regions and other structural variants that are generating false-positive heterozygosity are enriched on the X.

To further confirm that heterozygosity on the male X was largely due to repeated regions, we compared average depth at homozygous vs heterozygous loci. We standardised depth per individual and tested whether there was a significant difference using a paired t-test. We found that depth at heterozygous X-linked loci was higher than at homozygous X-linked loci in all 40 males (average: 2.8x higher; $t = -21.1$, $df = 39$, $p < 2.2^{-16}$; Figure S4B).

857
Figure S4 Repetitive regions generate false-positive heterozygosity on the X chromosome of males
and in parthenogenetically-produced offspring. **A** Relative heterozygosity (based solely on
polymorphic loci, see main text) on the X chromosome of males, on the X chromosome of
parthenogenetically-produced offspring and on the whole genome of parthenogenetically-produced
offspring. One point represents average heterozygosity across loci for one individual. Note that the
heterozygosity estimates are corrected for the effect of missing data as in Figure S1 (separately for
males and parthenogenetic offspring as the effect of missing data is expected to differ for monosomic
vs. diploid chromosomes). **B** Average depth (standardised per individual) of homozygous and
heterozygous loci on the X chromosome of males. Each dot is a male, depth values for homozygous
and heterozygous loci of the same individual are linked with a line.
868
869

Sexing sexually produced offspring

In order to sex the sexually produced hatchlings, we looked at heterozygosity on the X chromosome and depth ratio between the X chromosome and autosomes. Because they have only one copy of the X chromosome, males are expected to be homozygous at all X-linked loci. In addition, X-linked loci should have half the coverage of autosomes. However, we observed a higher amount of false-positive heterozygosity on the X than on autosomes, which was largely generated by repetitive regions (see previous section and Figure S4A). In addition to generating false-positive heterozygosity, these replicated regions also bias the X/autosome depth ratio. Both these effects are enhanced in individuals with large amounts of missing data because of low sequencing coverage (because loci with too low coverage are filtered out in all individuals). For this reason, we used a stricter cutoff for this analysis, retaining only individuals with fewer than 20% of missing data. 176 out of the 211 sexually produced offspring passed this cutoff, of which 95 were inferred to be males and 81 inferred to be females (Figure S5).

Figure S5 X / autosomes depth ratio and heterozygosity on the X, used to sex the sexually produced hatchlings (hatchlings cannot be sexed by morphology). Males and females collected in the field (and sexed by morphology) are indicated for comparison. Colours as in Figure 1. Hatchlings were considered males if they had an X / autosomes depth ratio lower than 0.9 (dotted line), and female otherwise. Note that the X/autosome depth ratios are higher than expected for both males and females, consistent with the X chromosome being enriched for repetitive content. Female 17-2162, the hybrid with a contribution from each of the three genetic lineages, stood out for both measures. Its X chromosome was more heterozygous than that of all field-collected adults, but its X/autosome depth ratio was lower than that of other females. While further research is required to explain this pattern, it could stem from a triploid constitution with only two X chromosomes.

Hatchling sex ratio

Figure S6: Sex ratio of sexually produced hatchlings per family. The grey dots represent a 50:50 sex ratio (the null hypothesis). The black lines are 95% binomial proportion confidence intervals around the observed sex ratio. All confidence intervals overlap with the expected value under mendelian inheritance, and the p-value of Fisher's exact tests for each family were higher than 0.5, providing no evidence for X-drive underlying biased sex ratios.