1

Title: Pharmacological manipulations of physiological arousal

and sleep-like slow waves modulate sustained attention

Authors: Elaine Pinggal[1], Paul M. Dockree[2], Redmond G. O'Connell[2], Mark A. Bellgrove[1]*
& Thomas Andrillon[3,4]*

*: equal contribution

Affiliations:

1. School of Psychological Sciences, Turner Institute for Brain and Mental Health, Monash University, Melbourne 3168, Victoria, Australia.
2. Trinity College Institute of Neuroscience & School of Psychology, Trinity College Dublin, Dublin, Ireland.
3. School of School of Philosophical, Historical, and International Studies, Centre for Consciousness and Contemplative Studies, Monash University, Melbourne 3168, Victoria, Australia.
4. Paris Brain Institute, Sorbonne Université, Inserm-CNRS, Paris, 75013, France.

Corresponding author: Thomas Andrillon.
Postal Address: Paris Brain Institute, Hôpital Pitié-Salpêtrière, 47 Bd de l'Hôpital, 75013 Paris, France.
Email: thomas.andrillon@icm-institute.org
ORCID: https://orcid.org/0000-0003-2794-8494

Abstract

Sustained attention describes our ability to keep a constant focus on a given task. This ability

is modulated by our physiological state of arousal. Although lapses of sustained attention have

been linked with dysregulations of arousal, the underlying physiological mechanisms remain

unclear. An emerging body of work proposes that the intrusion during wakefulness of sleep-

like slow waves, a marker of the transition toward sleep, could mechanistically account for

attentional lapses. This study aimed to expose, via pharmacological manipulations of the

monoamine system, the relationship between the occurrence of sleep-like slow waves and the

behavioural consequences of sustained attention failures. In a double-blind, randomised-

control trial, 32 healthy participants received methylphenidate, atomoxetine, citalopram or

placebo during four separate experimental sessions. During each session,

electroencephalography (EEG) was used to measure neural activity whilst participants

completed a visual task requiring sustained attention. Methylphenidate, which increases wake-

promoting dopamine and noradrenaline across cortical and subcortical areas, improved

behavioural performance whereas atomoxetine, which increases dopamine and noradrenaline

predominantly over frontal cortices, led to more impulsive responses. Additionally, citalopram,

which increases sleep-promoting serotonin, led to more missed trials. Based on EEG recording,

citalopram was also associated with an increase in sleep-like slow waves. Importantly,

compared to a classical marker of arousal such as alpha power, only slow waves differentially

predicted both slower and faster responses in a region-specific fashion. These results suggest

that a decrease in arousal can lead to local sleep intrusions during wakefulness which could be

mechanistically linked to impulsivity and sluggishness.

Introduction

Sustained attention refers to the ability to concentrate cognitive resources on a given task or specific stimuli for extended periods of time (van Schie et al., 2021). Previous research has emphasised the capacity limits of this ability, especially when the task at hand is easy and repetitive or when individuals are unmotivated (Esterman and Rothlein, 2019). This difficulty in maintaining a continuous attentional focus goes beyond lab studies, with an estimated 50% of our waking time spent not focusing on the task at hand (Killingsworth and Gilbert, 2010). Fluctuations of attention impact performance in a complex fashion, leading to a mixture of unresponsiveness, sluggishness and impulsivity (O'Connell et al., 2009; Oken et al., 2006; Riley et al., 2017). Fluctuating attention also impacts subjective experience, resulting in phenomena such as mind wandering (i.e. thinking about something else than the task at hand) (Smallwood and Schooler, 2015) or mind blanking (i.e. thinking about nothing at all) (Ward and Wegner, 2013).

Failures to maintain sustained attention vary with fluctuations in physiological arousal, defined here as the physiological state that allows encoding, processing, and responding to environmental stimuli. Indeed, on the one hand, attentional failures have been associated with hypo-arousal as they increase with fatigue and time spent awake (Riley et al., 2017) and are very sensitive to sleep deprivation (Doran et al., 2001). On the other hand, an excess of arousal (hyper-arousal) also appears detrimental to sustained attention (Mittner et al., 2016). For example, mind wandering has been associated with increased baseline pupil dilation (a marker of physiological arousal) compared to task-focused states (Smallwood et al., 2011). In addition, hypo- and hyper-arousal have been hypothesised to be linked as states of hyper-arousal could serve to transiently offset a baseline state of hypo-arousal. For example, individuals with attentional deficits (e.g. Attention Deficit Hyperactivity Disorder, ADHD) may experience low levels of arousal at rest (Satterfield and Dawson, 1971) and attentional deficits could stem from

a combination of a low-arousal baseline (e.g. low tonic dopaminergic activity) combined with an autoregulatory attempt at compensating for this low baseline (e.g. high phasic dopaminergic activity) (Geissler et al., 2014). At the level of cortical networks, intermediate levels of arousal could promote optimal levels of sustained attention by balancing levels of excitation and inhibition (Oken et al., 2006). Interestingly, the neuromodulators involved in the regulation of physiological arousal and sleep-wake transitions (Jones, 2005; Oken et al., 2006; Thiele and Bellgrove, 2018) are also involved in the regulation of sustained attention (Oken et al., 2006; Thiele and Bellgrove, 2018).

Arousal is regulated by a number of neuromodulators, which include (but is not limited to) noradrenaline, dopamine and serotonin (Thiele and Bellgrove, 2018). These neuromodulators have well-documented effects on attention by, for example, optimising neural responses to the environment (Ranjbar-Slamloo and Fazlali, 2020) or promoting prefrontal cortical processing (Robbins, 1997; Thiele and Bellgrove, 2018).

First, noradrenaline is one of the key neuromodulators allowing global transitions from wakefulness to sleep and vice versa (Jones, 2005). Noradrenaline release leads to broad cortical activation via the inhibition of sleep-promoting systems and the activation of wake-promoting systems. The widespread cortical release of noradrenaline by the Locus Coeruleus (LC) is maximal during wakefulness, decreases in Non-Rapid Eye Movement (NREM) sleep and vanishes in Rapid Eye Movement (REM) sleep. Additionally, noradrenaline is released following the occurrence of novel or salient stimuli and can decrease response thresholds and improve the efficiency of synaptic transmissions by modifying the cellular properties of target neurons (Robbins, 1997; Ross and Van Bockstaele, 2021; Sara, 2009). A previous report on the same dataset as we present here, showed that an increase of noradrenaline and dopamine via methylphenidate led to enhanced responses to visual stimuli and a reduction of alpha oscillations (Dockree et al., 2017). However, too much noradrenergic activity (as for example

during high stress situations) can also lead to an impairment of executive functions and behaviour (Arnsten, 2010, 2000). Finally, another recent study found that decreasing the concentration of noradrenaline through pharmacological challenge led to an increase in drowsiness, decreased performance in a visual task and weaker sensory responses whereas increasing noradrenaline had the opposite effect (Gelbard-Sagiv et al., 2018), confirming that the impact of noradrenaline on performance is U-shaped (Sara and Bouret, 2012).

Second, the relationship between dopamine and cognitive performance also follows an inverted U-shape, with the best cognitive performance being observed when dopamine levels are at an intermediate level (Maffei and Angrilli, 2018). However, the impact of dopamine on arousal is difficult to separate from that of noradrenaline (Lee and Dan, 2012; Robbins, 1997). Both noradrenaline and dopamine innervate related areas, share biosynthetic and intracellular signalling pathways, and are occasionally released simultaneously from noradrenergic terminals such as the LC (Ranjbar-Slamloo and Fazlali, 2020). An integrative model of the role of noradrenaline and dopamine posits that dopamine decreases the amount of noise whereas noradrenaline enhances signal, participating thus in a complementary process to increase the signal-to-noise ratio (Arnsten and Rubia, 2012). Importantly, like noradrenaline, dopamine also influences physiological arousal and promotes wakefulness (Boutrel and Koob, 2004). In fact, early treatments against narcolepsy included stimulant drugs such as methylphenidate, which increase the levels of dopamine and noradrenaline (Mignot, 2012) and is also used in the treatment of attentional disorders (Arnsten, 2010; Cortese, 2020).

Third, some evidence also suggests that serotonin may play a role in the regulation of arousal. For instance, serotonin may influence the regulation of sleep and wakefulness as the raphe nuclei releases serotonin maximally during wakefulness, less during NREM sleep and ceases firing during REM sleep (Jones, 2005). These effects have been interpreted as serotonin stimulating wakefulness (Jones, 2005), however, a recent study by Oikonomou et al. (2019)

suggests a role for serotonin in facilitating sleep onset. Accordingly, serotonin could be

involved in the homeostatic regulation of sleep and its high concentration during wakefulness

could be related to the build-up of sleep pressure during the day (Borbély et al., 2016; Jouvet,

1999). Serotonin also modulates attention directly or indirectly, by inhibiting, for example, the

cholinergic system which plays a key role in the maintenance of attention (Jones, 2005; Sparks

et al., 2018). Direct effects appear region or dose-specific (Thiele and Bellgrove, 2018) and

low tonic activity in serotonin could improve attention by reducing impulsivity while high tonic

activity could decrease the activation of prefrontal areas commonly activated in a sustained

attention task (Sturm and Willmes, 2001; Wingen et al., 2008) and increase distractibility.

Although the neuromodulators mentioned above play a role in both the regulation of

arousal and attention, these phenomena are often conceptualised and investigated separately.

Here, we sought to explore whether the regulation of sleep/wake dynamics could be directly

related to the modulation of sustained attention. For example, an increase in noradrenaline

and/or dopamine could result in increased levels of physiological arousal, leading to cortical

activation, improved signal-to-noise ratio of sensory responses, better intracortical

communication, and lower response thresholds (Thiele and Bellgrove, 2018). Conversely, a

decrease in arousal mediated by the same neuromodulators could negatively impact attention

through the promotion of cortical idling (Pfurtscheller, 1992) or even sleep-like activity and

cortical deactivation (Andrillon et al., 2019).

Electrophysiological (EEG) markers such as alpha oscillations have been used to

investigate the relationship between physiological arousal and attention (Cajochen et al., 1995;

Oken et al., 2006; Peylo et al., 2021). These oscillations are generated over the visual cortex

and consist of sustained oscillatory activity with a frequency between 7 and 13 Hz (Clayton et

al., 2018). A suppression of alpha oscillations has often been used as a proxy to estimate

cortical excitability (Romei et al., 2008), a key component of physiological arousal, although

the generalisation of this relationship outside visual areas has been questioned (Samaha et al.,

2017). A study by O'Connell et al. (2009) found that alpha oscillations were a strong predictor

of attentional lapses, with an increase in alpha activity predicting lapses up to 20 seconds before

their occurrence. Alpha oscillations exemplify how modulations of arousal could impact

attention as increased levels of alpha oscillations could favour lapses of attention (Dockree et

al., 2017; O'Connell et al., 2009) by increasing cortical deactivation (Pfurtscheller et al., 1996).

Conversely, a decrease in alpha oscillations could favour enhanced cortical activation and

promote information processing (Klimesch et al., 2011; Peylo et al., 2021). However, the

relationship between alpha oscillations and arousal is not monotonic as alpha oscillations

decrease when individuals become drowsy or fall asleep (Kalauzi et al., 2012). In addition, the

associations between alpha oscillations and attention are task-dependent as increases in alpha

power have also been associated with the suppression of task-irrelevant distractors (Klimesch,

2012; Klimesch et al., 2011) and improved auditory processing (Dimitrijevic et al., 2017).

Thus, while the level of alpha oscillations have been robustly linked to sustained attention, this

relationship appears to be dependent on the nature of the task, the brain region over which it is

measured (Jensen and Mazaheri, 2010), and the baseline arousal state. Similarly, the

relationship between alpha oscillations and mind wandering appears complex, with an

influential study reporting less alpha during mind wandering (Braboszcz and Delorme, 2011)

whereas others found the reverse pattern (Arnau et al., 2020; Compton et al., 2019; Macdonald

et al., 2011).

To overcome these limitations, we examined the relationship between physiological

arousal and attention using a novel electroencephalographic (EEG) marker of arousal: sleep-

like slow waves occurring within wakefulness (Andrillon et al., 2019). Indeed, while sleep and

wakefulness are usually presented as mutually exclusive states, recent discoveries have shown

that sleep and wake activity can be concomitantly observed in different brain regions (Nir et

al., 2017; Nobili et al., 2012; Vyazovskiy et al., 2011). In particular, when one becomes fatigued, high amplitude slow oscillations (<7Hz), analogous to delta or theta sleep slow waves but smaller in amplitude, can be detected through EEG recordings (Andrillon et al., 2021; Bernardi et al., 2015; Hung et al., 2013). At the neuronal level, these waves are paired with momentary and localised periods of neuronal silencing (Vyazovskiy et al., 2011). These occurrences have been interpreted as "local sleep episodes" (Vyazovskiy et al., 2011), and could perturb information processing (Andrillon et al., 2019), following similar mechanisms that limit sensory processing and responsiveness during sleep (Andrillon and Kouider, 2020). Accordingly, these local sleep episodes have been shown to negatively impact behaviour (Bernardi et al., 2015; Hung et al., 2013; Vyazovskiy et al., 2011), notably in the context of tasks requiring sustained attention, and increase the frequency of mind wandering and mind blanking events (Andrillon et al., 2021). Importantly, these sleep-like slow waves are not restricted to contexts of acute sleep deprivation (Bernardi et al., 2015; Hung et al., 2013) but can also be found in well-rested individuals (Andrillon et al., 2021; Quercia et al., 2018). We aimed here to test if these slow waves varied as a function of pharmacological manipulation of key monoamine neuromodulators in well-rested individuals and could explain the relationship between fluctuations of arousal and attention.

To expose the relationship between the physiological substrates of sustained attention and arousal, monoamine (dopamine, noradrenaline and serotonin) agonists were administered to healthy, non-sedated participants while they performed a sustained attention EEG task (Figure 1). We re-analysed a dataset of participants who completed the Continuous Temporal Expectancy Task (CTET) (Dockree et al., 2017), a monotonous task wherein participants are required to monitor visual checkerboard stimuli for intermittent targets defined by their longer presentation (O'Connell et al., 2009). Methylphenidate (MPH) was used to increase both noradrenaline and dopamine levels globally across the cortex whilst atomoxetine (ATM) was

used to increase noradrenaline and dopamine more specifically in the prefrontal cortex

(Bymaster, 2002; Chamberlain et al., 2009; Farr et al., 2014; Koda et al., 2010; Kowalczyk et

al., 2019). Lastly, citalopram (CIT) was administered to increase serotonin. Only the results

obtained with MPH have been previously reported, showing an improvement in sustained

attention, a decrease in alpha oscillations and an increase in target detection (indexed via the

P3 event-related potential), compared to placebo (Dockree et al., 2017). We expanded these

findings by examining the impact of both arousal-promoting (MPH, ATM) and arousal-

reducing (CIT) drugs. We also examined a more exhaustive set of EEG markers of attention

and arousal, including physiological markers of sleep intrusions such as slow waves. We

hypothesised that increasing noradrenaline and dopamine would lead to an increase in attention

(decrease in attentional lapses) and arousal (less slow waves and alpha oscillations), whereas

increasing serotonin would lead to impaired attention (increase in attentional lapses) and a

decrease in arousal (more slow waves and alpha oscillations). Additionally, we hypothesised

that slow waves and alpha oscillations would have differing roles in terms of their relationship

with attentional lapses. Specifically, we hypothesised that in contrast to alpha oscillations

which are generated within the occipital cortex, the behavioural impact of slow waves would

vary as a function of their spatial location, with frontal slow waves being associated with

impulsive behaviour (faster reaction times, more false alarms) and posterior slow waves being

associated with sluggishness (slower reaction times, more misses) (Andrillon et al., 2021).

Such a pattern of results would reinforce the idea that slow waves represent a

neurophysiological mechanism underlying attentional lapses, in contrast to alpha oscillations

which could more simply index an individual's level of arousal.

Figure 1. Experimental protocol.

a. Each experimental session started with participants being administered one out of four treatments (PLA: placebo; MPH: methylphenidate; ATM: atomoxetine; CIT: citalopram) and completing the first Visual Analogue Scale (VAS) (time=0). The administration of the treatments was randomised across sessions (within-subject design). A second VAS was completed before starting the Continuous Temporal Expectancy Task (CTET), 90 minutes after treatment administration. Finally, after the CTET (and another task not analysed here) and 180 minutes after treatment administration, participants completed a third VAS. High-density EEG was recorded during the CTET. b. The CTET task was used to test sustained attention. In the CTET, participants are instructed to monitor a stream of visual stimuli (black and white checkboards) displayed on a screen placed in front of them. There are two types of stimuli: target stimuli for which participants are instructed to press a response button, and non-target stimuli that do not require a response. The only distinguishing factor between the non-target and target stimuli is the duration of the stimulus presentation (non-target: 800ms; target: 1120ms).

Results

Subjective Sleepiness

To verify that the pharmacological interventions impacted subjective sleepiness, participants completed a visual analogue scale (VAS) three times during each experimental session: at drug administration (t=0 minutes), before the start of the sustained attention task (t=90 minutes), and after the task (t=180 minutes). A two-way repeated measures ANOVA was conducted to examine the effects of treatment (PLA, MPH, ATM, or CIT) and time (0, 90 and 180 minutes after drug administration; Figure S1) on subjective sleepiness. Thirty participants were included in this analysis after two subjects were excluded due to missing data (one experimental session missing in each participant, see Methods). This ANOVA revealed a main effect of time ($F(1.49, 43.28) = 22.4$, $p < .001$) and treatment ($F(3, 87) = 6.3$, $p < .001$) on sleepiness as well as a significant two-way interaction between treatment and time ($F(6, 174) = 6.5$, $p < .001$). Pairwise post-hoc comparisons by time points (t-tests, Bonferroni-corrected) indicated that there were no significant differences in sleepiness across drug treatment at the beginning of the protocol (drug administration: $F(3, 87) = 0.36$, $p_{adjusted} = 1$) but there were significant differences before and after the task ($F(3, 87) = 5.0$, $p_{adjusted} = .009$ and $F(3, 87) = 10.6$, $p_{adjusted} < .001$ resp.), confirming the effectiveness of the sustained attention protocol in engendering sleepiness. Pairwise post-hoc comparisons within treatments (t-tests, Bonferroni-corrected) indicated that this effect of treatment was driven by an absence of an increase in sleepiness for MPH, in keeping with the stimulant effect of MPH ($F(2, 58) = 2.0$, $p_{adjusted} = .59$), compared with PLA, CIT and ATM (all $p_{adjusted} < .01$). However, the administration of ATM or CIT did not significantly alter the subjective experience of sleepiness across the protocol compared with PLA.

Sustained Attention Performance

We next sought to examine the influence of the pharmacological manipulations on behavioural measures of sustained attention. To do so, we examined behavioural performance during the Continuous Temporal Expectancy Task (CTET), a task designed to monitor participants' ability to pay careful attention to a continuous stream of visual information (O'Connell et al., 2009). We focused on three behavioural variables: the proportion of missed target trials (misses), the proportion of incorrect responses to non-target trials (false alarms) and the reaction times for target trials (Figure 2).

To examine the performance across the entire CTET task (10 blocks), we fitted linear mixed effect models to predict each of these variables of interest, with treatment as a categorical fixed effect and subject identity as a categorical random effect (see Methods). We compared these models with a null model, including only subject identity as a random effect using the Likelihood Ratio Test, to test the effect of treatment on these behavioural variables. These model comparisons revealed a strong effect of treatment on misses ($\chi^2(3) = 207.2$, $p < .001$), false alarms ($\chi^2(3) = 52.1$, $p < .001$), and reaction times ($\chi^2(3) = 82.0$, $p < .001$).

Focusing on the winning models revealed that MPH, compared to PLA, was associated with an improvement in behavioural performance across all three variables: less misses (t = -10.5, $p < .001$), less false alarms (t = -3.5, $p < .001$) and faster reaction times (t = -7.2, $p < .001$). This means that, under MPH, participants were faster and better at detecting target trials without becoming too impulsive and responding to non-target trials. We applied the signal detection theory (Macmillan, 1993) to combine responses on target and non-target trials in one single index of behavioural performance (d'; Figure S2). MPH led to an increase in d' compared to PLA (t = 8.8, $p < .001$), indicating enhanced sensitivity.

Figure 2. Pharmacological modulations of behavioural performance.
Left panels show the sample and individual averages computed across all CTET blocks (N=10
blocks) for each treatment (PLA: N=32; CIT: N=31; ATM: N=31; and MPH: N=32) in the
form of raincloud plots (see Methods). Right panels show the sample averages for each drug
*treatment and block separately. Panel **a.** shows the percentages of missed target trials (misses),*
*panel **b.** shows the percentages of false alarms and panel **c.** shows the averaged reaction times*
on correctly detected targets. On the left panels, stars denote the significance level (ns: non-

Although ATM is also used to reduce ADHD symptoms, in contrast to MPH, it is a

non-stimulant. Results indicated that ATM was associated with more false alarms (t = 3.6,

$p < .001$) compared with PLA, whereas misses and reaction times did not differ compared with

PLA (t = -0.7, $p = .45$ and t = 0.0, $p = .99$ resp.). Thus, ATM decreased performance by

increasing impulsive responses (false alarms). However, given the low proportion of false

alarms, this did not result in a significant decrement of overall performance compared to

placebo, as assessed by d' (t = -1.7, $p = .09$; Figure S2).

We observed an increase in the proportion of misses under CIT compared with PLA,

(t = 4.0, $p < .001$), whereas the proportion of false alarms and the average reaction times did

not significantly differ (t = -1.5, $p = .13$ and t = 0.8, $p = .41$, respectively). Thus, contrary to

MPH, CIT led to a decrease in sustained attention performance, which was reflected in a

decrease of the d' (t = -2.5, $p = .01$; Figure S2). In addition, whereas ATM resulted in impulsive

responses, CIT resulted in the opposite behavioural effect with more misses. Thus, the three

pharmacological probes had unique effects on behavioural performance during the CTET,

either enhancing or impairing discrete aspects of sustained attention.

Finally, we also examined the effect of time-on-task on behavioural performance and

its interaction with the drug treatments. To do so, we fitted linear mixed effect models including

block number and treatment as fixed effects, along with their interaction component, and

subject identity as a categorical random effect. We tested the significance of the interaction

component by comparing this model with the same model minus the interaction component,

using the Likelihood Ratio Test. This approach showed that there was an interaction between

time and treatment for misses (Likelihood Ratio Test for the model comparison: $\chi^2(3) = 9.15$,

$p = 0.03$). This interaction was driven by MPH, which stabilised performance and prevented

the increase of misses with time-on-task (Figure 2a). CIT, on the other hand, resulted in a

constant increase in misses throughout the task compared to placebo. There was no interaction

between time and treatment for false alarms or reaction times ($\chi^2(3) = 2.38$, $p = 0.50$ and

$\chi^2(3) = 4.34$, $p = 0.23$, respectively). This analysis confirmed the opposing effects of MPH and

CIT on attentional lapses across time.

EEG indexes of visual processing: Event-Related Potentials (ERPs) and Steady-State
Visual Evoked Potentials (SSVEPs)

To further understand the impact of pharmacological treatment on the physiology of

sustained attention, we first focused on two EEG indices of visual processing: Event-related

potentials (ERPs) (Figure 3) and steady state visual evoked responses (SSVEPs) (Figure 4).

Comparing the ERPs associated with the offset of the Target and non-Target trials revealed a

strong P3 effect (Figure 3a), a larger positivity for Target trials, starting at stimulus offset and

maximal over parietal electrodes (Figure 3b). We thus extracted the amplitude of this P3

component and compared it across drug treatments using a cluster-permutation approach (see

Methods). This analysis revealed an increase of the P3 amplitude following MPH

administration compared to PLA (Figure 3c), in line with previous findings on this same dataset

(Dockree et al., 2017). Interestingly, now examining the impact of ATM and CIT as well, the

amplitude of the P3 was not significantly impacted by the administration of these drugs

compared with PLA (Figure 3c).

The rapid flickering of the visual checkerboard stimuli generated a SSVEP, allowing

us to examine the impact of drug treatment on an EEG index of visual processing. We observed

a clear peak at the flicker frequency of checkerboard (25 Hz) in the power spectrum (Figure 4a

showing the signal-to-noise-ratio (SNR) or power at each frequency bin normalised by

neighbouring bins, see Methods). We thus focused on the SNR at 25Hz and compared the

strength of this entrainment for each electrode across drug treatments (Figure 4b-c). This

analysis revealed an increase for MPH compared to PLA, maximal over central electrodes

(Figure 4c). Neither ATM nor CIT significantly modulated the strength of SSVEPs compared

to PLA (Figure 4c).

Figure 3. Event-related potentials.
a. Event-Related Potentials (ERPs) locked on stimulus offset for Target (continuous lines) and
non-Target (dashed lines) trials and split by drug treatment (PLA: grey, MPH: orange, ATM:
purple, CIT: green). ERPs are averaged across participants (PLA: N=32; CIT: N=31; ATM:
N=31; and MPH: N=32). ERPs computed on two electrodes are shown: Fz (Frontal, left) and
Pz (parietal, right). Coloured lines show the sample average and coloured areas the standard
error of the mean. The yellow area ([0, 0.3]s post-offset window) shows the interval of the
archetypal P3 used to compute the P3 amplitude (see Methods). b. Topographical maps of the
P3 amplitude (difference of the ERP amplitude between Target and non-Target trials and
average on a [0, 0.3]s window) for each drug treatment. c. Topographical maps of the
statistical differences in P3 amplitude between MPH and PLA (left), ATM and PLA (middle)
and CIT and PLA (right). The t-values obtained with paired t-tests for each electrode are
shown. Significant clusters of electrodes ($p_{cluster}$<.05) are shown with black dots (cluster-
permutation approach, see Methods). A significant cluster was found only for MPH.
369
370

Figure 4. Steady-State Visual Evoked Potentials.

Visual stimuli were flashed at a 25Hz rate on the screen, entraining neural activity at the same frequency (frequency tagging) and generating Steady-State Visual Evoked Potentials SSVEP). This frequency tagging can be observed when computing the power spectrum of the EEG signal and extracting the signal-to-noise ratio (SNR) of the frequency tag (see Methods). ***a.*** *SNR by frequency and drug treatment, computed for electrode Cz and across participants (PLA: N=32; CIT: N=31; ATM: N=31; and MPH: N=32). A clear peak at 25 Hz (highlighted in yellow) is present for all treatments (PLA: grey, MPH: orange, ATM: purple, CIT: green).* ***b.*** *Topographical maps for the SNR at 25 Hz for each drug treatment.* ***c.*** *Topographical maps of the statistical differences in 25 Hz SNR between MPH and PLA (left), ATM and PLA (middle) and CIT and PLA (right). The t-values obtained with paired t-tests for each electrode are shown. Significant clusters of electrodes ($p_{cluster}<.05$) are shown with black dots (cluster-permutation approach, see Methods). A significant cluster was found only for MPH.*

EEG indexes of arousal: Alpha Power and Sleep-Like Slow Waves

To better understand the impact of the drug manipulations on physiological arousal, we

examined a classical marker of arousal - alpha power. Inspection of the EEG power spectrum

revealed a bump, or deviation from the 1/f trend, in the alpha frequency band (8-11 Hz; Figure

5a). We thus extracted the average power in this band and compared this alpha power across

treatments at the electrode level (Figure 5b). A cluster permutation analysis revealed that MPH

was associated with reduced levels of alpha power in comparison to PLA, consistent with

previous reports (Dockree et al., 2017; Janssen et al., 2016) and the idea that alpha

desynchronisation is associated with improved attention. However, no differences were

observed for ATM and CIT compared with PLA (Figure 5c).

Figure 5. Power of alpha oscillations.
We computed the Power Spectral Density (PSD) to analyse the impact of treatments on alpha
oscillations, a common marker of physiological arousal. The power of alpha oscillations was
obtained by averaging the log power of the PSD on a [8, 11] Hz window and for each electrode,
participant and drug treatment (PLA: N=32; CIT: N=31; ATM: N=31; and MPH: N=32). ***a.***

*Average PSD for electrode Cz and for each drug treatment (PLA: grey, MPH: orange, ATM: purple, CIT: green). The alpha power window is highlighted in yellow. **b.** Topographical maps of alpha power for each drug treatment. **c.** Topographical maps of the statistical differences in alpha power between MPH and PLA (left), ATM and PLA (middle) and CIT and PLA (right). The t-values obtained with paired t-tests for each electrode are shown. Significant clusters of electrodes ($p_{cluster}$<.05) are shown with black dots (cluster-permutation approach, see Methods). A significant cluster was found only for MPH.*

Since only MPH demonstrated effects compared with PLA for the different EEG indices explored thus far (ERPs, SSVEPs and alpha power), these EEG markers are unable to account for the detrimental effects of CIT or ATM on behaviour (Figure 2). In addition, alpha power has a complex relationship with arousal, decreasing both when arousal increases and when individuals approach sleep. We therefore extracted another emerging EEG marker of low arousal, sleep-like slow waves, which has been proposed to index the transition of local networks toward sleep (Andrillon et al., 2019). We applied established algorithms to extract the number of sleep-like slow waves across electrodes and drug treatments (see Methods).

We compared the density of sleep-like slow waves (averaged across all electrodes) across sessions by fitting a linear fixed effect model comprising a fixed categorical effect of drug treatment and a random categorical effect of subject identity. We then compared this model with a model with only subject identity as a random effect (see Methods). A likelihood ratio test revealed that drug treatment had a significant effect on the density of sleep-like slow waves ($\chi^2(3) = 32.79$, $p < .001$). Examining the winning model revealed that this modulation was driven by an increase in slow-wave density for CIT compared to PLA (t = 5.13, $p < .001$; Figure 6a). No significant differences were observed for MPH and ATM (t = 0.19, $p = .85$ and t = 1.54, $p = .12$, resp.). We then explored the interaction between block number and treatment by comparing a model including treatment and experimental block as fixed effects, with a model including both fixed effects and their interaction component. A likelihood ratio test did not reveal a significant interaction between block number and treatment ($\chi^2(3) = 1.15$, $p = .77$,

Figure 6b). Rather, CIT was associated with more slow waves throughout the task, independent of any time-on-task effects. Finally, we examined topographical differences by extracting slow-wave density across electrodes and comparing it across treatments using linear mixed effects models with a single fixed effect of treatment (see Methods). A cluster-permutation analysis revealed a widespread increase in slow-wave density for CIT but no significant clusters for MPH and ATM (Figure 6c).

Figure 6. Sleep-like Slow Waves.
a. Slow-wave density (wave/min averaged across all electrodes) per treatment. The raincloud plot for each drug treatment (PLA: grey (N=32), MPH: orange (N=32), ATM: purple (N=31), CIT: green (N=31)) is shown in the left panel. Stars denote the significance level (ns: non-significant; (*): p < .05; (**): p < .01; (***): p < .001) of the difference between placebo and the three treatments used in this protocol (MPH vs PLA, ATM vs PLA, CIT vs PLA), as determined by linear mixed-effect models. In the right panel, slow-wave density split by

 treatment and block (N=10 blocks) showing the sample average (circle) and standard-error of
the mean (coloured areas) across participants is shown. A non-significant interaction (ns)
between block number and drug treatment was found, as determined by linear mixed-effect
*models (see Methods). **b.** Topographical maps of slow-wave density for each drug treatment.*
***c.** Topographical maps of the statistical differences in slow-wave density between MPH and*
PLA (left), ATM and PLA (middle) and CIT and PLA (right). The t-values obtained with mixed-
effect models for each electrode are shown (see Methods). Significant clusters of electrodes
($p_{cluster}$<.05) are shown with black dots (cluster-permutation approach, see Methods). A
significant cluster was found only for CIT.

Predicting behavioural errors with EEG indices of arousal

Given that alpha power decreased following the administration of MPH (Figure 5c) and

slow waves increased following the administration of CIT (Figure 6c), and since these two

treatments also showed the strongest modulations of behavioural performance (Figure 2), we

examined whether the amount of alpha power or the density of slow waves at the sensor level

could predict behavioural performance, irrespective of drug treatment, and the spatial

distributions of these effects. To do so, we fitted linear mixed effect models for each scalp

electrode using the proportion of misses, false alarms, or the average reaction times as predicted

variables, and either slow waves or alpha power as a single fixed effect. In all these models,

subject identity was fitted as a random categorical effect. We then used a cluster permutation

approach to identify clusters of electrodes for which alpha power or slow waves significantly

predicted behavioural variables (Figure 7, bottom).

This analysis revealed that an increase in alpha power was predictive of an increase in

misses in all electrodes. An increase in alpha power was also predictive of a slowing of

responses over centro-parietal electrodes. Finally, an increase in alpha power was negatively

associated with false alarms in frontal, left temporal and occipital electrodes. In all three of

these cases, topographical maps were homogeneous in the sense that, across all electrodes, the

association between alpha power and behavioural variables was always in the same direction

(positive for misses and reaction times, negative for false alarms).

Figure 7. Predicting behavioural errors with slow waves and alpha power.

We used slow waves (top row) and alpha power (bottom row), two complementary markers of physiological arousal, to predict behavioural variables: misses (left), false alarms (middle) and reaction times (right). For each predictor (slow waves or alpha power) and each behavioural variable (misses, false alarms, and reaction times), we fitted mixed-effect models at the electrode level (see Methods). The t-values derived from these models and estimating the influence of the predictor on the behavioural variable of interest are shown as topographical maps. Significant clusters of electrodes ($p_{cluster}$<.05) are shown with black dots (cluster-permutation approach, see Methods). Both alpha power and slow waves predict behavioural errors, but slow waves do so in a region-specific fashion.

Slow waves, on the other hand, had contrasting effects (Figure 7, top). An increase in temporo-occipital slow waves was predictive of more misses but an increase of frontal slow waves was associated with a reduction of missed targets. In addition, an increase in slow waves over frontal electrodes was associated with faster responses and, although there were no significant clusters over temporal or occipital electrodes, these electrodes showed a reversal in the direction of these effects (positive correlation between slow waves and reaction times). These effects suggest that slow waves can have opposite associations with behavioural errors depending on the spatial location of their occurrence.

Discussion

This study aimed to determine the relationship between physiological arousal and

sustained attention by manipulating key neuromodulators (noradrenaline, dopamine, and

serotonin) via the administration of three different drugs (MPH, ATM, and CIT) in a four-arm,

randomised, placebo-controlled trial (Figure 1).

Sustained attention is fine-tuned by neuromodulators

Although sustained attention performance classically decrements with time-on-task

(Parasuraman, 1979), we found that MPH decreased misses and false alarms, and led to faster

responses compared to placebo (Figure 2). These effects were paralleled by stable subjective

reports of sleepiness (Figure S1), meaning that participants did not experience the classical

build-up of fatigue with time-on-task. These results replicate previous findings in this dataset

(Dockree et al., 2017) and others within the literature (Bedard et al., 2015; Lufi et al., 2015;

Nandam et al., 2011; Solanto, 1998; Spencer et al., 2009). The beneficial effect of MPH could

stem from its combined effect on both dopamine and noradrenaline, neuromodulators involved

in cortical activation (Jones, 2005) and motivation (Robbins, 1997). More specifically, MPH

increases noradrenaline and dopamine levels in the prefrontal cortex (Berridge et al., 2006) and

the striatum (Volkow et al., 2012), brain regions involved in the execution and monitoring of

action (Arnsten and Li, 2005) and action selection and motivation (Liljeholm and O'Doherty,

2012) respectively.

Our results for ATM however show that increasing the concentrations of noradrenaline

and dopamine does not necessarily lead to behavioural improvements. Indeed, ATM led to

similar levels of sleepiness (Figure S1, $BF_{01}=5.8$), misses and reaction times (Figure 2, $BF_{01}>3$)

and an increase in false alarms (Figure 2), although this did not result in an overall decrease in

d' (Figure S2, $BF_{01}=1.9$). This pattern of results suggests that participants tended to become

impulsive under ATM, without making them more attentive to the target trials. This difference in results for ATM and MPH is somewhat surprising given that both target noradrenaline and dopamine. This inverted U-shaped pattern could stem from the particular responsiveness of the noradrenaline system (Graf et al., 2011; Ross and Van Bockstaele, 2021). Tailored administration of ATM doses could therefore have resulted in improvements in task performance, although this was not achieved with the fixed acute dose used here. Alternatively, the failure of ATM to improve performance in contrast with MPH could be due to the fact that ATM impacts the dorsolateral prefrontal cortex, in contrast to the broader effects of MPH on thalamo-cortical networks (Bymaster, 2002; Farr et al., 2014; Kowalczyk et al., 2019). Indeed, although MPH targets mostly prefrontal cortices, it has a more widespread cortical effect than ATM and, in particular, increases the concentration of dopamine in the striatum (Volkow et al., 2012). Thus, our results imply that increasing noradrenaline and dopamine may be more likely to improve performance when released across cortical and subcortical areas.

Finally, the administration of CIT led to a different pattern of results with participants missing more targets, without a slowing down of reaction times or an increase in false alarms (Figure 2, $BF_{01}>3$ for reaction times and false alarms). The increased misses led to a significant decrease in behavioural performance (d') compared to placebo (Figure S2) and can be interpreted as participants being more sluggish (which contrasts with the impulsivity following ATM administration). This interpretation is in line with previous work on another selective serotonin reuptake inhibitor (SSRI), paroxetine, which resulted in an increase in missed targets and reaction times (Schmitt et al., 2002). Our results may also provide indirect support for the theory that serotonin inhibits the activity of the noradrenergic and dopaminergic systems, causing impairments in behaviour (Robbins, 1997; Wingen et al., 2008).

Overall, these results show that optimal performance on a sustained attention task needs to be achieved by balancing activation and inhibition, impulsivity and sluggishness. Of course,

545 this optimal balance is achieved through the synergetic effects of various neuromodulators and

546 although the medications administered to participants were aimed at specific neuromodulatory

547 systems, additional downstream effects and interactions between these systems may play a

548 role.

Methylphenidate boosts visual processing

We investigated drug-induced changes in visual processing through two

complementary EEG indices: ERPs and SSVEPs. ERPs are a classical approach to decompose

and quantify the sub-processes involved in a particular cognitive task (Kappenman and Luck,

2011). It allows one to isolate components such as the P3, which has been associated with the

detection of rare, task-relevant events (Polich, 2007) and which is strongly elicited by the

CTET (Figure 3a). In agreement with a previous report on the same dataset (Dockree et al.,

2017), the administration of MPH robustly enhanced the amplitude of the P3 over central

electrodes (Figure 3b-c). Examining SSVEPs confirmed an enhancement of visual processing

with MPH (Figure 4b-c). This increase was maximal over central electrodes and overlapped

with the increase of the P3. This result differs this time from the previous report on the same

dataset (Dockree et al., 2017), which can be explained by the fact that the effect is observed

here over centro-parietal electrodes and not over occipital electrodes, which show the strongest

SSVEP and which were investigated in (Dockree et al., 2017). Finally, we report here that

ATM and CIT had no significant effect on the P3 or SSVEP amplitude.

Overall, these results indicate that MPH boosted visual processing, in accordance with

the improvement in behavioural performance. These results are consistent with previous

findings showing an enhancement of visual responses with increases of noradrenaline

(Gelbard-Sagiv et al., 2018). This enhancement could result from the improvement of neuronal

gain and signal-to-noise ratios with the increase in noradrenaline and dopamine levels (Thiele

and Bellgrove, 2018) . However, the fact that the effect on the frequency tag was maximal over

central electrodes, that is, away from the primary visual cortex generating the SSVEP, suggests

that higher-level, cognitive processes are at play. An increase in noradrenaline could have

increased participants' attentional focus, resulting in both stronger SSVEPs and P3 in the

cortical network tasked to detect target trials. An increase in dopamine could also heighten

motivation (Ranjbar-Slamloo and Fazlali, 2020; Thiele and Bellgrove, 2018) and lead to

similar behavioural improvements. Importantly, no significant modulation was observed for

the P3 or SSVEP for ATM and CIT, despite the impact of these drugs on performance. This

suggests that the detrimental effect of ATM on false alarms and of CIT on misses is unlikely

to result only from a degradation of the integration of visual information.

EEG markers of sleepiness are linked to changes in sustained attention

To better understand the neural mechanisms underlying the pharmacological effects on

behaviour, we examined how the different treatments impacted neural indices of arousal. We

first focused on alpha oscillations, a resting activity linked to the (active or passive) decrease

in visual processing over early visual cortices and which has been tied to modulations of

arousal. Here, we observed that alpha power was reduced following the administration of MPH.

This is in line with the effect of noradrenaline on the promotion of wakefulness (Lee and Dan,

2012; Robbins, 1997; Sara and Bouret, 2012) and the use of MPH in the treatment of

narcolepsy (Mignot, 2012). A positive effect of MPH on arousal through the increase in

noradrenaline is also in line with the stabilisation of subjective sleepiness following MPH

administration (Figure S1) and the positive behavioural and electrophysiological effects of

increased noradrenaline reported elsewhere (Gelbard-Sagiv et al., 2018).

However, the relationship between alpha power and vigilance is not monotonic and,

when individuals near sleep, alpha power also decreases. Alpha oscillations are thus a rather

ambiguous index of fatigue since a desynchronization of alpha oscillations can be associated

with both an increase and a decrease in alertness. To circumvent this limitation, we focused on

another, emerging index of fatigue: sleep-like slow waves (Andrillon et al., 2019; D'Ambrosio

et al., 2019). Indeed, recent research has shown that extended wakefulness and/or engagement

in a given task-set is associated with an increase in patterns of EEG activity reminiscent of

sleep slow waves (Bernardi et al., 2015; Hung et al., 2013; Vyazovskiy et al., 2011). These so-

called "local sleep" intrusions may result from a gradual transition of cortical activity towards

a bistable dynamic (Vyazovskiy et al., 2011). This bistable dynamic is characterised by the

occurrence of "down-states" in which neuronal assemblies are silent. These neuronal lapses,

when occurring during wakefulness, could impair the cognitive processes performed by the

impacted neuronal networks, leading to attentional lapses and errors (Andrillon et al., 2019).

To determine if sleep-like slow waves could partly explain the behavioural effects of

medications, we isolated these slow waves from the ongoing EEG activity while participants

performed the CTET. These detected slow waves had a spatio-temporal profile in line with

previous wake and sleep studies (Andrillon et al., 2021). Furthermore, we observed an increase

in the number of these sleep-like slow waves following the administration of CIT, a drug that

can be associated with sleepiness although it is less sedative than other classes of

antidepressants (Milne and Goa, 1991). Such an increase is also in line with recent findings

linking serotonin with the build-up of sleep pressure in wakefulness (Jouvet, 1999; Oikonomou

et al., 2019). Thus, an increase in serotonin levels through the administration of CIT could have

favoured the occurrence of these slow waves by shifting cortical dynamics towards bistability.

The fact that this increase was observed in nearly all scalp electrodes suggests a systemic

impact of serotonin.

Indexes of arousal can predict behavioural performance

A key aspect of the local sleep framework is that slow waves are predicted to be

mechanistically linked with the behavioural consequences of fatigue (Andrillon et al., 2019, p.

201). This is supported by their use-dependent increase in sleep deprivation studies as well as

by their region-specific consequences. Even in awake, non-sleep deprived individuals, the

cognitive consequences of slow waves depend on the location of these slow waves, with frontal

slow waves associated with impulsivity and posterior slow waves associated with sluggishness

(Andrillon et al., 2021). These consequences can be linked back to the respective involvement

of frontal cortices in executive control and parietal cortices in sensory integration and response

preparation. Thus, one can expect slow waves to be (1) predictive of behavioural errors (2) in

a region-specific fashion. Crucially, this is in contrast with alpha oscillations, which represent

a sensor and not an effector of the transition toward sleep and low levels of alertness.

Our present findings support these hypotheses as alpha power was predictive of only a

certain type of behavioural error (sluggishness) and in a homogeneous fashion (Figure 7): all

significant electrodes were positively associated with misses and slower reaction times, and

negatively associated with false alarms. Importantly, electrodes outside these clusters always

showed the same direction of effect compared with the electrodes within the clusters,

reinforcing the homogeneity of these statistical maps. In contrast, the relationship between slow

waves and behavioural errors showed regional specificities and slow waves accounted for both

more misses and shorter reaction times (Figure 7). In particular, a small cluster of frontal

electrodes showed a negative relationship between misses and reaction times, in contrast with

posterior electrodes which showed the reverse relationship with misses. Thus, according to our

hypotheses, a unique marker of hypo-arousal could account for the heterogeneous fingerprint

of attention failures and their combination of sluggishness and impulsivity. Indeed, the

paradoxical association between frontal slow waves and an improvement in behaviour could

be explained by a decrease of inhibition (through impairment of frontal cortices), which would facilitate fast (but potentially impulsive) responses. A similar antero-posterior contrast has been found in regards to subjective experience (larger slow waves in frontal electrodes being associated with more mind wandering and larger slow waves in posterior electrodes being associated with more mind blanking) in healthy well-rested, un-medicated individuals (Andrillon et al., 2021). However, we did not observe a cluster of electrodes with a significant correlation between the occurrence of slow-waves and false alarms. This could be related to the rare occurrence of both false alarms (<1%) and slow waves (<0.1 wave/second).

Conclusion

Sustained attention is fine-tuned by a combination of neuromodulators, possibly through the maintenance of an optimal level of physiological arousal and the balance of excitation and inhibition. The effect of MPH suggests that promoting arousal can lead to improved visual processing and behaviour through the combined effect of dopamine and noradrenaline. We also found that decreasing arousal through an increase in serotonin levels led to poorer behaviour in terms of increased misses. Importantly, this behavioural impairment seems associated with a state of hypo-arousal, characterised by the intrusion of sleep-like slow waves in wakefulness. These intrusions seem to result in local, region-specific consequences on behaviour and could represent a simple, unitary mechanism accounting for the various behavioural outcomes of inattention. This research paves the way for future studies seeking to examine the link between arousal and attention in clinical populations. These results also reinforce the notion that the pattern of behavioural impairments observed in disorders such as ADHD, might stem from a dysregulation of arousal (Owens, 2005).

Materials and methods

666 We re-analysed data from a previously published study (Dockree et al., 2017), adding new

667 analyses and focusing on two additional drugs. This re-analysis was approved by the Monash

668 University Human Research Ethics Committee (Project Number: 4663).

Participants

Forty (N=40) healthy participants were recruited in accordance with the ethical

guidelines of the University of Queensland (Project Number: 2008000463). Participants were

all right-handed Caucasian males, aged 18 to 45 years ($\mu = 24.3$ years, $\sigma = 5.6$ years) who were

non-smokers, had no history of neuropsychiatric disorder, and were not under the influence of

recreational drugs or psychoactive medication at the time of the study. They were screened for

psychiatric disorders using the Mini-International Neuropsychiatric Interview Screen (Sheehan

et al., 1998) and completed the Conners' Adult ADHD Rating Scales (Conners et al., 1999).

The mean for the total ADHD symptoms in the group was 49.73 ($\sigma = 11.89$), below the

threshold for ADHD diagnosis (threshold: 65). Eight (N=8) participants were discarded from

our analyses (contraindications to the study medications: 4; data loss: 1; technical issues: 3)

leaving 32 participants for our analyses. 30/32 participants completed all four versions and the

remaining two had one missing session (but completed the placebo session: PLA: N=32; CIT:

N=31; ATM: N=31; and MPH: N=32).

Procedure and Pharmacological Intervention

A randomised, double-blinded, placebo-controlled paradigm was used in which

participants partook in four different experimental sessions. Participants were required to fast

for at least one hour prior to each session. They were randomly administered a different drug,

90 minutes prior to the commencement of cognitive testing (Figure 1A). These drugs included

ATM (60 mg), MPH (30 mg), CIT (30 mg) or a placebo (dextrose). ATM inhibits the reuptake

of noradrenaline and dopamine, and thus, is commonly used to increase noradrenaline and

dopamine levels (Bedard et al., 2015; Bymaster, 2002; Ding et al., 2014). MPH inhibits

noradrenaline and dopamine transporters, resulting in an increase in cortical noradrenaline and

striatal dopamine levels (Farr et al., 2014; Kowalczyk et al., 2019; Volkow et al., 2012). CIT,

an SSRI, increases serotonin levels (McKie et al., 2005). Doses were selected according to the

reported effects on cognition from past studies (Aron et al., 2003; Graf et al., 2011; Nandam et

al., 2014). To avoid drug interactions, each session took place at least a week apart to allow the

washout of drugs. Sessions took place at the same time of the day and same day of the week

for each participant.

After ingesting the medication, participants waited for 60 minutes. They were then

equipped with a high-density electroencephalographic cap (64 active electrodes cap,

ActiveTwo BioSemi system), which took approximately 30 minutes. This ensured a 90 minute

delay between medication administration and the commencement of testing, allowing the three

treatments to reach peak plasma levels (Chamberlain et al., 2009; Kowalczyk et al., 2019;

Nandam et al., 2014; Sauer et al., 2005). EEG recording lasted for approximately 90 minutes

whilst the participants completed two tasks, the order of which was counterbalanced: a

Continuous Temporal Expectancy Task (CTET) (Connell et al., 2009; Dockree et al., 2017)

and a standard Eriksen flanker task (Barnes et al., 2014). We focused here on the CTET data.

For each session, participants first completed a visual analogue scale (VAS) (Bond and

Lader, 1974) to report their sleepiness, calmness and contentedness (Norris, 1971) across three

intervals: immediately prior to drug administration, prior to cognitive testing (+90 minutes),

and after cognitive testing (+180 minutes). We analysed here only the results from the

sleepiness scale.

Continuous Temporal Expectancy Task (CTET)

The CTET measures sustained attention. Participants were instructed to monitor a centrally presented patterned stimuli with a grey background (Figure 1B). The stimuli consisted of a square with an 8 cm^2 dimension. This was further divided into a 10 by 10 grid of squares, each diagonally split in half into white and black. Each time the frame changed, the stimuli randomly altered its orientation by 90° in a clockwise or counter-clockwise direction, producing four different arrangements. A white cross was present at the centre of the frame that participants were instructed to fixate on to limit eye movements. Each pattern was presented for a duration of either 800 milliseconds (standard stimuli) or 1120 milliseconds (target stimuli). Between two target stimuli, 7 to 15 non-target stimuli were pseudo-randomly presented. This equated to 5.6 to 12 seconds between each target presentation. Thus, this task required participants to continuously concentrate on the stream of visual information presented to them. In addition to the transition between patterns, each pattern flickered at 25 Hz during the stimulus presentation window. This allowed the generation of steady state visually evoked potentials (SSVEP). Participants were required to monitor the duration of each stimulus, pressing a button for each target stimulus detected. Prior to starting the CTET, participants had to practice the task until their performance level reached 100%. They then completed 10 experimental blocks, with each block containing 225 frames, 18 to 22 target stimuli presentations and lasting approximately 3 minutes. Participants were given a one-minute break in between blocks.

Behavioural Data

For the CTET, we computed the proportion of missed targets (target trials which were not associated with a response) and false alarms (non-target trials associated with a response). We also computed the reaction times of correctly detected target trials. A target stimulus was

considered missed if a response was not registered from 800ms after target onset and 1600ms

after the target offset (minimal reaction time: 0.8s; maximal reaction time: 2.72s). A false alarm

was defined by the presence of a response to a non-target stimulus without any target stimulus

in the two preceding trials. Reaction times for target trials were computed from the stimulus

onset. These behavioural variables were compared across drug treatments (Figure 2).

Indexes of participants' performance were extracted for each CTET block as follows:

percentage of missed targets, percentage of false alarms and average reaction times for

correctly detected targets (in seconds). To integrate the performance on both target and non-

target trials in one single variable, we also computed the d' at the block level (Figure S2)

following the Signal Detection Theory (Macmillan, 1993).

Electroencephalography

Data acquisition. EEG data were recorded using 64 active scalp electrodes and an

ActiveTwo BioSemi amplifier. EEG was sampled at 1024 Hz. Eye-movements were recorded

with pairs of electrodes (electrooculogram, EOG) placed above and below the left eye (vertical

eye-movements), and at the outer canthus of each eye (horizontal eye-movements). EEG data

were subsequently analysed using the Fieldtrip toolbox (Oostenveld et al., 2011) and custom

code.

Pre-processing of EEG Data. Raw EEG data were epoched on a [-0.5, 1.8]s window

around stimulus onset. The average voltage computed across the entire epoch was removed

and EEG data were lowpass-filtered below 40Hz (Butterworth filter at the 4th order) and notch-

filtered at 50Hz using a Discrete Fourier Transform to remove line noise. EEG data were then

down-sampled at 256 Hz and baseline corrected ([-0.5, 0]s). These pre-processed data were

then visualised to detect faulty electrodes and artefacted trials (based on the signal variance).

The former were interpolated using the "weighted" method from Fieldtrip and the latter were

discarded. Finally, EEG data were re-referenced to the average of all electrodes and an

Independent Component Analysis (ICA) decomposition was performed to visually identify

components associated with ocular artefacts (blinks and saccades). These components were

removed from the data epoched on trials.

In parallel, we also epoched raw EEG data on experimental blocks (from -2s after the

onset of the first trial to 1s after the offset of the last trial of the block). Data were demeaned,

lowpass-filtered below 40Hz (Butterworth, 4th order), highpass-filtered above 0.1Hz

(Butterworth, 4th order) and notch-filtered at 50Hz (Discrete Fourier Transform). Data were

then resampled at 256 Hz and baseline corrected ([-0.5, 0]s from block onset). Faulty electrodes

detected on trial-epoched data were interpolated and ocular ICA components (obtained from

the data epoched on trials) were removed. Prior to ICA removal, EEG data were re-referenced

to the average of all electrodes.

We used trial-epoched data for the analysis of Event-Related Potentials (ERP) and

block-epoched data for the analysis of the Power Spectrum and the detection of Slow Waves.

For topographical analyses, peripheral electrodes (Iz, P9 and P10) were removed, resulting in

61 electrodes being analysed.

Event related potentials. Event-Related Potentials were extracted for Target and Non-

Target trials separately. Data were re-aligned according to stimuli's offset ([-0.2, 0.8]s) in order

to take into account the different duration of Target and Non-Target trials. The EEG data were

baseline-corrected from -0.2s to 0s prior to the offset. For each participant, the corresponding

ERPs were averaged across trials for the different drug treatments. In Figure 3, we show the

corresponding ERPs for electrodes Fz and Pz. Based on these ERPs and previous findings

(Dockree et al., 2017), we extracted the average amplitude of the P3 potential typically elicited

by Target trials (in comparison with non-Target trials). The P3 amplitude was defined as the

difference in amplitude between the ERP associated with Target trials and the ERP associated

with non-Target trials averaged on a [0, 0.3]s window post-offset (Figure 3a).

Spectral Analysis. We computed the power spectral density (PSD) of each EEG

electrode using data epoched on experimental blocks. This was done using Welch's method

with windows of 10s, a 50% overlap between windows and a frequency resolution of 0.1 Hz

between 2 and 40 Hz. The PSD was log-transformed (base-10 log) for further analysis. We

extracted the alpha power by averaging for each participant, session, block, and electrode the

power within the [8, 11] Hz frequency band. This estimation of the strength of alpha oscillation

was compared across drug treatments (Figure 5) and used to predict behavioural performance

(Figure 7).

We also extracted the strength of the SSVEP frequency tag by computing the signal to

noise ratio (SNR) of each frequency compared to its neighbours. This was done to identify

sharp peaks in the power spectrum (associated with the frequency tagging), departing from the

1/f trend. In practice, the PSD for each frequency was divided by the average PSD on a [-0.5,

-0.1] Hz and [0.5, 0.1] Hz window around this frequency. This ratio was then log-transformed

for further analysis (base-10 log). We extracted the SNR at 25 Hz (frequency of the SSVEP)

for each participant, session, block, and electrode. This estimation of the strength of the

frequency tagging was then compared across drug treatments (Figure 3).

Sleep-like slow waves. The presence of sleep-like slow waves in the EEG was assessed

as in Andrillon et al. (2021), based on established algorithms used to detect local sleep in sleep

(Riedner et al., 2007). In practice, EEG data re-referenced to the average of the mastoid's

electrodes (TP7 and TP9) were bandpass filtered between 1 and 10Hz using a type-2

Chebyshev filter. For each block and electrode, we then centred the filtered signal around 0 by

removing the average voltage computed across the entire block. Individual waves were then

detected by locating all the negative peaks between two negative zero-crossings, which were

respectively defined as the beginning and end of the wave. The positive peak of the wave was

defined as the maximum voltage between this beginning and end. For each wave, we computed

its duration, frequency, peak-to-peak amplitude, downward slope (from the beginning to the

negative peak) and upward slope (from the negative peak to the positive peak). Waves with a

frequency above 7Hz (i.e. outside the delta/theta range: [1, 7] Hz), a positive peak above 75µV

or within 1s of a high-amplitude event (absolute amplitude >150µV) were discarded.

Finally, we focused on large-amplitude slow waves. To do so, we computed the

amplitude threshold of the top 10% of the largest waves recorded in the placebo session using

their peak-to-peak amplitude. This threshold was computed for each individual and each

electrode and was used to define large-amplitude slow waves in all sessions. Thus, the number

of slow waves can be compared within-subject, using the placebo session as a reference. We

computed the average temporal profile and spatial distribution of these waves (Figure S3) to

confirm their resemblance to sleep-like slow waves detected in wakefulness in previous studies

as well as typical slow waves detected during sleep. We finally computed the average number

of these slow waves per minute for each experimental block of each session (Figure 6).

Statistical Analyses

VAS sleepiness scores were analysed with a two-way repeated measure ANOVA, with

time points (t=0, +90, and +180 minutes post-drug administration, Figure 1A) and treatment

(MPH, ATM, CIT and PLA) coded as within-subject effects. We discarded two participants

who did not have all four experimental sessions from this analysis. The ANOVA was

performed with the 'anova_test' function from the 'rstatix' package in R. We then examined

separately post-hoc comparisons by time points and within treatments, and corrected the

corresponding *p*-values for multiple comparisons using the Bonferroni method.

Linear mixed effect models (LMEs) were implemented in R using the 'lme4' package

(Bates et al., 2015) to determine the effects of treatments (categorical fixed effect) on several

variables of interest: proportion of misses, proportion of false alarms, reaction times on target

trials and slow wave density (averaged across all electrodes). These variables were computed

for each block of each session. Subject identity was considered a random categorical effect.

We report the estimate and p-values for the comparison of the different drugs (MPH, ATM and

CIT) with the placebo session.

When testing the influence of the time spent on the task (Figure 2), we added a fixed

effect of block number as well as an interaction component with the effect of neuromodulation.

The significance level of the interaction was assessed by comparing the model with and without

the interaction component using the R Studio 'anova' function from the 'stat' package. We

report the Likelihood Ratio Test outcomes (chi-squared and p-values) for these tests.

Examining the influence of treatments on alpha power ([8, 11]Hz log-PSD), frequency

tagging (SNR at 25Hz) and event-related potentials ([0, 0.3]s ERP amplitude post offset) for

each electrode leads to the multiplication of non-independent statistical tests. We used a

cluster-permutation approach to mitigate this issue (Maris and Oostenveld, 2007). For each

electrode, we computed the t-value and p-value corresponding to the paired comparison (t-test)

for the variable of interests (alpha power, frequency tagging and ERPs) along contrasts of

interest (MPH vs PLA, ATM vs PLA, and CIT vs PLA). To examine the effect of treatment on

slow-wave density at the electrode level, we used a similar approach but using LMEs to

consider the non-normal distribution of slow-wave density. T-values were derived from the

LMEs (with treatments as a categorical fixed effect and subject identity was considered a

random categorical effect) for the same contrasts (MPH vs PLA, ATM vs PLA, and CIT vs

PLA). Clusters of neighbouring electrodes were defined as electrodes with p-values below .05

(cluster alpha). Once the clusters were defined, a comparison was performed with permuted

datasets (N=1000 permutations) and we used a Monte-Carlo *p*-value threshold of .05 (one-

sided test) to identify the significant clusters ($p_{cluster}$). Only clusters with at least 2 electrodes

were retained and shown in Figure 3-7.

861   When analysing the effect of slow-wave density and alpha power on behaviour

(proportion of misses, proportion of false alarms, and reaction times on target trials), we

implemented linear mixed effect models (LMEs). Slow-wave density or alpha power were used

as a fixed effect and subject identity was coded as a categorical random effect. We fitted one

model per electrode, and we extracted the *t*-value and *p*-value estimated by the model for the

effect of slow-wave density or alpha power. Clusters of neighbouring electrodes were defined

as electrodes with *p*-values below .05 (cluster alpha). Once the clusters were defined, a

comparison was again performed with permuted datasets, and we used a Monte-Carlo *p*-value

threshold of .05 (one-sided test) to identify the significant clusters ($p_{cluster}$).

870   Finally, to be able to interpret null results obtained for VAS scores or behavioural

indexes of performances (misses, false alarms and reaction times), we computed the Bayes

Factor for the null hypothesis (BF_{01}) using Bayesian ANOVAs (fixed effect of drug treatment)

implemented in JASP (JASP 0.16). A BF_{01} above 3 is interpreted as significant evidence for

the null hypothesis.

Graphical representation

In Figure 2, 6a, S1 and S2, we used 'raincloud plots' to show the distribution of our variable

of interests at the sample level. These plots show a combination of boxplots, smoothed density

and individual datapoints. For boxplots, the central thick horizontal line shows the median, the

lower and upper limit of the box show the 1st and 3rd quartiles respectively, and the lower and

upper limit of the vertical lines (whiskers) show the minimum and maximum values

respectively. On the left-hand side of the boxplots, individual averages are shown as circles. A

smoothed density distribution is also shown on the right-hand side of the boxplots. These plots

were created with the raincloud plot package for R (Allen et al., 2021).

Acknowledgements

We thank Jessica Barnes for data collection. T.A. was supported by a 'Long-Term

Fellowship' from the Human Frontier Science Program (LT000362/2018-L) and an 'Ideas

Grant' from the National Health and Medical Research Council (APP2002454). R.G.O. was

supported by a Horizon 2020 European Research Council Consolidator Grant IndDecision

(865474). PD was supported by an Irish Research Council Laureate Grant (201911).

Competing Interests

The authors have no competing interests to declare.

Data and Code Availability Statement

All data will be made available by contacting the corresponding author. All code used

for data analyses will be published online.

References

896 Allen M, Poggiali D, Whitaker K, Marshall TR, van Langen J, Kievit RA. 2021. Raincloud
897   plots: a multi-platform tool for robust data visualization. *Wellcome Open Res* **4**:63.
898   doi:10.12688/wellcomeopenres.15191.2

899 Andrillon T, Burns A, Mackay T, Windt J, Tsuchiya N. 2021. Predicting lapses of attention
900   with sleep-like slow waves. *Nat Commun* **12**:3657. doi:10.1038/s41467-021-23890-7

901 Andrillon T, Kouider S. 2020. The vigilant sleeper: neural mechanisms of sensory (de)coupling
902   during sleep. *Curr Opin Physiol* **15**:47–59. doi:10.1016/j.cophys.2019.12.002

903 Andrillon T, Windt J, Silk T, Drummond SPA, Bellgrove MA, Tsuchiya N. 2019. Does the
904   mind wander when the brain takes a break? Local sleep in wakefulness, attentional
905   lapses and mind-wandering. *Front Neurosci* **13**:949. doi:10.3389/fnins.2019.00949

906 Arnau S, Löffler C, Rummel J, Hagemann D, Wascher E, Schubert A. 2020. Inter-trial alpha
907   power indicates mind wandering. *Psychophysiology* **57**. doi:10.1111/psyp.13581

908 Arnsten AF. 2010. The use of α-2A adrenergic agonists for the treatment of attention-
909   deficit/hyperactivity disorder. *Expert Rev Neurother* **10**:1595–1605.
910   doi:10.1586/ern.10.133

911 Arnsten AFT. 2000. Through the Looking Glass: Differential Noradenergic Modulation of
912   Prefrontal Cortical Function. *Neural Plast* **7**:133–146. doi:10.1155/NP.2000.133

913 Arnsten AFT, Li B-M. 2005. Neurobiology of Executive Functions: Catecholamine Influences
914   on Prefrontal Cortical Functions. *Biol Psychiatry* **57**:1377–1384.
915   doi:10.1016/j.biopsych.2004.08.019

916 Arnsten AFT, Rubia K. 2012. Neurobiological Circuits Regulating Attention, Cognitive
917   Control, Motivation, and Emotion: Disruptions in Neurodevelopmental Psychiatric
918   Disorders. *J Am Acad Child Adolesc Psychiatry* **51**:356–367.
919   doi:10.1016/j.jaac.2012.01.008

920 Aron AR, Dowson JH, Sahakian BJ, Robbins TW. 2003. Methylphenidate improves response
921   inhibition in adults with attention-deficit/hyperactivity disorder. *Biol Psychiatry*
922   **54**:1465–1468. doi:10.1016/S0006-3223(03)00609-7

923 Barnes JJM, O'Connell RG, Nandam LS, Dean AJ, Bellgrove MA. 2014. Monoaminergic
924   modulation of behavioural and electrophysiological indices of error processing.
925   *Psychopharmacology (Berl)* **231**:379–392. doi:10.1007/s00213-013-3246-y

926 Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using
927   **lme4**. *J Stat Softw* **67**. doi:10.18637/jss.v067.i01

928 Bedard AC, Stein MA, Halperin JM, Krone B, Rajwan E, Newcorn JH. 2015. Differential
929   impact of methylphenidate and atomoxetine on sustained attention in youth with
930   attention-deficit/hyperactivity disorder. *J Child Psychol Psychiatry* **56**:40–48.
931   doi:10.1111/jcpp.12272

932 Bernardi G, Siclari F, Yu X, Zennig C, Bellesi M, Ricciardi E, Cirelli C, Ghilardi MF, Pietrini
933   P, Tononi G. 2015. Neural and behavioral correlates of extended training during sleep
934   deprivation in humans: evidence for local, task-specific effects. *J Neurosci Off J Soc*
935   *Neurosci* **35**:4487–4500. doi:10.1523/JNEUROSCI.4567-14.2015

936 Berridge CW, Devilbiss DM, Andrzejewski ME, Arnsten AFT, Kelley AE, Schmeichel B,
937   Hamilton C, Spencer RC. 2006. Methylphenidate Preferentially Increases
938   Catecholamine Neurotransmission within the Prefrontal Cortex at Low Doses that
939   Enhance Cognitive Function. *Biol Psychiatry* **60**:1111–1120.
940   doi:10.1016/j.biopsych.2006.04.022

941 Bond A, Lader M. 1974. The use of analogue scales in rating subjective feelings. *Br J Med*
942   *Psychol* **47**:211–218. doi:10.1111/j.2044-8341.1974.tb02285.x

Borbély AA, Daan S, Wirz-Justice A, Deboer T. 2016. The two-process model of sleep
regulation: a reappraisal. *J Sleep Res* **25**:131–143. doi:10.1111/jsr.12371

Boutrel B, Koob GF. 2004. What Keeps Us Awake: the Neuropharmacology of Stimulants and
Wakefulness Promoting Medications. *Sleep* **27**:1181–1194.
doi:10.1093/sleep/27.6.1181

Braboszcz C, Delorme A. 2011. Lost in thoughts: Neural markers of low alertness during mind
wandering. *NeuroImage* **54**:3040–3047. doi:10.1016/j.neuroimage.2010.10.008

Bymaster F. 2002. Atomoxetine Increases Extracellular Levels of Norepinephrine and
Dopamine in Prefrontal Cortex of Rat A Potential Mechanism for Efficacy in Attention
Deficit/Hyperactivity Disorder. *Neuropsychopharmacology* **27**:699–711.
doi:10.1016/S0893-133X(02)00346-9

Cajochen C, Brunner DP, Krauchi K, Graw P, Wirz-Justice A. 1995. Power density in
theta/alpha frequencies of the waking EEG progressively increases during sustained
wakefulness. *Sleep* **18**:890–894. doi:10.1093/sleep/18.10.890

Chamberlain SR, Hampshire A, Müller U, Rubia K, del Campo N, Craig K, Regenthal R,
Suckling J, Roiser JP, Grant JE, Bullmore ET, Robbins TW, Sahakian BJ. 2009.
Atomoxetine Modulates Right Inferior Frontal Activation During Inhibitory Control:
A Pharmacological Functional Magnetic Resonance Imaging Study. *Biol Psychiatry*
65:550–555. doi:10.1016/j.biopsych.2008.10.014

Clayton MS, Yeung N, Cohen Kadosh R. 2018. The many characters of visual alpha
oscillations. *Eur J Neurosci* **48**:2498–2508. doi:10.1111/ejn.13747

Compton RJ, Gearinger D, Wild H. 2019. The wandering mind oscillates: EEG alpha power is
enhanced during moments of mind-wandering. *Cogn Affect Behav Neurosci* **19**:1184–
1191. doi:10.3758/s13415-019-00745-9

Connell RG, Dockree PM, Robertson IH, Bellgrove MA, Foxe JJ, Kelly SP. 2009. Uncovering
the neural signature of lapsing attention: Electrophysiological signals predict errors up
to 20 s before they occur. *J Neurosci* **29**:8604–8611. doi:10.1523/JNEUROSCI.5967-
08.2009

Conners CK, Erhardt D, Epstein JN, Parker JDA, Sitarenios G, Sparrow E. 1999. Self-ratings
of ADHD symptoms in adults I: Factor structure and normative data. *J Atten Disord*
3:141–151. doi:10.1177/108705479900300303

Cortese S. 2020. Pharmacologic Treatment of Attention Deficit–Hyperactivity Disorder. *N*
Engl J Med **383**:1050–1056. doi:10.1056/NEJMra1917069

D'Ambrosio S, Castelnovo A, Guglielmi O, Nobili L, Sarasso S, Garbarino S. 2019. Sleepiness
as a Local Phenomenon. *Front Neurosci* **13**:1086. doi:10.3389/fnins.2019.01086

Dimitrijevic A, Smith ML, Kadis DS, Moore DR. 2017. Cortical alpha oscillations predict
speech intelligibility. *Front Hum Neurosci* **11**. doi:10.3389/fnhum.2017.00088

Ding YS, Naganawa M, Gallezot JD, Nabulsi N, Lin SF, Ropchan J, Weinzimmer D, McCarthy
TJ, Carson RE, Huang Y, Laruelle M. 2014. Clinical doses of atomoxetine significantly
occupy both norepinephrine and serotonin transports: Implications on treatment of
depression and ADHD. *NeuroImage* **86**:164–71.
doi:10.1016/j.neuroimage.2013.08.001

Dockree PM, Barnes JJ, Matthews N, Dean AJ, Abe R, Nandam LS, Kelly SP, Bellgrove MA,
O'Connell RG. 2017. The Effects of Methylphenidate on the Neural Signatures of
Sustained Attention. *Biol Psychiatry* **82**:687–694. doi:10.1016/j.biopsych.2017.04.016

Doran SM, Van Dongen HP, Dinges DF. 2001. Sustained attention performance during sleep
deprivation: evidence of state instability. *Arch Ital Biol* **139**:253–267.

Esterman M, Rothlein D. 2019. Models of sustained attention. *Curr Opin Psychol* **29**:174–180.
doi:10.1016/j.copsyc.2019.03.005

Farr OM, Zhang S, Hu S, Matuskey D, Abdelghany O, Malison RT, Li CR. 2014. The effects of methylphenidate on resting-state striatal, thalamic and global functional connectivity in healthy adults. *Int J Neuropsychopharmacol* **17**:1177–1191. doi:10.1017/S1461145714000674

Geissler J, Romanos M, Hegerl U, Hensch T. 2014. Hyperactivity and sensation seeking as autoregulatory attempts to stabilize brain arousal in ADHD and mania? *ADHD Atten Deficit Hyperact Disord* **6**:159–173. doi:10.1007/s12402-014-0144-z

Gelbard-Sagiv H, Magidov E, Sharon H, Hendler T, Nir Y. 2018. Noradrenaline Modulates Visual Perception and Late Visually Evoked Activity. *Curr Biol* **28**:2239-2249.e6. doi:10.1016/j.cub.2018.05.051

Graf H, Abler B, Freudenmann R, Beschoner P, Schaeffeler E, Spitzer M, Schwab M, Grön G. 2011. Neural Correlates of Error Monitoring Modulated by Atomoxetine in Healthy Volunteers. *Biol Psychiatry* **69**:890–897. doi:10.1016/j.biopsych.2010.10.018

Hung CS, Sarasso S, Ferrarelli F, Riedner B, Ghilardi MF, Cirelli C, Tononi G. 2013. Local experience-dependent changes in the wake EEG after prolonged wakefulness. *Sleep* **36**:59–72. doi:10.5665/sleep.2302

Janssen TWP, Bink M, Geladé K, van Mourik R, Maras A, Oosterlaan J. 2016. A randomized controlled trial into the effects of neurofeedback, methylphenidate, and physical activity on EEG power spectra in children with ADHD. *J Child Psychol Psychiatry* **57**:633–644. doi:10.1111/jcpp.12517

Jensen O, Mazaheri A. 2010. Shaping Functional Architecture by Oscillatory Alpha Activity: Gating by Inhibition. *Front Hum Neurosci* **4**. doi:10.3389/fnhum.2010.00186

Jones BE. 2005. From waking to sleeping: Neuronal and chemical substrates. *Trends Pharmacol Sci* **26**:578–586. doi:10.1016/j.tips.2005.09.009

Jouvet M. 1999. Sleep and Serotonin An Unfinished Story. *Neuropsychopharmacology* **21**:24S-27S. doi:10.1016/S0893-133X(99)00009-3

Kalauzi A, Vuckovic A, Bojić T. 2012. EEG alpha phase shifts during transition from wakefulness to drowsiness. *Int J Psychophysiol* **86**:195–205. doi:10.1016/j.ijpsycho.2012.04.012

Kappenman ES, Luck SJ. 2011. ERP components: The ups and downs of brainwave recordings. OUP.

Killingsworth MA, Gilbert DT. 2010. A Wandering Mind Is an Unhappy Mind. *Science* **330**:932–932. doi:10.1126/science.1192439

Klimesch W. 2012. Alpha-band oscillations, attention, and controlled access to stored information. *Trends Cogn Sci* **16**:606–617. doi:10.1016/j.tics.2012.10.007

Klimesch W, Fellinger R, Freunberger R. 2011. Alpha Oscillations and Early Stages of Visual Encoding. *Front Psychol* **2**. doi:10.3389/fpsyg.2011.00118

Koda K, Ago Y, Cong Y, Kita Y, Takuma K, Matsuda T. 2010. Effects of acute and chronic administration of atomoxetine and methylphenidate on extracellular levels of noradrenaline, dopamine and serotonin in the prefrontal cortex and striatum of mice: Acute and chronic treatment with ADHD drugs. *J Neurochem* no-no. doi:10.1111/j.1471-4159.2010.06750.x

Kowalczyk OS, Cubillo AI, Smith A, Barrett N, Giampietro V, Brammer M, Simmons A, Rubia K. 2019. Methylphenidate and atomoxetine normalise fronto-parietal underactivation during sustained attention in ADHD adolescents. *Eur Neuropsychopharmacol* **29**:1102–1116. doi:10.1016/j.euroneuro.2019.07.139

Lee S-H, Dan Y. 2012. Neuromodulation of Brain States. *Neuron* **76**:209–222. doi:10.1016/j.neuron.2012.09.012

Liljeholm M, O'Doherty JP. 2012. Contributions of the striatum to learning, motivation, and
performance: an associative account. *Trends Cogn Sci* **16**:467–475.
doi:10.1016/j.tics.2012.07.007
Lufi D, Bassin-Savion S, Rubel L. 2015. The effect of methylphenidate on sustained attention
among adolescents with attention-deficit hyperactivity disorder. *Neurocase* **21**:802–
808. doi:10.1080/13554794.2015.1023317
Macdonald JSP, Mathan S, Yeung N. 2011. Trial-by-Trial Variations in Subjective Attentional
State are Reflected in Ongoing Prestimulus EEG Alpha Oscillations. *Front Psychol* **2**.
doi:10.3389/fpsyg.2011.00082
Macmillan NA. 1993. Signal detection theory as data analysis method and psychological
decision model In: Keren G, Lewis C, editors. A Handbook for Data Analysis in the
Behavioral Sciences: Methodological Issues. Lawrence Erlbaum Associates, Inc. pp.
21–57.
Maffei A, Angrilli A. 2018. Spontaneous eye blink rate: An index of dopaminergic component
of sustained attention and fatigue. *Int J Psychophysiol* **123**:58–63.
doi:10.1016/j.ijpsycho.2017.11.009
Maris E, Oostenveld R. 2007. Nonparametric statistical testing of EEG- and MEG-data. *J
Neurosci Methods* **164**:177–190. doi:10.1016/j.jneumeth.2007.03.024
McKie S, Del-Ben C, Elliott R, Williams S, del Vai N, Anderson I, Deakin JFW. 2005.
Neuronal effects of acute citalopram detected by pharmacoMRI. *Psychopharmacology
(Berl)* **180**:680–686. doi:10.1007/s00213-005-2270-y
Mignot EJM. 2012. A Practical Guide to the Therapy of Narcolepsy and Hypersomnia
Syndromes. *Neurotherapeutics* **9**:739–752. doi:10.1007/s13311-012-0150-9
Milne RJ, Goa KL. 1991. Citalopram: A Review of its Pharmacodynamic and Pharmacokinetic
Properties, and Therapeutic Potential in Depressive Illness. *Drugs* **41**:450–477.
doi:10.2165/00003495-199141030-00008
Mittner M, Hawkins GE, Boekel W, Forstmann BU. 2016. A neural model of mind wandering.
Trends Cogn Sci **20**:570–578. doi:10.1016/j.tics.2016.06.004
Nandam LS, Hester R, Bellgrove MA. 2014. Dissociable and common effects of
methylphenidate, atomoxetine and citalopram on response inhibition neural networks.
Neuropsychologia **56**:263–270. doi:10.1016/j.neuropsychologia.2014.01.023
Nandam LS, Hester R, Wagner J, Cummins TDR, Garner K, Dean AJ, Kim BN, Nathan PJ,
Mattingley JB, Bellgrove MA. 2011. Methylphenidate But Not Atomoxetine or
Citalopram Modulates Inhibitory Control and Response Time Variability. *Biol
Psychiatry* **69**:902–904. doi:10.1016/j.biopsych.2010.11.014
Nir Y, Andrillon T, Marmelshtein A, Suthana N, Cirelli C, Tononi G, Fried I. 2017. Selective
neuronal lapses precede human cognitive lapses following sleep deprivation. *Nat Med*
23:1474–1480. doi:10.1038/nm.4433
Nobili L, De Gennaro L, Proserpio P, Moroni F, Sarasso S, Pigorini A, De Carli F, Ferrara M.
2012. Local aspects of sleep: Observations from intracerebral recordings in humans.
Prog Brain Res **199**:219–232. doi:10.1016/B978-0-444-59427-3.00013-7
Norris H. 1971. The action of sedatives on brain stem oculomotor systems in man.
Neuropharmacology **10**:181–191. doi:10.1016/0028-3908(71)90039-6
O'Connell RG, Dockree PM, Robertson IH, Bellgrove MA, Foxe JJ, Kelly SP. 2009.
Uncovering the Neural Signature of Lapsing Attention: Electrophysiological Signals
Predict Errors up to 20 s before They Occur. *J Neurosci* **29**:8604–8611.
doi:10.1523/JNEUROSCI.5967-08.2009
Oikonomou G, Altermatt M, Zhang RW, Coughlin GM, Montz C, Gradinaru V, Prober DA.
2019. The serotonergic raphe promote sleep in zebrafish and mice. *Neuron* **103**:686-
701.e8. doi:10.1016/j.neuron.2019.05.038

Oken BS, Salinsky MC, Elsas SM. 2006. Vigilance, alertness, or sustained attention: Physiological basis and measurement. *Clin Neurophysiol* **117**:1885–1901. doi:10.1016/j.clinph.2006.01.017

Oostenveld R, Fries P, Maris E, Schoffelen J-M. 2011. FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Comput Intell Neurosci* **2011**:1–9. doi:10.1155/2011/156869

Owens JA. 2005. The ADHD and Sleep Conundrum: A Review. *J Dev Behav Pediatr* **26**:312–322. doi:10.1097/00004703-200508000-00011

Parasuraman R. 1979. Memory Load and Event Rate Control Sensitivity Decrements in Sustained Attention. *Science* **205**:924–927. doi:10.1126/science.472714

Peylo C, Hilla Y, Sauseng P. 2021. Cause or consequence? Alpha oscillations in visuospatial attention. *Trends Neurosci* **44**:705–713. doi:10.1016/j.tins.2021.05.004

Pfurtscheller G. 1992. Event-related synchronization (ERS): an electrophysiological correlate of cortical areas at rest. *Electroencephalogr Clin Neurophysiol* **83**:62–69. doi:10.1016/0013-4694(92)90133-3

Pfurtscheller G, Stancák A, Neuper C. 1996. Event-related synchronization (ERS) in the alpha band - An electrophysiological correlate of cortical idling: A review. *Int J Psychophysiol* **24**:39–46.

Polich J. 2007. Updating P300: An integrative theory of P3a and P3b. *Clin Neurophysiol* **118**:2128–2148. doi:10.1016/j.clinph.2007.04.019

Quercia A, Zappasodi F, Committeri G, Ferrara M. 2018. Local use-dependent sleep in wakefulness links performance errors to learning. *Front Hum Neurosci* **12**:122. doi:10.3389/fnhum.2018.00122

Ranjbar-Slamloo Y, Fazlali Z. 2020. Dopamine and Noradrenaline in the Brain; Overlapping or Dissociate Functions? *Front Mol Neurosci* **12**:334. doi:10.3389/fnmol.2019.00334

Riedner BA, Vyazovskiy VV, Huber R, Massimini M, Esser S, Murphy M, Tononi G. 2007. Sleep homeostasis and cortical synchronization: III. A high-density EEG study of sleep slow waves in humans. *Sleep* **30**:1643–57.

Riley E, Esterman M, Fortenbaugh FC, DeGutis J. 2017. Time-of-day variation in sustained attentional control. *Chronobiol Int* **34**:993–1001. doi:10.1080/07420528.2017.1308951

Robbins TW. 1997. Arousal systems and attentional processes. *Biol Psychol* **45**:57–71. doi:10.1016/S0301-0511(96)05222-2

Romei V, Brodbeck V, Michel C, Amedi A, Pascual-Leone A, Thut G. 2008. Spontaneous Fluctuations in Posterior -Band EEG Activity Reflect Variability in Excitability of Human Visual Areas. *Cereb Cortex* **18**:2010–2018. doi:10.1093/cercor/bhm229

Ross JA, Van Bockstaele EJ. 2021. The Locus Coeruleus- Norepinephrine System in Stress and Arousal: Unraveling Historical, Current, and Future Perspectives. *Front Psychiatry* **11**:601519. doi:10.3389/fpsyt.2020.601519

Samaha J, Gosseries O, Postle BR. 2017. Distinct Oscillatory Frequencies Underlie Excitability of Human Occipital and Parietal Cortex. *J Neurosci* **37**:2824–2833. doi:10.1523/JNEUROSCI.3413-16.2017

Sara SJ. 2009. The locus coeruleus and noradrenergic modulation of cognition. *Nat Rev Neurosci* **10**:211–223. doi:10.1038/nrn2573

Sara SJ, Bouret S. 2012. Orienting and reorienting: The locus coeruleus mediates cognition through arousal. *Neuron* **76**:130–141. doi:10.1016/j.neuron.2012.09.011

Satterfield JH, Dawson ME. 1971. ELECTRODERMAL CORRELATES OF HYPERACTIVITY IN CHILDREN. *Psychophysiology* **8**:191–197. doi:10.1111/j.1469-8986.1971.tb00450.x

Sauer JM, Ring BJ, Witcher JW. 2005. Clinical pharmacokinetics of atomoxetine. *Clin Pharmacokinet* **44**:571–590. doi:10.2165/00003088-200544060-00002

Schmitt JAJ, Ramaekers JG, Kruizinga MJ, van Boxtel MPJ, Vuurman EFPM, Riedel WJ.
2002. Additional dopamine reuptake inhibition attenuates vigilance impairment
induced by serotonin reuptake inhibition in man. *J Psychopharmacol (Oxf)* **16**:207–
214. doi:10.1177/026988110201600303
Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, Hergueta T, Baker R,
Dunbar GC. 1998. The Mini-International Neuropsychiatric Interview (M.I.N.I.): The
development and validation of a structured diagnostic psychiatric interview for DSM-
IV and ICD-10. *J Clin Psychiatry* **59 Suppl 20**:22-33;quiz 34-57.
Smallwood J, Brown KS, Tipper C, Giesbrecht B, Franklin MS, Mrazek MD, Carlson JM,
Schooler JW. 2011. Pupillometric Evidence for the Decoupling of Attention from
Perceptual Input during Offline Thought. *PLoS ONE* **6**:e18298.
doi:10.1371/journal.pone.0018298
Smallwood J, Schooler JW. 2015. The Science of Mind Wandering: Empirically Navigating
the Stream of Consciousness. *Annu Rev Psychol* **66**:487–518. doi:10.1146/annurev-
psych-010814-015331
Solanto MV. 1998. Neuropsychopharmacological mechanisms of stimulant drug action in
attention-deficit hyperactivity disorder: a review and integration. *Behav Brain Res*
94:127–152. doi:10.1016/S0166-4328(97)00175-7
Sparks DW, Tian MK, Sargin D, Venkatesan S, Intson K, Lambe EK. 2018. Opposing
Cholinergic and Serotonergic Modulation of Layer 6 in Prefrontal Cortex. *Front Neural*
Circuits **11**:107. doi:10.3389/fncir.2017.00107
Spencer SV, Hawk LW, Richards JB, Shiels K, Pelham WE, Waxmonsky JG. 2009. Stimulant
Treatment Reduces Lapses in Attention among Children with ADHD: The Effects of
Methylphenidate on Intra-Individual Response Time Distributions. *J Abnorm Child*
Psychol **37**:805–816. doi:10.1007/s10802-009-9316-2
Sturm W, Willmes K. 2001. On the Functional Neuroanatomy of Intrinsic and Phasic
Alertness. *NeuroImage* **14**:S76–S84. doi:10.1006/nimg.2001.0839
Thiele A, Bellgrove MA. 2018. Neuromodulation of Attention. *Neuron* **97**:769–785.
doi:10.1016/j.neuron.2018.01.008
van Schie MKM, Lammers GJ, Fronczek R, Middelkoop HAM, van Dijk JG. 2021. Vigilance:
discussion of related concepts and proposal for a definition. *Sleep Med* **83**:175–181.
doi:10.1016/j.sleep.2021.04.038
Volkow ND, Wang G-J, Tomasi D, Kollins SH, Wigal TL, Newcorn JH, Telang FW, Fowler
JS, Logan J, Wong CT, Swanson JM. 2012. Methylphenidate-Elicited Dopamine
Increases in Ventral Striatum Are Associated with Long-Term Symptom Improvement
in Adults with Attention Deficit Hyperactivity Disorder. *J Neurosci* **32**:841–849.
doi:10.1523/JNEUROSCI.4461-11.2012
Vyazovskiy VV, Olcese U, Hanlon EC, Nir Y, Cirelli C, Tononi G. 2011. Local sleep in awake
rats. *Nature* **472**:443–447. doi:10.1038/nature10009
Ward AF, Wegner DM. 2013. Mind-blanking: When the mind goes away. *Front Psychol* **4**:650.
doi:10.3389/fpsyg.2013.00650
Wingen M, Kuypers KPC, van de Ven V, Formisano E, Ramaekers JG. 2008. Sustained
attention and serotonin: A pharmaco-fMRI study. *Hum Psychopharmacol Clin Exp*
23:221–230. doi:10.1002/hup.923
1184

Supplementary Figures

Supplementary Figure 1. Effect of treatments of subjective sleepiness

Participants were asked to complete a visual analogue scale (VAS) about their subjective sleepiness at three time points during each recording session: at drug administration (t=0 minutes), before the CTET (t=90 minutes) and after the CTET (t=180 minutes). The raincloud plot (see Methods) for each drug treatment (PLA: grey, MPH: orange, ATM: purple, CIT: green) and time is shown.

Supplementary Figure 2. Effect of treatments on behavioural performance (d')

*We computed an index of behavioural performance integrating correctness on both target and non-target trials: the d' (see Methods). **a**: Sample and individual averages (PLA: N=32; CIT: N=31; ATM: N=31; and MPH: N=32 individuals) computed across all CTET blocks (N=10 blocks) for each treatment. The raincloud plot (see Methods) for each drug treatment (PLA: grey, MPH: orange, ATM: purple, CIT: green) is shown. Stars denote the significance level (ns: non-significant; (*): p < .05; (**): p < .01; (***): p < .001) of the difference between placebo and the three treatments used in this protocol (MPH vs PLA, ATM vs PLA, CIT vs PLA), as determined by a linear mixed-effect model (see Methods). **b**: Sample averages (circles) and standard-error of the mean (coloured areas) across participants for each drug treatment and block (N=10 blocks). The star denotes the significance level (*: p < .05) of the interaction between block number and drug treatment, as determined by a linear mixed-effect model.*

Supplementary Figure 3. Slow waves' waveform and spatial distribution

a: *We computed the average waveforms of the detected slow waves for a selection of midline electrodes (Frontal (Fz), Central (Cz), Parietal (Pz), and Occipital (Oz)) on a [-0.5, 1]s window. Slow waves were aligned according to their first negative zero-crossing and baseline using a [-0.5, 0]s window. The temporal profiles shown here are the averages across all participants (N=32). **b**: Topographical map showing the spatial distribution of slow waves. We extracted slow-wave density (wave/min) per electrode and show here the average across participants.*