Conserved GYXLI motif of FlhA is involved in dynamic domain motions of FlhA required for flagellar protein export

Tohru Minamino[a], Miki Kinoshita[a], Yumi Inoue[a,*], Akio Kitao[b] and Keiichi Namba[a,c,d]

[a]Graduate School of Frontier Biosciences, Osaka University, 1-3 Yamadaoka, Suita, Osaka 565-0871, Japan

[b]School of Life Science and Technology, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro, Tokyo 152-8550, Japan

[c]RIKEN Center for Biosystems Dynamics Research and SPring-8 Center, 1-3 Yamadaoka, Suita, Osaka 565-0871, Japan

[d]JEOL YOKOGUSHI Research Alliance Laboratories, Osaka University, 1-3 Yamadaoka, Suita, Osaka 565-0871, Japan

Running title: Role of Gly-368 of FlhA in flagellar assembly

[†]Present address: Department of Ophthalmology and Visual Sciences, Kyoto University Graduate School of Medicine, Kyoto, 606-8507 Japan

Address correspondence to Tohru Minamino, tohru@fbs.osaka-u.ac.jp

ABSTRACT

Flagellar structural subunits are transported via the flagellar type III secretion system (fT3SS) and assemble at the distal end of the growing flagellar structure. The C-terminal cytoplasmic domain of FlhA (FlhA$_C$) serves as a docking platform for export substrates and flagellar chaperones and plays an important role in hierarchical protein targeting and export. FlhA$_C$ consists of domains D1, D2, D3, and D4 and adopts open and closed conformations. Gly-368 of *Salmonella* FlhA is located within the highly conserved GYXLI motif and is critical for the dynamic domain motions of FlhA$_C$. However, it remains unclear how it works. Here, we report that periodic conformational changes of the GYXLI motif induce a remodeling of hydrophobic side-chain interaction networks in FlhA$_C$ and promotes the cyclic open-close domain motions of FlhA$_C$. The temperature-sensitive *flhA(G368C)* mutation stabilized a completely closed conformation at 42ºC through strong hydrophobic interactions between Gln-498 of domain D1 and Pro-667 of domain D4 and between Phe-459 of domain D2 and Pro-646 of domain D4, thereby inhibiting flagellar protein export by the fT3SS. Its intragenic suppressor mutations reorganized the hydrophobic interaction networks in the closed FlhA$_C$ structure, restoring the protein export activity of the fT3SS to a significant degree. Furthermore, the conformational flexibility of the GYXLI motif was critical for flagellar protein export. We propose that the conserved GYXLI

1   **motif acts as a structural switch to induce the dynamic domain motions of FlhA$_C$**

2   **required for efficient and rapid protein export by the fT3SS.**

IMPORTANCE

Many motile bacteria employ the flagellar type III secretion system (fT3SS) to construct flagella beyond the cytoplasmic membrane. The C-terminal cytoplasmic domain of FlhA (FlhA$_C$), a transmembrane subunit of the fT3SS, provides binding-sites for export substrates and flagellar export chaperones to coordinate flagellar protein export with assembly. FlhA$_C$ undergoes cyclic open-close domain motions. The highly conserved Gly-368 residue of FlhA is postulated to be critical for dynamic domain motions of FlhA$_C$. However, it remains unknown how it works. Here, we carried out mutational analysis of FlhA$_C$ combined with molecular dynamics simulation and provide evidence that the conformational flexibility of FlhA$_C$ by Gly-368 is important for remodeling hydrophobic side-chain interaction networks in FlhA$_C$ to facilitate its cyclic open-close domain motions, allowing the fT3SS to transport flagellar structural subunits for efficient and rapid flagellar assembly.

KEYWORDS Bacterial flagellum, flagellar assembly, FlhA, molecular dynamics simulation, type III secretion system, *Salmonella*

INTRODUCTION

Many motile bacteria utilize flagella to swim in viscous liquids and move around on solid surfaces to migrate towards more favorable environments for their survival. The flagellum of *Salmonella enterica* serovar Typhimurium (hereafter referred to as *Salmonella*) is divided into three structural parts: the basal body, which acts as a bi-directional rotary motor; the filament, which functions as a helical propeller to produce thrust; and the hook, which is a universal joint connecting the basal body and filament and transmits torque produced by the motor to the filament. Flagellar assembly begins with the basal body, followed by the hook and finally the filament. To construct the flagellum on the cell surface, the flagellar type III secretion system (fT3SS) transports flagellar structural subunits from the cytoplasm to the distal end of the growing flagellar structure (1).

The fT3SS is located at the base of the flagellum and is composed of a transmembrane export gate complex with a stoichiometry of 9 FlhA, 1 FlhB, 5 FliP, 4 FliQ, and 1 FliR, and a cytoplasmic ATPase ring complex with a stoichiometry of 12 FliH, 6 FliI, and 1 FliJ (Fig. S1A in the Supplemental material). In addition, FlgN, FliS, and FliT act as export chaperones that escort their cognate substrates from the cytoplasm to the fT3SS (2,3). The export gate complex utilizes the transmembrane electrochemical gradient of protons (H^+) as the energy source to unfold and translocate export substrates across the cytoplasmic membrane (4,5). The export gate complex requires ATP hydrolysis by the cytoplasmic ATPase ring complex to become an active

H^+/protein antiporter that couples H^+ flow with protein translocation (6–8). The fT3SS

also has a Na^+-powered backup engine to maintain the flagellar assembly process

when the cytoplasmic ATPase ring complex does not function properly (9,10).

Salmonella FlhA is composed of an N-terminal transmembrane domain with eight

transmembrane helices (FlhA$_{TM}$, residues 1–327), a compactly folded cytoplasmic

domain (FlhA$_C$, residues 362–692) and a flexible linker (FlhA$_L$, residues 328–361)

connecting these two domains (11,12) (Fig. S1B). FlhA$_{TM}$ acts as a dual-ion channel

that conducts both H^+ and Na^+. An interaction between FlhA$_L$ and FliJ activates the ion

channel of FlhA$_{TM}$, allowing the export gate complex to couple either H^+ or Na^+ flow

with the translocation of export substrate across the cytoplasmic membrane (13).

FlhA$_C$ forms a homo-nonamer in the fT3SS (14). The FlhA$_C$ ring serves as a docking

platform for export substrates and flagellar chaperones along with the C-terminal

cytoplasmic domain of FlhB and plays an important role in hierarchical protein targeting

and export for efficient flagellar assembly (15–21).

FlhA$_C$ consists of domains D1, D2, D3, and D4 (Fig. S1B) and adopts open and

closed conformations (12,22). A highly conserved hydrophobic dimple is located at the

interface between domains D1 and D2 and is directly involved in substrate recognition

(16,17). Because a large cleft exits in the interface between domains D2 and D4 in the

open form of FlhA$_C$, but not in the closed form, the chaperone-substrate complexes

can bind to the conserved hydrophobic dimple in the open form but not in the closed

form (23,24). For filament assembly, the C-terminal region of FlhA$_L$ binds to domains

D1 and D2 of its neighboring FlhA$_C$ subunit in the FlhA$_C$ ring structure to stabilize the

open conformation, allowing the chaperones-substrate complexes to efficiently bind to

the FlhA$_C$ ring (25). During hook assembly, however, FlhA$_L$ binds to the conserved

hydrophobic dimple of the open form not only to suppress premature docking of the

chaperone-substrate complexes to FlhA$_C$ but also to facilitate the export of the hook

protein FlgE (26). These observations suggest that the open form of FlhA$_C$ reflects an

active state of the fT3SS. However, little is known about the role of the closed form of

FlhA$_C$ in flagellar protein export.

The *flhA(G368C)* mutation inhibits the protein transport activity of the fT3SS at a

restrictive temperature of 42ºC but not at a permissive temperature of 30ºC (27–29).

The temperature shift-up from 30ºC to 42ºC arrests the export of flagellar proteins by

the fT3SS with the *flhA(G368C)* mutation. Molecular dynamics (MD) simulation has

shown that the *flhA(G368C)* mutation restricts cyclic open-close domain motions of

FlhA$_C$ at 42ºC and stabilizes a completely closed conformation of FlhA$_C$ (24). Gly-368

is located within the highly conserved GYXLI motif (Fig. 1A). This leads to a plausible

hypothesis that the conserved GYXLI motif may be critical for such domain motions of

FlhA$_C$ coupled with flagellar protein export.

To clarify this hypothesis, we carried out mutational analysis of FlhA$_C$ combined

with MD simulation. We provide evidence that the *flhA(G368C)* mutation stabilizes

hydrophobic side-chain interactions between domains D1 and D3 and those between

domains D2 and D4, thereby suppressing dynamic open-close domain motions of

FlhA$_C$, and that its intragenic suppressor mutations induce the remodeling of the

hydrophobic interaction networks in FlhA$_C$, allowing FlhA$_C$ with the G368C mutation

(FlhA$_{C-G368C}$) to restore the dynamic open-close domain motions.

RESULTS

Effect of *flhA(G368C)* mutation on hydrophobic side-chain interaction networks

in FlhA$_C$. To address why the *flhA(G368C)* mutation stabilizes a completely closed

form of FlhA$_C$ at 42ºC, we first compared the interfaces between domains D1 and D3

and those between domains D2 and D4 in the open and closed forms of FlhA$_{C-G368C}$

obtained by MD simulation (Fig. 1B). Met-398 and Gln-498 of domain D1 make

hydrophobic contacts with Pro-550 of domain D3 and Pro-667 of domain D4,

respectively (left panel). Phe-459 of domain D2 makes a hydrophobic contact with Pro-

646 of domain D4 (left panel). Because the hydrophobic interactions between Phe-459

and Pro-646 and those between Gln-498 and Pro-667 are not seen in the open form

of FlhA$_{C-G368C}$ (right panel), we suggest that the *flhA(G368C)* mutation stabilizes these

two hydrophobic interactions at 42ºC, thereby suppressing cyclic open-close motions

of FlhA$_C$.

MD simulation of FlhA$_C$ with either G368C/K548C or G368C/F459C/K548C

mutation. The highly conserved Lys-548 residue of FlhA is located at the interface

between domains D1 and D3 and contributes to D1-D3 interactions (12) (Fig. 1B). The

flhA(K548C) mutation in domain D3 does not affect FlhA function at all. However, the

combination of *flhA(G368C)* and *flhA(K548C)* mutations results in a loss-of-function

phenotype even at 30ºC (Fig. 2A) (24). To clarify why the *flhA(G368C/K548C)* mutation

inhibits flagellar protein export at 30ºC, we analyzed the structure and dynamics of

FlhA$_{C-G368C/K548C}$ by MD simulation at 30ºC for 1.5 μs. Domains D1 and D2 behaved as

a rigid body in all cases. In contrast, domain D4 became closer to domain D2 through

conformational changes of two hinges, one between domains D1 and D3 and the other

between domains D3 and D4, when FlhA$_C$ switched from an open conformation to a

closed conformation. Therefore, we calculated the center-of-mass distance (d_{24})

between domains D2 and D4 during MD simulation (Fig. 2B).

Unlike FlhA$_{C-G368C}$, which changes its conformation back and forth between the

closed and open forms at 30ºC (24), FlhA$_{C-G368C/K548C}$ stopped the dynamic domain

motions at 30ºC after 0.3 μs and switched its conformation to a completely closed form

(Fig. 2B). When we compared the fully closed form of FlhA$_{C-G368C/K548C}$ with that of

FlhA$_{C-G368C}$, there was a significant difference in the orientation of domain D4 relative

to domain D3 (Fig. 2C, left panel). Unlike FlhA$_{C-G368C}$, in which hydrophobic interactions

between Gln-498 and Pro-667 and those between Phe-459 and Pro-646 stabilize the

closed conformation of FlhA$_{C-G368C}$ (Fig. 1B, left panel), the hydrophobic interactions of

Pro-667 with Phe-459 and Leu-461 locked the FlhA$_{C-G368C/K548C}$ structure into the fully

closed conformation at 30ºC (Fig. 2D).

It has been reported that the *flhA(F459C)* mutation restores motility of the

flhA(G368C/K548C) mutant to a considerable degree (Fig. 2A) (24), we analyzed the

effect of this mutation on the structure and dynamics of FlhA$_{C-G368C/K548C}$ by MD

simulation. The F459C substitution restored the dynamic open-close domain motions

of FlhA$_{C-G368C/K548C}$ at 30ºC (Fig. 2B). FlhA$_{C-G368C/F459C/K548C}$ also adopted a super-open

conformation (Fig. 2C, right panel) as seen in the wild-type structure at 42ºC (24).

Therefore, we conclude that the cyclic open-close domain motion of FlhA$_C$ is required

for efficient export of flagellar structural subunits by the fT3SS and that the completely

closed form of FlhA$_C$ reflects an inactive state of the fT3SS.

Isolation of pseudorevertants from the *flhA(G368C/K548C)* mutant. To clarify how

the *flhA(G368C/K548C)* double mutation locks FlhA$_C$ into the fully closed conformation

at 30ºC, pseudorevertants were isolated from the *flhA(G368C/K548C)* mutant by

streaking an overnight culture out on 0.35% soft agar plates, incubating them at 30ºC

for a few days and looking for motility halos emerging from the streak. In total five

motile colonies were purified from such halos. The motility of these pseudorevertants,

as illustrated with MMA2228KC-01 [*flhA(M365I/G368C/K548C)*] and MMA2228KC-03

[*flhA(G368C/A446E/K548C)*], was better than that of the *flhA(G368C/K548C)* strain at

30ºC (Fig. S2A in the Supplemental material). Consistently, the secretion levels of FlgD

(hook capping protein), FlgK (hook-filament junction protein), and FliC (filament protein,

flagellin) by these pseudorevertants were higher than those by the *flhA(G368C/K548C)*

mutant (Fig. S2B). All suppressor mutations were identified in FlhA$_C$: M365I and R370S

(isolated twice) in domain D1; A446E in domain D2; and P550S in domain D3 (Fig.

1B). These four mutations have also been isolated as gain-of-function mutations that

restore the motility defect of the *flhA(G368C)* mutant at 42ºC (28,29). Because Gly-

368 is located structurally far from Lys-548 (Fig. 1B), we hypothesize that the

flhA(G368C) mutation may induce remodeling of the hydrophobic side-chain

interaction networks in FlhA$_C$, thereby inhibiting the dynamic open-close domain

motions in FlhA$_{C-G368C/K548C}$.

Effect of *flhA(G368C)* and its intragenic suppressor mutations on the

hydrophobic side-chain interaction networks. There were structural differences in

domains D1 and D2 when the open and closed forms of FlhA$_{C-G368C}$ were compared

(Fig. S3 in the Supplemental material). Gly-368 forms the highly conserved GYXLI

motif along with Tyr-369, Arg-370, Leu-371, and Ile-372 (Fig. 1A), which forms a short

α-helix (Fig. S3). Cys-368 of FlhA$_{C-G368C}$ made much stronger hydrophobic contacts

with Arg-370, Leu-413 and Pro-415 of domain D1 in the completely closed form than

in its open form (Fig. 3A). As a result, the G368C mutation induced a significant

conformational change of this α-helix (Fig. S3). The R370S suppressor mutation is

located within the conserved GYXLI motif (Fig. 1A) and presumably weakened the

strong hydrophobic interactions among Cys-368, Leu-413 and Pro-415 (Fig. 3A),

thereby partially restoring the protein export activity of the fT3SS with the G368C

substitution. Therefore, we propose that the conserved GYXLI motif may act as a

structural switch to induce the cyclic open-close domain motion of FlhA$_C$.

To clarify how the *flhA(G368C)* mutation stabilizes the completely closed form of

FlhA$_C$ at 42ºC through the strong hydrophobic interactions between Gln-498 and Pro-

667 and those between Phe-459 and Pro-646, we next analyzed the effect of the

M365I, A446E, and P550S suppressor mutations on the conformational changes of

FlhA$_{C-G368C}$. Met-365 of domain D1 made a strong hydrophobic contact with Leu-492

of domain D1 in the closed form of FlhA$_{C-G368C}$ but not in its open form (Fig. 3B).

Therefore, the M365I substitution must have affected this hydrophobic interaction to

induce the conformational change of domain D1, thereby weakening the hydrophobic

interaction between Gln-498 and Pro-667 in FlhA$_{C-G368C}$. Ala-446 of domain D2

hydrophobically interacted with Gln-477 of domain D2 in both open and closed forms

of FlhA$_{C-G368C}$ (Fig. 3C), and so the A446E mutation would induce the formation of a

hydrogen bond between Glu-446 and Gln-447 as well as a hydrophobic contact. As a

result, this mutation may induce a conformational change of domain D2, affecting the

hydrophobic contact between Phe-459 and Pro-646. Pro-550 of domain D3 makes a

hydrophobic contact with Met-398 of domain D1 (Fig. 1B), and therefore the P550S

substitution should weaken the hydrophobic interactions between domains D1 and D3.

Taken all together, we hypothesize that a conformational change of the conserved

GYXLI motif may induce a remodeling of hydrophobic side-chain interaction networks

in FlhA$_C$ required for its dynamic open-close domain motion.

Mutational analysis of residues forming hydrophobic side-chain interaction networks in FlhA$_{C-G368C}$. To test whether the *flhA(G368C)* mutation affects the hydrophobic side-chain interaction networks in FlhA$_C$, we constructed *flhA(G368C)* mutants with either L413A, P415A, P434A, F459A, Q477A, L492A, Q498A, K548A, P646A or P667A substitution and analyzed the motility of these double mutants at both 30ºC and 42ºC (Fig. 4A). These double mutations did not affect the steady expression level of FlhA as judged by immunoblotting with polyclonal anti-FlhA$_C$ antibody (Fig. 4B),

The P415A, P434A, Q477A, Q498A, and K548A substitutions reduced the motility of the *flhA(G368C)* mutant even at 30ºC (Fig. 4A). Consistently, these five substitutions reduced the secretion levels of FlgD and FlgK by the fT3SS (Fig. 4B). Furthermore, they considerably reduced the motility of the *flhA(G368C)* mutant at 42ºC (Fig. 4A). The P415A, P434A, Q477A, Q498A and K548A substitutions themselves displayed no phenotype at both 30ºC and 42ºC (Fig. 5A). Therefore, we conclude that the *flhA(G368C)* mutation affects the hydrophobic side-chain interaction networks in FlhA$_C$, thereby inducing conformational changes of domains D1 and D2.

The L413A and L492A substitutions interfered with the motility of the *flhA(G368C)* mutant at both 30ºC and 42ºC (Fig. 4A). Consistently, these two substitutions inhibited the secretion of both FlgD and FlgK by the *flhA(G368C)* mutant grown at 30ºC (Fig. 4B). The L413A mutation itself did not inhibit motility at either 30ºC or 42ºC (Fig. 5A, upper panels). Because Cys-368 makes strong hydrophobic contacts with Arg-370,

Leu-413 and Pro-415 of domain D1 in the completely closed form of FlhA$_{C-G368C}$ (Fig.

3A), we suggest that the L413A mutation stabilizes these hydrophobic contacts even

at 30ºC, thereby inhibiting the protein transport activity of the fT3SS. In contrast to the

L413A mutation, the L492A substitution itself significantly reduced motility at 42ºC but

not at 30ºC (Fig. 5A, lower panels). Consistently, this substitution inhibited the

secretion of FlgD and FlgK by the fT3SS at 42ºC but not at 30ºC (Fig. 5B). These

results indicate that the temperature shift-up from 30ºC to 42ºC induces a

conformational change of domain D1 of FlhA$_C$ by the L492A substitution, thereby

reducing the protein export activity of the fT3SS. Because Leu-492 makes a

hydrophobic contact with Met-365 in the closed form of FlhA$_{C-G368C}$ but not in its open

form (Fig. 3B), we suggest that a proper switching of the hydrophobic side-chain

interaction between Met-365 and Leu-492 is critical for flagellar protein export by the

fT3SS.

Phe-459 of FlhA is located within a highly conserved hydrophobic dimple at the

interface between domains D1 and D2 (Fig. 1B) and is directly involved in substrate

recognition (16,17,23,24). The F459A mutation reduced the motility of the *flhA(G368C)*

mutant at both 30ºC and 42ºC (Fig. 4A, upper panels). This mutation reduced the level

of FlgK secretion by the *flhA(G368C)* mutant grown at 30ºC but not the level of FlgD

secretion (Fig. 4B), indicating that the F459A substitution affects the docking of the

FlgN-FlgK chaperone-export substrate complex to FlhA$_{C-G368C}$ but not that of FlgD, in

agreement with previous reports (17,21). Because the motility of the

flhA(G368C/F459A) was much worse than that of the *flhA(F459A)* mutant (Figs. 4 and

5), we suggest that the G368C mutation induces a conformational change of the

conserved hydrophobic dimple involved in the interaction with flagellar chaperones.

Furthermore, the temperature shift-up from 30ºC to 42ºC reduced the levels of both

FlgD and FlgK secretion by the *flhA(F459A)* mutant (Fig. 5B), thereby reducing motility

at 42ºC (Fig. 5A). Therefore, we propose that a remodeling of hydrophobic side-chain

interaction networks in the conserved hydrophobic dimple may be required for efficient

and stable binding of export substrates and chaperone-substrate complexes to FlhA$_C$.

The P646A and P667A mutations, which are located within domain D4, did not

affect the motility of the *flhA(G368C)* mutant at both 30ºC and 42ºC (Fig 3A, lower

panels). These two mutations themselves showed no phenotype (Fig4A, lower panels),

indicating that these two proline residues are not critical for the FlhA function. The

structural transition from the open conformation to the closed conformation occurs in

at least two steps (24). The first conformational change occurs at the interface between

domains D3 and D4, resulting in a 13° rotation of domain D4 in the direction towards

domain D2, and the second conformational change occurs at a flexible hinge between

domains D1 and D3, allowing FlhA$_C$ to adopt the closed form. Therefore, we suggest

that the *flhA(G368C)* mutation limits the hinge movement between domains D1 and

D3 through a remodeling of the hydrophobic side-chain interaction networks of domain

D1, thereby allowing Pro-646 and Pro-667 to make the strong hydrophobic contacts

with Phe-459 and Gln-498, respectively (Fig. 1B).

Discussion

The temperature-sensitive *flhA(G368C)* mutation limits the cyclic open-close domain motions of FlhA$_C$ at 42ºC, thereby reducing the flagellar protein export by the fT3SS and therefore the cell motility (24). Gly-368 is located within the highly conserved GYXLI motif of FlhA$_C$ (Fig. 1A). This suggests that the GYXLI motif is involved in such domain motions of FlhA$_C$. To clarify the role of Gly-368 in flagellar protein export, we performed mutational analysis of FlhA$_{C-G368C}$ combined with MD simulation and showed that the *flhA(G368C)* mutation induced a remodeling of hydrophobic side-chain interaction networks in FlhA$_C$ at 42ºC, allowing Gln-498 of domain D1 and Phe-459 of domain D2 to hydrophobically interact with Pro-667 and Pro-646 of domain D4, respectively (Fig. 1B). As a result, the *flhA(G368C)* mutation not only suppresses the dynamic open-close domain motions of FlhA$_C$ but also stabilizes the completely closed conformation at 42ºC. Its suppressor mutations affected the hydrophobic side-chain interaction networks in the closed FlhA$_C$ structure, thereby restoring the protein export activity of the fT3SS containing the *flhA(G368C)* mutation to a significant degree (Fig. 3). Cys-368 hydrophobically interacts with Arg-370, Leu-413 and Pro-415 of domain D1 in the closed form of FlhA$_{C-G368C}$ but not in its open form (Fig. 3A). Therefore, we conclude that the conformational flexibility of FlhA$_C$ by Gly-368 is required for cyclic conformational changes of the conserved GYXLI motif, allowing FlhA$_C$ to undergo the

cyclic open-close domain motions through the remodeling of the hydrophobic side-chain interaction networks in $FlhA_C$.

We showed that the completely closed form of $FlhA_C$ reflected an inactive state of the fT3SS and that the cyclic open-close domain motions of $FlhA_C$ was critical for efficient and rapid flagellar protein transport by the fT3SS (Fig. 2). The FlgN, FliS, and FliT chaperones in complex with their cognate substrates bind to $FlhA_C$ at a nanomolar affinity (17,30), and such strong interactions between the chaperones and $FlhA_C$ assist protein unfolding and export by the fT3SS (31,32). The flagellar chaperones bind to the open form of $FlhA_C$ but not to the closed form (23,24). Purified $FlhA_{C-G368C}$ prefers to adopt a closed conformation even at room temperature (24). Because pull-down assays by GST affinity chromatography have revealed that the *flhA(G368C)* mutation reduces the binding affinity of $FlhA_C$ for the chaperone-substrate complex (24), we propose that the structural transition of $FlhA_C$ from the open to the closed form may induce the dissociation of empty chaperone from $FlhA_C$ for the binding of a newly delivered chaperone-substrate complex for the export of next substrate.

FliJ binds to $FlhA_L$ with high affinity and opens both the H^+ and polypeptide channels of the transmembrane export gate complex, allowing the fT3SS to couple H^+ flow through the H^+ channel with the translocation of export substrate through the polypeptide channel (Fig. S1B in the Supplemental material) (6,15,26). FliJ also binds to $FlhA_C$ with low affinity (26). It has been shown that CdsO, a FliJ homolog, binds to $CdsV_C$, a $FlhA_C$ homolog, at a large cleft between the D4 domains of neighboring

CdsV$_C$ subunits in the CdsV$_C$ ring structure (33). This suggests that FliJ also binds to

the cleft between the D4 domains of neighboring FlhA$_C$ subunits in the FlhA$_C$ ring. It

has been reported that FliJ stabilizes the interaction between FlhA$_C$ and FliT in complex

with FliD (15). This raises the possibility that the interaction of FliJ with the cleft

between the D4 domains in the FlhA$_C$ ring may stabilize the open conformation of

FlhA$_C$ subunits in the ring. To clarify this possibility, we built the models of the FlhA$_C$

ring in complex with FliJ in the open and closed forms of FlhA$_C$ based on the crystal

structure of the CdsV$_C$ ring in complex with CdsO (PDB ID: 6WA9) (Fig. 6). FliJ can

bind to the cleft between the D4 domains of neighboring FlhA$_C$ subunits stably when

FlhA$_C$ adopts the open form in the ring structure (right panel) because FliJ can contact

both FlhA$_C$ subunits. However, the FliJ binding to of FlhA$_C$ in the closed form appears

to be unstable (left panel) because the distance between the neighboring D4 domains

is longer in the closed form than that in the open form, restraining the proper contacts

of FliJ with two FlhA$_C$ subunits (middle panel). Because the open and closed forms of

FlhA$_C$ represent the active and inactive states of the fT3SS, respectively (Fig. 2), we

propose that the cyclic open-close domain motion of FlhA$_C$ is important for efficient and

robust energy coupling mechanism of the fT3SS.

MATERIALS AND METHODS

Bacterial strains, P22-mediated transduction and DNA manipulations. *Salmonella*

strains used in this study are listed in Table 1. P22-mediated transductional crosses

were performed with P22HT*int* (34). DNA manipulations were performed using

standard protocols. Site-directed mutagenesis was carried out using Prime STAR Max

Premix as described in the manufacturer's instructions (Takara Bio). All of the *flhA*

mutations were confirmed by DNA sequencing (Eurofins Genomics).

Motility assay. Fresh colonies were inoculated onto soft agar plates [1% (w/v) triptone,

0.5% (w/v) NaCl, 0.35% (w/v) Bacto agar], and then the plates were incubated at 30°C.

Secretion assay. *Salmonella* cells were grown in 5 ml of L-broth [1% (w/v) Bacto-

tryptone, 0.5% (w/v) Bacto-yeast extract, 0.5% (w/v) NaCl] at either 30ºC or 42ºC with

shaking until the cell density reached an OD_{600} of ca. 1.2–1.4. Cultures were

centrifuged to obtain cell pellets and culture supernatants. The cell pellets were

resuspended in sodium dodecyl sulfate (SDS)-loading buffer solution [62.5 mM Tris-

HCl, pH 6.8, 2% (w/v) SDS, 10% (w/v) glycerol, 0.001% (w/v) bromophenol blue]

containing 1 µl of 2-mercaptoethanol. Proteins in the culture supernatants were

precipitated by 10% trichloroacetic acid and suspended in a Tris/SDS loading buffer

(one volume of 1 M Tris, nine volumes of 1 X SDS-loading buffer solution) containing

1 µl of 2-mercaptoethanol (35). Both whole cellular proteins and culture supernatants

were normalized to a cell density of each culture to give a constant number of

Salmonella cells. After boiling proteins in both whole cellular and culture supernatant

fractions at 95ºC for 3 min, these protein samples were separated by SDS–

polyacrylamide gel electrophoresis and transferred to nitrocellulose membranes

(Cytiva) using a transblotting apparatus (Hoefer). Then, immunoblotting with polyclonal

anti-FlgD, anti-FlgK, anti-FliC or anti-FlhA$_C$ antibody was carried using iBind Flex

Western Device (Thermo Fisher Scientific) as described in the manufacturer's

instructions. Detection was performed with Amersham ECL Prime western blotting

detection reagent (Cytiva). Chemiluminescence signals were captured by a

Luminoimage analyzer LAS-3000 (GE Healthcare). At least three measurements were

performed.

Multiple sequence alignment. Multiple sequence alignment was carried out using

CLUSTAL-Ω (http://www.ebi.ac.uk/Tools/msa/clustalo/).

MD simulation. MD simulations of FlhA$_C$ with either G368C/K548C or

G368C/F459C/K548C substitution were conducted as described previously (24). The

mutant structures were constructed based on the 3A5I structure of FlhA$_C$ (12). AMBER

ff14SB force field (36) was used for the proteins. Each mutant was initially solvated in

a cubic box of SPC/E$_b$ water molecules (37) and 0.17 M KCl (38) with a margin of at

least 12 Å from the proteins to the periodic box boundaries. The simulation was

conducted with the *pmemd.cuda* module (39) of AMBER20/21 (40). MD simulations of

the G368C/K548C and G368C/F459C/K548C mutants were independently conducted

at 300 K. After energy minimization and equilibration MD for 1 ns with positional

restraints for the protein main chain atoms at 300 K and 1 atm, MD simulation without restraints were conducted for 1.5 μs.

ACKNOWLEDGEMENTS

This work was supported in part by JSPS KAKENHI Grant Numbers JP20K15749 (to M.K.), JP19H03191, JP20H05439, and JP21H05510 (to A.K.), and JP19H03182 (to T.M.). This work has also been supported by Platform Project for Supporting Drug Discovery and Life Science Research (BINDS) from AMED under Grant Number JP19am0101117 to K.N., by the Cyclic Innovation for Clinical Empowerment (CiCLE) from AMED under Grant Number JP17pc0101020 to K.N. and by JEOL YOKOGUSHI Research Alliance Laboratories of Osaka University to K.N.

REFERENCES

1. Nakamura S, Minamino T. 2019. Flagella-driven motility of bacteria. Biomolecules 9:279.

2. Minamino T. 2014. Protein export through the bacterial flagellar type III export pathway. Biochim Biophys Acta 1843:1642–1648.

3. Minamino T, Kawamoto A, Kinoshita M, Namba K. 2020. Molecular organization and assembly of the export apparatus of flagellar type III secretion systems. Curr Top Microbiol Immunol 427:91–107.

4. Minamino T, Namba K. 2008. Distinct roles of the FliI ATPase and proton motive force in bacterial flagellar protein export. Nature 451:485–488.

5. Paul K, Erhardt M, Hirano T, Blair DF, Hughes KT. 2008. Energy source of flagellar type III secretion. Nature 451:489–492.

6. Minamino T, Morimoto YV, Hara N, Namba K. 2011. An energy transduction mechanism used in bacterial type III protein export. Nat Commun 2:475 (2011).

7. Morimoto YV, Kami-ike N, Miyata T, Kawamoto A, Kato T, Namba K, Minamino T. 2016. High-resolution pH imaging of living bacterial cell to detect local pH differences. mBio 7:01911-16.

8. Minamino T, Morimoto YV, Kinoshita, M., Aldridge PD, Namba K. 2014. The bacterial flagellar protein export apparatus processively transports flagellar proteins even with extremely infrequent ATP hydrolysis. Sci Rep 4:7579.

9. Minamino T, Morimoto YV, Hara N, Aldridge PD, Namba K. 2016. The bacterial flagellar type III export gate complex is a dual fuel engine that can use both H^+ and Na^+ for flagellar protein export. PLOS Pathog 12:e1005495.

10. Minamino T, Kinoshita M, Morimoto YV, Namba K. 2021. The FlgN chaperone activates the Na^+-driven engine of the *Salmonella* flagellar protein export apparatus. Commun Biol 4:335.

11. Minamino T, Iino T, Kutsukake K. 1994. Molecular characterization of the *Salmonella typhimurium flhB* operon and its protein products. J Bacteriol 176:7630–7637.

12. Saijo-Hamano Y, Imada K, Minamino T, Kihara M, Shimada M, Kitao A, Namba K.

2010. Structure of the cytoplasmic domain of FlhA and implication for flagellar type

III protein export. Mol Microbiol 76:260–268 (2010).

13. Minamino T, Morimoto YV, Kinoshita M, Namba K. 2021. Membrane voltage-

dependent activation mechanism of the bacterial flagellar protein export apparatus.

Proc Natl Acad Sci USA 118:e2026587118.

14. Abrusci P, Vergara-Irigaray M, Johnson S, Beeby MD, Hendrixson DR, Roversi P,

Friede ME, Deane JE, Jensen GJ, Tang CM, Lea SM. 2013. Architecture of the

major component of the type III secretion system export apparatus. Nat Struct Mol

Biol 20:99–104.

15. Bange G, Kümmerer N, Engel C, Bozkurt G, Wild K, Sinning I. 2010. FlhA provides

the adaptor for coordinated delivery of late flagella building blocks to the type III

secretion system. Proc Natl Acad Sci USA 107:11295–11300.

16. Minamino T, Kinoshita M, Hara N, Takeuchi S, Hida A, Koya S, Glenwright H,

Imada K, Aldridge PD, Namba K. 2012. Interaction of a bacterial flagellar

chaperone FlgN with FlhA is required for efficient export of its cognate substrates.

Mol Microbiol 83:775–788.

17. Kinoshita M, Hara N, Imada K, Namba K, Minamino, T. 2013. Interactions of

bacterial chaperone-substrate complexes with FlhA contribute to co-ordinating

assembly of the flagellar filament. Mol Microbiol 90:1249–1261.

18. Minamino T, Inoue Y, Kinoshita M, Namba K. 2020. FliK-driven conformational

rearrangements of FlhA and FlhB are required for export switching of the flagellar

protein export apparatus. J Bacteriol 202:e00637-19.

19. Kinoshita M, Namba K, Minamino T. 2021. A positive charge region of *Salmonella*

FliI is required for ATPase formation and efficient flagellar protein export. Commun

Biol 4:464.

20. Minamino T, Kinoshita M, Inoue Y, Morimoto YV, Ihara K, Koya S, Hara N,

Nishioka N, Kojima S, Homma M, Namba K. 2016. FliH and FliI ensure efficient

energy coupling of flagellar type III protein export in *Salmonella*. MicrobiologyOpen

5:424–435.

21. Inoue Y, Morimoto YV, Namba K, Minamino T. 2018. Novel insights into the

mechanism of well-ordered assembly of bacterial flagellar proteins in *Salmonella*.

Sci Rep 8:1787.

22. Moore SA, Jia Y. 2010. Structure of the cytoplasmic domain of the flagellar

secretion apparatus component FlhA from *Helicobacter pylori*. J Biol Chem

285:21060–21069.

23. Xing Q, Shi K, Portaliou A, Rossi P, Economou A, Kalodimos CG. 2018. Structure

of chaperone-substrate complexes docked onto the export gate in a type III

secretion system. Nat Commun 9:1773.

24. Inoue Y, Ogawa Y, Kinoshita M, Terahara N, Shimada M, Kodera N, Ando T,

Namba K, Kitao A, Imada K, Minamino T.2019. Structural insight into the substrate

specificity switching mechanism of the type III protein export apparatus. Structure 27:965–976.

25. Terahara N, Inoue Y, Kodera N, Morimoto YV, Uchihashi T, Imada K, Ando T, Namba K, Minamino T. 2018. Insight into structural remodeling of the FlhA ring responsible for bacterial flagellar type III protein export. Sci Adv 4:eaao7054.

26. Inoue Y, Kinoshita M, Kida M, Takekawa N, Namba K, Imada K, Minamino T. 2021. The FlhA linker mediates flagellar protein export switching during flagellar assembly. Commun Biol 4:646.

27. Saijo-Hamano Y, Minamino T, Macnab RM, Namba K. 2004. Structural and functional analysis of the C-terminal cytoplasmic domain of FlhA, an integral membrane component of the type III flagellar protein export apparatus in *Salmonella*. J Mol Biol 343:457–466.

28. Minamino T, Shimada M, Okabe M, Saijo-Hamano Y, Imada K, Kihara M, Namba K. 2010. Role of the C-terminal cytoplasmic domain of FlhA in bacterial flagellar type III protein export. J Bacteriol 192:1929–1936.

29. Shimada M, Saijo-Hamano Y, Furukawa Y, Minamino T, Imada K, Namba K. 2012. Functional defect and restoration of temperature-sensitive mutants of FlhA, a subunit of the flagellar protein export apparatus. J Mol Biol 415:855–865.

30. Kinoshita M, Nakanishi Y, Furukawa Y, Namba K, Imada K, Minamino T. 2016. Rearrangements of α-helical structures of FlgN chaperone control the binding affinity for its cognate substrates during flagellar type III export. Mol Microbiol

101:656–670.

31. Furukawa Y, Inoue Y, Sakaguchi A, Mori Y, Fukumura T, Miyata T, Namba K, Minamino T. 2016. Structural stability of flagellin subunit affects the rate of flagellin export in the absence of FliS chaperone. Mol Microbiol 102:405–416.

32. Minamino T, Morimoto YV, Kinoshita M, Namba K. 2021. Multiple roles of flagellar export chaperones for efficient and robust flagellar filament formation in Salmonella. Front Microbiol 12:756044.

33. Jensen J, Yamini S, Rietsch AR, Spiller BW. 2020. The structure of the type III secretion system export gate with CdsO, an ATPase lever arm. PLOS Pathog 16:e1008923.

34. Schmieger H. 1972. Phage P22 mutants with increased or decreased transduction abilities. Mol Gen Genet 119:75–88.

35. Minamino T, Macnab RM. 1999. Components of the *Salmonella* flagellar export apparatus and classification of export substrates. J Bacteriol 181:1388–1394 (1999).

36. Maier JA, Martinez C, Kasavajhala K, Wickstrom L, Hauser KE. 2015. Simmerling, C., ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. J Chem Theory Comput 11:3696–3713.

37. Takemura K, Kitao A. 2012. Water model tuning for improved reproduction of rotational diffusion and NMR spectral density. J Phys Chem B 116: 6279–6287.

38. Joung IS, Cheatham TE. 2008. Determination of alkali and halide monovalent ion

parameters for use in explicitly solvated biomolecular simulations. J Phys Chem B 112:9020–9041.

39. Gotz AW, Williamson MJ, Xu D, Poole D, Le Grand S, Walker RC. 2012. Routine microsecond molecular dynamics simulations with AMBER on GPUs. 1. generalized born. J Chem Theory Comput 8:1542–1555.

40. Case DA, Aktulga HM, Belfon K, Ben-Shalom IY, Brozell SR, Cerutti DS, Cheatham TE III, Cisneros GA, Cruzeiro VWD, Darden TA, Duke RE, Giambasu G, Gilson MK, Gohlke H, Goetz AW, Harris R, Izadi S, Izmailov SA, Jin C, Kasavajhala K, Kaymak MC, King E, Kovalenko A, Kurtzman T, Lee TS, LeGrand S, Li P, Lin C, Liu J, Luchko T, Luo R, Machado M, Man V, Manathunga M, Merz KM, Miao Y, Mikhailovskii O, Monard G, Nguyen H, O'Hearn KA, Onufriev A, Pan F, Pantano S, Qi R, Rahnamoun A, Roe DR, Roitberg A, Sagui C, Schott-Verdugo S, Shen J, Simmerling CL, Skrynnikov NR, Smith J, Swails J, Walker RC, Wang J, Wei H, Wolf RM, Wu X, Xue Y, York DM, Zhao S, Kollman PA. 2021. Amber 2021, University of California, San Francisco.

41. Yamaguchi S, Fujita H, Sugata K, Taira T, Iino T. 1984. Genetic analysis of *H2*, the structural gene for phase-2 flagellin in *Salmonella*. J Gen Microbiol 130:255–265.

42. Hara N, Namba K, Minamino T. 2011. Genetic characterization of conserved charged residues in the bacterial flagellar type III export protein FlhA. PLOS ONE 6:e22417.

43. Ohnishi K, Fan F, Schoenhals GJ, Kihara M, Macnab RM. 1997. The FliO, FliP,

FliQ, and FliR proteins of *Salmonella typhimurium*: putative components for

flagellar assembly. J Bacteriol 179:6092–6099.

44. Kihara M, Minamino T, Yamaguchi S, Macnab RM. 2001. Intergenic suppression

between the flagellar MS ring protein FliF of *Salmonella* and FlhA, a membrane

component of its export apparatus. J Bacteriol 183:1655–1662.

Table 1. Strains and plasmids used in this study

Strain/Plasmid	Relevant characteristics	References
Salmonella		
SJW1103	Wild-type for motility and chemotaxis	41
SJW2228	*flhA(G368C)*	27
NH001	Δ*flhA*	42
MMA2228KC	*flhA(G368C/K548C)*	This study
MMA2228FKC	*flhA(G368C/F459C/K548C)*	This study
MMA2228KC-01	*flhA(M365I/G368C/K548C)*	This study
MMA2228KC-02	*flhA(G368C/R370S/K548C)*	This study
MMA2228KC-03	*flhA(G368C/A446E/K548C)*	This study
MMA2228KC-04	*flhA(G368C/K548C/P550S)*	This study
Plasmid		
pTrc99AFF4	Expression vector	43
pMM130	pTrc99AFF4/ FlhA	44
pYI130(G368C)	pTrc99AFF4/ FlhA(G368C)	24
pMKM130(L413A)	pTrc99AFF4/ FlhA(L413A)	This study
pMKM130(G368C/L413A)	pTrc99AFF4/ FlhA(G368C/L413A)	This study
pMKM130(P415A)	pTrc99AFF4/ FlhA(L415A)	This study
pMKM130(G368C/P415A)	pTrc99AFF4/ FlhA(G368C/P415A)	This study
pMKM130(P434A)	pTrc99AFF4/ FlhA(P434A)	This study
pMKM130(G368C/P434A)	pTrc99AFF4/ FlhA(G368C/P434A)	This study
pMKM130(F459A)	pTrc99AFF4/ FlhA(F459A)	This study
pMKM130(G368C/F459A)	pTrc99AFF4/ FlhA(G368C/F459A)	This study
pMKM130(Q477A)	pTrc99AFF4/ FlhA(Q477A)	This study
pMKM130(G368C/Q477A)	pTrc99AFF4/ FlhA(G368C/Q477A)	This study
pMKM130(L492A)	pTrc99AFF4/ FlhA(L492A)	This study
pMKM130(G368C/L492A)	pTrc99AFF4/ FlhA(G368C/L492A)	This study
pMKM130(Q498A)	pTrc99AFF4/ FlhA(Q498A)	This study
pMKM130(G368C/Q498A)	pTrc99AFF4/ FlhA(G368C/Q498A)	This study
pMKM130(K548A)	pTrc99AFF4/ FlhA(K548A)	This study
pMKM130(G368C/K548A)	pTrc99AFF4/ FlhA(G368C/K548A)	This study
pMKM130(P646A)	pTrc99AFF4/ FlhA(P646A)	This study
pMKM130(G368C/P646A)	pTrc99AFF4/ FlhA(G368C/P646A)	This study
pMKM130(P667A)	pTrc99AFF4/ FlhA(P667A)	This study
pMKM130(G368C/P667A)	pTrc99AFF4/ FlhA(G368C/P667A)	This study

FIG 1. Role of a highly conserved Gly-368 residue of FlhA in dynamic domain motions of FlhA$_C$. (A) Multiple sequence alignments of the conserved GYXLI motif of FlhA homologs. Multiple sequence alignment was carried out by Clustal Omega. Conserved residues are highlighted in red. A highly conserved glycine residue is indicated by an asterisk. UniProt Accession numbers: *Salmonella*, P40729; *Vibrio*, A0A6F8WT44; *Helicobacter*, O06758, *Caulobacter*, Q03845; *Bacillus*, P35620. (B) The closed (left panel) and open (right panel) forms of FlhA$_{C\text{-}G368C}$ obtained by MD simulation. The Cα ribbon diagram of FlhA$_{C\text{-}368C}$ with residues of intragenic suppressor mutations that allow FlhA(G368C) to exert its export function at 42ºC. The intragenic suppressor mutations are highlighted in red. Phe-459 and Gln-498 make hydrophobic contacts with Pro-646 and Pro-667, respectively, in the closed from but not in the open form. The Cα backbone is color-coded from blue to red, going through the rainbow colors from the N- to the C-terminus.

FIG 2. MD simulation of FlhA$_C$ with either G368C/K548C or G368C/F459C/K548C.
(A) Motility of SJW1103 (wild type indicated as WT), SJW2228 [*flhA(G368C)* indicated as G368C], MMA2228KC [*flhA(G368C/K548C)* indicated as G368C/K548C] and MMA2228FKC [*flhA(G368C/F459C/K548C)* indicated as G368C/F459C/K548C] in soft agar. Plates were incubated at 30ºC for 6 hours. (B) Center-of-mass distance between domains D2 and D4 (d_{24}) during 1.5 µs MD simulation. (C) Structural comparisons between completely closed forms of FlhA$_{C-G368C}$ (magenta) and FlhA$_{C-G368C/K548C}$ (green) (left panels) and between the open forms of wild-type FlhA$_C$ (blue) and FlhA$_{C-G368C/F459C/K548C}$ (cyan). (D) Interfaces between domains D1 and D3 and between domains D2 and D4 of FlhA$_{C-G368C/K548C}$.

FIG 3. Structural comparison between closed (right panels) and open conformations (right panels) of FlhA$_{C-G368C}$ obtained by MD simulation. (A) Hydrophobic interactions of Cys-368 with Arg-370, Leu-413, and Pro-415 in the open and closed forms of FlhA$_{C-G368C}$. (B) Hydrophobic interactions between Met-365 and Leu-492. (C) Interaction between Gln-477 and Ala-446.

FIG 4. Effect of alanine substitution in residues involved in remodeling of hydrophobic side-chain interaction networks in FlhA$_{C\text{-}G368C}$ on flagellar protein export by the *flhA(G368C)* mutant. (A) Motility of NH001 ($\Delta flhA$) carrying pTrc99AFF4 (indicated as V), pMM130 (indicated as WT), pYI130(G368C) (indicated as G368C), pMKM130(G368C/L413A) (indicated as G368C/L413A), pMKM130(G368C/P415A) (indicated as G368C/P415A), pMKM130(G368C/P434A) (indicated as G368C/P434A), pMKM130(G368C/F459A) (indicated as G368C/F459A), pMKM130(G368C/Q477A) (indicated as G368C/Q477A), pMKM130(G368C/L492A) (indicated as G368C/L492A), pMKM130(G368C/Q498A) (indicated as G368C/Q498A), pMKM130(G368C/K548A) (indicated as G368C/K548A), pMKM130(G368C/P646A) (indicated as G368C/P646A) or pMKM130(G368C/P667A) (indicated as G368C/P667A) in 0.35% soft agar. Plates were incubated at 30ºC for 6.5 hours and at 42ºC for 6 hours. (B) Secretion assays of FlgD and FlgK. Immunoblotting, using polyclonal anti-FlgD (1st row), anti-FlgK (2nd row), or anti-FlhA$_C$ (3rd row) antibody, of whole cell proteins (Cell) and culture supernatants (Sup) prepared from the above strains, which were exponentially grown at 30ºC with shaking. The positions of FlgD, FlgK, and FlhA are indicated by arrows. Molecular mass markers (kDa) are shown on the left.

FIG 5. Effect of temperature on the protein export function of FlhA containing a single alanine substitution. (A) Motility of NH001 (Δ*flhA*) carrying pTrc99AFF4 (indicated as V), pMM130 (indicated as WT), pYI130(G368C) (indicated as G368C), pMKM130(L413A) (indicated as L413A), pMKM130(P415A) (indicated as P415A), pMKM130(P434A) (indicated as P434A), pMKM130(F459A) (indicated as F459A), pMKM130(Q477A) (indicated as Q477A), pMKM130(L492A) (indicated as L492A), pMKM130(Q498A) (indicated as Q498A), pMKM130(K548A) (indicated as K548A), pMKM130(P646A) (indicated as P646A)or pMKM130(P667A) (indicated as P667A) in 0.35% soft agar. Plates were incubated at 30ºC and 42ºC for 5 hours. (B) Secretion assays of FlgD and FlgK. Immunoblotting, using polyclonal anti-FlgD (1st row) or anti-FlgK (2nd row) antibody, of whole cell proteins (Cell) and culture supernatants (Sup) prepared from NH001 carrying pTrc99A, pMM130, pYI130(G368C), pMKM130(F459A), or pMKM130(L492A) which were exponentially grown at either 30ºC (left panels) or 42ºC (right panels). The positions of FlgD, and FlgK are indicated by arrows. Molecular mass markers (kDa) are shown on the left.

FIG. 6. Model for of the closed (left panel, magenta) and open (right panel, cyan) forms of the FlhA$_C$ ring with FliJ. The ring model was made by fitting domains D1 and D2 of the open form of FlhA$_C$ (PDB ID: 3A5I) and its closed form obtained by MD simulation to those of MxiA$_C$ in the nonameric ring structure (PDB ID: 4A5P). The CdsO-CdsV$_C$ complex structure (PDB ID: 6WA9) was superimposed on the FlhA$_C$ ring models, and then FliJ (green) (PDB ID: 3AJE) was superimposed on the 6WA9 structure to build the FlhA$_C$-FliJ ring complex. Two FlhA$_C$ subunits in the ring model are shown. FliJ can bind to a cleft between domains D4 of neighboring FlhA$_C$ subunits in the open form of the FlhA$_C$ ring (right panel). Because a distance between the D4 domains is longer in the closed form than in the open form (center panel), FliJ presumably cannot bind to the cleft in the closed form.