Paternity sharing in insects with female competition for nuptial gifts

Jessica H. Browne[1,2]

Darryl T. Gwynne[1,3]

[1] Department of Ecology and Evolutionary Biology, University of Toronto Mississauga,

Mississauga, Ontario L5L 1C6, Canada

[2] E-mail: jbrowne@mta.ca, Current address: Department of Biology
Mount Allison University, 62 York St. Sackville, NB, E4L 1E2, Canada
[3] E-mail: darryl.gwynne@utornto.ca

When males contribute costly investments to offspring (i.e., paternal care or nuptial

gifts), they are expected to have high confidence of paternity. While studies of species with

exclusive male parental care have provided support for this hypothesis, paternity may be more

uncertain in systems where males feed each mate. While mating co-occurs with each oviposition

in many male-care systems, allowing control over insemination, mating is separated from egg-

laying in nuptial-gift systems so that males typically compete more with rivals for fertilizations.

Given the costly gift investment, and long refractory periods that limit male ability to gain fitness

from frequent copulations, there should be strong selection to avoid cuckoldry, especially

complete paternity failure. Further, because females are expected to mate frequently to obtain

male-supplied nutrition prior to oviposition, males may be unable to avoid sperm competition

because virgins will be rare (and lower quality), and females are expected to undermine male

traits that limit remating or manipulate sperm stores. Thus, paternity sharing, in which males are

not excluded from siring offspring completely, is an expected outcome of sperm competition in

these systems. Using wild-caught females in two species (an orthopteran and a dipteran), in

which sexually selected females compete for important male nuptial gifts, we examined paternity

patterns and compared them to findings in other insects. Using microsatellite analysis of

offspring and sperm stores, we found evidence of shared paternity in both study species, where

very few males failed to sire offspring. Although paternity was not equal among sires, our

estimates of paternity bias were similar to other insects with valuable nuptial gifts and contrasted

with patterns in gryllid crickets where males supply little more than sperm and are frequently

excluded from siring offspring. Further support for paternity-sharing is the lack of last-male

sperm precedence in either of our study species.

Key words: parental investments, sperm competition, nuptial gifts, female competition, paternity

Introduction

When males provide their mates with nutrition or offspring care, females usually mate

multiply to obtain these benefits (Gwynne and Simmons 1990; Herridge et al. 2016;

Bonduriansky 2001). These investments can both have high costs for males, and be essential for

offspring production, leading females to compete for males and sometimes experience greater

sexual selection relative to males (Tobias et al. 2012; Hare and Simmons 2018) (Trivers 1972;

Thornhill 1979; Gwynne 1981; 1991; 2016; Gwynne and Simmons 1990; Simmons 1992;

Herridge et al. 2016). In species with costly paternal care (many fish and birds; Clutton-Brock

1991), female re-mating rates are limited not only by the cost of eggs but also by the availability

of parental males, which can drive fierce competition for the few mating opportunities. For

example, female Gulf pipefish (*Syngnathus scovelli*) compete for 0-4 mates (mean: 1.6-2.2) and

are subject to strong sexual selection on ornaments that attract males (Jones et al. 2001; Flanagan

et al. 2014). Caring males typically have high confidence in paternity of the offspring they invest

in, either due to control over insemination such as when unfertilized pipefish eggs are placed in

an enclosed male brood pouch (Jones and Avise 2001) or when there are repeated copulations

prior to each egg laid (*Abetus* water bugs; Smith 1979a, and sequentially polyandrous birds;

Møller and Birkhead 1991; Schamal et al. 2004; Owens et al. 1995; Delehanty et al. 1998). This

evidence has been central to parental investment theory (e.g., Trivers 1972; Westneat and

Sherman 1992; Møller and Birkhead 1993; Requena and Alonzo 2017), which suggests that high

paternity confidence facilitates the evolution of paternal care, given the fitness costs of cuckoldry

(investment without paternity).

Paternity confidence seems to be more uncertain in systems where males provide nuptial

gifts (common in insects: Thornhill 1976; Lewis et al. 2014) because there is less male control

over insemination. As insect copulation typically occurs separately from egg-laying, mate (and

thus paternity) monopolization by each of a female's mates does not co-occur with each

copulation/oviposition episode as in many male-care species. Thus, when female insects mate

multiply, each ejaculate competes with the spermatheca-stored sperm of rivals for fertilizations,

which can lead to biased paternity shares that is the essence of sperm competition theory (Parker

1970; Simmons 2001). Sperm competition risk and intensity might be especially high in species

with mating gifts, as females have a strong motivation to mate frequently (Gwynne and Simmons

1990; Herridge et al. 2016) and may be less constrained by male availability relative to species

with male care; while males providing nuptial gifts have refractory periods of a few days (e.g.,

the katydid *Requena verticalus*: Gwynne 1990), caring males typically have prolonged non-

mating periods: male pregnancy lasts several-weeks in pipefish (Berglund and Rosenqvist 2003)

and male *Abetus* water bugs carry eggs on their back for nearly a month until they hatch (Smith

1979b). Indeed, females have been observed mating with many males in several species with

nuptial gifts, while virgin females are rare (e.g., 1-7 mates in *Rhamphomyia longicauda* dance

flies; Herridge 2016, 0-6 mates in *Hemiandrus pallitarsis* weta (Browne 2021), and 1-6 in

Pholidoptera griseoaptera katydids; Parker et al. 2017).

Paternity bias, which may include unequal paternity among sires (paternity skew) and/or

paternity failure, in which some males are completely excluded from fertilizing offspring (e.g.,

only 51-66% of a female's mates sire offspring in gryllid crickets: *Laupala cerasina*; Turnell and

Shaw 2015), occurs when the ejaculates of certain males have an advantage over others. In

insects, paternity bias is frequently determined by mating order such that the last mating male

fertilizes more of a female's offspring than his rivals (Parker 1970; Lloyd 1979, Eberhard 1985;

Simmons 2001; Parker et al. 2017). This is due to processes such as (1) sperm stratification

within the spermatheca, layering of the sperm that results in the most recent sperm being used

first to fertilize eggs, (2) sperm displacement or removal where a male's sperm replaces that of

past mates, and (3) mating plugs or guarding that reduce the probability of female remating.

Compared to species where males contribute only sperm, males that invest heavily in mate

feeding will experience high costs in paternity losses (cuckoldry: Trivers 1972), especially if

their ability to mate again is limited by multi-day refractory periods (Gwynne 1990; Perry and

Tse 2013). Not only are the costs of cuckoldry high when males invest in costly nuptial gifts, but

female reliance on male gifts for survival or egg development probably undermine the ability of

males to inhibit female remating or manipulate her sperm stores. Furthermore, males supplying

valuable gifts are probably less able to avoid or reduce sperm competition by seeking unmated

females (see Simmons 2001; Thomas and Simmons 2007) given the scarcity, and potentially low

fecundity of virgins when females mate frequently to obtain nutritious gifts (Simmons 2007;

Herridge 2016, Browne 2021, Parker et al. 2017).

In contrast, if there is shared paternity (presumably via sperm mixing and/or female

control; Simmons 2001), each investing male will only sire a portion of a female's brood, and

there is a reduced chance of complete paternity loss. Given that the costs of siring no offspring

are expected to be high in paternally investing systems, shared or near-equal paternity may

benefit male fitness by increasing the probability of siring some of a female's lifetime offspring

production. While this is contrary to predicted increased paternity bias through last male sperm

precedence in species with larger nuptial gifts (Gwynne 1984a; Simmons 2001), this prediction

of shared paternity is not unlike paternity patterns in many parental care systems. Although

caring males typically have high confidence of paternity in the offspring they care for, this does

not represent a female's lifetime production of offspring; subsequent eggs she lays are fertilized

by other caring males. Indeed, female pipefish (*Syngnathus typhle*) distribute their eggs between

several males, often within the same laying bout (Berglund et al. 1988).

Although studies of paternity in wild arthropods are limited, microsatellite analysis of

offspring has revealed evidence of shared paternity in two nuptial gift-giving insects: a katydid,

Requena verticalis, and a ladybird beetle, *Adalia bipunctata*. In both species, paternity was

shared (multiple paternity) in nearly all broods, and importantly, it appeared that all males that

mated with a female sired offspring (Simmons 2007; Haddrill et al. 2008). However, despite the

low rates of paternity failure, paternity was still skewed in *R. verticalis* (not measured in *A.*

bipunctata), as males varied in the number of offspring sired (Simmons 2007). Similarly, in

Ephippiger ephippiger, a katydid species where females can compete for spermatophylax gifts

(Ritchie et al. 1998), each comprising about 20% of a male's weight (Vahed and Gilbert 1996),

paternity was distributed unequally among sires (Hockham et al. 2004). Although the authors

concluded there is a high degree of paternity bias in this species, there was no estimate of female

mate number, meaning they could not determine whether any males were excluded from

fathering offspring.

A high degree of paternity skew and paternity failure have been reported in several wild-

caught species of gryllid crickets (Simmons and Beveridge 2010; Turnell and Shaw 2015) that

lack costly nuptial gifts, further supporting the idea that shared paternity may be an adaptive

outcome of sperm competition when females compete for male-supplied nutrition. On the other

hand, a similar level of paternity bias was found in *Pholidoptera griseoaptera,* a katydid species

where males provide small yet substantial spermatophylax gifts (about 7% of their weight;

Vahed and Gilbert 1996). Using the number of spermatodose sperm sacs to estimate female

mating frequency, Parker et al. (2017) found unequal paternity shares among sires, and also a

large proportion of males that sired no offspring. However, paternity estimates in this study were

potentially limited. As *Pholidioptera* (as well as *Ephippiger*) require several winters for eggs laid

late in the season to hatch (Hartley and Warne 1972; Ingrisch 1986), paternity analyses were

conducted only on a portion of the offspring that reached the whole-embryo stage in the lab.

Eggs requiring a diapause to continue development (about 60% of viable eggs) would not have

been included (Parker et al. 2017) and this may have inflated the number of paternity-losing

males (Gerlach et al. 2012; Fritzche and Arnqvist 2013).

Our study measures the proportion of males failing to sire offspring and the skew among

sires in an orthopteran where, for the first time, we assess paternity in all of a female's eggs, her

lifetime reproduction. We also test for evidence of paternity sharing in a well-studied dance fly

species in which ornament-displaying females compete for costly male gifts of prey that are

necessary for offspring development.

In the New Zealand "short-tailed" ground weta, *Hemiandrus pallitarsis* (Orthoptera:

Anostostomatidae), females mate frequently and obtain spermatophylax meals (adhered to her

mid-abdomen, separate from the sperm-containing ampulla; Gwynne 2002; 2004), probably to

allow them survive a period of 5-6 months without food while caring for offspring before dying

(Gwynne 2004). At the end of a mating season females lay all their eggs in a brood chamber

(hence their "short-tail" reduced ovipositor) potentially reducing a male's chances of gaining

paternity via last male sperm precedence (most insects have multiple ovipositions). In *H.*

pallitarsis, there is evidence that females are under sexual selection, which may explain an

"ornamental" secondary genitalic device (accessory organ: Gwynne 2004, 2005), that is inserted

between two main parts of the male genitalia as the gift is deposited (Gwynne, unpublished).

Sexual selection on female accessory organs may be a result of males mating with longer

accessory organs (Browne 2021), possibly because it signals higher fecundity (Gwynne 2005).

Alternatively, longer organs may increase copulation duration thus allowing larger nuptial gifts

to be obtained, as suggested for the function of the "female penis" in certain psocopterans

(Yoshizawa et al. 2014; Yoshizawa et al. 2018) and the ridges on the subgenital plates of some

katydids with sexual competition among females (Gwynne 2001).

In the empid dance fly (Diptera: Empidae), *Rhamphomyia longicauda*, males provide

females with prey items (usually small insects) in exchange for mating. Females rely on these

mating gifts to obtain protein for egg development (Downes 1970; Hunter and Bussière 2019)

and mate multiply, with an estimated 1-7 males (mean: 2.5; Herridge 2016). Females possess

two sex-specific ornaments (pinnate scales on the legs and inflatable abdominal sacs) that

function in attracting prey-carrying males in mating swarms at dusk or dawn. Although studies

have not shown directional sexual selection on females in this species (See Wheeler et al. 2012;

Herridge 2016), there is evidence that males prefer females with larger ornaments (Funk and

Tallamy 2000; Murray et al. 2018), even though ornament size correlates weakly with egg

number and size (Wheeler 2008; Funk and Tallamy 2000). While we were unable to determine

paternity for a female's lifetime reproduction, *R. longicauda* is ideal for testing whether paternity

is shared, since females possess a single sclerotized spermatheca, creating high potential for

sperm displacement and thus biased paternity in favor of the last male (Simmons 2001). Further,

because females rely on prey gifts to complete ovarian development (Downes 1970; Hunter and

Bussière 2019), male behaviours that reduce the probability of future matings (gift acquisition)

by females are unlikely to be adaptive for either sex, due to the fitness of their offspring being

compromised. In both these species, we expect lower paternity bias and less last-male sperm

precedence compared to most insects. We predict similar paternity shares among sires, and

importantly, that few males will fail to fertilize offspring due to the fitness costs, given their

valuable investment.

Materials and Methods

Collection and rearing

We collected 17 *H. pallitarsis* pairs (females and their (last) mate guarding male) in 2017

from Kiriwhakapapa trail near Masterton, NZ (-40.807627, 175.546532) and two private gardens

in Palmerston North, NZ (-40.413909, 175.662814). Another 10 pairs were collected in 2002 by

D. Gwynne from the same locations. Males were preserved at -20°C for DNA analysis and live

females were transported to our lab in Canada to oviposit eggs. This was an extended process

that involved placing females in artificial brood chambers formed in potter's clay (2002) or soil

(2017) and allowing 2-3 months for females to lay eggs, as well as an additional 5-6 months for

eggs to mature. While in the brood chambers, we exposed females to typical winter soil

temperatures, which were based on New Zealand weather records for March to September. All

but 3 females laid eggs, however these only developed (hatched nymphs or visible eye spots) in

19 of the broods. After eggs began hatching, we collected the offspring from 17 broods

(excluding two where less than 5 offspring developed) and preserved them along with their

mothers at -20°C. We dissected spermathecae from females and isolated the sperm contents

using >70% ethanol, which causes sperm to harden into a pellet (Tripet et al. 2001) and stored

them at -20°C.

Similarly, we collected *R. longicauda* mating pairs (in copula, n=131) from mating swarms

near the Credit River, Glen Williams, Ontario, Canada (43.6865660, -79.9260960) from mid-

June to mid-July of 2017. Males were killed and preserved via freezing while females were

placed in individual plastic containers to lay eggs. They were provided with a variety of potential

oviposition substrates as oviposition location in the wild is unknown. Mortality was high (88

females died prior to oviposition), however, 43 females laid 1-2 batches of eggs (1-89 eggs

overall) on the sides of the containers before eventually dying. Dance fly eggs, which do not

require maternal care for survival, were collected and stored in petri dishes lined with damp filter

paper. They were left to develop at room temperature for up to six days, at which point those

showing signs of maturation (i.e., darkening of the chorion, visible mouthparts, hatching) were

frozen for genetic analysis. We removed female's spermathecae and again used strong ethanol

(>70%) to isolate the sperm pellet from female tissue (as in Tripet et al. 2001).

DNA Extraction and Microsatellite Analysis

DNA was extracted from *H. pallitarsis* adult and offspring tissue using a Proteinase K

based extraction method with multiple ethanol washes. DNA was extracted from sperm pellets

using a Qiagen DNeasy Blood and Tissue Extraction kit. We followed a modified version of the

recommended protocol, which involved adding 12μL DTT to each sample to improve lysis of

sperm cells and dissolving DNA in a reduced volume (50μL) of elution buffer. Unfortunately,

DNA extractions proved to be much more difficult in dance flies. While we were able to utilize

the same Proteinase K based method to extract DNA from the bodies of *R. longicauda* females

and their most recent mate, we had difficulty extracting sufficient quantities of DNA from

offspring (developed eggs or first instar larvae). We eventually had some success using a

Chelex-based extraction with an initial homogenization step; however, we were unable to extract

DNA from sperm pellets in quantities sufficient for PCR, despite using a QIAamp DNA Mirco

kit with the addition of DTT as in Herridge (2016).

In both ground weta and dance flies, we genotyped samples at four loci using fluorescent

labelled microsatellite markers. Markers for *H. pallitarsis* were developed by Genetic Marker

Services, Brighton, UK (table 1) and characterized in two populations (see Browne 2021). We

conducted PCR in two multiplex reactions, each containing 5.75μL sterile filtered water, 1μL

10X PCR reaction buffer, 0.2μL 10mM dNTP mixture, 0.5μL of forward primer (with

fluorescent labels HEX or 6-FAM), 0.5μL reverse primer, and 1μL of template DNA.

Thermocycling was done using an Eppendorf Mastercycler with the following temperature

regime: 2 minutes at 94°C followed by 35 cycles of 30 seconds at 94°C, 30 seconds at 60°C, and

15 seconds at 72°C, then an additional 3 minutes at 72°C.

Markers for *R. longicauda* were developed and characterized by E. J. Herridge and L.

Bussière at University of Stirling for use with the same study population (table 2; Herridge

2016). We again conducted PCR in two multiplex reactions, this time each containing 5µL

sterile filtered water, 1µL 10X PCR reaction buffer, 0.1µL 50M $MgCl_2$, 0.2µL 10mM dNTP

mixture, 0.75µL forward primer (with fluorescent label on 5' end), 0.75µL reverse primer, and

2µL of template DNA (1µL for adults). The thermocycling regime was adapted from Herridge

(2016) and included 5 minutes at 95°C followed by 40 cycles of 94°C for 30 seconds, 56 °C for

30 seconds, 72°C for 25 seconds, then an additional 72°C for 10 min. We sent all PCR products

to The Centre for Applied Genomics in Toronto, Ontario for fragment analysis on an Applied

Biosystems 3730xl capillary sequencer. The resulting electropherograms were examined using

GeneMarker V1.97, which allowed us to identify fluorescence peaks, representing alleles.

Paternity Analysis

Because of difficulties genotyping dance fly offspring, methods of paternity analysis

differed between the two species. In ground weta, we used data from 11-51 (mean: 28.8)

developed offspring (95% of those collected) to estimate the number of sires in each brood. We

first estimated the minimum sire number using GERUD 2.0 (Jones 2005), a parentage program

that computes the most likely minimum father combination given the offspring genotypes across

multiple loci. Offspring that shared an allele with a female's last mate at all loci were considered

to be fathered by this male. Additionally, we estimated the most likely number of sires in each

brood using Colony Version 2.0.6.6 (Jones and Wang 2010). This program uses population allele

frequencies (determined in Browne 2021) to determine the most likely parental configuration. A

female's last mate was included in Colony as a candidate father to determine which offspring

were most likely sired by him (See Turnell and Shaw 2015 for comparison of parentage

programs).

We then estimated the number of males that failed to sire offspring by identifying alleles

in a female's sperm storage organ that were not present in any of the offspring. We used allele

counting (assumes heterozygosity; number of unique paternal alleles present in offspring divided

by 2) to estimate the minimum number of additional males for each brood. In the dance flies, our

analysis was more limited. While we genotyped 11-58 (mean: 32.6) of a female's lab-laid eggs,

this only accounted for a portion (45% on average) of those collected, and in some cases only

included two loci. Further, we could not determine which male alleles remained in the

spermatheca, as we were unable to extract DNA from female sperm pellets. Using the available

offspring data for 11 dance fly broods, we first counted the unique paternal alleles present in

offspring and divided by 2 to estimate the minimum number of sires at each locus. Next, we

determined the most likely sire configuration using Colony (Jones and Wang 2010), again

including the female's last mate as a candidate father. In the absence of DNA from female sperm

stores, we compared our estimates of sire number to estimates of mate number for this

population (determined by Herridge 2016) to assess whether males likely experience paternity

loss. We used the estimate of minimum sire number (allele counting form offspring) so that this

would be comparable to Herridge's (2016) estimate of minimum mate number (allele counting

from spermathecal stores).

Statistical Analysis

In both ground weta and dance flies, we tested whether paternity patterns deviated from a

"fair raffle" scenario (Parker et al. 1990), in which males sire equal (or near equal; Herridge

2016) proportions of offspring. We measured the evenness of paternity shares in each brood

where females mated multiply by calculating the paternity skew and plotting this against sire

number. For this we used Starr's (1984) measure of paternity skew

($\sum$(proportion offspring sired)^2), in which a value of 1 represents complete paternity bias in

favor of a particular male. For both weta and dance flies, we determined whether the observed

paternity skew was significantly higher than the skew expected if offspring were distributed

evenly to all sires (as in Simmons and Beveridge 2010; Turnell and Shaw 2015) using the 95%

confidence intervals of the best fit line as well as a paired t-test. Finally, we tested for evidence

of last male sperm precedence in both *H. pallitarsis* and *R. longicauda* by determining whether a

female's last mate fathered a greater proportion of offspring than other males, or than would be

expected by equal shares.

Results

Weta: **Sire Number**

Among the 17 *H. pallitarsis* broods, the minimum sire number (GERUD) averaged 3 ±

1.6 SD (range 1-6). Estimates of most likely sire number (Colony) were much higher with an

average of 5.9 ± 2.4 SD (range 2-10). Regardless of the method used, the number of sires

detected did not change with the number of offspring tested (GERUD: ß=0.03, R^2=0.03,

p=0.525, Colony: ß=0.08, R^2=0.12, p=0.173). The majority of females mated multiply and had

offspring sired by at least 2 males (GERUD: 83% Colony: 100%)

Weta: Paternity Bias

Additional alleles, suggestive of males that mated but did not sire offspring, were found

in the spermathecae of 5 females, however, these rarely represented more than 1 male per brood.

On average, 85.4% (binomial 95% CI: 74.6%, 93.1%) of the males that inseminated a female

fathered offspring. The rate of paternity success was similar (mean: 88.6%, binomial 95% CI:

81.9%, 93.5%) when using Colony to estimate sire number. Despite this, paternity shares among

weta sires were still biased. Overall, the paternity skew (degree to which paternity is unevenly

distributed) was significantly higher than would be expected if all sires fathered an equal number

of offspring (one-tailed paired t-test, mean difference = 0.10, t = 3.72, df = 12, p = 0.001; fig. 3).

This result was consistent when using Colony to estimate the most likely sire number (one-tailed

paired t-test, mean difference = 0.09, t = 4.17, df = 16, p = 0.0004). Paternity skew declined

linearly with the number of sires (GERUD: ß= -0.10, R^2 = 0.61, p < 0.001; Colony: B= -0.03 R^2

= 0.42, p =0.003), although this was not a better fit than an inverse (non-linear, $y=\frac{1}{x}$) relationship

(GERUD: R^2 = 0.64, p < 0.001, δ AIC < 1.0; Colony: R^2 = 0.43, p =0.002, δ AIC < 1.0).

311
312
313   **Weta: Last male sperm precedence**
314
A female's last mate did not consistently have a fertilization advantage. Based on the

minimum father combination (GERUD), last males were seldom the most successful male (42%

of broods) but fathered offspring in all but two broods (Appendix). Overall, a female's last mate

did not father a significantly greater proportion of offspring than previous males (Welch two

sample t-test: t = 0.37, df = 16.89, p = 0.357) or that expected under a fair raffle scenario (equal

shares among sires) (one-tailed paired t-test: t = -0.70, df =11, p = 0.751; fig. 4).

When using Colony to reconstruct the mostly likely paternal configuration, the last male

was excluded from a further six broods (fathering the majority of offspring in only 24% of

broods; Appendix) and overall did not father a significantly greater proportion of offspring than

previous mates (Welch two sample t-test: t = 0.55, df = 17.75, p = 0.293) or that expected under

a fair raffle scenario (equal shares among sires) (one-tailed paired t-test: t = -0.40, df =16, p =

0.651; fig. 5).

Dance flies: Sire Number

We estimated sire number in dance flies using the 11 broods from which we were able to

genotype offspring. When we used the conservative method of allele counting, the number of

sires averaged 3.6 ± 1.3 SD and ranged from 2-6 sires. Estimates of sire number were much

higher when using Colony, which averaged 13.5 ± 5.9 SD sires and ranged from 7-26. In dance

flies, the number of offspring tested did influence the number of sires we were able to detect. We

found a significant positive relationship between the number of offspring genotyped and number

of sires detected for both of our estimates, however, this relationship was much stronger when

using Colony rather than allele counting (Allele counting: ß=0.07, R^2=0.69, p=0.002; Colony:

ß=0.34, R^2=0.87, p= <0.0001). Regardless of the method used to estimate sire number, all

females mated multiply, and had offspring fertilized by at least two sires.

Dance flies: Paternity Bias

Although we could not genotype the sperm stored in female's spermathecae to estimate

the proportion of failed matings, we found evidence of paternity skew among sires, similar to the

weta. Using the most likely paternal configuration (Colony), the paternity skew was determined

to be significantly higher than would be expected if all sires fathered an equal number of

offspring (one-tailed paired t-test, mean difference = 0.03, t = 6.09, df = 10, p = <0.0001; fig. 6).

Again, paternity skew declined linearly with number of sires (fig. 6; B= -0.004, R^2 = 0.59, p <

0.003), although this was not a better fit than an inverse (non-linear, $y = \frac{1}{x}$) relationship (R^2 = 0.57,

p = 0.004, δ AIC < 2.0).

Dance flies: Last male sperm precedence

There was no evidence of last male sperm precedence in our samples. In some cases, the last male did not fertilize any offspring (Appendix), however it is unclear whether this is a result of sperm competition or our sampling methods, which may have disrupted mating pairs prior to full insemination. Overall last males fathered a significantly lower proportion of offspring than previously mated males (Welch two sample t-test: t = -4.07, df = 13.58, p = 0.001) and that expected under a fair raffle scenario (equal shares among sires) (one-tailed paired t-test: t = -3.74, df =10, p = 0.004; fig. 7).

Discussion

As predicted we show evidence of shared paternity (i.e., reduced paternity bias relative to other orthopteran insects; Simmons and Beveridge 2010; Turnell and Shaw 2015) in two insect species where ornamented females appear to compete for important nutrition provided by males during mating. In one of the species, we used female lifetime reproduction. While both species showed some degree of paternity skew among sires, paternity was shared among multiple males in most broods, and there were few males excluded from fathering offspring.

In ground weta (*H. pallitarsis*), paternity estimates using lifetime offspring production revealed a level of paternity confidence in which most investing males sired offspring. We found multiple paternity in most broods, with an average of 3.0 ± 1.6 SD (most likely estimate: 5.9 ± 2.4 SD) sires, and importantly, nearly all of the males that inseminated a female fathered offspring.

In dance flies (*R. longicauda*), multiple paternity was observed in all clutches, with offspring being shared between a high number of sires (minimum mean: 3.6, most likely

mean:13.5). Although we could not measure the proportion of males that failed to sire offspring,

it was assessed indirectly by determining if the number of mates (estimated in Herridge 2016

using same study population) exceeds the number of sires. Using the conservative method of

allele counting, Herridge (2016) found that *R. longicauda* females mated with a mean of 2.5 ±

0.15 SD (range 1-6) males. Similarly, our estimate of sire number, which was determined by

counting the unique paternal alleles in each brood, averaged 3.6 ± 1.3 SD sires and ranged from

2-6. Based on this, the number of mates does not exceed the number of sires, suggesting that few

or no males experience paternity failure in *R. longicauda*.

Our results are similar to paternity patterns found in two other species with valuable

nuptial gifts: the ladybird beetle *A. bipunctata* (Haddrill et al. 2008) and orthopteran (katydid) *R.*

verticalis, both with nutritious spermatophore meals (Simmons 2007). These insects showed

high rates of multiple paternity (in 92.5%-100% of broods, respectively) and few males

experienced paternity failure. In *R. verticalis*, analysis of stored sperm revealed that all males

that inseminated a female sired offspring (Simmons 2007) and a similar pattern was evident for

A. bipunctata, as the number of mates did not exceed the number of sires (Haddrill et al. 2008)

(similar to our finding in dance flies).

By comparison, studies of insects that lack nuptial gifts and have no apparent sexual

competition among females show much greater rates of paternity failure, as only 60% of a

female's mates sire offspring in the gryllid cricket *Telogryllus commodus*, 75% in *Telogryllus*

oceanicus (Simmons and Beveridge 2010), and 51-66% in a third gryllid *L. cerasina* (Turnell

and Shaw 2015). Together, these data support the hypothesis that paternity bias is reduced in

species with valuable nuptial gifts. We suggest this is the best way of ensuring fitness (i.e.,

preventing complete paternity loss) when male mating frequency is constrained by long

refractory periods and the fertilization success of investing males is limited by an inability to

avoid sperm competition. Although shared paternity might reduce the number offspring for

individual males, fitness costs are potentially reduced because frequently mated (fed-by-males)

females may have higher fecundity or offspring quality (Browne 2021; Gwynne 1984b; reviewed

in Arnqvist and Nilsson 2000; Vahed 1998).

Contrary to our prediction, a high degree of paternity bias was found in the nuptial gift-

giving katydid, *P. griseoaptera* (Parker et al. 2017). However, as mentioned previously, this may

also be due to the low percentage of a female's lifetime production of offspring sampled. As the

authors sampled less than 40% of a female's lab-laid eggs, this may prevent successful sires from

being detected (especially those hatched in future seasons), thus inflating the apparent rate of

paternity failure.

In both our study species, there was significant paternity skew among sires, suggesting that

offspring are not divided equally among successful males. This result is not unexpected, as it has

been observed previously in gift giving insects *A. bipunctata* (Haddrill et al. 2008), *P.*

griseoaptera (Parker et al. 2017), *E. ephippiger* (Hockham et al. 2004), and *R. verticalis*

(Simmons 2007) as well as crickets that contribute little more than sperm when mating (*T.*

commodus, *T. oceanicus*; Simmons and Beveridge 2010, and *L. cerasina*; Turnell and Shaw

2015). There are several possibilities that may explain paternity skew despite the prediction of

shared paternity. First, random error during fertilization (e.g., slow or "sloppy" sperm mixing,

sperm loss; Simmons 2001) is likely to cause small differences in paternity success among males

that results in deviation from a perfect fair raffle (Herridge 2016). Because the degree of

paternity skew was weak in both the ground weta and dance fly, the observed skew may not

represent much more bias than would be expected at under a "noisy fair raffle" scenario

(Herridge 2016). Alternatively, the observed skew could be caused by variation in number of

sperm transferred (Simmons 2001; Parker et al. 2017). In *A. bipunctata*, Haddrill et al. (2008)

used in-lab studies to determine that the observed skew among sires was due to differences in

copulation duration. Differences in insemination duration may be especially pronounced in

ground weta, since sperm transfer ends in ensiferan Orthoptera when the externally placed sperm

ampulla (Brown and Gwynne 1997) is dislodged (male short-tailed ground weta may reduce

incidences of dislodging by guarding their mates after copulation; Gwynne 2004).

Finally, we found no evidence of last male sperm precedence, a common paternity pattern in

insects (Simmons 2001), in ground weta or dance flies. This further supports the hypothesis of

reduced paternity bias and suggests that any observed skew was not due to mating order. The

large, elastic sac of the spermathecae of ensiferan Orthoptera such as ground weta likely

facilitates sperm mixing, even with frequent mating (Simmons 2001). In contrast, dance flies

have a sclerotized, non-flexible spermathecae that would be expected to cause sperm

displacement, with sperm from previous males being indirectly flushed from the storage organ

(Simmons 2001). Indeed, Herridge 2016 found that stored sperm was dominated by a particular

male in *R. longicauda* but this was not related to mating order or the resulting paternity shares.

Importantly, one reason for the lack of last male precedence in dance flies observed in our study

may be that our collection methods prevented this male from inseminating the female before

uncoupling. However, at least one of the last male's alleles were present in each clutch of

offspring.

Additional constraints to our paternity analyses in dance flies are the incomplete genotypic

data. Unlike in the ground weta, we were unable to genotype a female's entire brood at all four

loci, potentially reducing the accuracy of the parental reconstruction, as this limits the amount of

information that can be used in the likelihood estimation (Jones and Wang 2010; Gerlach et al.

2012). This may explain the positive relationship between number of offspring analyzed and

sires detected for dance flies. Further, we were unable to incorporate population allele

frequencies into estimates of sire number (as in weta) because they did not include all alleles

present in offspring (likely a consequence of high microsatellite polymorphism). Given these

limitations, estimates of the most likely sire number in dance flies, including the offspring sired

by the last male, may be subject to additional error. This may help to explain the high the upper

limit on female mating rate (26 sires), however, this rate of polyandry is not unexpected given

the biology of dance flies: Individually marked female *R. longicauda*, which may live several

weeks (unpublished data), have been observed returning to (dawn and dusk) mating swarms, up

to two times per day (R. Murray, personal communication).

Using microsatellite paternity analysis of wild-caught species where females rely on

valuable nuptial gifts, we show that most males father offspring, resulting in low paternity bias

(paternity sharing). We demonstrate this in species from two insect orders (Orthoptera and

Diptera), using measures of lifetime female reproductive success in one to capture the final

outcome of sperm competition. While sperm sharing reduces the number offspring for individual

males, it reduces the chances of siring no offspring, which may be especially costly when males

supply nuptial gifts. Thus, shared paternity, in which few males are excluded from siring

offspring may make the evolution of large nuptial gifts more likely (Sakaluk 1986). These levels

of paternity sharing may be expected in other species where females rely on costly mating gifts

to produce offspring such as many katydids (Gwynne 1981; 1984b; 1985; 1988; Simmons and

Bailey 1990) and other species of empidine dance flies (Diptera: Empididae; Downes 1970;

Cumming 1994; Bussière et al. 2008; Wheeler et al. 2012; Murray et al. 2018).

Tables

Table 1. Microsatellite loci used to genotype *H. pallitarsis* (See Browne 2021 for details)

Locus	Repeat motif	Floro-label	Multiplex
wet80	(AG)14	HEX	1
wet81	(GA)16	HEX	2
wet89	(TC)41	6-FAM	1
wet83b	(AC)6	6-FAM	2

472   **Table 2**. Microsatellite markers used to analyze *R. longicauda* (See Herridge 2016)

Marker	Repeat Motif	Floro-label	Multiplex
RL1AXKU5	(TA)9	6-FAM	1
RL1BHDMD	(AT)9	HEX	1
RL1BWMXW	(TA)11	HEX	2
RL2F2Z0L	(AT)9	6-FAM	2

Figures

Figure 1. Female *H. pallitarsis* just after mating, eating part of the spermatophylax gift. One

lobe remains adhered to her abdomen and the globular sperm package is visible at the end of her

abdomen. Photo by Darryl Gwynne.

Figure 2. Female dance fly (*Rhamphomyia longicauda*) showing inflated abdomen and leg-scale

ornaments. Photo by Heather Proctor.

Figure 3. Paternity skew occurring in broods of multiply mated *H. pallitarsis* females, plotted

against the minimum number of sires (parental reconstruction determined using GERUD; n=13)

or most likely number of sires (Colony; n=17) estimated from microsatellite analysis of

offspring. The regression lines (solid) show the relationship between number of sires and

observed paternity skew with 95% confidence intervals showing deviation from the null skew

expected when all sires father an equal number of offspring (dashed line).

Figure 4. Proportion of offspring fathered by last male relative to all other competing males in

H. pallitarsis ground weta (using GERUD – minimum sire configuration). Data are shown

relative to the proportion of offspring each male would sire if paternity shares were equal (null

paternity).

Figure 5. Proportion of offspring fathered by last male relative to all other competing males in

H. pallitarsis ground weta (using COLONY – most likely sire configuration). Data are shown

relative to the proportion of offspring each male would sire if paternity shares were equal (the

null), indicated by the dashed line.

Figure 6. Paternity skew among offspring of female *R. longicauda* (n=11) plotted against the

most likely number of sires (determined from microsatellite analysis of offspring; Colony). The

regression line (solid) shows the linear relationship between number of sires and observed

paternity skew with 95% confidence intervals showing deviation from the null skew expected

when all sires father an equal number of offspring (dashed line). Size of points represents the

number of offspring analyzed for each estimate of paternity skew, which has a positive effect on

the number of sires detected (see main text).

Figure 7. Proportion of offspring fathered by last male relative to all other competing males in *R.*

longicauda dance flies (using COLONY – most likely sire configuration) Data are shown

relative to the proportion of offspring each male would sire if paternity shares were equal (the

null), indicated by the dashed line.

Literature Cited

Arnqvist, G. and T. Nilsson. 2000. The evolution of polyandry: multiple mating and female fitness in insects. Animal Behaviour 60, 145-164.

Berglund, A., G. Rosenqvist, and I. Svensson. 1988. Multiple matings and paternal brood care in the pipefish *Syngnathus typh*le. Oikos 51, 184-188.

Berglund, A. and G. Rosenqvist. 2003. Sex role reversal in pipefish. Advances in the Study of Behaviour 32, 131– 167.

Bonduriansky, R. 2001. The evolution of male mate choice in insects: a synthesis of ideas and evidence. Biological Reviews 76, 305-339.

Brown, W. D., and D. T. Gwynne. 1997. Evolution of mating in crickets, katydids, and wetas (Ensifera). Pages 279-312, *in* S. K. Gangwere, M. C. Mulalirangen and M. Muralirangen, eds. The Bionomics of Grasshoppers, Katydids and their Kin. CAB International, Oxford, UK.

Browne, J. H. 2021. Paternity and female ornaments in systems where females compete for nuptial gifts: A tale of long tails and short tails. PhD thesis. University of Toronto Mississauga, Ontario, CA.

Bussière, L. F., D. T. Gwynne, and R. Brooks. 2008. Contrasting sexual selectin on males and females in a role-reversed swarming dance fly, *Rhamphomyia longicauda*. Journal of Evolutionary Biology 21, 1683-1691.

Clutton-Brock, T. H. 1991. The evolution of parental care. Princeton, N.J: Princeton University Press.

Cumming, J. M. 1994. Sexual selection and the evolution of dance fly mating systems (Diptera: Empididae: Empidinae). The Canadian Entomologist 126, 907-920.

Delehanty, D. J., R. C. Fleischer, M. A. Colwell, and L. W. Oring. 1998. Sex-role reversal and the absence of extra-pair fertilization in Wilson's phalaropes. Animal Behaviour 55, 995-1002.

Downes, J. A. 1970. The feeding and mating behaviour of the specialized Empidinae (Diptera: Empididae: Empidinae). The Canadian Entomologist 126, 907–920.

Eberhard, W.G. 1985. Sexual selection and animal genitalia. Harvard University Press, Massachusetts, USA.

Flanagan, S. P., J. B. Johnson, E. Rose, and G. Jones. 2014. Sexual selection on female
ornaments in the sex-role-reversed pipefish (*Syngnathus scovelli*). Journal of
Evolutionary Biology 27, 2457-2467.

Fritzsche, K. and G. Arnqvist. 2013. Homage to Bateman: Sex roles predict sex differences in
sexual selection. Evolution 67,1926-1936.

Funk, D. H and D. W. Tallamy. 2000. Courtship role reversal and deceptive signals in the long-
tailed dance fly, *Rhamphomyia longicauda*. Animal Behaviour 59, 411-421.

Gerlach, N. M., J. W. McGlothlin, P. G. Parker, and E. D. Ketterson. 2012. Reinterpreting
Bateman gradients: multiple mating and selection in both sexes of a songbird species.
Behavioural Ecology 23, 1078-1088.

Gwynne, D. T. 1981. Sexual difference theory: Mormon Crickets show role reversal in mate
choice. Science 213, 779-780.

Gwynne, D. T. 1984a. Male mating effort, confidence of paternity, and insect sperm
competition, Pages 117-149. *in* R. L. Smith, ed. Sperm competition and the evolution of
animal mating systems, Academic Press Inc.

Gwynne, D. T. 1984b. Sexual selection and sexual differences in Mormon crickets (Orthoptera:
Tettigoniidae, *Anabrus simplex*). Evolution 38, 11011-1022.

Gwynne, D. T. 1985. Role-reversal in katydid: habitat influences reproductive behavior
(Orthoptera: Tettigoniidae, M*etaballus* sp.). Behavioural Ecology and Sociobiology 16,
355-361

Gwynne, D. T. 1988. Courtship feeding and the fitness of female katydids (Orthoptera:
Tettigonidae). Evolution 42, 545-555.

Gwynne, D. T. 1990. Testing parental investment and the control of sexual selection in
katydids: the operational sex ratio. American Naturalist 136, 474-484.

Gwynne, D. T. 1991. Sexual competition among females: What causes courtship-role
reversal? Trends in Ecology and Evolution 6, 118-121.

Gwynne, D. T. 2002. A secondary copulatory structure in a female insect: a clasp for a nuptial
meal? Naturwissenschaften 89, 125-127.

Gwynne, D. T. 2004. Reproductive behavior of ground weta (Orthoptera: Anostostomatidae):
Drumming behavior, nuptial feeding, post-copulatory guarding and maternal care.
Journal of the Kansas Entomological Society 77, 414-428.

Gwynne, D. T. 2005. The secondary copulatory organ in female ground weta (H*emiandrus*
pallitarsis, Orthoptera: Anostostomatidae): A sexually selected device in
females? Biological Journal of the Linnean Society 85, 463-469.

Gwynne, D. T. 2016. Sexual selection: roles evolving. Current Biology 26, 913-936.

Gwynne, D. T., and L. W. Simmons. 1990. Experimental reversal of courtship roles in an insect.
Nature 346, 172-174.

Haddrill, P. R., D. M. Shuker, W. Amos, M. E. N. Majerus, and S. Mayes. 2008. Female
multiple mating in wild and laboratory populations of the two-spot ladybird, *Adalia*
bipunctata. Molecular Ecology 17, 3189-3197.

Hare, R. M., and L. W. Simmons. 2018. Sexual selection and its evolutionary consequences in
female animals. Biological Reviews 94, 929-956.

Hartley, J. C. and A. C. Warne. 1972. The developmental biology of the egg stage of Western
European Tettigoniidae (Orthoptera). Journal of Zoology 168, 267-298.

Herridge, E. J. 2016. Polyandry rates and reproductive success in a nuptial gift giving dance fly
Rhamphomyia longicauda, PhD thesis. University of Stirling, Stirling, UK.

Herridge, E. J., R. L. Murray, D. T. Gwynne, and L. F. Bussière. 2016. Diversity in mating and
parental sex roles, Pages 453-458 *in* R. M. Kliman, ed. Encyclopedia of Evolutionary
Biology, Oxford Academic Press.

Hockham, L. R., J. A. Graves, and M. G. Ritchie. 2004. Sperm competition and the level of
polyandry in a bushcricket with large nuptial gifts. Behavioural Ecology and
Sociobiology 57, 149-154.

Hunter, F. D. L. and Bussière, L. F. 2019. Comparative evidence supports a role for
reproductive allocation in the evolution of female ornament diversity. Ecological
Entomology 44, 324-332.

Ingrisch, S. 1986. The plurennial life cycles of the European Tettigoniidae (Insecta: Orthoptera).
Oecologia 70, 606-616.

Jones, A. G. 2005. GERUD 2.0: A computer program for the reconstruction of parental
genotypes from half-sib progeny arrays with known or unknown parents. Molecular
Ecology Notes 5, 708-711.

Jones, A. G., J. C. Avise. 2001. Mating systems and sexual selection in male-pregnant pipefishes

and seahorses: Insights from microsatellite-based studies of maternity. Journal of

Heredity 92, 150-158.

Jones, A. G., D. Walker, and J. C. Avise. 2001. Genetic evidence for extreme polyandry and

extraordinary sex-role reversal in a pipefish. Proceedings of the Royal Society of London

B 268, 2531-2535.

Jones, O. R. and J. Wang. 2010. Colony: a program for parentage and sibship inference from

multilocus genotype data. Molecular Ecology Resources 10, 551-555.

Lewis, S. M., K. Vahed, J. M. Koene, L. Enggvist, L. Bussière, J. C. Perry, D. T. Gwynne, and

G. U. Lehmann. 2014. Emerging issues in the evolution of animal nuptial gifts. Biology

Letters 10, 20140336.

Lloyd, J. E. 1979. Mating behaviour and natural selection. The Florida Entomologist 62, 17-34.

Møller, A. P. and T. R. Birkhead. 1991. Frequent copulations and mate guarding as alternative

paternity guards in birds: a comparative study. Behaviour 118, 170-186.

Møller, A. P. and T. R. Birkhead. 1993. Certainty of paternity covaries with paternal care in

birds. Behavioral Ecology and Sociobiology 33, 261-268.

Murray, R. L., J. Wheeler, D. T. Gwynne, and L. F. Bussière. 2018. Sexual selection on multiple

female ornaments in dance flies. Proceedings of the Royal Society B 285, 20181525.

Owens, I. F. P., A. Dixon, T. Burke, and D. B. A. Thompson. 1995. Strategic paternity assurance

in the sex-role reversed Eurasian dotterel (*Charadrius morninellus*): Behavioral and

genetic evidence. Behavioral Ecology 6, 14-21.

Parker, G. A. 1970. Sperm competition and its evolutionary consequences in the insects.

Biological Reviews 45, 525-567.

Parker, G. A. 1990. Sperm competition games: Raffles and roles. Proceedings of the Royal

Society of London B 242, 120-126.

Parker, D. J., J. Zaborowska, M. G. Ritchie, and K. Vahed. 2017. Paternity analysis of wild-

caught females shows that sperm package size and placement influence fertilization

success in the bushcricket *Philidoptera griseoptera*. Molecular Ecology 26, 3050-3061.

Perry, J. C. and C. T. Tse. 2013. Extreme costs of mating for male two-spot ladybird beetles.

PLoS ONE 8, e81934.

Requena, G. S. and S. H. Alonzo. 2017. Sperm competition games when males invest in
paternal care. Proceedings of the Royal Society B 284, 20171266.

Ritchie, M. G., D. S. Sunter, and L. R. Hockham. 1998. Behavioral components of sex role
reversal in the tettigoniid bushcricket *Ephippiger ephippiger*. Journal of Insect Behavior,
11, 481-491.

Sakaluk, S. K. 1986. Sperm competition and the evlution of nuptial feeding behaviour in the
cricket *Gryllodes supplicans* (Walker). Evolution, 40, 584-593.

Schamel, D., D. M. Tracy, D. B. Lank, and D. F. Westneat. 2004. Mate guarding, copulation
strategies and paternity in the sex-role reversed, socially polyandrous red-necked
phalarope *Phalaropus lobatus*. Behavioral Ecology and Sociobiology 57, 110-118.

Simmons, L. W. 1992. Quantification of role reversal in relative parental investment in a bush
cricket. Nature 358, 61-63.

Simmons, L. W. 2001. Sperm competition and its evolutionary consequences in the insects.
Princeton University Press, New Jersey, USA.

Simmons, L. W. 2007. Polyandry in the wild: temporal changes in female mating frequency and
sperm competition intensity in natural populations of the tettigoniid *Requena verticalis*.
Molecular Ecology 16, 4613-4623.

Simmons, L. W. and W. J. Bailey. 1990. Resource influenced sex roles of Zaprochiline
Teettigoniids (Orthoptera: Tettigoniidae). Evolution 44, 1853-1868.

Simmons, L. W., and M. Beveridge. 2010. The strength of postcopulatory sexual selection
within natural populations of field crickets. Behavioural Ecology, 21:1179-1185.

Smith, R.L. 1979a. Repeated copulation and sperm precedence: paternity assurance for a male
brooding water bug. Science 205, 1029-1031.

Smith, R.L. 1979b. Paternity assurance and altered roles in the mating behaviour of a giant water
bug, *Abedus herberti* (Heteroptera: Belostomatidae). Animal Behaviour 27, 716-725

Starr, C. K. 1984. Sperm competition, kinship, and sociality in the aculeate Hymenoptera. Pages
427-464. *in* R. L. Smith, ed. Sperm Competition and the Evolution of Animal Mating
Systems. Academic Press, London, UK.

Thomas M. L. and L. W. Simmons. 2007. Male crickets adjust the viability of their sperm in
response to female mating status. American Naturalist 170, 190–195.

Thornhill, R. 1976. Sexual selection and paternal investment in insects. American Naturalist
110, 153-163.

Thornhill, R. 1979. Male and female sexual selection and the evolution of mating systems in
insects *in* M.S. Blum and N.A. Blum, eds. Sexual Selection and Reproductive
Competition in Insects. Academic Press, New York, NY

Tobias, J., R. Montgomerie, and B. E. Lyon. 2012. The evolution of female ornaments and
weaponry: social selection, sexual selection and ecological competition. Philosophical
transactions of The Royal Society B 367, 2274-2293.

Tripet, F., Y. T. Touré, C. E. Taylor, D. E. Norris, G. Dolo, and G. C. Lanzaro. 2001.
DNA analysis of transferred sperm reveals significant levels of gene flow between
molecular forms of *Anopheles gambiae*. Molecular Ecology 10, 1725–1732.

Trivers, R. L. 1972. Parental investment and sexual selection *in* B. Campbell, ed. Sexual
selection and the decent of man. Aldine, Chicago, USA

Turnell, B. R. and K. L. Shaw. 2015. Polyandry and postcopulatory sexual selection in a wild
population. Molecular Ecology 24, 6278-6288.

Vahed, K. 1998. The function of nuptial feeding in insects: a review of empirical studies.
Biological Reviews 73, 43-78.

Vahed, K. and F. S. Gilbert. 1996. Differences across taxa in nuptial gift size correlate with
differences in sperm number and ejaculate volume in bushcrickets (Orthoptera:
Tettigoniidae). Proceedings of the Royal Society B 263, 1257-1265.

Westneat, D. R. and P. W. Sherman. 1992. Parentage and the evolution of parental behaviour.
Behavioural Ecology 4, 66-77.

Wheeler, J. 2008. Sexual selection and female ornamentation in a role-reversed dance fly.
M.Sc. thesis. University of Toronto, Toronto, Canada.

Wheeler, J., D. T. Gwynne, and L. F. Bussière. 2012. Stabilizing sexual selection for female
ornaments in a dance fly. Journal of Evolutionary Biology 25, 1233-1242.

Yoshizawa, K., R. L. Ferreira, Y. Kamimura, and C. Lienhard. 2014. Female penis, male vagina,
and their correlated evolution in a cave insect. Current Biology 24, 1006-1010.

Yoshizawa, K., R. L. Ferreira, I. Yao, C. Lienhard, and Y. Kamimura. 2018. Independent origins

of female penis and its coevolution with male vagina in cave insects (*Psocodea:*

Prionoglarididae). Biology Letters 14, 20180533.

Acknowledgements:

The New Zealand Department of Conservation kindly gave permissions to collect specimens.

Genetic Marker Services developed of our microsatellite library and The Centre for Applied

Genomics conducted genetic analyses. Thanks to: the Marc Johnson lab for use of equipment

and other assistance, Lindsay Coome and Samreen Munim for their molecular work, Mary

Morgan-Richards and Steve Trewick for advice, hospitality, and allowing us to collect, as well as

Luc Bussière, Rosalind Murray, Marc Johnson, and Doug Currie) for comments on the research

(funded by a NSERC Discovery Grant to DTG)