Sexual selection on a female copulatory device in an insect with nuptial gifts

Jessica H. Browne[1,2]

Darryl T. Gwynne[1,3]

[1] Department of Ecology and Evolutionary Biology, University of Toronto Mississauga, Mississauga, Ontario L5L

1C6, Canada

[2] E-mail: jbrowne@mta.ca, Current address: Department of Biology

Mount Allison University, 62 York St. Sackville, NB, E4L 1E2, Canada

[3] E-mail: darryl.gwynne@utornto.ca

OCRID: 0000-0002-1849-5182

Abstract

Male genitalia are rapidly evolving structures, often driven sexual selection to increase fertilization success.
Although sexual selection on females can be strong in systems where males provide offspring care or feed their
mates, sometimes resulting in the evolution of female ornamentation, there are no actual estimates of direct sexual
selection on female genitalia. In a New Zealand ground weta, *Hemiandrus pallitarsis* (Orthoptera: Ensifera,
Anostostomatidae), females possess a genitalic device (the accessory organ) that is necessary for successful
copulation and the acquisition of glandular food-gifts from males. These nutritious gifts are known to result in
sexual competition among females in other ensiferan species. In ground weta, the gifts are probably important in
avoiding starvation during a months-long period when caring for (their lifetime production of) eggs and offspring.
Here, we test the hypothesis that the accessory organ is a sexually selected device in *H. pallitarsis* by measuring the
female Bateman gradient and directional sexual selection on the accessory organ. Using newly developed and
characterized microsatellite loci, we analyze offspring and/or stored sperm to estimate female mating frequency for
the first time in ground weta. As predicted, we found positive Bateman gradients for females, and some evidence of
directional sexual selection on accessory organ length. Although organ length does not correlate well with female
fecundity, it may increase mating success by indicating her condition and thus quality of her offspring care.

Keywords: female ornamentation, genital evolution, sexual selection, paternal investments, nuptial gifts,

microsatellite analysis

Statements and Declarations

The authors have no competing interests to declare

Acknowledgements and Funding Statement

We thank Mary Morgan Richards and Lindsay Coome for their help collecting data, Doug Currie, Marc Johnson,

Rosalind Murray, and Clint Kelly for comments on the research and manuscript, as well as the New Zealand

Department of Conservation (for permits). This research was supported by a Natural Sciences and Engineering

Research Council of Canada Discovery Grant to DTG.

Introduction

Genitalia and associated structures (secondary genitalia) are some of the most rapidly evolving animal

traits, often exhibiting extreme variation among closely related species (Eberhard 1985). While genital diversity has

been suggested to function in species isolation ("lock and key" hypothesis; Mayr 1963; Eberhard 1985), genital

evolution in males is thought to be driven mainly by sexual selection for increased fertilization success (Eberhard

1985). Because females are typically more selective when mating (Darwin 1871; Trivers 1972), elaborate male

genitalia may have evolved in the context of cryptic female choice, where males with the most "attractive"

stimulatory genitalia experience the greatest fertilization success (Lloyd 1979; Thornhill 1983; Eberhard 1985;

Simmons 2001). Alternatively, genitalia may function in manipulating female mating behaviour and/or her sperm

stores (e.g., grasping devices that coerce copulations or prevent female remating; Arnqvist and Rowe 2005).

Although much of the focus has been on male traits, females are not passive in this process (Sloan and Simmons

2019). In response to, for example, male grasping or piercing genitalia, there can be co-evolution of female genitalia

to reduce potential costs of mating (e.g., Arnqvist and Rowe 2005; Morrow and Arnqvist 2003) or to bias

fertilization success in favour of high-quality males (Eberhard 1985; Sloan and Simmons 2019).

In contrast to these examples of "typical" mating roles, the factors influencing male and female fitness are

expected to differ when males make valuable investments in paternal care or nuptial gifts. Females receiving these

direct benefits are expected to seek additional matings, while the potential for frequent mating by males can be

limited by investment cost, resulting in male preferences and/or direct sexual competition among females (Trivers

1972; Williams 1975; Parker and Simmons 1996; Bonduriansky 2001; Gwynne 2016; Herridge et al. 2016). In these

systems, especially those where females depend on male investments for offspring production, female competition

can cause strong sexual selection on females for traits that are attractive to males (Tobias et al. 2012; Herridge et al.

2016; Hare and Simmons 2018). Such traits are well-known both in certain vertebrates with paternal care (e.g., sex-

specific female ornaments in pipefish: Flanagan et al. 2014) and in insects such as empid dance flies (legs, wing, or

abdominal ornaments; Cumming 1994; Murray et al 2022) where females rely on male-provided nutrition (insect

prey items) to develop their eggs (Downes 1970). Also, in katydids (Orthoptera: Ensifera; Tettigoniidae) where

males supply large spermatophore gifts (Vahed and Gilbert 1996), there is sexual selection on females when they

rely on these gifts for nutrition, which impose additional mating costs for males (Gwynne 1981; 1984; Simmons and

Bailey 1990; Gwynne and Simmons 1990; Simmons 1992; Robson and Gwynne 2010). In *Kawanaphila nartee,*

females with larger ears are faster in the race to reach singing males (Gwynne and Bailey 1999) and there is positive

selection for ear size, a sexually dimorphic trait in this species, only when food is limited (Hare and Simmons 2021).

Virtually unstudied is direct sexual selection on sex-specific female genitalic traits where copulation is

associated with acquiring goods or services from males. As in males, female genital traits may be expected to evolve

either in the context of mate choice – such as stimulating males into copulating – or coercing males into supplying

more food. Suggested examples are (1) the interspecifically varying ridges on female subgenital plates of

Kawanaphila katydids (Rentz 1993) that grasp the male's genitalia, perhaps allowing longer copulation duration and

thus the delivery of a larger spermatophore gift (Gwynne 2001); and (2) a penis-like female genital modification in

two genera of psocopteran cave insects, *Neotrogla* and *Afrotrogla*. Male psocopterans can transfer nutritious semen

to females during copulation, which is thought to be an important source of nutrients (Wearing-Wilde 1996;

Yoshizawa et al. 2018). The penis-like gynosome, inserted into the male's genital opening, anchors him during

copulation (Yoshizawa et al. 2014), and is a device likely under sexual selection to allow females to prolong the

duration of copulation and thus the amount of food-gift received (Yoshizawa et al. 2014; Yoshizawa et al. 2018).

Our study species has female "secondary" genitalia, abdominal devices found in a clade of New Zealand

"short-tailed" ground weta (*Hemiandrus*, Anostostomatidae, Orthoptera: Trewick et al. 2020). As in *Neotroglea*

(Yoshizawa et al. 2014), male ground weta transfer nutrients to their mates, in this case, a glandular spermatophylax

part of the spermatophore (as in katydids; Gwynne 1995) that the mating male adheres to the female's abdomen

(separate from the sperm capsule) and is eaten by her after copulation (Gwynne 2004; Browne and Gwynne 2022).

The female device is an elongated abdominal modification on the 6^{th} sternite, known as an accessory organ (Salmon

1950). In contrast to other species of *Hemiandrus* that, like most ensiferan species, abandon their eggs after

ovipositing into the substrate (Gwynne 2004; Taylor Smith et al. 2013), short-tailed females care for their eggs and

newly-hatched offspring (their lifetime reproduction) in an underground chamber for a period of 5-6 months (hence

the "short-tail" reduced ovipositor; Gwynne 2004; Gwynne 2005; Taylor Smith 2015) and then die. Care-providing

females do not eat, so male mating gifts likely represent important nutrition to avoid starvation during the lengthy

care period, potentially leading females to compete for mating (feeding) opportunities (Gwynne 2004; Gwynne

2005). Because of this, the accessory organ, positioned close to where the sticky spermatophylax gift is adhered, is

expected to be involved in the acquisition of gifts.

Male genitalia in many weta (Anostostomatidae) and some related ensiferans, include genital hooks (falci)

on the dorsal side of the 10^{th} segment (Johns 2001) which hook onto the edge of one of the of the female's sternites

during copulation (the male is positioned beneath the female: Gwynne 2002). This is likely an ancestral behaviour,

used to grip the female for stability or to coerce her into copulation (e.g., Weissman 2001). While some weta species

have sclerotised cuticle on the sternite edge that receives the male hooks, female short-tailed ground weta are unique

in possessing the accessory-organ modifications on the 6^{th} sternite, to which the male genitalia attach, but do not

hook (D. Gwynne, personal observation). Females of our study species, *H. pallitarsis,* possess the most elaborate

accessory organ (fig. 1), which protrudes from the abdomen. After pairing (female mounted above male), males first

make genitalic contact with the accessory organ then move their genitalia back to attach to the female's primary

genitalia when they insert the sperm ampulla (Gwynne 2004; Gwynne 2005). Following this, the male detaches from

her primary genitalia, but his genitalia remain attached to her abdominal accessory organ while he extends paired

phalli that deposit the bilobed spermatophylax gift (Gwynne 2002; Gwynne 2004; Gwynne 2005). During gift

transfer, the male's paired falci press onto the dorsal surface of the accessory organ, while his paired epiprocts

appear to be inserted ventral to the organ, and the distal edge of the male's 8^{th} abdominal segment pushes into the

region between the female's 5^{th} and 6^{th} sternites (photographic evidence: DTG unpublished). The placement of the

spermatophylax gift anterior to the accessory organ suggests that it is not a holding device (see Gwynne 2002) for

the gift (Gwynne 2005). However, it is necessary for mating (Gwynne 2005) as copulation fails when the female

112 organ is experimentally removed (Gwynne 2005). Thus, this device has not evolved to thwart copulation attempts

113 (as in some gerrid water striders: Arnqvist and Rowe 2005).

116 **Figure 1.** Lateral view of the extended female secondary genitalic accessory organ (6[th] sternite, located between the

117 hind and middle legs) in A) a newly-mated female *H. pallitarsis* and B) a close-up of the organ. Photos by Darryl

118 Gwynne.

120  Although the female accessory organ is involved in transfer of the ejaculate (sperm and the gift in ground

121 weta), there is some evidence that it is sexually selected to secure matings as envisioned by Eberhard (1990) for

122 male genitalia (examples: fish, Brooks and Caithness 1995; lizards, Bohme 1983; flies,   . 2021; Briceño

123 et al. 2010; and katydids, Wulff and Lehmann 2016). First, extensive interspecific variation in the accessory organs

124 of short-tailed ground weta (Johns 2001; Trewick et al. 2020) reflects the typical pattern in male animals (Gwynne

125 2005) where external genitalia vary greatly across closely related species. Variation in female accessory organs is

126 probably not related to species isolation ("lock and key" hypothesis; Mayr 1963; Eberhard 1985) since there is

greater support for sexual selection on complex genitalia in male insects (Eberhard 1985; Gwynne 2005) and, for

ground weta, mating interactions between species are more likely prevented by the species-specific precopulatory

drumming behaviour of males and females that precede pairing (Gwynne 2004; Gwynne 2005; Chappell et al.

2012). Furthermore, there is little overlap in ground weta species ranges (Trewick and Bland 2011; Trewick et al.

2020), which makes species mis-pairing unlikely. Also consistent with the sexual selection hypothesis, male ground

weta make close contact with the accessory organ prior to mating and have been observed to reject females after

probing her mid-abdomen (Gwynne 2005). As females with longer accessory organs tend to have a greater number

of eggs that develop to the nymphal stage (female body size controlled; Gwynne 2005), this trait may help females

obtain matings (gifts) by signalling quality to males (as in Andersson 1994; Berglund et al. 1997; Barry 2015). Thus,

one hypothesis is that this structure evolved via male choice in the context of female mate/gift acquisition (Gwynne

2005),

As the female organ is inserted between two parts of the male genitalia, another possibility is that it has

evolved by sexual selection on females to anchor between the male genital falci and epiprocts, and thus manipulate

mating or spermatophylax-gift transfer, in a similar manner to the anchoring female penis of *Neotroglea* bark lice

suggested by Yoshizawa et al. (2014) as a sexually-selected organ to obtain nutritious ejaculates. Gwynne's (2005)

finding that accessory organ length correlates with the number of eggs that develop to the nymphal stage in *H.*

pallitarsis is also consistent with this hypothesis if accessory organs result in a longer copulation and more nuptial

food. However, there is no direct evidence that any female copulation devices such as accessory organs or female

penes are ornaments subject to sexual selection. Here, we test this by measuring sexual selection on female ground

weta *H. pallitarsis*. Using estimates of female mating success from microsatellite analysis of stored sperm and

measures of lifetime female reproductive success from females that cared for their broods in the lab, we determine

the degree of sexual selection on females (Bateman gradient) and estimate directional sexual selection on several

accessory organ traits. If the accessory organ is sexually-selected in the context of gift acquisition, we predict

directional selection on accessory organ size (length; Gwynne 2005) and that this trait will correlate with female

fecundity. Characteristic of sexual selection on a sex (Bateman 1948; Arnold 1994), we also expect there to be a

positive relationship between female mating success and reproductive success (Bateman gradient).

Methods

Specimen collection: Bateman gradients

To determine Bateman gradients for female ground weta *(H. pallitarsis)*, females were collected in 2002 (n=10)

and 2017 (n=17) from two main sites on the North Island of New Zealand: Kiriwhakapapa trail near Masterton (-

40.807627, 175.546532) and private gardens in Palmerston North (-40.413909, 175.662814). Collections took place

from mid-February to early March, nearing the end of the ground weta mating season (Gwynne 2004). Live

specimens were transported back to our lab in Toronto, Canada to oviposit and obtain measures of lifetime

reproductive success from caring females. This was a lengthy process that involved placing females in artificial

brood chambers formed from potter's clay (2002) or soil (2017) and allowing 2-3 months for them to lay eggs, as

well as an additional 5-6 months for eggs to develop, during which time females were not fed. Several females died

during this period, four of which were excluded from further analysis because both her and her eggs decayed rapidly

(final n=23). Once eggs began hatching, mothers and their broods were preserved at -20°C to prevent consumption

of dead or developing tissue. We counted the total number of eggs laid as well as the number of hatched (1st instar

nymphs) or nearly hatched (eggs with visible eye spots) offspring, which we considered "developed eggs". To

obtain the sperm from mates that did not fertilize offspring, we dissected female sperm storage organs, isolated the

sperm pellet (using >70% ethanol; Tripet et al. 2001), and stored at -20°C. We then photographed females under a

dissecting scope fitted with a digital camera and measured pronotum length as an estimate of body size (as in Taylor

Smith 2015).

Specimen collection: Sexual selection on accessory organs

To measure sexual selection on accessory organs, additional *H. pallitarsis* females (n=58) were collected in

2012, 2016, 2017, and 2019 near the end of the mating period from the same two sites: Kiriwhakapapa and

Palmerston North. Because our primary interest here was not reproductive output, these females were frozen

immediately following capture and stored in strong ethanol (70-95%). This method allowed us to avoid a time-

consuming and risky offspring-rearing period and ensured that both female phenotype and the quality of sperm

stores were not compromised. For this same reason, we did not include the 23 females that laid eggs (used for

Bateman gradient estimates) in our measures of selection on accessory organs. We photographed preserved females

under a dissecting scope and measured pronotum length (body size; Taylor Smith 2015) as well as accessory organ

length and width (fig. 2) using ImageJ software (Schneider et al. 2012). Prior to measurement, all accessory organs

were removed and soaked in a 5% Decon 90 solution for 1 hour at 40ºC to soften and restore the structures (Banks

and Williams 1972), as many were bent and brittle after prolonged storage in ethanol. We then dissected the

spermathecae from all females, isolated the sperm contents using strong ethanol (Tripet et al. 2001), and stored them

at -20°C. During the dissection, we also counted the number of developing ova in the female's abdomen.

Figure 2. Modified 6[th] sternite, the accessory organ, on the abdomen of female *H. pallitarsis* ground weta. Image

shows the structure removed from the body and the two measurements taken: A) accessory organ length, from the

point where the organ attaches to the body to the furthest point on the distal end and B) accessory organ width, left

to right on the widest point of the distal end.

Microsatellite analysis

Prior to conducting genetic analysis, we assessed amplification quality as well the degree of polymorphism

for the nine candidate loci in our microsatellite library (developed by Genetic Marker Services, Brighton, UK for *H.*

pallitarsis; provided in Online Resource 1: Table S1) We used polymerase chain reaction (PCR) to amplify DNA

samples extracted from adult hind leg tissue, including the females in the current study plus an additional 24 males.

Using GENEPOP version 4.2 (Raymond and Rousset 1995), we determined that the degree of genetic differentiation

(F_{st}) was low between our two study sites (populations) and identified three polymorphic loci (>20 alleles) with a

low frequency (< 20%; Dakin and Avise 2004) of null alleles (Dempsters EM method; heterozygote deficiency test)

and non-specific amplification (wet80, wet 81, and wet89; details in Online Resource 1: Table S2). Targeting these

loci, we PCR-amplified samples of stored sperm and offspring. We extracted DNA from stored sperm using the

Qiagen DNeasy Blood and Tissue Extraction kit (Qiagen, Hilden, Germany), and added 12μL DTT to each sample

to improve lysis of sperm cells. We extracted DNA from offspring tissue (nymph or egg) using a Proteinase K based

extraction method with multiple ethanol washes. The resulting concentration of DNA was much higher for offspring

relative to sperm and so we diluted offspring samples to roughly 50ng/μL. Each PCR reaction contained 5.75μL

sterile filtered water, 1μL 10X PCR reaction buffer, 0.2μL 10mM dNTP mixture, 0.5μL of forward primer (each

with fluorescent labels), 0.5μL reverse primer, and 1μL of template DNA. Thermocycling was done using an

Eppendorf Mastercycler with the following temperature regime: 2 minutes at 94°C followed by 35 cycles of 30

seconds at 94°C, 30 seconds at 60°C, and 15 seconds at 72°C, then an additional 3 minutes at 72°C. PCR products

were sent to The Centre for Applied Genomics at Sick Kids Hospital in Toronto, ON for fragment analysis on an

Applied Biosystems 3730xl capillary sequencer. The resulting electropherograms were examined using GeneMarker

V1.97, which allowed us to identify fluorescence peaks, representing alleles.

Estimating female mating success

For measures of the Bateman gradient, we used a combination of paternity data and the remaining stored

sperm to estimate female mating frequency. Based on offspring genotypes, we first determined the minimum

number of sires for each brood using the parentage program GERUD 2.0 (Jones 2005). Any alleles found in

female's sperm stores that were not present in the offspring genotypes were assumed to be additional mates and

were factored into a female's total mating frequency. We estimated the most probable number of additional mates at

each locus using a method described by Bretman and Tregenza (2005). This method (See Bretman and Tregenza

2005; Herridge 2016 thesis for details) makes use of known allele frequencies in the population to determine the

probability of observing a particular set of alleles given different numbers of contributing males. This provides a less

conservative estimate than simple allele counting, which assumes heterozygosity (two alleles indicate one

individual). Because estimates of females mating success incorporated offspring data, we could not remain blind to

reproductive success (the response) when collecting these data. One of our specimens was known to have mated, as

she was found with remnants of a spermatophylax gift in her mouthparts; however, the male (who remains close by

while she consumes the gift) failed to inseminate this female as his genotype did not appear in her sperm stores, nor

her offspring. Because acquisition of male-provided materials (in this case the spermatophylax gift) is the key factor

affecting female fitness (Clutton-Brock 2007), an additional mate was added to this female's total mate number for

our estimates of the Bateman Gradient.

To measure directional sexual selection on accessory organs, we estimated female mating frequency from

stored sperm alone. Using all male alleles detected in stored sperm, we determined the most probable mate number

(see Bretman and Tregenza 2005) obtained by each female. Unfortunately, due to poor amplification at all loci, we

were unable to estimate the mate number for four females. In cases where an allele was found in sperm stores that

was not previously detected in the population, it was assumed to be rare and assigned a frequency equal to the least

common allele known in that population. We were blind to estimates of female mating frequency as they were made

without knowledge of phenotype.

Statistical analysis: Bateman gradients

We measured the Bateman gradient for females using our estimates of mating frequency from stored sperm

and the number of offspring in single broods that each female cared for before dying (lifetime reproductive success).

We calculated standardized Bateman gradients (β'_{ss}; Bateman 1948; Arnold 1994) using an ordinary least squares

(OLS) regression in which relative number of mates (mean-standardized) was the predictor variable and relative

reproductive success (mean-standardized) was the response variable. A separate regression model was run for each

of our measures of reproductive success: the number of eggs laid and the number of eggs that developed to the

nymphal stage (determined by hatching or visible eye spots). We measured pronotum length as an indicator of body

size (e.g., Taylor Smith 2015), however decomposition of several females in their brooding chambers prevented us

from obtaining data for all specimens. Using the available pronotum data (n=20), we included body size in our

regression model to determine whether this explained significant variation in reproductive success or altered

measures of the Bateman gradients Due to our low sample size, and deviations from normality (assessed visually

with Q-Q plots), we used permutation tests to obtain p-values. We ran 5000 permutations to generate a null

distribution of possible Bateman gradients (regression coefficients) from our data and compared this to our observed

values to determine if they were significantly different from those expected by chance. Due to apparent variation

between our collection sites and years, we assessed whether measures of the Bateman gradient interacted with either

of these variables. Using separate regression models for each variable, we included site and year as interaction terms

and used p values derived from multiple least squares regression to assess significance. We did not use permutation

tests to obtain p values for interaction terms, as this approach does not preserve important associations in the data,

leading to inflated error rates (Bužková et al. 2011; Bužková 2016; Foster et al. 2016).

Statistical analysis: Sexual selection on female accessory organs

We measured sexual selection acting directly on female accessory organs using female phenotype (body

size, accessory organ length, accessory organ width) and estimates of mating frequency from stored sperm. We

obtained linear selection gradients (indicative of directional selection) using multiple least squares regression (MLS)

in which trait values (z-score standardized) were predictor variables and relative (mean-standardized) number of

mates was the fitness response variable. Low sample size limited the complexity of our regression model; thus, we

were unable to include quadratic and correlational terms (Lande and Arnold 1983) in our analysis. To determine

whether observed selection gradients were statistically significant, we again used permutation tests with 5000

randomizations (as in Kelly & Gwynne 2023). Due to apparent variation in the effect of female body size as well as

accessory organ width between our two collection sites, we tested for interactions using MLS regression. We ran

individual models for each female trait, in which the interaction with site was included as a predictor, and used this

to derive both effect sizes and p-values for interaction terms (see Bužková et al. 2011; Bužková 2016; Foster et al.

2016 for limitations of permutation tests)

Finally, in order to assess whether the accessory organ has the potential to signal female fecundity, we ran

another MLS regression model with z-score standardized trait values (body size, accessory organ length, accessory

organ width), but used relative (mean-standardized) number of developing eggs in the abdomen (not yet laid) as the

response variable. Again, we used permutation tests with 5000 randomizations to obtain p-values and tested for

interactions between site and either body size or accessory organ width (due to visual variation) using MLS

regression.

Results

Bateman gradients

Consistent with our prediction that female ground weta are subject to sexual selection, we found evidence of

significant positive Bateman gradients in one of our two collection years (table 1). Although Bateman gradients

using pooled data were found to be positive, they were not statistically significant (table 1) and showed considerable

variation across our two collection sites and years, especially when using the total eggs laid (fig. 3). While there

were no significant interactions when using the number of developed eggs, there was a significant interaction with

year when using eggs laid (table 1), leading us to analyze data separately for each year. The Bateman gradient was

high in the year 2017 and appears to drive the overall positive trend found between mating success and the number

of eggs laid (table 1; fig. 3).

Estimates of β'_{ss} were similar regardless of whether we included the available pronotum data in our

regression model ($\beta'_{ss} = 0.155 \pm 0.264$, F(2, 16) = 1.46, p = 0.280 for eggs laid; $\beta'_{ss} = 0.433 \pm 0.310$, F(2, 17) =

1.16, p = 0.090 for eggs developed) and in line with Gwynne (2005), female body size had no significant effect on

the number of developed eggs ($\beta' = 2.343 \pm 2.982$, $R^2 = 0.120$, F(2, 17) = 1.16, p = 0.214), but a marginally

significant effect on the number of eggs laid ($\beta' = 4.07 \pm 2.490$, $R^2 = 0.154$, F(2, 16) = 1.46, p = 0.043).

Table 1 Standardized Bateman Gradients (β'_{ss}) for wild-caught female ground weta, *H. pallitarsi*s

Response Variable	Predictors	$\beta'_{ss} \pm SE$	R^2	F	df	p-value
Eggs laid	**Mates**	**0.117 ± 0.231**	**0.013**	**0.2548**	**1, 20**	**0.313**
	Mates:site	-0.545 ± 0.413	0.310,	2.69	3,18	0.203[a]
	Mates:year	0.950 ± 0.393	0.420	4.34	3,18	0.0267[a] *
	Mates (2002 samples)	-0.121 ± 0.185	0.067	0.432	1, 6	0.249
	Mates (2017 samples)	1.032 ± 0.358	0.4096	8.326	1, 12	0.0084*
Developed Eggs	**Mates**	**0.392 ± 0.272**	**0.090**	**2.071**	**1, 21**	**0.084 .**
	Mates:site	-0.123 ± 0.576	0.126	2.69	3,19	0.833[a]
	Mates:year	0.570 ± 0.525	0.335	3.189	3,19	0.291[a]

β'_{ss} = Standardized Bateman gradient (variables mean-standardized)

SE = Standard error

Estimates obtained from separate OLS regression analyses; p-values from Permutation tests with 5000

randomizations

[a] p values obtained from separate multiple OLS regressions

* Statistically significant p-value

308
Figure 3. Raw Bateman Gradients for female *H. pallitarsis* showing relationship between number of mates and

either A) number of eggs laid or B) number of developed eggs with 95% confidence intervals. Variation between

collection sites and years are shown with dashed lines (associated values provided in Online Resource 1: Table S3).

Sexual selection on accessory organs

Female mating success ranged from 0-6 mates (mean: 2.3 +/- 1.4 SD) and increased significantly with

accessory organ length (table 2). Larger body size showed a non-significant trend with increased mating success

(table 2). Selection did not vary significantly between collection sites (MLS interaction terms; body size: β'=0.319 ±

0.247, $F(3, 50)$ = 1.39, p=0.202; accessory organ width: β'= -0.331 ± 0.276, $F(3,50)$ = 0.537, p=0.236), but we note

that (non-significant) trends tended to vary between our two populations of ground weta (visual provided in Online

Resource 1: Fig. S4).

None of the accessory organ traits measured significantly predicted the number of eggs in a female's

abdomen. Although we found a statistically significant effect of accessory organ width (table 3), this trend appears

to be driven almost entirely by differences over time (visual provided in Online Resource 1: Fig. S5). Indeed, when

collection year is added as a random effect in a general linear mixed model, the significant effect of accessory organ

width disappears ($\beta'=0.127 \pm 0.138$, $p=0.238$). The relationship between female traits and fecundity did not vary

significantly between collection sites (OLS interaction terms; body size: $\beta'=0.191 \pm 0.342$, $F(3, 53) = 1.172$,

$p=0.579$; accessory organ width: $\beta'= -0.331 \pm 0.359$, $F(3,53) = 2.09$, $p=0.361$, but again (non-significant) trends

varied between sites (visual provided in Online Resource: Fig. S6).

Table 2 Relationship between phenotype (traits z-score standardized) and mating success (relative number of mates)

in female ground weta, *H. pallitarsis*.

Female traits	β (SE)	p-value
Body size	0.106 (0.093)	0.2094
Accessory organ length	0.195 (0.097)	0.0426 *
Accessory organ width	-0.023 (0.095)	0.7776

331 $R^2 = 0.105$, $F (3, 50) = 1.96$
332 $\beta=$ standardized directional selection gradients; regression coefficients
333 SE = Standard error
334 Estimates obtained from multiple OLS regression; p-values from Permutation tests with 5000 randomizations
335 * Statistically significant p-value
336

337 **Table 3** Relationship between female phenotype (traits z-score standardized) and relative number of eggs in the

338 abdomen in female ground weta, *H. pallitarsis*.

Female traits	β (SE)	p-value
Body size	0.001 (0.126)	1.00
Accessory organ length	-0.043 (0.118)	0.7026
Accessory organ width	0.248 (0.128)	0.0306 *

339 $R^2 = 0.089$, $F(3, 53) = 1.72$
340 β = standardized regression coefficients
341 SE = Standard error
342 Estimates obtained from multiple OLS regression; p-values from Permutation tests with 5000 randomizations

343　* Statistically significant p-value
344

Discussion

In *H. pallitarsis* ground weta, sexual competition for male-provided nutrition (spermatophylaxes) prior to

an extended maternal care period without food probably creates an opportunity for sexual selection on females when

they "forage" for mating gifts. Morphological evidence that females are subject to sexual selection comes from the

existence of an apparent female ornament, a secondary-genitalic device known as an accessory organ (Gwynne

2002; Gwynne 2005) that varies greatly between species (Johns 2001; Trewick et al. 2020). Here, we provide some

evidence that *H. pallitarsis* females are subject to sexual selection, by showing 1) a significant positive relationship

between female mating success and lifetime reproductive success (Standardized Bateman gradient; β'_{ss}) and 2) a

significant relationship between the length of the female accessory organ and mating success.

Although our estimates of the Bateman gradient were not consistent across collection years, they reached

values in 2017 that are similar to other species where sexually selected females compete for mates, such as bean

beetles (β'_{ss} =0.62-0.87; Fritzsche & Arnqvist 2013) or pipefish (β'_{ss} = 0.86-1.04; Mobley & Jones 2012; Aronsen et

al. 2013). While pooled estimates of the Bateman gradient were not statistically significant, they were higher than

the estimates found in female *K. nartee* (β'_{ss} = 0.029-0.034; Hare and Simmons 2022) as well as the relationship

between number of sires and absolute reproductive success found for females of another anostostomatid weta,

Hemideina crassidens, where males are sexually competitive (they fight using very large mandibular weaponry) (β=

0.47; Nason and Kelly 2020).

Further evidence of sexual selection on female ground weta comes from our finding that accessory organs

appear to be subject to directional selection. There was a marginally significant positive relationship between the

length of the accessory organ and female mating success, supporting the hypothesis that this this structure functions

as a device that ensures successful mating (there was also a non-significant trend for larger females to mate more

frequently). In all our measures of sexual selection, there was considerable variation between study sites. This is

interesting given the lack of genetic differentiation between these populations (F_{st} low compared to typical values in

insects; Ward et al 1992) and their close proximity (roughly 100km) relative to the total species range.

Morphological differences have been reported between populations of *H. pallitarsis*, with Chappell et al. (2012)

finding that some populations differed in body size as well as the drumming pattern performed by males prior to

mating. While this variation was not congruent with genetic differences among populations (suggesting that widely

separated *H. pallitarsis* populations comprise one interbreeding species; Chappell et al. 2012), patterns of selection

and mate attraction may differ slightly between populations of ground weta in Palmerston North compared to

Kiriwhakapapa, possibly because the habitats differ; overgrown gardens (with apparently high population density)

in the former site and wild vegetation in the latter. Regardless of this variation, female ground weta appear to be

subject to sexual selection, and the length of accessory organ plays a consistent role in acquiring mates in both

populations.

Despite this finding, neither accessory organ length, nor any of the female traits measured were related to

the number of eggs in a female's abdomen (fecundity). However, this does not mean the accessory organ does not

signal attractiveness to males. One reason for the non-significant finding may be that the number of eggs in the

abdomen is not the main indicator of female quality for the mating male. Because females oviposit all their eggs and

then engage in a long maternal care period, during which the mother does not eat (Gwynne 2004; Browne and

Gwynne 2022), choosy males may prioritise female condition and thus the potential quality of care she is able to

provide, as this will determine the number of surviving offspring. This is supported by Gwynne's (2005) study

which showed a positive relationship between accessory organ length and number of hatched nymphs rather than

eggs in the abdomen and might also help to explain why our measures of the Bateman gradients tended to be more

consistent (and possibly stronger) when using the number of developed eggs to estimate reproductive success rather

than just the number of eggs laid (fig. 3; Supplementary information 3). Similar results might also be expected

however, if there is variation in quality or size of the spermatophylax gift received, and thus we cannot rule out the

possibility that females with longer accessory organs also obtain more nutrients for successful offspring production,

due, for example, to male preferences (reviewed in Simmons 2001) or a manipulative anchoring function similar to

Neotroglea (Yoshizawa et al. 2014; Yoshizawa et al. 2018).

As noted previously, the accessory organ's role in competing successfully for copulations does not exclude

its mechanical function in successful ejaculate transfer (male genitalia facilitate sperm transfer as well as potentially

stimulating mates; Eberhard 1990). Given the presumed ancestral state where males used genitalic hooks to latch

onto a female's unmodified 6[th] sternite, the accessory organ could have initially evolved in the context of mate

assessment (in the region that interacts closely with sensitive male genitalia) or as an grasping device between the

male genital falci and epiprocts, that facilitates spermatophylax-gift transfer. The association between paired male

genitalic structures (falci, paraprocts, and the end of the 8[th] sternite; D. Gwynne personal observation) almost

certainly achieve stability while first the sperm capsule (ampulla) and then the spermatophyax gift is delivered.

However, it is possible, that like many male structures (Eberhard 1990), different parts of the accessory organ serve

different functions. We note, for example, that the distal end of the accessory organ has a very prominent bilobed

shape that appears to align with the lobes of the males' paired paraprocts (in several *Hemiandrus* species: Gwynne

2002). Because our measures of the accessory organ were simple linear ones, we have not captured the full

complexity of this structure that may be important to successful mating. Future studies should focus on quantifying

the shape of this structure and its role in mating.

Another limitation of the study is that we were only able to measure directional selection on accessory

organs. Due to our limited sample size, we did not compute estimates of non-linear or correlational selection,

meaning we may have an incomplete understanding of the type of selection acting on these structures (Lande and

Arnold 1983). In particular, it would be valuable to test for stabilizing sexual selection on female accessory organs

as preference for intermediately ornamented females has been predicted in some species (typically those with more

elaborate and costly traits) due to the potential costs (e.g., in egg production) of investing in these traits (Fitzpatrick

1995; Chenoweth et al. 2006; Wheeler et al. 2012). Furthermore, genital structures may be expected to be under

stabilizing sexual selection in cases where they have evolved to fit the average male (Eberhard et al. 1998).

In conclusion, we show some evidence of sexual selection on female ground weta, *H. pallitarsis*, and also

on a unique secondary genitalic structure they possess. Directional selection for longer accessory organs suggests

this device functions like a sexual ornament to secure matings. Although this structure does not correlate with

measures of female fecundity, it may well signal female condition (and thus perhaps her genitalic grasping ability)

or quality of care as important predictors of female reproductive success as females engage in a long and costly

maternal care period.

References

Andersson M (1994) Sexual selection. Princeton University Press, New Jersey

Arnold SJ (1994) Bateman's principles and the measurement of sexual selection in plants and
animals. Am Nat 144:126–149. https://doi.org/10.1086/285656

Arnqvist G, Rowe L (2005) Sexual Conflict. Princeton University Press, New Jersey

Aronsen T, Berglund A, Mobley KB, Ratikainen II, Rosenqvist G (2013). Sex ration and density affect sexual
selection in a sex-role reversed fish. Evolution 67:3243–3257. https://doi.org/10.1111/evo.12201

Banks HJ, Williams DJ (1972) Use of the surfactant, Decon 90, in the preparation of coccids and other insects for
microscopy. J Aust Entomol Soc 11:347-448.

Barry KL (2015) Sexual deception in a cannibalistic mating system? Testing the Femme Fatale hypothesis. Proc R
Soc B 282:20141428. https://doi.org/10.1098/rspb.2014.1428

Bateman AJ (1948) Intra-sexual selection in *Drosophila*. Heredity 2:349–368. https://doi.org/10.1038/hdy.1948.21

Berglund A, Rosenqvist G, Bernet P (1997) Ornamentation predicts reproductive success in female pipefish. Behav
Ecol Sociobiol 40:145–150. https://www.jstor.org/stable/4601312

Bohme W (1983) The Tucano Indians of Columbia and the iguanid lizard *Plica plica*: Ethnological Herpetological
and ethological implications. Biotropica 15:148-150. https://doi.org/10.2307/2387961

Bonduriansky R (2001) The evolution of male mate choice in insects: a synthesis of ideas and evidence. Biol Rev
76:305-339. https://doi.org/10.1017/S1464793101005693

Bretman A, Tregena T (2005) Measuring polyandry in wild population: a case study using promiscuous crickets.
Mol Ecol 14:2169-2179. https://doi.org/10.1111/j.1365-294X.2005.02556.x

Briceño R, Wegrzynek DD, Chinea-Cano E, Eberhard WG, dos Santos Rolo T (2010) Movements and morphology
under sexual selection: tsetse fly genitalia. Ethol Ecol Evol 22:385-391.
https://doi.org/10.1080/03949370.2010.505581

Brooks R, Caithness N (1995) Female choice in a feral guppy population: are there multiple cues? Anim Behav
50:301-307. https://doi.org/10.1006/anbe.1995.0246

Browne JH, Gwynne DT (2022) Paternity sharing in insects with female competition for nuptial gifts. Ecol Evol
12:E9463 https://doi.org/10.1002/ece3.9463

Buzkova P (2016) Interaction testing: Residuals-based permutations and parametric bootstrap in continuous count,

and binary data. Epidemiol Methods 5:119-128. https://doi.org/10.1515/em-2015-0010

Buzkova P, Lumley T, Rice K (2011) Permutation and parametric bootstrap tests for gene-gene and gene-

environment interactions. Ann Hum Genet 75:36-45. https://doi.org/10.1111/j.1469-1809.2010.00572.x

Chappell EM., Trewick SA, Morgan-Richards M (2012) Shape and sound reveal genetic cohesion not speciation in

the New Zealand orthopteran, *Hemiandrus pallitarsis*, despite high mitochondrial DNA divergence. Biol J

Linn Soc 105:169-186. https://doi.org/10.1111/j.1095-8312.2011.01777.x

Chenoweth SF, Doughty P, Kokko H (2006) Can non-directional male mating preferences facilitate honest female

ornamentation? Ecol Lett 9:179-184. https://doi.org/10.1111/j.1461-0248.2005.00867.x

Clutton-Brock T (2007) Sexual selection in males and females. Science 318:1882-1885.

https://www.science.org/doi/10.1126/science.1133311

Cumming JM (1994) Sexual selection and the evolution of dance fly mating systems (Diptera: Empididae:

Empidinae). Can Entomol 126:907-920. https://doi.org/10.4039/Ent126907-3

Dakin EE, Avise JC (2004) Microsatellite null alleles in parentage analysis. Heredity 93:504–509.

https://doi.org/10.1038/sj.hdy.6800545

Darwin C (1871) The decent of man and selection in relation to sex. John Murray, London

Downes JA (1970) The feeding and mating behaviour of the specialized Empidinae (Diptera); observations on four

species of *Rhamphomyia* in the high arctic and general discussion. Can Entomol 102:769-791.

https://doi.org/10.4039/Ent102769-7

Eberhard WG (1985) Sexual selection and animal genitalia. Harvard University Press, Massachusetts

Eberhard WG (1990) Animal genitalia and female choice. Am Sci 78:134-141. https://www.jstor.org/stable/2462660

Eberhard WG, Huber BA, Rodriguez RL, Briceño RD, Salas I, Rodriguez V (1998) One size fits all? Relationships

between the size and degree of variation in genitalia and other body parts in twenty species of insects and

spiders. Evolution 52:415-431. https://doi.org/10.2307/2411078

Fitzpatrick S, Berglund A, Rosenqvist G (1995) Ornaments or offspring: costs to reproductive success restrict sexual

selection processes. Biol J Linn Soc 55:251-260. https://doi.org/10.1111/j.1095-8312.1995.tb01063.x

Flanagan SP, Johnson JB, Rose E, Jones G (2014) Sexual selection on female ornaments in the sex-role-reversed

pipefish (*Syngnathus scovelli*). J Evol Biol 27:2457-2467. https://doi.org/10.1111/jeb.12487

Foster JC, Nan B, Shen L, Kaciroti N, Taylor JMG (2016) Permutation testing for treatment-covariate interactions

and subgroup identification. Stat in Biosci 8:77-98. https://doi.org/10.1007/s12561-015-9125-9

Fritzsche K, Arnqvist G (2013) Homage to Bateman: Sex roles predict sex differences in sexual selection. Evolution

67:1926–1936. https://doi.org/10.1111/evo.12086

Gwynne DT (1981) Sexual difference theory: Mormon crickets show role reversal in mate choice. Science

213:779-780. https://www.science.org/doi/10.1126/science.213.4509.779

Gwynne DT (1984) Sexual selection and sexual differences in Mormon crickets (Orthoptera: Tettigoniidae,

Anabrus simplex). Evolution 38:1011-1022. https://doi.org/10.2307/2408435

Gwynne DT (1995) Phylogeny of the Ensifera (Orthoptera): A hypothesis supporting multiple origins of acoustical

signalling, complex spermatophores and maternal care in crickets, katydids and weta. JOR 4:203-218.

https://doi.org/10.2307/3503478

Gwynne DT (2001) Katydids and bush-crickets: reproductive behavior and evolution of the Tettigoniidae. Cornell

University Press, New York

Gwynne DT (2002) A secondary copulatory structure in a female insect: a clasp for a nuptial meal?

Naturwissenschaften 89:125-127. https://doi.org/10.1007/s00114-002-0298-y

Gwynne DT (2004) Reproductive behavior of ground weta (Orthoptera: Anostostomatidae): Drumming behavior,

nuptial feeding, post-copulatory guarding and maternal care. J Kans Entomol Soc 77:414-428.

https://www.jstor.org/stable/25086232

Gwynne DT (2005) The secondary copulatory organ in female ground weta (H*emiandrus pallitarsis,* Orthoptera:

Anostostomatidae): A sexually selected device in females? Biol J Linn Soc 85:463-

469. https://doi.org/10.1111/j.1095-8312.2005.00510.x

Gwynne DT (2016) Sexual selection: roles evolving. Curr Biol 26:913-936.

https://doi.org/10.1016/j.cub.2016.08.063

Gwynne DT, Bailey WJ (1999) Female-female competition in katydids: Sexual selection for increased sensitivity to

a male signal? Evolution 53:546– 551. https://doi.org/10.1111/j.1558-5646.1999.tb03789.x

Gwynne DT, Simmons LW (1990) Experimental reversal of courtship roles in an insect. Nature 346:172-174.

https://doi.org/10.1038/346172a0

Hare RM, Simmons LW (2018) Sexual selection and its evolutionary consequences in female animals. Biol Rev

94:929-956. https://doi.org/10.1111/brv.12484

Hare RM, Simmons LW (2021) Sexual selection maintains a female-specific character in a species with dynamic

sex roles. Behav Ecol 32:609-616. https://doi.org/10.1093/beheco/arab005

Hare RM, Simmons LW (2022) Bateman gradients reflect variation in sexual selection in a species with dynamic

sex roles. J Evol Biol 35:1206-1217. https://doi.org/10.1111/jeb.14070

Herridge EJ (2016) Polyandry rates and reproductive success in a nuptial gift giving dance fly *Rhamphomyia*

longicauda, Dissertation. University of Stirling

Herridge EJ, Murray RL, Gwynne DT, Bussière LF (2016) Diversity in mating and parental sex roles. In: Kliman

RM (ed) Encyclopedia of evolutionary biology. Elsevier, Oxford, pp 453-458

Johns PM (2001) Distribution and conservation status of ground weta, *Hemiandrus* species (Orthoptera:

Anostostomatidae). Science for Conservation Report 180, Department of Conservation, Wellington, New

Zealand

Jones AG (2005) GERUD 2.0: A computer program for the reconstruction of parental genotypes from half-sib

progeny arrays with known or unknown parents. Mol Ecol Notes 5:708-711.

https://doi.org/10.1111/j.1471-8286.2005.01029.x

Kelly CD, Gwynne DT (2023) Mating assortment and the strength of sexual selection in a polyandrous population

of Cook Strait giant weta. Behav Ecol 34:506–513. https://doi.org/10.1093/beheco/arad017

Lande R, Arnold ST (1983) The measurement of selection on correlated characters. Evolution 37:1210-1226.

https://doi.org/10.2307/2408842

Lefèvre BM, Catté D, Courtier-Orgogoza V, Lang M (2021) Male genital lobe morphology affects the chance to

copulate in *Drosophila pachea*. BMC Ecol Evol 21:23. https://doi.org/10.1186/s12862-021-01759-z

Lloyd JE (1979) Mating behaviour and natural selection. Fla Entomol 62:17-34. https://doi.org/10.2307/3494039

Mayr E (1963) Animal species and evolution. Harvard University Press, Cambridge

Mobley KB, Jones AG (2012) Overcoming statistical bias to estimate genetic mating systems in open populations: A

comparison of Bateman's principles between the sexes in a sex-role-reversed pipefish. Evolution 67:646–

660. https://doi.org/10.1111/j.1558-5646.2012.01819.x

Morrow EH, Arnqvist G (2003) Costly traumatic insemination and female counter-adaptation in bed bugs. Proc R

Soc B 270: 2377-2381. https://doi.org/10.1098/rspb.2003.2514

Murray RL, Gwynne DT, Bussière LF (2022) Mating and sexual selection in empidine dance flies. Insects 13:839.

https://doi.org/10.3390/insects13090839

Nason SE, Kelly CD (2020) Benefits of multiple mating in a sexually dimorphic polygynandrous insect. Anim

Behav 164:65-72. https://doi.org/10.1016/j.anbehav.2020.03.018

Parker GA, Simmons LW (1996) Parental investment and the control of sexual selection: predicting the direction of

sexual competition. Proc R Soc. 263:315–321. https://www.jstor.org/stable/50614

Raymond M, Rousset F (1995) GENEPOP (version 1.2): population genetics software from exact tests and

ecumenicism. J Hered 86:248-249. https://doi.org/10.1093/oxfordjournals.jhered.a111573

Rentz DCF (1993) Tettigoniidae of Australia Volume 2: The Austrosaginae, Zaprochilinae and Phasmodinae.

CSIRO, Canberra

Robson LJ, Gwynne DT (2010) Measuring sexual selection on females in sex-role reversed Mormon crickets

(*Anabrus simplex*, Orthoptera: Tettigoniidae). J Evol Ecol 23:1528-1537. https://doi.org/10.1111/j.1420-

9101.2010.02021.x

Salmon JT (1950) A revision of the of the New Zealand weta Anostostomatidae (Orthoptera: Stenopelmatidae).

Dominion Museum Records in Entomology, ISSN 0373-7233

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. Nat Methods

9:671-675. https://doi.org/10.1038/nmeth.2089

Simmons LW (1992) Quantification of role reversal in relative parental investment in a bush cricket. Nature

358:61-63. https://doi.org/10.1038/358061a0

Simmons LW (2001) Sperm competition and its evolutionary consequences in the insects. Princeton University

Press, New Jersey

Simmons LW, Bailey WJ (1990). Resource influenced sex roles of zaprochiline tettigoniids (orthoptera:

Tettigoniidae). Evolution 44:1853–1868. https://doi.org/10.1111/j.1558-5646.1990.tb05255.x

Sloan NS, Simmons LW (2019) The evolution of female genitalia. J Evol Biol 00:1-18.

https://doi.org/10.1111/jeb.13503

Taylor Smith BL, Morgan-Richards M, Trewick SA (2013) New Zealand ground weta (Anostostomatidae: *Hemiandrus*): descriptions of two new species with notes on their biology. N Z J Zool 40:314-329. https://doi.org/10.1080/03014223.2013.804422

Taylor Smith BL (2015) Evolution of diversity: analysis of species and speciation in *Hemiandrus* ground weta. Dissertation, Massey University

Thornhill R (1983) Cryptic female choice and its implications in the scorpionfly *Harpobittacus nigriceps*. Am Nat 122:765-788. https://www.jstor.org/stable/2460916

Tobias J, Montgomerie R, Lyon BE (2012) The evolution of female ornaments and weaponry: social selection, sexual selection and ecological competition. Philos Trans R Soc B 367:2274-2293. https://doi.org/10.1098/rstb.2011.0280

Trewick SA, Bland KJ (2011) Fire and slice: palaeogeography for biogeography at New Zealand's North Island/South Island juncture. JRSNZ 42:153-183. https://doi.org/10.1080/03036758.2010.549493

Trewick SA, Taylor-Smith B, Morgan-Richards M (2020) Ecology and systematics of the wine weta and allied species, with description of four new *Hemiandrus* species. N Z J Zool 48:47-80. https://doi.org/10.1080/03014223.2020.1790396

Tripet F, Touré YT, Taylor CE, Norris DE, Dolo G, Lanzaro GC (2001) DNA analysis of transferred sperm reveals significant levels of gene flow between molecular forms of *Anopheles gambiae*. Mol Ecol 10:1725–1732. https://doi.org/10.1046/j.0962-1083.2001.01301.x

Trivers RL (1972) Parental investment and sexual selection. In: Campbell B (ed) Sexual selection and the decent of man. Aldine, Chicago, USA, pp 136-179

Vahed K, Gilbert FS (1996) Differences across taxa in nuptial gift size correlate with differences in sperm number and ejaculate volume in bushcrickets (Orthoptera: Tettigoniidae). Proc R Soc B 263:1257-1265. https://doi.org/10.1098/rspb.1996.0185

Ward RD, Skibinski DOF, Woodwark M (1992) Protein heterozygosity, protein structure and taxonomic differentiation. Evol Biol 26:73–159. http://hdl.handle.net/102.100.100/249083?index=1

Wearing-Wilde J (1996) Mate choice and competition in the barklouse *Lepinotus patruelis* (Psocoptera: Trogiidae): The effect of diet quality and sex ratio. J Insect Behav 9:599-612. https://doi.org/10.1007/BF02213883

Weissman DB (2001) Communication and reproductive behaviour in North American Jerusalem crickets. In: Field

LH (ed) The biology of wetas, king crickets and their allies. CAB International, New York, pp 351-373

Wheeler J, Gwynne DT, Bussière LF (2012) Stabilizing sexual selection for female ornaments in a dance fly. J

Evol Biol 25:1233-1242. https://doi.org/10.1111/j.1420-9101.2012.02522.x

Williams GC (1975) Sex and Evolution. Princeton University Press, New Jersey

Wulff NC, Lehmann GUC (2016) Function of male genital titillators in mating and spermatophore transfer in the

tettigoniid bushcricket *Metrioptera roeselii*. Biol J Linn Soc 117:206-216. https://doi.org/10.1111/bij.12661

Yoshizawa K, Ferreira RL, Kamimura Y, Lienhard C (2014) Female penis, male vagina, and their correlated

evolution in a cave insect. Curr Biol 24:1006-1010. https://doi.org/10.1016/j.cub.2014.03.022

Yoshizawa K, Ferreira RL, Yao I, Lienhard C, Kamimura Y (2018) Independent origins of female penis and its

coevolution with male vagina in cave insects (*Psocodea: Prionoglarididae*). Biol Lett 14:20180533.

https://doi.org/10.1098/rsbl.2018.0533