Sexual selection on a female copulatory device in an insect with nuptial gifts

Jessica H. Browne[1,2]

Darryl T. Gwynne[1,3]

[1] Department of Ecology and Evolutionary Biology, University of Toronto Mississauga,

Mississauga, Ontario L5L 1C6, Canada

[2] E-mail: jbrowne@mta.ca, Current address: Department of Biology

Mount Allison University, 62 York St. Sackville, NB, E4L 1E2, Canada

ORCID : 0000-0002-1849-5182

[3] E-mail: darryl.gwynne@utornto.ca

Keywords: sexual selection, female genitalia, ground weta, nuptial gifts, female ornaments

Male genitalia are rapidly evolving structures, often driven by sexual conflict and/or

sexual selection to increase fertilization success. Although sexual selection on females can be

strong in systems where males provide offspring care or feed their mates, sometimes resulting in

the evolution of female ornaments to acquire male investments, there are no studies of sexual

selection on female genitalia. In New Zealand ground weta, *Hemiandrus pallitarsis*, females

possess a genitalic device (the accessory organ) that is necessary for successful copulation and

the acquisition of spermatophylax glandular gifts, which are probably important in avoiding

starvation during a months-long period when caring for (their lifetime production of) eggs. Here,

we test the hypothesis that the accessory organ is a sexually selected device in *Hemiandrus*

pallitarsis by measuring the female Bateman gradient and directional sexual selection on the

accessory organ (female mating frequency estimated from microsatellite analysis of offspring

and/or stored sperm). As predicted, there was both positive Bateman gradients for females, and

evidence of directional sexual selection on accessory organ length. Although organ length did

not correlate well with female fecundity (eggs in abdomen), it may increase mating success by

indicating maternal condition and thus quality of her offspring care.

Introduction

Genitalia and associated structures (secondary genitalia) are some of the most rapidly

evolving animal traits, often exhibiting extreme variation among closely related species

(Eberhard 1985; Sloan and Simmons 2019). While genital diversity has been suggested to be a

result of species isolation ("lock and key" hypothesis; Mayr 1963; Eberhard 1985; Sloan and

Simmons 2019), genital evolution in males is thought to be driven mainly by sexual selection for

increased fertilization success (Eberhard 1985; Sloan and Simmons 2019). Because females are

typically more selective when mating (Darwin 1871; Trivers 1972), elaborate male genitalia may

have evolved in the context of cryptic female choice, where males with the most "attractive"

stimulatory genitalia experience the greatest fertilization success (Lloyd 1979; Thornhill 1983;

Eberhard 1985; Simmons 2001; Sloan and Simmons 2019). Alternatively, genitalia may function

in manipulating female mating behaviour and/or her sperm stores (e.g., grasping devices that

coerce copulations or prevent her from remating; Arnqvist and Rowe 2005). Although much of

the focus has been on male traits, females are not passive in this process. In response to, for

example, male grasping genitalia, females undergo rapid genital co-evolution to reduce potential

costs of mating (Arnqvist and Rowe 2005) or to bias fertilization success in favour of high-

quality males (Eberhard 1985; Sloan and Simmons 2019). A well-known example is the

abdominal structure of female bedbugs (*Cimex lectularius)* that minimize the harm of traumatic

insemination (penile piercing of her abdomen) by males (Morrow and Arnqvist 2003).

In contrast to these examples of "typical" mating roles, the factors influencing male and

female fitness are expected to differ when males make large investments in reproduction, such as

paternal care or nuptial gifts. Females receiving these direct benefits are expected to seek

additional matings while the potential for frequent mating by males can be limited by investment

cost, resulting in male preferences (Trivers 1972; Gwynne 1990; 1991; 2016; Gwynne and

Simmons 1990; Simmons 1992; Herridge et al. 2016) for high quality females that increase male

reproductive success (Bonduriansky 2001; Herridge et al. 2016). In these systems, especially

those where females depend on male investments for offspring production, female competition

can cause strong sexual selection on females for traits that are attractive to males (Tobias et al.

2012; Herridge et al. 2016; Hare and Simmons 2018). Well-known examples are sex-specific

female ornaments in pipefish with paternal care (abdominal color pattern; Flanagan et al. 2014)

and in empid dance flies (legs, wings, or abdomens; Cumming 1994; Murray et al 2018) where

females rely on male-provided nutrition (insect prey items) to develop their eggs (Downes 1970).

Virtually unstudied is direct sexual selection on sex specific female genitalic traits where

copulation is associated with acquisition of goods or services from males. Much like males,

female genital traits may be expected to evolve either in the context of male mate choice where

certain female genital traits have evolved such as in the context of stimulating males to invest or

to coerce the male into supplying more food. Suggested examples are (1) the interspecifically

varying ridges on female subgenital plates of Australian *Kawanaphila* katydids that grasp the

male's genitalia, perhaps allowing longer copulation duration and thus the delivery of a larger

spermatophylax gift (Gwynne 2001); and (2) a penis-like female genital modification in two

genera of psocopteran cave insects, *Neotrogla* and *Afrotrogla* where males transfer nutritious

semen to females during copulation, thought to be an important source of food (Yoshizawa et al.

2018). The penis-like gynosome, is inserted into the male's genital opening and anchors him

during copulation (Yoshizawa et al. 2014), allowing females to prolong the duration of

copulation and thus the amount of food-gift received (Yoshizawa et al. 2014; Yoshizawa et al.

2018).

Our study species has female "secondary" genitalia, abdominal devices found in a clade

of New Zealand "short-tailed" ground weta (*Hemiandrus*, Anostostomatidae, Orthoptera:

Trewick et al. 2020). Similar to *Neotroglea*, male ground weta feed their mates, in this case with

a glandular spermatophylax that the mating male adheres to the female's abdomen (separate

from the sperm capsule) and is eaten by her after copulation (Gwynne 2004). The female device

is an elongated abdominal modification on the 6th sternite, known as an accessory organ (Salmon

1950). In contrast to other species of *Hemiandrus* that, like most crickets and relatives (suborder:

Ensifera), abandon their eggs after injecting them into the substrate using long ovipositors

(Gwynne 2004; Taylor Smith et al. 2013), "short-tailed" females care for their eggs and newly-

hatched offspring (their lifetime reproduction) in an underground chamber for a period of 5-6

months (hence the "short-tailed" reduced ovipositor; Gwynne 2004; Gwynne 2005; Taylor Smith

2015). Care-providing females do not eat so male mating gifts likely represent important

nutrition to avoid starvation during the lengthy care period, potentially leading females to

compete for mating (feeding) opportunities (Gwynne 2004; Gwynne 2005). Because of this, the

accessory organ, positioned close to where the sticky spermatophylax gift is adhered, is expected

to be involved in the acquisition of gifts. This structure is not used as a holding device (as

suggested in Gwynne 2002) for the gift (Gwynne 2005).

Male genitalia in many weta (Anostostomatidae) and some related ensiferans, include genital hooks (falci) on the dorsal side of the 10^{th} segment (Johns 2001) which hook onto the edge of one of the of the female's sternites during copulation (the male is positioned beneath the female: Gwynne 2002). This is likely an ancestral behaviour, used to anchor the female for stability or to coerce her into copulation (D. Gwynne, personal communication). While some weta species have sclerotised cuticle on the sternite edge that receives the male hooks, female short-tailed ground weta are unique in possessing the accessory-organs modifications on the 6^{th} sternite, to which the male genitalia attach (D. Gwynne, personal communication). Females of our study species, *H. pallitarsis,* possess the most elaborate accessory organ (fig. 1), which protrudes from the abdomen. After pairing (female mounted above male), males first make genitalic contact with the accessory organ then move their genitalia back to attach to the female's primary genitalia when they insert the sperm ampulla (Gwynne 2004; Gwynne 2005). Following this, the male detaches from her primary genitalia, but his genitalia remain attached to her abdominal accessory organ while he extends paired phalli that deposit the bilobed spermatophylax gift (Gwynne 2002; Gwynne 2004; Gwynne 2005). The accessory organ is positioned between paired males falci pressed onto its dorsal surface, while his paired epiprocts appear to be inserted ventral to the female organ, and the distal edge of the male's 8^{th} abdominal segment pushes into the region between the female's 5^{th} and 6^{th} sternites (photographic evidence: D. Gwynne unpublished). The female accessory organ has not evolved via sexual conflict to thwart copulation attempts (as in some gerrid water striders: Arnqvist and Rowe 2005) because it is apparently necessary for mating: copulation fails when this device is experimentally removed (Gwynne 2005).

Extensive interspecific variation in the accessory organ of short-tailed species (Johns

2001; Trewick et al. 2020) suggests they are sexually-selected devices in females, possibly

driven by competition for male gifts. This variation is opposite to the usual Darwinian pattern,

where male external genitalia vary greatly across closely related species (Gwynne 2005).

Interspecific variation in female accessory organs is probably not related to species isolation

("lock and key" hypothesis; Mayr 1963; Eberhard 1985) since there is greater support for sexual

selection on complex genitalia in males (Eberhard 1985; Gwynne 2005) and, for ground weta,

mating interactions between species are more likely prevented by the species-specific

precopulatory drumming behaviour of males and females that precede pairing (Gwynne 2004;

Gwynne 2005; Chappell et al. 2012). Furthermore, there is little overlap in ground weta species

ranges (Trewick and Bland 2011; Trewick et al. 2020), which makes species mis-pairing

unlikely.

The accessory organ functions in the transfer of sperm and the gift but may also be under

sexual selection to secure mates as envisioned by Eberhard (1990) for males. Examples are the

attractive genital structures (e.g., long penes in guppies, *Poecilia reticulata*; Brooks and

Caithness 1995, brightly colored penes in some lizards; Bohme 1983) or stimulatory devices

(e.g., genital lobes in *Drosophila pachea*; Lefèvre et al. 2021, pneumopophyses in the tsetse fly

Glossina pallidipes; Briceño et al. 2012, and genital titillators in tettigoniid bushcrickets; Wulff

and Lehmann 2016) that increase female receptivity to mating or fertilization. Support for this

hypothesis for females in ground weta is that males make close contact with the accessory organ

prior to mating and have been observed to reject females after probing her mid-abdomen

(Gwynne 2005). As females with longer accessory organs tend to have a greater number of eggs

that develop to the nymphal stage (female body size controlled; Gwynne 2005), this trait may

help females obtain matings (gifts) by signalling quality to males (as in Andersson 1994;

Berglund et al. 1997; Barry 2015). Thus, this structure may have evolved via male choice in the

context of female mate/gift acquisition (Gwynne 2005),

As the female organ is inserted between two parts of the male genitalia, an alterative

hypothesis is that it has evolved to anchor between the male genital falci and epiprocts, and thus

assure spermatophylax-gift transfer, in a similar manner to the anchoring female penis of

Neotroglea bark lice that was suggested by Yoshizawa et al. (2014) as a sexually-selected organ

to obtain a nutritious sperm capsule. Gwynne's (2005) finding that accessory organ length

correlates with the number of eggs that develop to the nymphal stage is also consistent with this

hypothesis, as it may alternatively suggest that *H. pallitarsis* accessory organs result in longer

copulation and more nuptial food. Despite this, there is no direct evidence that any female

copulation devices such as accessory organs or female penes are ornaments subject to sexual

selection. Here, we test this hypothesis by measuring sexual selection on female ground weta *H.*

pallitarsis. Using estimates of female mating success from microsatellite analysis of stored

sperm and measures of lifetime female reproductive success, we determine the degree of sexual

selection on females (Bateman gradient) and estimate directional sexual selection on several

accessory organ traits. If the accessory organ is sexually-selected in the context of gift

acquisition, we predict both directional selection on accessory organ size (length; Gwynne 2005)

and that this trait will correlate with female fecundity. Characteristic of sexual selection on a sex

(Bateman 1948; Arnold 1994), we also expect there to be a positive relationship between female

mating success and reproductive success (Bateman gradient).

Methods

Specimen Collection: Bateman Gradient

To determine Bateman gradients for female ground weta *(H. pallitarsis)*, females were collected in 2002 (n=10) and 2017 (n=17) from two main sites on the North Island of New Zealand: Kiriwhakapapa trail near Masterton (-40.807627, 175.546532) and private gardens in Palmerston North (-40.413909, 175.662814). Collections took place from mid-February to early March, nearing the end of the ground weta mating season (Gwynne 2004). Live specimens were transported back to our lab in Toronto, Canada to oviposit and obtain measures of lifetime reproductive success. This was a lengthy process that involved placing females in artificial brood chambers formed from potter's clay (2002) or soil (2017) and allowing 2-3 months for them to lay eggs, as well as an additional 5-6 months for eggs to mature. Once eggs began hatching, offspring were counted (including the total number of eggs laid and the number of developed eggs: hatched nymphs or eggs with visible eye spots), and mothers were preserved with their broods at -20°C. We then dissected females to remove the spermatheca, containing the sperm stored after each mating. We isolated the sperm contents from the spermatheca using >70% ethanol, which causes sperm to harden into a pellet (Tripet et al. 2001) and stored them at -20°C. We then imaged females under a dissecting scope fitted with a digital camera and measured pronotum length as an estimate of body size (as in Taylor Smith 2015). Unfortunately, four broods were excluded from the study because females died and rapidly decayed, preventing any further analysis.

Specimen Collection: Sexual Selection on Accessory organs

To measure sexual selection on accessory organs, an additional *H. pallitarsis* females (n=58) were collected in the years 2012, 2016, 2017, and 2019 from the same two sites: Kiriwhakapapa

and Palmerston North. Following capture, specimens were killed via freezing and placed in

strong ethanol (70-95%) to be taken to Toronto, Canada. We then imaged all females under a

dissecting scope and measured them using ImageJ software. We measured pronotum length

(body size; Taylor Smith 2015) as well as accessory organ length and width (fig. 2). Prior to

measurement, all accessory organs were removed and soaked in a 5% Decon 90 solution for 1

hour at 40ºC. This was done in order to soften and restore the structures (Banks and Williams

1972), as prolonged preservation caused many to become bent and brittle. We then dissected the

spermathecae from all females, isolated the sperm contents using strong ethanol (Tripet et al.

2001), and stored them at -20°C. During the dissection, we also counted the number of

developing ova in the female's abdomen.

Locus Characterization & Microsatellite Analysis

Our microsatellite library for *H. pallitarsis* was developed by Genetic Marker Services

(Brighton, UK) and consisted of nine candidate loci. Each locus was characterized using samples

from both populations, including the females in the current study plus an additional 24 males.

After extracting DNA from hind leg tissue of adults, we conducted PCR, each reaction

containing 5.75μL sterile filtered water, 1μL 10X PCR reaction buffer, 0.2μL 10mM dNTP

mixture, 0.5μL of forward primer (each with fluorescent labels), 0.5μL reverse primer, and 1μL

of template DNA. Thermocycling was done using an Eppendorf Mastercycler with the following

temperature regime: 2 minutes at 94°C followed by 35 cycles of 30 seconds at 94°C, 30 seconds

at 60°C, and 15 seconds at 72°C, then an additional 3 minutes at 72°C. PCR products were sent

to The Centre for Applied Genomics in Sick Kids Hospital in Toronto, ON for fragment analysis

on an Applied Biosystems 3730xl capillary sequencer. The resulting electropherograms were

examined using GeneMarker V1.97, which allowed us to identify fluorescence peaks,

representing alleles. For each of the microsatellite loci, we assessed amplification quality and the

degree of polymorphism. Using GENEPOP version 4.2 (Raymond and Rousset 1995), we

determined if loci deviated from Hardy-Weinberg equilibrium (heterozygote deficiency test), and

computed estimates of null allele frequencies (Dempsters EM method) overall and for each

population separately. Finally, we assessed the degree of genetic differentiation between our two

study populations by determining F_{st} values (GENEPOP version 4.2) for each locus.

After identifying three suitable microsatellite loci (wet80, wet81, and wet89), we

conducted PCR on samples of stored sperm and offspring using the same protocol as above. We

extracted DNA from stored sperm using the Qiagen DNeasy Blood and Tissue Extraction kit.

We followed a modified version of the recommended protocol, which involved adding 12μL

DTT to each sample to improve lysis of sperm cells and dissolving DNA in a reduced volume

(50μL) of elution buffer. We extracted DNA from offspring tissue (nymph or egg) using a

Proteinase K based extraction method with multiple ethanol washes. The resulting concentration

of DNA was much higher for offspring relative to sperm and so we diluted offspring samples to

roughly 50ng/μL.

Female mating success

For measures of the Bateman gradient, we used a combination of paternity data and

analysis of stored sperm to estimate female mating frequency. Based on offspring genotypes, we

first determined the minimum number of sires in each brood using the parentage program

GERUD 2.0 (Jones 2005). Any alleles found in female's sperm stores that were not present in

the offspring genotypes were assumed to be additional mates and were factored into a female's

total mating frequency. We used a method described by Bretman and Tregenza (2005) to

estimate the most probable number of mates at each locus, given the additional alleles observed

in stored sperm (See Bretman and Tregenza 2005; Herridge 2016 thesis for details). This is

considered a less conservative method than allele counting (which assumes heterozygosity)

although it is still limited by an inability to compute multi-locus data (Bretman and Tregenza

2005; Herridge 2016) and so we used the highest estimate. One female was known to have

mated, based on the observed consumption of a spermatophylax gift; however, the male failed to

inseminate the female as his genotype did not appear in her sperm stores, nor her offspring.

Because acquisition of male-provided materials (in this case the spermatophylax gift) is the key

factor affecting female fitness (Clutton-Brock 2007), an additional mate was added to this

female's total mate number.

To measure directional sexual selection on accessory organs, we again estimated female

mating frequency using the probable mate number (Bretman and Tregenza 2005), this time using

all the male alleles detected in stored sperm. In cases where an allele was found in sperm stores

that was not previously detected in the population, it was assumed to be rare and assigned a

frequency equal to the least common allele in the population. We were unable to estimate the

mate number for four females due to poor amplification at all three loci.

Statistical Analysis: Bateman Gradients

We measured the Bateman gradient for female *H. pallitarsis* using our estimates of

mating frequency and the number of offspring in single broods each female cares for before

dying (lifetime reproductive success). We calculated standardized Bateman gradients (β'_{ss};

Bateman 1948; Arnold 1994) using separate ordinary least squares (OLS) regressions of the

relative number of mates and relative number of offspring- both the number of eggs laid, and the

number of eggs that developed to the nymphal stage (determined by hatching or visible eye

spots). Decomposition of several females prevented us from controlling for body size in our

regression model, however, we used the remaining females to test whether the inclusion of

pronotum length (an indicator of body size e.g., Taylor Smith 2015) altered our results and

whether body size alone explained significant variation in reproductive success. Significance of

β'_{ss} was determined using a generalized linear model (GLM) with Poisson error distribution of

the unstandardized traits (similar to Gerlach et al. 2012; Martin et al. 2015). We tested whether

measures of sexual selection varied by collection year or population by including interaction

terms in our regression model.

Statistical Analysis: Sexual selection on female accessory organs

We assessed the relationship between phenotype (body size, accessory organ length,

accessory organ width) and number of mates in female ground weta using multiple least squares

regression. We first z-score standardized all trait values and converted the fitness response

variable to relative mating success by dividing by the mean. Low sample size limited the

complexity of our regression model; thus, we were unable to include quadratic and correlational

terms (Lande and Arnold 1983) in our analysis. Our estimates of sexual selection only included

linear selection gradients, indicative of directional selection. Apparent differences in the strength

of sexual selection between the two collection sites led us to examine sexual selection in each

population separately. We re-standardized the data within each population and again regressed

the trait values against female mating success. Additionally, we tested if the accessory organ has

the potential to signal female fecundity by measuring the relationship between female phenotype

and the number of developing eggs in their abdomen.

Results

Locus Characterization

All microsatellite loci used in the study were polymorphic, with wet80, wet 81, and

wet89 all showing more than 20 distinct alleles (table 1). Several of the loci deviated from

Hardy-Weinberg Equilibrium (wet18, wet21, wet81, wet83, wet89), however the frequency of

null alleles was estimated to be low (< 20%; Dakin and Avise 2004) for both wet81 and wet89

(table 1), consistent across populations (table 2). Issues with unreliable or non-specific

amplification were present at three loci (wet18, wet14, wet22, and wet27), limiting their use in

future studies. Finally, the degree of genetic differentiation (F_{st}) was low between our two study

populations, ranging from 0 to 0.06 across all loci tested (table 2).

Bateman Gradients

Consistent with our prediction that female ground weta are subject to sexual selection, the

standardized Bateman gradient (table 3; Appendix) was positive for both measures of female

reproductive success, but only reached significance when using the number of developed eggs.

Estimates of β'_{ss} were similar (0.16 for eggs laid; 0.43 for eggs laid and developed) when we

controlled for body size using the available pronotum data. Consistent with Gwynne (2005),

female body size alone had no significant effect on either measure of reproductive success ($R^2 =$

0.136, p = 0.120 for eggs laid; $R^2 = 0.09$, p = 0.563 for eggs developed). When using number of

eggs to estimate reproductive success, Bateman gradients varied significantly between

populations (interaction term from GLM; p<0.0001). While mating success apparently had no

effect on the number of eggs laid had in the Palmerston North population, this relationship was

much stronger in Kiriwhakapa. Notably, estimates of β'_{ss} were significant in Kiriwhakapa

regardless of the measure of reproductive success used (eggs laid or eggs developed; table 3;

Appendix)

Sexual selection on accessory organs

Female mating success ranged from 0-6 mates (mean: 2.3 +/- 1.4 SD) and increased

significantly with accessory organ length (table 4). Larger body size and narrower accessory

organs were also associated with increased mating success; however, these trends were non-

significant. Although there was no significant interaction between phenotype and population

(OLS; body size p=0.112; accessory organ length p=0.749; accessory organ width p=0.129), we

found that selection appeared to be stronger on all traits in Palmerston North when the two

populations were analyzed separately (table 4). Although the magnitude of selection was higher

in Palmerston North, the best predictor of female mating success was accessory organ length in

both populations, followed by body size and accessory organ width (table 4; Appendix).

Interestingly, females with wider accessory organs had lower mating success in the Palmerston

North population, but marginally higher mating success in the Kiriwhakapapa population.

Accessory organs as a signal of fecundity

None of the measured traits significantly predicted the number of eggs in a female's

abdomen although those with wider accessory organs tended to be more fecund (table 5). Again,

trends differed between collection sites. In Palmerston North, larger females tended to have

somewhat greater fecundity and in Kiriwhakapapa, wider accessory organs were associated with

higher fecundity (table 5).

Discussion

In *H. pallitarsis* ground weta, sexual competition for male-provided nutrition

(spermatophylaxes) prior to an extended maternal care period without food probably creates an

opportunity for sexual selection on females when they "forage" for mating gifts. Morphological

evidence that females are subject to sexual selection comes from the existence of an apparent

female ornament, a secondary-genitalic device known as an accessory organ (Gwynne 2002;

Gwynne 2005) that varies greatly between species (Johns 2001; Trewick et al. 2020). Here, we

provide empirical evidence that *H. pallitarsis* females are subject to sexual selection, by showing

a significant positive relationship between female mating success and lifetime reproductive

success (Standardized Bateman gradient; β'_{ss}). Although estimates of the Bateman gradient were

not as high as some species where sexually selected females compete for mates, such as bean

beetles or pipefish (reviewed in Hare and Simmons 2018; Fromonteil et al. unpublished), they

were higher than the non-significant relationship between number of sires and absolute hatching

success found for females of another anostostomatid weta, *Hemideina crassidens,* where males

are sexually competitive (males fight using very large mandibular weaponry) ($\beta = 0.04$; Nason

and Kelly 2020). Measures of β'_{ss} for female *H. pallitarsis* ground weta tended to be stronger

when using the number of developed eggs to estimate reproductive success rather than just the

number of eggs laid. Because these females do not eat for several months while providing care,

we suggest that the nutritional benefits from each mating may have greater effects on female

fitness via effects on her survival and quality of care rather than on fecundity (total eggs laid). A

comparable example is found in birds, where mate-feeding increases hatching success by

reducing a mother's energetic constraints and thus improving her nest attentiveness throughout

the incubation period (Galván and Sanz 2011).

Further evidence of sexual selection on female ground weta comes from our finding that

accessory organs are subject to directional selection. There was a significant positive relationship

between the length of the accessory organ and female mating success, supporting the hypothesis

that this this structure functions as a device that ensures successful mating (there was also a non-

significant trend for larger females to mate more frequently). Sexual selection on accessory

organs appeared to be greater at the Palmerston North site, as all female traits had a stronger

effect on mating success than in Kiriwhakapapa. Despite this, none of these estimates were

significant, likely due to the small sample size in the Palmerston North population. Notably,

body size appeared to have very little effect on mating success in Kiriwhakapapa and although

non-significant, accessory organ width had an opposite (positive) effect on female mating

success relative to Palmerston North. Given the apparent lack of genetic differentiation between

our study populations (F_{st} low compared to typical values in insects; Ward et al 1992) and their

proximity (roughly 100km) relative to the total species range, it is interesting that the strength of

sexual selection, and potentially even the traits subject to sexual selection differ. There are

morphological differences between populations of *H. pallitarsis* (Chappell et al. 2012), with

Chappell et al. (2012) finding that some populations differed in size as well as the drumming

pattern performed by males prior to mating. While this variation was not congruent with genetic

differences among populations (suggesting that widely separated *H. pallitarsis* populations

comprise one interbreeding species; Chappell et al. 2012), it is possible that patterns of selection

and mate attraction differ slightly between populations of ground weta in Palmerston North

compared to Kiriwhakapapa. Regardless, the accessory organ appears to be involved in acquiring

mates and in particular, the length of the device appears to play a role in mating success in both

populations.

None of the traits measured, including female body size, were related to the number of

eggs in a female's abdomen (fecundity). However, this does not mean the accessory organ does

not signal attractiveness to males. One reason for the non-significant finding may be that the

number of eggs in the abdomen is not the main indicator of female quality for the mating male.

Because females oviposit all their eggs and then engage in a long maternal care period, during

which the mother does not eat (Gwynne 2004; J. Browne personal observation), choosy males

may prioritise female condition and thus the potential quality of care she is able to provide, as

this will determine the number of surviving offspring. Some support comes from Gwynne's

(2005) study showing a positive relationship between accessory organ length and number of

hatched nymphs rather than eggs in the abdomen. Alternatively, these females are able to

prolong copulation and thus obtain more nutrients for offspring production.

As noted previously, the accessory organ's role in competing successfully for copulations

does not exclude a mechanical function for this device (many male genitalia function in sperm

transfer as well as stimulating mates; Eberhard 1990). Given the presumed ancestral state where

males used genitalic hooks to latch onto a female's unmodified 6th sternite, the accessory organ

could have either evolved in the context of mate assessment (in the region that interacts closely

with sensitive male genitalia) or as an anchoring device between the male genital falci and

epiprocts, that assures full spermatophylax-gift transfer. The association between paired male

genitalic structures (falci, paraprocts, and the end of his 8th sternite; D. Gwynne personal

communication) almost certainly achieve stability while first the sperm capsule (ampulla) and

then the spermatophyax gift is delivered. It is possible, however, that like many male structures

(Eberhard 1990), different parts of the accessory organ serve different functions. We note for

example that the distal end of the accessory organ has a very prominent bilobed shape that

appears to align with the lobes of the males' paired paraprocts (in several *Hemiandrus* species:

Gwynne 2002). Because our measures of the accessory organ were simple linear ones, we may

not have captured the full complexity of this trait that may be important to successful mating.

Future studies should focus on quantifying the shape of this structure and its role in mating.

Another limitation of the study is that we were only able to measure directional selection

on accessory organs. Due to our limited sample size, we did not compute estimates of non-linear

or correlational selection, meaning we may have an incomplete understanding of the type of

selection acting on these structures (Lande and Arnold 1983). In particular, it would be valuable

to test for stabilizing sexual selection on female accessory organs as preference for

intermediately ornamented females has been predicted in some species (typically those with

more elaborate and costly traits) due to the energetic costs of investing in these traits (Fitzpatrick

1995; Chenoweth et al. 2006; Wheeler et al. 2012). Further, genital structures may be expected

to be under stabilizing sexual selection in cases where they have evolved to fit the average male

(Eberhard 1998).

In conclusion, we show evidence of sexual selection on female ground weta, *H.*

pallitarsis generally, and also on a secondary genitalic structure known as the accessory organ.

Directional selection for longer accessory organs suggests this device is a sexual ornament or a

coercive device used to increase mating success, however the strength of selection differed

between the two collection sites. Although this structure does not correlate with measures of

female fecundity, it may well signal female condition (and thus perhaps her genitalic grasping

ability) or quality of care as important predictors of female reproductive success as females

engage in a long and costly maternal care period.

Acknowledgements and Funding Statement

We thank Mary Morgan Richards and Lindsay Coome for their help collecting data, Doug

Currie, Marc Johnson, Rosalind Murray, and Clint Kelly for comments on the research, as well

as the New Zealand Department of Conservation (for permits). This research was supported by

an NSERC Discovery Grant (#458057) to DTG.

436   **Tables and Figures**

438
439
440   **Figure 1.** Lateral view of the extended female secondary genitalic accessory organ (6th sternite,

441   located between the hind and middle legs) in A) a newly-mated female *H. pallitarsis* and B) a

442   close-up of the organ. Photos by Darryl Gwynne.

443
444 **Figure 2.** Modified 6ᵗʰ sternite, the accessory organ, on the abdomen of female *H. pallitarsis*

445 ground weta. Image shows the structure removed from the body and the two measurements

446 taken: A) accessory organ length, from the point where the organ attaches to the body to the

447 furthest point on the distal end and B) accessory organ width, left to right on the widest point of

448 the distal end.

458   **Table 1.** Characterization of microsatellite loci developed for ground weta *H. pallitarsis*

Locus	Primer Sequence (5'-3')	bp	label	N	A	H_E	H_O	f_{null}
wet14	F: TTT TCC AAG TGG GTT ATC GTG R: CGG TAG CTA ATG GTC GGT GT	169-183	HEX	44	6	0.434	0.500	0.025
wet18	F: TGC GAT GTT ACC AAA ACA TCA T R: AGG CGT CCG ATA TCT CCT TT	96-114	HEX	32	9	0.640	0.312*	0.280
wet21	F: TAG TCC ATT ATT CAA TAC CC R: GAG AAT AAG AAG TAG CAA AAT AA	91-127	6-FAM	28	12	0.822	0.357*	0.270
wet22	F: GTA CTG CAT CTT GTC TGA GAG C R: AAG AAG TGT CCA AGG CCA TC	155-235	6-FAM	25	18	0.892	0.840	0.032
wet27	F: TCG CAA GAG TAG GCT ACG TG R: CGT GCG AGT AGG TGT TTG AG	85-97	6-FAM	38	7	0.177	0.184	0.000
wet80	F: TCA ATG AAA TGT TTC AGA CAG ACA R: GCA CAA AGA GCC ATC TGT CA	124-172	HEX	73	22	0.911	0.890	0.013
wet81	F: ACG ACG AGG AAA TGA AAT GT R: GTT GAC AAA ACT CGG GCA GT	158-226	HEX	72	24	0.917	0.736*	0.103
wet83	F: GGT AGA CAA AGC GGT TGA TG R: CAA CAT CCT CCC GTA GTT TT	120-158	6-FAM	56	5	0.650	0.411*	0.156
wet89	F: CCC TTG TTC CGT CCT ACA AG R: CAA AAC CCC ACC ATG ACT TT	180-258	6-FAM	75	21	0.920	0.773*	0.092

459   **bp**= allele size range, **N**= number of individuals tested, **A**= number of alleles detected, H_E =

460   Expected heterozygosity, H_O = Observed heterozygosity * indicates significant deviation from

461   Hardy-Weinberg equilibrium, F_{null} = frequency of null alleles

462   All estimates were obtained from GENEPOP version 4.2 (Raymond and Rousset 1995)

473 **Table 2.** Characterization of microsatellite loci for ground weta *H. pallitarsis* by population.

Locus	Kiriwhakapapa					Palmerston North					F_{st}
	N	A	H_E	H_O	f_{null}	N	A	H_E	H_O	f_{null}	
wet14	22	3	0.21	0.23	0.000	22	5	0.58	0.77	0.019	0.16
wet18	17	4	0.53	0.18*	0.230	15	7	0.71	0.47*	0.302	0.06
wet21	13	9	0.80	0.31*	0.303	15	7	0.82	0.40*	0.221	0.01
wet22	7	8	0.87	0.86	0.051	18	13	0.88	0.83	0.000	0.04
wet27	19	5	0.20	0.21	0.000	19	4	0.15	0.16	0.000	0.00
wet80	42	17	0.91	0.93	0.002	31	14	0.85	0.84	0.000	0.06
wet81	36	11	0.89	0.56*	0.172	36	20	0.92	0.92	0.007	0.03
wet83b	31	4	0.66	0.52*	0.084	25	5	0.61	0.28*	0.244	0.036
wet89	44	16	0.90	0.77*	0.085	31	12	0.88	0.77*	0.051	0.05

N= number of individuals tested**, A**= number of alleles detected, **H_E** = Expected heterozygosity,

H_o = Observed heterozygosity * indicates significant deviation from Hardy-Weinberg

equilibrium**, f_{null}** = estimated frequency of null alleles

All calculations were done using GENEPOP version 4.2 (Raymond and Rousset 1995)

Table 3. Standardized Bateman Gradients (β'_{ss}) for female ground weta, *H. pallitarsis*. Estimates

are shown overall and in each population.

Reproductive success	β'_{ss}	p-value	Kiriwhakapapa (n=12)		Palmerston North (n=11)	
			β'_{ss}	p-value	β'_{ss}	p-value
Number of eggs laid	0.12	0.072	0.44	<0.001*	-0.10	0.294
Number of developed eggs	0.39	<0.001*	0.49	<0.001*	0.36	0.001*

481 β'_{ss} = Standardized Bateman gradient

482 *Denotes significant p-values, determined using GLM with Poisson error.

Table 4. Relationship between female phenotype (traits z-score standardized) and mating success (relative number of mates) in ground weta (*H. pallitarsis*). Shown overall and by population.

Female traits	β	p-value	Kiriwhakapapa n=35		Palmerston North n=19	
			β	p-value	β	p-value
Body size	0.106	0.262	0.037	0.692	0.165	0.449
Accessory organ length	0.195	0.049*	0.103	0.301	0.234	0.318
Accessory organ width	-0.023	0.807	0.031	0.732	-0.079	0.711

β= standardized directional selection gradients

* Denotes significant p-value determined by OLS regression

Table 5. Relationship between female phenotype (traits z-score standardized) and relative number of eggs in the abdomen. Shown overall and by population.

Female traits	β_R	p-value	Kiriwhakapapa n=37		Palmerston North n=21	
			β_R	p-value	β_R	p-value
Body size	0.001	0.992	-0.077	0.596	0.147	0.610
Accessory organ length	-0.043	0.719	-0.054	0.731	-0.004	0.987
Accessory organ width	0.248	0.057	0.256	0.092	0.006	0.983

β_R = standardized regression coefficients

* Denotes significant p-value determined by OLS regression

References

Andersson, M. 1994. Sexual selection. Princeton University Press, Princeton, NJ

Arnold, S. J. 1994. Bateman's principles and the measurement of sexual selection in plants and
animals. American Naturalist 144, 126–149.

Arnqvist, G. and L. Rowe. 2005. Sexual Conflict, Princeton university Press, Princeton, NJ.

Banks, H. J. and D. J. Williams. 1972. Use of the surfactant, Decon 90, in the preparation of
coccids and other insects for microscopy. Journal of the Australian Entomological
Society 11, 347-448.

Barry, K. L. 2015. Sexual deception in a cannibalistic mating system? Testing the Femme Fatale
hypothesis. Proceedings of the Royal Society B 282, 20141428.

Bateman A. J. 1948. Intra-sexual selection in *Drosophila*. Heredity. 2: 349–368.

Berglund, A., G. Rosenqvist, and P. Bernet. 1997. Ornamentation predicts reproductive
success in female pipefish. Behavioral Ecology & Sociobiology 40, 145 - 150

Bohme, W. 1983. The Tucano Indians of Columbia and the iguanid lizard Plica plica:
Ethnological Herpetological and ethological implications. Biotropica 15, 148-150.

Bonduriansky 2001

Bretman, A. and T. Tregena. 2005. Measuring polyandry in wild population: a case study using
promiscuous crickets. Molecular Ecology 14, 2169-2179.

Briceño, R. D., D. Wegrzynek, E. Chinea-Cano, W. G. Eberhard, and T. dos Santos Rolo. 2010.
Movements and morphology under sexual selection: tsetse fly genitalia. Ethology
Ecology and Evolution 22, 385-391.

Brooks, R. and N. Caithness. 1995. Female choice in a feral guppy population: are there multiple
cues? Animal Behaviour 50, 301-307.

Chappell, E. M., S. A. Trewick, and M. Morgan-Richards. 2012. Shape and sound reveal genetic
cohesion not speciation in the New Zealand orthopteran, *Hemiandrus pallitarsis*, despite
high mitochondrial DNA divergence. Biological Journal of the Linnean Society 105, 169-
186.

Chenoweth, S. F., P. Doughty, and H. Kokko. 2006. Can non-directional male mating
preferences facilitate honest female ornamentation? Ecology Letters 9, 179-184.

Clutton-Brock, T. 2007. Sexual selection in males and females. Science 318: 1882-1885.

Cumming, J. M. 1994. Sexual selection and the evolution of dance fly mating systems

(Diptera: Empididae: Empidinae). The Canadian Entomologist 126, 907-920.

Dakin, E. E. and J. C. Avise. 2004. Microsatellite null alleles in parentage analysis. Heredity 93,

504–509.

Darwin, C. 1871. The decent of man and selection in relation to sex. London: John Murray.

Downes, J. A. 1970. The feeding and mating behaviour of the specialized Empidinae (Diptera:

Empididae: Empidinae). The Canadian Entomologist 126, 907–920.

Eberhard, W. G. 1985. Sexual selection and animal genitalia. Harvard University Press,

Massachusetts, USA.

Eberhard, W. G. 1990. Animal genitalia and female choice. American Scientist 78, 134-141.

Eberhard, W. G., B. A. Huber, R. L. Rodriguez, R. D. Briceño, I. Salas, and V. Rodriguez. 1998.

One size fits all? Relationships between the size and degree of variation in genitalia and

other body parts in twenty species of insects and spiders. Evolution 52, 415-431.

Fitzpatrick, S., A. Berglund, and G. Rosenqvist. 1995. Ornaments or offspring: costs to

reproductive success restrict sexual selection processes. Biological Journal of the Linnean

Society 55, 251-260.

Flanagan, S. P., J. B. Johnson, E. Rose, and G. Jones. 2014. Sexual selection on female

ornaments in the sex-role-reversed pipefish (*Syngnathus scovelli*). Journal of

Evolutionary Biology 27, 2457-2467.

Fromonteil, S., L. Winkler, L. Marie-Orleach, and T. Janicke. Unpublished. Sexual selection in

females across the animal tree of life. bioRxiv 2021.05.25.445581.

Funk, D. H and D. W. Tallamy. 2000. Courtship role reversal and deceptive signals in the long-

tailed dance fly, *Rhamphomyia longicauda*. Animal Behaviour 59, 411-421.

Galvan and Sanz 2011

Gerlach, N. M., J. W. McGlothlin, P. G. Parker, and E. D. Kettersona. 2012. Reinterpreting

Bateman gradients: multiple mating and selection in both sexes of a songbird species.

Behavioral Ecology 23, 1078–1088.

Gwynne, D. T. 1990. Testing parental investment and the control of sexual selection in

katydids: the operational sex ratio. American Naturalist 136, 474-484.

Gwynne, D. T. 1991. Sexual competition among females: What causes courtship-role

reversal? Trends in Ecology and Evolution 6, 118-121.

Gwynne, D. T. 2002. A secondary copulatory structure in a female insect: a clasp for a nuptial meal? Naturwissenschaften 89, 125-127.

Gwynne, D. T. 2004. Reproductive behavior of ground weta (Orthoptera: Anostostomatidae): Drumming behavior, nuptial feeding, post-copulatory guarding and maternal care. Journal of the Kansas Entomological Society 77, 414-428.

Gwynne, D. T. 2005. The secondary copulatory organ in female ground weta (H*emiandrus pallitarsis,* Orthoptera: Anostostomatidae): A sexually selected device in females? Biological Journal of the Linnean Society 85, 463-469

Gwynne, D. T. 2016. Sexual selection: roles evolving. Current Biology 26, 913-936.

Gwynne, D. T., and L. W. Simmons. 1990. Experimental reversal of courtship roles in an insect. Nature 346, 172-174.

Hare, R. M., and L. W. Simmons. 2018. Sexual selection and its evolutionary consequences in female animals. Biological Reviews 94, 929-956.

Herridge, E. J. 2016. Polyandry rates and reproductive success in a nuptial gift giving dance fly *Rhamphomyia longicauda*, PhD thesis. University of Stirling, Stirling, UK.

Herridge, E. J., R. L. Murray, D. T. Gwynne, and L. F. Bussière. 2016. Diversity in mating and parental sex roles, Pages 453-458 *in* R. M. Kliman, ed. Encyclopedia of Evolutionary Biology, Oxford Academic Press.

Johns, P. M. 2001. Distribution and conservation status of ground weta, *Hemiandrus* species (Orthoptera: Anostostomatidae). Science for Conservation Report 180, Department of Conservation, Wellington, New Zealand.

Jones, A. G. 2005. GERUD 2.0: A computer program for the reconstruction of parental genotypes from half-sib progeny arrays with known or unknown parents. Molecular Ecology Notes 5, 708-711.

Lande, R. and S. T. Arnold. 1983. The measurement of selection on correlated characters. Evolution 37, 1210-1226

Lefèvre, B. M., D. Catté, V. Courtier-Orgogoza, and M. Lang. 2021. Male genital lobe morphology affects the chance to copulate in *Drosophila pachea*. BMC Ecology and Evolution 21.

Lloyd, J. E. 1979. Mating behaviour and natural selection. The Florida Entomologist 62, 17-34.

588   Martin, A. M., M. Festa-Bianchet, D. W. Coltman, and F. Pelletier. 2015. Comparing measures

589   of breeding inequality and opportunity for selection with sexual selection on a

590   quantitative character in bighorn rams. Journal of Evolutionary Biology 28, 223-230.

591   Mayr, E. 1963. Animal species and evolution. Harvard University Press, Cambridge.

592   Morrow, E. H. and G. Arnqvist. 2003. Costly traumatic insemination and female counter-

593   adaptation in bed bugs. Proceedings of the Royal Society B 270, 2377-2381

594   Murray, R. L., J. Wheeler, D. T. Gwynne, and L. F. Bussière. 2018. Sexual selection on multiple

595   female ornaments in dance flies. Proceedings of the Royal Society B 285, 20181525

596   Raymond, M. and F. Rousset. 1995. GENEPOP (version 1.2): population genetics software from

597   exact tests and ecumenicism. Journal of Heredity 86, 248-249.

598   Salmon, J. T. 1950. A revision of the of the New Zealand weta Anostostomatidae (Orthoptera:

599   Stenopelmatidae). Dominion Museum Records in Entomology, Wellington 1, 121-177.

600   Simmons, L. W. 1992. Quantification of role reversal in relative parental investment in a bush

601   cricket. Nature 358, 61-63.

602   Simmons, L. W. 2001. Sperm competition and its evolutionary consequences in the insects.

603   Princeton University Press, New Jersey, USA.

604   Sloan, N. S. and L. W. Simmons. 2019. The evolution of female genitalia. Journal of

605   Evolutionary Biology 00, 1-18.

606   Taylor Smith, B. L., M. Morgan-Richards, and S. A. Trewick. 2013. New Zealand ground weta

607   (Anostostomatidae: *Hemiandrus*): descriptions of two new species with notes on their

608   biology. New Zealand Journal of Zoology 40, 314-329.

609   Taylor Smith, B. L. 2015. Evolution of diversity: analysis of species and speciation in

610   *Hemiandrus* ground weta. PhD thesis. Massey University, Palmerston North, NZ

611   Thornhill, R. 1983. Cryptic female choice and its implications in the scorpionfly *Harpobittacus*

612   *nigriceps*. The American Naturalist 122, 765-788.

613   Tobias, J., R. Montgomerie, and B. E. Lyon. 2012. The evolution of female ornaments and

614   weaponry: social selection, sexual selection and ecological competition. Philosophical

615   transactions of The Royal Society B 367, 2274-2293.

616   Trewick, S. A. and K. J. Bland. 2011. Fire and slice: palaeogeography for biogeography at New

617   Zealand's North Island/South Island juncture. Journal of the Royal Society New Zealand

618   iFirst, 1-31.

Trewick, S. A., B. Taylor-Smith, and M. Morgan-Richards. 2020. Ecology and systematics of the
wine weta and allied species, with description of four new *Hemiandrus* species. New
Zealand Journal of Zoology 48, 47-80.
Tripet, F., Y. T. Touré, C. E. Taylor, D. E. Norris, G. Dolo, and G. C. Lanzaro. 2001.
DNA analysis of transferred sperm reveals significant levels of gene flow between
molecular forms of *Anopheles gambiae*. Molecular Ecology 10, 1725–1732.
Trivers, R. L. 1972. Parental investment and sexual selection *in* B. Campbell, ed. Sexual
selection and the decent of man. Aldine, Chicago, USA
Ward, R. D., D. O. F. Skibinski, and M. Woodwark. 1992. Protein heterozygosity, protein
structure and taxonomic differentiation. Evolutionary Biology 26, 73–159
Wheeler, J., D. T. Gwynne, and L. F. Bussière. 2012. Stabilizing sexual selection for female
ornaments in a dance fly. Journal of Evolutionary Biology 25,1233-1242.
Wulff, N. C. and G. U. C. Lehmann. 2016. Function of male genital titillators in mating and
spermatophore transfer in the tettigoniid bushcricket *Metrioptera roeselii*. Biological
Journal of the Linnean Society 117, 206-216.
Yoshizawa, K., R. L. Ferreira, Y. Kamimura, and C. Lienhard. 2014. Female penis, male vagina,
and their correlated evolution in a cave insect. Current Biology 24, 1006-1010.
Yoshizawa, K., R. L. Ferreira, I. Yao, C. Lienhard, and Y. Kamimura. 2018. Independent origins
of female penis and its coevolution with male vagina in cave insects (*Psocodea:*
Prionoglarididae). Biology Letters 14, 20180533.

654   **Appendix:** Visualization of directional sexual selection on female *H. pallitarsis*

656   **Figure A1** Raw Bateman Gradients for female *H. pallitarsis* showing relationship between

657   number of mates and either A) number of eggs laid or B) number of developed eggs across

658   populations

660
661 **Figure A2** Thin-plate spline visualization of female body size and accessory organ length and

662 their effect on relative mating success in female ground weta (*H. pallitarsis*) collected from

663 Kiriwhakapapa. Relative mating success was estimated from the number of alleles found in a

664 female's stored sperm.

665
666 **Figure A3** Thin-plate spline visualization of female body size and accessory organ length and

667 their effect on relative mating success in female ground weta (*H. pallitarsis*) collected from

668 Palmerston North. Relative mating success was estimated from the number of alleles found in a

669 female's stored sperm.